Bounded H_{∞} -Calculus for Differential Operators on Conic Manifolds with Boundary

S. Coriasco, E. Schrohe, and J. Seiler

ABSTRACT. We derive conditions that ensure the existence of a bounded H_{∞} -calculus in weighted L_p -Sobolev spaces for closed extensions \underline{A}_T of a differential operator A on a conic manifold with boundary, subject to differential boundary conditions T. In general, these conditions ask for a particular pseudodifferential structure of the resolvent $(\lambda - \underline{A}_T)^{-1}$ in a sector $\Lambda \subset \mathbb{C}$. In case of the minimal extension they reduce to parameter-ellipticity of the boundary value problem $\binom{A}{T}$. Examples concern the Dirichlet and Neumann Laplacians.

Contents

1.	Introduction	1
2.	Differential operators on smooth manifolds with boundary	3
3.	Boundary value problems on conic manifolds	5
4.	H_{∞} -calculus of cone differential operators	10
5.	Parameter-ellipticity of the minimal extension	15
6.	Example: The Dirichlet and Neumann problems for the Laplacian	19
References		22

1. Introduction

Establishing the existence of a bounded H_{∞} -calculus is an important tool in the modern analysis of nonlinear partial differential equations. The H_{∞} -calculus was introduced by McIntosh in [18]. We refer the reader to Denk, Hieber, Prüss [5] or Kunstmann, Weis [17] for recent and extensive surveys.

Let $A : \mathcal{D}(A) \subset Y \to Y$ be a closed and densely defined operator in a Banach space Y. In the sequel, $\Lambda = \Lambda(\theta)$ will denote a sector

(1.1)
$$\Lambda = \Lambda(\theta) = \{\lambda = re^{i\varphi} \mid r \ge 0, \ \theta \le \varphi \le 2\pi - \theta\}, \qquad 0 < \theta < \pi,$$

in the complex plane. By $H = H(\theta)$ we denote the space of all holomorphic functions $\mathbb{C} \setminus \Lambda \to \mathbb{C}$ for which $|f(\lambda)| \leq c(|\lambda|^{\delta} + |\lambda|^{-\delta})^{-1}$ for some $\delta > 0$. If $\Lambda \setminus \{0\}$ is contained in the resolvent set of A and $\|\lambda(\lambda - A)^{-1}\|_{\mathcal{L}(Y)}$ is uniformly bounded on $\Lambda \setminus \{0\}$, then

(1.2)
$$f(A) := \frac{1}{2\pi i} \int_{\partial \Lambda} f(\lambda) (\lambda - A)^{-1} d\lambda, \qquad f \in H,$$

converges absolutely and defines an element in $\mathcal{L}(Y)$. By definition, the operator A has a bounded H_{∞} -calculus with respect to $\mathbb{C} \setminus \Lambda$, if

(1.3)
$$||f(A)||_{\mathcal{L}(Y)} \le M ||f||_{\infty} \quad \forall f \in H$$

with a constant M not depending on $f \in H$ (here, $||f||_{\infty}$ denotes the supremum norm of f). The name H_{∞} -calculus originates from the fact that (1.3) then allows the definition of f(A) in $\mathcal{L}(Y)$ for any bounded holomorphic function f on $\mathbb{C} \setminus \Lambda$. In particular, the choice of $f(\lambda) = \lambda^{it}, t \in \mathbb{R}$, implies the boundedness of purely imaginary powers, $A^{it} \in \mathcal{L}(Y)$, and $||A^{it}|| \leq Me^{|t|\theta}$ for all $t \in \mathbb{R}$. Establishing (1.3) requires a thorough understanding of the resolvent of A that goes well beyond proving the boundedness of $||\lambda(\lambda - A)^{-1}||_{\mathcal{L}(Y)}$.

In view of its importance for nonlinear parabolic equations, there is a vast literature concerned with the question of the existence of a bounded H_{∞} -calculus for operators A in quite different contexts. For example, Amann, Hieber, Simonett [1] treat differential operators on \mathbb{R}^n and on compact manifolds with little regularity in the coefficients; Duong in [8] considers boundary value problems on smooth manifolds, extending Seeley's work [24] on bounded imaginary powers; Denk, Dore, Hieber, Prüss, and Venni [6] then investigate boundary value problems of little regularity; in [9], Escher and Seiler consider the Dirichlet-Neumann operator for domains of low regularity; [3] shows boundedness of the imaginary powers for differential operators on manifolds with conical singularities.

In the present paper we study realizations (i.e. closed extensions) \underline{A}_T of a μ -th order cone differential operator A subject to lower order differential boundary conditions T on a manifold \mathbb{D} with conical singularities, where the boundary value problem $\binom{A}{T}$ is assumed to be Shapiro-Lopatinskii elliptic. More precisely (for an explanation of the following notation see Section 3), we consider A as an unbounded operator in a weighted L_p -space $\mathcal{H}_p^{0,\gamma}(\mathbb{D}, E)$ of sections of a vector bundle E, initially defined on a space of smooth sections that vanish under the boundary condition T. In general, Awill have a large number of closed extensions with domain contained in $\mathcal{H}_p^{\mu,\gamma}(\mathbb{D}, E)_T$, the space of sections of smoothness μ vanishing under T. We let \underline{A}_T denote one of these extensions. All closed extensions can be described explicitly, cf. [4].

In Theorem 4.1 we show that \underline{A}_T admits a bounded H_{∞} -calculus provided the resolvent of \underline{A}_T has the structure of an element of (a version of) the parameter-dependent cone calculus for pseudodifferential boundary value problems. In Section 5 we next establish conditions on $\begin{pmatrix} A \\ T \end{pmatrix}$ which are more easily verified and guarantee that the resolvent of the minimal extension $A_{T,\min}$ has the desired structure. In essence, these conditions ask for the invertibility of all parameter-dependent principal symbols associated with $\begin{pmatrix} \lambda - A \\ T \end{pmatrix}$, for then the cone calculus developed by Schulze (see e.g. [15] for a presentation) allows the construction of a parametrix which yields the resolvent to $A_{T,\min}$. Based on the results on resolvents of closed extensions of cone differential operators in [10], [11] by Gil, Krainer, and Mendoza, in [16] by Krainer, and in [22], we expect to obtain such natural conditions also for extensions different from the minimal one. This analysis, however, is not the focus of the present paper.

In Section 6 we apply these results to the minimal extension of the Laplacian on a manifold with straight conical singularity, subject to Dirichlet or Neumann conditions. We find conditions on the dimension of \mathbb{D} and on the weight γ such that $\Delta_{D,\min}$ and $\Delta_{N,\min}$ in $\mathcal{H}_p^{0,\gamma}(\mathbb{D})$ satisfy all the above assumptions. As a consequence, we derive maximal regularity for the associated initial boundary value problem.

2. Differential operators on smooth manifolds with boundary

In this section we want to recall classical results of [23], [24] and [8] on the resolvent of differential operators on manifolds with boundary and their H_{∞} -calculus.

In the following let A be a differential operator of positive order $\mu \in \mathbb{N}$ on a smooth manifold with boundary X, acting on sections into a vector bundle E,

$$A: \mathcal{C}^{\infty}(X, E) \longrightarrow \mathcal{C}^{\infty}(X, E).$$

Moreover, let $T = (T_0, \ldots, T_{\mu-1})$ be a tuple of (normal) boundary conditions, i.e.

$$T_j = \gamma_0 \circ B_j : \mathcal{C}^{\infty}(X, E) \longrightarrow \mathcal{C}^{\infty}(\partial X, F_j);$$

here, the B_j are differential operators of order $\mu - j$ acting from sections into E to sections into some hermitian vector bundle F_j and γ_0 is the operator of restriction to the boundary of X. It is allowed that some of the F_j are zero dimensional, i.e. the corresponding boundary condition T_j is void.

For a function space \mathcal{F} on X let us set

$$\mathcal{F}_T = \{ u \in \mathcal{F} \mid Tu = 0 \},\$$

provided application of T to elements of \mathcal{F} makes sense. Let us now consider A as the unbounded operator

(2.1)
$$A: \mathcal{C}^{\infty}(X, E)_T \subset L_p(X, E) \longrightarrow L_p(X, E)$$

with (a fixed) $1 . If the boundary value problem <math>\binom{A}{T}$ is (Shapiro-Lopatinskii) elliptic, the closure A_T of this operator can be shown to be defined by the action of A on the domain

(2.2)
$$\mathcal{D}(A_T) = H^{\mu}_p(X, E)_T$$

2.1. Elements of Boutet de Monvel's calculus for boundary value problems. In [2] Boutet de Monvel introduced a pseudodifferential calculus on manifolds with boundary which allowed the construction of parametrices to elliptic problems $\binom{A}{T}$. This calculus has a corresponding parameter-dependent version, some of whose elements we describe now. For a short presentation see for example [4].

Let $\Sigma \subset \mathbb{C}$ be a closed sector with vertex in the origin. We recall the definition of the space $B^{\nu,0}(X;\Sigma,E), \nu \in \mathbb{R}$, of operator-families on X,

(2.3)
$$A(\eta) = P_+(\eta) + G(\eta),$$

depending on $\eta \in \Sigma$ as parameter. The first term is a pseudodifferential operator

$$P_+(\eta) = r_+ P(\eta) e_+;$$

here, $e_+ : \mathcal{C}^{\infty}(X, E) \to \mathcal{C}^{\infty}(2X, 2E)$ denotes the operator of extending sections on X by zero to sections on the double 2X (or any other smooth closed manifold containing X as submanifold), while $r_+ : \mathcal{C}^{\infty}(2X, 2E) \to \mathcal{C}^{\infty}(X, E)$ is the operator of restriction. $P(\eta)$ is a classical, parameterdependent pseudodifferential operator of order ν on 2X. Furthermore it is required that $P(\eta)$ satisfies the *transmission condition* with respect to ∂X . Since differential operators as well as their parametrices always satisfy the transmission condition, we shall skip the details.

The second term on the right-hand side of (2.3) belongs to $B_G^{\nu,0}(X; \Sigma, E)$, the space of (singular) Green operators. Modulo regularizing operators, i.e. integral operators having a kernel in $S(\Sigma, \mathcal{C}^{\infty}(X \times X))$, each Green operator is localized in a collar neighborhood of the boundary. By standard use of a partition of unity on X, this localized part is determined by operators on the half space $\overline{\mathbb{R}}^n_+ = \{(x', x_n) \mid x' \in \mathbb{R}^{n-1}, x_n \geq 0\}$ of the form

$$\operatorname{op}'(d)(\eta) : \mathcal{C}^{\infty}_{\operatorname{comp}}(\overline{\mathbb{R}}^n_+, \mathbb{C}^q) \longrightarrow \mathcal{C}^{\infty}(\overline{\mathbb{R}}^n_+, \mathbb{C}^q), \qquad q = \dim E$$

defined by

(2.4)
$$(\operatorname{op}'(d)(\eta)u)(x,t) = \int_{\mathbb{R}^{n-1}} e^{ix\xi} \widetilde{d}(x,\xi,\eta;s,t) (\mathcal{F}_{y\to\xi}u)(\xi,s) \, ds d\xi$$

with a symbol kernel

(2.5)
$$d(x,\xi,\eta;s,t) = \widetilde{d}(x,\xi,\eta;[\xi,\eta]s,[\xi,\eta]t),$$

where \tilde{d} is a $(q \times q)$ -matrix whose components are symbols

(2.6)
$$\widetilde{d}_{ij}(x,\xi,\eta;s,t) \in S_{\rm cl}^{\nu+1}(\mathbb{R}^{n-1} \times \mathbb{R}^{n-1} \times \Sigma, \mathcal{S}(\overline{\mathbb{R}}_+ \times \overline{\mathbb{R}}_+)).$$

In (2.5) and the sequel, $[\xi, \eta]$ denotes a smoothed norm function, i.e. a smooth positive function in (ξ, η) that coincides with $|(\xi, \eta)|$ outside the unit ball. Roughly speaking, op'(d) acts as an integral operator with smooth kernel in the direction normal to the boundary, while it is a pseudodifferential operator of order ν in the tangential direction.

Both $B_G^{\nu,0}(X;\Sigma,E)$ and $B^{\nu,0}(X;\Sigma,E)$ are Fréchet spaces in a natural way, since the spaces of local symbols and symbol kernels as well as the regularizing operators carry Fréchet topologies.

2.2. Parameter-ellipticity and the resolvent of A_T . Let $\Lambda = \Lambda(\theta)$ be a sector in \mathbb{C} as introduced in (1.1). We shall call A_T , cf. (2.1) and (2.2), (parameter-)elliptic with respect to Λ if two conditions are satisfied. First, the principal symbol of A has no spectrum in Λ for non-zero covariables, i.e. in local coordinates,

(2.7)
$$\det \left(\lambda - \sigma_{\psi}^{\mu}(A)(x,\xi)\right) \neq 0 \qquad \forall \ 0 \neq (\xi,\lambda) \in \mathbb{R}^{n}_{\xi} \times \Lambda.$$

The second condition is the invertibility of the boundary symbol of $\binom{\lambda-A}{T}$; in local coordinates (x', x_n) near the boundary of X this means that, for fixed (x', ξ', λ) ,

$$\begin{pmatrix} \lambda - \sigma_{\psi}^{\mu}(A)(x', 0, \xi', D_{x_n}) \\ \left(\gamma_0 \circ \sigma_{\psi}^{j}(B_j)(x', 0, \xi', D_{x_n})\right)_{j=0,\dots,\mu-1} \end{pmatrix} : \ \mathcal{S}(\overline{\mathbb{R}}_+, \mathbb{C}^q) \longrightarrow \begin{array}{c} \mathcal{S}(\mathbb{R}_+, \mathbb{C}^q) \\ \oplus \\ \mathbb{C}^{m_0} \oplus \dots \oplus \mathbb{C}^{m_{\mu-1}} \end{array}$$

is an isomorphism whenever $(\xi', \lambda) \neq 0$. Here, $m_j = \dim F_j$ and γ_0 denotes evaluation at 0 of functions on $\overline{\mathbb{R}}_+$. Now let

(2.8)
$$\Sigma = \Sigma(\theta, \mu) = \left\{ \eta = se^{i\alpha} \mid s \ge 0, \ \frac{\theta}{\mu} \le \alpha \le \frac{2\pi - \theta}{\mu} \right\}.$$

Then Σ is a sector and $\Lambda = \{\eta^{\mu} \mid \eta \in \Sigma\}$ for Λ as in (1.1).

THEOREM 2.1. Let A_T be elliptic with respect to Λ . Then

$$\lambda - A_T : H^{\mu}_p(X, E)_T \subset L_p(X, E) \longrightarrow L_p(X, E)$$

is invertible for large $\lambda \in \Lambda$ and there exists an operator-family $P_+(\eta) + G(\eta) \in B^{-\mu,0}(X; \Sigma, E)$ such that, for large $\eta \in \Sigma$,

$$(\eta^{\mu} - A_T)^{-1} = P_+(\eta) + G(\eta).$$

This theorem was essentially proven by Seeley [23] but without using the 'language' of Boutet de Monvel's calculus. For other proofs we refer to [12] and [13].

COROLLARY 2.2. If A_T is elliptic with respect to Λ then there exists a constant $c_p \geq 0$ such that for all large $\lambda \in \Lambda$

$$\|(\lambda - A_T)^{-1}\|_{L_p(X,E)} \le c_p \frac{1}{|\lambda|}.$$

Using the above resolvent structure, [24] shows the existence of bounded imaginary powers, while [8] proves a bounded H_{∞} -calculus:

THEOREM 2.3. If A_T is elliptic with respect to Λ then there exists a constant $c \ge 0$ such that $A_T + c$ has a bounded H_{∞} -calculus with respect to $\mathbb{C} \setminus \Lambda$ (simultaneously for all 1).

3. Boundary value problems on conic manifolds

The main objective of this section is the description of boundary value problems on a manifold with conical singularities and of a parameter-dependent Boutet de Monvel calculus adapted to this situation. For simplicity we shall confine the description to the case of one conical singularity.

3.1. Cone differential operators. Let int \mathbb{D} be an (n+1)-dimensional riemannian manifold with boundary having a cylindrical end, i.e. there exists a compact subset C such that int $\mathbb{D} \setminus C$ is isometric to the product $]0,1[\times X$ for a smooth compact (not necessarily connected) riemannian manifold X with boundary. We fix the coordinate in]0,1[in such a way that every neighborhood of C in int \mathbb{D} has nonempty intersection with $]\frac{1}{2},1[\times X$. We complete int \mathbb{D} with the help of the riemannian metric. If \mathbb{D} is the resulting space, then $\mathbb{D} \setminus C$ can be identified with $[0,1[\times X]$. We next denote by int \mathbb{B} the boundary of int \mathbb{D} and by \mathbb{B} its completion in the metric inherited from int \mathbb{D} . Then $\mathbb{B} \setminus C$ can be identified with $[0,1[\times \partial X]$ and \mathbb{B} itself is a smooth manifold with boundary.

In the sequel, a vector bundle over \mathbb{D} will be a smooth hermitian vector bundle over $\operatorname{int} \mathbb{D}$ such that $E|_{]0,1[\times X}$ is isometric to the pull-back under the canonical projection $]0,1[\times X \to X \text{ of a}]$ hermitian bundle E_0 over X.

That we call \mathbb{D} a manifold with conical singularity is due to the class of differential operators we consider on it. A μ -th order differential operator A on int \mathbb{D} with smooth coefficients, acting between sections of E, is called a *cone differential operator* if

(3.1)
$$A = t^{-\mu} \sum_{j=0}^{\mu} a_j(t) (-t\partial_t)^j, \qquad a_j \in \mathcal{C}^{\infty}([0,1[,\mathrm{Diff}^{\mu-j}(X;E_0))])$$

on $]0,1[\times X]$; here we use the canonical coordinates (t, x) with $0 \le t < 1$, $x \in X$, and $\text{Diff}^k(X; E_0)$ denotes the space of k-th order differential operators on X with smooth coefficients, acting between sections of E_0 .

3.2. Function spaces. We first introduce some function spaces on \mathbb{D} (the corresponding spaces on \mathbb{B} are obtained analogously). Patching together two copies of \mathbb{D} , we obtain a smooth manifold with boundary $2\mathbb{D}$ containing \mathbb{D} . Let ω be an arbitrary fixed cut-off function on \mathbb{D} (extended by 1 to $2\mathbb{D}$).

DEFINITION 3.1. Let $\gamma \in \mathbb{R}$. Then $\mathcal{C}^{\infty,\gamma}(\mathbb{D})$ denotes the space of all smooth functions u for which $(1-\omega)u \in \mathcal{C}^{\infty}(2\mathbb{D})$ and

(3.2)
$$t \mapsto t^{\frac{n+1}{2}-\gamma} (\log t)^k (t\partial_t)^l (\omega u)(t,\cdot)$$

is a bounded function on]0,1[with values in $\mathcal{C}^{\infty}(X)$ for all $k, l \in \mathbb{N}_0$. Moreover, we set

$$\mathcal{C}^{\infty,\infty}(\mathbb{D}) = \bigcap_{\gamma \in \mathbb{R}} \mathcal{C}^{\infty,\gamma}(\mathbb{D}).$$

DEFINITION 3.2. For $s \in \mathbb{N}_0$, $\gamma \in \mathbb{R}$, and $1 , we let <math>\mathcal{H}_p^{s,\gamma}(\mathbb{D})$ be the space of all functions u such that $(1 - \omega)u \in H_p^s(2\mathbb{D})$ and

$$t^{\frac{n+1}{2}-\gamma}(t\partial_t)^l\partial_x^{\alpha}(\omega u)(t,x) \in L_p(]0,1[\times X,\frac{dt}{t}dx) \qquad \forall l+|\alpha| \le s$$

where dx refers to some metric on the manifold with boundary X.

These definitions naturally extend to sections into E and to real numbers s (however, we shall only need the case of nonnegative integers). Then we write $\mathcal{C}^{\infty,\gamma}(\mathbb{D}, E)$, $\mathcal{C}^{\infty,\infty}(\mathbb{D}, E)$, and $\mathcal{H}_{p}^{s,\gamma}(\mathbb{D}, E)$.

3.3. Elliptic boundary value problems. Let A be a μ -th order cone differential operator on \mathbb{D} as described in the introduction,

$$A: \ \mathcal{C}^{\infty,\infty}(\mathbb{D},E) \longrightarrow \mathcal{C}^{\infty,\infty}(\mathbb{D},E),$$

and $T = (T_0, \ldots, T_{\mu-1})$ be a tuple of (normal) boundary conditions, i.e.

(3.3)
$$T_j = \gamma_0 \circ B_j : \ \mathcal{C}^{\infty,\infty}(\mathbb{D}, E) \longrightarrow \mathcal{C}^{\infty,\infty}(\mathbb{B}, F_j)$$

with cone differential operators B_j of order j, and γ_0 being the operator of restriction to the boundary \mathbb{B} of \mathbb{D} . Setting $F = F_0 \oplus \ldots \oplus F_{\mu-1}$ we thus have

(3.4)
$$\begin{pmatrix} A \\ T \end{pmatrix} : \mathcal{C}^{\infty,\infty}(\mathbb{D}, E) \longrightarrow \bigoplus_{\substack{ \mathcal{C}^{\infty,\infty}(\mathbb{B}, F) \\ \mathcal{C}^{\infty,\infty}(\mathbb{B}, F)}}^{\mathcal{C}^{\infty,\infty}(\mathbb{D}, E)}$$

Let us shortly sketch what \mathbb{D} -ellipticity of $\binom{A}{T}$ means; for details we refer to Section 3.2 of [4]. First, A is elliptic on int \mathbb{D} in the standard sense, i.e. the principal symbol $\sigma_{\psi}^{\mu}(A)$ is everywhere (i.e. for non-zero covariables) invertible. In the splitting of coordinates $(t, x) \in [0, 1] \times X$ near t = 0 we have

$$\sigma^{\mu}_{\psi}(A)(t, x, \tau, \xi) = t^{-\mu} \sum_{j=0}^{\mu} \sigma^{\mu-j}_{\psi}(a_j)(t, x, \xi)(-it\tau)^j.$$

Observe a characteristic "degenerate" structure: There is the singular factor $t^{-\mu}$ and the covariable τ appears only in the form $t\tau$. Removing this degeneracy and freezing the coefficients in t = 0 we obtain the so-called *rescaled symbol*

$$\widetilde{\sigma}^{\mu}_{\psi}(A)(x,\tau,\xi) = \sum_{j=0}^{\mu} \sigma^{\mu-j}_{\psi}(a_j)(0,x,\xi)(-i\tau)^j.$$

We require this symbol to be invertible for all x and $(\tau, \xi) \neq 0$. Since $\binom{A}{T}$ is a usual boundary value problem on int \mathbb{D} , we may associate with it the standard principal boundary symbol. Ellipticity asks the invertibility of this symbol. Since the boundary conditions are also given by cone differential operators, the boundary symbol again has a degenerate structure. Removing the degeneracy and freezing coefficients in t = 0 yields the *rescaled boundary symbol* which is also required to be invertible.

3.4. Realizations of A with respect to T. For a function space \mathcal{F} on \mathbb{D} we use again the notation $\mathcal{F}_T = \{u \in \mathcal{F} \mid Tu = 0\}$. Let us now consider the unbounded operator

(3.5)
$$A: \ \mathcal{C}^{\infty,\infty}(\mathbb{D},E)_T \subset \mathcal{H}^{0,\gamma}_p(\mathbb{D},E) \longrightarrow \mathcal{H}^{0,\gamma}_p(\mathbb{D},E)$$

with $\gamma \in \mathbb{R}$ and $1 (it would be more precise to write <math>A_{\gamma,p}$ but for convenience we exclude γ and p from the notation). We shall assume that $\binom{A}{T}$ is \mathbb{D} -elliptic in the sense of Section 3.3. We write $A_{T,\min}$ for the closure of A and define $A_{T,\max}$ by the action of A on the space

$$\mathcal{D}(A_{T,\max}) = \{ u \in \mathcal{H}_p^{\mu,\gamma}(\mathbb{D}, E)_T \mid Au \in \mathcal{H}_p^{0,\gamma}(\mathbb{D}, E) \}.$$

By abuse of notation, $A_{T,\max}$ is not the true maximal extension of A, but it is the largest one that still 'feels' the boundary condition T. The closed extensions between the minimal and the maximal are usually called the *realizations* of A with respect to T. In [4], Theorem 5.12, we have shown:

THEOREM 3.3. There exists a finite dimensional space $\mathcal{E} = \mathcal{E}_{A,T} \subset \mathcal{C}^{\infty,\gamma}(\mathbb{D}, E)$ of smooth functions on int \mathbb{D} such that

$$\mathcal{D}(A_{T,\max}) = \mathcal{D}(A_{T,\min}) \oplus \mathcal{E}$$

As a consequence, any realization \underline{A}_T is determined by a subspace $\underline{\mathcal{E}}$ of \mathcal{E} , i.e.

$$\mathcal{D}(\underline{A}_T) = \mathcal{D}(A_{T,\min}) \oplus \underline{\mathcal{E}}$$

In [4] we also have characterized the domain of the closure, namely

$$\mathcal{D}(A_{T,\min}) = \left\{ u \in \bigcap_{\varepsilon > 0} \mathcal{H}_p^{\mu,\gamma+\mu-\varepsilon}(\mathbb{D}, E)_T \mid Au \in \mathcal{H}_p^{0,\gamma}(\mathbb{D}, E) \right\}$$

In particular, $\mathcal{H}_{p}^{\mu,\gamma+\mu}(\mathbb{D}, E)_{T} \subset \mathcal{D}(A_{T,\min}) \subset \mathcal{H}_{p}^{\mu,\gamma+\mu-\varepsilon}(\mathbb{D}, E)_{T}$ for any $\varepsilon > 0$. If in addition A_{T} is *conormally elliptic* with respect to the weight $\gamma + \mu$, see (E3) in Section 5 for an explanation, we even have

(3.6)
$$\mathcal{D}(A_{T,\min}) = \mathcal{H}_p^{\mu,\gamma+\mu}(\mathbb{D}, E)_T.$$

3.5. Parameter-dependent operators. A generalization of Boutet de Monvel's calculus to boundary value problems on manifolds with conical singularities was introduced in [20] and [21]. A corresponding parameter-dependent version can be found e.g. in [15]. We give here a somewhat simplified presentation of some of the elements of this calculus, focusing on the structures that are necessary for the description of resolvents. For convenience, we shall assume $E = \mathbb{C}$, since the general case is a straightforward extension of this situation.

3.5.1. Green operators. With $X^{\wedge} := \mathbb{R}_+ \times X$ and for $\gamma \in \mathbb{R}$ let us set

$$(3.7) \qquad \mathcal{H}_{p}^{0,\gamma}(X^{\wedge}) = L_{p}(X^{\wedge}, t^{(\frac{n+1}{2}-\gamma)p}\frac{dt}{t}dx) = t^{\gamma-\frac{n+1}{2}}L_{p}(X^{\wedge}, \frac{dt}{t}dx),$$

$$(3.8) \qquad \mathcal{S}_{0}^{\gamma}(X^{\wedge}) = \{u \in \mathcal{C}^{\infty}(X^{\wedge}) \mid \omega u \in \mathcal{C}^{\infty,\gamma}(\mathbb{D}) \text{ and } (1-\omega)u \in \mathcal{S}(\mathbb{R}, \mathcal{C}^{\infty}(X))\}.$$

The latter is a Fréchet space in a natural way.

Let Σ be a closed subsector of \mathbb{C} with vertex in 0 and $\nu, \gamma \in \mathbb{R}$. Then we denote by $R_G^{\nu,0}(X^{\wedge}; \Sigma, \gamma)$ the space of all operator families $G(\eta), \eta \in \Sigma$, such that

$$\left(G(\eta)u \right)(t,x) = \left\langle k(\eta;t,x), \overline{u} \right\rangle_{\mathcal{H}_{2}^{0,0}(X^{\wedge})} = \int_{X^{\wedge}} k(\eta;t,x,t',x')u(t',x') t'^{n} dt' dx',$$

where the integral kernel k has the form

(3.9)
$$k(\eta; t, x, t', x') = \widetilde{k}(\eta; [\eta]t, x, [\eta]t', x')$$

with

(3.10)
$$\widetilde{k}(\eta; t, x, t', x') \in S^{\nu+n+1}\left(\Sigma, \mathcal{S}_0^{\gamma+\varepsilon}(X^{\wedge}_{(t,x)})\widehat{\otimes}_{\pi} \mathcal{S}_0^{-\gamma+\varepsilon}(X^{\wedge}_{(t',x')})\right)$$

for some $\varepsilon > 0$ (depending on G).

3.5.2. Mellin pseudodifferential operators. For $\beta \in \mathbb{R}$ let $\Gamma_{\beta} = \{z \in \mathbb{C} \mid \text{Re } z = \beta\}$. We denote by $M_{\beta}^{\nu,0}(X; \Sigma), \nu \in \mathbb{R}$, the space of all functions $h: \Gamma_{\beta} \times \Sigma \to B^{\nu,0}(X)$ such that

$$h(\beta + i\tau, \eta) \in B^{\nu,0}(X; \mathbb{R}_{\tau} \times \Sigma_{\eta})$$

(cf. Section 2.1 with parameter space $\mathbb{R} \times \Sigma$ instead of Σ). A *Mellin symbol* is a function $h = h(t, \beta + i\tau, \eta) \in \mathcal{C}^{\infty}(\mathbb{R}_+, M^{\nu,0}_{\beta}(X; \Sigma))$; it induces the Mellin pseudodifferential operator $\operatorname{op}_M^{\frac{1}{2}-\beta}(h)(\eta) : \mathcal{C}^{\infty}_{\operatorname{comp}}(X^{\wedge}) \to \mathcal{C}^{\infty}(X^{\wedge})$ by

$$\left(\operatorname{op}_{M}^{\frac{1}{2}-\beta}(h)(\eta)u\right)(t) = \int_{-\infty}^{\infty} t^{-\beta-i\tau} h(t,\beta+i\tau,\eta)(\mathcal{M}u)(\beta+i\tau)\,d\tau;$$

here we have identified $\mathcal{C}^{\infty}(X^{\wedge})$ with $\mathcal{C}^{\infty}(\mathbb{R}_+, \mathcal{C}^{\infty}(X))$, and \mathcal{M} denotes the Mellin transform, i.e. $(\mathcal{M}u)(z) = \int_0^{\infty} t^z u(t) \frac{dt}{t} \text{ for } z \in \mathbb{C}.$

3.5.3. The full class and a norm estimate. We define $C^{\nu,0}(\mathbb{D}; \Sigma, \gamma)$ with $\nu \leq 0$ and $\gamma \in \mathbb{R}$ as the set of all operator-families $A(\eta)$ of the form

(3.11)
$$A(\eta) = \sigma_0 \left\{ t^{-\nu} \mathrm{op}_M^{\gamma - \frac{n}{2}}(h)(\eta) + G(\eta) \right\} \sigma_1 + (1 - \sigma_3) P(\eta)(1 - \sigma_4) + G_\infty(\eta),$$

where $\sigma_j \in \mathcal{C}^{\infty}_{\text{comp}}([0,1[) \text{ are cut-off functions and }$

- a) $h(t,z,\eta) = \widetilde{h}(t,z,t\eta)$ with $\widetilde{h} \in \mathcal{C}^{\infty}(\overline{\mathbb{R}}_+, M^{\nu,0}_{\frac{n+1}{2}-\gamma}(X;\Sigma))$, cf. Section 3.5.2,
- b) $P(\eta) \in B^{\nu,0}(2\mathbb{D}; \Sigma)$, cf. Section 2.1,
- c) $G(\eta) \in R_G^{\nu,0}(X^{\wedge}; \Sigma, \gamma)$, cf. Section 3.5.1,
- d) $G_{\infty} \in \mathcal{S}(\Sigma, \mathcal{L}(\mathcal{H}_{p}^{s,\gamma}(\mathbb{D}), \mathcal{H}_{p}^{s',\gamma}(\mathbb{D})))$ for all $1 and <math>s, s' \in \mathbb{R}$ with $s > -1 + \frac{1}{p}$.

Initially, each such operator is defined on smooth functions, i.e. $A(\eta) : \mathcal{C}^{\infty}_{\text{comp}}(\text{int }\mathbb{D}) \to \mathcal{C}^{\infty}(\text{int }\mathbb{D})$, where multiplications by σ_1 and σ_0 are viewed as operators $\mathcal{C}^{\infty}_{\text{comp}}(\text{int }\mathbb{D}) \to \mathcal{C}^{\infty}_{\text{comp}}(X^{\wedge})$ and $\mathcal{C}^{\infty}_{\text{comp}}(X^{\wedge}) \to \mathcal{C}^{\infty}_{\text{comp}}(\text{int }\mathbb{D})$, respectively. Similarly, $1 - \sigma_3$ and $1 - \sigma_4$ act between smooth functions on \mathbb{D} and $2\mathbb{D}$.

Let $A(\eta) \in C^{\nu,0}(\mathbb{D}; \Sigma, \gamma)$ as in (3.11). We deduce from (2.3) that there is a decomposition of \tilde{h} as

(3.12)
$$h(t, \frac{n+1}{2} - \gamma + i\tau, \eta) = \widetilde{p}_+(t, \tau, \eta) + \widetilde{g}(t, \tau, \eta)$$

where $t \mapsto \tilde{p}(t,\tau,\eta)$ is a smooth (up to t=0) family of parameter-dependent pseudodifferential operators of order ν on 2X with parameter space $\mathbb{R} \times \Sigma$, while $\tilde{g}(t,\tau,\eta)$ belongs to $\mathcal{C}^{\infty}(\overline{\mathbb{R}}_+, B_G^{\nu,0}(X; \mathbb{R} \times \Sigma))$. In local coordinates,

(3.13)
$$\widetilde{p}_{+}(t,\tau,\eta) = \operatorname{op}_{+}(\widetilde{a})(t,\tau,\eta) \text{ with } \widetilde{a}(t,x,\tau,\xi,\eta) \in S_{\operatorname{cl}}^{\nu}(\overline{\mathbb{R}}_{+} \times \mathbb{R}_{x}^{n} \times \mathbb{R}_{(\tau,\xi)}^{1+n} \times \Sigma)$$

and $\widetilde{g}(t,\tau,\eta)$ is defined via a symbol kernel

(3.14)
$$\widetilde{d}(t, x', \tau, \xi', \eta; u, v) = d(t, x', \tau, \xi', \eta; [\tau, \xi', \eta] u, [\tau, \xi', \eta] v)$$

with

(3.15)
$$d(t, x', \tau, \xi', \eta; u, v) \in S^{\nu+1}(\overline{\mathbb{R}}_+ \times \mathbb{R}^{n-1}_{x'} \times \mathbb{R}^n_{(\tau, \xi')} \times \Sigma; \mathcal{S}(\overline{\mathbb{R}}_+ \times \overline{\mathbb{R}}_+)).$$

The non-local, smoothing terms arising are as above in d).

THEOREM 3.4. Let $A(\eta) \in C^{\nu,0}(\mathbb{D}; \Sigma, \gamma), \nu \leq 0$. Then $A(\eta)$ extends for each η to a bounded operator in $\mathcal{H}^{0,\gamma}_p(\mathbb{D})$ and there exists a constant $c_p \geq 0$ such that

$$\|A(\eta)\|_{\mathcal{L}(\mathcal{H}_{p}^{0,\gamma}(\mathbb{D}))} \leq c_{p} \langle \eta \rangle^{\nu} \qquad \forall \eta \in \Sigma.$$

PROOF. The proof is a combination of that for the boundaryless case (cf. Proposition 1 in [3]) and the results of [14] on the L_p -continuity of operators from Boutet's calculus. To give an idea, we shall derive the desired norm estimate for the operator $\sigma_0 t^{-\nu} \operatorname{op}_M^{\gamma-\frac{n}{2}}(g) \sigma_1$ with $g(t,\tau,\eta) = \tilde{g}(\tau,t\eta)$ as in (3.12) (the *t*-independence of \tilde{g} can always be reached by a tensor product argument). By conjugation with *t*-powers, it is no restriction to assume that $\gamma = (n+1)(\frac{1}{2} - \frac{1}{p})$. With this choice of γ , we have $\mathcal{H}_p^{0,\gamma}(X^{\wedge}) = L_p(X^{\wedge}, t^n dt dx)$ and the operators

$$\kappa_{\varrho} \in \mathcal{L}(\mathcal{H}_{p}^{0,\gamma}(X^{\wedge})), \text{ defined by } (\kappa_{\varrho}u)(t,x) = \varrho^{\frac{n+1}{p}}u(\varrho t,x),$$

provide bijective isometries for each $\rho > 0$. Moreover, defining $(Su)(r, x) = e^{(\frac{n}{2} - \gamma)r}u(e^{-r}, x)$ for functions u on X^{\wedge} , we obtain an isometric isomorphism

$$S: \mathcal{H}^{0,\gamma}_p(X^{\wedge}) \longrightarrow L_p(\mathbb{R} \times X, drdx).$$

Thus we are done if we can prove that

$$\langle \eta \rangle^{-\nu} \| S \kappa_{\langle \eta \rangle}^{-1} t^{-\nu} \operatorname{op}_{M}^{\gamma - \frac{n}{2}}(g)(\eta) \kappa_{\langle \eta \rangle} S^{-1} \|_{\mathcal{L}(L_{p}(\mathbb{R} \times X, drdx))} \leq c_{p},$$

uniformly in η . Now note that conjugation of $\operatorname{op}_{M}^{\gamma-\frac{n}{2}}(g)(\eta)$ by $\kappa_{\langle\eta\rangle}$ amounts to replacing $g(t, z, \eta)$ by $g(\frac{t}{\langle\eta\rangle}, z, \eta)$, and conjugation with S transforms a Mellin operator on X^{\wedge} with symbol $h(t, \frac{n+1}{2} - \gamma + i\tau)$ into a pseudodifferential operator on $\mathbb{R} \times X$ with symbol $a(r, \varrho) = h(e^{-r}, \frac{n+1}{2} - \gamma + i\varrho)$. Passing to the local situation, cf. (3.14) and (3.15), we have to show that $\|\operatorname{op}'(d_0)(\eta)\|_{\mathcal{L}(L_p(\mathbb{R}^{n+1}_+))} \leq c_p$ for the symbol kernel

$$d_0(r, x', \tau, \xi', \eta; u, v) = e^{r\nu} \widetilde{d}(x', \tau, \xi', e^{-r} \frac{\eta}{\langle \eta \rangle}; [\tau, \xi', e^{-r} \frac{\eta}{\langle \eta \rangle}] u, [\tau, \xi', e^{-r} \frac{\eta}{\langle \eta \rangle}] v),$$

see (2.4) for the definition of op'. By a straightforward calculation,

$$\|u^{k'}D_{u}^{k}v^{l'}D_{v}^{l}D_{r}^{m'}D_{\tau}^{m}D_{x'}^{\alpha'}D_{\xi'}^{\alpha}d_{0}(r,x',\tau,\xi',\eta;u,v)\|_{L^{2}(\mathbb{R}_{+,u}\times\mathbb{R}_{+,v})} \leq C \langle \tau,\xi' \rangle^{-|\alpha|-m+k-k'+l-l'},$$

¹By definition, $S_{cl}^{\nu}(\overline{\mathbb{R}}_{+} \times \mathbb{R}_{x}^{n} \times \mathbb{R}_{(\tau,\xi)}^{l+n} \times \Sigma)$ denotes the space of all functions *b* that are smooth on $\overline{\mathbb{R}}_{+} \times \mathbb{R}_{x}^{n} \times \mathbb{R}_{(\tau,\xi)}^{l+n} \times \Sigma$ satisfy there uniform estimates $|\partial_{t}^{l}\partial_{x}^{\beta}\partial_{(\tau,\xi,\eta)}^{\alpha}b(t,x,\tau,\xi,\eta)| \leq c \langle \tau,\xi,\eta \rangle^{\nu-|\alpha|}$ for any order of derivatives and that have asymptotic expansions into components that are positively homogeneous in (τ,ξ,η) .

uniformly in $(r, x', \tau, \xi', \eta)$. Now Theorem 4.1.(5) in [14] gives the continuity of $\operatorname{op}'(d_0)(\eta)$ in $L_p(\overline{\mathbb{R}}^{n+1}_+)$, with operator norm uniformly bounded in η .

The local symbols \tilde{a} in (3.13) are defined for $(\tau, \xi, \eta) \in \mathbb{R}^{1+n} \times \Sigma$. The symbols that arise in the analysis of resolvents extend – holomorphically in η – to larger subsets of \mathbb{C} . We shall need this property in particular for the principal part and define a corresponding class:

DEFINITION 3.5. For $c_0 > 0$ let $\Omega_{(\tau,\xi)} = \{\eta \in \mathbb{C} \mid |\eta| \ge c_0 \langle \tau, \xi \rangle\}$. The set $C_{\text{hol}}^{\nu,0}(\mathbb{D}; \Sigma, \gamma)$ consists of all operator-families $A(\eta) \in C^{\nu,0}(\mathbb{D}; \Sigma, \gamma)$ such that the local symbols \tilde{a} from (3.13) admit a decomposition $\tilde{a} = \tilde{a}_0 + \tilde{a}_1$ with the following properties:

a) \tilde{a}_0 extends holomorphically to $\eta \in \Omega_{(\tau,\xi)}$ (for some c_0) and satisfies the estimates

$$(3.16) \qquad \qquad |\partial_t^l \partial_x^\beta \partial_\tau^k \partial_\xi^\alpha \partial_\eta^\gamma \,\widetilde{a}_0(t, x, \tau, \xi, \eta)| \le c \,\langle \tau, \xi, \eta \rangle^{\nu-k-|\alpha|-|\gamma|}$$

uniformly in $(\tau,\xi) \in \mathbb{R}^{1+n}$ and $\eta \in \Omega_{(\tau,\xi)} \cup \Sigma$ for any order of derivatives, b) $\widetilde{a}_1(t,x,\tau,\xi,\eta) \in S_{\mathrm{cl}}^{\nu-1}(\overline{\mathbb{R}}_+ \times \mathbb{R}^n_x \times \mathbb{R}^{1+n}_{(\tau,\xi)} \times \Sigma_\eta).$

For the pseudodifferential symbols of $P(\eta)$ in (3.11) we require the corresponding structure.

4. H_{∞} -calculus of cone differential operators

The aim of this section is to prove the following theorem:

THEOREM 4.1. If the resolvent set of the realization \underline{A}_T contains $\{\lambda \in \Lambda \mid |\lambda| \geq R\}$ for some $R \geq 0$ and there exists an $A(\eta) \in C^{-\mu,0}(\mathbb{D}; \Sigma, \gamma)$ such that

$$(\eta^{\mu} - \underline{A}_T)^{-1} = A(\eta) \qquad \forall \ |\eta| \ge R^{1/\mu},$$

then there exists a constant $c \ge 0$ such that $c + \underline{A}_T$ admits a bounded H_∞ -calculus with respect to $\mathbb{C} \setminus \Lambda$ (simultaneously for all 1).

Before going into details, let us give an outline of the proof. In large parts it follows the proof of Theorem 5.1 in [3], where we showed the existence of bounded imaginary powers for operators on conic manifolds without boundary.

By replacing from the very beginning the differential operator A by c + A we can assume that the resolvent set contains all of Λ and

$$A(\eta) := (\eta^{\mu} - \underline{A}_T)^{-1} \in C^{-\mu,0}(\mathbb{D}; \Sigma, \gamma).$$

We then show that \underline{A}_T itself has a bounded H_{∞} -calculus. Let us remark that, for $f \in H$,

$$f(\underline{A}_T) = \int_{\partial \Lambda} f(\lambda) (\lambda - \underline{A}_T)^{-1} d\lambda = \int_{\partial \Sigma} f(\eta^{\mu}) A(\eta) \eta^{\mu - 1} d\eta.$$

Inserting the explicit formula for $A(\eta)$, cf. (3.11), we obtain four integrals over the boundary of Σ which have to be estimated from above by $M||f||_{\infty}$ with a constant M independent of $f \in H$, cf.(1.3).

Obviously, the integral associated with $G_{\infty}(\eta)$ can be estimated as desired, since $\eta^{\mu-1}G_{\infty}(\eta)$ is reapidly decreasing in η , hence integrable.

For the integral associated with $(1 - \sigma)P(\eta)(1 - \sigma_1)$ we can apply Theorem 2.3, since away from the singularity the Sobolev spaces and operator classes coincide with the usual ones on the smooth manifold with boundary 2D. In Propositions 4.3, 4.4, and 4.5 we shall treat the term

(4.1)
$$\sigma_0 \int_{\partial \Sigma} f(\eta^{\mu}) t^{\mu} \mathrm{op}_M^{\gamma - \frac{n}{2}}(h)(\eta) \eta^{\mu - 1} d\eta \,\sigma_1 = \mathrm{op}_M^{\gamma - \frac{n}{2}} \Big(\sigma_0 t^{\mu} \int_{\partial \Sigma} f(\eta^{\mu}) h(\eta) \eta^{\mu - 1} d\eta \Big) \sigma_1.$$

First, however, we shall study the term induced by $G(\eta)$. To this end note that multiplication by a cut-off function $\sigma \in \mathcal{C}^{\infty}_{\text{comp}}([0,1[)$ induces continuous operators $\mathcal{H}^{0,\gamma}_p(\mathbb{D}) \to \mathcal{H}^{0,\gamma}_p(X^{\wedge})$ and $\mathcal{H}^{0,\gamma}_p(X^{\wedge}) \to \mathcal{H}^{0,\gamma}_p(\mathbb{D})$, cf. (3.7). Thus it suffices to show the following:

PROPOSITION 4.2. Let $G(\eta) \in R_G^{-\mu,0}(X^{\wedge}; \Sigma, \gamma)$ and $G_f = \sigma_0 \int_{\partial \Sigma} f(\eta^{\mu}) G(\eta) \eta^{\mu-1} d\eta \sigma_1$ for $f \in H$. Then $G_f \in \mathcal{L}(\mathcal{H}_p^{0,\gamma}(X^{\wedge}))$, and there exists a constant $M_p \ge 0$ such that

(4.2)
$$\|G_f\|_{\mathcal{L}(\mathcal{H}_p^{0,\gamma}(X^{\wedge}))} \le M_p \|f\|_{\infty} \quad \forall f \in H.$$

PROOF. By conjugation with t^{γ} we can assume that $\gamma = 0$. Due to the symmetry of $\partial \Sigma$ and the fact that $\eta^{\mu-1}G(\eta)$ is integrable on compact parts of $\partial \Sigma$, we may confine ourselves to the integration over the set

$$\mathcal{C} = \{ \varrho e^{i\alpha} \mid 1 \le \varrho < \infty \}.$$

According to Section 3.5.1, $G(\eta)$ is an integral operator (with respect to the scalar product in $\mathcal{H}_2^{0,0}(X^{\wedge})$). Suppressing, for notational simplicity, the dependence on the variables x, x', the kernel of $G(\eta), |\eta| \geq 1$ is $k(\eta, t, s) = \tilde{k}(\eta, |\eta| t, |\eta| s)$, where, for some $\varepsilon > 0$,

$$\widetilde{k}(\eta, t, s) \in S^{-\mu+n+1}(\Sigma) \widehat{\otimes}_{\pi} \mathcal{S}_{0}^{\varepsilon}(X^{\wedge}) \widehat{\otimes}_{\pi} \mathcal{S}_{0}^{\varepsilon}(X^{\wedge}).$$

Then G_f the an integral operator with kernel

(4.3)
$$k_f(t,s) = \sigma_0(t) \,\sigma_1(s) \int_{\mathcal{C}} f(\eta^{\mu}) k(\eta,t,s) \,\eta^{\mu-1} \,d\eta$$

Writing $\tilde{k}(\eta, t, s) = (\chi(t) + (1 - \chi)(t))\tilde{k}(\eta, t, s)(\chi(s) + (1 - \chi)(s))$ with the characteristic function χ of [0, 1], the proposition will be true, if we can show that in each one of the four cases

(4.4)
$$k_f(t,s) = \sigma_0(t) \,\sigma_1(s) \int_{\mathcal{C}} f(\eta^{\mu}) \chi(|\eta| \, t) k(\eta, t, s) \chi(|\eta| \, s) \, \eta^{\mu-1} \, d\eta$$

(4.5)
$$k_f(t,s) = \sigma_0(t) \,\sigma_1(s) \int_{\mathcal{C}} f(\eta^{\mu}) \chi(|\eta| \, t) k(\eta,t,s) (1-\chi)(|\eta| \, s) \, \eta^{\mu-1} \, d\eta$$

(4.6)
$$k_f(t,s) = \sigma_0(t) \,\sigma_1(s) \int_{\mathcal{C}} f(\eta^{\mu}) (1-\chi) (|\eta| \, t) k(\eta,t,s) \chi(|\eta| \, s) \, \eta^{\mu-1} \, d\eta$$

(4.7)
$$k_f(t,s) = \sigma_0(t) \,\sigma_1(s) \int_{\mathcal{C}} f(\eta^{\mu}) (1-\chi) (|\eta| \, t) k(\eta,t,s) (1-\chi) (|\eta| \, s) \, \eta^{\mu-1} \, d\eta$$

the associated integral operators satisfies (4.2). To begin with case (4.4) we use the fact that, for some fixed $\varepsilon > 0$,

$$|k(\eta, t, s)| \le c|\eta|^{-\mu + 2\varepsilon} t^{-\frac{n+1}{2} + \varepsilon} s^{-\frac{n+1}{2} + \varepsilon},$$

uniformly in $\eta \in \mathcal{C}$ and t, s > 0, cf. (3.8) and (3.2). Then

$$|k_f(t,s)| \le c \, \|f\|_{\infty} \, \sigma_0(t) \, \sigma_1(s) \, t^{-\frac{n+1}{2}+\varepsilon} s^{-\frac{n+1}{2}+\varepsilon} \int_1^\infty \varrho^{-1+2\varepsilon} \chi(\varrho t) \chi(\varrho s) \, d\varrho$$
$$\le \|f\|_{\infty} \, \frac{c}{2\varepsilon} \, \min\left(\frac{1}{t}, \frac{1}{s}\right)^{2\varepsilon} \sigma_0(t) \, \sigma_1(s) \, t^{-\frac{n+1}{2}+\varepsilon} s^{-\frac{n+1}{2}+\varepsilon}.$$

Since the factor $\frac{c}{2\varepsilon}\sigma_0(t)\sigma_1(s)$ can be estimated from above by a constant, it remains to consider the kernel $t^{-\frac{n+1}{2}+\varepsilon}s^{-\frac{n+1}{2}+\varepsilon}\min(\frac{1}{t},\frac{1}{s})^{2\varepsilon}$. Because this kernel is symmetric in s and t, indeed it suffices to treat

(4.8)
$$k(t,s) = \begin{cases} t^{-\frac{n+1}{2} - \varepsilon} s^{-\frac{n+1}{2} + \varepsilon} & : s \le t \\ 0 & : s > t. \end{cases}$$

Recalling the Hardy inequality

$$\int_0^\infty \left(\int_0^t g(s)\,ds\right)^p t^{-1-r}\,dt \le \left(\frac{p}{r}\right)^p \int_0^\infty g(t)^p t^{p-1-r}\,dt,$$

which holds for any non-negative function g on \mathbb{R}_+ and r > 0 (cf. [26], Lemma 3.14, page 196), and denoting by G the integral operator with kernel (4.8), we have

$$\begin{split} \|Gu\|_{\mathcal{H}_{p}^{0,0}(X^{\wedge})}^{p} &\leq \int_{0}^{\infty} \left(\int_{0}^{\infty} k(t,s)|u(s)|\,s^{n}ds\right)^{p} t^{\frac{n+1}{2}p-1} \, dt = \int_{0}^{\infty} \left(\int_{0}^{t} s^{\frac{n-1}{2}+\varepsilon}|u(s)|\,ds\right)^{p} t^{-1-p\varepsilon} \, dt \\ &\leq \left(\frac{p}{p\varepsilon}\right)^{p} \int_{0}^{\infty} |u(t)|^{p} \, t^{\frac{n+1}{2}p-1} dt = \left(\frac{1}{\varepsilon}\right)^{p} \|u\|_{\mathcal{H}_{p}^{0,0}(X^{\wedge})}^{p}, \end{split}$$

which completes the proof for the case (4.4). The proofs for the cases (4.5), (4.6) and (4.7) can be obtained similarly, cf. [3]. \Box

Let us now turn our attention to the analysis of (4.1). We make use of the structure of h as described in (3.13), (3.14), (3.15), and Definition 3.5. Then we have to handle two terms involving \tilde{a}_0 and \tilde{a}_1 , and a term coming from the symbol kernel \tilde{d} .

Let us recall that any Mellin symbol $h \in MS^{\nu}(\mathbb{R}_+ \times \mathbb{R}^n \times \Gamma_{\beta} \times \mathbb{R}^n)$, i.e. any smooth function satisfying, for any order of derivatives,

$$|(t\partial_t)^k \partial_x^{\alpha'} \partial_\tau^l \partial_\xi^\alpha h(t, x, \beta + i\tau, \xi)| \le c \langle \tau, \xi \rangle^{\nu - l - |\alpha|},$$

induces continuous operators from $\mathcal{H}_p^{s,\frac{n+1}{2}-\beta}(\mathbb{R}_+\times\overline{\mathbb{R}}_+^n)$ to $\mathcal{H}_p^{s-\nu,\frac{n+1}{2}-\beta}(\mathbb{R}_+\times\overline{\mathbb{R}}_+^n)$.

PROPOSITION 4.3. Let $\tilde{a}_1 \in S^{-\mu-1}(\overline{\mathbb{R}}_+ \times \mathbb{R}^n \times \mathbb{R}^{1+n} \times \Sigma)$ be compactly supported in (t, x) and

$$h_f(t, x, \frac{n+1}{2} - \gamma + i\tau, \xi) = t^{\mu} \int_{\partial \Sigma} f(\eta^{\mu}) \,\widetilde{a}_1(t, x, \tau, \xi, t\eta) \, \eta^{\mu-1} \, d\eta$$

with $f \in H$. This defines a symbol $h_f \in MS^{-1}(\mathbb{R}_+ \times \mathbb{R}^n \times \Gamma_{\frac{n+1}{2}-\gamma} \times \mathbb{R}^n)$ and the symbol estimates of $\|f\|_{\infty}^{-1} h_f$ are uniform in $0 \neq f \in H$. Consequently,

$$\|\mathrm{op}_{M}^{\gamma - \frac{\mu}{2}} \mathrm{op}_{+}(h_{f})\|_{\mathcal{L}(\mathcal{H}_{p}^{0,\gamma}(\mathbb{R}_{+} \times \overline{\mathbb{R}}_{+}^{n}))} \leq M_{p} \|f\|_{\infty} \qquad \forall f \in H$$

for a suitable constant $M_p \geq 0$.

PROOF. Without loss of generality, we consider the case $\gamma = \frac{n+1}{2}$. We have to show that

(4.9)
$$\left|\partial_{\tau}^{l}(t\partial_{t})^{k}\partial_{\xi}^{\alpha}\partial_{x}^{\beta}h_{f}(t,x,i\tau,\xi)\right|\langle\tau,\xi\rangle^{1+l+|\alpha|} \|f\|_{\infty}^{-1}$$

is uniformly bounded for $t > 0, x \in \mathbb{R}^n, \tau \in \mathbb{R}$ and $0 \neq f \in H$. Observing that $t\partial_t t^\mu = \mu t^\mu$ and

$$t\partial_t \left(\widetilde{a}_1(t, x, i\tau, \xi, t\eta) \right) = (t\partial_t \widetilde{a}_1)(t, x, i\tau, \xi, t\eta) + (\eta\partial_\eta \widetilde{a}_1)(t, x, i\tau, \xi, t\eta)$$

we see that the totally characteristic derivative gives rise to terms of the same type as \tilde{a}_1 . Since the derivatives with respect to x, τ and ξ can be taken under the integral sign, we may assume from the beginning that $\tilde{a}_1 \in S^{-\mu-1-j}(\overline{\mathbb{R}}_+ \times \mathbb{R}^n \times \mathbb{R}^{1+n} \times \Sigma)$ and show that

$$|h_f(t, x, i\tau, \xi)| \le c \langle \tau, \xi \rangle^{-1-j} \, \|f\|_{\infty}$$

uniformly in $t > 0, x \in \mathbb{R}^n$ and $f \in H$. By hypothesis, we have

$$|h_f(t, x, i\tau, \xi)| \le c \, \|f\|_{\infty} \, t^{\mu} \int_0^{+\infty} \langle \tau, \xi, t\varrho \rangle^{-\mu - 1 - j} \, \varrho^{\mu - 1} \, d\varrho,$$

The transformation $\rho = t^{-1} \langle \tau, \xi \rangle \sigma$ together with the identity $\langle \tau, \xi, \langle \tau, \xi \rangle \sigma \rangle = \langle \tau, \xi \rangle \langle \sigma \rangle$ yields

$$\begin{aligned} |h_f(t,x,i\tau,\xi)| &\leq c \, \|f\|_{\infty} \int_0^{+\infty} \langle \tau,\xi \rangle^{-\mu-1-j} \, \langle \sigma \rangle^{-\mu-1-j} \, \langle \tau,\xi \rangle^{\mu-1} \, \sigma^{\mu-1} \, \langle \tau,\xi \rangle \, d\sigma \\ &\leq c \, \|f\|_{\infty} \, \langle \tau,\xi \rangle^{-1-j} \int_0^{+\infty} \langle \sigma \rangle^{-2} \, d\sigma. \end{aligned}$$

This finishes the proof.

PROPOSITION 4.4. Let $\widetilde{a}_0 \in S^{-\mu}(\overline{\mathbb{R}}_+ \times \mathbb{R}^n \times \mathbb{R}^{1+n} \times \Sigma)$ be as described in Definition 3.5 with compact (t, x)-support. Define

$$h_f(t, x, \frac{n+1}{2} - \gamma + i\tau, \xi) = t^{\mu} \int_{\partial \Sigma} f(\eta^{\mu}) \,\widetilde{a}_0(t, x, \tau, \xi, t\eta) \, \eta^{\mu-1} \, d\eta$$

with $f \in H$. Then $h_f \in MS^0(\mathbb{R}_+ \times \mathbb{R}^n \times \Gamma_{\frac{n+1}{2}-\gamma} \times \mathbb{R}^n)$, and the symbol estimates of $||f||_{\infty}^{-1} h_f$ are uniform in $0 \neq f \in H$. Therefore, for a suitable constant $M_p \geq 0$,

$$\|\mathrm{op}_{M}^{\gamma-\frac{n}{2}}\mathrm{op}_{+}(h_{f})\|_{\mathcal{L}(\mathcal{H}_{p}^{0,\gamma}(\mathbb{R}_{+}\times\overline{\mathbb{R}}_{+}^{n}))} \leq M_{p} \|f\|_{\infty} \qquad \forall f \in H.$$

PROOF. It is sufficient to show the symbol estimate for h_f ; the estimates for derivatives of h_f are obtained similarly, arguing as in the proof of Proposition 4.3. The change of variables $\eta \mapsto t^{-1}\eta$ yields

(4.10)
$$h_f(t, x, \frac{n+1}{2} - \gamma + i\tau, \xi) = \int_{\partial \Sigma} f(t^{-\mu} \eta^{\mu}) \, \tilde{a}_0(t, x, \tau, \xi, \eta) \, \eta^{\mu - 1} \, d\eta.$$

Let us now denote by $C_{(\tau,\xi)}$ the natural parametrization of the boundary of $\{\eta \in \mathbb{C} \setminus \Sigma \mid |\eta| < c_0 \langle \tau, \xi \rangle\}$, where c_0 is associated with \tilde{a}_0 as in Definition 3.5. As the integrand in (4.10), for fixed (t, x, τ, ξ) , is holomorphic in η outside $\Sigma \cup \{|\eta| < c_0 \langle \tau, \xi \rangle\}$ and decays there as $|\eta|^{-1-\varepsilon}$ for some $\varepsilon > 0$ (recall the decay property of functions in H), we may replace the integration over $\partial \Sigma$ in (4.10) by integration over $C_{(\tau,\xi)}$ and obtain

$$|h_f(t, x, \frac{n+1}{2} - \gamma + i\tau, \xi)| \le c \, \|f\|_{\infty} \operatorname{length}(\mathcal{C}_{(\tau,\xi)}) \, \langle \tau, \xi \rangle^{-\mu} \, \langle \tau, \xi \rangle^{\mu-1} \le c \, \|f\|_{\infty},$$

since the length of $\mathcal{C}_{(\tau,\xi)}$ is less than $(2+2\pi)c_0 \langle \tau, \xi \rangle$.

PROPOSITION 4.5. Let d be a symbol kernel as in (3.14), (3.15), with compact support in (t, x'). Define, for $f \in H$, the symbol kernel

$$h_f(t, x', \frac{n+1}{2} - \gamma + i\tau, \xi'; u, v) = t^{\mu} \int_{\partial \Sigma} f(\eta^{\mu}) \, d(t, x', \tau, \xi'; u, v) \, \eta^{\mu - 1} \, d\eta.$$

Then there exists a constant $M_p \ge 0$ such that

(4.11)
$$\|\mathrm{op}_{M}^{\gamma-\frac{n}{2}}\mathrm{op}'(h_{f})\|_{\mathcal{L}(\mathcal{H}_{p}^{0,\gamma}(\mathbb{R}_{+}\times\overline{\mathbb{R}}_{+}^{n}))} \leq M_{p} \|f\|_{\infty} \quad \forall f \in H.$$

PROOF. Without loss of generality, $\gamma = \frac{n+1}{2} - \frac{1}{p}$. For this choice $\mathcal{H}_p^{0,\gamma}(\mathbb{R}_+ \times \overline{\mathbb{R}}_+^n) = L_p(\mathbb{R}_+ \times \mathbb{R}_+^n)$. We are now going to show that

(4.12)
$$|\partial_{\tau}^{k} \partial_{\xi'}^{\alpha}(t\partial_{t})^{l} \partial_{x'}^{\beta} h_{f}(t, x', \frac{1}{p} + i\tau, \xi'; u, v)| \leq c ||f||_{\infty} \frac{\langle \tau, \xi' \rangle^{-k - |\alpha|}}{u + v}$$

for any $k, l \in \mathbb{N}_0$ and $\alpha, \beta \in \mathbb{N}_0^{n-1}$. It is enough to consider the case k = l = 0 and $\alpha = \beta = 0$, since the terms for higher order derivatives are of the same kind, cf. the proof of Proposition 4.3. Inserting the explicit form of d, cf. (3.14), we obtain

$$h_f(t, x', \frac{1}{p} + i\tau, \xi'; u, v) = t \int_{\partial \Sigma} f(\eta^{\mu}) \, \widetilde{d}(x', \xi', t\eta; [\xi', \tau, t\eta] u, [\xi', \tau, t\eta] v) \, (t\eta)^{\mu - 1} \, d\eta.$$

Since $[\cdot] \sim \langle \cdot \rangle$ and d is rapidly decreasing in (u, v), hence in u + v, we can estimate

$$\begin{aligned} |h_f(t, x', \frac{1}{p} + i\tau, \xi'; u, v)| &\leq c \|f\|_{\infty} \int_0^\infty (\varrho t)^{\mu - 1} \left\langle (u + v) \left\langle \xi', \tau, \varrho t \right\rangle \right\rangle^{-2} \left\langle \xi', \tau, \varrho t \right\rangle^{1 - \mu} t \, d\varrho \\ &= c \|f\|_{\infty} \int_0^\infty \sigma^{\mu - 1} \left\langle (u + v) \left\langle \tau, \xi' \right\rangle \left\langle \sigma \right\rangle \right\rangle^{-2} \left\langle \sigma \right\rangle^{1 - \mu} \left\langle \tau, \xi' \right\rangle \, d\sigma; \end{aligned}$$

for the last identity we made use of the change of variables $\rho = t^{-1} \langle \tau, \xi' \rangle \sigma$. The change of variables $r = (u + v) \langle \tau, \xi' \rangle (1 + \sigma)$ then yields

$$\begin{aligned} |h_f(t, x', \frac{1}{p} + i\tau, \xi'; u, v)| &\leq c ||f||_{\infty} \int_0^\infty \sigma^{\mu - 1} \left\langle (u + v) \left\langle \tau, \xi' \right\rangle (1 + \sigma) \right\rangle^{-2} \left\langle \sigma \right\rangle^{1 - \mu} \left\langle \tau, \xi' \right\rangle \, d\sigma \\ &\leq c ||f||_{\infty} \frac{1}{u + v} \int_0^\infty \left\langle r \right\rangle^{-2} \, dr, \end{aligned}$$

i.e. (4.12). Using the continuity of the Hilbert transform,

$$u \mapsto Hu = \int_0^\infty \frac{u(s)}{\cdot + s} \, ds : L_p(\mathbb{R}_+) \longrightarrow L_p(\mathbb{R}_+),$$

and the $L_p(\mathbb{R}^n)$ -continuity of standard zero order pseudodifferential operators, assertion (4.11) is obtained as follows: Let us write for short y = (t, x') and $\operatorname{op}_y = \operatorname{op}_{M,t}^{\gamma - \frac{n}{2}} \operatorname{op}_{x'}$. Then, for $\varphi = \varphi(u, y) \in \mathcal{C}^{\infty}_{\operatorname{comp}}(\mathbb{R}_+ \times \mathbb{R}^n_+)$,

$$\left(\mathrm{op}_{M}^{\gamma-\frac{n}{2}}\mathrm{op}'(h_{f})\varphi\right)(u,y) = \int_{0}^{\infty} \left(\mathrm{op}_{y}(h_{f}(u,v))\varphi\right)(v,y)\,dv,$$

and therefore

$$\begin{aligned} \|\mathrm{op}'(h_f)\varphi\|_{L_p(\mathbb{R}_+\times\mathbb{R}^n_+)}^p &\leq \int_0^\infty \int \Big(\int_0^\infty |\big(\mathrm{op}_y(h_f(u,v))\varphi\big)(u,y)|dv\Big)^p \, dy du \\ &\leq \int_0^\infty \Big(\int_0^\infty \Big(\int |\big(\mathrm{op}_y(h_f(u,v))\varphi\big)(u,y)|^p \, dy\Big)^{1/p} \, dv\Big)^p \, du \end{aligned}$$

The second estimate is due to Minkowski's inequality for integrals. Thus

$$\begin{aligned} \|\mathrm{op}'(h_f)\varphi\|_{L_p(\mathbb{R}_+\times\mathbb{R}^n_+)}^p &\leq c\|f\|_{\infty}^p \int_0^\infty \left(\int_0^\infty \frac{1}{u+v}\|\varphi(v,y)\|_{L_p(\mathbb{R}_+\times\mathbb{R}^n_{+,y})}\,dv\right)^p du \\ &\leq c\|f\|_{\infty}^p \int_0^\infty \|\varphi(u,y)\|_{L_p(\mathbb{R}_+\times\mathbb{R}^n_{+,y})}^p\,du = c\|f\|_{\infty}^p \|\varphi\|_{L_p(\mathbb{R}_+\times\mathbb{R}^n_+)}^p.\end{aligned}$$

This finishes the proof.

5. Parameter-ellipticity of the minimal extension

In Theorem 4.1 we showed the existence of a bounded H_{∞} -calculus for a closed extension \underline{A}_T , assuming that the resolvent exists in the sector Λ and has a suitable structure. An obvious problem is now to find conditions on A and T which are more easily checked and which ensure all the required assumptions on \underline{A}_T . In this section we shall give such conditions for the case $\underline{A}_T = A_{T,\min}$. In fact, these conditions are obtained by combining the concept of parameter-ellipticity in Schulze's cone calculus and the observations from Section 3.2 of [4].

As described in Section 3.3, \mathbb{D} -ellipticity of $\binom{A}{T}$ is characterized by the invertibility of the principal symbol $\sigma^{\mu}_{\psi}(A)$, the rescaled symbol $\tilde{\sigma}^{\mu}_{\psi}(A)$, the boundary symbol $\sigma^{\mu}_{\partial}\binom{A}{T}$, and the rescaled boundary symbol $\tilde{\sigma}^{\mu}_{\partial}\binom{A}{T}$. We shall now pose stronger conditions, which we call *parameter-ellipticity* with respect to the sector Λ . The first condition is an analog of condition (2.7):

(E1) Both $\sigma^{\mu}_{\psi}(A)$ and $\tilde{\sigma}^{\mu}_{\psi}(A)$ pointwise do not have spectrum in Λ (for non-zero covariables). For $\begin{pmatrix} A \\ T \end{pmatrix}$ as in (3.4) we consider the boundary symbol and rescaled boundary symbol

$$\sigma^{\mu}_{\partial} \begin{pmatrix} A \\ T \end{pmatrix} : \mathcal{S}(\overline{\mathbb{R}}_{+}, E') \longrightarrow \begin{pmatrix} \mathcal{S}(\overline{\mathbb{R}}_{+}, E') \\ \oplus \\ F' \end{pmatrix}, \qquad \widetilde{\sigma}^{\mu}_{\partial} \begin{pmatrix} A \\ T \end{pmatrix} : \mathcal{S}(\overline{\mathbb{R}}_{+}, E'') \longrightarrow \begin{pmatrix} \mathcal{S}(\overline{\mathbb{R}}_{+}, E'') \\ \oplus \\ F'' \end{pmatrix},$$

where E', E'' and F', F'' are the corresponding pull-backs of E and F to T^* int \mathbb{B} and $T^*\partial \mathbb{B}$, respectively. We require that

(E2) Both
$$\binom{\lambda}{0} - \sigma^{\mu}_{\partial} \binom{A}{T}$$
 and $\binom{\lambda}{0} - \tilde{\sigma}^{\mu}_{\partial} \binom{A}{T}$ are pointwise invertible on $(T^* \text{int } \mathbb{B} \times \Lambda) \setminus \{0\}$
and $(T^* \partial \mathbb{B} \times \Lambda) \setminus \{0\}$, respectively.

Mainly to ensure the identity (3.6), we pose a condition on the so-called *conormal symbol* $\sigma_M^{\mu}\begin{pmatrix} A\\T \end{pmatrix}(z) =$

 $\begin{pmatrix} \sigma_M^{\mu}(A)(z) \\ \sigma_M^{\mu}(T)(z) \end{pmatrix}$. This is a holomorphic family in $z \in \mathbb{C}$ of boundary value problems on X obtained in the following way: Using the representation of A as in (3.1),

$$\sigma_M^{\mu}(A)(z) := \sum_{j=0}^{\mu} a_j(0) z^j$$

is a holomorphic family of differential operators on X. Similarly, using the notation from (3.3), one defines $\sigma_M^{\mu}(T) = (\sigma_M^0(T_0), \dots, \sigma_M^{\mu-1}(T_{\mu-1}))$ by

$$\sigma_M^j(T_j)(z) = \gamma_0 \circ \sigma_M^j(B_j)(z).$$

It can be shown that $\sigma_M^{\mu}\begin{pmatrix}A\\T\end{pmatrix}$ is meromorphically invertible in case $\begin{pmatrix}A\\T\end{pmatrix}$ is \mathbb{D} -elliptic (in the sense of Section 3.3). Then the condition is that

(E3)
$$\sigma_M^{\mu}\begin{pmatrix}A\\T\end{pmatrix}(z): \mathcal{C}^{\infty}(X, E) \longrightarrow \begin{array}{c} \mathcal{C}^{\infty}(X, E)\\ \oplus\\ \mathcal{C}^{\infty}(\partial X, F) \end{array}$$
 is invertible for each z with $\operatorname{Re} z = \frac{n+1}{2} - \gamma - \mu$.

REMARK 5.1. The invertibility of the conormal symbol from (E3) is equivalent to that of

$$\sigma_{M}^{\mu}\begin{pmatrix}A\\T\end{pmatrix}(z): H_{p}^{s}(X, E) \longrightarrow \bigoplus_{\substack{j=0\\j=0}}^{H_{p}^{s-\mu}(X, E)} \mathcal{B}_{pp}^{s-j-1/p}(\partial X, F_{j})$$

for $s \ge 0$ and 1 . In fact, according to a result of Grubb [12, Theorem 1.12], invertibilityfor one choice of s and p implies that the inverse also is an element in Boutet de Monvel's calculusand thus infers the invertibility for every other choice. Equivalence with the invertibility on spacesof smooth functions then follows from the fact that the kernel and the cokernel of an elliptic operatorconsist of smooth functions.

For A as in (3.1), we define the so-called model cone operator \widehat{A} on $X^{\wedge} = \mathbb{R}_+ \times X$ as

$$\widehat{A} = t^{-\mu} \sum_{j=0}^{\mu} a_j(0) (-t\partial_t)^j.$$

Similarly, we define $\widehat{T} = (\widehat{T}_0, \ldots, \widehat{T}_{\mu-1})$ by $\widehat{T}_j = \gamma_0 \circ \widehat{B}_j$, cf. (3.3), where now γ_0 denotes the restriction to $\partial X^{\wedge} = \mathbb{R}_+ \times \partial X$. For the analysis of \widehat{A} , one uses a special scale of Sobolev spaces $\mathcal{K}_p^{s,\gamma}(X^{\wedge})$ on X^{\wedge} with $s \in \mathbb{R}$ and 1 , namely

$$\mathcal{K}_p^{s,\gamma}(X^{\wedge}) = \{ u \in H_{p,\text{loc}}^s(X^{\wedge}) \mid \omega u \in \mathcal{H}_p^{s,\gamma}(\mathbb{D}) \text{ and } (1-\omega)u \in H_{p,\text{cone}}^s(X^{\wedge}) \};$$

here ω is a cut-off function located near t = 0, and the subscript 'cone' indicates that we do not consider X^{\wedge} with its product structure, but as an *SG*-manifold, cf. Section 4.2 in [**21**] for more details. If E^{\wedge} is the pull-back of $E|_X$ to X^{\wedge} under the canonical projection $X^{\wedge} \to X$, this definition also extends to sections, i.e. we may define $\mathcal{K}_p^{s,\gamma}(X^{\wedge}, E^{\wedge})$.

Analogously to Section 3.4, we then consider \widehat{A} as an unbounded operator

(5.1)
$$\widehat{A}: \mathcal{S}^{\infty}(X^{\wedge}, E^{\wedge})_{\widehat{T}} \subset \mathcal{K}^{0,\gamma}_p(X^{\wedge}, E^{\wedge}) \longrightarrow \mathcal{K}^{0,\gamma}_p(X^{\wedge}, E^{\wedge}),$$

where $S^{\infty}(X^{\wedge}, E^{\wedge})$ are the smooth sections of E^{\wedge} that vanish rapidly for $t \to \infty$ and vanish to infinite order in t = 0. The main ingredients for the analysis of the closed extensions of \widehat{A} are:

- i) \hat{T} has a right-inverse that belongs to the cone calculus for boundary value problems for the infinite cone (cf. for example [15]).
- ii) For a fixed $0 \neq \lambda_0 \in \Lambda$, $\begin{pmatrix} \lambda_0 A \\ \widehat{T} \end{pmatrix}$ is an elliptic element in the cone calculus and one can construct a parametrix $\begin{pmatrix} \widehat{R} & \widehat{K} \end{pmatrix}$ inverting it modulo finite rank operators and such that $\widehat{T}\widehat{R} = 0$.

For the analysis of closed extensions of A on \mathbb{D} both the corresponding right-inverse as well as the parametrix were constructed in Lemma 3.4 and Propositions 3.3, 3.7 of [4] relying on results of [12] for boundary value problems on smooth manifolds. Both constructions extend to X^{\wedge} .

THEOREM 5.2. Under conditions (E1) to (E3) the following statements hold:

- a) $\mathcal{S}^{\infty}(X^{\wedge}, E^{\wedge})_{\widehat{T}}$ is a dense subspace of $\mathcal{K}_p^{s,\gamma}(X^{\wedge}, E^{\wedge})_{\widehat{T}}$ for any $1 and <math>s, \gamma \in \mathbb{R}$ with $s > \mu - 1 + \frac{1}{n}$.
- b) The domain of the closure of \widehat{A} from (5.1), which we denote by $\widehat{A}_{\widehat{T},\min}$, coincides with $\mathcal{K}_{p}^{\mu,\gamma+\mu}(X^{\wedge}, E^{\wedge})_{\widehat{T}}$.

PROOF. a) Using i) above, we obtain a projection $P : \mathcal{K}_p^{s,\gamma}(X^{\wedge}, E^{\wedge}) \to \mathcal{K}_p^{s,\gamma}(X^{\wedge}, E^{\wedge})_{\widehat{T}}$ within the cone calculus on the infinite cone X^{\wedge} (cf. Section 2.2.3 in [15]). Then one argues as in the proof of Corollary 3.10 in [4].

b) The continuity of \widehat{A} together with a) implies that $\mathcal{K}_{p}^{\mu,\gamma+\mu}(X^{\wedge}, E^{\wedge})_{\widehat{T}} \subset \mathcal{D}(\widehat{A}_{\widehat{T},\min})$. The reverse inclusion follows as in the proof of Proposition 4.2 in [4] with the special parametrix from ii). \Box

Our next – and final – requirement is that

(E4) $\widehat{A}_{\widehat{T},\min}$ does not have spectrum in $\Lambda \setminus \{0\}$.

As \widehat{A} and \widehat{T} are invariant under dilations, the spectrum of $\widehat{A}_{\widehat{T},\min}$ is automatically a conical subset of \mathbb{C} . The next proposition shows that the spectrum of $\widehat{A}_{\widehat{T},\min}$ does not depend on the choice of 1 :

PROPOSITION 5.3. Assume that conditions (E1) to (E3) hold and fix a $\lambda_0 \neq 0$. Let us denote, for the moment, by A_p the minimal extension of $\lambda_0 - \hat{A}$ subject to \hat{T} in $\mathcal{K}_p^{0,\gamma}(X^{\wedge}, E^{\wedge})$. Suppose that for some $1 < p_0 < \infty$ the operator A_{p_0} is invertible. Then A_p is invertible for all 1 .

PROOF. By Theorem 5.2, the domain of A_p is $\mathcal{K}_p^{\mu,\gamma+\mu}(X^{\wedge}, E^{\wedge})_{\widehat{T}}$. Then the invertibility of A_p is equivalent to the invertibility of

$$\widehat{\mathcal{A}} := \begin{pmatrix} \lambda_0 - \widehat{A} \\ \widehat{T} \end{pmatrix} : \mathcal{K}_p^{\mu,\gamma+\mu}(X^\wedge, E^\wedge) \longrightarrow \begin{array}{c} \mathcal{K}_p^{0,\gamma}(X^\wedge, E^\wedge) \\ \oplus \\ \bigoplus_{j=0}^{\mu-1} \mathcal{B}_{pp}^{\mu-j-1/p,\gamma+\mu-j-1/2}(\partial X^\wedge, F_j^\wedge) \end{array}$$

see e.g. Corollary 7.2 in [4]. Since $\widehat{\mathcal{A}}$ is an elliptic element in the cone calculus on X^{\wedge} , we find a parametrix \mathcal{B} to $\widehat{\mathcal{A}}$ such that

(5.2)
$$\widehat{\mathcal{A}}\mathcal{B} = I + \mathcal{R}_1 \text{ and } \mathcal{B}\widehat{\mathcal{A}} = I + \mathcal{R}_2.$$

where \mathcal{R}_1 and \mathcal{R}_2 are operators of order $-\infty$ and types 0 and μ , respectively. They have the following mapping properties:

$$\mathcal{K}_{p}^{s,\gamma}(X^{\wedge}, E^{\wedge}) \qquad \qquad \mathcal{S}^{\gamma}(X^{\wedge}, E^{\wedge}) \\ \oplus \\ \bigoplus_{j=0}^{\mu-1} \mathcal{B}_{pp}^{s+\mu-j-1/p,\gamma+\mu-j-1/2}(\partial X^{\wedge}, F_{j}^{\wedge}) \qquad \qquad \bigoplus_{j=0}^{\mu-1} \mathcal{S}^{\gamma+\mu-j-1/2}(\partial X^{\wedge}, F_{j}^{\wedge})$$

for any $1 and <math>s > -1 + \frac{1}{p}$, and

$$\mathcal{R}_2: \mathcal{K}_p^{s,\gamma+\mu}(X^\wedge, E^\wedge) \longrightarrow \mathcal{S}^{\gamma+\mu}(X^\wedge, E^\wedge), \qquad 1 \mu - 1 + \frac{1}{p}.$$

In fact, in Section 2.1.6 of [15] these mapping properties are shown for the case p = 2; for the extension to arbitrary p we use [14]. In case $p = p_0$, multiplying (5.2) by $\widehat{\mathcal{A}}^{-1}$ yields that

(5.3)
$$\widehat{\mathcal{A}}^{-1} = \mathcal{B} + \mathcal{B}\mathcal{R}_1 - \mathcal{R}_2\widehat{\mathcal{A}}^{-1}\mathcal{R}_1.$$

For each 1 the right hand side extends to a bounded map

$$\begin{array}{c} \mathcal{K}^{0,\gamma}_{p}(X^{\wedge}, E^{\wedge}) \\ \oplus \\ \bigoplus_{j=0}^{\mu-1} \mathcal{B}^{\mu-j-1/p,\gamma+\mu-j-1/2}_{pp}(\partial X^{\wedge}, F^{\wedge}_{j}) \end{array} \longrightarrow \mathcal{K}^{\mu,\gamma+\mu}_{p}(X^{\wedge}, E^{\wedge})$$

and moreover restricts to a continuous map

$$\begin{array}{c} \mathcal{S}_{0}^{\gamma}(X^{\wedge}, E^{\wedge}) \\ \oplus \\ \bigoplus_{j=0}^{\mu-1} \mathcal{S}_{0}^{\gamma+\mu-j-1/2}(\partial X^{\wedge}, F_{j}^{\wedge}) \end{array} \longrightarrow \mathcal{S}_{0}^{\gamma+\mu}(X^{\wedge}, E^{\wedge}).$$

By density, the right hand side of (5.3) therefore furnishes an inverse to $\widehat{\mathcal{A}}$ for arbitrary p.

THEOREM 5.4. Let $\begin{pmatrix} A \\ T \end{pmatrix}$ satisfy the conditions (E1) to (E4). Then

$$A_{T,\min}: \mathcal{H}_p^{\mu,\gamma+\mu}(\mathbb{D},E)_T \subset \mathcal{H}_p^{0,\gamma}(\mathbb{D},E) \longrightarrow \mathcal{H}_p^{0,\gamma}(\mathbb{D},E)$$

fulfills the assumptions of Theorem 4.1. Hence there exists a $c \geq 0$ such that $c + A_{T,\min}$ has a bounded H_{∞} -calculus with respect to $\mathbb{C} \setminus \Lambda$.

PROOF. Let us choose parameter-dependent order reductions in the cone algebra on \mathbb{B} ,

$$R_j(\eta) \in C^{\mu-j}(\mathbb{B}, \Sigma; \gamma + \mu - j, \gamma, \theta; F_j, F_j), \qquad j = 0, \dots, \mu - 1,$$

where the F_j are the bundles from (3.3) and $\theta \in \mathbb{N}$ is arbitrary. Let $R(\eta) = \text{diag}(R_0(\eta), \ldots, R_{\mu-1}(\eta))$. Now the conditions (E1) to (E4) are chosen in such a way that

$$\mathcal{A}(\eta) := \begin{pmatrix} 1 & 0 \\ 0 & R(\eta) \end{pmatrix} \begin{pmatrix} \eta^{\mu} - A \\ T \end{pmatrix} = \begin{pmatrix} \eta^{\mu} - A \\ R(\eta)T \end{pmatrix} \in C^{\mu,\mu}(\mathbb{D}, \Sigma; \gamma + \mu, \gamma, \theta; E; E, F)$$

with $F := F_0 \oplus \ldots \oplus F_{\mu-1}$ is a parameter-elliptic element the cone calculus for boundary value problems. It follows that there exists a parametrix $\mathcal{B}(\eta) = \begin{pmatrix} B(\eta) & K(\eta) \end{pmatrix}$ and that, for p = 2,

(5.4)
$$\mathcal{A}(\eta): \mathcal{H}_{p}^{\mu,\gamma+\mu}(\mathbb{D},E) \longrightarrow \begin{array}{c} \mathcal{H}_{p}^{0,\gamma}(\mathbb{D},E) \\ \oplus \\ \mathcal{B}_{pp}^{-\frac{1}{2},\gamma-\frac{1}{2}}(\mathbb{B},F) \end{array}$$

is bijective for sufficiently large $|\eta|$, and the inverse coincides with the parametrix. Also it follows that $B(\eta) \in C^{-\mu,0}(\mathbb{D}, \Sigma; \gamma; E)$ in the sense of Section 3.5.3. As we shall show below even $B(\eta) \in C^{-\mu,0}_{\text{hol}}(\mathbb{D}, \Sigma; \gamma; E)$. Next, we note that invertibility of (5.4) is equivalent to the invertibility of

(5.5)
$$\eta^{\mu} - A : \mathcal{H}_{p}^{\mu,\gamma+\mu}(\mathbb{D}, E)_{T} \longrightarrow \mathcal{H}_{p}^{0,\gamma}(\mathbb{D}, E);$$

the inverse of (5.5) is just $B(\eta)$. By Theorem 3.4 we can conclude that the invertibility of (5.5) then also is true for arbitrary $1 , and the inverse again coincides with <math>B(\eta)$.

Now let $B(\eta)$ be as in (3.11). Decomposing \tilde{h} as in (3.12) and (3.13) yields local symbols $\tilde{a} \in S_{\rm cl}^{-\mu}(\overline{\mathbb{R}}_+ \times \mathbb{R}_x^n \times \mathbb{R}^{1+n} \times \Sigma)$. By parametrix construction, the leading term is given by inversion of the parameter-dependent principal symbol, i.e. $\tilde{a} = \tilde{a}_0 + \tilde{a}_1$ with $\tilde{a}_1(t, x, \tau, \xi, \eta) \in S_{\rm cl}^{-\mu-1}(\overline{\mathbb{R}}_+ \times \mathbb{R}_x^n \times \mathbb{R}^{1+n}_{(\tau,\xi)} \times \Sigma)$ and

$$\widetilde{a}_0(t, x, \tau, \xi, \eta) = \chi(|(\tau, \xi, \eta)|)(\eta^\mu - \widetilde{\sigma}^\mu_\psi(A)(t, x, \tau, \xi))^{-1},$$

where χ is a 0-excision function. By ellipticity assumption (E1), $(\eta^{\mu} - \tilde{\sigma}^{\mu}_{\psi}(A)(t, x, \tau, \xi))^{-1}$ is defined for $0 \neq (\tau, \xi, \eta) \in \mathbb{R}^{1+n} \times \Sigma$. By homogeneity, it is clear that

$$\operatorname{spec}(\widetilde{\sigma}^{\mu}_{\psi}(A)(t,x,\tau,\xi)) \subset \{\lambda \in \mathbb{C} \mid \frac{1}{c_0} | (\tau,\xi) |^{\mu} < |\lambda| < c_0 | (\tau,\xi) |^{\mu} \}$$

for a suitable constant $c_0 > 1$. Thus $(\eta^{\mu} - \widetilde{\sigma}^{\mu}_{\psi}(A)(t, x, \tau, \xi))^{-1}$ is defined on

$$\Omega := \{ (\tau, \xi, \eta) \mid (\tau, \xi, \eta) \in (\mathbb{R}^{1+n} \times \Sigma) \setminus \{0\} \text{ or } |\eta| \ge c_0 |\tau, \xi| \}$$

As $(\eta^{\mu} - \tilde{\sigma}^{\mu}_{\psi}(A)(t, x, \tau, \xi))^{-1}$ is positively homogeneous of degree $-\mu$ in $(\eta, \tau, \xi) \in \Omega$, and $\Omega \cap \partial U_1(0)$ is compact $(U_1(0)$ denoting the unit ball), the estimates (3.16) with $\nu = -\mu$ follow.

6. Example: The Dirichlet and Neumann problems for the Laplacian

We equip \mathbb{D} with a straight conical metric, i.e. a metric that coincides with $dt^2 + t^2g$ on $]0, 1[\times X$ for a fixed metric g on X. The associated scalar Laplacian $-\Delta$ is a Fuchs-type operator. Near t = 0 it can be written in the form

(6.1)
$$-\Delta = -t^{-2} \left\{ (t\partial_t)^2 + (n-1)t\partial_t + \Delta_X \right\}, \qquad n = \dim X,$$

where Δ_X denotes the Laplacian on X with respect to g. We let

$$\mathcal{A}_D := \begin{pmatrix} -\Delta \\ \gamma_0 \end{pmatrix}$$
 and $\mathcal{A}_N := \begin{pmatrix} -\Delta \\ t^{-1}\gamma_1 \end{pmatrix}$

be the Dirichlet and Neumann boundary value problems for $-\Delta$, respectively. We denote by $-\Delta_D$ and $-\Delta_N$ the unbounded operators in $\mathcal{H}_p^{0,\gamma}(\mathbb{D})$, acting as $-\Delta$ on the domains $\mathcal{C}^{\infty,\infty}(\mathbb{D})_{\gamma_0}$ and $\mathcal{C}^{\infty,\infty}(\mathbb{D})_{\gamma_1}$, respectively. In the sequel we will write $-\Delta_{D/N}$ to address both operators. It is easy to see that both satisfy the ellipticity conditions (E1) and (E2).

Given a function space \mathcal{F} we will use the notation \mathcal{F}_D and \mathcal{F}_N in place of \mathcal{F}_{γ_0} and \mathcal{F}_{γ_1} to denote the closed subspace of \mathcal{F} where γ_0 and γ_1 , respectively, vanish.

6.1. Closed extensions of the Dirichlet and Neumann Laplacians on \mathbb{D} . According to the definition in Section 5, the principal conormal symbols of \mathcal{A}_D and \mathcal{A}_N are

$$\sigma_M^2(\mathcal{A}_D)(z) = \begin{pmatrix} -z^2 + (n-1)z - \Delta_X \\ \gamma_0 \end{pmatrix} \text{ and } \sigma_M^2(\mathcal{A}_N)(z) = \begin{pmatrix} -z^2 + (n-1)z - \Delta_X \\ \gamma_1 \end{pmatrix}.$$

They are invertible, unless $-z^2 + (n-1)z = \lambda_j$ for one of the eigenvalues $\lambda_0 > \lambda_1 > \ldots$ of the boundary problems Δ_{X,γ_0} and Δ_{X,γ_1} , respectively (recall that $\lambda_0 < 0$ for the Dirichlet problem, while $\lambda_0 = 0$ for the Neumann problem). This is the case for $z = q_i^+$ or $z = q_i^-$ with

$$q_j^{\pm} = \frac{n-1}{2} \pm \sqrt{\left(\frac{n-1}{2}\right)^2 - \lambda_j}, \qquad j \in \mathbb{N}_0$$

We shall now study the minimal extension of $\Delta_{D/N}$ in $\mathcal{H}^{0,\gamma}_p(\mathbb{D})$. We shall require that

(6.2)
$$1 - \sqrt{\left(\frac{n-1}{2}\right)^2 - \lambda_0} < \gamma < -1 + \sqrt{\left(\frac{n-1}{2}\right)^2 - \lambda_0}$$

Of course, this only makes sense, if

(6.3)
$$\left(\frac{n-1}{2}\right)^2 - \lambda_0 > 1.$$

THEOREM 6.1. Assume (6.2) and (6.3). For $1 , the minimal and maximal extensions coincide both for the Dirichlet and the Neumann Laplacians on <math>\mathcal{H}_p^{0,\gamma}(\mathbb{D})$, and their domain is $\mathcal{H}_p^{2,\gamma+2}(\mathbb{D})_{D/N}$. In case $\gamma = 0$ and p = 2, the (minimal) extension is self-adjoint.

PROOF. This is an immediate consequence of Proposition 6.1 of [4], since condition (6.2) implies that the conormal symbols of both \mathcal{A}_D and \mathcal{A}_N are invertible for all z with $\frac{n+1}{2} - \gamma - 2 \leq \operatorname{Re} z \leq \frac{n+1}{2} - \gamma$. The self-adjointness is a consequence of Theorem 4.5 in [4], since the adjoint problems of \mathcal{A}_D and \mathcal{A}_N are $\begin{pmatrix} -\Delta \\ -i\gamma_0 \end{pmatrix}$ and $\begin{pmatrix} -\Delta \\ -it^{-1}\gamma_1 \end{pmatrix}$, respectively.

- REMARK 6.2. (a) Assumption (6.2) implies that both $-\Delta_D$ and $-\Delta_N$, considered as unbounded operators on $\mathcal{H}^{0,\gamma}_n(\mathbb{D})$, satisfy the ellipticity condition (E3).
 - (b) In the Dirichlet case, condition (6.3) is always true for n ≥ 3; depending on λ₀ (i.e. on X and g), it might also hold for n = 1 or n = 2. In the Neumann case, condition (6.3) only holds for n > 3.

6.2. Domains of the model cone operator. Recall that $\widehat{\Delta}_{D/N,\min}$ denotes the closure in $\mathcal{K}_p^{0,\gamma}(X^{\wedge})$ of the model cone operator $\widehat{\Delta}_{D/N}$ considered with domain $\mathcal{S}^{\infty}(X^{\wedge})_{D/N}$, while

$$\mathcal{D}(-\widehat{\Delta}_{D/N,\max}) = \{ u \in \mathcal{K}_p^{2,\gamma}(X^{\wedge})_{D/N} \mid -\widehat{\Delta}u \in \mathcal{K}_p^{0,\gamma}(X^{\wedge}) \}$$

defines the closed operator $\widehat{\Delta}_{D/N,\min}$. Note that always $\mathcal{K}_p^{2,\gamma+2}(X^{\wedge})_{D/N} \subset \mathcal{D}(-\widehat{\Delta}_{D/N,\min})$.

LEMMA 6.3. Under the assumptions of Theorem 6.1, the minimal and maximal extensions coincide both for the model Dirichlet and Neumann Laplacians on $\mathcal{K}_p^{0,\gamma}(X^{\wedge})$ and

$$\mathcal{D}(-\widehat{\Delta}_{D/N,\min}) = \mathcal{D}(-\widehat{\Delta}_{D/N,\max}) = \mathcal{K}_p^{2,\gamma+2}(X^{\wedge})_{D/N}.$$

PROOF. For $u \in \mathcal{D}(-\widehat{\Delta}_{D,\max})$ we have $u \in \mathcal{K}_p^{2,\gamma}(X^{\wedge})_D$ and $-\widehat{\Delta}u \in \mathcal{K}_p^{0,\gamma}(X^{\wedge})$. Then $(1 - \omega)u \in \mathcal{K}_p^{2,\infty}(X^{\wedge})_D$, and $-\widehat{\Delta}(1 - \omega)u \in \mathcal{K}_p^{0,\infty}(X^{\wedge})$. Hence, $(1 - \omega)u \in \mathcal{D}(-\widehat{\Delta}_{D,\max})$, thus $\omega u \in \mathcal{D}(-\widehat{\Delta}_{D,\max})$. As we may consider ωu as an element of $\mathcal{H}_p^{2,\gamma}(\mathbb{D})$, and since Δ and $\widehat{\Delta}$ have the same form close to t = 0, we also have $\omega u \in \mathcal{D}(-\Delta_{D,\max})$. By Theorem 6.1, $\omega u \in \mathcal{D}(-\Delta_{D,\min}) = \mathcal{H}_p^{2,\gamma+2}(\mathbb{D})_D$. Therefore, $u = \omega u + (1 - \omega)u \in \mathcal{K}_p^{2,\gamma+2}(X^{\wedge})_D \subset \mathcal{D}(-\widehat{\Delta}_{D,\min})$. We conclude that $\mathcal{D}(-\widehat{\Delta}_{D,\min}) = \mathcal{D}(-\widehat{\Delta}_{D,\max}) = \mathcal{K}_p^{2,\gamma+2}(X^{\wedge})_D$, and the proof of the statement for $-\widehat{\Delta}_D$ is complete. The argument for $-\widehat{\Delta}_N$ is the same.

COROLLARY 6.4. For $\gamma = 0$ and p = 2 the closure $\Delta_{D/N,\min}$ is self-adjoint.

PROOF. Let $\widehat{\Delta}_F$ denote the Friedrichs extension of $-\widehat{\Delta}_{D/N}$. By construction, one has

$$\mathcal{D}(\widehat{\Delta}_{D/N,\min}) \subset \mathcal{D}(\widehat{\Delta}_F) \subset \mathcal{D}((\widehat{\Delta}_{D/N,\min})^*).$$

Let $u \in \mathcal{D}((\widehat{\Delta}_{D/N,\min})^*)$ be given. Using the special parametrix \widehat{R} from ii) in Section 5 (for $\widehat{A} = -\widehat{\Delta}, T = D/N$, and $\lambda_0 = -1$), and the fact that $R^*(\widehat{\Delta}_{D/N,\min})^* \subset (\widehat{\Delta}_{D/N}R)^*$, we deduce that $u \in \mathcal{K}_2^{2,0}(X^{\wedge})$. Then, arguing as in the proof of Theorem 4.5 in [4] to verify there the identity (4.5), we conclude that $u \in \mathcal{D}(\widehat{\Delta}_{D/N,\max})$. It follows from Lemma 6.3 that $\widehat{\Delta}_F = \widehat{\Delta}_{D/N,\min}$.

THEOREM 6.5. Assume (6.2) and (6.3). Then, for 1 , both the Dirichlet and Neumann Laplacian fulfill the ellipticity conditions (E1) to (E4) of Section 5.

PROOF. It was noted above that (E1), (E2), and (E3) hold for $-\Delta_{D/N}$. It remains to check (E4). By spectral invariance, cf. Proposition 5.3, we may assume p = 2. For $\gamma = 0$, the Laplacians $-\widehat{\Delta}_{D/N,\min}$ are self-adjoint with domain $\mathcal{K}_2^{2,2}(X^{\wedge})_{D/N}$. For $\lambda \notin \mathbb{R}_+$, we therefore conclude that

(6.4)
$$-\widehat{\Delta}_{D/N} - \lambda : \ \mathcal{K}_2^{2,2}(X^{\wedge})_{D/N} \longrightarrow \mathcal{K}_2^{0,0}(X^{\wedge})$$

is invertible. Clearly, (6.4) shows the injectivity of $-\widehat{\Delta}_{D/N} - \lambda$ on $\mathcal{K}_2^{2,\gamma+2}(X^{\wedge})_{D/N} \subset \mathcal{K}_2^{2,2}(X^{\wedge})_{D/N}$ for $\gamma > 0$.

Next suppose $\gamma < 0, u \in \mathcal{K}_2^{2,\gamma+2}(X^{\wedge})_{D/N}$ and $u \in \ker\{-\widehat{\Delta}_{D/N} - \lambda \colon \mathcal{K}_2^{2,\gamma+2}(X^{\wedge})_{D/N} \to \mathcal{K}_2^{0,\gamma}(X^{\wedge})\}$. Then u is in the maximal domain of $-\widehat{\Delta}_{D/N} - \lambda$, considered as an unbounded operator on $\mathcal{K}_2^{0,\gamma+2}(X^{\wedge})$. Since condition (E4) holds, u also is in the associated minimal domain, which is $\mathcal{K}_2^{2,4+\gamma}(X^{\wedge})_{D/N}$. We can iterate this argument j times, until $\gamma + 2j > 0$. Then, we conclude that u = 0, by the previous step. Thus $-\widehat{\Delta}_{D/N} - \lambda$ is injective on $\mathcal{K}_2^{2,\gamma+2}(X^{\wedge})_{D/N}$ for all γ satisfying the hypotheses and $\lambda \notin \mathbb{R}_+$. Finally, we note that the adjoint of

$$-\widehat{\Delta}_{D/N} - \lambda : \ \mathcal{K}_2^{2,\gamma+2}(X^{\wedge})_{D/N} \longrightarrow \mathcal{K}_2^{0,\gamma}(X^{\wedge})$$

with respect to the scalar product of $\mathcal{K}_2^{0,0}(X^\wedge)$ is

$$-\widehat{\Delta}_{D/N} - \overline{\lambda} : \ \mathcal{K}_2^{2,-\gamma+2}(X^{\wedge})_{D/N} \longrightarrow \mathcal{K}_2^{0,-\gamma}(X^{\wedge}).$$

Hence, also the adjoint is injective for $\lambda \notin \overline{\mathbb{R}}_+$, so $-\widehat{\Delta}_{D/N} - \lambda$ is bijective, as claimed.

6.3. Maximal L_p regularity of the Cauchy problem for Dirichlet and Neumann Laplacians. As a consequence of Theorem 6.5 we get the following result on the solvability of the Cauchy Problem for the Dirichlet and Neumann Laplacians.

THEOREM 6.6. Let Δ be the Laplacian as described above, 1 , and assume (6.2) and (6.3).Then the initial boundary value problems

(6.5)
$$u'(\tau) - \Delta u(\tau) = f(\tau), \quad 0 \le \tau \le T; \quad u(0) = 0, \quad \gamma_j u = 0,$$

j = 0, 1, have a unique solution

$$u \in W_r^1\left([0,T], \mathcal{H}_p^{0,\gamma}(\mathbb{D})\right) \cap L_r\left([0,T], \mathcal{H}_q^{2,\gamma+2}(\mathbb{D})_{D/N}\right)$$

for each

$$f \in L_r\left([0,T], \mathcal{H}^{0,\gamma}_p(\mathbb{D})_{D/N}\right), \qquad 1 < r < \infty$$

Furthermore, u, u', and Δu depend continuously on f.

PROOF. Solving (6.5) is equivalent to solving $v'(\tau) - (\Delta_{D/N} - c)v(\tau) = e^{c\tau}f(\tau), v(0) = 0,$ $\gamma_j v = 0$, for some c > 0.

The operators $-\Delta_{D/N}$ are closed with minimal (and maximal) domain equal to $\mathcal{H}_p^{2,\gamma+2}(\mathbb{D})_{D/N}$. According to Theorem 6.5, they satisfy conditions (E1)-(E4) for each sector Λ not containing \mathbb{R}_+ . Applying Theorem 5.4, we deduce that $-\Delta_{D/N} + c$ has a bounded \mathcal{H}_{∞} -calculus for sufficiently large c, and Theorem 6.6 immediately follows from Dore and Venni's theorem, cf. Theorem 3.2 in [7].

Nazarov [19] has studied the Dirichlet and the Neumann problem for the Laplacian on infinite cones and wedges in Euclidean space. He shows results on maximal regularity in weighted Sobolev spaces using the explicit Green's function. He obtains restrictions on the weight which are similar to (6.2). They do not coincide, however, since he works on a different scale of spaces.

References

- [1] H. Amann, M. Hieber, G. Simonett. Bounded H_{∞} -calculus for elliptic operators. *Diff. Integral Eq.* 7: 613-653, 1994.
- [2] L. Boutet de Monvel. Boundary value problems for pseudo-differential operators. Acta Math. 126: 11-51, 1971.
- [3] S. Coriasco, E. Schrohe, J. Seiler. Bounded imaginary powers of cone differential operators. Math. Z. 244: 235-269, 2003.
- [4] S. Coriasco, E. Schrohe, J. Seiler. Realizations of differential operators on manifolds with boundary. *Preprint*, 2004. math.AP/0401395
- [5] R. Denk, M. Hieber, J. Prüss. *R*-boundedness, Fourier multipliers and problems of elliptic and parabolic type. *Mem. Amer. Math. Soc.* 166 (2003), no. 788.
- [6] R. Denk, G. Dore, M. Hieber, J. Prüss, A. Venni. New thoughts on old results of R.T. Seeley. Math. Ann. 328 (2004), 545-583.
- [7] G. Dore, A. Venni. On the closedness of the sum of two closed operators. Math. Z. 196: 189-201, 1987.
- [8] X.T. Duong. H_∞ functional calculus of elliptic operators with C[∞] coefficients on L_p spaces of smooth domains. J. Austral. Math. Soc. 48 (1990), 113-123.
- [9] J. Escher, J. Seiler. Bounded H_∞-calculus for pseudodifferential operators and applications to the Dirichlet-Neumann operator for C^{1+r}-domains. Preprint, 2005.
- [10] J.B. Gil, Th. Krainer, G. Mendoza. Resolvents of elliptic cone operators. Preprint, 2004. math.AP/0410176
- J.B. Gil, Th. Krainer, G. Mendoza. Geometry and spectra of closed extensions of elliptic cone operators. Preprint, 2004. math.AP/0410178
- [12] G. Grubb. Functional Calculus of Pseudodifferential Boundary Problems, 2nd edition. Progress in Math. 65, Birkhäuser, 1996.
- [13] G. Grubb. Parameter-elliptic and parabolic pseudodifferential boundary problems in global L_p Sobolev spaces. Math. Z. 218 (1995), 43-90.
- [14] G. Grubb, N. Kokholm. A global calculus of parameter-dependent pseudodifferential boundary problems in L_p Sobolev spaces. Acta Math. 171: 165-229, 1983.
- [15] D. Kapanadze, B.-W. Schulze. Crack Theory and Edge Singularities. Math. and its Appl. 561, Kluwer, 2003.
- [16] Th. Krainer. Resolvents of elliptic boundary problems on conic manifolds. Preprint, 2005. math.AP/0503021
- [17] P.C. Kunstmann, L. Weis. Maximal L_p -regularity for parabolic equations, Fourier multiplier theorems and H^{∞} -functional calculus. Preprint, 2004.
- [18] A. McIntosh. Operators which have an H_∞-calculus. In B. Jeffries et al. (eds.), Miniconference on Operator Theory and Partial Differential Equations, Proc. Center Math. Anal. A.N.U. 14, 1986.
- [19] A. Nazarov. L_p -estimates for a solution to the Dirichlet problem and to the Neumann problem for the heat equation in a wedge with edge of arbitrary codimension. J. Math. Sciences **106** (2001), 2989-3014.
- [20] E. Schrohe, B.-W. Schulze. Boundary value problems in Boutet de Monvel's calculus for manifolds with conical singularities I. In M. Demuth et al. (eds.), *Pseudo-Differential Operators and Mathematical Physics*, Math. Topics 5: Advances in Part. Diff. Equ., Akademie Verlag, 1994.
- [21] E. Schrohe, B.-W. Schulze. Boundary value problems in Boutet de Monvel's calculus for manifolds with conical singularities II. In M. Demuth et al. (eds.), Boundary Value Problems, Schrödinger Operators, Deformation Quantization, Math. Topics 8: Advances in Part. Diff. Equ., Akademie Verlag, 1995.
- [22] E. Schrohe, J. Seiler. The resolvent of closed extensions of cone differential operators. Can. J. Math., to appear.
- [23] R. Seeley. The resolvent of an elliptic boundary problem. Amer. J. Math. 91 (1969), 889-920.
- [24] R. Seeley. Norms and domains of the complex powers A_B^z . Amer. J. Math. 93 (1971), 299-309.
- [25] S. Sohr. Beschränkter H_{∞} -Funktionalkalkül für elliptische Randwertprobleme. PhD-thesis, University of Kassel, 1999.
- [26] E.M. Stein, G. Weiss. Introduction to Fourier Analysis on Euclidean Spaces. Princeton University Press, 1971.

$H_\infty\text{-}\mathrm{CALCULUS}$ FOR CONE DIFFERENTIAL OPERATORS

Universita di Torino, Dipartimento di Matematica, Via Carlo Alberto 10, 10123 Torino, Italy

UNIVERSITÄT HANNOVER, INSTITUT FÜR MATHEMATIK, WELFENGARTEN 1, 30167 HANNOVER, GERMANY

UNIVERSITÄT HANNOVER, INSTITUT FÜR ANGEWANDTE MATHEMATIK, WELFENGARTEN 1, 30167 HANNOVER, GER-MANY

 ${\it E-mail\ address:\ \tt sandro.coriasco@unito.it,\ schrohe@math.uni-hannover.de,\ seiler@ifam.uni-hannover.de}$