

# SUPERTRACE AND SUPERQUADRATIC LIE STRUCTURE ON THE WEYL ALGEBRA, WITH APPLICATIONS TO FORMAL INVERSE WEYL TRANSFORM

GEORGES PINCZON, ROSANE USHIROBIRA

ABSTRACT. Using the Moyal  $\star$ -product and orthosymplectic supersymmetry, we construct a natural non trivial supertrace and an associated non degenerate invariant supersymmetric bilinear form for the Lie superalgebra structure of the Weyl algebra  $W$ . We decompose adjoint and twisted adjoint actions. We define a renormalized supertrace and a formal inverse Weyl transform in a deformation quantization framework and develop some examples.

## 0. INTRODUCTION

Since Killing, it is known that the existence of a non degenerate invariant symmetric bilinear form is a crucial property for a Lie algebra. Let us call such a Lie algebra a *quadratic* Lie algebra. These Lie algebras are of interest, not only when they are finite dimensional, but infinite dimensional as well [10]. A corresponding notion of a *superquadratic* Lie superalgebra exists: namely, it is a Lie superalgebra endowed with a non degenerate invariant supersymmetric bilinear form. Besides, invariant bilinear forms are often constructed from Traces or Supertraces (though it is not always the case), even in the infinite dimensional case.

The main result of the present paper is:

**Theorem 1 :** The Weyl algebra is a superquadratic Lie superalgebra with (essentially unique) bilinear form derived from a supertrace.

To explain the origin of this statement, let us consider the Weyl algebra  $W$  in  $2n$  generators realized as the polynomial algebra in  $2n$  indeterminates endowed with the Moyal  $\star$ -product denoted by  $\star$ . Briefly, in a deformation quantization framework (see [3, 17]), we quantize the natural Poisson bracket in an explicit way (Moyal  $\star$ -product). The main advantage of this quantization is the fact of being invariant with respect to the natural embedding of the Lie superalgebra  $\mathfrak{osp}(1, 2n)$  in  $W$  ([7, 13, 8]), i.e. with respect to the orthosymplectic supersymmetry. The conjunction of both arguments, explicit Moyal product and supersymmetry, is indeed a powerful machinery that allows to deduce algebraic properties of the Weyl algebra, as we shall show in this paper.

---

*Date:* February 13, 2018.

*2000 Mathematics Subject Classification.* 53D55, 17B05, 17B10, 17B20, 17B60, 17B65.

*Key words and phrases.* Deformation quantization, supersymmetry, Weyl algebra, supertrace, renormalization, formal inverse Weyl transform.

The Weyl algebra  $W$  is a Lie algebra with bracket denoted by  $[\cdot, \cdot]_{\mathcal{L}}$ , and since it is naturally  $\mathbb{Z}_2$ -graded,  $W$  is a Lie superalgebra with bracket denoted by  $[\cdot, \cdot]$ .

We define the supertrace on the Weyl algebra  $W$  as the usual evaluation at 0:

$$\text{Str}(F) := F(0), \forall F \in W.$$

and a bilinear form  $\kappa$  on  $W$  as:

$$\kappa(F, G) := \text{Str}(F \star G), \forall F, G \in W.$$

Then we prove:

**Theorem 2:**

- (1)  $\text{Str}$  is a supertrace on  $W$ , that is  $\text{Str}([F, G]) = 0$ , for all  $F, G \in W$  and one has  $\ker(\text{Str}) = [W, W]$ .
- (2) The bilinear form  $\kappa$  is supersymmetric, non degenerate and invariant under the adjoint representation of the Lie superalgebra  $W$ .

Theorem 1 is a consequence of Theorem 2. Let us mention that obviously there is no non trivial trace on the Lie algebra  $W$ .

As another consequence of Theorem 2, we deduce a decomposition of  $W$ :

$$W = \mathbb{K} \oplus [W, W],$$

recovering in a natural way a nice result of I. M. Musson [13].

Concerning the bilinear form  $\kappa$ , given the decomposition of  $W$  into homogeneous factors (for the commutative product),  $W = \bigoplus_{k \geq 0} S^k$ , we show that

$$\kappa(S^k, S^\ell) = \{0\}, k \neq \ell,$$

so that the restriction of  $\kappa$  to each  $S^k$  is non degenerate, and provides an explicit (orthogonal or symplectic, according to the parity of  $k$ ) invariant bilinear form for the standard simple action of  $\mathfrak{sp}(2n)$  on  $S^k$ .

We study in addition, the decomposition of the adjoint  $\text{ad}$  and twisted adjoint  $\text{ad}'$  actions of the Lie superalgebra  $W$  on itself:

**Theorem 3:**

- (1) Under the adjoint action,  $W$  decomposes as  $W = \mathbb{K} \oplus [W, W]$ . Besides,  $[W, W]$  is a simple  $\text{ad}(W)$ -module.
- (2)  $W$  is a simple  $\text{ad}'(W)$ -module and has a non degenerate invariant supersymmetric bilinear form.

The bilinear form associated to the  $\text{ad}'$ -action is deduced from  $\kappa$ , and extends the bilinear form used to construct the natural embedding of  $\mathfrak{osp}(1, 2n)$  in  $W$ . As a corollary:

**Corollary:**  $[W, W]$  is a simple superquadratic Lie superalgebra.

Let us quote that the simplicity of  $[W, W]$  was known some time ago, as a combination of a result by S. Montgomery [12] proving that  $[W, W]/\mathbb{K}$  is simple and a result by I. M. Musson [13] proving that  $W = \mathbb{K} \oplus [W, W]$ . Our proof using a supertrace and the Moyal product is direct and completely different of the initial proofs in [12] and [13].

Finally, we reinterpret the supertrace in a Renormalization Theory context. In a few words, let  $\mathcal{P}$  be the polynomial algebra in  $n$  indeterminates: the Weyl algebra  $W$  acts on  $\mathcal{P}$  as differential operators with polynomial coefficients. It has been shown in [14] that any linear operator on  $\mathcal{P}$  is in fact a differential operator, eventually of infinite order. A slight improvement provides a very explicit remarkable formula:

**Theorem 4:** If  $T \in \mathcal{L}(\mathcal{P})$ , then

$$T = \sum_N \frac{1}{N!} (m \circ (T \otimes \mathcal{S}) \circ \Delta(x^N)) \frac{\partial^N}{\partial x^N}$$

where  $m$  is the product,  $\Delta$  the coproduct and  $\mathcal{S}$  the antipode of  $\mathcal{P}$ .

Let  $\text{Str}$  be the ordinary supertrace defined on the ideal of finite rank operators  $\mathcal{L}_f(\mathcal{P})$  and  $\text{Str}_W$  be the supertrace previously defined on  $W$ . We show the following theorem:

**Theorem 5:** Let  $T = \sum_I \alpha_I(Q) \star P^I$ . If  $T \in \mathcal{L}_f(\mathcal{P})$ , then:

$$\text{Str}(T) = \frac{1}{2^n} \sum_I \text{Str}_W(\alpha_I(Q) \star P^I).$$

Therefore we can construct a natural extension of  $\text{Str}$  to  $\mathcal{L}_f(\mathcal{P}) \oplus W$  (in fact, to a bigger subspace that will not be discussed here). For this renormalized extension, that we note  $\text{RStr}$ , one has:

$$\text{RStr}(\text{Id}) = \frac{1}{2^n},$$

a formula that renormalizes  $\text{Str}(\text{Id}) = \infty - \infty$  obtained by wildly applying the definition of  $\text{Str}$ . Moreover, the power  $n$  in this formula is precisely the dimension of the underlying variety so our renormalization of supertrace has a geometric flavor. At last, we give some ideas of what a formal inverse Weyl transform could be, and compute examples.

We have tried to make this paper as self-contained as possible. For instance, we give a short introduction to the Moyal  $\star$ -product in Section 1, and in Section 2, an introduction to the embedding of  $\mathfrak{osp}(1, 2n)$  in  $W$ , together with the decompositions of the corresponding adjoint and twisted adjoint actions. Not only because all this material provides tools used all along this paper, but also for the convenience of the reader, who is not forced to be an expert in deformation quantization theory, orthosymplectic supersymmetry, etc.

### Remarks.

- (1) Theorem 2 results from Proposition 1.9 and Theorem 4.1. Theorem 3 corresponds to Theorem 3.7. Theorem 5.2 gives Theorem 4 and the proof of Theorem 5 is Theorem 5.4.
- (2) In Sections 1 to 4 of this paper,  $\mathbb{K}$  denotes a field of characteristic zero (not necessarily algebraically closed). In Section 5,  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ .

- (3) There are so many references on deformation quantization that we cannot quote all of them. The reader should refer to the beautiful paper [3] that is the beginning (and much more) of this theory and to [17] for a complete history with 164 references.

### 1. MOYAL $\star$ -PRODUCTS

Moyal  $\star$ -products are the first examples of Deformation Quantization of Poisson Brackets (see e.g. [3, 17]). In this Section, we recall their well-known properties, giving proofs for the convenience of the reader.

Let  $V$  be a vector space with a basis  $\{P_1, Q_1, \dots, P_n, Q_n\}$  and  $\{p_1, q_1, \dots, p_n, q_n\}$  its dual basis.

We denote by  $S$  the commutative algebra  $S := \text{Sym}(V^*)$  with the usual grading  $S = \bigoplus_{k \geq 0} S^k$  and the Poisson bracket:

$$\{F, G\} = \sum_{i=1}^n \left( \frac{\partial F}{\partial p_i} \frac{\partial G}{\partial q_i} - \frac{\partial F}{\partial q_i} \frac{\partial G}{\partial p_i} \right), \quad \forall F, G \in S.$$

Denoting by  $\{X_1, \dots, X_{2n}\}$  the given basis of  $V$  and by  $\{x_1, \dots, x_{2n}\}$  its dual basis, we introduce a duality that identifies  $S^*$  and the commutative algebra of formal power series  $\mathcal{F} := \mathbb{K}[[X_1, \dots, X_{2n}]]$  by:

$$\langle x^I \mid X^J \rangle := \delta_{I,J} I!$$

where  $I, J \in \mathbb{N}^{2n}$ ,  $x^I = x_1^{i_1} \dots x_{2n}^{i_{2n}}$ ,  $X^J = X_1^{j_1} \dots X_{2n}^{j_{2n}}$  and  $I! = i_1! \dots i_{2n}!$ .

The following properties result from a straightforward verification:

**Proposition 1.1.**

- (1) For all  $v \in V$  and  $F \in S$ ,  $\langle F \mid e^v \rangle = F(v)$ .  
(2) For all  $v \in V$  and  $F \in S$ ,

$$\langle F \mid X^I e^v \rangle = \left\langle \frac{\partial^I F}{\partial x^I} \mid e^v \right\rangle = \frac{\partial^I F}{\partial x^I}(v),$$

$$\text{where } \frac{\partial^I F}{\partial x^I} := \frac{\partial^{i_1 + \dots + i_{2n}} F}{\partial x_1^{i_1} \dots \partial x_{2n}^{i_{2n}}}.$$

Notice that (1) is Taylor's formula and (2) means that the transpose of the operator  $\frac{\partial^I}{\partial x^I}$  of  $S$  is the operator of multiplication by  $X^I$  of  $\mathcal{F}$ .

Define an operator  $\wp: S \otimes S \rightarrow S \otimes S$  by:

$$(1) \quad \wp(F \otimes G) := \sum_{i=1}^n \left( \frac{\partial F}{\partial p_i} \otimes \frac{\partial G}{\partial q_i} - \frac{\partial F}{\partial q_i} \otimes \frac{\partial G}{\partial p_i} \right), \quad \forall F, G \in S.$$

Since  $S \otimes S = \mathbb{K}[p_1, q_1, \dots, p_n, q_n, p'_1, q'_1, \dots, p'_n, q'_n]$ , we have:

$$(S \otimes S)^* = \mathbb{K}[[P_1, Q_1, \dots, P_n, Q_n, P'_1, Q'_1, \dots, P'_n, Q'_n]].$$

If  $D = {}^t \wp$ , from Proposition 1.1 it follows:

$$D(A \otimes B) = \left( \sum_{i=1}^n (P_i \otimes Q_i - Q_i \otimes P_i) \right) \cdot A \otimes B, \quad \forall A, B \in \mathcal{F},$$

so that  $D$  is the operator of multiplication by  $d(\mathcal{X}, \mathcal{X}') = \sum_{i=1}^n (P_i Q'_i - Q_i P'_i)$  of  $(S \otimes S)^*$  where  $\mathcal{X} = (P_1, Q_1, \dots, P_n, Q_n)$  and  $\mathcal{X}' = (P'_1, Q'_1, \dots, P'_n, Q'_n)$ .

Denote by  $m$  the product of  $S$ . We can now define:

**Definition 1.2.** For  $F, G \in S$ , the *Moyal  $\star$ -product* is defined as:

$$F \star_t G := \left( m \circ \sum_{k \geq 0} \frac{t^k}{2^k k!} \delta^k \right) (F \otimes G).$$

Let  $\Delta$  be the coproduct of  $\mathcal{F}$  associated to  $m$ . With the usual obvious abuse of notation, one can write:

$$\Delta(A) = A(\mathcal{X} + \mathcal{X}'), \forall A \in \mathcal{F}.$$

So if we denote by  $m_\star$  the Moyal  $\star$ -product and by  $\Delta_\star$  the associated coproduct  $\Delta_\star = {}^t m_\star$ , we get:

$$\Delta_\star(A) = \sum_{k \geq 0} \frac{t^k}{2^k k!} D^k(A(\mathcal{X} + \mathcal{X}')) = e^{\frac{t}{2} d(\mathcal{X}, \mathcal{X}')} A(\mathcal{X} + \mathcal{X}'), \forall A \in \mathcal{F}.$$

**Proposition 1.3.**

- (1)  $\Delta_\star$  is coassociative.
- (2)  $m_\star$  is associative.

*Proof.* Let  $A \in \mathcal{F}$ . One has:

$$\begin{aligned} (\text{Id} \circ \Delta_\star)(\Delta_\star(A)) &= (\text{Id} \circ \Delta_\star) \left( e^{\frac{t}{2} d(\mathcal{X}, \mathcal{X}')} A(\mathcal{X} + \mathcal{X}') \right) \\ &= e^{\frac{t}{2} d(\mathcal{X}', \mathcal{X}'')} e^{\frac{t}{2} d(\mathcal{X}, \mathcal{X}' + \mathcal{X}'')} A(\mathcal{X} + \mathcal{X}' + \mathcal{X}'') \end{aligned}$$

$$\begin{aligned} (\Delta_\star \circ \text{Id})(\Delta_\star(A)) &= e^{\frac{t}{2} d(\mathcal{X}, \mathcal{X}')} \Delta_\star(A)(\mathcal{X} + \mathcal{X}', \mathcal{X}'') \\ &= e^{\frac{t}{2} d(\mathcal{X}, \mathcal{X}')} e^{\frac{t}{2} d(\mathcal{X} + \mathcal{X}', \mathcal{X}'')} A(\mathcal{X} + \mathcal{X}' + \mathcal{X}'') \end{aligned}$$

and since  $d$  is bilinear, one obtains  $(\text{Id} \circ \Delta_\star) \circ \Delta_\star = (\Delta_\star \circ \text{Id}) \circ \Delta_\star$ , so  $\Delta_\star$  is coassociative. From  $\Delta_\star = {}^t m_\star$ , it results  $m_\star$  associative.  $\square$

We introduce the notation:

$$(II) \quad C_k(F, G) = \frac{1}{2^k k!} \left( m \circ \delta^k \right) (F \otimes G), \forall F, G \in S.$$

Then we can write:

$$F \star_t G = FG + \frac{t}{2} \{F, G\} + t^2 C_2(F, G) + \dots, \forall F, G \in S.$$

Thus the Moyal  $\star$ -product is a deformation of the commutative product of  $S$  and also a deformation quantization of the Poisson bracket.

We remark that  $C_k$  is a bidifferential operator of order  $(k, k)$ , so:

**Proposition 1.4.** For all  $F \in \mathcal{S}^f$  and  $G \in \mathcal{S}^g$ ,  $C_k(F, G) \in \mathcal{S}^{f+g-2k}$  (with  $\mathcal{S}^\ell = \{0\}$  when  $\ell < 0$ ) and

$$F \star_t G = \sum_{k=0}^{\min(f,g)} t^k C_k(F, G).$$

Consider  $\tau$  the twist  $\tau(F \otimes G) := G \otimes F$ ,  $\forall F, G \in \mathcal{S}$ . Then  $\wp \circ \tau = -\tau \circ \wp$ , so  $\wp^k \circ \tau = (-1)^k \tau \circ \wp^k$  and one deduces a useful parity property of the coefficients  $C_k$ :

$$(III) \quad C_k(G, F) = (-1)^k C_k(F, G), \quad \forall F, G \in \mathcal{S}.$$

As a consequence, the Lie bracket associated to the Moyal  $\star$ -product contains only odd terms:

$$(IV) \quad [F, G]_{\wp_t} = 2 \sum_{p \geq 0} t^{2p+1} C_{2p+1}(F, G), \quad \forall F, G \in \mathcal{S}.$$

Now, from Proposition 1.4, we obtain the following useful property:

**Proposition 1.5.** Let  $F \in \mathcal{S}$  of degree  $\leq 2$ . Then  $[F, G]_{\wp_t} = t\{F, G\}$ , for all  $G \in \mathcal{S}$ .

Therefore  $[p_i, q_i]_{\wp_t} = t$ , for  $i = 1, \dots, n$  and all other brackets between  $p_i$ 's and  $q_j$ 's vanish.

Next we shall relate the Moyal  $\star$ -product to the Weyl algebra. For this purpose, we need a Lemma:

**Lemma 1.6.** One has

$$(1) \text{ For all } \varphi \in V^*, \varphi \star_t^k \varphi = \underbrace{\varphi \star_t \dots \star_t \varphi}_{k \text{ times}} = \varphi^k.$$

$$(2) \text{ For all } \varphi_1, \dots, \varphi_k \in V^*, \varphi_1 \dots \varphi_k = \frac{1}{k!} \sum_{\sigma \in \mathfrak{S}_k} \varphi_{\sigma(1)} \star_t \dots \star_t \varphi_{\sigma(k)}.$$

*Proof.* To prove (1): by Proposition 1.4, one has  $\varphi \star_t \varphi = \varphi^2 + \frac{t}{2}\{\varphi, \varphi\} = \varphi^2$ . Also  $\varphi \star_t \varphi \star_t \varphi = \varphi^3 + \frac{t}{2}\{\varphi, \varphi^2\} = \varphi^3$  and so on.

Then  $(\lambda_1 \varphi + \dots + \lambda_k \varphi_k) \star_t^k = (\lambda_1 \varphi + \dots + \lambda_k \varphi_k)^k$ , for all  $\lambda_1, \dots, \lambda_k \in \mathbb{K}$ . Identifying the coefficient of the term  $\lambda_1 \dots \lambda_k$  on each side, one obtains:

$$\sum_{\sigma \in \mathfrak{S}_k} \varphi_{\sigma(1)} \star_t \dots \star_t \varphi_{\sigma(k)} = \sum_{\sigma \in \mathfrak{S}_k} \varphi_{\sigma(1)} \dots \varphi_{\sigma(k)}$$

and (2) follows.  $\square$

*Remark 1.7.* Consequently, since  $\mathcal{S}^k$  is linearly generated by  $\{\varphi^k \mid \varphi \in V^*\}$ , we can conclude that  $(\mathcal{S}, m_\star)$  is generated by  $V^*$  as an algebra.

Let us denote by  $W(n)$ , or  $W$  when there is no ambiguity, the Weyl algebra generated by  $\{p_1, q_1, \dots, p_n, q_n\}$  with relations  $p_i q_j - q_j p_i = \delta_{ij}$ ,  $p_i p_j - p_j p_i = q_i q_j - q_j q_i = 0$ , for  $i, j = 1, \dots, n$  (see [6]). A well-known fact is that  $W$  is the algebra of polynomial differential operators on  $\mathbb{K}[y_1, \dots, y_n]$  with  $p_i = \frac{\partial}{\partial y_i}$  and

$q_i = y_i \times \dots$ . We can also realize  $W$  as the quotient algebra  $T(V^*)/I$  where  $I$  is the ideal in the tensor algebra  $T(V^*)$  generated by the above relations. Therefore we can consider that  $V^* \subset W$  as will be done in the sequel.

Denote by  $\rho : S \rightarrow W$  the *symmetrization map* defined as:

$$\rho(\varphi_1 \dots \varphi_k) = \frac{1}{k!} \sum_{\sigma \in \mathfrak{S}_k} \varphi_{\sigma(1)} \varphi_{\sigma(2)} \dots \varphi_{\sigma(k)}$$

where the right hand side products are computed in  $W$ . Notice that  $\rho$  is well defined since it is the composition map of the usual symmetrization map from  $S = \text{Sym}(V^*)$  into  $T(V^*)$  with the canonical map from  $T(V^*)$  onto  $W = T(V^*)/I$ .

Using the symmetrization  $\rho$ , we pull back the product of  $W$  on  $S$  by:

$$F \times_{\rho} G := \rho^{-1}(\rho(F) \rho(G)), \quad \forall F, G \in S.$$

We fix the value  $t = 1$  of the parameter, and denote by  $\star$  the  $\star_1$ -Moyal product.

**Proposition 1.8.** *One has  $\times_{\rho} = \star$ . Therefore  $F \star G = \rho^{-1}(\rho(F) \rho(G))$ , for all  $F, G \in S$  and the algebra  $(S, \star)$  is isomorphic to the Weyl algebra  $W$ .*

*Proof.* It is clear from the definition that  $\varphi \times_{\rho}^k = \varphi^k$  for all  $\varphi \in V^*$ ,  $k \in \mathbb{N}$ . As in the proof of Lemma 1.6, one can write  $\varphi_1 \dots \varphi_k = \frac{1}{k!} \sum_{\sigma \in \mathfrak{S}_k} \varphi_{\sigma(1)} \times_{\rho} \dots \times_{\rho} \varphi_{\sigma(k)}$

and using the same Lemma once more, one has  $\varphi \times_{\rho}^k = \varphi^{\star k}$ ,  $\forall \varphi \in V^*$ ,  $k \in \mathbb{N}$ , and  $\varphi_1 \times_{\rho} \dots \times_{\rho} \varphi_k = \varphi_1 \star \dots \star \varphi_k$ ,  $\forall \varphi_1, \dots, \varphi_k \in V^*$ . It results that  $(S, \times_{\rho})$  is generated by  $V^*$  as an algebra and the same holds for  $(S, \star)$  thanks to Remark 1.7. So in order to prove that  $\times_{\rho} = \star$ , we need only to prove that they do coincide on the linear generators of  $S$ , i.e. on the monomials  $\varphi_1 \times_{\rho} \dots \times_{\rho} \varphi_k = \varphi_1 \star \dots \star \varphi_k$ , for all  $\varphi_1, \dots, \varphi_k \in V^*$ . But this is trivial.  $\square$

As a consequence of Proposition 1.8, we can identify the Weyl algebra  $W$  and the Moyal algebra  $S$  endowed with the  $\star$ -product. Such an identification reveals to be useful, since the Moyal  $\star$ -product provides an explicit formula for the product of the Weyl algebra, i.e. for the product of polynomial differential operators. To illustrate, when  $n = 1$  one has the following explicit formula:

$$F \star_t G = \sum_{k \geq 0} \frac{t^k}{2^k k!} \left( \sum_{r+s=k} (-1)^s \binom{k}{s} \frac{\partial^k F}{\partial p_1^r \partial q_1^s} \frac{\partial^k G}{\partial p_1^s \partial q_1^r} \right).$$

Consider now the basis  $\{q_1^i \star p_1^j \mid (i, j) \in \mathbb{N}^2\}$  of  $W(1)$ . By an easy computation, one obtains the following expression in terms of classical orthogonal polynomials:

$$(V) \quad q_1^i \star p_1^j = \begin{cases} (-1)^j \frac{j!}{2^j} L_j^{(i-j)}(2p_1 q_1) q_1^{i-j}, & \text{if } i \geq j \\ (-1)^i \frac{i!}{2^i} L_i^{(j-i)}(2p_1 q_1) p_1^{j-i}, & \text{if } i \leq j \end{cases}.$$

where  $L_\beta^{(\alpha)}$  is the Laguerre polynomial (see e.g. [18]). Notice that this gives also the expression of the basis  $\{q_1^{i_1} \star p_1^{i_1} \star \dots \star q_n^{i_n} \star p_n^{i_n} \mid (i_1, \dots, i_n) \in \mathbb{N}^n\}$  of  $W(n)$  since  $q_1^{i_1} \star p_1^{i_1} \star \dots \star q_n^{i_n} \star p_n^{i_n} = q_1^{i_1} \star p_1^{i_1} \cdot \dots \cdot q_n^{i_n} \star p_n^{i_n}$ .

Let us now define a filtration and also a  $\mathbb{Z}_2$ -grading of  $W$ . For the filtration, we keep the filtration of  $S$ , that is, we set:

$$W_k := \bigoplus_{r \leq k} S^r.$$

Using (II) and Proposition 1.4, it is clear that this is indeed a filtration of  $W$  and that the associated graded algebra is  $S$ .

What about the  $\mathbb{Z}_2$ -grading of  $W$ ? It can be defined in two ways: first one, consider  $W$  as the algebra of polynomial differential operators on  $\mathbb{K}[y_1, \dots, y_n]$ . Define an element of  $W$  to be *even* if it maps even polynomials into even polynomials and odd polynomials into odd ones. An *odd* element takes even polynomials into odd polynomials and vice versa. Evidently, this defines a  $\mathbb{Z}_2$ -grading on  $W$  and one has  $\deg_{\mathbb{Z}_2}(p_1^{r_1} \star q_1^{s_1} \star \dots \star p_n^{r_n} \star q_n^{s_n}) \equiv (\sum_{i=1}^n r_i + \sum_{i=1}^n s_i) \pmod{2}$ .

On the other hand, thanks to Proposition 1.4, one can define a  $\mathbb{Z}_2$ -grading on  $W$  by

$$W_{\bar{0}} := \bigoplus_{k \geq 0} S^{2k} \text{ and } W_{\bar{1}} := \bigoplus_{k \geq 0} S^{2k+1}.$$

This  $\mathbb{Z}_2$ -grading is exactly the preceding one. Therefore  $W$  can be endowed with two Lie structures:

- a Lie algebra structure given by

$$[F, G]_{\mathcal{L}} := F \star G - G \star F, \text{ for all } F, G \in W.$$

From (IV), one has  $[W_k, W_\ell]_{\mathcal{L}} \subset W_{k+\ell-2}$ , so  $W$  is a filtered Lie algebra for the shifted filtration  $W_k^{\mathcal{L}} := W_{k+2}$  and its associated graded Lie algebra is  $S$  endowed with the Poisson bracket (using (IV) once again). A widely known fact is that  $[W, W]_{\mathcal{L}} = W$  (we shall give a proof in Section 3), so there does not exist a non zero trace map on  $W$  satisfying  $\text{Tr}(F \star G) = \text{Tr}(G \star F)$ , for  $F, G \in W$ , i.e.  $H^1(W) = \{0\}$  for the Lie algebra cohomology.

- a Lie superalgebra structure given by

$$[F, G] := F \star G - (-1)^{fg} G \star F, \text{ for all } F, G \in W.$$

with  $f = \deg_{\mathbb{Z}_2}(F)$  and  $g = \deg_{\mathbb{Z}_2}(G)$ .

A significant difference between the Lie algebra and the Lie superalgebra structures is set in the following Proposition:

**Proposition 1.9.** *Let  $\text{Str}$  be the evaluation at 0 of the commutative algebra  $S$ ,*

$$\text{Str}(F) := F(0), \text{ for all } F \in S.$$

*Then  $\text{Str}$  is a supertrace on  $W$ , that is,  $\text{Str}$  is homogeneous and satisfies*

$$\text{Str}([F, G]) = 0, \text{ for all } F, G \in W.$$

*Proof.* From Proposition 1.4,

$$F \star G = \sum_{k=0}^{\min(f,g)} C_k(F, G)$$

with  $C_k(F, G) \in \mathbb{S}^{f+g-2k}$  for  $F \in \mathbb{S}^f$  and  $G \in \mathbb{S}^g$ . One has  $C_k(F, G) \in \mathbb{K}$  if and only if  $k = \frac{f+g}{2}$ . Since  $k$  runs from 0 to  $\min(f, g)$ , one can deduce: either  $f \neq g$  and then  $\text{Str}(F \star G) = 0$ , or  $f = g$  and  $\text{Str}(F \star G) = C_f(F, G)$ . In this later case, if  $f$  is even, then by (III),  $C_f(F, G) = C_f(G, F)$ , so  $\text{Str}(F \star G) = \text{Str}(G \star F)$ . If  $f$  is odd,  $C_f(F, G) = -C_f(G, F)$  (III), so  $\text{Str}(F \star G) = -\text{Str}(G \star F)$ . Either way,  $\text{Str}([F, G]) = 0$ . Finally,  $\text{Str}(F) = 0$  if  $F$  is odd, so  $\text{Str}$  is homogeneous.  $\square$

*Remark 1.10.* Assume that  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ . The Moyal  $\star$ -product is clearly not restricted to live only on polynomial functions: obviously, assuming that  $t$  is a formal parameter, Definition 1.2 defines an associative deformation of  $\mathcal{C}^\infty(V)$ . All above properties are true, in the formal sense, thanks to the density of polynomials inside  $\mathcal{C}^\infty(V)$  (with its usual Fréchet topology, see e.g. [19]).

On the other hand, if  $F, G \in \mathcal{S}(V)$  (fast decreasing smooth functions), one has  $\int F \star G = \int F G$ , so one can define a trace by  $\text{Tr}(F) = \int F$  and this has very important consequences (see e.g. [5, 17]).

## 2. EMBEDDING $\mathfrak{sp}(2n)$ AND $\mathfrak{osp}(1, 2n)$ INTO $W$

In the sequel, we denote by  $\mathfrak{g}$  the Lie superalgebra  $\mathfrak{osp}(1, 2n)$  and by  $\mathfrak{g}_0$  its even part, i.e.  $\mathfrak{g}_0 = \mathfrak{sp}(2n)$ . The Weyl algebra  $W$  is endowed with the super bracket  $[F, G] = F \star G - (-1)^{fg} G \star F$ , for  $F \in \mathbb{S}^f, G \in \mathbb{S}^g$ . Denote by  $\text{ad}$  the corresponding adjoint representation.

There exists a well-known embedding of  $\mathfrak{g}_0$  into  $W$  given by:

**Proposition 2.1.** *Given  $X \in S^2$ , denote by  $\text{ad}_p(X) := \{X, \cdot\}$ . Then*

$$\text{ad}_p(X)|_{S^1} = \text{ad}(X)|_{S^1} \in \mathfrak{sp}(2n).$$

*Moreover  $[S^2, S^2] = \{S^2, S^2\} \subset S^2$ , so that  $S^2$  is a Lie algebra for the Poisson bracket or the  $\star$ -bracket and the map*

$$X \mapsto \text{ad}_p(X)|_{S^1}$$

*is a Lie algebra isomorphism from  $S^2$  onto  $\mathfrak{sp}(2n)$ .*

*Proof.* Set a non degenerated skew symmetric bilinear form  $\Phi$  on  $S^1$  by  $\Phi(\varphi, \psi) = \frac{1}{2}\{\varphi, \psi\} = \frac{1}{2}[\varphi, \psi]_{\mathcal{S}}$ , for  $\varphi, \psi \in S^1 = V^*$ . Notice that when  $X \in S^2$ , one has  $\text{ad}_p(X) = \text{ad}(X)$  by Proposition 1.5. Then  $[S^2, S^2] = \{S^2, S^2\} \subset S^2$  which implies that  $S^2$  is a Lie algebra for the Poisson bracket (=  $\star$ -bracket). Since  $\text{ad}_p(X)$  is a derivation for the Poisson bracket, the bilinear form  $\Phi$  is invariant and that means  $\text{ad}_p(X)|_{S^1} \in \mathfrak{sp}(2n)$ . Let  $\theta$  be the map from  $S^2$  to  $\mathfrak{sp}(2n)$  defined as  $\theta(X) = \text{ad}_p(X)|_{S^1}$ . So  $\theta$  is clearly a Lie algebra homomorphism, which is injective since  $\text{ad}_p(X)|_{S^1} = 0$  implies  $X \in \mathbb{K}$ , thus  $X = 0$ . Besides  $\theta$  is onto since  $\dim(S^2) = \dim(\mathfrak{sp}(2n))$ .  $\square$

In the remaining of the paper, we shall identify  $\mathfrak{g}_{\bar{0}} = \mathfrak{sp}(2n)$  and  $S^2$  via the isomorphism in Proposition 2.1.

Let us define an embedding of  $\mathfrak{g}$  into the Weyl algebra  $W$  extending the previous embedding of  $\mathfrak{g}_{\bar{0}}$  into  $W$ . Long ago, this embedding was already well-known among mathematical physicists (e.g. [7]) and it was used for instance to develop singleton Anti-de-Sitter theories [7]. This embedding was also described in [13] and [8]. We give the proof in [8]:

**Definition 2.2.** The *twisted adjoint action* of the Lie superalgebra  $W$  on itself is defined as:

$$\text{ad}'(F)(G) := F \star G - (-1)^{f(g+1)} G \star F, \forall F \in S^f, G \in S^g.$$

Such a twisted action is typical of supersymmetry (see [2, 9]). Remark that if  $F \in W_{\bar{0}}$ , then  $\text{ad}'(F) = \text{ad}(F)$ .

**Proposition 2.3.** Let  $\mathfrak{s} = S^1 \oplus S^2$  with  $\mathbb{Z}_2$ -grading induced by  $W$ . Then  $\mathfrak{s}$  is a subalgebra of the Lie superalgebra  $W$ . Moreover  $\mathfrak{s} \simeq \mathfrak{osp}(1, 2n)$ .

*Proof.* If  $\varphi, \psi \in S^1$ , one has  $\varphi \star \psi = \varphi\psi + \frac{1}{2}\{\varphi, \psi\}$ , so  $[\varphi, \psi] = 2\varphi\psi \in S^2$ . If  $\varphi \in S^1$  and  $X \in S^2$ , one has  $[\varphi, X] = \{\varphi, X\}$  (by Proposition 1.5), and  $\{\varphi, X\} \in S^1$ . So  $\mathfrak{s}$  is a subalgebra of the Lie superalgebra  $W$ .

Set  $V = V_{\bar{0}} \oplus V_{\bar{1}}$  where  $V_{\bar{0}} = \mathbb{K}$  and  $V_{\bar{1}} = S^1$ . It is easy to check that  $\text{ad}'(\mathfrak{s})(V) \subset V$ . For example, if  $\varphi, \psi \in S^1$ , one has  $\text{ad}'(\varphi)(\psi) = [\varphi, \psi]_{\mathcal{L}} = \{\varphi, \psi\} \in \mathbb{K}$  and  $\text{ad}'(\varphi)(1) = 2\varphi$ .

Define a bilinear form  $\Theta$  on  $V$  by  $\Theta(\varphi, \psi) = \frac{1}{2}\{\varphi, \psi\}$  and  $\Theta(1, 1) = -1$ . It is clear that  $\Theta$  is a supersymmetric form, i.e.  $\Theta(A, B) = (-1)^{ab}\Theta(B, A)$  for all  $A \in V_a$  and  $B \in V_b$ . Straightforward computations show that  $\Theta$  is  $\text{ad}'$ -invariant. Therefore we obtain a Lie superalgebra homomorphism  $\phi$  that maps  $X$  to  $\text{ad}'(X)|_V$  from  $\mathfrak{s}$  into  $\mathfrak{osp}(1, 2n)$ . Remark that  $\text{ad}'(X)|_V = 0$  for  $X \in S^1$  implies that  $\text{ad}'(X)(1) = 2X = 0$ , so  $\phi$  is injective and since  $\dim(\mathfrak{s}) = \dim(\mathfrak{osp}(1, 2n))$ ,  $\phi$  is an isomorphism.  $\square$

In the remaining of the paper, we shall identify

$$\mathfrak{g} = \mathfrak{osp}(1, 2n) \quad \text{and} \quad \mathfrak{s} = S^1 \oplus S^2$$

via the isomorphism given in Proposition 2.3. Then

$$\mathfrak{g} = S^1 \oplus S^2, \quad \mathfrak{g}_{\bar{1}} = S^1 \quad \text{and} \quad \mathfrak{g}_{\bar{0}} = S^2 = [\mathfrak{g}_{\bar{1}}, \mathfrak{g}_{\bar{1}}] = \mathfrak{sp}(2n).$$

The root system of  $\mathfrak{g}$  is easily deduced: consider

$$H_i = -\frac{1}{2} [p_i, q_i] \text{ for } 1 \leq i \leq n.$$

Set  $\mathfrak{h} = \text{span}\{H_1, \dots, H_n\}$ . It turns out that  $\mathfrak{h}$  is a Cartan subalgebra of  $\mathfrak{g}_{\bar{0}}$  and  $\mathfrak{g}$ . Let  $\{\omega_1, \dots, \omega_n\}$  be the dual basis. The root vectors and corresponding roots are:

- in  $\mathfrak{g}_{\bar{1}}$ :  
 $p_i$  with root  $\omega_i$  (positive),  $q_i$  with root  $-\omega_i$ .

- in  $\mathfrak{g}_{\bar{0}}$ :
  - $[p_i, q_j]$  ( $i \neq j$ ) with root  $\omega_i - \omega_j$  (positive if  $i < j$ ),
  - $[p_i, p_j]$  with root  $\omega_i + \omega_j$  (positive),
  - $[q_i, q_j]$  with root  $-(\omega_i + \omega_j)$ .

The fundamental root system is  $\{\omega_i - \omega_{i+1} \ (i = 1, \dots, n-1), \omega_n\}$  with corresponding root vectors  $[p_i, q_{i+1}] \ (i = 1, \dots, n-1)$  and  $p_n$ . These vectors generate the subalgebra  $\mathfrak{n}^+$  of  $\mathfrak{g}$  with basis the positive root vectors. Any simple finite dimensional  $\mathfrak{g}$ -module has a *highest weight vector*  $v$  of weight  $\lambda \in \mathfrak{h}^*$  satisfying if  $Hv = \lambda(H)v$  for all  $H \in \mathfrak{h}$  and  $\mathfrak{n}^+v = 0$ .

**Proposition 2.4.** *The  $\mathfrak{g}$ -module  $\text{ad}_{\mathbb{P}}(\mathfrak{g}_{\bar{0}})|_{S^k}$  is a simple module with highest weight vector  $p_1^k$  of weight  $k\omega_1$ .*

*Proof.* See [16] or [13] for a proof using Weyl's formula. We provide here a short direct proof: it is clear that  $\text{ad}_{\mathbb{P}}(\mathfrak{g}_{\bar{0}})(S^k) \subset S^k$ . Now given  $\varphi \in S^1$ ,  $\varphi \neq 0$ , one can construct a Darboux basis with respect to  $\Phi$  such that  $\varphi$  is the first basis vector. It results that there exists  $A \in \text{Sp}(2n)$  such that  $\varphi = A(p_1)$ . Denoting by  $A$  as well the corresponding isomorphism of the commutative algebra  $S$ , one has  $\varphi^k = A(p_1^k)$ . Since  $S^k = \text{span}\{\varphi^k \mid \varphi \in S^1\}$ , one has  $\text{span}(\text{Sp}(2n))(p_1^k) = S^k$ . One has a representation of  $\text{Sp}(2n)$  in  $S^k$ ,  $A \mapsto A|_{S^k}$  with differential  $\text{ad}_{\mathbb{P}}(\mathfrak{g}_{\bar{0}})|_{S^k}$ . As a consequence, the  $\mathfrak{g}_{\bar{0}}$ -submodule of  $S^k$  generated by  $p_1^k$  is  $S^k$  itself. Since  $p_1^k$  is a highest weight vector of weight  $k\omega_1$ ,  $S^k$  is a simple  $\mathfrak{g}_{\bar{0}}$ -module with highest weight  $k\omega_1$ .  $\square$

The next Proposition was proved in [15] for  $\mathfrak{osp}(1, 2)$  and in [13] for  $\mathfrak{osp}(1, 2n)$ . We give a simplified proof that uses the Moyal  $\star$ -product.

**Proposition 2.5.** *Let  $A_k = S^{2k-1} \oplus S^{2k}$ ,  $k \geq 1$  and  $A_0 = \{0\}$ . Then  $A_k$  is stable by  $\text{ad}(\mathfrak{g})$ . Moreover,  $A_k$  is a simple  $\mathfrak{g}$ -module with highest weight vector  $p_1^{2k}$  of weight  $2k\omega_1$ .*

*Proof.* Let  $X \in \mathfrak{g}_{\bar{1}}$  and  $F \in S^{2k}$ . Then  $\text{ad}(X)(F) = \{X, F\}$  by Proposition 1.5, hence  $\text{ad}(X)(F) \in S^{2k-1}$ . Now assume that  $F \in S^{2k-1}$ , then  $\text{ad}(X)(F) = X \star F + F \star X = 2XF \in S^{2k}$ . So  $A_k$  is  $\text{ad}(\mathfrak{g})$ -stable. By Proposition 2.4,  $A_k = S^{2k-1} \oplus S^{2k}$  is its decomposition into isotypic components under the action of  $\text{ad}(\mathfrak{g}_{\bar{0}})$ . Any  $\mathfrak{g}$ -submodule  $U$  of  $A_k$  must be decomposed as  $U = (U \cap S^{2k}) \oplus (U \cap S^{2k-1})$  and since  $S^{2k}$  and  $S^{2k-1}$  are simple  $\mathfrak{g}_{\bar{0}}$ -modules, one of them is contained in  $U$  if  $U \neq \{0\}$  from what one deduces that  $U = A_k$  using the beginning of the proof. Moreover,  $p_1^{2k}$  is clearly a highest vector of weight  $2k\omega_1$ .  $\square$

Next we decompose the twisted adjoint representation of  $\mathfrak{g}$  in  $W$ :

**Proposition 2.6.** *Let  $B_k = S^{2k} \oplus S^{2k+1}$ ,  $k \geq 0$ . Then  $B_k$  is stable by  $\text{ad}'(\mathfrak{g})$ . Moreover,  $B_k$  is a simple  $\mathfrak{g}$ -module with highest vector  $p_1^{2k+1}$  of weight  $(2k+1)\omega_1$ .*

*Proof.* Let  $X \in \mathfrak{g}_{\bar{1}}$  and  $F \in S^{2k}$ . Then  $\text{ad}'(X)(F) = 2XF \in S^{2k}$  and if  $F \in S^{2k+1}$ , then  $\text{ad}'(X)(F) = \{X, F\} \in S^{2k}$ , so  $B_k$  is  $\text{ad}'(\mathfrak{g})$ -stable. By the same arguments in the proof of the previous Proposition, one obtains easily that  $B_k$  is a simple  $\mathfrak{g}$ -module and  $p_1^{2k+1}$  is clearly a highest weight vector of weight  $(2k+1)\omega_1$ .  $\square$

*Remark 2.7.* From Propositions 2.5 and 2.6, it results that the only homomorphism of  $\mathfrak{g}$ -modules from  $(W, \text{ad}(\mathfrak{g}))$  into  $(W, \text{ad}'(\mathfrak{g}))$  (or the other way around) is zero.

### 3. DECOMPOSITION OF ADJOINT AND TWISTED ADJOINT $W$ -MODULES

Let us recall our conventions: the Weyl algebra  $W$  is a Lie algebra with bracket denoted by

$$[A, B]_{\mathcal{L}} = A \star B - B \star A, \quad \forall A, B \in W.$$

Denote by  $\text{ad}_{\mathcal{L}}$  the corresponding adjoint representation. Thanks to the  $\mathbb{Z}_2$ -gradation of the associative algebra  $W$ , there is a twisted adjoint representation of the Lie algebra  $W$  defined as:

$$\text{ad}'_{\mathcal{L}}(A)(B) = A \star B - (-1)^a B \star A, \quad \forall A \in W_a, B \in W.$$

Note that  $\text{ad}'_{\mathcal{L}}(W_{\bar{0}})(W_i) \subset W_i$  ( $i = \bar{0}, \bar{1}$ ) and  $\text{ad}'_{\mathcal{L}}(W_{\bar{1}})$  takes  $W_{\bar{0}}$  into  $W_{\bar{1}}$  and vice versa.

The Weyl algebra is also a Lie superalgebra with bracket denoted by

$$[A, B] = A \star B - (-1)^{ab} B \star A, \quad \forall A \in W_a, B \in W_b.$$

Denote by  $\text{ad}$  the corresponding adjoint representation. There is a twisted adjoint representation of the Lie superalgebra  $W$  defined as:

$$\text{ad}'(A)(B) = A \star B - (-1)^{a(b+1)} B \star A, \quad \forall A \in W_a, B \in W_b.$$

This twisted adjoint representation was used in Proposition 2.4 to prove the embedding of  $\mathfrak{g} = \mathfrak{osp}(1, 2n)$  in  $W$ .

The decomposition of  $\text{ad}_{\mathcal{L}}(\mathfrak{g}_{\bar{0}}) = \text{ad}'_{\mathcal{L}}(\mathfrak{g}_{\bar{0}})$  is given in Proposition 2.4, the decomposition of  $\text{ad}(\mathfrak{g})$  is given in Proposition 2.5 and in Proposition 2.6 for  $\text{ad}'(\mathfrak{g})$ .

We will now examine  $\text{ad}_{\mathcal{L}}$ ,  $\text{ad}'_{\mathcal{L}}$  as Lie algebra representations of  $W$ ,  $\text{ad}$  and  $\text{ad}'$  as Lie superalgebra representations of  $W$ . The main technical argument is given by the Theorem:

**Theorem 3.1.** *Consider the  $\text{ad} = \text{ad}_{\mathcal{L}}$ -action of  $\mathfrak{g}_{\bar{0}}$  on  $W$ . One has:*

- (1) *The map  $F \otimes G \mapsto F \star G$  is a homomorphism of  $\mathfrak{g}_{\bar{0}}$ -modules from  $S^\ell \otimes S^m$  onto  $\bigoplus_{k=0}^{\min(\ell, m)} S^{\ell+m-2k}$ , so one has*

$$S^\ell \star S^m = \bigoplus_{k=0}^{\min(\ell, m)} S^{\ell+m-2k}.$$

- (2) *The map  $F \otimes G \mapsto [F, G]_{\mathcal{L}}$  is a homomorphism of  $\mathfrak{g}_{\bar{0}}$ -modules from  $S^\ell \otimes S^m$  onto  $\bigoplus_{2p+1 \leq \min(\ell, m)} S^{\ell+m-2(2p+1)}$ , so one has*

$$[S^\ell, S^m]_{\mathcal{L}} = \bigoplus_{2p+1 \leq \min(\ell, m)} S^{\ell+m-2(2p+1)}.$$

- (3) *The map  $F \otimes G \mapsto [F, G]$  is a homomorphism of  $\mathfrak{g}_{\bar{0}}$ -modules from  $S^\ell \otimes S^m$  onto*

- $\bigoplus_{2p+1 \leq \min(\ell, m)} S^{\ell+m-2(2p+1)}$  if  $\bar{\ell} \bar{m} \equiv \bar{0}$ , so in this case one has

$$[S^\ell, S^m] = \bigoplus_{2p+1 \leq \min(\ell, m)} S^{\ell+m-2(2p+1)}.$$

- $\bigoplus_{2p+1 \leq \min(\ell, m)} S^{\ell+m-4p}$  if  $\bar{\ell} \bar{m} \equiv \bar{1}$ , so in this case one has

$$[S^\ell, S^m] = \bigoplus_{2p \leq \min(\ell, m)} S^{\ell+m-4p}.$$

- (4) The map  $C_k$  is a homomorphism of  $\mathfrak{g}_{\bar{0}}$ -modules from  $S^\ell \otimes S^m$  into  $S^{\ell+m-2k}$ , so one has

$$\begin{cases} C_k(S^\ell, S^m) = S^{\ell+m-2k}, & \text{if } 0 \leq k \leq \min(\ell, m) \\ 0, & \text{otherwise.} \end{cases}$$

To prove the above Theorem, we need the following Lemma:

**Lemma 3.2.** Let  $\mathfrak{h}$  be a Lie algebra,  $\mathcal{U} = \mathcal{U}(\mathfrak{h})$  its enveloping algebra and  $V$  a semisimple  $\mathfrak{h}$ -module which decomposes as  $V = \bigoplus_{s=1}^r V_s$  with  $V_s$  simple and two by two non isomorphic. Let  $v = \sum_{s=1}^r v_s$  with  $v_s \in V_s$  and  $v_s \neq 0$ . Then  $V = \mathcal{U}(\mathfrak{h})v$ .

*Proof.* The  $\mathcal{U}$ -module  $V$  is semisimple and  $V = \bigoplus_{s=1}^r V_s$  is its decomposition into isotypic components. Let  $\pi_s$  be the projection of  $V$  into  $V_s$ . Then  $\pi_s$  is an element of the bicommutant of the  $\mathcal{U}$ -module  $V$  and by the Jacobson density theorem, there exists  $u \in \mathcal{U}$  such that  $v_s = \pi_s(v) = uv$  and the result follows from the simplicity of each  $V_s$ .  $\square$

*Proof.* (Proof of Theorem 3.1)

For all  $X \in \mathfrak{g}_{\bar{0}}$ ,  $\mathfrak{g}_{\bar{0}}$  acts by  $\text{ad}(X)$ , which is a derivation of the  $\star$ -product. One has  $F \star G = \sum_{k=0}^{\min(\ell, m)} C_k(F, G)$ , for all  $F \in S^\ell$ ,  $G \in S^m$  with  $C_k(F, G) \in S^{\ell+m-2k}$  which is irreducible under the  $\text{ad}(\mathfrak{g}_{\bar{0}})$ -action. It results that the map  $F \otimes G \mapsto F \star G$  is a homomorphism of  $\mathfrak{g}_{\bar{0}}$ -modules and so does any of the maps  $C_k$ .

If  $L = \bigoplus_{k=0}^{\min(\ell, m)} S^{\ell+m-2k}$ , denote by  $\rho$  the map  $\rho: S^\ell \otimes S^m \rightarrow L$ ,  $\rho(F \otimes G) = F \star G$ . We compute  $\rho(p_1^\ell \otimes q_1^m)$ : using (I), one has  $\rho(p_1^\ell \otimes q_1^m) = \ell m p_1^{\ell-1} \otimes q_1^{m-1}$ , so

$$\rho^k(p_1^\ell \otimes q_1^m) = \ell(\ell-1) \dots (\ell-k+1)m(m-1) \dots (m-k+1) p_1^{\ell-k} \otimes q_1^{m-k},$$

if  $k \leq \min(\ell, m)$  and 0 otherwise. Then

$$C_k(p_1^\ell, q_1^m) = \frac{k!}{2^k} \binom{\ell}{k} \binom{m}{k} p_1^{\ell-k} q_1^{m-k},$$

if  $k \leq \min(\ell, m)$  and 0 otherwise. Thus

$$\rho(p_1^\ell \otimes q_1^m) = \sum_{k=0}^{\min(\ell, m)} \frac{k!}{2^k} \binom{\ell}{k} \binom{m}{k} p_1^{\ell-k} q_1^{m-k}.$$

We can apply Lemma 3.2 to the  $\mathfrak{g}_{\bar{0}}$ -submodule  $\rho(S^\ell \otimes S^m)$  of  $L$  with  $v = \rho(p_1^\ell \otimes q_1^m)$  to obtain that  $\rho(S^\ell \otimes S^m) = L$ .

But  $C_k$  is a homomorphism of  $\mathfrak{g}_{\bar{0}}$ -modules from  $S^\ell \otimes S^m$  into  $S^{\ell+m-2k}$  and one has  $C_k(S^\ell, S^m) = \{0\}$  if  $k > \min(\ell, m)$ . When  $k \leq \min(\ell, m)$ ,  $C_k(S^\ell, S^m)$  is a non zero submodule of the simple module  $S^{\ell+m-2k}$  since  $C_k(p_1^\ell, q_1^m) \neq 0$ . Therefore  $C_k(S^\ell, S^m) = S^{\ell+m-2k}$ .

The proof of (2) is completely similar, using (IV). The same reasons can also be used to prove (3): the definition of  $[\cdot, \cdot]$  implies that  $[F, G] = 2 \sum_{p \geq 0} C_{2p+1}(F, G)$  if  $\overline{f\bar{g}} \equiv \bar{0}$  and  $[F, G] = 2 \sum_{p \geq 0} C_{2p}(F, G)$  if  $\overline{f\bar{g}} \equiv \bar{1}$ .  $\square$

*Remark 3.3.*

- (1) Let us consider the case  $n = 1$ ,  $W = W(1)$ . Then  $\mathfrak{g}_{\bar{0}} = \mathfrak{sl}(2)$  and  $S^\ell$  is the simple  $(\ell + 1)$ -dimensional  $\mathfrak{g}_{\bar{0}}$ -module, denoted by  $D\left(\frac{\ell}{2}\right)$ . Then  $\dim(S^\ell \otimes S^m) = \dim\left(\bigoplus_{k=0}^{\min(\ell, m)} S^{\ell+m-2k}\right)$ . So the map  $F \otimes G \mapsto F \star G$  of Proposition 3.1(1) is an isomorphism from  $D\left(\frac{\ell}{2}\right) \otimes D\left(\frac{m}{2}\right)$  onto

$$D\left(\frac{\ell}{2} + \frac{m}{2}\right) \oplus D\left(\frac{\ell}{2} + \frac{m}{2} - 1\right) \oplus \cdots \oplus D\left(\left|\frac{\ell}{2} - \frac{m}{2}\right|\right),$$

providing an explicit and very handy formula for the computation of Clebsch-Gordan coefficients. Such a formula was used for instance in [1] to compute the commutation rules of high dimensional Lie algebras when usual Clebsch-Gordan formulas were hopeless. Unfortunately, it is easy to check that the  $\star$ -product will provide only a partial decomposition of the tensor product  $S^\ell \otimes S^m$  when  $n \geq 2$ .

- (2) Using Proposition 3.1, one has the following identities:

$$\begin{aligned} \text{(VI)} \quad [S^1, S^k]_{\mathcal{L}} &= S^{k-1}, \quad [S^2, S^k]_{\mathcal{L}} = S^k \quad (k \geq 1), \\ [S^3, S^k]_{\mathcal{L}} &= S^{k-3} \oplus S^{k+1} \quad (k \geq 3), \\ [S^3, S^2]_{\mathcal{L}} &= S^3 \quad \text{and} \quad [S^3, S^1]_{\mathcal{L}} = S^2. \end{aligned}$$

$$\begin{aligned} \text{(VII)} \quad [S^1, S^{2k}] &= S^{2k-1} \quad (k \geq 1), \quad [S^1, S^{2k+1}] = S^{2k+2} \quad (k \geq 0), \\ [S^3, S^{2k-1}] &= S^{2k-2} \oplus S^{2k+2} \quad (k \geq 2), \quad [S^3, S^1] = S^4, \\ [S^3, S^{2k}] &= S^{2k-3} \oplus S^{2k+1} \quad (k \geq 2) \quad \text{and} \quad [S^3, S^2] = S^3. \end{aligned}$$

These identities turn out to be quite useful.

**Proposition 3.4.** *One has*

$$\ker(\text{Str}) = [\mathfrak{g}_{\bar{0}}, W] = [W, W] = \bigoplus_{k \geq 1} S^k.$$

*Proof.* We need the following obvious result:

*If  $\mathfrak{h}$  is a Lie algebra and  $U$  a non trivial simple  $\mathfrak{h}$ -module, then  $\mathfrak{h}U = U$ .*

Applying it to  $\mathfrak{g}_{\bar{0}}$  and the simple  $\mathfrak{g}_{\bar{0}}$ -module  $S^k$  ( $k \geq 1$ ) we get that  $[\mathfrak{g}_{\bar{0}}, S^k] = S^k$  (alternatively,  $[S^2, S^k] = S^k$  by Formula (VI)). As a consequence,  $[\mathfrak{g}_{\bar{0}}, \bigoplus_{k \geq 1} S^k] = \bigoplus_{k \geq 1} S^k$ . By Proposition 1.9, one obtains  $[W, W] \subset \ker(\text{Str})$ , so finally we conclude that  $\bigoplus_{k \geq 1} S^k \subset [W, W] \subset \ker(\text{Str})$ , but since  $\text{codim}\left(\bigoplus_{k \geq 1} S^k\right) = \text{codim}(\ker(\text{Str})) = 1$ , the result is proved.  $\square$

**Corollary 3.5.** [13] *One has  $W = \mathbb{K} \oplus [W, W]$ .*

*Remark 3.6.* The use of the supertrace  $\text{Str}$  is quite natural and provides a real simplification of Musson's proof which does not use  $\text{Str}$ . Moreover, it provides an enlightening on the origin of the result.

**Theorem 3.7.**

- (1) *The representation  $(\text{ad}_{\mathcal{L}}, W)$  of the Lie algebra  $W$  is indecomposable with Jordan-Hölder series  $\{0\} \subset \mathbb{K} \subset W$  and one has  $\text{ad}_{\mathcal{L}}(W)(W) = W$ .*
- (2) *The representation  $(\text{ad}'_{\mathcal{L}}, W)$  of the Lie algebra  $W$  is indecomposable,  $[W, W]$  is a simple subrepresentation,  $W/[W, W]$  is the trivial representation and there exists a non trivial  $[W, W]$ -valued cocycle  $\xi$  defined by  $\xi(F) = 0$  if  $F$  is even, and  $\xi(F) = 2F$  if  $F$  is odd.*
- (3) *The representation  $(\text{ad}, [W, W])$  of the Lie superalgebra  $W$  is simple. Moreover, the  $\text{ad}(W)$ -module  $W$  decomposes as  $W = \mathbb{K} \oplus [W, W]$ .*
- (4) *The representation  $(\text{ad}', W)$  of the Lie superalgebra  $W$  is simple.*

*Proof.*

- (1) Let  $M$  be a non trivial  $\text{ad}_{\mathcal{L}}(W)$ -module. Then  $M$  can be decomposed into isotypic components under the  $\text{ad}_{\mathcal{L}}(\mathfrak{g}_{\bar{0}})$ -action. By Proposition 2.4, there exists  $k_0$  such that  $S^{k_0} \subset M$ . If  $k_0 > 0$ , using (VI) one has  $[S^1, S^{k_0}]_{\mathcal{L}} = S^{k_0-1} \subset M$  implying that  $\bigoplus_{k \leq k_0} S^k \subset M$ . Using  $\text{ad}_{\mathcal{L}}(S^3)$  and (VI), one deduces that  $S^{k_0+1} \subset M$  and repeating the argument, that  $M = W$ .

If the only  $k$  such that  $S^k \subset M$  is 0, then  $M = S^0 = \mathbb{K}$ . Therefore, there is exactly one non trivial invariant subspace namely  $\mathbb{K}$ . It results that  $W/\mathbb{K}$  is simple, that the representation  $(\text{ad}_{\mathcal{L}}, W)$  is indecomposable, with Jordan-Hölder series  $\{0\} \subset \mathbb{K} \subset W$ . To finish, notice that  $\text{ad}_{\mathcal{L}}(W)(W)$  is invariant, contains strictly  $\mathbb{K}$ , so  $\text{ad}_{\mathcal{L}}(W)(W) = W$ .

- (2) We start by proving that  $\text{Str}(\text{ad}'_{\mathcal{L}}(F)(G)) = 0, \forall F, G \in W$ . If  $F$  is even, or if  $F$  and  $G$  are odd, then  $\text{ad}'_{\mathcal{L}}(F)(G) = \text{ad}(F)(G)$ , so we apply Proposition 1.9. Now assume that  $F \in S^k, k$  odd and  $G \in S^\ell, \ell$  even. Then  $\text{ad}'_{\mathcal{L}}(F)(G) = F \star G + G \star F = 2 \sum_{2s \leq \min(k, \ell)} C_{2s}(F, G)$  and  $C_{2s}(F, G) \in S^{k+\ell-4s}$ . But  $k + \ell - 4s > 0$ , so  $\text{ad}'_{\mathcal{L}}(F)(G) \in \bigoplus_{r > 0} S^r = \ker(\text{Str})$ .

It results that  $\text{ad}'_{\mathcal{L}}(W)(W) \subset \ker(\text{Str}) = [W, W]$  and a fortiori,  $[W, W]$  is stable. The quotient  $W/[W, W]$  is the trivial representation. If  $F$  is invariant under  $\text{ad}'_{\mathcal{L}}(W)$ , it is invariant under  $\text{ad}_{\mathcal{L}}(\mathfrak{g}_{\bar{0}})$  and has to be a constant. But  $\text{ad}'_{\mathcal{L}}(G)(1) = 2G$ , if  $G$  is odd, so  $F = 0$ . Therefore the extension

$$0 \rightarrow [W, W] \rightarrow W \rightarrow \mathbb{K} = W/[W, W] \rightarrow 0$$

is non trivial with corresponding non trivial  $[W, W]$ -valued cocycle  $\xi$  defined by:

$$\xi(F) = \text{ad}'_{\mathcal{L}}(F)(1) = \begin{cases} 0, & \text{if } F \text{ is even.} \\ 2F, & \text{if } F \text{ is odd.} \end{cases}$$

Assume now that  $M$  is a non zero  $\text{ad}'_{\mathcal{L}}(W)$ -invariant subspace. As in (1), there exists there exists  $k_0$  such that  $S^{k_0} \subset M$ . Let  $X \in S^1$  and  $F \in S^{k_0}$ . One has  $\text{ad}'_{\mathcal{L}}(X)(F) = 2XF$ , so  $\text{ad}'_{\mathcal{L}}(S^1)(S^{k_0}) = S^{k_0+1}$  and  $\bigoplus_{k \geq k_0} S^k \subset M$ . Take  $X \in S^3$ , then  $\text{ad}'_{\mathcal{L}}(X)(F) = 2XF + 2C_2(X, F)$ . As in the proof of Theorem 3.2, it results that  $\text{ad}'_{\mathcal{L}}(S^3)(S^{k_0}) = S^{k_0+3} \oplus S^{k_0-1}$  if  $k_0 > 1$ , so  $S^{k_0-1}$  is contained in  $M$  and then  $\bigoplus_{k \geq 1} S^k \subset M$ . Thus, either  $M = \bigoplus_{k \geq 1} S^k = [W, W]$  or  $M = W$ .

- (3) Thanks to Proposition 3.4, it remains to show that the representation  $(\text{ad}, [W, W])$  is simple. This is an easy consequence of (VII): let  $M$  be a non zero invariant subspace in  $[W, W]$ . Since  $M$  is  $\text{ad}(\mathfrak{g})$ -stable, it decomposes into the isotypic components of  $[W, W] = \bigoplus_{k \geq 1} A_k$  with  $A_k = S^{2k-1} \oplus S^{2k}$  (see Proposition 2.5). Hence  $M = \bigoplus_{A_k \subset M} A_k$ . Take  $k_0$  to be the smallest  $k$  in this decomposition. There are two cases:
- if  $k_0 = 1$ , then  $\mathfrak{g} = A_1 \subset M$  and  $[\mathfrak{g}\mathfrak{g}, \bigoplus_{k \geq 1} S^k] = \bigoplus_{k \geq 1} S^k \subset M$ , one has  $M = [W, W]$  by Proposition 3.4.
  - if  $k_0 > 1$ , then  $[S^3, A_{k_0}] \subset M$ . By (VII), we deduce that  $S^{2k_0-3}, S^{2k_0-2}, S^{2k_0+1}$  and  $S^{2k_0+2}$  are contained in  $M$ . But it results that  $A_{k_0-1} = S^{2k_0-3} \oplus S^{2k_0-2}$  is contained in  $M$ , a contradiction.
- (4) The proof of (4) is completely similar: one uses the decomposition of the representation  $(\text{ad}, W)$  into isotypic components,  $W = \bigoplus_{k \geq 0} B_k$  where  $B_k = S^{2k} \oplus S^{2k+1}$  (see Proposition 2.6).

□

#### 4. INVARIANT BILINEAR FORMS FOR THE ADJOINT AND TWISTED ADJOINT ACTIONS OF $W$

There is a bilinear form  $\kappa$  canonically associated to the supertrace on the Weyl algebra  $W$ , namely:

$$\kappa(F, G) := \text{Str}(F \star G), \quad \forall F, G \in W.$$

By Proposition 1.9,  $\kappa$  is supersymmetric:

$$\kappa(G, F) = (-1)^{fg} \kappa(F, G), \quad \forall F \in S^f, G \in S^g.$$

and from its very definition,  $\kappa$  is invariant under the adjoint representation:

$$\kappa(\text{ad}(F)(G), H) + (-1)^{fg} \kappa(G, \text{ad}(F)(H)) = 0, \quad \forall F \in S^f, G \in S^g.$$

Now,  $\kappa$  will be really interesting if it is non degenerate, and indeed this is the case:

**Theorem 4.1.** *The bilinear form  $\kappa$  on the Lie superalgebra  $W$  is non degenerate supersymmetric and ad-invariant. Moreover,*

$$\kappa(S^\ell, S^m) = \{0\}, \quad \text{if } \ell \neq m,$$

*and the restriction of  $\kappa$  to  $S^\ell$  is non degenerate. If  $\kappa'$  is another ad-invariant bilinear form on  $W$ , there exists  $\alpha \in \mathbb{K}$  such that  $\kappa' = \alpha \kappa$  on  $[W, W]$ .*

*Proof.* Given  $F \in S^\ell, G \in S^m$  with  $\ell < m$ , then  $\kappa(F, G) = \text{Str}(\sum_{k=0}^{\ell} C_k(F, G))$  by Proposition 1.4, with  $C_k(F, G) \in S^{\ell+m-2k}$  and since  $\ell + m - 2k > 0$ ,  $\kappa(F, G) = 0$ . Indeed,  $\kappa(F, G) = 0$  if  $\bar{\ell} + \bar{m} = \bar{1}$ .

To prove that  $\kappa$  is non degenerate, we have to show it on each component  $S^\ell$ . Using Proposition 1.4, if  $F, G \in S^\ell$ , then  $\kappa(F, G) = C_\ell(F, G)$ . Define  $\phi: S^\ell \rightarrow (S^\ell)^*$  as  $\phi(F)(G) = \kappa(F, G)$ , for all  $F, G \in S$ . Since  $\kappa$  is  $\mathfrak{g}_{\bar{0}}$ -invariant,  $\phi$  is homomorphism from the  $\mathfrak{g}_{\bar{0}}$ -module  $S^\ell$  into its contragredient module  $(S^\ell)^*$ . Both are simple  $\mathfrak{g}_{\bar{0}}$ -modules and  $\phi$  is non zero since  $C_\ell(S^\ell, S^\ell) = S^0 = \mathbb{K}$  by Theorem 3.1(4), so  $\phi$  is an isomorphism by Schur's Lemma and this proves that  $\kappa$  is non degenerate.

Assume now that  $\kappa'$  is an ad-invariant bilinear form. Then the map  $F \mapsto \kappa'|_F$  where  $\kappa'_F(G) = \kappa(F, G)$  is a homomorphism of  $\mathfrak{g}_{\bar{0}}$ -modules from  $S^\ell$  into  $(S^m)^*$ . When  $\ell \neq m$ ,  $S^\ell$  and  $(S^m)^* \cong S^m$  are not isomorphic, so  $\kappa'(S^\ell, S^m) = \{0\}$  by Schur's Lemma. If  $\ell = m$ ,  $S^\ell$  is a simple highest weight  $\mathfrak{g}_{\bar{0}}$ -module, so it is Schur irreducible, and it results that there is on  $S^\ell$  only one invariant bilinear form, up to a scalar, and consequently there exists  $\alpha_\ell \in \mathbb{K}$  such that  $\kappa'|_{S^\ell \times S^\ell} = \alpha_\ell \kappa|_{S^\ell \times S^\ell}$ . Next we want to prove that  $\alpha_\ell = \alpha_{\ell+1}$  for all  $\ell \geq 0$ . First we prove that  $\alpha_{2\ell-1} = \alpha_{2\ell}$  for all  $\ell \geq 1$ . Note that  $\text{ad}(p_1)(q_1^{2\ell}) = 2\ell q_1^{2\ell-1}$ , then from  $\kappa'(\text{ad}(p_1)(q_1^{2\ell}), F) = -\kappa'(q_1^{2\ell}, \text{ad}(p_1)(F))$ , for all  $F \in S^{2\ell-1}$ , we deduce that:

$$(\alpha_{2\ell-1} - \alpha_{2\ell})\kappa(\text{ad}(p_1)(q_1^{2\ell}), F) = 0,$$

and since  $\kappa$  is non degenerate on  $S^{2\ell-1} \times S^{2\ell-1}$ , one can conclude that  $\alpha_{2\ell-1} = \alpha_{2\ell}$  for all  $\ell \geq 1$ .

We then show that  $\alpha_{2\ell} = \alpha_{2\ell+1}$  for all  $\ell \geq 0$ . One has  $\kappa'(\text{ad}(p_1^3)(q_1^{2\ell}), F) = -\kappa'(q_1^{2\ell}, \text{ad}(p_1^3)(F))$ , for all  $F \in S^{2\ell+1}$  and that implies

$$(\alpha_{2\ell+1} - \alpha_{2\ell})\kappa(\text{ad}(p_1^3)(q_1^{2\ell}), F) = 0,$$

for all  $F \in S^{2\ell+1}$ . But  $\text{ad}(p_1^3)(q_1^{2\ell})$  has component  $6\ell p_1^2 q_1^{2\ell-1}$  on  $S^{2\ell+1}$  so  $\alpha_{2\ell} = \alpha_{2\ell+1}$  as wanted.

Now, starting from  $\ell = 1$ , we conclude that  $\kappa' = \alpha_1 \kappa$  on  $[W, W]$ .  $\square$

**Corollary 4.2.**  *$W$  and  $[W, W]$  are superquadratic Lie superalgebras.*

*Remark 4.3.* Consider the adjoint representation of  $\mathfrak{g}$  in  $W$ . By Proposition 2.5, it decomposes into isotypic components  $W = \bigoplus_{k \geq 0} A_k$ , with  $A_0 = \mathbb{K}$ ,  $A_k = S^{2k-1} \oplus S^{2k}$ ,  $k \geq 1$ . Each  $A_k$  has an explicit invariant supersymmetric bilinear form, namely  $\kappa|_{A_k \times A_k}$ , so the ad-representation of  $\mathfrak{g}$  in  $A_k$  is valued in

$$\mathfrak{osp}(S^{2k}, S^{2k-1}) = \mathfrak{osp}\left(\begin{pmatrix} 2n+2k-1 \\ 2k \end{pmatrix}, \begin{pmatrix} 2n+2k-2 \\ 2k-1 \end{pmatrix}\right).$$

Also the ad-representation of  $\mathfrak{g}_{\bar{0}}$  in  $S^k$  is orthogonal or symplectic, according to the parity of  $k$ :  $S^{2k}$  is orthogonal,  $S^{2k+1}$  is symplectic, and the corresponding bilinear form is explicitly computable using the Moyal  $\star$ -product.

What about the ad'-representation of  $W$ ? In what follows, we shall prove that it has also a non degenerate supersymmetric invariant bilinear form. Actually, this bilinear form extends the one defined on  $S^0 \oplus S^1$  when embedding  $\mathfrak{osp}(1, 2n)$  in  $W$  (see Proposition 2.4), so it is directly related to orthosymplectic supersymmetry.

**Theorem 4.4.** *Let  $B(F, G) = (-1)^{fg+1} \kappa(F, G)$ , for all  $F \in S^f$ ,  $G \in S^g$ . Then  $B$  is a non degenerate supersymmetric bilinear form on  $\mathbb{W}$ . Moreover,  $B(S^\ell, S^m) = \{0\}$ , if  $\ell \neq m$ ,  $B_{S^\ell \times S^\ell}$  is non degenerate and  $B$  is invariant under the  $\text{ad}'$ -representation of  $\mathbb{W}$ . If  $B'$  is an  $\text{ad}'$ -invariant bilinear form on  $\mathbb{W}$ , there exists  $\beta \in \mathbb{K}$  such that  $B' = \beta B$ .*

*Proof.* it is easy to check that  $B$  is supersymmetric. Let us prove that  $B$  is  $\text{ad}'$ -invariant: consider  $I = B(\text{ad}'(A)(F), G) + (-1)^{af} B(F, \text{ad}'(A)(G))$  with  $\deg_{\mathbb{Z}_2}(A) = a$ ,  $\deg_{\mathbb{Z}_2}(F) = f$  and  $\deg_{\mathbb{Z}_2}(G) = g$ . If  $f + g + a = \bar{1}$ , then  $I = 0$ . If  $f + g + a = \bar{0}$ , then

$$\begin{aligned}
I &= (-1)^{(f+a)g+1} \kappa(\text{ad}'(A)(F), G) + (-1)^{af} (-1)^{(f+a)g+1} \kappa(F, \text{ad}'(A)(G)) \\
&= (-1)^f \left( (-1)^{a+1} \kappa(\text{ad}'(A)(F), G) + (-1)^{af+1} \kappa(F, \text{ad}'(A)(G)) \right) \\
&= (-1)^f \text{Str} \left( (-1)^{a+1} (A \star F - (-1)^{a(f+1)} F \star A) \star G + \right. \\
&\quad \left. (-1)^{af+1} F \star (A \star G - (-1)^{a(g+1)} G \star A) \right) \\
&= (-1)^{f+a+1} \text{Str} \left( A \star F \star G - (-1)^{a(f+g)} F \star G \star A \right) \\
&= (-1)^{f+a+1} \text{Str}([A, F \star G]) = 0
\end{aligned}$$

Since  $\kappa$  is non degenerate,  $B$  is non degenerate. Finally, in order to prove the uniqueness of two  $\text{ad}'$ -invariant bilinear forms modulo  $\mathbb{K}$ , one proceeds as in the proof of Theorem 4.1, so we leave out the proof.  $\square$

## 5. RENORMALIZED SUPERTRACE AND FORMAL INVERSE WEYL TRANSFORM

In this Section, we assume that  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ . Let  $\mathcal{P}$  be the algebra  $\mathbb{K}[x_1, \dots, x_n]$ ,  $\mathcal{F}$  be the algebra  $\mathbb{K}[[X_1, \dots, X_n]]$ ,  $V$  be the space  $V = \text{span}\{X_1, \dots, X_n\}$  and  $V^*$  be its dual,  $V^* = \text{span}\{x_1, \dots, x_n\}$  with  $\langle x_i, X_j \rangle = \delta_{ij}$ .

There is a one to one mapping (the Laplace transform) from  $\mathcal{P}^*$  onto  $\mathcal{F}$  defined by the duality  $\langle x^I, X^J \rangle = \delta_{IJ} I!$ , where  $x^I := x_1^{i_1} \dots x_n^{i_n}$ ,  $X^J := X_1^{j_1} \dots X_n^{j_n}$ ,  $I! := i_1! \dots i_n!$ . So the spaces  $\mathcal{P}^*$  and  $\mathcal{F}$  can be identified and as a consequence, the Dirac distributions  $\partial_v$ ,  $v \in V$ ,  $\partial_v(P) = P(v)$  become formal power series  $e^v$  so that:

$$P(v) = \langle P | e^v \rangle, \forall P \in \mathcal{P}, v \in V \text{ (Taylor's Formula).}$$

Also one has :

$$\left\langle \frac{\partial^I P}{\partial x^I} \middle| F \right\rangle = \langle P | X^I F \rangle, \forall P \in \mathcal{P}, F \in \mathcal{F}$$

and that means  ${}^t \left( \frac{\partial^I}{\partial x^I} \right)$  is the multiplication by  $X^I$  in  $\mathcal{F}$ .

The algebra  $\mathcal{P}$  has a Hopf algebra structure with coproduct  $\Delta(P) := P(x + x')$  if one identifies  $\mathcal{P} \otimes \mathcal{P} = \mathbb{K}[x_1, \dots, x_n, x'_1, \dots, x'_n]$ , and antipode  $\mathcal{S}(P)(x) = P(-x)$ .

Next we endow  $\mathcal{P}$  with its natural topology, as defined in [4] and  $\mathcal{F} = \mathcal{P}^*$  with the strong dual topology which is exactly the product topology  $\prod_{k \geq 0} \mathcal{F}^k$ , where  $\mathcal{F}^k$  denotes the set of homogeneous polynomials of degree  $k$ . Then the transposition map induces on  $\mathcal{F}$  a topological Hopf algebra structure, which is exactly the

usual structure (see [4]) with the identification  $\mathcal{F} \widehat{\otimes} \mathcal{F} = \mathbb{K}[[X_1, \dots, X_n, X'_1, \dots, X'_n]]$  (where  $\widehat{\otimes}$  is the projective tensor product, see [19]).

Any linear operator  $T: \mathcal{P} \rightarrow \mathcal{P}$  is continuous for the natural topology ([4]) and  ${}^tT: \mathcal{F} \rightarrow \mathcal{F}$  is continuous. Denote by  $\mathcal{L}(\mathcal{P})$  the space of linear operators of  $\mathcal{P}$ , and by  $\mathcal{L}(\mathcal{F})$  the space of continuous linear operators of  $\mathcal{F}$ . Then one has:

$$\mathcal{L}(\mathcal{P}) = \mathcal{P}^* \widehat{\otimes} \mathcal{P} = \mathcal{P}^* \widehat{\otimes} \mathcal{P}^{**} = \mathcal{L}(\mathcal{F}) \quad (\text{see [19]}).$$

Let us quickly explain how it works: given  $T \in \mathcal{L}(\mathcal{P})$ , then  $T = \sum_K \frac{1}{K!} P_K \otimes X^K$  with  $P_K = T(x^K)$  and one has  $T(P) = \sum_K \frac{1}{K!} P_K \langle X^K | P \rangle$  for all  $P \in \mathcal{P}$ . Now one has  ${}^tT = \sum_K \frac{1}{K!} P_K \otimes X^K$  and

$$(VIII) \quad {}^tT(F) = \sum_K \frac{1}{K!} \langle P_K | F \rangle X^K \text{ for all } F \in \mathcal{F}.$$

Given a polynomial function  $d: V \rightarrow \mathcal{F}$  defined as  $d(v) = \sum_K D_K(v) X^K$ ,  $D_K \in \mathcal{P}$ , there is an associated operator  $D: \mathcal{P} \rightarrow \mathcal{P}$  defined by

$$D = \sum_K D_K(v) \frac{\partial^K}{\partial x^K}$$

and one has:

$$D(P)(v) = \langle P | d(v) e^v \rangle, \quad \forall P \in \mathcal{P}, v \in V.$$

Since  $\text{span}\{e^v | v \in V\}$  is dense in  $\mathcal{F} = \mathcal{P}^*$  by the Hahn-Banach Theorem, we deduce:

$$(IX) \quad \langle D(P) | F \rangle = \langle P | m \circ (d \otimes \text{Id}) \circ \Delta(F) \rangle,$$

noticing that  $\Delta(e^v) = e^v \otimes e^v$  and extending  $d$  to  $\mathcal{F}$  by  $d(F) = \sum_K \langle D_K | F \rangle X^K$ ,  $F \in \mathcal{F}$  (so  $d(e^v) = d(v)$ ).

The operator  $D$  is what we shall call a *differential operator* of  $\mathcal{P}$ . We shall say that  $D$  is *differential operator for finite order* if  $d(V) \subset \mathcal{F}_k = \{F \in \mathcal{F} | \text{deg}(F) \leq k\}$ .

A fundamental property of  $\mathcal{P}$  is established by:

**Lemma 5.1.** ([14]) *Any linear operator of  $\mathcal{P}$  is a differential operator.*

*Proof.* Given  $T \in \mathcal{L}(\mathcal{P})$ , then  ${}^tT \in \mathcal{L}(\mathcal{F})$ , so we have to find a polynomial map from  $V$  to  $\mathcal{F}$  satisfying  ${}^tT = m \circ (d \otimes \text{Id}) \circ \Delta$  (due to (IX)) From the density of  $\text{span}\{e^v | v \in V\}$ , it is enough to prove the last identity on this set. Let  $d = \sum_k D_k X^k$ ,  $D_k \in \mathcal{P}$ , then  ${}^tT(e^v) = (m \circ (d \otimes \text{Id}) \circ \Delta)(e^v) = \sum_K \langle D_K | e^v \rangle X^K e^v$  gives  $\sum_K \langle D_K | e^v \rangle X^K = {}^tT(e^v) e^{-v}$ , hence  $d(v) = {}^tT(e^v) e^{-v}$ .  $\square$

Let us give an explicit formula: starting from  $T \in \mathcal{L}(\mathcal{P})$ ,  $T = \sum_I \frac{1}{I!} P_I \otimes X^I$  with  $P_I \in \mathcal{P}$ . Then  ${}^tT(e^v) = \sum_I \frac{1}{I!} P_I(v) X^I$  by (VIII) so if  $v = x_1 X_1 + \dots + x_n X_n$  and

$$\begin{aligned}
|I| &= i_1 + \cdots + i_n, \\
{}^i T(e^v) e^{-v} &= \\
&= \left( \sum_I \frac{1}{I!} P_I(v) X^I \right) \left( \sum_j \frac{(-1)^j v^j}{j!} \right) \\
&= \left( \sum_I \frac{1}{I!} P_I(v) X^I \right) \left( \sum_j \frac{(-1)^j}{j!} \sum_{i_1 + \cdots + i_n = j} \frac{(i_1 + \cdots + i_n)!}{i_1! \cdots i_n!} x_1^{i_1} \cdots x_n^{i_n} X_1^{i_1} \cdots X_n^{i_n} \right) \\
&= \left( \sum_I \frac{1}{I!} P_I(v) X^I \right) \left( \sum_I \frac{(-1)^{|I|}}{I!} x^I X^I \right) \\
&= \sum_N \frac{1}{N!} \left( \sum_{R+S=N} (-1)^{|S|} \frac{N!}{R!S!} P_R(v) x^S \right) X^N.
\end{aligned}$$

Finally

$$T = \sum_N \frac{1}{N!} \left( \sum_{R+S=N} (-1)^{|S|} \frac{N!}{R!S!} P_R x^S \right) \frac{\partial^N}{\partial x^N}.$$

Recall that  $P_R = T(x^R)$ , therefore:

$$T = \sum_N \frac{1}{N!} \left( \sum_{R+S=N} (-1)^{|S|} \frac{N!}{R!S!} T(x^R) x^S \right) \frac{\partial^N}{\partial x^N}.$$

But  $\Delta(x^N) = \sum_{R+S=N} \frac{N!}{R!S!} x^R \otimes x^S$ , so this last formula can be written:

**Theorem 5.2.**

$$T = \sum_N \frac{1}{N!} (m \circ (T \otimes \mathcal{S}) \circ \Delta(x^N)) \frac{\partial^N}{\partial x^N}.$$

where  $\mathcal{S}$  is the antipode of  $\mathcal{P}$ .

For instance, when  $n = 1$ , consider the operator  $T$  defined as  $T(x^i) = x^j$  for fixed  $i$  and  $j$ , and 0 otherwise. Then;

$$T = \frac{x^j}{i!} \sum_{\ell \geq 0} (-1)^\ell \frac{x^\ell}{\ell!} \frac{d^{i+\ell}}{dx^{i+\ell}}.$$

Using the notation  $P^I = p_1^{i_1} \cdots p_n^{i_n}$  and  $Q^J = q_1^{j_1} \cdots q_n^{j_n}$ , consider the (formal) completion  $\overline{W} = \mathbb{K}[Q][[P]]$  of the  $\star$ -algebra  $W$  ( $\overline{W}$  is an algebra with the Moyal  $\star$ -product). Elements of  $\overline{W}$  are formal power series  $\tilde{F} = \sum_I \alpha_I(Q) \star P^I$  with  $\alpha_i \in \mathbb{K}[Q]$ . Define a map  $\mathcal{W} : \overline{W} \rightarrow \mathcal{L}(\mathcal{P})$  as:

$$\mathcal{W}(\tilde{F}) = \sum_I \alpha_i(x) \frac{\partial^I}{\partial x^I}.$$

Remark that  $\mathcal{W}$  is simply the extension to  $\overline{W}$  of the natural  $W$ -module structure of  $\mathcal{P}$  defined by  $p_i \cdot P = \frac{\partial P}{\partial x_i}$ ,  $q_i \cdot P = x_i P$ ,  $i = 1, \dots, n$ , hence  $\mathcal{W}$  is an algebra homomorphism.

Lemma 5.1 can now be reinterpreted as:

**Proposition 5.3.** *The map  $\mathscr{W}$  is an isomorphism of algebras.*

One should be careful that the domain of  $\mathscr{W}$ , i.e.  $\overline{W}$ , is not at all identical to the formal completion  $\overline{S} = \mathbb{K}[Q][[P]]$  endowed with an Abelian product and with elements  $\widehat{F} = \sum_I \alpha_i(Q) P^I$ .

Given  $\widetilde{F} = \sum_I \alpha_i(Q) \star P^I \in \overline{W}$ , we can try to define its supertrace using the supertrace of  $W$  that we will denote by  $\text{Str}_W$ . A natural candidate would be  $\text{Str}_{\overline{W}}(\widetilde{F}) := \sum_I \text{Str}_W(\alpha_i(Q) \star P^I)$  but it is clear that this series happens to diverge. So  $\text{Str}_{\overline{W}}$  has a domain denoted by  $\text{Dom}(\text{Str}_{\overline{W}})$ . Evidently,  $W \subset \text{Dom}(\text{Str}_{\overline{W}})$ . On the other hand,  $\mathcal{L}(\mathcal{P})$  is a  $\mathbb{Z}_2$ -graded algebra since  $\mathcal{P}$  is  $\mathbb{Z}_2$ -graded, so we can define a supertrace at least on the ideal  $\mathcal{L}_f(\mathcal{P})$  of finite rank operators. Note that  $\mathcal{L}_f(\mathcal{P}) \cap W = \{0\}$  since  $W$  is a simple algebra. So we now have two supertraces, living apparently on different domains, and we wish to compare these supertraces. This is done by the following Theorem:

**Theorem 5.4.**

*One has  $\mathcal{L}_f(\mathcal{P}) \subset \text{Dom}(\text{Str}_{\overline{W}})$  and if  $\widetilde{F} \in \mathcal{L}_f(\mathcal{P})$ ,*

$$\text{Str}(\widetilde{F}) = \frac{1}{2^n} \text{Str}_{\overline{W}}(\widetilde{F}).$$

To prove Theorem 5.4, we need precise formulas for  $\text{Str}_W$ :

**Proposition 5.5.**

(1) *Using the natural isomorphism  $W(n) = W(1) \otimes \cdots \otimes W(1)$ , one has*

$$\text{Str}_W(F_1 \otimes \cdots \otimes F_n) = \text{Str}_W(F_1) \cdots \text{Str}_W(F_n).$$

(2) *One has  $\text{Str}_W(P^I \star Q^J) = \delta_{IJ} \frac{I!}{2^{|I|}}$ .*

*Proof.*

(1) Recall that the isomorphism  $W(n) = W(1) \otimes \cdots \otimes W(1)$  is defined by  $F_1 \otimes \cdots \otimes F_n = F_1 \dots F_n$ . Then

$$\begin{aligned} \text{Str}_W(F_1 \otimes \cdots \otimes F_n) &= \text{Str}_W(F_1 \dots F_n) = (F_1 \dots F_n)(0) = F_1(0) \dots F_n(0) \\ &= \text{Str}_W(F_1) \cdots \text{Str}_W(F_n). \end{aligned}$$

(2) We start with the case  $n = 1$ . Then  $\text{Str}_W(p_1^{i_1} \star q_1^{j_1}) = \kappa(p_1^{i_1}, q_1^{j_1}) = 0$  if  $i_1 \neq j_1$  by Proposition (4.1). Furthermore,  $\text{Str}_W(p_1^{i_1} \star q_1^{i_1}) = C_{i_1}(p_1^{i_1}, q_1^{i_1}) = \frac{i_1!}{2^{i_1}}$ .

Compute now:

$$\begin{aligned} \text{Str}_W(p_1^{i_1} \dots p_n^{i_n} \star q_1^{j_1} \dots q_n^{j_n}) &= \text{Str}_W(p_1^{i_1} \star q_1^{j_1} \otimes p_2^{i_2} \star q_2^{j_2} \otimes \cdots \otimes p_n^{i_n} \star q_n^{j_n}) \\ &= \text{Str}_W(p_1^{i_1} \star q_1^{j_1}) \cdots \text{Str}_W(p_n^{i_n} \star q_n^{j_n}) = 0, \end{aligned}$$

$$\text{if } I \neq J, \text{ and } \frac{I!}{2^{|I|}} \text{ if } I = J.$$

□

As a practical case of Theorem 5.4, we prove:

**Proposition 5.6.** *Let  $T$  be the (finite rank) operator defined as  $T(x^I) = x^J$  for fixed  $I, J$  and 0 otherwise. Then*

$$T = \frac{x^J}{I!} \sum_S (-1)^S \frac{x^S}{S!} \frac{\partial^{I+S}}{\partial x^{I+S}},$$

and

$$\text{Str}_{\overline{\mathbb{W}}}(T) = \begin{cases} 2^n \text{Str}(T) = 2^n (-1)^{|I|}, & \text{if } I = J \\ 0 & \text{otherwise} \end{cases}.$$

Thus  $T \in \text{Dom}(\text{Str}_{\overline{\mathbb{W}}})$  and its usual supertrace is, up to a factor  $\frac{1}{2^n}$ , its  $\overline{\mathbb{W}}$ -supertrace.

*Proof.* The formula for  $T$  is obtained by applying Theorem 5.2:

$$T = \sum_S (-1)^{|S|} \frac{Q^{J+S}}{I!S!} \star P^{I+S}.$$

Then  $\text{Str}_{\overline{\mathbb{W}}}(T) = \sum_S \frac{(-1)^{|S|}}{I!S!} \text{Str}_{\mathbb{W}}(Q^{J+S} \star P^{I+S}) = 0$ , if  $J \neq I$ , by Proposition

5.5. When  $I = J$ ,  $\text{Str}_{\overline{\mathbb{W}}}(Q^{I+S} \star P^{I+S}) = (-1)^{|I+S|} \frac{(I+S)!}{2^{|I+S|}}$ , so

$$\text{Str}_{\overline{\mathbb{W}}}(T) = \frac{(-1)^{|I|}}{2^{|I|}} \sum_S \frac{(I+S)!}{I!S!} \frac{1}{2^{|S|}}.$$

If  $|\tau| < 1$ , one has  $\frac{1}{(1-\tau)^{i+1}} = \sum_s \binom{i+s}{s} \tau^s$ , therefore:

$$\frac{1}{(1-\tau_1)^{i_1+1} \dots (1-\tau_n)^{i_n+1}} = \sum_{s_1, \dots, s_n} \binom{i_1+s_1}{s_1} \dots \binom{i_n+s_n}{s_n} \tau_1^{s_1} \dots \tau_n^{s_n},$$

if  $|\tau_i| < 1$ , for all  $i$ . It results that:

$$\sum_S \frac{(I+S)!}{I!S!} \frac{1}{2^{|S|}} = \frac{1}{(1-\frac{1}{2})^{i_1+1} \dots (1-\frac{1}{2})^{i_n+1}} = 2^{|I|+n}.$$

At last, we obtain

$$\text{Str}_{\overline{\mathbb{W}}}(T) = (-1)^{|I|} 2^n = \text{Str}(T) 2^n.$$

□

*Proof.* (Proof of Theorem 5.4)

This proof is completely similar to the previous one and for this reason, we omit it (note that one can restrict to  $T = \varphi \otimes x^k$ ,  $\varphi \in \mathcal{P}^*$  since  $\mathcal{L}_f(\mathcal{P}) = \mathcal{P}^* \otimes \mathcal{P}$ ). □

*Remark 5.7.* Let us give some more interpretation about supertraces. We want to show how one can define a renormalized supertrace from Theorem 5.4. We begin with  $\mathcal{L}_f(\mathcal{P})$  and its natural supertrace  $\text{Str}$ . Notice that this supertrace is of intrinsic nature, since defined by  $\text{Str}(\varphi \otimes P) = (-1)^{\deg_{\mathbb{Z}_2}(\varphi) \deg_{\mathbb{Z}_2}(P)} \langle \varphi | P \rangle$ ,  $\varphi \in \mathcal{P}^*$ ,  $P \in \mathcal{P}$ .

Secondly, we have the supertrace  $\text{Str}_{\overline{W}}$  defined on its domain  $\text{Dom}(\text{Str}_{\overline{W}})$  which contains  $W$  and  $\mathcal{L}_f(\mathcal{P})$ . On  $\mathcal{L}_f(\mathcal{P})$ , one has

$$\text{Str}(T) = \frac{1}{2^n} \text{Str}_{\overline{W}}(T)$$

by Theorem 5.4. So we can extend  $\text{Str}$  to  $\text{Dom}(\text{Str}_{\overline{W}})$  and define a *renormalized supertrace*, denoted by  $\text{RStr}$ , as:

$$\text{RStr}(T) = \frac{1}{2^n} \text{Str}_{\overline{W}}(T).$$

This extension is indeed a renormalized extension of  $\text{Str}$ : for instance, with the usual definition of the supertrace:  $\text{Str}(\text{Id}) = \infty - \infty$ , a rather bad result, but with the renormalization:

$$\text{RStr}(\text{Id}) = \frac{1}{2^n}.$$

Notice that  $n = \dim(V)$  is the dimension of the underlying variety.

Next, we will clarify what we mean by a *formal inverse Weyl transform*. Recall that we can identify  $\overline{W} = \mathcal{L}(\mathcal{P})$  thanks to Proposition 5.3. So, given  $T \in \mathcal{L}(\mathcal{P})$ , one can write:

$$T = \sum_I \alpha_I(Q) \star P^I \in \overline{W}.$$

Consider the Abelian algebra  $\mathbb{K}[[P, Q]]$ , and denote by  $\mathbb{K}[[P, Q]]^k$  the space of homogeneous polynomials of degree  $k$ . We endow  $\mathbb{K}[[P, Q]]$  with the product topology  $\mathbb{K}[[P, Q]] = \prod_{k \geq 0} \mathbb{K}[[P, Q]]^k$  which is Fréchet. Now  $\sum_I \alpha_I(Q) \star P^I \in \overline{W}$  is a series (of polynomials) in  $\mathbb{K}[[P, Q]]$ , it converges in  $\mathbb{K}[[P, Q]]$  if and only if for any  $k$ , the series obtained by taking the  $k$ th components converge in  $\mathbb{K}[[P, Q]]^k$ .

**Definition 5.8.** When  $\sum_I \alpha_I(Q) \star P^I$  converges in  $\mathbb{K}[[P, Q]]$ , we say that  $T$  has a *formal inverse Weyl transform* denoted by  $\text{IW}(T)$  and defined as the sum of the series.

Remark that taking  $k = 0$ , we obtain that if  $T$  has a formal inverse Weyl transform, the series

$$\sum_I (\alpha_I(Q) \star P^I)(0) = \sum_I \text{Str}_W(\alpha_I(Q) \star P^I) = \text{Str}_{\overline{W}}(T)$$

must converge, so:

**Proposition 5.9.** *The existence of  $\text{RStr}(T)$  is a necessary condition for the existence of the formal inverse Weyl transform of  $T$ . When  $\text{IW}(T)$  exists, one has:*

$$\text{RStr}(T) = \frac{1}{2^n} \text{IW}(T)(0).$$

As an operator from  $\mathcal{L}(\mathcal{P})$  into  $\mathbb{K}[[P, Q]]$ , the formal inverse Weyl transform has a domain  $\mathcal{D} \subsetneq \mathcal{L}(\mathcal{P})$  containing  $W$  and naturally, it would be nice to have a characterization of  $\mathcal{D}$ . This will be done elsewhere, rather let us develop here some examples. We take  $n = 1$  and consider the elementary operators  $E_{ij}$  of  $\mathcal{P}$  defined by  $E_{ij}(x^k) = \delta_{jk}x^i$ . One has  $E_{ij} = \frac{1}{j!} \sum_{\ell \geq 0} \frac{(-1)^\ell}{\ell!} q^{\ell+i} \star p^{\ell+j}$  and some computation shows that

**Lemma 5.10.** *The formal inverse Weyl transform of  $E_{ij}$  is*

$$\text{IW}(E_{ij}) = \begin{cases} (-1)^j 2^{i-j+1} L_j^{(i-j)}(4pq) e^{-2pq} q^{i-j}, & \text{if } j \leq i \\ (-1)^i 2^{j-i+1} \frac{i!}{j!} L_i^{(j-i)}(4pq) e^{-2pq} p^{j-i}, & \text{if } j \geq i \end{cases}.$$

where  $L_\beta^{(\alpha)}$  is the Laguerre polynomial (see [18]).

Consider now the operators  $S_\lambda$  defined as  $S_\lambda(x^k) = \lambda^k x^k$ ,  $\lambda \in \mathbb{K}$ . Then by Theorem 5.2, one has:

$$S_\lambda = \sum_{\ell \geq 0} \frac{(-1)^\ell}{\ell!} (1 - \lambda)^\ell q^\ell \star p^\ell.$$

By (V) in Section 1,  $q^\ell \star p^\ell = (-1)^\ell \frac{\ell!}{2^\ell} L_\ell(2pq)$ , so using the generating function of Laguerre polynomials  $L_\beta^{(0)} := L_\beta$ , one finds:

**Lemma 5.11.** *The formal inverse Weyl transform of  $S_\lambda$  is*

$$\text{IW}(S_\lambda) = \frac{2}{1 + \lambda} \exp\left(2 \frac{\lambda - 1}{\lambda + 1} pq\right), \text{ if } |1 - \lambda| < 2.$$

Notice that  $S_\lambda$  has a renormalized supertrace given by:

$$\text{RStr}(S_\lambda) = \frac{1}{\lambda + 1}.$$

When  $|\lambda| < 1$ , this is exactly  $\text{Str}(S_\lambda)$ .

Here are some cases of interest:

First, take  $\lambda = e^\tau$ , then  $S_\lambda = \exp\left(\tau x \frac{d}{dx}\right) = \exp(\tau q \star p)$  and one finds that its formal inverse Weyl transform is:

$$\text{IW}\left(\exp\left(\tau x \frac{d}{dx}\right)\right) = \frac{e^{-\frac{\tau}{2}}}{\cosh\left(\frac{\tau}{2}\right)} \exp\left(2pq \tanh\left(\frac{\tau}{2}\right)\right), \text{ if } |1 - e^\tau| < 2.$$

Take  $\tau = i\theta$ ,  $\theta \in \mathbb{R}$  with  $\theta \neq (2s + 1)\pi$ ,  $s \in \mathbb{Z}$ . Then  $S_{e^{i\theta}} = \exp(i\theta q \star p)$  has formal inverse Weyl transform:

$$\text{IW}\left(\exp\left(i\theta x \frac{d}{dx}\right)\right) = \frac{e^{-i\frac{\theta}{2}}}{\cos\left(\frac{\theta}{2}\right)} \exp\left(2ipq \tan\left(\frac{\theta}{2}\right)\right),$$

and an interesting case is  $S_i$  with formal inverse Weyl transform  $(1 - i) \exp(2ipq)$  (see [3], where similar formulas are obtained, in the context of quantization of the harmonic oscillator). Note that the value  $\lambda = -1$  is critical: indeed  $S_{-1} = P$  is the parity operator, which has divergent renormalized supertrace and cannot have a formal inverse Weyl transform. So the estimation in (5.11) is the best one.

Let us now justify our construction somewhat. There is a natural question : quantum mechanics is built from operators in a Hilbert space, so where is the Hilbert space in that picture? Here is the answer: let  $\mathcal{H}$  be the Hilbert space of entire functions  $f(x)$ ,  $x = ai + b$  such that  $\int e^{-|x|^2} |f(x)|^2 da db < \infty$  with scalar product  $\langle f | g \rangle = \frac{1}{\pi} \int e^{-|x|^2} f(x) \overline{g(x)} da db$ . Then  $\mathcal{P} \subset \mathcal{H}$ , as a dense subspace and we can apply our algebraic formalism as follows: given  $T$  an operator of  $\mathcal{H}$ ,

whose domain contains  $\mathcal{P}$ , assume that  $T(\mathcal{P}) \subset \mathcal{P}$  and denote by  $T_{\mathcal{P}}$  the restriction to  $\mathcal{P}$ ; to have one to one  $T \mapsto T_{\mathcal{P}}$ , we have to assume more, for instance either  $T$  bounded, or  $T_{\mathcal{P}}$  essentially self-adjoint, this is assumed in the foregoing. We can now define the formal inverse Weyl transform of  $T$  to be the formal inverse Weyl transform of  $T_{\mathcal{P}}$  and the renormalized supertrace of  $T$  to be the renormalized supertrace of  $T_{\mathcal{P}}$ . For instance, consider  $S_i$  which is a unitary operator of  $\mathcal{H}$ , it has renormalized supertrace  $\text{RStr}(S_i) = \frac{1}{2}(1-i)$  and formal inverse Weyl transform  $(1-i)\exp(2ipq)$ . We are actually working in the so called ‘‘coherent states formalism’’, and we can easily translate in terms of the usual ‘‘metaplectic formalism’’. Define an operator  $H: \mathcal{H} \rightarrow \mathcal{L}^2(\mathbb{R})$  by

$$H(f)_t = \frac{e^{-\frac{t^2}{2}}}{\pi^{\frac{3}{4}}} \int f(\sqrt{2}(t+i\xi)) e^{-\xi^2} d\xi,$$

let  $\phi_n$  be the orthonormal basis of Hermite functions in  $\mathcal{L}^2(\mathbb{R})$ , and  $Z_n = \frac{x^n}{\sqrt{n!}}$  the orthonormal basis of  $\mathcal{H}$ , then one has  $H(Z_n) = \phi_n, \forall n$ , so  $H$  is a unitary isomorphism from  $\mathcal{H}$  to  $\mathcal{L}^2(\mathbb{R})$ . The operator  $H$  maps the operator  $x$  (resp.  $\frac{d}{dx}$ ) or  $\mathcal{H}$  onto the operator  $\frac{1}{\sqrt{2}}(t - \frac{d}{dt})$  (resp.  $\frac{1}{\sqrt{2}}(t + \frac{d}{dt})$ ) and we recover the usual metaplectic formalism (see [11] for details). now we can explain our interest for  $S_i$ : indeed  $H$  maps  $S_i$  onto the Fourier transform  $\mathfrak{F}$  of  $\mathcal{L}^2(\mathbb{R})$  and since ‘‘coherent states formalism’’ and ‘‘metaplectic formalism’’ are completely equivalent by  $H$ , we have a formal inverse Weyl transform of  $\mathfrak{F}$ , namely the formal inverse Weyl transform of  $S_i$ , i.e.

$$\text{IW}(\mathfrak{F}) = (1-i)\exp(2ipq)$$

(attention to the interpretation of  $p$  and  $q$  in terms of  $t$ ), and also a renormalized supertrace

$$\text{RStr}(\mathfrak{F}) = \frac{1}{2}(1-i).$$

So we think that our formalism might be of interest. Next step will be a Wigner’s type formula: this is already done and will be explained in a subsequent paper.

#### REFERENCES

- [1] Arnal, D.; Benamor, H.; Benayadi, S.; Pinczon, G., Une algèbre de Lie non semi-simple rigide et sympathique:  $H^1(\mathfrak{g}) = H^2(\mathfrak{g}) = H^0(\mathfrak{g}, \mathfrak{g}) = H^1(\mathfrak{g}, \mathfrak{g}) = H^2(\mathfrak{g}, \mathfrak{g}) = \{0\}$ , *C. R. Acad. Sci.*, Paris, Sér. I **315**, No.3, 261-263 (1992).
- [2] Arnaudon, D.; Bauer, M.; Frappat, L., On Casimir’s ghost, *Commun. Math. Phys.*, **187**, No.2, 429-439 (1997).
- [3] Bayen, F.; Flato, M.; Frønsdal, C.; Lichnerowicz, A.; Sternheimer, D., Deformation theory and quantization. I,II. Deformations of symplectic structures, Physical applications, *Ann. Phys.*, **111**, 61-110, 111-151 (1978).
- [4] Bonneau, P.; Flato, M.; Gerstenhaber, M.; Pinczon, G., The hidden group structure of quantum groups: Strong duality, rigidity and preferred deformations, *Commun. Math. Phys.*, **161**, No.1, 125-156 (1994).
- [5] Connes, A.; Flato, M.; Sternheimer, D., Closed star products and cyclic cohomology, *Lett. Math. Phys.*, **24**, No.1, 1-12 (1992).
- [6] Dixmier, J., Sur les algèbres de Weyl, *Bull. Soc. Math. Fr.* 96, 209-242 (1968). Sur les algèbres de Weyl II, *Bull. Sci. Math.*, Sér. **94**, 289-301 (1970).

- [7] Fronsdal, C.; Flato, M.; Hirai, T. (Eds.), Essays on supersymmetry, *Mathematical Physics Studies*, Vol. **8**, D. Reidel Publishing Co., Dordrecht, (1986).
- [8] Gié, P-A.; Pinczon, G.; Ushirobira, R., Back to the Amitsur–Levitzki theorem: a super version for the orthosymplectic Lie superalgebra  $\mathfrak{osp}(1, 2n)$ , *Lett. Math. Phys.*, **66**, No.1-2, 141-155 (2003).
- [9] Gorelik, M., On the ghost centre of Lie superalgebras, *Ann. Inst. Fourier* **50**, No.6, 1745-1764 (2000).
- [10] Kac, V. G., Infinite dimensional Lie algebras, *Cambridge University Press.*, (1990).
- [11] Lesimple, M.; Pinczon, G., Deformations of the metaplectic representations, *J. Math. Phys.* **42**, no. 4, 1887–1899 (2001).
- [12] Montgomery, S., Constructing simple Lie superalgebras from associative graded algebras, *J. Algebra*, **195**, No.2, 558-579 (1997).
- [13] Musson, I. M., Some Lie superalgebras associated to the Weyl algebras, *Proc. Am. Math. Soc.* **127**, No.10, 2821-2827 (1999).
- [14] Pinczon, G., On the equivalence between continuous and differential deformation theories, *Lett. Math. Phys.*, **39**, No.2, 143-156 (1997).
- [15] Pinczon, G., The enveloping algebra of the Lie superalgebra  $\mathfrak{osp}(1, 2)$ , *J. Algebra*, **132**, no. 1, 219–242 (1990).
- [16] Pinczon, G.; Simon, J., Non-linear representations of inhomogeneous groups, *Lett. Math. Phys.* **2**, 499-504 (1978).
- [17] Sternheimer, D., Deformation quantization: Twenty years after, Rembielinski, J. (ed.), *Particles, fields, and gravitation*, Papers from the conference dedicated to the memory of Ryszard Raczka, Łódź, Poland, April 15-19, 1998. Woodbury, NY: American Institute of Physics. AIP Conf. Proc. **453**, 107-145 (1998).  
Dito, G.; Sternheimer, D., Deformation quantization: genesis, developments and metamorphoses, Deformation quantization (Strasbourg, 2001), 9–54, *IRMA Lect. Math. Theor. Phys.*, **1**, de Gruyter, Berlin (2002).
- [18] Szegő, G., Orthogonal Polynomials, *American Mathematical Society Colloquium Publications*, v. **23**. American Mathematical Society, New York, (1939).
- [19] Trèves, F., Topological vector spaces, distributions and kernels, *New York-London: Academic Press* (1967).

INSTITUT DE MATHÉMATIQUES DE BOURGOGNE, UNIVERSITÉ DE BOURGOGNE, B.P. 47870,  
F-21078 DIJON CEDEX, FRANCE

*E-mail address:* gpinczon, rosane@u-bourgogne.fr