SUBALGEBRAS AND FREE PRODUCT STRUCTURES OF A GRAPH W^* -PROBABILITY SPACE

ILWOO CHO

ABSTRACT. Let G be a countable directed graph. Then we can construct the graph W^{*}-algebra $W^*(G)$ and its diagonal subalgebra D_G . By defining the conditional expectation $E: W^*(G) \to D_G$, we have the grarph W^{*}-probability space over D_G , $(W^*(G), E)$, as an amalgamated W^* -probability space over D_G . The amalgamated freeness on $(W^*(G), E)$ is defined in the sense of Speicher. In this papaer, we will define the D_G -semicircular system and D_G valued R-diagonal system. If the graph G contains N-mutually diagram**distinct** loops, then we can construct the D_G -semicircular system L_N in $(W^*(G), E)$ and the D_G -semicircular algebra $W^*(L_N, D_G)$ generated by L_N and D_G , characterized by $(W^*(L_N, D_N), E_N) \otimes (D_G, \mathbf{1})$, where $D_N \leq D_G$ is a subalgebra determined by L_N . If we have N-mutually distinct finite paths $w_1, ..., w_N \notin loop(G)$, then we can construct the D_G -valued R-diagonal system $R = \{L_{w_1}, L_{w_1}^*, ..., L_{w_N}, L_{w_N}^*\}$ satisfying that $\{L_{w_1}, L_{w_1}^*\}, ..., \{L_{w_N}, L_{w_N}^*\}$ are D_G -free from each other. The subalgebra $W^*(R, D_G)$ is observed. Precisely, we show that the graph W^* -algebra $W^*(G)$ is the D_G -free product of D_G -free building blocks D_G and $W^*(\{L_e\}, D_G)$, for all $e \in E(G)$. So, we can observe the D_G -free product structure of $(W^*(G), E)$ by these building blocks. Also, we can see that

 $(W^*(G), E) = (D_G, \mathbf{1}) *_{D_G} (*_{D_G} (W^*(\{L_e\}, D_G), E)) \\ \underset{e \in E(G)}{\overset{\bullet}{=}} (W^*(\{L_e\}, D_G), E))$

In [16], we constructed the graph W^* -probability spaces. The graph W^* -probability theory is one of the good example of Speicher's combinatorial free probability theory with amalgamation. In [16], we observed how to compute the certain operatorvalued moments and cumulants of an arbitrary operator-valued random variables in the graph W^* -probability space and observed the amalgamated freeness on the graph W^* -probability space, with respect to the given conditional expectation. Also, in [17], we consider certain operator-valued random variables of the graph W^* -probability space, for example, semicircular elements, even elements and Rdiagonal elements. This shows that the graph W^* -probability spaces contain the rich free probabilistic objects.

Throughout this paper, let G be a countable directed graph and let $\mathbb{F}^+(G)$ be the free semigroupoid of G, in the sense of Kribs and Power. i.e., it is a collection of all vertices of the graph G as units and all admissible finite paths, under the admissibility. The admissible product between two elements in the set $\mathbb{F}^+(G)$ is

Key words and phrases. Graph W*-Probability Sapces over the Diagonal Subalgebras, D_G -valued Moments and Cumulants, D_G -Freeness, D_G -Semicircular Systems, D_G -Semicircular Subalgebras, D_G -valued R-diagonal Systems,

 $^{{\}cal D}_G\text{-valued}$ R-diagonal Subalgebras.

the binary operation on $\mathbb{F}^+(G)$. As a set, the free semigroupoid $\mathbb{F}^+(G)$ can be decomposed by

$$\mathbb{F}^+(G) = V(G) \cup FP(G),$$

where V(G) is the vertex set of the graph G and FP(G) is the set of all admissible finite paths. Trivially the edge set E(G) of the graph G is properly contained in FP(G), since all edges of the graph can be regarded as finite paths with their length 1. Kribs and Power defined the graph Hilbert space $H_G = l^2$ ($\mathbb{F}^+(G)$) with its Hilbert basis $\{\xi_w : w \in \mathbb{F}^+(G)\}$. In [16] and [17], we defined the creation operator L_w by the multiplication operator with its symbol ξ_w and its adjoint L_w^* , the annihilation operator on H_G . They have the following relation; if $w = v_1 w v_2$ in $\mathbb{F}^+(G)$ with $v_1, v_2 \in V(G)$, then

- $\begin{array}{ll} ({\rm i}) & L_w = L_{v_1}L_wL_{v_2} \\ ({\rm ii}) & L_wL_w^* = L_{v_1} \quad {\rm and} \quad L_w^*L_w = L_{v_2} \\ ({\rm iii}) & L_w^2 = L_w = L_w^*, \ {\rm if} \ w = www \in V(G) \\ ({\rm iv}) & L_wL_w^*L_w = L_w, \ {\rm if} \ w \in FP(G). \end{array}$

We define a graph W^* -algebra of G by

$$W^*(G) \stackrel{def}{=} \overline{\mathbb{C}[\{L_w, L_w^* : w \in \mathbb{F}^+(G)\}]}^w.$$

Notice that the creation operators induced by vertices are projections and the creation operators induced by finite paths are partial isometries. We can define the W^* -subalgebra D_G of $W^*(G)$, which is called the diagonal subalgebra by

$$D_G \stackrel{def}{=} \overline{\mathbb{C}[\{L_v : v \in V(G)\}]}^w.$$

Then each element a in the graph W^* -algebra $W^*(G)$ is expressed by

(0.1)
$$a = \sum_{v \in V(G)} p_v L_v + \sum_{w \in FP(G)} \left(p_w L_w + p_w^* L_w^* \right),$$

for $p_v, p_w, p_w^* \in \mathbb{C}$. Here, p_w^* is just a complex number. Remark that p_w^* is not a conjugate $\overline{p_w}$ of p_w in \mathbb{C} . The above expression of a is said to be the Fourier expansion of a. Define the support $\mathbb{F}^+(G:a)$ of a by

$$\mathbb{F}^+(G:a) = V(G:a) \cup FP(G:a)$$

where

$$V(G:a) = \{ v \in V(G) : p_v \neq 0 \text{ in } (0.1) \}$$

and

$$FP(G:a) = \{ w \in FP(G) : p_w \neq 0 \text{ in } (0.1) \} \\ \cup \{ w' \in FP(G) : p_w^* \neq 0 \text{ in } (0.1) \}.$$

Notice that if $V(G:a) \neq \emptyset$, then $\sum_{v \in V(G:a)} p_v L_v$ is contained in the diagonal subalgebra D_G . Thus we have the canonical conditional expectation $E: W^*(G) \to D_G$, defined by

$$E(a) = \sum_{v \in V(G:a)} p_v L_v,$$

for all a in $W^*(G)$ with its Fourier expansion (0.1). Then the algebraic pair $(W^*(G), E)$ is a W^* -probability space with amalgamation over D_G (See [16]). This structure is called the graph W^* -probability space over its diagonal subalgebra D_G , and all elements in $(W^*(G), E)$ are said to be D_G -valued random variables. It is easy to check that the conditional expectation E is faithful in the sense that if $E(a^*a) = 0_{D_G}$, for $a \in W^*(G)$, then $a = 0_{D_G}$.

In [16] and [17], we computed the D_G -valued moments and cumulants of an arbitrary D_G -valued random variable a having its Fourier expansion (0.1). In particular, by using the D_G -cumulant formula, we got the D_G -valued mixed cumulants of D_G -valued random variables a_1 and a_2 and then we could find the D_G -freeness characterization which is so abstract to use. However, by using this characterization, we can characterize the D_G -freeness of generators L_w 's ($w \in \mathbb{F}^+(G)$) D_G -freeness condition;

 L_{w_1} and L_{w_2} are free over D_G in $(W^*(G), E)$ $\iff w_1$ and w_2 in $F^+(G)$ are diagram-distinct,

in the sense that w_1 and w_2 have different diagram on the graph G, graphically. Based on this D_G -freeness condition, in this paper, we will observe the D_G -free structure of the given graph W^* -algebra $W^*(G)$.

Define the subset loop(G) of FP(G), by the collection of all loop finite paths in FP(G). We will consider the family in loop(G)

 $\mathcal{F} = \{l_j \in loop(G) : \text{mutually diagram-distinct}\}_{j=1}^N$

and construct the corresponding D_G -semicircular system

$$\mathcal{L}_N = \left\{ L_{l_j} + L_{l_j}^* : l_j \in F \right\}.$$

Indeed, the elements in \mathcal{L}_N are free from each other over D_G , by the diagramdistinctness of \mathcal{F} and they are all D_G -semicircular, by [17]. So, the family \mathcal{L}_N is the D_G -semicircular system in $(W^*(G), E)$. We show that the free product structure of the W^* -subalgebra $W^*(\mathcal{L}_N, D_G)$ generated by the D_G -semicircular system \mathcal{L}_N and the diagonal subalgebra D_G ;

$$(W^*(\mathcal{L}_N, D_G), E) \simeq (W^*(\mathcal{L}_N, D_N) \otimes D_G, E_N \otimes \mathbf{1}) \simeq (W^*(\mathcal{L}_N, D_N), E_N) \otimes (D_G, \mathbf{1}) = *_{D_G}^N ((W^*(\{L_{l_j}\}, D_N), E_N) \otimes (D_G, \mathbf{1})) i=1$$

where $D_N = \overline{\mathbb{C}[L_{v_j} : l_j = v_j l_j v_j, l_j \in F]}^w$, $E_N = E_{D_N}^{D_G} \circ E$, and **1** is the identity map on D_G .

Also we define a new family

$$\mathcal{R} = \{ L_{w_j}, L_{w_j}^* : w_j \in loop^c(G) \}_{j=1}^N,$$

where $loop^{c}(G) = FP(G) \setminus loop(G)$. Then since the distinctness of non-loop finite paths is the diagram-distinctness on them, $\{L_{w_j}, L_{w_j}^*\}$'s, for j = 1, ..., N, in \mathcal{R} are free from each other over D_G . Moreover, by [17], L_{w_j} and $L_{w_j}^*$ are D_G -valued R-diagonal. We call this family \mathcal{R} , the D_G -valued R-diagonal system. Similar to the D_G -semicircular system case, we will define a W^* -subalgebra $W^*(\mathcal{R}, D_G)$ of the graph W^* -algebra $W^*(G)$, called the R-diagonal subalgebra of $(W^*(G), E)$. We can see that

$$(W^*(\mathcal{R}, D_G), E_{\mathcal{R}}) = \left(\begin{array}{c} {}^N_{*D_{\mathcal{R}}} \left(W^*(\{L_{w_j}\}, D_{\mathcal{R}}), E_{D_{\mathcal{R}}}^{D_G} \circ E \right) \right) \otimes (D_G, \mathbf{1}),$$

where $D_{\mathcal{R}} = \overline{\mathbb{C}[\{L_{v_1}, L_{v_2} : w_j = v_1 w_j v_2\}]}^w$ and $E_{\mathcal{R}} = E|_{W^*(\mathcal{R}, D_G)}$.

We will also define the D_G -free building blocks of the graph W^* -algebra $W^*(G)$ and we prove that

$$(W^{*}(G), E) = (D_{G}, E)$$

$$*_{D_{G}} \begin{pmatrix} *_{D_{G}} \\ l \in Loop(G) \end{pmatrix} (W^{*}(\{L_{l}\}, D_{G}), E) \end{pmatrix}$$

$$*_{D_{G}} \begin{pmatrix} *_{D_{G}} \\ w \in loop^{c}(G) \end{pmatrix} (W^{*}(\{L_{w}\}, D_{G}), E) \end{pmatrix},$$

where Loop(G) is the set of all basic loops in loop(G). A loop l is basic if there is no loop w and a natural number $k \in \mathbb{N} \setminus \{1\}$ such that $l = w^k$. This free product structure of $(W^*(G), E)$ is nice for studying the subalgebras of $W^*(G)$.

Finally, by considering the subalgebra inclusion, we can get the following free product structure of $(W^*(G), E)$,

SUBALGEBRAS OF $(W^*(G), E)$

$$(W^*(G), E) = (D_G, E) *_{D_G} \left(*_{D_G} (W^*(\{L_e\}, D_G), E) \right)$$

1. Semicircular System

1.1. The D_G -Semicircular System. In this chapter, we will consider the amalgamated semicircular system observed by Shlyaktenko (See [10]), in our graph structure. Throughout this chapter, let G be a countable directed graph and $(W^*(G), E)$, the graph W^* -probability space over the diagonal subalgebra D_G .

Definition 1.1. Let B be a von Neumann algebra and A, a von Neumann algebra over B and let $F : A \to B$ be a conditional expectation. Then (A, F) is the amalgamated W^* -probability space over B. The B-valued random variable $x \in (A, F)$ is said to be a B-semicircular element if x is self-adjoint and if the only nonvanishing B-cumulant of x is the second B-cumulant of x. Let $x_1, ..., x_s$ be self-adjoint B-valued random variables in (A, F), where $s \in \mathbb{N}$. We say that the set $S = \{x_1, ..., x_s\}$ is a B-semicircular family if all x_j 's are B-semicircular, for j = 1, ..., s. The B-semicircular family S is said to be a B-semicircular system if $x_1, ..., x_s$ are free from each other over B, in (A, F). The algebra generated by a B-semicircular system and B is called the B-semicircular (sub)algebra of A.

Assume that we have a one-vertex directed graph H. Then the diagonal subalgebra $D_H = \mathbb{C}$. So, in this case, the D_H -semicircularity is same as the Voiculescu's semicircularity. We will define the lattice path model LP_n^* ,

 $LP_n^* = \{L : \text{lattice path having the *-axis-property}\}$

(See [16] and [17]). Take $L \in LP_n^*$. Then we have a (non-unique) corresponding lattice path $l_{w_1,\ldots,w_n}^{u_1,\ldots,u_n}$ of the D_G -valued random variable $L_{w_1}^{u_1} \ldots L_{w_n}^{u_n}$, where $u_j \in \{1,*\}$, in some graph W^* -probability space $(W^*(G), E)$ over D_G .

Theorem 1.1. (See [17]) The D_G -valued random variables $a_l = L_l + L_l^*$ are D_G -semicircular, for all $l = vlv \in loop(G)$, with $v \in V(G)$. In particular, we have that

$$k_n \left(a_l,...,a_l \right) = \begin{cases} 2L_v & \text{if } n = 2 \\ \\ 0_{D_G} & \text{otherwise,} \end{cases}$$

and

$$E\left(a_{l}^{n}\right) = \begin{cases} c_{\frac{n}{2}} \cdot \left(2L_{v}\right)^{\frac{n}{2}} & \text{if } n \text{ is even} \\ 0_{D_{G}} & \text{if } n \text{ is odd,} \end{cases}$$

for all $n \in \mathbb{N}$, where $c_{k} = \frac{1}{k+1} \begin{pmatrix} 2k \\ k \end{pmatrix}$ is the k-th Catalan number. \Box

We will consider the D_G -semicircular family $\{a_1, ..., a_N\}$, for $N \in \mathbb{N}$. Let $l_j = v_j l_j v_j \in loop(G)$, for j = 1, ..., N, with $v_j \in V(G)$. Assume that loops $l_1, ..., l_N$ are mutually **diagram-distinct**. Define D_G -valued random variables $a_1, ..., a_N$,

$$a_j \stackrel{def}{=} L_{l_j} + L_{l_j}^*, \text{ for all } j = 1, ..., N.$$

Again, remark that we assumed that l_j 's are mutually diagram-distinct. So, $a_1, ..., a_N$ are free from each other over D_G in $(W^*(G), E)$ and hence the D_G semicircular family $\{a_1, ..., a_N\}$ is a D_G -semicircular system in $(W^*(G), E)$. So, we have the D_G -semicircular system, in $W^*(G)$, induced by the mutually diagramdistinct loops in FP(G). i.e., the set

 $\mathcal{L}_N = \{a_j : l_j \text{'s are diagram-distinct in } loop(G)\}_{j=1}^N$

is the D_G -semicircular system in $(W^*(G), E)$.

1.2. D_G -Semicircular Subalgebra of $(W^*(G), E)$.

Now, we will construct the D_G -semicircular algebra $W^*(\mathcal{L}_N, D_G)$, as a W^* -subalgebra of the graph W^* -algebra, generated by \mathcal{L}_N and D_G . Let

$$\mathcal{F} = \{l_j \in loop(G) : j = 1, ..., N\}$$

be a collection of mutually diagram-distinct loops in FP(G) and let

$$\mathcal{L}_N = \{a_j = L_{l_j} + L_{l_j}^* : l_j \in \mathcal{F}\}.$$

Then the family \mathcal{L}_N is a D_G -semicircular system in $(W^*(G), E)$ and the W^* -subalgebra $W^*(\mathcal{L}_N, D_G)$ is the D_G -semicurcular subalgebra of $W^*(G)$. The D_G -semicircular subalgebra $W^*(\mathcal{L}_N, D_G)$ have the following free product structure which is very natural by the very definition.

Lemma 1.2. Let $(W^*(G), E)$ be a graph W^* -probability space over the diagonal subalgebra D_G and let

$$\mathcal{L}_N = \{a_j = L_{l_j} + L_{l_j}^* : l_j \text{ 's are diagram-distinct in } loop(G)\}_{j=1}^N.$$

Then the W^* -subalgebra $W^*(\mathcal{L}_N, D_G)$ of $W^*(G)$ is a D_G -semicircular algebra satisfies that

$$W^*\left(\mathcal{L}_N, D_G\right) \simeq *_{D_G}^N W^*(a_j, D_G).$$

Let \mathcal{L}_N be give as above. Assume that $l_j = v_j l_j v_j$, for $v_j \in V(G)$. (It is possible that $v_i = v_k$, for some i, k in $\{1, ..., N\}$.) Define the subalgebra D_N of the diagonal subalgebra D_G by

$$D_N = \overline{\mathbb{C}[\{L_{v_j} : j = 1, ..., N\}]}^w.$$

Trivially, $D_N \leq D_G$, as von Neumann algebras.

Proposition 1.3. (Also See [18]) Let \mathcal{L}_N be the given D_G -semicircular system and let D_N be defined as above. As amalgamated W^* -probability spaces,

$$\begin{array}{ll} \left(W^*\left(\mathcal{L}_N, D_G\right), E\right) &\simeq \left(W^*\left(\mathcal{L}_N, D_N\right) \otimes D_G, E_N \otimes \mathbf{1}\right) \\ &\simeq \left(W^*(\mathcal{L}_N), E_N\right) \otimes \left(D_G, \mathbf{1}\right), \end{array}$$

where $E_N : W^*(\mathcal{L}_N) \to D_N$ is the conditional expectation defined by $E_N = E_{D_N}^{D_G}$ $\circ E$ and $\mathbf{1}(d) = d, \forall d \in D_G$.

Proof. As W^* -algebras, $W^*(\mathcal{L}_N, D_G) \simeq W^*(\mathcal{L}_N, D_N) \otimes D_G$. Indeed, without loss of generality, take $a \in W^*(\mathcal{L}_N, D_G)$ by

$$a = d_1 a_{l_{i_1}}^{k_1} d_2 a_{l_{i_2}}^{k_2} \dots d_n a_{l_{i_n}}^{k_n}$$
 and $a_{l_j} = L_{l_j} + L_{l_j}^*$

where $d_1, ..., d_n \in D_G, k_1, ..., k_n \in \mathbb{N}$ and $(i_1, ..., i_n) \in \{1, ..., N\}^n, n \in \mathbb{N}$. Observe that, for any $j \in \{1, ..., N\}$, we have that

$$a_{l_j}^k = \left(L_{l_j} + L_{l_j}^*\right)^k = L_{l_j^k} + L_{l_j^k}^* + Q\left(L_{l_j}, L_{l_j}^*\right),$$

where $Q \in \mathbb{C}[z_1, z_2]$. Also, observe that $L_{l_j}^{k_1} L_{l_j}^{* k_2}$, for any $k_1, k_2 \in \mathbb{N}$, satisfies that

$$L_{l_j}^{k_1} L_{l_j}^{* \ k_2} = L_{l_j^{k_1}} L_{l_j^{k_2}}^{*} = \begin{cases} L_{l_j^{k_1 - k_2}} = L_{v_j} L_{l_j^{k_1 - k_2}} & \text{if } k_1 > k_2 \\ L_{l_j^{k_2 - k_1}}^{*} = L_{v_j} L_{l_j^{k_2 - k_1}}^{*} & \text{if } k_1 < k_2 \\ L_{v_j} & \text{if } k_1 = k_2, \end{cases}$$

and similarly,

$$L_{l_j}^{* k_1} L_{l_j}^{k_2} = L_{l_j}^{*} L_{l_j}^{k_2} = \begin{cases} L_{l_j}^{*} L_{l_j}^{k_1 - k_2} = L_{v_j} L_{l_j}^{*} L_{l_j}^{k_1 - k_2} & \text{if } k_1 > k_2 \\ L_{l_j}^{k_2 - k_1} = L_{v_j} L_{l_j}^{k_2 - k_1} & \text{if } k_1 < k_2 \\ L_{v_j} & \text{if } k_1 = k_2 \end{cases}$$

So,

$$Q(L_{l_j}, L_{l_j}^*) = L_{v_j} \left(Q(L_{l_j}, L_{l_j}^*) \right) L_{v_j},$$

for all j = 1, ..., N. Thus

(1.1)
$$a_{l_j}^k = L_{v_j} a_{l_j}^k = L_{v_j} a_{l_j}^k L_{v_j}$$
, for all $j = 1, ..., N$.

Now, consider that

$$d_j = d_j^N + d'_j, \, \forall j = 1, ..., N.$$

where $d_j^N = \sum_{j=1}^N L_{v_j} d_j L_{v_j}$ and $d'_j = d_j - d_j^N$ in D_G . So, we can rewrite that $a = (d_1^N + d'_1) a_{l_{i_1}}^{k_1} (d_2^N + d'_2) a_{l_{i_2}}^{k_2} \dots (d_n^N + d'_n) a_{l_{i_n}}^{k_n}$ $= d_1^N a_{l_{i_1}}^{k_1} d_2^N a_{l_{i_2}}^{k_2} \dots d_n^N a_{l_{i_n}}^{k_n} + d'_1 a_{l_{i_1}}^{k_1} d'_2 a_{l_{i_2}}^{k_2} \dots d'_n a_{l_{i_n}}^{k_n}$ $= d_1^N a_{l_{i_1}}^{k_1} d_2^N a_{l_{i_2}}^{k_2} \dots d_n^N a_{l_{i_n}}^{k_n},$

by (1.1). This shows that $a = a \otimes 1 \in W^*(\mathcal{L}_N, D_N) \otimes 1$ and

$$E(a) = E_{D_N}^{D_G} \circ E(a) = E_N(a) = E_N \otimes \mathbf{1}(a \otimes 1)$$

Trivially, if $a \in D_G \subset W^*(\mathcal{L}_N, D_G)$, then $a = 1 \otimes a \in 1 \otimes D_G$. Futhermore, if $a \in D_G$ in $W^*(\mathcal{L}_N, D_G)$, then

$$E(a) = a = 1 \otimes a = E_N \otimes \mathbf{1}(1 \otimes a).$$

Now, consider $W^*(\mathcal{L}_N, D_N)$. By the previous lemma, similarly, we have that

(1.2)
$$W^*(\mathcal{L}_N, D_N) = W^*(\{a_1\}, D_N) *_{D_N} \dots *_{D_N} W^*(\{a_N\}, D_N).$$

Indeed, the D_G -semicircular elements a_i and a_j in L_N are free over D_N in $W^*(L_N, D_N)$. Clearly, since $D_N \subset D_G$ and since a_i and a_j are free over D_G , they are free over D_N . Therefore, the formula (3.1.2) holds true with respect to the (compressed) conditional expectation

$$E_N = E \mid_{W^*(L_N, D_N)} = E_{D_N}^{D_G} \circ E,$$

on $W^*(\mathcal{L}_N, D_N)$.

Corollary 1.4. Let $\mathcal{F} = \{l_j : l_j = v_0 l_j v_0\}_{j=1}^N$ be a family of mutually diagramdistinct loops in loop(G), where $v_0 \in V(G)$. If the collection

$$\mathcal{L}_N = \{ \frac{1}{\sqrt{2}} a_{l_j} : l_j \in \mathcal{F} \}$$

is a D_G -semicircular system induced by the family \mathcal{F} , then

$$(W^*(\mathcal{L}_N, D_G), E) \simeq (L(F_N), tr) \otimes (D_G, \mathbf{1}),$$

in the sense of Voiculescu, where tr is the canonical II₁-trace of the free group factor $L(F_K)$, $\forall K \in \mathbb{N}$, and where $\mathbf{1}(d) = d$, $\forall d \in D_G$. \Box

By the previous proposition and corollary, we can have the following fact ;

Theorem 1.5. Let $\mathcal{F} = \{l_{k1}, ..., l_{kn_k} : l_{kj} = v_k l_{kj} v_k, j = 1, ..., n_k\}_{k=1}^N$ be the collection of $\sum_{k=1}^N n_k$ -mutually diagram-distinct loops in FP(G). Assume that $v_{k_1} \neq v_{k_2}$, for all pair $(k_1, k_2) \in \mathbb{N}^2$. If

and

$$D_{\mathcal{L}} = \overline{\mathbb{C}[\{L_{v_k} : k = 1, ..., N\}]}^w$$

 $E_{\mathcal{L}} = E_{D_L}^{D_G} \circ E,$

 $\mathcal{L} = \{\frac{1}{\sqrt{2}}a_{l_{k_j}} : j = 1, ..., n_k\}_{k=1}^N$

are the corresponding D_G -semicircular system and the subalgebra of D_G by L, respectively, and

then

$$(W^*(\mathcal{L}, D_G), E) \simeq \begin{pmatrix} N \\ *D_{\mathcal{L}} \\ k=1 \end{pmatrix} ((L(F_{n_k}), tr) \otimes (D_{\mathcal{L}}, \mathbf{1})) \otimes (D_G, \mathbf{1})$$

Corollary 1.6. Let $\mathcal{F}_1 = \{l_1^1, ..., l_N^1 : l_j^1 = v_1 l_j^1 v_1\}$ and $\mathcal{F}_2 = \{l_1^2, ..., l_N^2 : l_j^2 = v_2 l_j^2 v_2\}$ be the N-mutually diagram-distinct families of loops in FP(G), where $v_1 \neq v_2 \in V(G)$ are fixed. Let

$$\mathcal{L}_{1} = \left\{ a_{k} = \frac{1}{\sqrt{2}} \left(L_{l_{k}^{1}} + L_{l_{k}^{1}}^{*} \right) : k = 1, ..., N \right\}$$

and
$$\mathcal{L}_{2} = \left\{ b_{k} = \frac{1}{\sqrt{2}} \left(L_{l_{k}^{2}} + L_{l_{k}^{2}}^{*} \right) : k = 1, ..., N \right\}$$

be the corresponding D_G -semicircular systems, respectively. Then two D_G -semicircular subalgebras $(W^*(\mathcal{L}_1, D_G), E)$ and $(W^*(\mathcal{L}_2, D_G), E)$ are free over D_G and they are isomorphic, as amalgamated W^* -probability spaces. \Box

The above corollary shows us how to construct the isomorphic semicircular subalgebras in the graph W^* -probability space from two vertices having the same number of loops. (Assume that the vertex v_1 has its loops $l_1^1, \ldots, l_{n_1}^1$ and v_2 has its loops $l_1^2, \ldots, l_{n_2}^2$ and suppose that $n_1 < n_2$. Then we can choose n_1 -loops of $v_2, l_{i_1}^2, \ldots, l_{i_{n_1}}^2$. And then we can apply the above corollary for them.)

One of the most interesting example of D_G -semicircular subalgebra is as follows ;

Example 1.1. Let G be a directed graph with

$$V(G) = \{v\} and E(G) = \{l_1, ..., l_N : l_j = vl_jv\}$$

Note that $D_G = \Delta_1 = \mathbb{C}$. Also, note that the projection $L_v = 1_{D_G} = 1_{\mathbb{C}} = 1 \in \mathbb{C}$. We have the graph W^* -probability space $(W^*(G), E)$ over \mathbb{C} . Consider the W^* -subalgebra $W^*(\mathcal{L}_N, D_G) = W^*(\mathcal{L}_N)$, where

$$\mathcal{L}_N = \left\{ \frac{1}{\sqrt{2}} a_j : a_j = L_{l_j^{n_j}} + L_{l_j^{n_j}}^* \right\}_{j=1}^N,$$

where $n_j \in \mathbb{N}$, j = 1, ..., N. Then \mathcal{L}_N is a D_G -semicircular system, too. Definitely it is a D_G -semicircular family. Since $\mathcal{F} = \{l_j^{n_j} : j = 1, ..., N\}$ is consists of mutually diagram-distinct loops in FP(G), a_j 's are free from each other and hence \mathcal{L}_N is a D_G -semicircular system. We have that

$$\begin{aligned} k_2 \left(\frac{1}{\sqrt{2}} a_j, \frac{1}{\sqrt{2}} a_j \right) &= \frac{1}{2} k_2 \left(a_j, a_j \right) \\ &= \frac{1}{2} k_2 \left(L_{l_j^{n_j}} + L_{l_j^{n_j}}^*, L_{l_j^{n_j}} + L_{l_j^{n_j}}^* \right) \\ &= \frac{1}{2} \sum_{(r_1, r_2) \in \{1, *\}^2} k_2 \left(L_{l_j^{n_j}}^{r_1}, L_{l_j^{n_j}}^{r_2} \right) \\ &= \frac{1}{2} \left(\mu_{l_j^{n_j}, l_j^{n_j}}^{1, *} \cdot E \left(L_{l_j^{n_j}}, L_{l_j^{n_j}}^* \right) + \mu_{l_j^{n_j}, l_j^{n_j}}^{*, 1} \cdot E \left(L_{l_j^{n_j}}^*, L_{l_j^{n_j}}^* \right) \right) \\ &= \frac{1}{2} \left(L_v + L_v \right) = \frac{1}{2} \cdot 2 = 1, \end{aligned}$$

for all j = 1, ..., N. So, we can get that

$$(W^*(L_N), E) = (W^*(\mathcal{L}_N, D_G), E)$$

= $(W^*(\mathcal{L}_N, D_N), E_N) \otimes (D_G, \mathbf{1})$
= $(W^*(\mathcal{L}_N), E_N) \otimes (\mathbb{C}, \mathbf{1})$
= $(L(F_N), tr),$

where $tr : L(F_N) \to \mathbb{C}$ is the canonical II_1 -trace on the free group factor $L(F_K)$, $\forall K \in \mathbb{N}$. So, the graph W^* -probability space $(W^*(G), E)$ contains the free group factor $L(F_N)$ which is isomorphic to the D_G -semicircular subalgebra $W^*(\mathcal{L}_N)$, generated by \mathcal{L}_N .

2. R-DIAGONAL SYSTEMS

In this chapter, similar to Chapter 2, we will consider the special W^* -subalgebra of the graph W^* -probability space $(W^*(G), E)$ over the diagonal subalgebra D_G . As we defined the D_G -semicircular systems in $W^*(G)$, we will define the $(D_G$ -valued) R-diagonal systems in $W^*(G)$. Recall that if $w \in loop^c(G)$ is a (non-loop) finite path in $\mathbb{F}^+(G)$, then the D_G -valued random variables L_w and L_w^* are R-diagonal over D_G (See [17]). Take a finite family

$$\mathcal{F} = \{w_j : w_j \in loop^c(G)\}_{j=1}^N, N \in \mathbb{N}.$$

Define a (D_G -valued) R-diagonal family induced by \mathcal{F} by

$$R = \{L_w, L_w^* : w \in \mathcal{F}\}.$$

Notice that, by the (diagram-)distinctness of w_j 's in \mathcal{F} , the subfamilies $\{L_{w_1}, L_{w_1}^*\}$, ..., $\{L_{w_N}, L_{w_N}^*\}$ are free from each other over D_G in $(W^*(G), E)$. We will observe that the R-diagonal subalgebra $W^*(R, D_G)$ satisfies that

$$(W^*(R, D_G), E) = *_{D_R}^N \left(\left(W^*(\{L_{w_j}\}, D_R), E_R \right) \otimes (D_G, \mathbf{1}) \right),$$

where D_R is the subalgebra generated by the projections

$$\{L_{v_1}, L_{v_2}: w = v_1 w v_2, \forall w \in \mathcal{F}\}.$$

and $E_R = E \mid_{W^*(\{L_{w_j}\}_{j=1}^N, D_R)}$ is the restricted conditional expectation onto D_R .

2.1. D_G -valued R-diagonal Systems.

Let G be a countable directed graph and $(W^*(G), E)$, the graph W^* -probability space over the diagonal subalgebra D_G . Throughout this section, we will fix the following finite family,

$$\mathcal{F} = \{ w_j : w_j \in loop^c(G) \}_{j=1}^N \subset FP(G),$$

where $N \in \mathbb{N}$. Notice that all elements in the family \mathcal{F} are non-loop finite paths and hence the corresponding D_G -valued random variables are D_G -valued R-diagonal elements (See [17]).

Definition 2.1. Let $a \in (W^*(G), E)$ be a D_G -valued random variable. The D_G -valued random variable is said to be an $(D_G$ -valued) R-diagonal element if it has the only nonvanishing D_G -valued cumulants having their forms of

$$k_n(a, a^*, ..., a, a^*)$$
 and $k_n(a^*, a, ..., a^*, a)$,

for all $n \in 2\mathbb{N}$ (See [19] and [17]). Clearly, if a is a R-diagonal, then automatically a^* is R-diagonal. (In other words, the pair (a, a^*) is a D_G -valued R-diagonal pair. Also, see [19].) Suppose we have a collection

$$R = \{a_1, \dots, a_N : a_j \text{ is } R \text{-} diagonal \text{ over } D_G\}.$$

Then the family R is called the $(D_G$ -valued) R-diagonal system. The subalgebra $W^*(R, D_G)$ is called the $(D_G$ -valued) R-diagonal subalgebra, induced by R in $(W^*(G), E)$.

In [17], we showed the following theorem;

Theorem 2.1. Let $w \in FP(G)$. Then the D_G -valued random variables L_w and L_w^* are *R*-diagonal over D_G in $(W^*(G), E)$. \Box

Notice that all D_G -semicircular elements are R-diagonal, by definition. But we will restrict our interests to the R-diagonal systems consisting of the non-loop finite paths. Recall that if $w_1 \neq w_2 \in loop^c(G)$, then the D_G -valued R-diagonal elements L_{w_1} and L_{w_2} are free over D_G , in $(W^*(G), E)$, since the distinctness of non-loop finite paths is equivalent to the diagram-distinctness of them.

2.2. D_G -valued R-diagonal Subalgebras.

In this section, we will consider the D_G -valued R-diagonal subalgebras of $W^*(G)$, generated by the fixed R-diagonal systems and D_G . As before, let \mathcal{F} be a finite family consisting of N-mutually diagram-distinct non-loop finite paths and let R = $\{L_w : w \in \mathcal{F}\}\$. As we saw in the previous section, the family R is the R-diagonal system in $(W^*(G), E)$. Therefore, we can get the following result ;

Proposition 2.2. Let \mathcal{F} and R be given as before and let $W^*(R, D_G)$ be the R-diagonal subalgebra of $W^*(G)$. Then

$$(W^*(R, D_G), E_R) = *_{D_G}^N (W^*(\{L_{w_j}\}, D_G), E_j),$$

where $E_R = E \mid_{W^*(R,D_G)}$ and $E_j = E \mid_{W^*(\{L_{w_j}\},D_G)}$, for all j = 1, ..., N. \Box

Similar to the previous chapter, we will define

$$D_R \stackrel{def}{=} \overline{\mathbb{C}[\{L_{v_1}, L_{v_2} : w = v_1 w v_2, w \in \mathcal{F}\}]}^w.$$

Notice that, for the given R-diagonal system R, the von Neumann algebra D_R should not be \mathbb{C} , because \mathcal{F} is consists of all non-loop finite paths which are mutually distinct. For example, if the family $\mathcal{F} = \{w_0 = v_1 \ w_0 \ v_2\}$, where $v_1 \neq v_2$ in V(G), and $R = \{L_{w_0} : w_0 \in \mathcal{F}\}$. Then

$$D_R = \overline{\mathbb{C}[\{L_{v_1}, L_{v_2}\}]}^w \simeq \Delta_2,$$

where Δ_2 is the subalgebra of the matricial algebra $M_2(\mathbb{C})$ generated by all diagonal matrices. Notice that, for the inclusion $D_R \subset D_G$, there exists the well-determined canonical conditional expectation $E_{D_R}^{D_G}: D_G \to D_R$. Also, notice that $D_R \subset W^*(R)$.

Proposition 2.3. Let \mathcal{F} and R be given as before and let $W^*(R, D_G)$ is the R-diagonal subalgebra of $W^*(G)$. Then

$$(W^*(R, D_G), E_R) = \left(W^*(R, D_R), E_{D_R}^{D_G} \circ E\right) \otimes (D_G, \mathbf{1}),$$

where $E_R = E \mid_{W^*(R,D_G)}$ and **1** is the identity map on D_G .

Proof. Let $a \in (W^*(R, D_G), E_R)$ be the nonzero D_G -valued random variable such that

$$a = d_1 L_{w_{i_1}}^{r_1} d_2 L_{w_2}^{r_2} \dots d_n L_{w_n}^{r_n} \quad \text{or} \ a \in D_G,$$

where $d_1, ..., d_n \in D_G$, $r_1, ..., r_n \in \{1, *\}$ and $(i_1, ..., i_n) \in \{1, ..., N\}^n$, $n \in \mathbb{N}$.

Let
$$d_j = \sum_{v_j \in V(G:d_j)} q_{v_j} L_{v_j}$$
, for all $j = 1, ..., n$. Then

$$a = \sum_{(v_1, \dots, v_n) \in \Pi_{j=1}^n V(G:d_j)} (\Pi_{j=1}^n q_{v_j}) \left(L_{v_1} L_{w_{i_1}}^{r_1} L_{v_2} L_{w_2}^{r_2} \dots L_{v_n} L_{w_n}^{r_n} \right)$$
$$= \sum_{(v_1, \dots, v_n) \in \Pi_{j=1}^n V(G:d_j)} (\Pi_{j=1}^n q_{v_j}) (\Pi_{j=1}^n \delta_{(v_j, x_j, y_j:r_j)}) \left(L_{w_{i_1}}^{r_1} L_{w_2}^{r_2} \dots L_{w_n}^{r_n} \right),$$

where $\delta_{(v_j,x_j,y_j:r_j)} = \delta_{v_j,x_j}$ if $r_j = 1$ and $\delta_{(v_j,x_j,y_j:r_j)} = \delta_{v_j,y_j}$ if $r_j = *$. Now, assume that there exists at least one $j \in \{1, ..., n\}$ such that $v_j \notin D_R$. Then $\delta_{(v_j,x_j,y_j:r_j)} = 0$, where either $r_j = 1$ or $r_j = *$. So, to make *a* be nonzero, the elements $d_1, ..., d_n$ should be chosen in $D_R \subset D_G$. This shows that the arbitrary element *x* in the R-diagonal subalgebra $W^*(R, D_G)$, we have that

$$x = x \otimes 1 \in W^*(R, D_R) \otimes 1$$

or
$$x = 1 \otimes a \in 1 \otimes D_G$$

or
$$x = x_1 \otimes x_2 \in W^*(R, D_R) \otimes D_G.$$

Also, if $a = d_1 L_{w_{i_1}}^{r_1} d_2 L_{w_2}^{r_2} \dots d_n L_{w_n}^{r_n} \in W^*(R, D_R)$, in $W^*(R, D_G)$, for $d_1, \dots, d_n \in D_R$, then

$$E_R \left(d_1 L_{w_{i_1}}^{r_1} d_2 L_{w_2}^{r_2} \dots d_n L_{w_n}^{r_n} \right) = E_{D_R}^{D_G} \circ E \left(d_1 L_{w_{i_1}}^{r_1} d_2 L_{w_2}^{r_2} \dots d_n L_{w_n}^{r_n} \right)$$
$$= E_{D_R}^{D_G} \circ E \left(d_1 L_{w_{i_1}}^{r_1} d_2 L_{w_2}^{r_2} \dots d_n L_{w_n}^{r_n} \right).$$

Trivially, if $a \in D_G$ in $W^*(R, D_G)$, then

$$E_R(a) = E(a) = a = \mathbf{1}(a),$$

where **1** is the identity map on D_G .

By the above proposition, we can conclude this section by the following theorem

Theorem 2.4. Let \mathcal{F} and R be given as before and let $W^*(R, D_G)$ be the R-diagonal subalgebra of $W^*(G)$. Then

$$(W^*(R, D_G), E_R) = \left(\begin{array}{c} N \\ *_{D_R} \left(W^*(\{L_{w_j}\}, D_R), E_{D_R}^{D_G} \circ E \right) \right) \otimes (D_G, \mathbf{1}).$$

Proof. Notice that if two D_G -valued random variables x and y are free over D_G in $(W^*(G), E)$, then they are free over D_R , since $D_R \subset D_G$. i.e., since all mixed

14

;

 D_G -valued cumulants of $p(x, x^*)$ and $q(y, y^*)$ vanish, for all $p, q \in \mathbb{C}[z_1, z_2]$, all their mixed D_R -valued cumulants vanish, again. Thus, we have that

$$\left(W^*(R, D_R), E_{D_R}^{D_G} \circ E\right) = *_{D_G}^N \left(W^*(\{L_{w_j}\}, D_G), E_{D_R}^{D_G} \circ E\right).$$

Therefore, by the previous proposition, we can get that

$$(W^*(R, D_G), E_R) = \left(W^*(R, D_R), E_{D_R}^{D_G} \circ E\right) \otimes (D_G, \mathbf{1})$$
$$= \left(\begin{array}{c} N\\ *D_R\\ j=1 \end{array} \left(W^*(\{L_{w_j}\}, D_R), E_{D_R}^{D_G} \circ E\right) \right) \otimes (D_G, \mathbf{1}).$$

Now, we will provide the following fundamental examples ;

Example 2.1. Let G be a directed graph with $V(G) = \{v_1, v_2\}$ and $E(G) = \{e = v_1 e v_2\}$. Let

$$\mathcal{F} = \{e\} \quad and \quad R = \{L_e, L_e^*\}.$$

We can construct the R-diagonal subalgebra $W^*(R, D_G)$. It is easy to check that

$$W^*(R, D_G) = W^*(\{L_e, L_e^*\}, D_G)$$

= $W^*(\{L_e, L_e^*\}, D_R) \otimes D_G$
= $W^*(\{L_e, L_e^*, L_{v_1}, L_{v_2}\}).$

By the result from the final chapter, later, we can conclude that $W^*(R, D_G) = W^*(G)$. So, the graph W^* -algebra $W^*(G)$ is same as the R-diagonal subalgebra $W^*(R, D_G)$ of it, where R is consists of all generators of $W^*(G)$ induced by the edge.

Example 2.2. Let G be a directed graph with $V(G) = \{v_1, v_2\}$ and $E(G) = \{e_j = v_1e_jv_2\}_{j=1}^N$. Let $n \leq N$. Then we have a R-diagonal system,

$$R = \{L_{e_1}, L_{e_1}^*, \dots, L_{e_n}, L_{e_n}^*\}, \,\forall n \le N.$$

The R-diagonal subalgebra $W^*(R, D_G)$ is trivially a W^* -subalgebra of $W^*(G)$. Also,

$$(W^*(R, D_G), E_R) = *_{D_G}^n (W^*(\{L_{e_j}\}, D_G), E_j)$$

$$= \left(*_{D_R}^n \left(W^*(\{L_{e_j}\}, D_R), E_j \right) \right) \otimes (D_G, \mathbf{1}).$$

Example 2.3. Let G be a directed graph with $V(G) = \{v_1, v_2, v_3\}$ and

$$\begin{split} E(G) &= \{ e_j = v_1 e_j v_2, \ e_k' = v_2 e_k' v_3 \\ &: j = 1, ..., n \ \mathcal{C} \ k = 1, ..., m \}. \end{split}$$

Take

where

$${\cal F}={\cal F}_1{\cup}{\cal F}_2,$$

 $\mathcal{F}_1 = \{e_1, ..., e_n\} \text{ and } \mathcal{F}_2 = \{e'_1, ..., e'_m\}.$

If we construct the R-diagonal systems R_1 and R_2 , induced by \mathcal{F}_1 and \mathcal{F}_2 , respectively, then R_1 and R_2 are free over D_G in $(W^*(G), E)$, because they are totally disjoint (See [16]). Therefore, the R-diagonal subalgebra generated by $R_1 \cup R_2$ is

$$W^* (R_1 \cup R_2, D_G) = W^* (R_1, D_G) *_{D_G} W^* (R_2, D_G)$$
$$= \left(\left(\left(*_{D_{R_1}}^n \left(W^* (\{L_{e_j}\}, D_{R_1}), E_{e_j} \right) \right) \otimes (D_G, \mathbf{1}) \right) \right)$$
$$*_{D_G} \left(\left(\left(*_{D_{R_2}}^m \left(W^* (\{L_{e_j'}\}, D_{R_2}), E_{e_j'} \right) \right) \otimes (D_G, \mathbf{1}) \right) \right).$$

3. Free Product Structures of $(W^*(G), E)$

Throughout this chapter, let G be a countable directed graph and $(W^*(G), E)$, the graph W^* -probability space over the diagonal subalgebra D_G . In this chapter, we will consider the building blocks of $W^*(G)$. Notice that if $w_1 \neq w_2 \in FP(G)$ and if $w_1w_2 \in \mathbb{F}^+(G)$, then we can construct the D_G -valued random variable $L_{w_1w_2}$ which is same as $L_{w_1}L_{w_2}$. Also notice that L_{w_1} and L_{w_2} are not free over D_G , in general. But under the diagram-distinctness of w_1 and w_2 , L_{w_1} and L_{w_2} are free over D_G .

3.1. The D_G -Free Product Structures of $(W^*(G), E)$ I..

16

In this section, we will consider the D_G -semicircular algebras and D_G -valued R-diagonal algebras in $W^*(G)$, more in detail. Recall that the loop l is basic if there is no other loop w such that $l = w^k$, for some $k \in \mathbb{N} \setminus \{1\}$. Define

$$Loop(G) \stackrel{def}{=} \{l \in loop(G) : l \text{ is basic}\}.$$

The following lemma is easily proved, by the very definition of basic loops ;

Lemma 3.1. Let $w \in loop(G)$ with $w = l^k$, for some $l \in Loop(G)$ and $k \in \mathbb{N} \setminus \{1\}$. Then

$$W^*(\{L_w\}, D_G) \le W^*(\{L_l\}, D_G).$$

Remark that since w and l are not diagram-distinct, L_w and L_l are not free over D_G in $(W^*(G), E)$. In fact, $W^*(\{L_l\}, D_G)$ contains all $W^*(\{L_w\}, D_G)$, if $l \in Loop(G)$ and $w = l^k$, $\forall k \in \mathbb{N}$.

Proposition 3.2. $(W^*(\{L_l : l \in loop(G)\}, D_G), E) = \underset{l \in Loop(G)}{*} M^*(\{L_l\}, D_G), E).$

Proof. Let

$$\mathcal{L} = \{L_l : l \in loop(G)\}, \\ \mathcal{L}_l = \{L_{l^k} : l \in Loop(G), k \in \mathbb{N}\}$$

and

Then

$$\mathcal{L}_0 = \{ L_l : l \in Loop(G) \}.$$

$$\mathcal{L} = \bigcup_{l \in Loop(G)} \mathcal{L}_l = \bigcup_{L_l \in \mathcal{L}_0} \left(\bigcup_{k=1}^\infty \{ L_l^k \} \right).$$

Thus

$$(W^* \left(\{ L_l : l \in loop(G) \}, D_G \right), E \right) = (W^* (\mathcal{L}, D_G), E)$$
$$= \left(W^* \left(\bigcup_{l \in Loop(G)} \mathcal{L}_l, D_G \right), E \right)$$
$$= \left(W^* \left(\bigcup_{L_l \in \mathcal{L}_0} \left(\bigcup_{k=1}^{\infty} \{ L_l^k \} \right), D_G \right), E \right)$$
$$= \sum_{\substack{k \in \mathcal{L}_0 \\ L_l \in \mathcal{L}_0}} \left(W^* \left(\left(\bigcup_{k=1}^{\infty} \{ L_l^k \} \right), D_G \right), E \right)$$

by the fact that if $L_{l_1} \neq L_{l_2}$ in \mathcal{L}_0 , then they are free over D_G , by the diagramdistinctness of $l_1 \neq l_2 \in Loop(G)$

$$= *_{D_g} (W^*((\{L_l\}), D_G), E),$$

since $W^*((\bigcup_{k=1}^{\infty} \{L_l^k\}), D_G) = W^*((\{L_l\}), D_G).$ Therefore,
 $(W^*(\{L_l : l \in loop(G)\}, D_G), E) = *_{D_g} (W^*((\{L_l\}), D_G), E).$

Finally, we can have the D_G -free product structure of the graph W^* -probability space $(W^*(G), E)$ over its diagonal subalgebra D_G . By considering the D_G -freeness of generators of $W^*(G)$, we can characterize the free product structure of $(W^*(G), E)$.

Theorem 3.3. Let G be a countable directed graph and $(W^*(G), E)$, the graph W^* -probability space over its diagonal subalgebra D_G . Then

$$(W^{*}(G), E) = (D_{G}, E)$$

$$*_{D_{G}} \begin{pmatrix} *_{D_{G}} \\ l \in Loop(G) \end{pmatrix} (W^{*}(\{L_{l}\}, D_{G}), E) \end{pmatrix}$$

$$*_{D_{G}} \begin{pmatrix} *_{D_{G}} \\ w \in loop^{c}(G) \end{pmatrix} (W^{*}(\{L_{w}\}, D_{G}), E) \end{pmatrix}.$$

Proof. Recall that D_G -valued random variables L_{w_1} and L_{w_2} are free over D_G in $(W^*(G), E)$ if and only if w_1 and w_2 are diagram-distinct. So, for any loop l and non-loop finite path w, L_l and L_w are free over D_G . So,

and

$$W^*\left(\{L_w: w \in loop^c(G)\}, D_G\right)$$

 $W^*\left(\{L_l: l \in loop(G)\}, D_G\right)$

are free over D_G in $(W^*(G), E)$. Denote the above subalgebras by \mathbf{L} and \mathbf{R} , respectively. Therefore, we have that the free product space

$$D_G \quad *_{D_G} W^* \left(\{ L_l : l \in loop(G) \}, D_G \right)$$
$$*_{D_G} W^* \left(\{ L_w : w \in loop^c(G) \}, D_G \right)$$

is contained in $W^*(G)$. Since the generators of $W^*(G)$ and those of $D_G *_{D_G} \mathcal{L} *_{D_G} \mathcal{R}$ are same, we can conclude that

$$(W^*(G), E) = (D_G *_{D_G} \mathbf{L} *_{D_G} \mathbf{R}, E).$$

But, by the previous proposition, we obtained that

18

$$(\mathbf{L}, E) = *_{D_G}_{l \in Loop(G)} (W^*(\{L_l\}, D_G), E).$$

Now, we will observe that

$$(\mathbf{R}, E) = *_{D_G}_{w \in loop^c(G)} (W^*(\{L_w\}, D_G), E).$$

Assume that $L_{w_1}, L_{w_2} \in (\mathbf{R}, E)$ are the generators. Then

$$W^*(\{L_{w_1}, L_{w_2}\}, D_G) = W^*(\{L_{w_1}\}, D_G) *_{D_G} W^*(\{L_{w_2}\}, D_G).$$

Therefore, we can get

$$(\mathbf{R}, E) \leq *_{D_G}_{w \in loop^c(G)} (W^*(\{L_w\}, D_G), E).$$

The subalgebra inclusion " \geq " is clear. So,

$$(W^{*}(G), E) = (D_{G} *_{D_{G}} \mathbf{L} *_{D_{G}} \mathbf{R}, E)$$

= $(D_{G}, E) *_{D_{G}} (\mathbf{L}, E) *_{D_{G}} (\mathbf{R}, E)$
= $(D_{G}, E) *_{D_{G}} \left(*_{D_{G}} (W^{*}(\{L_{l}\}, D_{G}), E) \right)$
 $*_{D_{G}} \left(*_{D_{G}} (W^{*}(\{L_{w}\}, D_{G}), E) \right).$

3.2. D_G -Free Building Blocks of $(W^*(G), E)$.

In this section, we will construct the D_G -free building blocks of the graph W^* -probability space $(W^*(G), E)$ over its diagonal subalgebra D_G . Recall that

$$\begin{aligned} (W^*(G), E) &= & (D_G, E) \\ & *_{D_G} \begin{pmatrix} *_{D_G} \\ l \in Loop(G) \end{pmatrix} \\ & *_{D_G} \begin{pmatrix} *_{D_G} \\ w \in loop^c(G) \end{pmatrix} (W^*(\{L_w\}, D_G), E) \end{pmatrix}, \end{aligned}$$

where Loop(G) is the collection of all basic loops contained in loop(G). Notice that, even though we have a finite directed graph, Loop(G) and $loop^{c}(G)$ may contain countably many elements. So, $(W^{*}(G), E)$ is, in general, the D_{G} -free product of infinitely many D_{G} -free W^{*} -subalgebras. But, in the final chapter, we will show that this infinite free product of algebras can be contained in the finite free product of algebras.

Definition 3.1. Let G be a countable directed graph and let $W^*(G)$ be the graph W^* -algebra. The diagonal subalgebra D_G and W^* -subalgebras, $W^*(\{L_l\}, D_G)$ and $W^*(\{L_w\}, D_G)$ for all $l \in Loop(G)$ and $w \in loop^c(G)$ are D_G -free building blocks of $W^*(G)$.

As we observed in Chapter 1 and Chapter 2, we have that if l is a loop, then

$$(3.1) \quad (W^*(\{L_l\}, D_G), E) = (W^*(\{L_l\}), tr) \otimes (D_G, \mathbf{1}),$$

where $tr = E \mid_{W^*(\{L_l\})}$ is a tracial linear functional on $W^*(\{L_l\})$. We also have that if w is a non-loop finite path, then

(3.2)
$$(W^*(\{L_w\}, D_G), E) = (W^*(\{L_w\}, D_w), E_2) \otimes (D_G, \mathbf{1}),$$

where $D_w = \overline{\mathbb{C}[\{L_{v_1}, L_{v_2} : w = v_1 w v_2\}]}^w$ is the W^* -subalgebra of D_G and $E_2 = E_{D_w}^{D_G} \circ E$. By (3.2), we can get the following proposition which shows us the vector space property of the non-loop D_G -free building blocks of $W^*(G)$.

Proposition 3.4. Let $w \in loop^{c}(G)$ be a non-loop finite path and let $W^{*}(\{L_{w}\}, D_{G})$ be the corresponding free building block. Then, as a topological vector space,

$$W^*(\{L_w\}, D_G) = \overline{\mathbb{C}\{d, \, p(L_w, L_w^*) : d \in D_G, \, p \in \mathbb{C}_1[z_1, z_2]\}}^w,$$

where

$$\mathbb{C}_1[z_1, z_2] \stackrel{def}{=} \{ p \in \mathbb{C}[z_1, z_2] : p(z_1, z_2) = \alpha_0 + \alpha_1 z_1 + \alpha_2 z_2 \},\$$

for $\alpha_0, \alpha_1, \alpha_2 \in \mathbb{C}$. \Box

Proof. In general, if $w \in FP(G)$ is a finite path, then the free building block $W^*(\{L_w\}, D_G)$ is a W^* -subalgebra of the graph W^* -algebra $W^*(G)$ such that

$$W^*(\{L_w\}, D_G) = \overline{span\{d, \, p(L_w, L_w^*) : d \in D_G, \, p \in \mathbb{C}[z_1, z_2]\}^w},$$

as a topological vector space. Let $w \in loop^c(G)$ be a non-loop finite path. Then $w^k \notin \mathbb{F}^+(G)$, for all $k \in \mathbb{N} \setminus \{1\}$. In other words, if $k \neq 1$, then w^k is not a admissible finite path of the graph G. Thus $L_w^k = L_w^{*k} = 0_{D_G}$, for all $k \in \mathbb{N} \setminus \{1\}$. Therefore, if $q \in \mathbb{C}_1[z_1, z_2]$, then

20

$$q\left(L_{w}, L_{w}^{*}\right) = \begin{cases} q_{1}(L_{w}, L_{w}^{*}) & \text{or} \\ \alpha \in \mathbb{C}, \end{cases}$$

in general, where $q_1 \in \mathbb{C}_1[z_1, z_2]$. i.e.e,

(i) if
$$q(z_1, z_2) = \alpha_0 + \sum_{k=2}^{\infty} (\alpha_k^1 z_1^k + \alpha_k^2 z_2^k)$$
, then
 $q(L_w, L_w^*) = \alpha_0.$

(ii) if $q(z_1, z_2) = \alpha_0 + \sum_{k=1}^{\infty} \left(\alpha_k^1 z_1^k + \alpha_k^2 z_2^k \right)$, then

$$q(L_w, L_w^*) = \alpha_0 + (\alpha_1^1 L_w + \alpha_1^2 L_w^*).$$

So, if we define $q_1 \in \mathbb{C}_1[z_1, z_2]$ by

 $q_1(z_1, z_2) = \alpha_0 + \left(\alpha_1^1 z_1 + \alpha_1^2 z_2\right),$

then

and

$$q\left(L_{w}, L_{w}^{*}\right) = q_{1}\left(L_{w}, L_{w}^{*}\right)$$

3.3. The D_G -Free Product Structure of $(W^*(G), E)$ II..

By Section 3.1 and by (3.1) and (3.2), we can get the following theorem;

Theorem 3.5. Let G be a countable directed graph and $(W^*(G), E)$, the graph W^* -probability space over its diagonal subalgebra D_G . Then

$$\begin{aligned} (W^*(G), E) &= D_G \\ &*_{D_G} \left(*_{D_G} ((W^*(\{L_l\}), tr_l) \otimes (D_G, \mathbf{1})) \right) \\ &*_{D_G} \left(*_{D_G} ((W^*(\{L_w\}, D_w), E_w) \otimes (D_G, \mathbf{1})) \right), \end{aligned}$$

where $tr_l = E_{D_l}^{D_G} \circ E$ is a trace on $W^*(\{L_l\})$ and $E_w = E_{D_w}^{D_G} \circ E$ is a conditional expectation from $W^*(G)$ onto D_w (E_B^A means the conditional expectation from A onto B) and

 $D_{l} = \overline{\mathbb{C}[\{L_{v}: l = vlv\}]}^{w} = \mathbb{C}$ $D_{w} = \overline{\mathbb{C}[\{L_{v_{1}}, L_{v_{2}}: w = v_{1}wv_{2}\}]}^{w} = \Delta_{2}.$

4. More About the Free Product Structure of $(W^*(G), E)$

In this chapter, we will complete to observe the free product structure of the graph W^* -probability spaces. Let G be a countable directed graph and let $(W^*(G), E)$ be the graph W^* -probability space over its diagonal subalgebra D_G . In Chapter 3, we showed that

$$(W^{*}(G), E) = (D_{G}, \mathbf{1})$$

$$*_{D_{G}} \left(*_{D_{G}} ((W^{*}(\{L_{l}\}), tr) \otimes (D_{G}, \mathbf{1})) \right)$$

$$*_{D_{G}D_{G}} \left(*_{D_{G}} ((W^{*}(\{L_{w}\}, D_{w}), E_{w}) \otimes (D_{G}, \mathbf{1})) \right).$$

In the previous chapter, we emphasize the roles of free building blocks and tried to consider each free building block. By using the characterization of the free building blocks, we could get the above free product structure of the graph W^* -probability space $(W^*(G), E)$. Without considering the structure of each free building blocks of $(W^*(G), E)$, by Section 3.1, we can rewrite the above formula as

$$(W^{*}(G), E) = (D_{G}, E)$$

$$*_{D_{G}} \begin{pmatrix} *_{D_{G}} \\ l \in Loop(G) \end{pmatrix} (W^{*}(\{L_{l}\}, D_{G}), E) \end{pmatrix}$$

$$*_{D_{G}} \begin{pmatrix} *_{D_{G}} \\ w \in loop^{c}(G) \end{pmatrix} (W^{*}(\{L_{w}\}, D_{G}), E) \end{pmatrix}.$$

In this chapter, we will show that (4.2)

$$(W^{*}(G), E) = (D_{G}, E)$$

$$*_{D_{G}} \left(*_{D_{G}}_{l \in ELoop(G)} (W^{*}(\{L_{l}\}, D_{G}), E) \right)$$

$$*_{D_{G}} \left(*_{D_{G}}_{w \in Eloop^{c}(G)} (W^{*}(\{L_{w}\}, D_{G}), E) \right).$$

where

$$ELoop(G) \stackrel{def}{=} E(G) \cap Loop(G)$$

and

$$Eloop^{c}(G) \stackrel{def}{=} E(G) \cap loop^{c}(G).$$

Equivalently, we will show that

(4.3)
$$(W^*(G), E) = (D_G, \mathbf{1}) *_{D_G} \left(*_{D_G}_{e \in E(G)} (W^*(\{L_e\}, D_G), E) \right).$$

First, we will concentrate on proving the formula (4.1) is equivalent to the formula (4.2). For the convenience, define

$$\mathcal{L}_{edge}(G) \stackrel{def}{=} *_{D_G} W^*(\{L_l\}, D_G),$$

$$\mathcal{L}^{c}_{edge}(G) \stackrel{def}{=} *_{D_{G}} W^{*}(\{L_{w}\}, D_{G}),$$

$$\mathcal{L}(G) \stackrel{def}{=} *_{D_G} W^*(\{L_l\}, D_G),$$

and

$$\mathcal{L}^{c}(G) \stackrel{def}{=} *_{D_{G}} W^{*}(\{L_{w}\}, D_{G}).$$

Theorem 4.1. Let G be a countable directed graph and let $(W^*(G), E)$ be the graph W^* -probability space over its diagonal subalgebra D_G . Then

$$(W^*(G), E) = (D_G, E) *_{D_G} \left(\mathcal{L}_{edge}(G), E \right) *_{D_G} \left(\mathcal{L}^c_{edge}(G), E \right).$$

Proof. Let $ELoop(G) = E(G) \cap Loop(G)$ and $Eloop^{c}(G) = E(G) \cap loop^{c}(G)$. By (4.1),

$$(W^*(G), E) = (D_G, E) *_{D_G} (\mathcal{L}(G), E) *_{D_G} (\mathcal{L}^c(G), E)$$

Since $Eloop(G) \subset Loop(G)$ and $Eloop^{c}(G) \subset loop^{c}(G)$, we have that

 $\mathcal{L}_{edge}(G) \leq \mathcal{L}(G)$ and $\mathcal{L}^{c}_{edge}(G) \leq \mathcal{L}^{c}(G).$

Therefore, we have the following subalgebra inclusion " \geq ";

(4.4)
$$(W^*(G), E) \ge (D_G, E) *_{D_G} (\mathcal{L}_{edge}(G), E) *_{D_G} (\mathcal{L}^c_{edge}(G), E).$$

So, it suffices to show that we have the reverse subalgebra inclusion " \leq ".

(Case I) Assume that $l \in Loop(G)$. If $l \in ELoop(G)$, then

$$(4.5) W^*(\{L_l\}, D_G) < \mathcal{L}_{edge}(G).$$

If $l = e_1, ..., e_n$ with $e_1, ..., e_n \in E(G)$, such that $e_j \in Eloop^c(G)$, for all j = 1, ..., n, n > 1, then we have that

$$W^*(\{L_l\}, D_G) \le *_{D_G}^n W^*(\{L_{e_j}\}, D_G).$$

 $j=1$

Therefore, by the assumption that $e_i \in Eloop^c(G)$, we have that

(4.6)
$$W^*\left(\{L_l\}, D_G\right) < \mathcal{L}^c_{edge}(G).$$

By (4.5) and (4.6), we can conclude that if $l \in Loop(G)$, then the subalgebra $W^*(\{L_l\}, D_G)$ of $W^*(G)$ is the subalgebra of $\mathcal{L}_{edge}(G) *_{D_G} \mathcal{L}^c_{edge}(G)$. i.e.,

$$(4.7) \quad (W^*(\{L_l\}, D_G)) \le (\mathcal{L}_{edge}(G), E) *_{D_G} \left(\mathcal{L}^c_{edge}(G), E \right),$$

for all $l \in Loop(G)$. Therefore,

(4.8)
$$(\mathcal{L}(G), E) \leq (\mathcal{L}_{edge}(G), E) *_{D_G} \left(\mathcal{L}^c_{edge}(G), D_G \right).$$

(Case II) Now, assume that $w \in loop^{c}(G)$. Suppose that $w \in Eloop^{c}(G)$. Then, clearly,

(4.9)
$$W^*(\{L_w\}, D_G) < \mathcal{L}^c_{edge}(G).$$

Now, assume that $w = e_1 \dots e_k \in loop^c(G)$ with $e_1, \dots, e_k \in E(G)$ are edges satisfying that the initial vertex of e_1 and the final vertex of e_k are different. Then

(4.10)
$$W^*(\{L_w\}, D_G) \le \underset{j=1}{\overset{k}{*} D_G} W^*(\{L_{e_j}\}, D_G).$$

This also shows that

(4.11)
$$W^*\left(\{L_w\}, D_G\right) < \mathcal{L}^c_{edge}(G).$$

By (4.10) and (4.11), we can conclude that if $w \in loop^c(G)$, then the subalgebra $W^*(\{L_w\}, D_G)$ of $W^*(G)$ is the subalgebra of $\mathcal{L}^c_{edge}(G)$. i.e.,

(4.12)
$$(W^*(\{L_w\}, D_G), E) \le \left(\mathcal{L}^c_{edge}(G), E\right),$$

for all $w \in loop^{c}(G)$. Therefore, we have that

(4.13)
$$(\mathcal{L}^{c}(G), E) \leq \left(\mathcal{L}^{c}_{edge}(G), E\right).$$

As we considered in the previous two cases, we can get that

$$(4.14) \qquad \mathcal{L}(G) *_{D_G} \mathcal{L}^c(G) \le \mathcal{L}_{edge}(G) *_{D_G} \mathcal{L}^c_{edge}(G).$$

Therefore, by the relation (4.14), we can conclude that

$$(W^*(G), E) \le (D_G, E) *_{D_G} (\mathcal{L}_{edge}(G), E) *_{D_G} (\mathcal{L}^c_{edge}(G), E)$$

The above theorem provides us that

$$\begin{aligned} (W^*(G), E) &= (D_G, E) \\ &*_{D_G} \left(\underset{l \in E(G) \cap Loop(G)}{*_{D_G}} (W^*(\{L_l\}, D_G), E) \right) \\ &*_{D_G} \left(\underset{w \in E(G) \cap loop^c(G)}{*_{D_G}} (W^*(\{L_w\}, D_G), E) \right) \end{aligned}$$

Therefore, we can get the following simple free product structure of a graph W^* -probability space $(W^*(G), E)$;

Corollary 4.2. Let G be a countable directed graph and let $(W^*(G), E)$ be the graph W^* -probability space over its diagonal subalgebra D_G . Then $(W^*(G), E)$ has the following free product structure ;

$$(W^*(G), E) = (D_G, E) *_{D_G} \left(*_{D_G}_{e \in E(G)} (W^*(\{L_e\}, D_G), E) \right).$$

Proof. Notice that the edge set E(G) of the graph G satisfies that

 $E(G) = (E(G) \cap Loop(G)) \cup (Eloop^{c}(G)).$

Assume that if $e \in E(G)$ is a loop-edge in loop(G), then e is a basic loop in Loop(G). i.e.e,

$$E(G) \cap Loop(G) = Eloop(G).$$

Thus we have that

$$E(G) = (E(G) \cap Loop(G)) \cup (E(G) \cap loop^{c}(G))$$
$$= (E(G) \cap loop(G)) \cup (E(G) \cap loop^{c}(G))$$
$$= E(G) \cap (loop(G) \cup loop^{c}(G))$$
$$= E(G) \cap FP(G) = E(G).$$

The above theorem and corollary shows us that the free product structure of a graph W^* -probability space $(W^*(G), E)$ is totally depending on the admissibility on edges of the graph G.

In the rest of this section, we will consider the several examples ;

Example 4.1. Let G be a directed one-vertex graph with $V(G) = \{v\}$ and $E(G) = \{l_1, ..., l_N\}$, where l_j 's are all loop edges, j = 1, ..., N. Then, by the previous theorem, we can have that

$$(W^*(G), E) = (D_G, E) *_{D_G} \left(\sum_{j=1}^N (W^*(\{L_{e_j}\}, D_G), E) \right).$$

Notice that $D_G = \mathbb{C}$ and E = tr, where tr is the canonical trace, induced by the given conditional expectation. Therefore,

$$(W^*(G), tr) = *_{D_G}^N \left(W^*(\{L_{e_j}\}), tr \right).$$

Trivially, it contains the free group factor $(L(F_N), \tau)$, where $\tau = tr \mid_A$, where

$$A = W^* \left(\{ L_{e_j} + L_{e_j}^* : j = 1, ..., N \} \right).$$

Example 4.2. Suppose that we have a directed graph G with

26

$$V(G) = \{v_1, v_2\} \text{ and } E(G) = \{l_1^1, l_2^1, l_2^2, l_3^2, e\},\$$

$$l_j^1 = v_1 l_j^1 v_1, \text{ for } j = 1, 2, \quad e = v_1 e v_2$$

where

and

$$l_j^1 = v_1 l_j^1 v_1$$
, for $j = 1, 2, \quad e = v_1 e^{-1}$
 $l_j^2 = v_2 l_j^2 v_2$, for $j = 1, 2, 3.$

Then $D_G = \Delta_2 \subset M_2(\mathbb{C})$ and

$$(W^{*}(G), E) = D_{G} *_{D_{G}} \left(*_{D_{G}}^{2} \left(W^{*}(\{L_{l_{i}^{1}}\}) \otimes D_{G}, tr \otimes \mathbf{1} \right) \right) \\ *_{D_{G}} \left(*_{D_{G}}^{3} \left(W^{*}(\{L_{l_{j}^{2}}\}) \otimes D_{G}, tr \otimes \mathbf{1} \right) \right) \\ *_{D_{G}} \left((W^{*}(\{L_{e}\}, \Delta_{2}), E_{2}) \otimes (D_{G}, \mathbf{1}) \right).$$

This shows that our graph W^* -probability space $(W^*(G), E)$ contains free group factors

and

$$L(F_2) \le *_{i=1}^2 \left(W^*(\{L_{l_i^1}\}), tr \right)$$
$$L(F_3) \le *_{j=1}^3 \left(W^*(\{L_{l_j^2}\}), tr \right)$$

(See Section 4.1). Notice that these free group factors $L(F_2)$ and $L(F_3)$ are free **over** D_G in $(W^*(G), E)$. Therefore, $(W^*(G), E)$ contains $L(F_2) *_{D_G} L(F_3)$. Remark that

$$L(F_2) *_{D_G} L(F_3) \neq L(F_2) * L(F_3) = L(F_5).$$

Example 4.3. Let C_N be the circular graph with $V(C_N) = \{v_1, ..., v_N\}$ and

$$E(C_N) = \{e_j : e_j = v_j e_j v_{j+1}, \ j = 1, \dots, N-1, \ e_N = v_N e_N v_1\}.$$

Then $D_G = \Delta_N$. Thus we have that

$$Loop(G) = \{l := e_1...e_N\}$$

equivalently,

$$loop^{c}(G) = \{ w \in FP(G) : w \neq l^{k}, k \in \mathbb{N} \}.$$

So, for the canonical conditional expectation E, we have that

$$(W^*(C_N), E) = (\Delta_N, \mathbf{1})$$
$$*_{\Delta_N} (W^*(\{L_l\}) \otimes \Delta_N, E \otimes \mathbf{1})$$

$$*_{\Delta_N} \left(*_{D_G}_{w \in loop^c(G)} (W^*(\{L_w\}, \Delta_2) \otimes \Delta_N, E_2 \otimes \mathbf{1}) \right)$$
$$= (\Delta_N, E) *_{\Delta_N} \left(*_{\Delta_N}^N (W^*(\{L_{e_j}\}, \Delta_N), E) \right).$$

Reference

[1] A. Nica, R-transform in Free Probability, IHP course note, available at www.math.uwaterloo.ca/~anica.

[2] A. Nica and R. Speicher, R-diagonal Pair-A Common Approach to Haar Unitaries and Circular Elements, (1995), www.mast.queensu.ca/~speicher.

[3] B. Solel, You can see the arrows in a Quiver Operator Algebras, (2000), preprint

[4] A. Nica, D. Shlyakhtenko and R. Speicher, R-cyclic Families of Matrices in Free Probability, J. of Funct Anal, 188 (2002), 227-271.

[5] D. Shlyakhtenko, Some Applications of Freeness with Amalgamation, J. Reine Angew. Math, 500 (1998), 191-212.

[6] D.Voiculescu, K. Dykemma and A. Nica, Free Random Variables, CRM Monograph Series Vol 1 (1992).

[7] D. Voiculescu, Operations on Certain Non-commuting Operator-Valued Random Variables, Astérisque, 232 (1995), 243-275.

[10] D. Shlyakhtenko, A-Valued Semicircular Systems, J. of Funct Anal, 166 (1999), 1-47.

[10] D.W. Kribs and M.T. Jury, Ideal Structure in Free Semigroupoid Algebras from Directed Graphs, preprint

[10] D.W. Kribs and S.C. Power, Free Semigroupoid Algebras, preprint

[11] I. Cho, Amalgamated Boxed Convolution and Amalgamated R-transform Theory, (2002), preprint.

[12] I. Cho, The Tower of Amalgamated Noncommutative Probability Spaces, (2002), Preprint.

[13] I. Cho, Free Perturbed R-transform Theory, (2003), Preprint.

[14] I. Cho, Compatibility of a Noncommutative Probability Space and a Noncommutative Probability Space with Amalgamation, (2003), Preprint

[15] I. Cho, An Example of Scalar-Valued Moments, Under Compatibility, (2003), Preprint.

[16] I. Cho, Graph W^* -Probability Theory, (2004), Preprint.

[17] I. Cho, Random Variables in Graph W^* -Probability Spaces, (2004), Preprint.

[18] I. Cho, Amalgamated Semicircular Systems in Graph W^* -Probability Spaces, (2004), Preprint.

[19] I. Cho, Amalgamated R-diagonal Pairs, (2004), Preprint.

[20] I. Cho, Compressed Random Variables in Graph $W^{\ast}\text{-}\mathsf{Probability}$ Spaces, (2004), Preprint.

[21] P.Śniady and R.Speicher, Continuous Family of Invariant Subspaces for R-diagonal Operators, Invent Math, 146, (2001) 329-363.

[22] R. Speicher, Combinatorial Theory of the Free Product with Amalgamation and Operator-Valued Free Probability Theory, AMS Mem, Vol 132, Num 627, (1998).

[23] R. Speicher, Combinatorics of Free Probability Theory IHP course note, available at www.mast.queensu.ca/~speicher.

UNIV. OF IOWA, DEP. OF MATH, IOWA CITY, IA, U. S. A *E-mail address*: ilcho@math.uiowa.edu