# Hyperbolic 2-Dimensional Manifolds with 3-Dimensional Automorphism Groups I\*†

#### A. V. Isaev

Let M be a Kobayashi-hyperbolic 2-dimensional complex manifold and  $\operatorname{Aut}(M)$  the group of holomorphic automorphisms of M. We showed earlier that if  $\dim\operatorname{Aut}(M)=3$ , then  $\operatorname{Aut}(M)$ -orbits are closed submanifolds in M of (real) codimension 1 or 2. In this paper we classify all connected Kobayashi-hyperbolic 2-dimensional manifolds with 3-dimensional automorphism groups in the case when every orbit has codimension 1. A classification in the case when codimension 2 orbits are present will appear elsewhere.

# 0 Introduction

If M is a connected n-dimensional Kobayashi-hyperbolic complex manifold, then the group  $\operatorname{Aut}(M)$  of holomorphic automorphisms of M is a (real) Lie group in the compact-open topology, of dimension d(M) not exceeding  $n^2+2n$ , with the maximal value occurring only for manifolds holomorphically equivalent to the unit ball  $B^n \subset \mathbb{C}^n$  [11], [9]. We are interested in describing hyperbolic manifolds with lower values of d(M).

In [7], [4] we completely classified manifolds with  $n^2 \leq d(M) < n^2 + 2n$  (partial classifications for  $d(M) = n^2$  were also obtained in [3] and [10]). Note that for  $d(M) = n^2$  the manifold M may not be homogeneous, which makes this case substantially more difficult than the case  $d(M) > n^2$ , where homogeneity always takes place [9]. A further decrease in d(M) almost immediately leads to unclassifiable cases. Indeed, no reasonable classification exists for n = 2, d(M) = 2, in which case  $d(M) = n^2 - 2$  (observe, for example, that the automorphism group of a generic Reinhardt domain in  $\mathbb{C}^2$  is 2-dimensional). While it is possible that there is some classification for  $n \geq 3$ ,  $d(M) = n^2 - 2$ , as well as for particular pairs n, d(M) with  $d(M) < n^2 - 2$  (see [3] for the special case of Reinhardt domains), the case

<sup>\*</sup>Mathematics Subject Classification: 32Q45, 32M05, 32M10.

 $<sup>^{\</sup>dagger}$  **Keywords and Phrases:** Kobayashi-hyperbolic manifolds, holomorphic automorphism groups.

 $d(M) = n^2 - 1$  is probably the only remaining candidate to investigate for the existence of an explicit classification for every  $n \ge 2$ . It turns out that all hyperbolic manifolds with  $n \ge 2$ ,  $d(M) = n^2 - 1$  indeed can be explicitly described. The case  $n \ge 3$  was treated in [5]. In this paper we deal with the far more difficult case n = 2.

For simplicity we will call connected 2-dimensional hyperbolic manifolds with 3-dimensional automorphism group (2,3)-manifolds. As we will see below, the classification of such manifolds is the richest among all previously obtained classifications. We will work with the group  $G(M) := \operatorname{Aut}(M)^c$ , the connected identity component of Aut(M). Since the G(M)-action on M is proper [9], for every  $p \in M$  its isotropy subgroup  $I_p := \{ f \in G(M) : f(p) = \}$ p} is compact in G(M) and the orbit  $O(p) := \{f(p) : f \in G(M)\}$  is a connected closed submanifold in M. It follows from Propositions 1.1 and 4.1 of [5] (see Proposition 1.1 below) that for every  $p \in M$  the orbit O(p) has (real) codimension 1 or 2 in M (in the latter case O(p) is either a complex curve or a totally real submanifold). In this paper we obtain a complete classification of (2,3)-manifolds M under the additional assumption that every G(M)-orbit is of codimension 1, and hence is either strongly pseudoconvex or Levi-flat (see Theorems 2.1 and 3.2). Due to the substantial length of this article we decided to publish our classification of (2,3)-manifolds with codimension 2 orbits elsewhere. We note, however, that the codimension 1 case presented here contains the vast majority of all possible types of (2,3)-manifolds.

The paper is organized as follows. In Section 1 (see Proposition 1.1) we list some basic properties of G(M)-orbits. The most important properties for the purposes of this paper are: every orbit has codimension 1 or 2, and every Levi-flat codimension 1 orbit is foliated by complex curves holomorphically equivalent to the unit disk (that we will always realize as the right half-plane  $\mathcal{P} := \{z \in \mathbb{C} : \operatorname{Re} z > 0\}$ ). The last fact will be extensively used in Section 3.

In Section 2 we deal with the case when every orbit is strongly pseudoconvex and classify all (2,3)-manifolds with this property in Theorem 2.1. An important ingredient in the proof of Theorem 2.1 is E. Cartan's classification of 3-dimensional homogeneous strongly pseudoconvex CR-manifolds [2], together with the explicit determination of all covers of the non-simply connected hypersurfaces on Cartan's list [6].

In Section 3 we first classify Levi-flat orbits together with all possible actions of G(M) (see Proposition 3.1). In the proof of Proposition 3.1 we show, in particular, that if the Lie algebra of G(M) is isomorphic to  $\mathfrak{su}_{1,1}$ ,

then there are no Levi-flat orbits in M. It is interesting to remark, however, that during the course of proof we discovered actions by CR-transformations of groups with Lie algebras isomorphic to  $\mathfrak{su}_{1,1}$ , on the Levi-flat hypersurfaces  $\mathcal{P} \times \mathbb{R}$  and  $\mathcal{P} \times S^1$  in  $\mathbb{C}^2$  that do not extend to actions by holomorphic transformations on any neighborhood of either hypersurface. These actions are of independent interest, and, as far as we are aware, they were previously unknown. Proposition 3.1 is used to classify (2,3)-manifolds with Levi-flat orbits in Theorem 3.2.

#### 1 Initial Classification of Orbits

In this section we will list some initial facts about G(M)-orbits that follow from the results of [5]. For  $p \in M$  let  $L_p := \{d_p f : f \in I_p\}$  be the linear isotropy subgroup of p, where  $d_p f$  is the differential of a map f at p. The group  $L_p$  is a compact subgroup of  $GL(T_p(M), \mathbb{C})$  isomorphic to  $I_p$  by means of the isotropy representation

$$I_p \to L_p, \quad f \mapsto d_p f.$$

Propositions 1.1, and 4.1 of [5] imply the following

**Proposition 1.1** Let M be a (2,3)-manifold. Fix  $p \in M$  and let  $V_p := T_p(O(p))$ . Then the following holds:

- (i) The orbit O(p) is either a closed real hypersurface, or a closed complex curve, or a closed totally real 2-dimensional submanifold of M.
- (ii) If O(p) is a real Levi-flat hypersurface, it is foliated by complex curves holomorphically equivalent to  $\Delta$ , and there exist coordinates on  $T_p(M)$  such with respect to the orthogonal decomposition  $T_p(M) = (V_p \cap iV_p)^{\perp} \oplus (V_p \cap iV_p)$  we have  $L_p \subset \{\pm id\} \times L'_p$ , where  $L'_p$  is a finite subgroup of  $U_1$ .
- (iii) If O(p) is a complex curve, it is holomorphically equivalent to  $\Delta$ . Furthermore, there exist holomorphic coordinates (z, w) on  $T_p(M)$  in which  $V_p = \{z = 0\}$  and the identity component  $L_p^c$  of  $L_p$  is given by either the matrices

$$\left(\begin{array}{cc} a^{\frac{k_1}{k_2}} & 0\\ 0 & a \end{array}\right),\,$$

for some  $k_1, k_2 \in \mathbb{Z}$ ,  $(k_1, k_2) = 1$ ,  $k_2 \neq 0$ , or the matrices

$$\left(\begin{array}{cc} a & 0 \\ 0 & 1 \end{array}\right),\,$$

where |a| = 1.

(iv) If n=2 and O(p) is totally real, then  $T_p(M)=V_p\oplus iV_p$ , and there are coordinates on  $V_p$  such that every transformation from  $L_p^c$  has the form:  $v_1+iv_2\mapsto Av_1+iAv_2,\ v_1,v_2\in V_p$ , where  $A\in SO_2(\mathbb{R})$ .

Below we only consider (2,3)-manifolds without codimension 2 orbits.

# 2 The Case of Strongly Pseudoconvex Orbits

In this section we will give a complete classification of (2,3)-manifolds M for which every G(M)-orbit is a strongly pseudoconvex real hypersurface in M. We will start with a large number of examples of such manifolds.

(1) Fix  $b \in \mathbb{R}$  such that  $|b| \geq 1$ ,  $b \neq 1$ , and choose  $0 \leq s < t \leq \infty$ . Define

$$R_{b,s,t} := \left\{ (z, w) \in \mathbb{C}^2 : s \left( \operatorname{Re} z \right)^b < \operatorname{Re} w < t \left( \operatorname{Re} z \right)^b, \operatorname{Re} z > 0 \right\}.$$

All these domains are obviously hyperbolic and the group  $\operatorname{Aut}(R_{b,s,t}) = G(R_{b,s,t})$  consists of all the maps

$$\begin{array}{ccc}
z & \mapsto & \lambda z + i\beta, \\
w & \mapsto & \lambda^b w + i\gamma,
\end{array}$$
(2.1)

where  $\lambda > 0$  and  $\beta, \gamma \in \mathbb{R}$ . The  $G(R_{b,s,t})$ -orbits are the following pairwise CR-equivalent hypersurfaces:

$$O_{\alpha}^{R_b} := \left\{ (z, w) \in \mathbb{C}^2 : \operatorname{Re} w = \alpha \left( \operatorname{Re} z \right)^b, \operatorname{Re} z > 0 \right\}, \quad s < \alpha < t,$$

and we set

$$\tau_b := O_1^{R_b}. \tag{2.2}$$

(2) For  $0 \le s < t \le \infty$  with either s > 0 or  $t < \infty$  define

$$U_{s,t} := \left\{ (z, w) \in \mathbb{C}^2 : \operatorname{Re} w \cdot \ln \left( s \operatorname{Re} w \right) < \operatorname{Re} z < \operatorname{Re} w \cdot \ln \left( t \operatorname{Re} w \right), \, \operatorname{Re} w > 0 \right\}.$$

All these domains are clearly hyperbolic and the group  $\operatorname{Aut}(U_{s,t}) = G(U_{s,t})$  consists of all the maps

$$z \mapsto \lambda z + (\lambda \ln \lambda)w + i\beta, w \mapsto \lambda w + i\gamma,$$
 (2.3)

where  $\lambda > 0$  and  $\beta, \gamma \in \mathbb{R}$ . The  $G(U_{s,t})$ -orbits are the following pairwise CR-equivalent hypersurfaces:

$$O_{\alpha}^{U} := \left\{ (z,w) \in \mathbb{C}^{2} : \operatorname{Re}z = \operatorname{Re}w \cdot \ln\left(\alpha \operatorname{Re}w\right), \, \operatorname{Re}w > 0 \right\}, \quad s < \alpha < t,$$

and we set

$$\xi := O_1^U. \tag{2.4}$$

(3) For  $0 \le s < t < \infty$  define

$$\mathfrak{S}_{s,t} := \{(z, w) \in \mathbb{C}^2 : s < (\operatorname{Re} z)^2 + (\operatorname{Re} w)^2 < t\}.$$

All these domains are clearly hyperbolic and the group  $\operatorname{Aut}(\mathfrak{S}_{s,t})$  consists of all maps of the form

$$\begin{pmatrix} z \\ w \end{pmatrix} \mapsto A \begin{pmatrix} z \\ w \end{pmatrix} + i \begin{pmatrix} \beta \\ \gamma \end{pmatrix}, \tag{2.5}$$

where  $A \in O_2(\mathbb{R})$  and  $\beta, \gamma \in \mathbb{R}$ . The subgroup  $G(\mathfrak{S}_{s,t})$  is given by the condition  $A \in SO_2(\mathbb{R})$ . The  $G(\mathfrak{S}_{s,t})$ -orbits are the following pairwise CR-equivalent hypersurfaces:

$$O_{\alpha}^{\mathfrak{S}} := \left\{ (z, w) \in \mathbb{C}^2 : (\operatorname{Re} z)^2 + (\operatorname{Re} w)^2 = \alpha \right\}, \quad s < \alpha < t,$$

and we set

$$\chi := O_1^{\mathfrak{S}}. \tag{2.6}$$

(4) For more details on this example see [6]. Let  $\Phi_{\chi}^{(\infty)}: \mathbb{C}^2 \to \mathbb{C}^2$  be the following map:

$$z \mapsto \exp(\operatorname{Re} z)\cos(\operatorname{Im} z) + i\operatorname{Re} w,$$
  
 $w \mapsto \exp(\operatorname{Re} z)\sin(\operatorname{Im} z) + i\operatorname{Im} w.$ 

It is easy to see that  $\Phi_{\chi}^{(\infty)}$  is an infinitely-sheeted covering map onto  $\mathbb{C}^2 \setminus \{\text{Re } z = 0, \text{ Re } w = 0\}$ . Introduce on the domain of  $\Phi_{\chi}^{(\infty)}$  a complex structure defined by the condition that the map  $\Phi_{\chi}^{(\infty)}$  is holomorphic (the pull-back complex structure under  $\Phi_{\chi}^{(\infty)}$ ), and denote the resulting manifold by  $M_{\chi}^{(\infty)}$ .

Next, for an integer  $n \geq 2$  consider the map  $\Phi_{\chi}^{(n)}$  from  $\mathbb{C}^2 \setminus \{\operatorname{Re} z = 0, \operatorname{Re} w = 0\}$  onto itself defined as follows:

$$z \mapsto \operatorname{Re}\left(\left(\operatorname{Re}z + i\operatorname{Re}w\right)^{n}\right) + i\operatorname{Im}z,$$
  
 $w \mapsto \operatorname{Im}\left(\left(\operatorname{Re}z + i\operatorname{Re}w\right)^{n}\right) + i\operatorname{Im}w.$ 

Denote by  $M_{\chi}^{(n)}$  the domain of  $\Phi_{\chi}^{(n)}$  with the pull-back complex structure under  $\Phi_{\chi}^{(n)}$ .

For  $0 \le s < t < \infty$  define

$$\mathfrak{S}_{s,t}^{(\infty)} := \left\{ (z, w) \in M_{\chi}^{(\infty)} : (\ln s)/2 < \operatorname{Re} z < (\ln t)/2 \right\},$$

$$\mathfrak{S}_{s,t}^{(n)} := \left\{ (z, w) \in M_{\chi}^{(n)} : s^{1/n} < (\operatorname{Re} z)^2 + (\operatorname{Re} w)^2 < t^{1/n} \right\}, \quad n \ge 2.$$

The domains  $\mathfrak{S}_{s,t}^{(\infty)}$  and  $\mathfrak{S}_{s,t}^{(n)}$  are respectively an infinite- and n-sheeted cover of the domain  $\mathfrak{S}_{s,t}$  introduced in (3) and hence are hyperbolic.

The group  $\operatorname{Aut}\left(\mathfrak{S}_{s,t}^{(\infty)}\right)$  is generated by  $G\left(\mathfrak{S}_{s,t}^{(\infty)}\right)$  that consists of the maps

$$\begin{array}{ccc}
z & \mapsto & z + i\beta, \\
w & \mapsto & w + a,
\end{array}$$
(2.7)

where  $\beta \in \mathbb{R}$ ,  $a \in \mathbb{C}$ , and the map

$$\begin{array}{ccc}
z & \mapsto & \overline{z}, \\
w & \mapsto & \overline{w}
\end{array} \tag{2.8}$$

The  $G\left(\mathfrak{S}_{s,t}^{(\infty)}\right)$ -orbits are the following pairwise CR-equivalent hypersurfaces:

$$O_{\alpha}^{\mathfrak{S}^{(\infty)}} := \{(z, w) \in M_{\chi}^{(\infty)} : \operatorname{Re} z = \alpha\}, \quad (\ln s)/2 < \alpha < (\ln t)/2,$$

and we set

$$\chi^{(\infty)} := O_0^{\mathfrak{S}^{(\infty)}}.\tag{2.9}$$

The group Aut  $\left(\mathfrak{S}_{s,t}^{(n)}\right)$  for  $n \geq 2$  is generated by  $G\left(\mathfrak{S}_{s,t}^{(n)}\right)$  that consists of the maps

$$z \mapsto \cos \varphi \cdot \operatorname{Re} z + \sin \varphi \cdot \operatorname{Re} w + i \Big( \cos(n\varphi) \cdot \operatorname{Im} z + \sin(n\varphi) \cdot \operatorname{Im} w + \beta \Big),$$

$$w \mapsto -\sin \varphi \cdot \operatorname{Re} z + \cos \varphi \cdot \operatorname{Re} w + i \Big( -\sin(n\varphi) \cdot \operatorname{Im} z + \cos(n\varphi) \cdot \operatorname{Im} w + \gamma \Big),$$

$$(2.10)$$

where  $\varphi, \beta, \gamma \in \mathbb{R}$ , and the map

$$\begin{array}{ccc}
z & \mapsto & z, \\
w & \mapsto & -w.
\end{array}$$
(2.11)

The  $G\left(\mathfrak{S}_{s,t}^{(n)}\right)$ -orbits are the following pairwise CR-equivalent hypersurfaces:

$$O_{\alpha}^{\mathfrak{S}^{(n)}} := \{ (z, w) \in M_{\chi}^{(n)} : (\operatorname{Re} z)^2 + (\operatorname{Re} w)^2 = \alpha \}, \quad s^{1/n} < \alpha < t^{1/n},$$

and we set

$$\chi^{(n)} := O_1^{\mathfrak{S}^{(n)}}. \tag{2.12}$$

(5) Fix b > 0 and for  $0 < t < \infty$ ,  $e^{-2\pi b}t < s < t$  consider

$$V_{b,s,t} := \{(z, w) \in \mathbb{C}^2 : se^{b\varphi} < r < te^{b\varphi} \},$$

where  $(r, \varphi)$  denote the polar coordinates in the (Re z, Re w)-plane with  $\varphi$  varying from  $-\infty$  to  $\infty$  (thus, the boundary of  $V_{b,t,s}$  consists of two spirals accumulating to the origin and infinity). All these domains are hyperbolic and Aut $(V_{b,s,t}) = G(V_{b,s,t})$  consists of all maps of the form

$$\begin{pmatrix} z \\ w \end{pmatrix} \mapsto e^{b\psi} \begin{pmatrix} \cos\psi & \sin\psi \\ -\sin\psi & \cos\psi \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix} + i \begin{pmatrix} \beta \\ \gamma \end{pmatrix}, \tag{2.13}$$

where  $\psi, \beta, \gamma \in \mathbb{R}$ . The  $G(V_{b,s,t})$ -orbits are the following pairwise CR-equivalent hypersurfaces:

$$O_{\alpha}^{V_b} := \left\{ (z, w) \in \mathbb{C}^2 : r = \alpha e^{b\varphi} \right\}, \quad s < \alpha < t,$$

and we set

$$\rho_b := O_1^{V_b}. \tag{2.14}$$

(6) For  $1 \le s < t < \infty$  define

$$E_{s,t} := \left\{ (z:w:\zeta) \in \mathbb{CP}^2 : s|z^2 + w^2 + \zeta^2| < |z|^2 + |w|^2 + |\zeta|^2 < t|z^2 + w^2 + \zeta^2| \right\},$$

The domain  $E_{s,t}$  is hyperbolic for each t since it is covered in a 2-to-1 fashion by the manifold

$$\Big\{(z,w,\zeta)\in\mathbb{C}^3: s<|z|^2+|w|^2+|\zeta|^2< t,\ z^2+w^2+\zeta^2=1\Big\},$$

which is clearly hyperbolic (the covering map is  $(z, w, \zeta) \mapsto (z : w : \zeta)$ ). The group  $\operatorname{Aut}(E_{s,t}) = G(E_{s,t})$  is given by

$$\begin{pmatrix} z \\ w \\ \zeta \end{pmatrix} \mapsto A \begin{pmatrix} z \\ w \\ \zeta \end{pmatrix}, \tag{2.15}$$

where  $A \in SO_3(\mathbb{R})$ . The orbits of the action of the group  $G(E_{s,t})$  on  $E_{s,t}$  are the following pairwise CR non-equivalent hypersurfaces:

$$\mu_{\alpha} := \{ (z : w : \zeta) \in \mathbb{CP}^2 : |z|^2 + |w|^2 + |\zeta|^2 = \alpha |z^2 + w^2 + \zeta^2| \}, \quad (2.16)$$

(7) For more details on this example see [6]. Let  $\mathcal{Q}_+$  be the variety in  $\mathbb{C}^3$  given by

$$z_1^2 + z_2^2 + z_3^2 = 1.$$

Consider the map  $\Phi_{\mu}: \mathbb{C}^2 \setminus \{0\} \to \mathcal{Q}_+$  defined by the formulas

$$z_1 = -i(z^2 + w^2) + i\frac{z\overline{w} - w\overline{z}}{|z|^2 + |w|^2},$$

$$z_2 = z^2 - w^2 - \frac{z\overline{w} + w\overline{z}}{|z|^2 + |w|^2},$$

$$z_3 = 2zw + \frac{|z|^2 - |w|^2}{|z|^2 + |w|^2}.$$

This map was first introduced in [12]. It is straightforward to verify that  $\Phi_{\mu}$  is a 2-to-1 covering map onto  $\mathcal{Q}_{+} \setminus \mathbb{R}^{3}$ . Introduce on the domain of  $\Phi_{\mu}$  the pull-back complex structure under  $\Phi_{\mu}$  and denote the resulting complex manifold by  $M_{\mu}^{(4)}$ .

For  $1 \le s < t < \infty$  define

$$E_{s,t}^{(2)} := \{(z_1, z_2, z_3) \in \mathbb{C}^3 : s < |z_1|^2 + |z_2|^2 + |z_3|^2 < t\} \cap \mathcal{Q}_+,$$

$$E_{s,t}^{(4)} := \{(z, w) \in M_{\mu}^{(4)} : \sqrt{(s-1)/2} < |z|^2 + |w|^2 < \sqrt{(t-1)/2}\}.$$

These domains are respectively 2- and 4-sheeted covers of the domain  $E_{s,t}$  introduced in (6) and hence are hyperbolic.

The group Aut  $\left(E_{s,t}^{(2)}\right)$  consists of the maps

$$\begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix} \mapsto A \begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix}, \tag{2.17}$$

where  $A \in O_3(\mathbb{R})$ , and  $G\left(E_{s,t}^{(2)}\right)$  is given by the condition  $A \in SO_3(\mathbb{R})$ . The  $G\left(E_{s,t}^{(2)}\right)$ -orbits are the following pairwise non-equivalent hypersurfaces:

$$\mu_{\alpha}^{(2)} := \{(z_1, z_2, z_3) \in \mathbb{C}^3 : |z_1|^2 + |z_2|^2 + |z_3|^2 = \alpha\} \cap \mathcal{Q}_+, \ s < \alpha < t. \ (2.18)$$

The group Aut  $\left(E_{s,t}^{(4)}\right)$  is generated by  $G\left(E_{s,t}^{(4)}\right)$  that consists of the maps

$$\begin{pmatrix} z \\ w \end{pmatrix} \mapsto A \begin{pmatrix} z \\ w \end{pmatrix}, \tag{2.19}$$

where  $A \in SU_2$ , and the map

$$z \mapsto i \frac{z(|z|^2 + |w|^2) - \overline{w}}{\sqrt{1 + (|z|^2 + |w|^2)^2}},$$

$$w \mapsto i \frac{w(|z|^2 + |w|^2) + \overline{z}}{\sqrt{1 + (|z|^2 + |w|^2)^2}}.$$
(2.20)

The  $G\left(E_{s,t}^{(4)}\right)$ -orbits are the following pairwise CR non-equivalent hypersurfaces:

$$\mu_{\alpha}^{(4)} := \left\{ (z, w) \in M_{\mu}^{(4)} : |z|^2 + |w|^2 = \sqrt{(\alpha - 1)/2} \right\}, \quad s < \alpha < t. \quad (2.21)$$

(8) For  $-1 \le s < t \le 1$  define

$$\Omega_{s,t} := \left\{ (z, w) \in \mathbb{C}^2 : s|z^2 + w^2 - 1| < |z|^2 + |w|^2 - 1 < t|z^2 + w^2 - 1| \right\}.$$

Clearly,  $\Omega_{s,t}$  is bounded if t < 1. Further,  $\Omega_{s,1}$  is hyperbolic since it is contained in the hyperbolic product domain

$$\left\{(z,w)\in\mathbb{C}^2:z,w\not\in(-\infty,-1]\cup[1,\infty)\right\}.$$

The group  $\operatorname{Aut}(\Omega_{s,t})$  for every t consists of the maps

$$\begin{pmatrix} z \\ w \end{pmatrix} \mapsto \frac{\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}}{c_1 z + c_2 w + d}, \tag{2.22}$$

where

$$Q := \begin{pmatrix} a_{11} & a_{12} & b_1 \\ a_{21} & a_{22} & b_2 \\ c_1 & c_2 & d \end{pmatrix} \in SO_{2,1}(\mathbb{R}). \tag{2.23}$$

The group  $\operatorname{Aut}(\Omega_{s,t})$  has two connected components (that correspond to the connected components of  $SO_{2,1}(\mathbb{R})$ ), and its identity component  $G(\Omega_{s,t})$  is given by the condition  $a_{11}a_{22} - a_{12}a_{21} > 0$ . The orbits of  $G(\Omega_{s,t})$  on  $\Omega_{s,t}$  are the following pairwise CR non-equivalent hypersurfaces:

$$\nu_{\alpha} := \{ (z, w) \in \mathbb{C}^2 : |z|^2 + |w|^2 - 1 = \alpha |z^2 + w^2 - 1| \} \setminus \{ (x, u) \in \mathbb{R}^2 : x^2 + u^2 = 1 \}, \quad s < \alpha < t.$$
 (2.24)

(9) For  $1 \le s < t \le \infty$  define

$$D_{s,t} := \left\{ (z, w) \in \mathbb{C}^2 : s|1 + z^2 - w^2| < 1 + |z|^2 - |w|^2 < t|1 + z^2 - w^2|, \right.$$
$$\left. \operatorname{Im} \left( z(1 + \overline{w}) \right) > 0 \right\}.$$

All these domains lie in the hyperbolic product domain

$$\{(z, w) \in \mathbb{C}^2 : \operatorname{Im} z > 0, \ w \notin (-\infty, -1] \cup [1, \infty)\},\$$

hence they are hyperbolic. For every matrix  $Q \in SO_{2,1}(\mathbb{R})^c$  as in (2.23) consider the map

$$\begin{pmatrix} z \\ w \end{pmatrix} \mapsto \frac{\begin{pmatrix} a_{22} & b_2 \\ c_2 & d \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix} + \begin{pmatrix} a_{21} \\ c_1 \end{pmatrix}}{a_{12}z + b_1w + a_{11}}.$$
 (2.25)

The group  $\operatorname{Aut}(D_{s,t}) = G(D_{s,t})$  consists of all such maps. The orbits of  $G(D_{s,t})$  on  $D_{s,t}$  are the following pairwise CR non-equivalent hypersurfaces:

$$\eta_{\alpha} := \left\{ (z, w) \in \mathbb{C}^2 : 1 + |z|^2 - |w|^2 = \alpha |1 + z^2 - w^2|, \\
\operatorname{Im}(z(1 + \overline{w})) > 0 \right\}, \quad s < \alpha < t.$$
(2.26)

(10) For more details on this example see [6]. Let  $\mathcal{Q}_{-}$  be the variety in  $\mathbb{C}^{3}$  given by

$$z_1^2 + z_2^2 - z_3^2 = 1,$$

and let

$$\Sigma := \{(z, w) \in \mathbb{C}^2 : |z|^2 - |w|^2 > 0\}.$$

For every integer  $n \geq 2$  consider the map  $\Phi^{(n)}$  from  $\Sigma$  to  $\mathcal{Q}_-$  defined as follows:

$$z_1 = -i(z^n + z^{n-2}w^2) - i\frac{z\overline{w} + w\overline{z}}{|z|^2 - |w|^2},$$

$$z_2 = z^n - z^{n-2}w^2 + \frac{z\overline{w} - w\overline{z}}{|z|^2 - |w|^2},$$

$$z_3 = -2iz^{n-1}w - i\frac{|z|^2 + |w|^2}{|z|^2 - |w|^2}.$$

The above maps were introduced in [6]. Further, set

$$\Omega_{\nu}^{(n)} := \{(z, w) \in \mathbb{C}^2 : 0 < |z|^n - |z|^{n-2}|w|^2 < 1\}, 
\Omega_{\eta}^{(n)} := \{(z, w) \in \mathbb{C}^2 : |z|^n - |z|^{n-2}|w|^2 > 1\}.$$

Clearly,  $\Omega_{\nu}^{(n)} \subset \Sigma$  and  $\Omega_{\eta}^{(n)} \subset \Sigma$  for all  $n \geq 2$ . Let  $\Phi_{\nu}^{(n)}$  and  $\Phi_{\eta}^{(n)}$  be the restrictions of  $\Phi^{(n)}$  to  $\Omega_{\nu}^{(n)}$  and  $\Omega_{\eta}^{(n)}$ , respectively. A straightforward calculation

shows that  $\Phi_{\nu}^{(n)}$  and  $\Phi_{\eta}^{(n)}$  are n-to-1 covering maps onto

$$\Omega_{\nu} := \left\{ (z_1, z_2, z_3) \in \mathbb{C}^3 : -1 < |z_1|^2 + |z_2|^2 - |z_3|^2 < 1, \right.$$

$$\operatorname{Im} z_3 < 0 \right\} \cap \mathcal{Q}_{-}$$

and

$$\Omega_{\eta} := \left\{ (z_1, z_2, z_3) \in \mathbb{C}^3 : |z_1|^2 + |z_2|^2 - |z_3|^2 > 1, \\
\operatorname{Im}(z_2(\overline{z_1} + \overline{z_3})) > 0 \right\} \cap \mathcal{Q}_-,$$

respectively. We now introduce on  $\Omega_{\nu}^{(n)}$ ,  $\Omega_{\eta}^{(n)}$  the pull-back complex structures under the maps  $\Phi_{\nu}^{(n)}$ ,  $\Phi_{\eta}^{(n)}$ , respectively, and denote the resulting complex manifolds by  $M_{\nu}^{(n)}$ ,  $M_{\eta}^{(n)}$ .

Further, let  $\Lambda: \overset{\cdot}{\mathbb{C}\times \Delta} \to \Sigma$  be the following covering map:

$$\begin{array}{ccc} z & \mapsto & e^z, \\ w & \mapsto & e^z w, \end{array}$$

where  $z \in \mathbb{C}$ ,  $w \in \Delta$  and  $\Delta$  is the unit disk. Define

$$\begin{array}{lcl} U_{\nu} &:= & \{(z,w) \in \mathbb{C}^2 : |w| < 1, \ \exp(2\operatorname{Re}z)(1-|w|^2) < 1\}, \\ U_{\eta} &:= & \{(z,w) \in \mathbb{C}^2 : |w| < 1, \ \exp(2\operatorname{Re}z)(1-|w|^2) > 1\}. \end{array}$$

Denote by  $\Lambda_{\nu}$ ,  $\Lambda_{\eta}$  the restrictions of  $\Lambda$  to  $U_{\nu}$ ,  $U_{\eta}$ , respectively. Clearly,  $U_{\nu}$  covers  $M_{\nu}^{(2)}$  by means of  $\Lambda_{\nu}$ , and  $U_{\eta}$  covers  $M_{\eta}^{(2)}$  by means of  $\Lambda_{\eta}$ . We now introduce on  $U_{\nu}$ ,  $U_{\eta}$  the pull-back complex structures under the maps  $\Lambda_{\nu}$ ,  $\Lambda_{\eta}$ , respectively, and denote the resulting complex manifolds by  $M_{\nu}^{(\infty)}$ ,  $M_{\eta}^{(\infty)}$ .

For  $-1 \le s < t \le 1$  we now define

$$\Omega_{s,t}^{(n)} := \left\{ (z,w) \in M_{\nu}^{(n)} : \sqrt{(s+1)/2} < |z|^n - |z|^{n-2}|w|^2 < \sqrt{(t+1)/2} \right\}, \quad n \ge 2,$$

$$\Omega_{s,t}^{(\infty)} := \left\{ (z,w) \in M_{\nu}^{(\infty)} : \sqrt{(s+1)/2} < \exp\left(\operatorname{Re}z\right)(1-|w|^2) < \sqrt{(t+1)/2} \right\}.$$

The domain  $\Omega_{s,t}^{(n)}$ ,  $n \geq 2$ , is an *n*-sheeted cover of the domain  $\Omega_{s,t}$  introduced in (8) and the domain  $\Omega_{s,t}^{(\infty)}$  is its infinitely-sheeted cover. In particular, each of  $\Omega_{s,t}^{(n)}$ ,  $\Omega_{s,t}^{(\infty)}$  is hyperbolic.

The group Aut  $\left(\Omega_{s,t}^{(n)}\right)$  is generated by  $G\left(\Omega_{s,t}^{(n)}\right)$  that consists of all maps of the form

$$z \mapsto z \sqrt[n]{(a+bw/z)^2},$$

$$w \mapsto z \frac{\overline{b} + \overline{a}w/z}{a+bw/z} \sqrt[n]{(a+bw/z)^2},$$
(2.27)

where  $\sqrt[n]{}$  is any branch of the *n*th root and  $|a|^2 - |b|^2 = 1$ , and the map

$$z \mapsto \overline{z} \left( \sqrt[n]{\frac{\left(1 + z^{n-1}w(1 - |w|^2/|z|^2)\right)^2}{1 - |z|^{2n}(1 - |w|^2/|z|^2)^2}} \right)',$$

$$w \mapsto -\frac{\overline{w}/\overline{z} + z^n(1 - |w|^2/|z|^2)}{1 + z^{n-1}w(1 - |w|^2/|z|^2)} \times$$

$$\overline{z} \left( \sqrt[n]{\frac{\left(1 + z^{n-1}w(1 - |w|^2/|z|^2)\right)^2}{1 - |z|^{2n}(1 - |w|^2/|z|^2)^2}} \right)',$$
(2.28)

for some branch  $\binom{n}{\sqrt{}}'$  of  $\sqrt[n]{}$ . The  $G\left(\Omega_{s,t}^{(n)}\right)$ -orbits are the following pairwise CR non-equivalent hypersurfaces:

$$\nu_{\alpha}^{(n)} := \left\{ (z, w) \in M_{\nu}^{(n)} : |z|^{n} - |z|^{n-2}|w|^{2} = \sqrt{(\alpha + 1)/2} \right\}, \tag{2.29}$$

The group  $\operatorname{Aut}\left(\Omega_{s,t}^{(\infty)}\right)$  is generated by  $G\left(\Omega_{s,t}^{(\infty)}\right)$  that consists of the following maps:

$$z \mapsto z + \ln(a + bw),$$

$$w \mapsto \frac{\overline{b} + \overline{a}w}{a + bw},$$
(2.30)

where ln is any branch of the logarithm and  $|a|^2 - |b|^2 = 1$ , and the map

$$z \mapsto \overline{z} + \ln' \left( -\frac{1 + e^{2z} w (1 - |w|^2)}{\sqrt{1 - \exp(4\operatorname{Re}z)(1 - |w|^2)^2}} \right),$$

$$w \mapsto -\frac{\overline{w} + e^{2z} (1 - |w|^2)}{1 + e^{2z} w (1 - |w|^2)},$$
(2.31)

for some branch  $\ln'$  of the logarithm. The  $G\left(\Omega_{s,t}^{(\infty)}\right)$ -orbits are the following pairwise CR non-equivalent hypersurfaces:

$$\nu_{\alpha}^{(\infty)} := \left\{ (z, w) \in M_{\nu}^{(\infty)} : \exp(2\operatorname{Re} z) (1 - |w|^2) = \sqrt{(\alpha + 1)/2} \right\}, \quad (2.32)$$

Next, for  $1 \le s < t \le \infty$  we define

$$D_{s,t}^{(2)} := \left\{ (z_1, z_2, z_3) \in \mathbb{C}^3 : s < |z_1|^2 + |z_2|^2 - |z_3|^2 < t, \\ \operatorname{Im}(z_2(\overline{z_1} + \overline{z_3})) > 0 \right\} \cap \mathcal{Q}_-,$$

$$D_{s,t}^{(2n)} := \left\{ (z, w) \in M_{\eta}^{(n)} : \sqrt{(s+1)/2} < |z|^n - |z|^{n-2}|w|^2 < \sqrt{(t+1)/2} \right\}, \quad n \ge 2,$$

$$D_{s,t}^{(\infty)} := \left\{ (z, w) \in M_{\eta}^{(\infty)} : \sqrt{(s+1)/2} < \exp\left(\operatorname{Re} z\right) (1 - |w|^2) < \sqrt{(t+1)/2} \right\}.$$

The domain  $D_{s,t}^{(2n)}$ ,  $n \geq 1$ , is an 2n-sheeted cover of the domain  $D_{s,t}$  introduced in (9) and the domain  $D_{s,t}^{(\infty)}$  is its infinitely-sheeted cover. To obtain an n-sheeted cover of  $D_{s,t}$  for odd  $n \geq 3$ , the domain  $D_{s,t}^{(4n)}$  must be factored by the action of the cyclic group of four elements generated by the following

automorphism of  $M_{\eta}^{(2n)}$ :

$$z \mapsto iz^{2}\overline{z} \left( \sqrt[n]{\frac{1 - |w|^{2}/|z|^{2} + z^{-2n}\overline{w}/\overline{z}}{\sqrt{|z|^{4n}(1 - |w|^{2}/|z|^{2})^{2} - 1}}} \right)',$$

$$w \mapsto i\frac{1 + z^{2n-1}w(1 - |w|^{2}/|z|^{2})}{\overline{w}/\overline{z} + z^{2n}(1 - |w|^{2}/|z|^{2})} \times$$

$$z^{2}\overline{z} \left( \sqrt[n]{\frac{1 - |w|^{2}/|z|^{2} + z^{-2n}\overline{w}/\overline{z}}{\sqrt{|z|^{4n}(1 - |w|^{2}/|z|^{2})^{2} - 1}}} \right)',$$

for some branch  $\binom{n}{\sqrt{}}'$  of  $\sqrt[n]{}$ . Let  $\Pi_n$  denote the corresponding factorization map. Then  $D_{s,t}^{(n)} := \Pi_n(D_{s,t}^{(2n)})$  is an n-sheeted cover of  $D_{s,t}$ . Clearly, each of  $D_{s,t}^{(n)}$ ,  $D_{s,t}^{(\infty)}$  is hyperbolic.

The group  $G\left(D_{s,t}^{(2)}\right)$  consists of all maps of the form (2.17) with  $A \in SO_{2,1}(\mathbb{R})^c$ , and the full group  $\operatorname{Aut}\left(D_{s,t}^{(2)}\right)$  is generated by  $G\left(D_{s,t}^{(2)}\right)$  and the map

$$z_1 \mapsto -z_1,$$

$$z_2 \mapsto -z_2,$$

$$z_3 \mapsto -z_3.$$

$$(2.33)$$

The  $G\left(D_{s,t}^{(2)}\right)$ -orbits are the following pairwise non-equivalent hypersurfaces:

$$\eta_{\alpha}^{(2)} := \left\{ (z_1, z_2, z_3) \in \Omega_{\eta} : |z_1|^2 + |z_2|^2 - |z_3|^2 = \sqrt{(\alpha + 1)/2}, \right\}, \quad (2.34)$$

For  $n \geq 2$  the group  $G\left(D_{s,t}^{(2n)}\right)$  consists of all maps of the form (2.27),

and the full group Aut  $\left(D_{s,t}^{(2n)}\right)$  is generated by  $G\left(D_{s,t}^{(2n)}\right)$  and the map

$$z \mapsto z^2 \overline{z} \left( \sqrt[n]{ \frac{\left(1 - |w|^2/|z|^2 + z^{-n} \overline{w}/\overline{z}\right)^2}{|z|^{2n} (1 - |w|^2/|z|^2)^2 - 1}} \right)',$$

$$w \mapsto \frac{1 + z^{n-1}w(1 - |w|^2/|z|^2)}{\overline{w}/\overline{z} + z^n(1 - |w|^2/|z|^2)} \times$$
 (2.35)

$$z^{2}\overline{z}\left(\sqrt[n]{\frac{\left(1-|w|^{2}/|z|^{2}+z^{-n}\overline{w}/\overline{z}\right)^{2}}{|z|^{2n}(1-|w|^{2}/|z|^{2})^{2}-1}}\right)',$$

for some branch  $\binom{n}{\sqrt{}}'$  of  $\binom{n}{\sqrt{}}$ . The  $G\left(D_{s,t}^{(2n)}\right)$ -orbits are the following pairwise CR non-equivalent hypersurfaces:

$$\eta_{\alpha}^{(2n)} := \left\{ (z, w) \in M_{\eta}^{(n)} : |z|^n - |z|^{n-2}|w|^2 = \sqrt{(\alpha + 1)/2} \right\},$$

$$s < \alpha < t, n > 2.$$
(2.36)

Next, the group Aut  $\left(D_{s,t}^{(n)}\right)$  for odd  $n \geq 3$  is connected and consists of all lifts from the domain

$$\mathfrak{D} := \Big\{ (z, w) \in \mathbb{C}^2 : 1 + |z|^2 - |w|^2 > |1 + z^2 - w^2|, \, \operatorname{Im}(z(1 + \overline{w})) > 0 \Big\}.$$

to  $M_n := \Pi_n\left(M_{\eta}^{(2n)}\right)$  of maps of the form (2.25). The  $G\left(D_{s,t}^{(n)}\right)$ -orbits are the following pairwise CR non-equivalent hypersurfaces:

$$\eta_{\alpha}^{(n)} := \Pi_n \left( \eta_{\alpha}^{(4n)} \right). \tag{2.37}$$

Finally, the group  $G\left(D_{s,t}^{(\infty)}\right)$  consists of all maps of the form (2.30), and the full group  $\operatorname{Aut}\left(D_{s,t}^{(\infty)}\right)$  is generated by  $G\left(D_{s,t}^{(\infty)}\right)$  and the map

$$z \mapsto 2z + \overline{z} + \ln' \left( i \frac{1 - |w|^2 + e^{-2z} \overline{w}}{\sqrt{\exp(4\operatorname{Re}z)(1 - |w|^2)^2 - 1}} \right),$$

$$w \mapsto \frac{1 + e^{2z} w(1 - |w|^2)}{\overline{w} + e^{2z}(1 - |w|^2)}.$$
(2.38)

The  $G\left(D_{s,t}^{(\infty)}\right)$ -orbits are the following pairwise CR non-equivalent hypersurfaces:

$$\eta_{\alpha}^{(\infty)} := \left\{ (z, w) \in M_{\eta}^{(\infty)} : \exp(2\operatorname{Re} z) (1 - |w|^2) = \sqrt{(\alpha + 1)/2} \right\}, \quad (2.39)$$

We are now ready to formulate the main result of this section.

**THEOREM 2.1** Let M be a (2,3)-manifold. Assume that the G(M)-orbit of every point in M is a strongly pseudoconvex real hypersurface. Then M is holomorphically equivalent to one of the following manifolds:

- (i)  $R_{b,s,t}, b \in \mathbb{R}, |b| \geq 1, b \neq 1, \text{ for } 0 \leq s < t \leq \infty;$
- $U_{s,t}$ , for  $0 \le s < t \le \infty$ , with either s > 0 or  $t < \infty$ ; (ii)
- $\mathfrak{S}_{s,t}, \ 0 \le s < t < \infty;$ (iii)
- $\mathfrak{S}_{s,t}^{(\infty)}, \ 0 \le s < t < \infty;$ (iv)
- $\mathfrak{S}_{s,t}^{(n)}, \ 0 \le s < t < \infty, \ n \ge 2;$ (v)
- $V_{b,s,t}, b > 0, t > 0, e^{-2\pi b}t < s < t;$ (vi)
- $E_{s,t}$ ,  $1 \le s < t < \infty$ ; (vii)
- (viii)  $E_{s,t}^{(4)}, 1 \le s < t < \infty;$
- $E_{s,t}^{(2)}, 1 \le s < t < \infty;$ (ix)
- $\Omega_{s,t}, -1 \le s < t \le 1;$ (X)
- (xi)
- $\Omega_{s,t}^{(\infty)}, -1 \le s < t \le 1;$   $\Omega_{s,t}^{(n)}, -1 \le s < t \le 1, n \ge 2;$ (xii)
- (xiii)  $D_{s,t}, 1 \leq s < t \leq \infty;$
- (xiv)  $D_{s,t}^{(\infty)}, 1 \le s < t \le \infty;$ (xv)  $D_{s,t}^{(n)}, 1 \le s < t \le \infty;$ (xv)  $D_{s,t}^{(n)}, 1 \le s < t \le \infty, n \ge 2.$

**Proof:** In [2] E. Cartan classified all homogeneous 3-dimensional strongly pseudoconvex CR-manifolds. Since the G(M)-orbit of every point in M is such a manifold, every G(M)-orbit is equivalent to a manifold on Cartan's list. We reproduce Cartan's classification below together with the corresponding groups of CR-automorphisms. Note that all possible covers of the

hypersurfaces  $\chi$ ,  $\mu_{\alpha}$ ,  $\nu_{\alpha}$  and  $\eta_{\alpha}$  appear below as explicitly calculated in [6].

```
(a)
```

(b)  $\mathcal{L}_{m} := S^{3}/\mathbb{Z}_{m}, m \in \mathbb{N}, m \geq 2,$ (c)  $\sigma := \{(z, w) \in \mathbb{C}^{2} : \operatorname{Re} w = |z|^{2}\},$ (d)  $\varepsilon_{b} := \{(z, w) \in \mathbb{C}^{2} : \operatorname{Re} z = |w|^{b}, w \neq 0\}, b > 0,$ 

 $\omega := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z = \exp(\operatorname{Re} w)\},\$ 

 $\delta := \{(z, w) \in \mathbb{C}^2 : |w| = \exp(|z|^2)\},\$ (f)

(g)  $\tau_b, b \in \mathbb{R}, |b| \ge 1, b \ne 1 \text{ (see (2.2))},$ 

(h)  $\xi$  (see (2.4)),

 $\chi$  (see (2.6)), (j)

 $\chi^{(\infty)}$  (see (2.9)), (j')

 $\chi^{(n)}, n \ge 2 \text{ (see (2.12))},$ 

(k)  $\rho_b, b > 0 \text{ (see (2.14))},$ 

 $\mu_{\alpha}, \, \alpha > 1 \text{ (see (2.16))},$ (1)

 $\mu_{\alpha}^{(4)}, \, \alpha > 1 \text{ (see (2.21))},$ (1')

 $\mu_{\alpha}^{(2)}, \, \alpha > 1 \text{ (see (2.18))},$ 

 $\nu_{\alpha}$ ,  $-1 < \alpha < 1$ , (see (2.24)), (m)

(m')  $\nu_{\alpha}^{(\infty)}$ ,  $-1 < \alpha < 1$ , (see (2.32)),

(m")  $\nu_{\alpha}^{(n)}$ ,  $-1 < \alpha < 1$ ,  $n \ge 2$  (see (2.29)),

 $\eta_{\alpha}$ ,  $\alpha > 1$ , (see (2.26)), (n)

(n')  $\eta_{\alpha}^{(\infty)}$ ,  $\alpha > 1$ , (see (2.39)),

(n")  $\eta_{\alpha}^{(n)}$ ,  $\alpha > 1$ , n > 2 (see (2.34), (2.36), (2.37)).

The above hypersurfaces are pairwise CR non-equivalent. The corresponding groups of CR-automorphisms are as follows:

# (a) $\operatorname{Aut}_{CR}(S^3)$ :

$$\begin{pmatrix} z \\ w \end{pmatrix} \mapsto \frac{\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}}{c_1 z + c_2 w + d},$$

with  $Q \in SU_{2,1}$ , where

$$Q := \begin{pmatrix} a_{11} & a_{12} & b_1 \\ a_{21} & a_{22} & b_2 \\ c_1 & c_2 & d \end{pmatrix};$$

#### (b) $\operatorname{Aut}_{CR}(\mathcal{L}_m), m \geq 2$ :

$$\left[ \left( \begin{array}{c} z \\ w \end{array} \right) \right] \mapsto \left[ U \left( \begin{array}{c} z \\ w \end{array} \right) \right],$$

where  $U \in U_2$  and  $[(z, w)] \in \mathcal{L}_m$  denotes the equivalence class of  $(z, w) \in S^3$  under the action of  $\mathbb{Z}_m$  embedded in  $U_2$  as a subgroup of scalar matrices;

#### (c) $\operatorname{Aut}_{CR}(\sigma)$ :

$$\begin{array}{rcl}
z & \mapsto & \lambda e^{i\varphi}z + a, \\
w & \mapsto & \lambda^2 w + 2\lambda e^{i\varphi}\overline{a}z + |a|^2 + i\gamma,
\end{array} (2.40)$$

where  $\lambda \in \mathbb{R}^*$ ,  $\varphi, \gamma \in \mathbb{R}$ ,  $a \in \mathbb{C}$ ;

### $(d) \operatorname{Aut}_{CR}(\varepsilon_b)$ :

$$z \mapsto \frac{\lambda z + i\beta}{i\mu z + \kappa},$$

$$w \mapsto \frac{e^{i\varphi}}{(i\mu z + \kappa)^{2/b}} w,$$
(2.41)

where  $\lambda, \beta, \mu, \kappa, \varphi \in \mathbb{R}$ ,  $\lambda \kappa + \mu \beta = 1$ , and for  $t \in \mathbb{C} \setminus (-\infty, 0]$  we set  $t^{2/b} = \exp(2/b \ln t)$  with  $\ln$  being any branch of the logarithm in  $\mathbb{C} \setminus (-\infty, 0]$  (for  $\mu = 0$  we assume that  $\kappa > 0$ );

# $\underline{\text{(e) Aut}_{CR}(\omega)}$ :

$$z \mapsto \frac{\lambda z + i\beta}{i\mu z + \kappa},$$
 (2.42)

$$w \mapsto w - 2\ln(i\mu z + \kappa) + i\gamma,$$

where  $\lambda, \beta, \mu, \kappa, \gamma \in \mathbb{R}$ ,  $\lambda \kappa + \mu \beta = 1$ , and ln is any branch of the logarithm in  $\mathbb{C} \setminus (-\infty, 0]$  (for  $\mu = 0$  we assume that  $\kappa > 0$ );

# $\underline{\text{(f) Aut}_{CR}(\delta)}$ :

$$\begin{array}{ccc} z & \mapsto & e^{i\varphi}z + a, \\ w & \mapsto & e^{i\psi}\exp\Bigl(2e^{i\varphi}\overline{a}z + |a|^2\Bigr)w, \end{array}$$

where  $\varphi, \psi \in \mathbb{R}, a \in \mathbb{C}$ ;

- (g)  $\operatorname{Aut}_{CR}(\tau_b)$  : see (2.1);
- (h)  $Aut_{CR}(\xi)$  : see (2.3);
- (j)  $\operatorname{Aut}_{CR}(\chi)$  : see (2.5);
- $\underline{(\mathbf{j}')\operatorname{Aut}_{CR}\left(\chi^{(\infty)}\right)}$ : see (2.7), (2.8);
- (j") Aut<sub>CR</sub>  $(\chi^{(n)})$ ,  $n \ge 2$ : see (2.10), (2.11);
- (k)  $Aut_{CR}(\rho_b)$  : see (2.13);
- $\underline{\text{(l)} \operatorname{Aut}_{CR}(\mu_{\alpha})}$  : see (2.15);
- $(1') \operatorname{Aut}_{CR} \left( \mu_{\alpha}^{(4)} \right)$  : see (2.19), (2.20);
- (l")  $\operatorname{Aut}_{CR}\left(\mu_{\alpha}^{(2)}\right)$  : see (2.17);
- $\underline{\text{(m) Aut}_{CR}(\nu_{\alpha})}$  : see (2.22);
- $\underline{\text{(m') Aut}_{CR}\left(\nu_{\alpha}^{(\infty)}\right)}$ : see (2.30), (2.31);
- (m")  $\operatorname{Aut}_{CR}\left(\nu_{\alpha}^{(n)}\right) \ n \ge 2 : \operatorname{see} (2.27), (2.28);$
- $\underline{\text{(n)} \operatorname{Aut}_{CR}(\eta_{\alpha})}$ : see (2.25);
- (n') Aut<sub>CR</sub>  $\left(\eta_{\alpha}^{(\infty)}\right)$  : see (2.30), (2.38);
- $\underline{\text{(n")} \operatorname{Aut}_{CR}\left(\eta_{\alpha}^{(2)}\right)} : \operatorname{see} (2.17) \text{ with } A \in SO_{2,1}(\mathbb{R})^c, (2.33);$

(n") 
$$\operatorname{Aut}_{CR}\left(\eta_{\alpha}^{(2n)}\right) \ n \ge 2$$
: see (2.27), (2.35);

 $(n") \operatorname{Aut}_{CR} \left( \eta_{\alpha}^{(n)} \right), n \geq 3 \text{ is odd} : \text{this group is connected and consists of all lifts from the domain } \mathfrak{D} \text{ to } M_n \text{ of maps of the form } (2.25) \text{ (see (10))}.$ 

We will now show that the presence of an orbit of a particular kind in M determines the group G(M) as a Lie group. Fix  $p \in M$  and suppose that O(p) is CR-equivalent to  $\mathfrak{m}$ , where  $\mathfrak{m}$  is one of the model hypersurfaces (a)–(n") listed above. In this case we say that  $\mathfrak{m}$  is the model for O(p). Since G(M) acts effectively on O(p), the CR-equivalence induces an isomorphism between G(M) and a (possibly non-closed) connected 3-dimensional subgroup  $R_{\mathfrak{m}}$  of  $\operatorname{Aut}_{CR}(\mathfrak{m})$  that acts transitively on  $\mathfrak{m}$  (we will often say that the CR-equivalence transforms G(M) into  $R_{\mathfrak{m}}$ ). The subgroup  $R_{\mathfrak{m}}$  a priori depends on the choice of CR-equivalence between O(p) and  $\mathfrak{m}$ , but, as we will see below, this dependence is insignificant.

We will now list all possible groups  $R_{\mathfrak{m}}$  for each model (a)-(n"). In the following lemma  $\mathcal{P}$ , as before, denotes the right half-plane  $\{z \in \mathbb{C} : \operatorname{Re} z > 0\}$ .

#### Lemma 2.2 We have

- (A)  $R_{\mathfrak{m}} = Aut_{CR}(\mathfrak{m})^c$ , if  $\mathfrak{m}$  is one of (g)-(n");
- (B)  $R_{S^3}$  is conjugate in  $Aut_{CR}(S^3)$  to  $SU_2$ ,
- (C)  $R_{\mathcal{L}_m} = SU_2/(SU_2 \cap \mathbb{Z}_m), m \geq 2;$
- (D)  $R_{\sigma}$  is the Heisenberg group, that is, it consists of all elements of  $Aut_{CR}(\sigma)$  with  $\lambda = 1$ ,  $\varphi = 0$  in formula (2.40);
- (E)  $R_{\varepsilon_b}$  is either the subgroup of  $Aut_{CR}(\varepsilon_b)$  corresponding to a subgroup of  $Aut(\mathcal{P})$ , conjugate in  $Aut(\mathcal{P})$  to the subgroup  $\mathcal{T}$  given by

$$z \mapsto \lambda z + i\beta,$$
 (2.43)

where  $\lambda > 0$ ,  $\beta \in \mathbb{R}$ , or is the subgroup  $V_b$  given by  $\varphi = 0$  in formula (2.41);

- (F)  $R_{\omega}$  is either the subgroup of  $Aut_{CR}(\omega)$  corresponding to a subgroup of  $Aut(\mathcal{P})$  conjugate in  $Aut(\mathcal{P})$  to the subgroup  $\mathcal{T}$  specified in (E), or is the subgroup  $V_{\infty}$  given by  $\gamma = 0$  in formula (2.42);
- (G)  $R_{\delta}$  coincides with the subgroup of  $Aut_{CR}(\delta)$  given by  $\varphi = 0$ .

**Proof:** Statement (A) is clear since in cases (g)-(n") we have dimAut<sub>CR</sub>( $\mathfrak{m}$ ) = d(M) = 3. Further, statements (B)-(D), (G) are obtained as in Lemma 2.4 of [5].

To prove statements (E) and (F) note that every (a priori non-closed) codimension 1 subgroup of  $\operatorname{Aut}(\mathcal{P})$  is conjugate in  $\operatorname{Aut}(\mathcal{P})$  to the subgroup  $\mathcal{T}$  defined in (2.43).

The lemma is proved.

Lemma 2.2 implies, in particular, that if for some point  $p \in M$  the model for O(p) is  $S^3$ , then M admits an effective action of  $SU_2$ . Therefore, M is holomorphically equivalent to one of the manifolds listed in [8]. However, none of the hyperbolic manifolds with 3-dimensional automorphism group on the list has a spherical orbit. Hence  $S^3$  is never the model of an G(M)-orbit.

We now observe that each of the groups  $R_{\mathfrak{m}}$  acts by holomorphic transformations on a certain complex manifold  $M_{\mathfrak{m}}$  containing  $\mathfrak{m}$ , and every orbit of this action either is CR-equivalent to  $\mathfrak{m}$  (cases (b)–(k)), or belongs to the same group to which  $\mathfrak{m}$  belongs and the orbits are pairwise CR non-equivalent (cases (l)–(n")). The manifolds  $M_{\mathfrak{m}}$  are as follows:

(b) 
$$M_{\mathcal{L}_m} = \mathbb{C}^2 \setminus \{0\}/\mathbb{Z}_m, m \geq 2;$$

(c) 
$$M_{\sigma} = \mathbb{C}^2$$
;

(d) 
$$M_{\varepsilon_b} = \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0, w \neq 0\};$$

(e) 
$$M_{\omega} = \{(z, w) \in \mathbb{C}^2 : \text{Re } z > 0\};$$

(f) 
$$M_{\delta} = \{(z, w) \in \mathbb{C}^2 : w \neq 0\};$$

(g) 
$$M_{\tau_b} = \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0, \operatorname{Re} w > 0\};$$

(h) 
$$M_{\xi} = \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} w > 0\};$$

(j) 
$$M_{\chi} = \mathbb{C}^2 \setminus \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z = 0, \operatorname{Re} w = 0\};$$

(j') 
$$M_{\chi^{(\infty)}} = M_{\chi}^{(\infty)}$$
 (see **(4)**);

(j") 
$$M_{\chi^{(n)}} = M_{\chi}^{(n)}, n \ge 2 \text{ (see (4))};$$

(k) 
$$M_{\rho_b} = \mathbb{C}^2 \setminus \{(z, w) \in \mathbb{C}^2 : \text{Re } z = 0, \text{ Re } w = 0\};$$

(1) 
$$M_{\mu_{\alpha}} = \bigcup_{\alpha > 1} \mu_{\alpha} = \mathbb{CP}^2 \setminus \mathbb{RP}^2;$$

(l') 
$$M_{\mu_{\alpha}^{(4)}} = \bigcup_{\alpha>1} \mu_{\alpha}^{(4)} = M_{\mu}^{(4)} \text{ (see (7))};$$

(l") 
$$M_{\mu_{\alpha}^{(2)}} = \bigcup_{\alpha>1} \mu_{\alpha}^{(2)} = \mathcal{Q}_{+} \setminus \mathbb{R}^{3} \text{ (see (7))};$$

(m) 
$$M_{\nu_{\alpha}} = \bigcup_{-1<\alpha<1} \nu_{\alpha} = \left\{ (z, w) \in \mathbb{C}^2 : -|z^2 + w^2 - 1| < |z|^2 + |w|^2 - 1 < |z^2 + w^2 - 1| \right\};$$

(m') 
$$M_{\nu_{\alpha}^{(\infty)}} = \bigcup_{-1 < \alpha < 1} \nu_{\alpha}^{(\infty)} = M_{\nu}^{(\infty)} \text{ (see (10))};$$

(m") 
$$M_{\nu_{\alpha}^{(n)}} = \bigcup_{-1 < \alpha < 1} \nu_{\alpha}^{(n)} = M_{\nu}^{(n)}, \ n \ge 2 \text{ (see (10))};$$

(n) 
$$M_{\eta_{\alpha}} = \bigcup_{\alpha > 1} \eta_{\alpha} = \mathfrak{D}$$
 (see **(10)**);

(n') 
$$M_{\eta_{\alpha}^{(\infty)}} = \bigcup_{\alpha>1} \eta_{\alpha}^{(\infty)} = M_{\eta}^{(\infty)}$$
 (see **(10)**);

(n") 
$$M_{\eta_{\alpha}^{(2)}} = \bigcup_{\alpha>1} \eta_{\alpha}^{(2)} = \Omega_{\eta} \text{ (see (10))};$$

(n") 
$$M_{\eta_{\alpha}^{(2n)}} = \bigcup_{\alpha>1} \eta_{\alpha}^{(2n)} = M_{\eta}^{(n)}, n \geq 2 \text{ (see (10))};$$

(n") 
$$M_{\eta_{\alpha}^{(n)}} = \bigcup_{\alpha > 1} \eta_{\alpha}^{(n)} = M_n, \ n \ge 3 \text{ is odd (see (10))}.$$

In cases (b)–(k) every two  $R_{\mathfrak{m}}$ -orbits are CR-equivalent (and equivalent to  $\mathfrak{m}$ ) by means of automorphisms of  $M_{\mathfrak{m}}$  of the simple forms specified below:

- (b)  $[(z, w)] \mapsto [(az, aw)], a > 0;$
- (c)  $z \mapsto z, w \mapsto w + a, a \in \mathbb{R}$ ;
- (d)  $z \mapsto az, w \mapsto w, a > 0$ ;
- (e) as in (d);
- (f)  $z \mapsto z, w \mapsto aw, a > 0$ ;

(g) as in (f); 
$$(2.44)$$

- (h)  $z \mapsto az, w \mapsto aw, a > 0$ ;
- (j) as in (h);
- (j')  $z \mapsto z + a, w \mapsto w, a \in \mathbb{R};$
- (j) as in (h);
- (k) as in (h).

We will now show how the models can be glued together to form hyperbolic manifolds. The gluing procedure described below is similar to that used in [4], [5]. The procedure comprises the following steps:

(I). Start with a real hypersurface orbit O(p) with model  $\mathfrak{m}$  and consider a real-analytic CR-isomorphism  $f:O(p)\to\mathfrak{m}$ . Clearly, f satisfies

$$f(gq) = \varphi(g)f(q), \tag{2.45}$$

for all  $g \in G(M)$  and  $q \in O(p)$ , where  $\varphi : G(M) \to R_{\mathfrak{m}}$  is an isomorphism.

(II). Observe that f can be extended to a biholomorphic map from a G(M)-invariant connected neighborhood of O(p) in M onto an  $R_{\mathfrak{m}}$ -invariant neighborhood of  $\mathfrak{m}$  in  $M_{\mathfrak{m}}$ . First of all, extend f to some neighborhood U of O(p)

to a biholomorphic map onto a neighborhood of  $\mathfrak{m}$  in  $M_{\mathfrak{m}}$ . Fix  $s \in U$  and  $s_0 \in O(s)$ . Choose  $h_0 \in G(M)$  such that  $s_0 = h_0 s$  and define  $f(s_0) := \varphi(h_0) f(s)$ . To see that f is well-defined at  $s_0$ , suppose that for some  $h_1 \in G(M)$ ,  $h_1 \neq h_0$ , we have  $s_0 = h_1 s$ , and show that  $\varphi(h)$  fixes f(s), where  $h := h_1^{-1} h_0$ . Indeed, for every  $g \in G(M)$  identity (2.45) holds for  $g \in U_g$ , where  $g \in U_g$  is the connected component of  $g^{-1}(U) \cap U$  containing  $g \in U_g$ . Since  $g \in U_g$  we have  $g \in U_g$  and the application of (2.45) to  $g \in U_g$  and  $g \in U_g$  is the extended map satisfies (2.45) for all  $g \in G(M)$  and  $g \in U'$ .

(III). Consider a maximal G(M)-invariant domain  $D \subset M$  from which there exists a biholomorphic map f onto an  $R_{\mathfrak{m}}$ -invariant domain in  $M_{\mathfrak{m}}$  satisfying (2.45) for all  $g \in G(M)$  and  $g \in D$ . The existence of such a domain is guaranteed by the previous step. Assume that  $D \neq M$  and consider  $x \in \partial D$ . Let  $\mathfrak{m}_1$  be the model for O(x) and let  $f_1:O(x)\to\mathfrak{m}_1$  be a real-analytic CR-isomorphism satisfying (2.45) for all  $g \in G(M)$ ,  $q \in O(x)$  and some isomorphism  $\varphi_1: G(M) \to R_{\mathfrak{m}_1}$  in place of  $\varphi$ . As in (II), extend  $f_1$  to a biholomorphic map from a connected G(M)-invariant neighborhood V of O(x)onto an  $R_{\mathfrak{m}_1}$ -invariant neighborhood of  $\mathfrak{m}_1$  in  $M_{\mathfrak{m}_1}$ . The extended map satisfies (2.45) for all  $g \in G(M)$ ,  $q \in V$  and  $\varphi_1$  in place of  $\varphi$ . Consider  $s \in V \cap D$ . The maps f and  $f_1$  take O(s) onto an  $R_{\mathfrak{m}}$ -orbit  $\mathfrak{m}'$  in  $M_{\mathfrak{m}}$  and an  $R_{\mathfrak{m}_1}$ -orbit  $\mathfrak{m}'_1$  in  $M_{\mathfrak{m}_1}$ , respectively. Then  $F:=f_1\circ f^{-1}$  establishes a CR-isomorphism between  $\mathfrak{m}'$  and  $\mathfrak{m}'_1$ . Hence  $M_{\mathfrak{m}} = M_{\mathfrak{m}_1}$  and  $R_{\mathfrak{m}} = R_{\mathfrak{m}_1}$ . Moreover, F is either an element of  $\operatorname{Aut}_{CR}(\mathfrak{m}')$  (if  $\mathfrak{m}'=\mathfrak{m}'_1$ ), or is a composition of an element of  $\operatorname{Aut}_{CR}(\mathfrak{m}')$  and a non-trivial map from list (2.44) that takes  $\mathfrak{m}'$  onto  $\mathfrak{m}'_1$  (if  $\mathfrak{m}' \neq \mathfrak{m}'_1$ ). Therefore, F extends to a holomorphic automorphism of  $M_{\mathfrak{m}}$ .

(IV). Since O(x) is strongly pseudoconvex and closed in M, for V small enough we have  $V = V_1 \cup V_2 \cup O(x)$ , where  $V_j$  are open connected non-intersecting sets. For each j,  $D \cap V_j$  is a union of G(M)-orbits and therefore is mapped by f onto a union W of  $R_{\mathfrak{m}}$ -orbits in  $M_{\mathfrak{m}}$ . If W is disconnected, so is D which contradicts the definition of D. Thus,  $D \cap V_j$  is connected for j = 1, 2, and, if V is sufficiently small, then each  $V_j$  is either a subset of D or disjoint from it. Suppose first that there is only one j for which  $D \cap V_j \neq \emptyset$ . In this case  $D \cap V$  is connected and  $V \setminus (D \cup O(x)) \neq \emptyset$ . Setting now

$$\tilde{f} := \begin{cases} f & \text{on } D, \\ F^{-1} \circ f_1 & \text{on } V, \end{cases}$$
 (2.46)

we obtain a biholomorphic extension of f to  $D \cup V$ . By construction,  $\tilde{f}$  satisfies (2.45) for  $g \in G(M)$  and  $q \in D \cup V$ . Since  $D \cup V$  is strictly larger than D, we obtain a contradiction with the maximality of D. Thus, in this case D = M, and hence M is holomorphically equivalent to an  $R_{\mathfrak{m}}$ -invariant domain in  $M_{\mathfrak{m}}$ .

Suppose now that  $V_j \subset D$  for j=1,2. Applying formula (2.46) we can extend  $f|_{V_1}$  and  $f|_{V_2}$  to biholomorphic maps  $\hat{f}_1$ ,  $\hat{f}_2$ , respectively, from a neighborhood of O(x) into  $M_{\mathfrak{m}}$ ; each of these maps satisfies (2.45). Let  $\hat{\mathfrak{m}}_j := \hat{f}_j(O(x))$ , j=1,2. Clearly,  $\hat{\mathfrak{m}}_1 \neq \hat{\mathfrak{m}}_2$ . The map  $\hat{F} := \hat{f}_2 \circ \hat{f}_1^{-1}$  is a CR-isomorphism from  $\hat{\mathfrak{m}}_1$  onto  $\hat{\mathfrak{m}}_2$ . In this case  $M_{\mathfrak{m}}$  is one of the manifolds occurring in (b)-(k), and  $\hat{F}$  is a composition of an element of  $\operatorname{Aut}_{CR}(\hat{\mathfrak{m}}_1)$  and a non-trivial map from the corresponding form on list (2.44) that takes  $\hat{\mathfrak{m}}_1$  onto  $\hat{\mathfrak{m}}_2$ . Let  $\Gamma$  denote the cyclic group of automorphisms of  $M_{\mathfrak{m}}$  generated by  $\hat{F}$ . It follows from the explicit form of F and  $M_{\mathfrak{m}}$  in each of the cases (b)-(k) that  $\Gamma$  acts freely properly discontinuously on  $M_{\mathfrak{m}}$ . Let  $M_{\Gamma}$  be the complex manifold obtained by factoring  $M_{\mathfrak{m}}$  by the action of  $\Gamma$  and let  $\Pi_{\Gamma}: M_{\mathfrak{m}} \to M_{\Gamma}$  be the factorization map. Then the map  $\Pi_{\Gamma} \circ \mathcal{F}$ , where

$$\mathcal{F} := \begin{cases} f & \text{on } D, \\ \hat{f}_1 & \text{on } O(x), \end{cases}$$
 (2.47)

establishes holomorphic equivalence between M and  $M_{\Gamma}$ . Observe, however, that in cases (b)–(f), (h)–(k) the manifold  $M_{\Gamma}$  is not hyperbolic, since it is covered by  $M_{\mathfrak{m}}$  and none of the manifolds  $M_{\mathfrak{m}}$  in these cases is hyperbolic. Further, it is straightforward to see that in the remaining case (g) the manifold  $M_{\Gamma}$  is isomorphic to the product of either an annulus or a punctured disk and  $\Delta$  and hence  $d(M_{\Gamma})=4$ . These contradictions show that in all cases (b)–(k) exactly one of  $V_j$ , j=1,2, is a subset of D, and hence M is holomorphically equivalent to an  $R_{\mathfrak{m}}$ -invariant domain in  $M_{\mathfrak{m}}$ .

(V). In each of cases (b)-(n") the determination of  $R_{\mathfrak{m}}$ -invariant domains is straightforward; they are as follows:

(b) 
$$\{(z, w) \in \mathbb{C}^2 : s < |z|^2 + |w|^2 < t\} / \mathbb{Z}_m, \ 0 \le s < t < \infty, \ m \ge 2;$$

(c) 
$$\{(z, w) \in \mathbb{C}^2 : s + |z|^2 < \text{Re } w < t + |z|^2\}, -\infty < s < t \le \infty;$$

(d) 
$$\mathfrak{R}_{b,s,t} := \{(z,w) \in \mathbb{C}^2 : s|w|^b < \operatorname{Re} z < t|w|^b, w \neq 0\}, \text{ for } 0 \leq s < t \leq \infty, \text{ with either } s > 0 \text{ or } t < \infty;$$

(e) 
$$\mathfrak{R}_{s,t} := \{(z, w) \in \mathbb{C}^2 : s \exp(\operatorname{Re} w) < \operatorname{Re} z < t \exp(\operatorname{Re} w)\}, \text{ for } 0 < s < t < \infty, \text{ with either } s > 0 \text{ or } t < \infty;$$

(f) 
$$\{(z, w) \in \mathbb{C}^2 : s \exp(|z|^2) < |w| < t \exp(|z|^2)\}, \ 0 < s < t \le \infty;$$

(g) 
$$R_{b,s,t}$$
, with  $0 \le s < t \le \infty$ ;

(h) 
$$U_{s,t}$$
, for  $0 \le s < t \le \infty$ , with either  $s > 0$  or  $t < \infty$ ;

(j) 
$$\mathfrak{S}_{s,t}$$
,  $0 \le s < t < \infty$ ;

(j') 
$$\mathfrak{S}_{s,t}^{(\infty)}$$
,  $0 \le s < t < \infty$ ;

(j") 
$$\mathfrak{S}_{s,t}^{(n)}$$
,  $0 \le s < t < \infty$ ,  $n \ge 2$ ;

(k) 
$$V_{b,s,t}$$
,  $0 < t < \infty$ ,  $e^{-2\pi b}t < s < t$ ;

(1) 
$$E_{s,t}$$
,  $1 \le s < t < \infty$ ;

(l') 
$$E_{s,t}^{(4)}$$
,  $1 \le s < t < \infty$ ;

(l") 
$$E_{s,t}^{(2)}$$
,  $1 \le s < t < \infty$ ;

(m) 
$$\Omega_{s,t}, -1 \le s < t \le 1;$$

(m') 
$$\Omega_{s,t}^{(\infty)}$$
,  $-1 \le s < t \le 1$ ;

(m") 
$$\Omega_{s,t}^{(n)}$$
,  $-1 \le s < t \le 1$ ,  $n \ge 2$ ;

(n) 
$$D_{s,t}$$
,  $1 \le s < t \le \infty$ ;

(n') 
$$D_{s,t}^{(\infty)}$$
,  $1 \le s < t \le \infty$ ;

(n") 
$$D_{s,t}^{(2)}$$
,  $1 \le s < t \le \infty$ ;

(n") 
$$D_{s,t}^{(2n)}$$
,  $1 \le s < t \le \infty$ ,  $n \ge 2$ ;

(n") 
$$D_{s,t}^{(n)}$$
,  $1 \le s < t \le \infty$ ,  $n \ge 3$ ,  $n$  is odd.

This concludes our gluing procedure. Note that in each of cases (d) and (e) we have two non-isomorphic possibilities for  $R_{\mathfrak{m}}$ . Each of the possibilities leads to the same set of  $R_{\mathfrak{m}}$ -invariant domains.

To finish the proof of the theorem it now remains to observe that the automorphism groups of  $R_{\mathfrak{m}}$ -invariant domains that appear in cases (b)–(f) have dimension at least 4 rather than 3. The remaining domains form list (i)–(xv) as stated in the theorem.

The proof of the theorem is complete.

# 3 Levi-Flat orbits

In this section we deal with Levi-flat orbits. First, we will classify such orbits. In order to formulate our orbit classification result, we introduce the following groups of transformations.

For every  $b \in \mathbb{R}$  let  $G_b$  be the group of all maps of the form (2.1),  $G'_0$  be the group of all maps of the form

$$\begin{array}{ccc}
z & \mapsto & \lambda z + i\beta, \\
w & \mapsto & e^{i\varphi}w,
\end{array} \tag{3.1}$$

where  $\lambda > 0$ ,  $\beta, \varphi \in \mathbb{R}$ , and G the group of all maps of the form (2.3).

We will now prove the following proposition. Observe that the proposition applies to an arbitrary (2,3)-manifold (possibly containing codimension 2 orbits).

**Proposition 3.1** Let M be a (2,3)-manifold. Assume that for a point  $p \in M$  its orbit O(p) is Levi-flat. Then O(p) is equivalent to one of the following hypersurfaces:

$$\begin{array}{lcl} O_1 &:= & \left\{ (z,w) \in \mathbb{C}^2 : \operatorname{Re} z > 0, \, \operatorname{Re} w = 0 \right\}, \\ O_2 &:= & \left\{ (z,w) \in \mathbb{C}^2 : \operatorname{Re} z > 0, \, |w| = 1 \right\}, \end{array}$$

by means of a real-analytic CR-map. The CR-equivalence can be chosen so that it transforms G(M) into either the group  $G_b$  for some  $b \in \mathbb{R}$  or the group G in the first case, and into the group  $G'_0$  in the second case.

**Proof:** Recall that the hypersurface O(p) is foliated by complex manifolds holomorphically equivalent to  $\mathcal{P}$  (see (ii) of Proposition 1.1). Denote by  $\mathfrak{g}$  the Lie algebra of vector fields on O(p) arising from the action of G(M). Clearly,  $\mathfrak{g}$  is isomorphic to the Lie algebra of G(M). For  $q \in O(p)$  we consider the leaf  $M_q$  of the foliation passing through q and the subspace  $\mathfrak{l}_q \subset \mathfrak{g}$  of all vector fields tangent to  $M_q$  at q. Since vector fields in  $\mathfrak{l}_q$  remain tangent to  $M_q$  at each point in  $M_q$ , the subspace  $\mathfrak{l}_q$  is in fact a Lie subalgebra of  $\mathfrak{g}$ . It follows from the definition of  $\mathfrak{l}_q$  that dim  $\mathfrak{l}_q = 2$ .

Denote by  $H_q$  the (possibly non-closed) connected subgroup of G(M)with Lie algebra  $l_q$ . It is straightforward to verify that the group  $H_q$  acts on  $M_q$  by holomorphic transformations. If some element  $g \in H_q$  acts trivially on  $M_q$ , then  $g \in I_q$ . If for every non-identical element of  $L_q$  its projection to  $L'_q$  is non-identical (see (ii) of Proposition 1.1), then every non-identical element of  $I_q$  acts non-trivially on  $M_q$  and thus g = id; if  $L_q$  contains a non-identical element with an identical projection to  $L'_q$  and  $g \neq id$ , then  $g = g_q$ , where  $g_q$  denotes the element of  $I_q$  corresponding to the non-trivial element in  $\mathbb{Z}_2$  (see (ii) of Proposition 1.1). Thus, either  $H_q$  or  $H_q/\mathbb{Z}_2$  acts effectively on  $M_q$  (the former case occurs if  $g_q \notin H_q$ , the latter if  $g_q \in H_q$ ). Since dim  $H_q = 2$ , we obtain that either  $H_q$  or  $H_q/\mathbb{Z}_2$  is isomorphic to a 2-dimensional (possibly non-closed) subgroup of  $Aut(\mathcal{P})$ . As we noted in the proof of Lemma 2.2, every codimension 1 subgroup of  $Aut(\mathcal{P})$  is conjugate in  $\operatorname{Aut}(\mathcal{P})$  to the subgroup  $\mathcal{T}$  (see (2.43)). The Lie algebra of this subgroup is isomorphic to the 2-dimensional Lie algebra  $\mathfrak{h}$  given by two generators X and Y satisfying [X,Y]=X. Therefore,  $\mathfrak{l}_q$  is isomorphic to  $\mathfrak{h}$  for every  $q\in O(p)$ .

It is not difficult to determine all 3-dimensional Lie algebras containing a subalgebra isomorphic to  $\mathfrak{h}$ . Every such algebra has generators X, Y, Z that satisfy one of the following sets of relations:

$$\begin{array}{lll} (\mathrm{R1}) & [X,Y] = X, & [Z,X] = 0, & [Z,Y] = bZ, & b \in \mathbb{R}, \\ (\mathrm{R2}) & [X,Y] = X, & [Z,X] = 0, & [Z,Y] = X+Z, \\ (\mathrm{R3}) & [X,Y] = X, & [Z,X] = Y, & [Z,Y] = -Z. \end{array}$$

Suppose first that  $\mathfrak{g}$  is given by relations (R1). In this case  $\mathfrak{g}$  is isomorphic to the Lie algebra of the simply connected Lie group  $G_b$ . Indeed, the Lie algebra of  $G_b$  is isomorphic to the Lie algebra of vector fields on  $\mathbb{C}^2$  with the generators

$$X_1 := i \partial/\partial z,$$
  
 $Y_1 := z \partial/\partial z + bw \partial/\partial w,$   
 $Z_1 := i \partial/\partial w,$ 

that clearly satisfy (R1).

Assume first that  $b \neq 0$ . In this case the center of  $G_b$  is trivial, and hence  $G_b$  is the only (up to isomorphism) connected Lie group whose Lie algebra is given by relations (R1). Therefore, G(M) is isomorphic to  $G_b$ . Assume further that  $b \neq 1$ . In this case, it is straightforward to observe that every subalgebra of  $\mathfrak{g}$  isomorphic to  $\mathfrak{h}$  is generated by either X and  $Y + \nu Z$ , or by Z and  $\nu X + Y$  for some  $\nu \in \mathbb{R}$ . The connected subgroup of  $G_b$  with Lie algebra generated by  $X_1$  and  $Y_1 + \nu Z_1$  is conjugate in  $G_b$  to the closed subgroup  $G_b^1$  given by  $\gamma = 0$ ; similarly, the connected subgroup of with Lie algebra generated by  $Z_1$  and  $\nu X_1 + Y_1$  is conjugate to the closed subgroup  $G_b^2$  given by  $\beta = 0$  (see (2.1)). Moreover, the conjugating element can be chosen to have the form

$$\begin{array}{ccc}
z & \mapsto & z \\
w & \mapsto & w + i\gamma, & \gamma \in \mathbb{R},
\end{array} \tag{3.3}$$

in the first case, and the form

$$\begin{array}{ccc}
z & \mapsto & z + i\beta, & \gamma \in \mathbb{R}, \\
w & \mapsto & w,
\end{array} \tag{3.4}$$

in the second case.

Thus we have shown that (upon identification of G(M) and  $G_b$ ), the subgroup  $H_q$  for every  $q \in O(p)$  is conjugate to either  $G_b^1$  or to  $G_b^2$  by a map of either the form (3.3) or the form (3.4), respectively. In particular,  $H_q$  is isomorphic to  $\mathcal{T}$  and hence does not have subgroups isomorphic to  $\mathbb{Z}_2$ . Therefore,  $H_q$  acts effectively on  $M_q$ . Since the subgroups  $H_q$  are conjugate to each other, it follows that either  $H_q$  is conjugate to  $G_b^1$  for every q, or  $H_q$ is conjugate to  $G_b^2$  for every q. Suppose, for example, that the former holds. Since the normalizer of  $G_b^1$  in  $G_b$  coincides with  $G_b^1$ , for every  $q_1, q_2 \in O(p)$ there exists a unique  $g \in G(M)$  of the form (3.3) such that  $gM_{q_1} = M_{q_2}$ . Fix  $q_0 \in O(p)$  and let  $f: M_{q_0} \to \mathcal{P}$  be a holomorphic equivalence that transforms  $H_{q_0}|_{M_{q_0}}$  into the group  $\mathcal T$  and such that  $\hat X_1|_{M_{q_0}}$  and  $\hat Y_1|_{M_{q_0}}$  are transformed into the vector fields  $i\partial/\partial z|_{\mathcal{P}}$  and  $z\partial/\partial z|_{\mathcal{P}}$ , respectively, where  $X_1, Y_1$  are the vector fields on O(p) corresponding to  $X_1, Y_1$ . For every  $q \in O(p)$  find the unique map g of the form (3.3) such that  $gM_{q_0} = M_q$ and define  $F(q) := (f(g^{-1}(q)), i\gamma) \in \mathbb{C}^2$ , with  $\gamma$  corresponding to g as in formula (3.3). Clearly, F is a real-analytic CR-isomorphism between O(p)and  $O_1$  that transforms G(M) into  $G_b$ . The case when  $H_q$  is conjugate to  $G_b^2$  for every  $q \in O(p)$  is treated similarly; arguing as above we construct a real-analytic CR-isomorphism between O(p) and

$$O_1' := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z = 0, \operatorname{Re} w > 0\},$$
 (3.5)

that transforms G(M) into  $G_b$ . Further, interchanging the variables turns  $O'_1$  into  $O_1$  and  $G_b$  into  $G_{1/b}$ .

Suppose now that b=1. In this case, in addition to the subalgebras analogous to those arising for  $b \neq 1$ , a subalgebra of  $\mathfrak{g}$  isomorphic to  $\mathfrak{h}$  can also be generated by  $X + \eta Z$  and  $Y + \nu Z$  for some  $\eta, \nu \in \mathbb{R}, \eta \neq 0$ . The connected subgroup of  $G_1$  corresponding to this subalgebra is conjugate in  $G_1$  to the subgroup  $G_{1,\eta}$  of all maps of the form (2.1) with  $b=1, \gamma=\beta\eta$ . Moreover, the conjugating element can be chosen to have the form (3.3). Thus, upon identification of G(M) and  $G_1$ , the subgroup  $H_q$  for every  $q \in$ O(p) is conjugate to either  $G_1^1$  or to  $G_1^2$ , or to  $G_{1,\eta}$  for some  $\eta \neq 0$  (all these subgroups are closed). In particular,  $H_q$  is isomorphic to  $\mathcal{T}$  and hence acts effectively on  $M_q$ . Since the subgroups  $H_q$  are conjugate to each other, it follows that either  $H_q$  is conjugate to  $G_1^1$  for every q, or  $H_q$  is conjugate to  $G_1^2$  for every q, or  $H_q$  is conjugate to  $G_{1,\eta}$  for every q and a fixed  $\eta$ . The first two cases are treated as for  $b \neq 1$ . Suppose that  $H_q$  is conjugate to  $G_{1,\eta}$  for every  $q \in O(p)$ . Since the normalizer of  $G_{1,\eta}$  in  $G_1$  coincides with  $G_{1,\eta}$ , for every  $q_1,q_2\in O(p)$  there exists a unique  $g\in G(M)$  of the form (3.3) such that  $gM_{q_1} = M_{q_2}$ . Fix  $q_0 \in O(p)$  and let  $f: M_{q_0} \to N_{\eta}$ , with  $N_{\eta} := \{(z, w) \in \mathbb{C}^2 : w = \eta z, \operatorname{Re} z > 0\}, \text{ be a holomorphic equivalence that}$ transforms  $H_{q_0}|_{M_{q_0}}$  into the group  $G_{1,\eta}|_{N_{\eta}}$  and such that  $X_1|_{M_{q_0}}$  and  $Y_1|_{M_{q_0}}$  are transformed into the vector fields  $i\partial/\partial z + i\eta\partial/\partial w|_{N_{\eta}}$  and  $z\partial/\partial z + w\partial/\partial w|_{N_{\eta}}$ , respectively. For every  $q \in O(p)$  find the unique map g of the form (3.3) such that  $gM_{q_0} = M_q$  and define  $F(q) := (f(g^{-1}(q)), i\gamma)$ , with  $\gamma$  corresponding to g as in formula (3.3). Clearly, F is a real-analytic CR-isomorphism between O(p) and

$$O_{1,\eta} := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} w = \eta \operatorname{Re} z, \operatorname{Re} z > 0 \},$$

that transforms G(M) into  $G_1$ . Finally, the transformation

$$\begin{array}{ccc} z & \mapsto & z, \\ w & \mapsto & w - \eta z \end{array}$$

turns  $O_{1,\eta}$  into  $O_1$  and preserves the group  $G_1$ .

Suppose now that b=0. In this case there are exactly two (up to isomorphism) connected Lie groups with Lie algebra  $\mathfrak{g}$ :  $G_0$  and  $G'_0$ . It is straightforward to see that every subalgebra of  $\mathfrak{g}$  isomorphic to  $\mathfrak{h}$  is generated by X and  $Y + \nu Z$  for some  $\nu \in \mathbb{R}$ . Clearly, the connected subgroup of  $G_0$  with Lie algebra generated by  $X_1$  and  $Y_1 + \nu Z_1$  coincides with the closed normal subgroup  $G_{0,\nu}$  given by  $\lambda = e^t$ ,  $\gamma = \nu t$ ,  $t \in \mathbb{R}$  (see (2.1)). It then follows that if G(M) is isomorphic to  $G_0$ , there exists  $\nu \in \mathbb{R}$ , such that, identifying G(M) and  $G_0$ , we have  $H_q = G_{0,\nu}$  for every  $q \in O(p)$ . Further, realize the Lie algebra of  $G'_0$  as the Lie algebra generated by the following vector fields on  $\mathbb{C}^2$ :

 $X'_1 := i \partial/\partial z,$   $Y'_1 := z \partial/\partial z,$  $Z'_1 := i w \partial/\partial w,$ 

which clearly satisfy (R1) of (3.2). Therefore, the connected subgroup of  $G'_0$  with Lie algebra generated by  $X'_1$  and  $Y'_1 + \nu Z'_1$  coincides with the closed normal subgroup  $G'_{0,\nu}$  of  $G'_0$  given by  $\lambda = e^t$ ,  $\varphi = \nu t$ ,  $t \in \mathbb{R}$  (see (3.1)). It then follows that if G(M) is isomorphic to  $G'_0$ , there exists  $\nu \in \mathbb{R}$ , such that, identifying G(M) and  $G'_0$ , we have  $H_q = G'_{0,\nu}$  for every  $q \in O(p)$ .

Thus, if b=0, every subgroup  $H_q$  is normal, closed, isomorphic to  $\mathcal{T}$ (hence acts effectively on  $M_q$ ). In particular, all these subgroups coincide for  $q \in O(p)$ . Denote by H the coinciding subgroups  $H_q$ . The group H acts properly on O(p), and the orbits of this action are the leaves  $M_q$  of the foliation on O(p). Further, we have  $G(M) = H \times L$ , where L is either the subgroup given by  $\lambda = 1$ ,  $\beta = 0$  in formula (2.1) where we set b = 0, or to the subgroup given by  $\lambda = 1$ ,  $\beta = 0$  in formula (3.1), and hence isomorphic to either  $\mathbb{R}$  or  $S^1$ . For every  $q \in O(p)$  let  $S_q := \{g \in L : gM_q = M_q\}$ . Since  $M_q$  is closed,  $S_q$  is a closed subgroup of L. Clearly, for every  $g \in S_q$  there is  $h \in H$  such that  $hg \in I_q$ . The elements g and h lie in the projections of  $I_q$  to L and H, respectively. Since H is isomorphic to  $\mathcal{T}$ , it does not have non-trivial finite subgroups, hence the projection of  $I_q$  to H is trivial, and therefore  $S_q = I_q$ . Since all isotropy subgroups are contained in the Abelian subgroup L and are conjugate to each other in G(M), they are in fact identical. The effectiveness of the action of G(M) on M now implies that all isotropy subgroups are trivial and hence every  $S_q$  is trivial as well.

Thus, we have shown that for every  $q_1, q_2 \in O(p)$  there is a unique  $g \in L$ , such that  $gM_{q_1} = M_{q_2}$ . Fix  $q_0 \in O(p)$  and let  $f: M_{q_0} \to \mathcal{P}$  be a holomorphic equivalence that transforms  $H_{q_0}|_{M_{q_0}}$  into the group  $\mathcal{T}$ . For every  $q \in O(p)$  find the unique map  $g \in L$  such that  $gM_{q_0} = M_q$  and define

 $F(q) := (f(g^{-1}(q)), u)$ , where u corresponds to g as follows (see formula (2.1) with b = 0 and formula (3.1)):  $u = i\gamma$  if  $L \simeq \mathbb{R}$  and  $u = e^{i\varphi}$  if  $L \simeq S^1$ . Clearly, if  $G(M) \simeq G_0$ , the map F takes O(p) onto  $O_1$  and transforms G(M) into  $G_0$ , and if  $G(M) \simeq G'_0$ , it takes O(p) onto  $O_2$  and transforms G(M) into  $G'_0$ .

Suppose next that  $\mathfrak{g}$  is given by relations (R2) (see (3.2)). In this case  $\mathfrak{g}$  is isomorphic to the Lie algebra of the simply connected Lie group G. Indeed, the Lie algebra of G is isomorphic to the Lie algebra of holomorphic vector fields on  $\mathbb{C}^2$  with the following generators:

$$\begin{array}{lll} X_2 &:= & i\,\partial/\partial z, \\ Y_2 &:= & (z+w)\,\partial/\partial z + w\,\partial/\partial w, \\ Z_2 &:= & i\,\partial/\partial w, \end{array}$$

that clearly satisfy (R2). It is straightforward to observe that the center of G is trivial, and hence G is the only (up to isomorphism) connected Lie group whose Lie algebra is given by relations (R2). Therefore, G(M) is isomorphic to G. In this case every subalgebra of  $\mathfrak{g}$  isomorphic to  $\mathfrak{h}$  is generated by X and  $Y + \nu Z$  for some  $\nu \in \mathbb{R}$ . The connected subgroup of G with Lie algebra generated by  $X_2$  and  $Y_2 + \nu Z_2$  is conjugate in G to the closed subgroup G given by G (see (2.3)). Moreover, the conjugating element can be chosen to have the form (3.3).

Thus – upon identification of G(M) and G – the subgroup  $H_q$  for every  $q \in O(p)$  is conjugate to Q by a map of the form (3.3). Further, since the normalizer of Q in G coincides with Q, we proceed as in the case of relations (R1) with  $b \neq 0, 1$  and obtain that there exists a real-analytic CR-isomorphism between O(p) and  $O_1$  that transforms G(M) into G.

Suppose finally that  $\mathfrak{g}$  is given by relations (R3) (see (3.2)). We will show that this is in fact impossible. In this case  $\mathfrak{g}$  is isomorphic to the algebra  $\mathfrak{su}_{1,1}$ . All connected Lie groups with Lie algebra isomorphic to  $\mathfrak{su}_{1,1}$  are described as follows: the simply connected group (the universal cover of  $SU_{1,1}$ ) is the group  $V_{\infty}$ , and any non-simply connected group is isomorphic to  $V_n$  with  $n \in \mathbb{N}$ , where  $V_{\infty}$  and  $V_n$  are the Lie groups defined in Lemma 2.2. Clearly, the set  $\mathcal{C} := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0\}$  is  $V_{\infty}$ -invariant, and the set  $\mathcal{C}' := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0, w \neq 0\}$  is  $V_n$ -invariant for  $n \in \mathbb{N}$ .

Consider in  $V_j$  three one-parameter subgroups  $H_X$ ,  $H_Y$ ,  $H_Z$ 

for 
$$j = n \in \mathbb{N}$$
: 
$$z \mapsto z - \frac{i}{2}\beta, \quad w \mapsto w,$$

$$z \mapsto \lambda z, \qquad w \mapsto \lambda^{1/n}w,$$

$$z \mapsto \frac{z}{i\mu z + 1}, \quad w \mapsto \frac{1}{(i\mu z + 1)^{2/n}}w.$$

where  $\lambda > 0$ ,  $\beta, \mu \in \mathbb{R}$ ,  $t^{2/n} = \exp(2/n \ln_0 t)$  for  $t \in \mathbb{C} \setminus (-\infty, 0]$ , and  $\ln_0$  is the branch of the logarithm in  $\mathbb{C} \setminus (-\infty, 0]$  defined by the condition  $\ln_0 1 = 0$ . The vector fields on  $\mathbb{C}^2$  corresponding to these subgroups generate the Lie algebras of  $V_j$  for  $j \in \{1, 2, ..., \infty\}$  and are as follows:

for 
$$j = \infty$$
:  

$$X_3 := -\frac{i}{2} \partial/\partial z,$$

$$Y_3 := z \partial/\partial z + \partial/\partial w,$$

$$Z_3 := -iz^2 \partial/\partial z - 2iz \partial/\partial w,$$
for  $j = n \in \mathbb{N}$ :  

$$X_3 := -\frac{i}{2} \partial/\partial z,$$

$$Y_3 := z \partial/\partial z + \frac{1}{n} \partial/\partial w,$$

$$Z_3 := -iz^2 \partial/\partial z - \frac{2izw}{n} \partial/\partial w.$$
(3.6)

One can verify that these vector fields satisfy relations (R3).

Next, it is straightforward to observe that any subalgebra of  $\mathfrak{g}$  isomorphic to  $\mathfrak{h}$  is generated by either  $X + \eta Y - \eta^2/2 Z$ ,  $Y - \eta Z$ , with  $\eta \in \mathbb{R}$ , or by Y,

Z. For every  $j \in \{1, 2, ..., \infty\}$  the connected subgroups of  $V_j$  corresponding to the subalgebras generated by  $X_3 + \eta Y_3 - \eta^2/2 Z_3$ ,  $Y_3 - \eta Z_3$ , with  $\eta \in \mathbb{R}$ , or by  $Y_3$ ,  $Z_3$  are all conjugate to each other in  $V_j$  by elements of the subgroup  $W_j$  that we will describe below. Let first  $j = \infty$ . Transform  $\mathcal{C}$  into  $\{(z, w) : |z| < 1\}$  by means of the map  $\Phi_{\infty}$ 

$$z \mapsto \frac{z-1}{z+1},$$

$$w \mapsto w-2\ln_0((z+1)/2),$$
(3.7)

and let  $W_{\infty}$  be the subgroup of  $V_{\infty}$  that transforms under  $\Phi_{\infty}$  into the group of maps

$$\begin{array}{ccc} z & \mapsto & e^{it}z, \\ w & \mapsto & w+it, \end{array}$$

where  $t \in \mathbb{R}$ . Let now j = n. Transform C' into  $\{(z, w) : |z| < 1, w \neq 0\}$  by means of the map  $\Phi_n$ 

$$z \mapsto \frac{z-1}{z+1},$$

$$w \mapsto \left(\frac{2}{z+1}\right)^{2/n} w,$$
(3.8)

and let  $W_n$  be the subgroup of  $V_n$  that transforms under  $\Phi_n$  into the group of maps

$$\begin{array}{ccc} z & \mapsto & e^{it}z, \\ w & \mapsto & e^{it/n}w, \end{array}$$

where  $0 \leq t < 2\pi n$ . In particular, for every  $q \in O(p)$  the groups  $H_{q'}$  are pairwise distinct for q' sufficiently close to q and  $M_{q'}$  pairwise distinct, and every group  $H_q$  is isomorphic to  $\mathcal{T}$  (hence it acts effectively on  $M_q$ ).

Observe that – upon identification of G(M) and  $V_j$  for a particular value of j – for every  $q \in O(p)$  every element  $g \in G(M)$  can be written in the form g = g'g'' with  $g' \in W_j$ ,  $g'' \in H_q$ . Hence for every two points  $q_1, q_2 \in O(p)$  there exists  $g \in W_j$  such that  $gM_{q_1} = M_{q_2}$ . If for some  $q \in O(p)$  there is a non-trivial  $g \in W_j$  such that  $gM_q = M_q$ , then  $gH_qg^{-1} = H_q$  and hence g has

the form

for 
$$j=\infty$$
: 
$$z \mapsto z, \\ w \mapsto w+2\pi i k_0, \quad k_0 \in \mathbb{Z} \setminus \{0\},$$

for 
$$j = n \in \mathbb{N}$$
: 
$$z \mapsto z,$$
 
$$w \mapsto e^{2\pi i k_0/n} w, \quad 0 < k_0 \le n - 1.$$

Find  $h \in H_q$  such that  $gh \in I_q$ . However, every element of  $I_q$  has finite order (see (ii) of Proposition 1.1). If  $G(M) = V_{\infty}$ , then gh is clearly of infinite order. Assume that  $G(M) = V_n$  for some  $n \in \mathbb{N}$ . In this case gh can only be of finite order if  $h = \mathrm{id}$ . Hence  $g \in I_q$  and, since this argument can be applied to any point in  $M_q$ , we obtain that g fixes every point in  $M_q$ . Hence  $g = g_q$ , where, as before,  $g_q$  denotes the element of  $I_q$  corresponding to the non-trivial element in  $\mathbb{Z}_2$  (see (ii) of Proposition 1.1). Then for a pair of points  $q_1$ ,  $q_2$  close to q and such that  $M_{q_1} \neq M_{q_2}$  we obtain that  $H_{q_1} = g_q H_{q_2} g_q^{-1} = H_{q_2}$  which is impossible. The above contradictions show that in fact  $gM_q \neq M_q$  for every  $q \in O(p)$  and every non-trivial  $g \in W_j$ .

Suppose that  $G(M) = V_{\infty}$ . Fix  $q_0 \in O(p)$  and let  $f: M_{q_0} \to \mathcal{P}$  be a holomorphic equivalence that transforms  $H_{q_0}|_{M_{q_0}}$  into the group  $\mathcal{T}$  and such that  $\hat{X}_3|_{M_{q_0}}$  and  $\hat{Y}_3|_{M_{q_0}}$  are transformed into the vector fields  $i\partial/\partial z|_{\mathcal{P}}$  and  $z\partial/\partial z|_{\mathcal{P}}$ , respectively, where  $\hat{X}_3$ ,  $\hat{Y}_3$  are the vector fields on O(p) corresponding to  $X_3$ ,  $Y_3$ . For every  $q \in O(p)$  find the unique map  $g \in W_{\infty}$  such that  $gM_{q_0} = M_q$  and define  $F_{\infty}(q) := \left(f(g^{-1}(q)), it\right) \in \mathbb{C}^2$ , where t is the parameter value corresponding to g. Clearly,  $F_{\infty}$  is a real-analytic CR-isomorphism between O(p) and  $O_1$  that transforms  $\hat{X}_3 - 1/2 \hat{Z}_3$  into  $i\partial/\partial w|_{O_1}$ , where  $\hat{Z}_3$  is the vector field on O(p) corresponding to  $Z_3$ .

Denote by  $\tilde{X}$ ,  $\tilde{Y}$ ,  $\tilde{Z}$  the vector fields on  $O_1$  into which  $F_{\infty}$  transforms  $\hat{X}_3$ ,  $\hat{Y}_3$ ,  $\hat{Z}_3$ , respectively. Since  $F_{\infty}$  is real-analytic, it extends to a biholomorphic map from a neighborhood of O(p) in M onto a neighborhood of  $O_1$  in  $\mathbb{C}^2$ . Clearly,  $\hat{X}_3$ ,  $\hat{Y}_3$ ,  $\hat{Z}_3$  extend to holomorphic vector fields on all of M and hence  $\tilde{X}$ ,  $\tilde{Y}$ ,  $\tilde{Z}$  extend to holomorphic vector fields defined in a neighborhood of  $O_1$ . Since  $\tilde{X}$  and  $\tilde{Y}$  are tangent to  $O_1$  and since their restrictions to  $\mathcal{P} \times \{0\} \subset O_1$ 

are  $i\partial/\partial z$  and  $z\partial/\partial z$ , respectively, they have the form

$$\tilde{X} = (i + \rho(z, w))\partial/\partial z + \sigma(z, w)\partial/\partial w, 
\tilde{Y} = (z + \mu(z, w))\partial/\partial z + \nu(z, w)\partial/\partial w,$$
(3.9)

where  $\rho, \sigma, \mu, \nu$  are functions holomorphic near  $O_1$  such that

$$\rho(z,0) \equiv \sigma(z,0) \equiv \mu(z,0) \equiv \nu(z,0), \tag{3.10}$$

and

$$\operatorname{Re} \sigma(z, iv) \equiv \operatorname{Re} \nu(z, iv) \equiv 0.$$
 (3.11)

Since  $[X_3-1/2 Z_3, X_3] = -1/2 Y_3$  and  $[X_3-1/2 Z_3, Y_3] = 2X_3 - (X_3-1/2 Z_3)$ , we obtain

$$[\mathcal{Z}, \tilde{X}] = -\frac{1}{2}\tilde{Y},$$

$$[\mathcal{Z}, \tilde{Y}] = 2\tilde{X} - \mathcal{Z},$$
(3.12)

where  $\mathcal{Z} := i\partial/\partial w$ .

Conditions (3.12) together with (3.9) and (3.10) uniquely determine the functions  $\rho, \sigma, \mu, \nu$  as follows:

$$\rho = \frac{i}{4} \Big( (z+2)e^w - (z-2)e^{-w} \Big) - i,$$

$$\sigma = -\frac{i}{4} \Big( e^w + e^{-w} - 2 \Big),$$

$$\mu = \frac{1}{2} \Big( (z+2)e^w + (z-2)e^{-w} \Big) - z,$$

$$\nu = -\frac{1}{2} \Big( e^w - e^{-w} \Big).$$

The functions  $\sigma$ ,  $\nu$  automatically satisfy conditions (3.11).

Integrating  $\tilde{X}$ ,  $\tilde{Y}$ ,  $\mathcal{Z}$  we respectively obtain the following one-parameter

subgroups of holomorphic transformations of  $O_1$ :

$$z \mapsto \frac{1}{4} \left( z \left[ \left( 2 + e^w + e^{-w} \right) e^t + \left( 2 - e^w - e^{-w} \right) e^{-t} \right] + 2 \left( e^w - e^{-w} \right) \left( e^t - e^{-t} \right) \right),$$

$$H_{\tilde{Y}}:$$

$$w \mapsto \ln^w \frac{e^w + 1 + (e^w - 1)e^{-t}}{e^w + 1 - (e^w - 1)e^{-t}},$$

$$H_{\mathcal{Z}}: \begin{array}{ccc} z & \mapsto & z, \\ \\ w & \mapsto & w+it, \end{array}$$

where  $t \in \mathbb{R}$  and  $\ln^w := \ln'_0 + 2\pi i k$  for  $2\pi k < \operatorname{Im} w < 2\pi (k+1)$ , and  $\ln^{2\pi i k} \equiv 2\pi i k$ ,  $k \in \mathbb{Z}$ , with  $\ln'_0$  being the branch of the logarithm defined in  $\mathbb{C} \setminus [0, \infty)$  by the condition  $\ln'_0(-1) = \pi i$ .

We note that the group  $\tilde{V}_{\infty}$  of CR-transformations of  $O_1$  generated by  $H_{\tilde{X}}$ ,  $H_{\tilde{Y}}$  and  $H_Z$  is a Lie group isomorphic to  $V_{\infty}$  and that the action of  $\tilde{V}_{\infty}$  on  $O_1$  does not extend to an action on any neighborhood of  $O_1$  in  $\mathbb{C}^2$ ; more precisely, for every point  $p_0 = (z_0, w_0) \in \mathbb{C}^2$  with  $\operatorname{Re} w_0 \neq 0$  there exists an element of  $\tilde{V}_{\infty}$  that has a singularity at  $p_0$ .

Clearly, there exist a neighborhood W of O(p) in M (not necessarily G(M)-invariant) such that O(q) is of codimension 1 in M for every  $q \in W$  (it follows from [1] that in the presence of a codimension 1 orbit there may be at most two codimension 2 orbits in M). Choosing W sufficiently small, we can assume that  $F_{\infty}$  extends to W to a biholomorphic map onto a neighborhood

 $\mathcal{W}'$  of  $O_1$  in  $\mathbb{C}^2$ . Suppose first that for every  $q \in \mathcal{W}$  the orbit O(q) is Levi-flat. Let  $\hat{q} := F_{\infty}^{-1}(1,0)$  and  $q_m := F_{\infty}^{-1}(1,\epsilon_m)$  with  $\epsilon_m \to 0$  as  $m \to \infty$  (we assume that m is sufficiently large to ensure that  $(1,\epsilon_m) \in \mathcal{W}'$ ). Let  $M_{q_m}^{O(q_m)}$  be the leaf of the foliation on  $O(q_m)$  passing through  $q_m$ . It now follows that for every m there exists  $g_m \in W_{\infty}$ , such that  $g_m H_{\hat{q}} g_m^{-1}$  is the connected subgroup of G(M) corresponding to the subalgebra of vector fields on  $O(q_m)$  tangent to  $M_{q_m}^{O(q_m)}$ . Clearly, we can choose the sequence  $\{g_m\}$  so that  $g_m \to \mathrm{id}$  as  $m \to \infty$ . At the same time, conjugating the one-parameter subgroups  $H_{\tilde{X}}$  and  $H_{\tilde{Y}}$  by the element of  $H_{\mathcal{Z}}$  corresponding to  $g_m$  with  $t = t_m$  and computing the vector fields corresponding to the conjugated subgroups, we see that these vector fields at the points  $(1, \epsilon_m)$  span all of  $\mathbb{C}^2$  rather than a complex line in  $\mathbb{C}^2$  (the span coincides with that of the pair of vectors  $(3e^{\epsilon_m+it_m}, 1-e^{\epsilon_m+it_m})$  and  $(-1, 1-e^{\epsilon_m+it_m})$  which are linearly independent for large m).

Thus, for some  $q \in \mathcal{W}$  the orbit O(q) is strongly pseudoconvex. Since G(M) is isomorphic to  $V_{\infty}$ , the orbit O(q) is CR-equivalent to either the hypersurface  $\omega$  or a hypersurface  $\varepsilon_b$  for  $b \notin \mathbb{Q}$  (see Lemma 2.2). Suppose first that O(q) is CR-equivalent to  $\omega$  and let  $\Lambda$  be the equivalence map. Then  $\Lambda$  induces an automorphism of  $V_{\infty}$  (recall that G(M) is identified with  $V_{\infty}$ ), and hence an isomorphism  $\varphi_{\Lambda}$  between the Lie algebra  $\hat{\mathfrak{v}}_{\infty}$  generated by the vector fields  $\hat{X}_1$ ,  $\hat{Y}_1$ ,  $\hat{Z}_3$  on O(q) and the Lie algebra  $\mathfrak{v}_{\infty}$  of  $V_{\infty}$  (both these algebras are isomorphic to  $\mathfrak{su}_{1,1}$ ). Every automorphism of  $\mathfrak{v}_{\infty}$  is either an inner automorphism corresponding to the conjugation by an element of  $V_{\infty}$ , or is a composition of such an inner automorphism and the automorphism  $\varphi_0$  given by

$$X_3 \mapsto -X_3, \quad Y_3 \mapsto Y_3, \quad Z_3 \mapsto -Z_3.$$

Hence  $\Lambda$  can be chosen so that  $\varphi_{\Lambda}(\hat{X}_3-1/2\hat{Z}_3)=\pm(X_3-1/2Z_3)$ . Arguing as in the proof of Theorem 2.1, we see that  $\Lambda$  can be extended to a holomorphic equivalence between a neighborhood of O(q) and the domain  $\mathfrak{R}_{s,t}$  for some 0 < s < t. Then the map  $F_{\infty} \circ \Lambda^{-1}$  transforms the vector field  $X_3 - 1/2Z_3$  on an open subset of  $\mathfrak{R}_{s,t}$  into the vector field  $\pm i\partial/\partial w$ . It is not difficult to show that the general form of a locally defined biholomorphic map with this property is the composition  $\Gamma_{\infty} \circ \Psi_{\infty} \circ \Phi_{\infty}$ , where  $\Phi_{\infty}$  is defined in (3.7),  $\Psi_{\infty}$  is given by

$$\begin{array}{ccc} z & \mapsto & ze^{-w}, \\ w & \mapsto & w, \end{array}$$

and  $\Gamma_{\infty}$  is of the form

$$z \mapsto r(z),$$
  
 $w \mapsto \pm (w + \ln(Cz)),$ 

for some branch of the logarithm, holomorphic function r and  $C \in \mathbb{C}$ . A straightforward calculation shows, however, that the composition  $\Gamma_{\infty} \circ \Psi_{\infty} \circ \Phi_{\infty}$  does not transform any of the vector fields from  $\mathfrak{v}_{\infty}$  into any of the vector fields  $\tilde{X}$ ,  $\tilde{Y}$  on any open subset of  $\mathfrak{R}_{s,t}$ .

Suppose now that O(q) is CR-equivalent to  $\varepsilon_b$  for some  $b \notin \mathbb{Q}$ , and let  $\Lambda$  be the equivalence map. Then  $\Lambda$  induces an isomorphism  $V_{\infty} \to V_b$ , and hence an isomorphism  $\varphi_{\Lambda}$  between the respective Lie algebras  $\hat{\mathfrak{v}}_{\infty}$  and  $\mathfrak{v}_b$ . The algebra  $\mathfrak{v}_b$  is generated by the vector fields  $X_{3,b}, Y_{3,b}, Z_{3,b}$  defined as in (3.6) where n is replaced with b. Every automorphism of  $\mathfrak{v}_b$  is either an inner automorphism corresponding to the conjugation by an element of  $V_b$ , or is a composition of such an inner automorphism and the automorphism given by

$$X_{3,b} \mapsto -X_{3,b}, \quad Y_{3,b} \mapsto Y_{3,b}, \quad Z_{3,b} \mapsto -Z_{3,b}.$$

Hence  $\Lambda$  can be chosen so that  $\varphi_{\Lambda}(\hat{X}_3 - 1/2 \hat{Z}_3) = \pm (X_{3,b} - 1/2 Z_{3,b})$ . Again, arguing as in the proof of Theorem 2.1, we see that a neighborhood of O(q) is holomorphically equivalent to the domain  $\mathfrak{R}_{b,s,t}$  for some 0 < s < t. Then the map  $F_{\infty} \circ \Lambda^{-1}$  transforms the vector field  $X_{3,b} - 1/2 Z_{3,b}$  on an open subset of  $\mathfrak{R}_{b,s,t}$  into the vector field  $\pm i\partial/\partial w$ . It is not difficult to show that the general form of a locally defined biholomorphic map with this property is the composition  $\Gamma_{\infty} \circ \Psi_{\infty} \circ \Phi_{\infty} \circ \Theta_b$ , where  $\Gamma_{\infty}$ ,  $\Psi_{\infty}$ ,  $\Phi_{\infty}$  are as above and  $\Theta_b$  is given by

$$\begin{array}{ccc} z & \mapsto & z, \\ w & \mapsto & b \ln w. \end{array}$$

As before, one can verify by a straightforward calculation that  $\Gamma_{\infty} \circ \Psi_{\infty} \circ \Phi_{\infty} \circ \Theta_b$  does not transform any of the vector fields from  $\mathfrak{v}_b$  into any of the vector fields  $\tilde{X}$ ,  $\tilde{Y}$  on any open subset of  $\mathfrak{R}_{b,s,t}$ .

Suppose that  $G(M) = V_n$  for  $n \in \mathbb{N}$ . Fix  $q_0 \in O(p)$  and let  $f: M_{q_0} \to \mathcal{P}$  be a holomorphic equivalence that transforms  $H_{q_0}|_{M_{q_0}}$  into the group  $\mathcal{T}$  and such that  $\hat{X}_3|_{M_{q_0}}$  and  $\hat{Y}_3|_{M_{q_0}}$  are transformed into the vector fields  $i\partial/\partial z$  and  $z\partial/\partial z$ , respectively, where  $\hat{X}_3$ ,  $\hat{Y}_3$  are the vector fields on O(p) corresponding to  $X_3$ ,  $Y_3$ . For every  $q \in O(p)$  find the unique map  $g \in W_n$  such that  $gM_{q_0} = M_q$  and define  $F_n(q) := \left(f(g^{-1}(q)), e^{it/n}\right) \in \mathbb{C}^2$ , where t is the parameter

value corresponding to g. Clearly,  $F_n$  is a real-analytic CR-isomorphism between O(p) and  $O_2$  that transforms  $\hat{X}_3 - 1/2 \hat{Z}_3$  into  $iw/n \partial/\partial w|_{O_2}$ , where  $\hat{Z}_3$  is the vector field on O(p) corresponding to  $Z_3$ .

Denote by  $\tilde{X}$ ,  $\tilde{Y}$ ,  $\tilde{Z}$  the vector fields on  $O_2$  into which  $F_n$  transforms  $\hat{X}_3$ ,  $\hat{Y}_3$ ,  $\hat{Z}_3$ , respectively. Since  $F_n$  is real-analytic, it extends to a biholomorphic map from a neighborhood of O(p) in M onto a neighborhood of  $O_2$  in  $\mathbb{C}^2$ . Clearly,  $\hat{X}_3$ ,  $\hat{Y}_3$ ,  $\hat{Z}_3$  extend to holomorphic vector fields on all of M and hence  $\tilde{X}$ ,  $\tilde{Y}$ ,  $\tilde{Z}$  extend to holomorphic vector fields defined in a neighborhood of  $O_2$ . Since  $\tilde{X}$  and  $\tilde{Y}$  are tangent to  $O_2$  and since their restrictions to  $\mathcal{P} \times \{1\} \subset O_2$  are  $i\partial/\partial z$  and  $z\partial/\partial z$ , respectively, they have the form (3.9), where  $\rho$ ,  $\sigma$ ,  $\mu$ ,  $\nu$  are functions holomorphic near  $O_2$  such that

$$\rho(z,1) \equiv \sigma(z,1) \equiv \mu(z,1) \equiv \nu(z,1), \tag{3.13}$$

and

$$\sigma \overline{w} + \overline{\sigma} w \equiv \nu \overline{w} + \overline{\nu} w \equiv 0, \quad \text{for } |w| = 1.$$
 (3.14)

As before, we obtain

$$\left[\frac{iw}{n}\partial/\partial w, \tilde{X}\right] = -\frac{1}{2}\tilde{Y},$$

$$\left[\frac{iw}{n}\partial/\partial w, \tilde{Y}\right] = 2\tilde{X} - \frac{iw}{n}\partial/\partial w.$$
(3.15)

Conditions (3.15) together with (3.9) and (3.13) uniquely determine the functions  $\rho, \sigma, \mu, \nu$  as follows:

$$\rho = \frac{i}{4} ((z+2)w^n - (z-2)w^{-n}) - i,$$

$$\sigma = -\frac{i}{4n} (w^{n+1} + w^{1-n} - 2w),$$

$$\mu = \frac{1}{2} ((z+2)w^n + (z-2)w^{-n}) - z,$$

$$\nu = -\frac{1}{2n} (w^{n+1} - w^{1-n}).$$

The functions  $\sigma$ ,  $\nu$  automatically satisfy conditions (3.14).

Integrating  $\tilde{X}$ ,  $\tilde{Y}$ ,  $\mathcal{Z}$  we respectively obtain the following one-parameter subgroups of holomorphic transformations of  $O_2$ :

$$z \mapsto \frac{1}{4} \left( z \left[ \left( 2 + w^n + w^{-n} \right) e^t + \left( 2 - w^n - w^{-n} \right) e^{-t} \right] + 2 \left( w^n - w^{-n} \right) \left( e^t - e^{-t} \right) \right),$$

$$H_{\tilde{Y}}:$$

$$w \mapsto \left( \frac{w^n + 1 + (w^n - 1)e^{-t}}{w^n + 1 - (w^n - 1)e^{-t}} \right)^{1/n},$$

$$H_{\mathcal{Z}}: \begin{array}{ccc} z & \mapsto & z, \\ & & & \\ w & \mapsto & e^{it/n}w \end{array}$$

where  $t \in \mathbb{R}$ ,  $\tau^{1/n} = \exp(1/n \widetilde{\ln^w} \tau)$ , where  $\widetilde{\ln^w} := \ln_0' + 2\pi i k$  for  $2\pi i k/n < \arg w < 2\pi i (k+1)/n$ , and  $\widehat{\ln^{2\pi i k/n}} \equiv 2\pi i k$ ,  $0 \le k \le n-1$ , with  $\ln_0'$  defined as above.

We note that the group  $\tilde{V}_n$  of CR-transformations of  $O_2$  generated by  $H_{\tilde{X}}$ ,  $H_{\tilde{Y}}$  and  $H_{\mathcal{Z}}$  is a Lie group isomorphic to  $V_n$  and that the action of  $\tilde{V}_n$  on  $O_2$  does not extend to an action on any neighborhood of  $O_2$  in  $\mathbb{C}^2$  since for every point  $p_0 = (z_0, w_0) \in \mathbb{C}^2$  with  $|w_0| \neq 1$  there exists an element of  $\tilde{V}_n$  that has a singularity at  $p_0$ .

As before, there exist a neighborhood W of O(p) in M such that O(q) is a real hypersurface in M for every  $q \in W$ . Choosing W sufficiently small, we can assume that  $F_n$  extends to W to a biholomorphic map onto a neighborhood W' of  $O_2$  in  $\mathbb{C}^2$ . Suppose first that for every  $q \in W$  the orbit O(q) is

Levi-flat. Let  $\hat{q} := F_n^{-1}(1,1)$  and  $q_m := F_n^{-1}(1,\epsilon_m)$  with  $\epsilon_m \to 1$  as  $m \to \infty$ . Let  $M_{q_m}^{O(q_m)}$  be the leaf of the foliation on  $O(q_m)$  passing through  $q_m$ . It now follows that for every m there exists  $g_m \in W_n$ , such that  $g_m H_{\hat{q}} g_m^{-1}$  is the connected subgroup of G(M) corresponding to the subalgebra of vector fields on  $O(q_m)$  tangent to  $M_{q_m}^{O(q_m)}$ . The sequence  $\{g_m\}$  can be chosen so that  $g_m \to \mathrm{id}$  as  $m \to \infty$ .

At the same time, conjugating the one-parameter subgroups  $H_{\tilde{X}}$  and  $H_{\tilde{Y}}$  by the element of  $H_{\mathcal{Z}}$  corresponding to  $g_m$  with  $t=t_m$  and computing the vector fields corresponding to the conjugated subgroups, we see that these vector fields at the point  $(1, \epsilon_m)$  span all of  $\mathbb{C}^2$  rather than a complex line in  $\mathbb{C}^2$  (the span coincides with that of the pair of vectors  $(3\epsilon_m^n e^{it_m}, \epsilon_m/n - \epsilon_m^{n+1} e^{it_m}/n)$ ) and  $(-1, \epsilon_m/n - \epsilon_m^{n+1} e^{it_m}/n)$  which are linearly independent for large m).

Thus, for some  $q \in \mathcal{W}$  the orbit O(q) is strongly pseudoconvex. Since G(M) is isomorphic to  $V_n$ , the orbit O(q) is CR-equivalent to a hypersurface  $\varepsilon_{n/k}$ , where  $k \in \mathbb{N}$  is such that (n,k)=1 (see Lemma 2.2). Suppose first that O(q) is CR-equivalent to  $\omega$ . Let  $\Lambda$  be the equivalence map. Then  $\Lambda$  induces an isomorphism  $V_n \to V_{n/k}$ , and hence an isomorphism  $\varphi_{\Lambda}$  between the respective Lie algebras  $\hat{\mathfrak{v}}_n$  and  $\mathfrak{v}_{n/k}$ . Here  $\hat{\mathfrak{v}}_n$  is generated by the vector fields  $\hat{X}_3, \hat{Y}_3, \hat{Z}_3$  on O(q) and  $\mathfrak{v}_{n/k}$  is generated by the vector fields  $X_{3,n/k}, X_{3,n/k}, Z_{3,n/k}$  defined as in (3.6) where n is replaced with n/k. Every automorphism of  $\mathfrak{v}_{n/k}$  is either an inner automorphism corresponding to the conjugation by an element of  $V_{n/k}$ , or is a composition of such an inner automorphism and the automorphism given by

$$X_{3,n/k} \mapsto -X_{3,n/k}, \quad Y_{3,n/k} \mapsto Y_{3,n/k}, \quad Z_{3,n/k} \mapsto -Z_{3,n/k}.$$

Hence  $\Lambda$  can be chosen so that  $\varphi_{\Lambda}(\hat{X}_3 - 1/2\hat{Z}_3) = \pm (X_{3,n/k} - 1/2Z_{3,n/k})$ . Arguing as in the proof of Theorem 2.1, we see that a neighborhood of O(q) is holomorphically equivalent to the domain  $\mathfrak{R}_{n/k,s,t}$  for some 0 < s < t. Then the map  $F_n \circ \Lambda^{-1}$  transforms the vector field  $X_{3,n/k} - 1/2Z_{3,n/k}$  on an open subset of  $\mathfrak{R}_{n/k,s,t}$  into the vector field  $\pm iw\partial/\partial w$ . It is not difficult to show that the general form of a locally defined map with this property is the composition  $\Gamma_n \circ \Psi_n \circ \Phi_n \circ \Pi_k$ , where  $\Phi_n$  is defined in (3.8),  $\Psi_n$  is given by

$$\begin{array}{cccc} z & \mapsto & zw^{-n}, \\ w & \mapsto & w, \end{array}$$

 $\Gamma_n$  is of the form

$$\begin{array}{ccc} z & \mapsto & r(z), \\ w & \mapsto & \pm (Cz)^{1/n} w, \end{array}$$

for some branch of r, holomorphic function r and  $C \in \mathbb{R}$ , and  $\Pi_k$  is given by

$$\begin{array}{ccc} z & \mapsto & z, \\ w & \mapsto & w^{1/k}. \end{array}$$

It is straightforward to verify, however, that the composition  $\Gamma_n \circ \Psi_n \circ \Phi_n \circ \Pi_k$ does not transform any of the vector fields from  $\mathfrak{v}_{n/k}$  into any of the vector fields  $\tilde{X}$ ,  $\tilde{Y}$  on any open subset of  $\mathfrak{R}_{n/k,s,t}$ .

Thus, g cannot be given by relations (R3), which completes the proof of the proposition. 

We will now prove the following theorem that finalizes the classification of (2,3)-manifolds in the case when every orbit is a real hypersurface.

**THEOREM 3.2** Let M be a (2,3)-manifold. Assume that the G(M)-orbit of every point in M has codimension 1 and that at least one orbit is Levi-flat. Then M is holomorphically equivalent to one of the following domains:

- $S+i\mathbb{R}^2$ , where S is an angle with vertex at the origin of size less than  $\pi$  in the  $(\operatorname{Re} z, \operatorname{Re} w)$ -plane;
- (ii)  $\left\{ (z,w) \in \mathbb{C}^2 : s \left( \operatorname{Re} z \right)^b < \operatorname{Re} w < t \left( \operatorname{Re} z \right)^b, \operatorname{Re} z > 0 \right\}$ , where  $|b| \geq 1$ ,  $b \neq 1, -\infty \leq s < 0 < t \leq \infty$  and at least one of s, t is finite; (iii)  $\left\{ (z,w) \in \mathbb{C}^2 : \operatorname{Re} w > r_{s,t}(z) \left| \operatorname{Re} z \right|^b \right\}$ , where b > 1,

$$r_{s,t}(z) := \begin{cases} s & \text{if } \operatorname{Re} z > 0, \\ t & \text{if } \operatorname{Re} z < 0, \\ 0 & \text{if } \operatorname{Re} z = 0, \end{cases}$$

and  $s, t \in \mathbb{R}$  are such that either  $s \cdot t < 0$ , or  $s \ge 0$ ,  $t \ge 0$ ,  $s^2 + t^2 > 0$ , and in the latter case  $s \neq t$  for b = 2;

(iv) 
$$\{(z, w) \in \mathbb{C}^2 : 0 < Rew < r(z) | Rez|^b \}$$
, where  $b \le -1$ ,  $r(z)$  as in (iii), and  $s \ge 0$ ,  $t \ge 0$ ,  $s^2 + t^2 > 0$ ; (v)  $\{(z, w) \in \mathbb{C}^2 : Rez < R_{s,t}(w) \}$ , where

$$(v) \quad \{(z,w) \in \mathbb{C}^2 : \operatorname{Re} z < R_{s,t}(w)\}, \text{ where}$$

$$R_{s,t}(w) := \begin{cases} \operatorname{Re} w \cdot \ln(s\operatorname{Re} w) & \text{if } \operatorname{Re} w > 0, \\ \operatorname{Re} w \cdot \ln(-t\operatorname{Re} w) & \text{if } \operatorname{Re} w < 0, \\ 0 & \text{if } \operatorname{Re} w = 0, \end{cases}$$

with s, t > 0.

**Proof:** We will use the notation introduced earlier in this section. Observe that the set  $\mathfrak{L} := \{p \in M : O(p) \text{ is Levi-flat}\}$  is closed in M. Hence, if  $\mathfrak{L}$  is also open, then every orbit in M is Levi-flat. Let  $p \in \mathfrak{L}$  and suppose first that there exists a CR-isomorphism  $f: O(p) \to O_1$  that transforms G(M) into the group  $G_0$ . Since  $G_0$  acts on all of  $\mathcal{C}$ , arguing as on step (II) of the orbit gluing procedure introduced in the proof of Theorem 2.1, we extend f to a biholomorphic map between a G(M)-invariant neighborhood U of O(p) and a  $G_0$ -invariant neighborhood of  $O_1$  in  $\mathcal{C}$  that satisfies (2.45) for all  $g \in G(M)$  and  $g \in U$ , where  $g : G(M) \to G_0$  is an isomorphism. Since every  $G_0$ -orbit in  $\mathcal{C}$  has the form

$$\mathfrak{b}_r := \{(z, w) \in \mathcal{C} : \operatorname{Re} w = r\},\$$

for  $r \in \mathbb{R}$ , and hence is Levi-flat, the set  $\mathfrak{L}$  is open, and thus every orbit in M is Levi-flat. Since the group  $G_0$  is not isomorphic to any of the groups  $G_b$  for  $b \in \mathbb{R}^*$ , G and  $G'_0$ , it follows that every orbit in M is CR-equivalent to  $O_1$  by means of a map that transforms G(M) into  $G_0$ .

We will now further utilize the orbit gluing procedure from the proof of Theorem 2.1. Consider a maximal G(M)-invariant domain  $D \subset M$  from which there exists a biholomorphic map f onto a  $G_0$ -invariant domain in  $\mathcal{C}$  satisfying (2.45) for all  $g \in G(M)$  and  $q \in D$ . Assume that  $D \neq M$  and consider  $x \in \partial D$ . Let  $f_1 : O(x) \to O_1$  be a real-analytic CR-isomorphism satisfying (2.45) for all  $g \in G(M)$ ,  $q \in O(x)$  and an isomorphism  $\varphi : G(M) \to G_0$ . As before, extend  $f_1$  to a biholomorphic map from a connected G(M)-invariant neighborhood V of O(x) onto a  $G_0$ -invariant neighborhood of  $G_1$  in  $\mathcal{C}$ . The extended map satisfies (2.45) for all  $g \in G(M)$  and  $g \in V$ . Consider  $g \in V \cap D$ . The maps  $g \in V \cap D$  and  $g \in V \cap D$  are stablishes a  $G_0$ -invariant neighborhood  $g_0$ -orbits  $g_0$ -orbit

and  $g \in \operatorname{Aut}_{CR}(\mathfrak{b}_{r_1})$ . Each of the maps f and  $f_1$  transforms the group G(M) into the group  $G_0$ , and therefore g lies in the normalizer of  $G_0$  in  $\operatorname{Aut}_{CR}(\mathfrak{b}_{r_1})$ . The general form of an element of  $\operatorname{Aut}_{CR}(\mathfrak{b}_{r_1})$  is

$$(z, r_1 + iv) \mapsto (a_v(z), r_1 + i\mu(v)),$$
 (3.16)

where  $v \in \mathbb{R}$ ,  $a_v \in \operatorname{Aut}(\mathcal{P})$  for every v, and  $\mu$  is a diffeomorphism of  $\mathbb{R}$ . Considering g in this form, we obtain that  $a_v \in \mathcal{T}$  for every v and that  $a_{v_1}aa_{v_1}^{-1} = a_{v_2}aa_{v_2}^{-1}$  for all  $a \in \mathcal{T}$  and all  $v_1, v_2 \in \mathbb{R}$ . Therefore,  $a_{v_1}a_{v_2}^{-1}$  lies in the center of  $\mathcal{T}$ , which is trivial. Hence we obtain that  $a_{v_1} = a_{v_2}$  for all  $v_1, v_2$ . In addition, there exists  $k \in \mathbb{R}^*$  such that  $\mu^{-1}(v) + c \equiv \mu^{-1}(v + kc)$ , for  $c \in \mathbb{R}$ . Differentiating this identity with respect to c at 0 gives

$$\mu^{-1}(v) = v/k + t_0 \tag{3.17}$$

for some  $t_0 \in \mathbb{R}$ . Therefore, F extends to a holomorphic automorphism of C as the following map:

$$\begin{array}{ccc}
z & \mapsto & \lambda z + i\beta, \\
w & \mapsto & kw + \sigma - ikt_0,
\end{array}$$
(3.18)

where  $\lambda > 0$ ,  $\beta, \sigma \in \mathbb{R}$ .

Since  $O_1$  splits  $\mathcal{C}$ , for V small enough we have  $V = V_1 \cup V_2 \cup O(x)$ , where  $V_j$  are open connected non-intersecting sets. For each j,  $D \cap V_j$  is connected, and, if V is sufficiently small, each  $V_j$  is either a subset of D or disjoint from it. Suppose first that there is only one j for which  $D \cap V_j \neq \emptyset$ . In this case  $D \cap V$  is connected and  $V \setminus (D \cup O(x)) \neq \emptyset$ . Then the map  $\tilde{f}$  defined by formula (2.46) gives a biholomorphic extension of f to  $D \cup V$ . By construction,  $\tilde{f}$  satisfies (2.45) for  $g \in G(M)$  and  $q \in D \cup V$ . Since  $D \cup V$  is strictly larger than D, we obtain a contradiction with the maximality of D. Thus, in this case D = M, and hence M is holomorphically equivalent to a  $G_0$ -invariant domain in  $\mathcal{C}$ .

Suppose now that  $V_j \subset D$  for j = 1, 2. Applying formula (2.46) we can extend  $f|_{V_1}$  and  $f|_{V_2}$  to biholomorphic maps  $\hat{f}_1$ ,  $\hat{f}_2$ , respectively, from a neighborhood of O(x) into C; each of these maps satisfies (2.45). Let  $\hat{f}_j(O(x)) = \mathfrak{b}_{\rho_j}$ , j = 1, 2. Clearly,  $\rho_1 \neq \rho_2$ . The map  $\hat{F} := \hat{f}_2 \circ \hat{f}_1^{-1}$  is a CR-isomorphism from  $\mathfrak{b}_{\rho_1}$  onto  $\mathfrak{b}_{\rho_2}$ . It follows, as before, that  $\hat{F}$  has the form (3.18) with  $\sigma \neq \rho_1(1-k)$ . Let  $\Gamma$  denote the cyclic group of automorphisms of C generated by  $\hat{F}$ . This group acts freely properly discontinuously on C.

Let  $C_{\Gamma}$  be the complex manifold obtained by factoring C by the action of  $\Gamma$  and let  $\Pi_{\Gamma}: C \to C_{\Gamma}$  be the factorization map. Then the map  $\Pi_{\Gamma} \circ \mathcal{F}$ , where  $\mathcal{F}$  is given by formula (2.47) establishes holomorphic equivalence between M and  $C_{\Gamma}$ . Observe, however, that  $C_{\Gamma}$  is not hyperbolic, since it is covered by C which is not hyperbolic. This contradiction shows that in fact exactly one of  $V_j$ , j = 1, 2, is a subset of D, and hence M is holomorphically equivalent to a  $G_0$ -invariant domain in C, that is, to a domain of the form

$$\{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0, r < \operatorname{Re} w < R\},$$

for some  $0 \le r < R \le \infty$ . Thus, M is equivalent to  $\Delta^2$  which is impossible, since  $d(\Delta^2) = 6$ .

Next, if for  $p \in \mathfrak{L}$  there exists a CR-isomorphism between O(p) and  $O_2$  that transforms G(M) into the group  $G'_0$ , a similar argument gives that M is holomorphically equivalent to the product of  $\Delta \times A$ , where A is either an annulus or a punctured disk. This is impossible either since  $d(\Delta \times A) = 4$ .

Let now  $p \in \mathfrak{L}$  and suppose that there exists a CR-isomorphism  $f: O(p) \to O_1$  that transforms G(M) into the group  $G_1$ . The group  $G_1$  acts on

$$\mathcal{D} := \mathbb{C}^2 \setminus \left\{ (z, w) \in \mathbb{C}^2 : \operatorname{Re} z = \operatorname{Re} w = 0 \right\},\,$$

and, as as before, we can extend f to a biholomorphic map between a G(M)-invariant neighborhood U of O(p) and a  $G_1$ -invariant neighborhood of  $O_1$  in  $\mathcal{D}$  that satisfies (2.45) for all  $g \in G(M)$  and  $q \in U$ , where  $\varphi : G(M) \to G_1$  is an isomorphism. A  $G_1$ -orbit in  $\mathcal{D}$  is either of the form

$$\{(z, w) \in \mathbb{C}^2 : \operatorname{Re} w = r \operatorname{Re} z, \operatorname{Re} z > 0\},$$

or of the form

$$\{(z, w) \in \mathbb{C}^2 : \operatorname{Re} w = r \operatorname{Re} z, \operatorname{Re} z < 0\},\$$

for  $r \in \mathbb{R}$ , or coincides with either  $O'_1$  (see (3.5)), or

$$O_1'' := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z = 0, \operatorname{Re} w < 0\},$$
 (3.19)

and hence is Levi-flat. Therefore the set  $\mathfrak{L}$  is open, and thus every orbit in M is Levi-flat. Since the group  $G_1$  is not isomorphic to any of the groups  $G_b$  for  $b \in \mathbb{R}^*$ ,  $b \neq 1$ , and G, it follows that every orbit in M is CR-equivalent to  $O_1$  by means of a map that transforms G(M) into  $G_1$ .

As before, consider a maximal G(M)-invariant domain  $D \subset M$ ,  $x \in \partial D$ , and a CR-isomorphism  $f_1: O(x) \to O_1$ . Extend  $f_1$  to a biholomorphic map from a connected G(M)-invariant neighborhood V of O(x) onto a  $G_1$ -invariant neighborhood of  $O_1$  in D and consider  $s \in V \cap D$ . The maps f and  $f_1$  take O(s) onto some  $G_1$ -orbits  $o_1$ ,  $o_2$ , respectively. Clearly,  $F := f_1 \circ f^{-1}$  has the form  $F = \nu \circ g$ , where  $\nu$  is a map of one of the forms

with  $\sigma \in \mathbb{R}$ , and  $g \in \operatorname{Aut}_{CR}(o_1)$ . Any two  $G_1$ -orbits are equivalent by means of a map of one of the forms (3.20).

As before, g lies in the normalizer of  $G_1$  in  $\operatorname{Aut}_{CR}(o_1)$ . Let  $\mathfrak{X}$  be a map of the form (3.20) that transforms  $o_1$  into  $O_1$  and  $g_{\mathfrak{X}} := \mathfrak{X} \circ g \circ \mathfrak{X}^{-1}$ . Considering  $g_{\mathfrak{X}}$  in the general form (3.16) with  $r_1 = 0$  we see that for every  $\lambda > 0$ ,  $\beta, \gamma \in \mathbb{R}$ , the composition  $a_{\lambda v + \gamma} \circ a^{\lambda, \beta} \circ a_v^{-1}$ , where  $a^{\lambda, \beta}(z) := \lambda z + i\beta$ , belongs to  $\mathcal{T}$  and is independent of v. This implies that  $a_v(z) = \lambda_0 z + i(C_1 v + C_2)$  for some  $\lambda_0 > 0$ ,  $C_1, C_2 \in \mathbb{R}$ . Also, for every  $\lambda > 0$ ,  $\gamma \in \mathbb{R}$  there exist  $\lambda_1 > 0$ ,  $\gamma_1 \in \mathbb{R}$  such that  $\mu(\lambda \mu^{-1}v + \gamma) = \lambda_1 v + \gamma_1$ , which gives that  $\mu^{-1}$ , as before, has the form (3.17) for some  $k \in \mathbb{R}^*$ ,  $t_0 \in \mathbb{R}$ . It then follows that  $g_{\mathfrak{X}}$  extends to a holomorphic automorphism of  $\mathcal{D}$  as follows:

$$\begin{array}{rcl}
z & \mapsto & \lambda_0 z + C_1 w + i C_2, \\
w & \mapsto & k w - i k t_0,
\end{array}$$
(3.21)

and thus F extends to an automorphism of  $\mathcal{D}$  as well.

As before, choose a neighborhood V of O(x) such that  $V = V_1 \cup V_2 \cup O(x)$  and suppose first that there is only one j for which  $D \cap V_j \neq \emptyset$ . Arguing as in the case of the group  $G_0$ , we now see that M is holomorphically equivalent to a  $G_1$ -invariant domain in  $\mathcal{D}$ . Every such a domain clearly has the form  $S + i\mathbb{R}^2$ , where S is an angle with vertex at the origin in the  $(\operatorname{Re} z, \operatorname{Re} w)$ -plane. Such a domain is only hyperbolic if the size of S is less than  $\pi$ , and we obtain (i) of the theorem.

Suppose now that  $V_j \subset D$  for j = 1, 2. As before, applying formula (2.46) we can extend  $f|_{V_1}$  and  $f|_{V_2}$  to biholomorphic maps  $\hat{f}_1$ ,  $\hat{f}_2$ , respectively, from a neighborhood of O(x) into  $\mathcal{D}$ , and let  $\hat{f}_j(O(x)) = \hat{o}_j$ , j = 1, 2. Clearly, the  $G_1$ -orbits  $\hat{o}_1$  and  $\hat{o}_2$  are distinct. The map  $\hat{F} := \hat{f}_2 \circ \hat{f}_1^{-1}$  is a CR-isomorphism from  $\hat{o}_1$  onto  $\hat{o}_2$ , and it follows that  $\hat{F}$  has the same form as the map F introduced above, where  $\nu$  is non-trivial. Then the cyclic group

 $\Gamma$  generated by  $\hat{F}$  acts freely properly discontinuously on  $\mathcal{D}$ , which gives, as before, that M is biholomorphically equivalent to  $\mathcal{D}_{\Gamma} := \mathcal{D}/\Gamma$ . However,  $\mathcal{D}_{\Gamma}$  is not hyperbolic since it is covered by the non-hyperbolic domain  $\mathcal{D}$ , and thus exactly one of  $V_j$ , j=1,2, is a subset of D; hence M is holomorphically equivalent to a  $G_1$ -invariant domain in  $\mathcal{D}$ , as specified above.

Let now  $p \in \mathfrak{L}$  and suppose that there exists a CR-isomorphism  $f: O(p) \to O_1$  that transforms G(M) into the group  $G_b$  for some  $b \in \mathbb{R}^*$ ,  $b \neq 1$ . The group  $G_b$  acts on  $\mathcal{D}$ , and, as as before, we can consider an extension of f. Every  $G_b$ -orbit in  $\mathcal{D}$  is either strongly pseudoconvex and has one of the forms

$$\left\{ (z, w) \in \mathbb{C}^2 : \operatorname{Re} w = r \left( \operatorname{Re} z \right)^b, \operatorname{Re} z > 0 \right\},$$
$$\left\{ (z, w) \in \mathbb{C}^2 : \operatorname{Re} w = r \left( -\operatorname{Re} z \right)^b, \operatorname{Re} z < 0 \right\},$$

for  $r \in \mathbb{R}^*$ , or coincides with one of  $O_1$ ,  $O'_1$ ,  $O''_1$  (see (3.5), (3.19)), and

$$O_1''' := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z < 0, \operatorname{Re} w = 0\}.$$
 (3.22)

It then follows that every Levi-flat orbit in M has a G(M)-invariant neighborhood in which every other orbit is strongly pseudoconvex. Among the groups  $G_c$  for  $c \in \mathbb{R}^*$ ,  $c \neq 1$ , and G, the only group isomorphic to  $G_b$  is  $G_{1/b}$ . Thus, it follows that every Levi-flat orbit in M is CR-equivalent to  $O_1$  by means of a map that transforms G(M) into either  $G_b$  or  $G_{1/b}$ . In the latter case interchanging the variables we obtain a map that takes the Levi-flat orbit into  $O'_1$  and transforms G(M) into  $G_b$ . Next, it follows from Lemma 2.2 that every strongly pseudoconvex orbit is CR-equivalent to  $\tau_b$  (see (2.2)) by means of a CR-map that transforms G(M) into  $G_b$ .

As before, consider a maximal G(M)-invariant domain  $D \subset M$ ,  $x \in \partial D$  and a CR-map  $f_1: O(x) \to O$ , where O is either  $O_1$  or  $O'_1$  or  $\tau_b$ . As before, extend  $f_1$  to a biholomorphic map from a connected G(M)-invariant neighborhood V of O(x) onto a  $G_b$ -invariant neighborhood of O in  $\mathcal{D}$ , and consider  $s \in V \cap D$  such that O(s) is strongly pseudoconvex. The maps f and  $f_1$  take O(s) onto some strongly pseudoconvex  $G_b$ -orbits. Then  $F := f_1 \circ f^{-1}$  establishes a CR-isomorphism between the orbits, and thus has the form  $F = \nu \circ g$ , where  $\nu$  is a map of the form

$$\begin{array}{ccc}
z & \mapsto & \pm z, \\
w & \mapsto & kw
\end{array} \tag{3.23}$$

with  $k \in \mathbb{R}^*$ , and  $g \in G_b$ . Therefore, F extends to an automorphism of  $\mathcal{D}$ .

As before, choose a neighborhood V of O(x) such that  $V = V_1 \cup V_2 \cup O(x)$ . If there is only one j for which  $D \cap V_j \neq \emptyset$ , then, as before, we obtain that M is holomorphically equivalent to a  $G_b$ -invariant hyperbolic domain in  $\mathcal{D}$ . This leads to (ii), (iii), (iv) of the theorem (note that interchanging the variables transforms  $G_b$ -invariant domains into  $G_{1/b}$ -invariant domains).

Suppose now that  $V_j \subset D$  for j=1,2. As before, consider the CR-isomorphism  $\hat{F}: \hat{o}_1 \to \hat{o}_2$ . If O(x) is strongly pseudoconvex, then  $\hat{F} = \nu \circ g$ , where  $\nu$  is a non-trivial map of the form (3.23), and  $g \in G_b$ . Suppose now that O(x) is Levi-flat. Then  $\hat{F} = \nu \circ g$ , where  $\nu$  is a non-trivial map of one of the forms

and  $g \in \operatorname{Aut}_{CR}(\hat{o}_1)$ . In this case, either g lies in the normalizer of  $G_b$  in  $\operatorname{Aut}_{CR}(\hat{o}_1)$ , or  $gG_bg^{-1} = G_{1/b}$  and  $gG_{1/b}g^{-1} = G_b$ . Transforming  $\hat{o}_1$  into  $O_1$  by a map  $\mathfrak{X}$  of one of the forms (3.24) and arguing similarly to the case of the group  $G_1$ , we obtain that  $g_{\mathfrak{X}} := \mathfrak{X} \circ g \circ \mathfrak{X}^{-1}$  extends to a holomorphic automorphism of  $\mathcal{D}$  as a map of the form (3.21) with  $C_1 = 0$ .

It now follows that the cyclic group  $\Gamma$  generated by  $\hat{F}$  acts freely properly discontinuously on  $\mathcal{D}$ , which leads, as before, to holomorphic equivalence between M and the non-hyperbolic manifold  $\mathcal{D}/\Gamma$ . This contradiction shows that in fact exactly one of  $V_j$ , j=1,2, is a subset of D, and hence M is holomorphically equivalent to a  $G_b$ -invariant domain in  $\mathcal{D}$ .

Let now  $p \in \mathfrak{L}$  and suppose that there exists a CR-isomorphism  $f: O(p) \to O_1$  that transforms G(M) into the group G. The group G acts on  $\mathcal{D}$ , and we can consider an extension of f. Every G-orbit in  $\mathcal{D}$  is either strongly pseudoconvex and has one of the forms

$$\left\{ (z,w) \in \mathbb{C}^2 : \operatorname{Re} z = \operatorname{Re} w \ln (r \operatorname{Re} w), \operatorname{Re} w > 0 \right\},$$
 
$$\left\{ (z,w) \in \mathbb{C}^2 : \operatorname{Re} z = \operatorname{Re} w \ln (-r \operatorname{Re} w), \operatorname{Re} w < 0 \right\},$$

for r > 0, or coincides with one of  $O_1$ ,  $O_1'''$  (see (3.22)).

It then follows that every Levi-flat orbit in M has a G(M)-invariant neighborhood in which every other orbit is strongly pseudoconvex, that every Levi-flat orbit in M is CR-equivalent to  $O_1$  by means of a map that transforms G(M) into G and that every strongly pseudoconvex orbit is CR-equivalent to  $\xi$  by means of a map that transforms G(M) into G.

Consider a maximal G(M)-invariant domain  $D \subset M$ ,  $x \in \partial D$  and a CR-map  $f_1: O(x) \to O$ , where O is either  $O_1$  or  $\xi$ . As before, extend  $f_1$  to a biholomorphic map from a connected G(M)-invariant neighborhood V of O(x) onto a G-invariant neighborhood of O in D, and consider  $s \in V \cap D$  such that O(s) is strongly pseudoconvex. The maps f and  $f_1$  take O(s) onto some strongly pseudoconvex G-orbits. Then  $F := f_1 \circ f^{-1}$  establishes a CR-isomorphism between the orbits, and thus has the form  $F = \nu \circ g$ , where  $\nu$  is a map of the form

$$\begin{array}{ccc}
z & \mapsto & kz, \\
w & \mapsto & kw
\end{array} \tag{3.25}$$

with  $k \in \mathbb{R}^*$ , and  $g \in G$ . Therefore, F extends to an automorphism of  $\mathcal{D}$ .

As before, choose a neighborhood V of O(x) such that  $V = V_1 \cup V_2 \cup O(x)$ . If there is only one j for which  $D \cap V_j \neq \emptyset$ , then, as before, we obtain that M is holomorphically equivalent to a G-invariant hyperbolic domain in  $\mathcal{D}$ . This leads to (v) of the theorem.

Suppose now that  $V_j \subset D$  for j=1,2. As before, consider the CR-isomorphism  $\hat{F}: \hat{o}_1 \to \hat{o}_2$ . If O(x) is strongly pseudoconvex, then  $\hat{F} = \nu \circ g$ , where  $\nu$  is a non-trivial map of the form (3.25), and  $g \in G$ . Suppose now that O(x) is Levi-flat. Then  $\hat{F} = \nu \circ g$ , where  $\nu$  is map (3.25) with k=-1, and  $g \in \operatorname{Aut}_{CR}(\hat{o}_1)$ . The element g lies in the normalizer of G in  $\operatorname{Aut}_{CR}(\hat{o}_1)$ . Transforming  $\hat{o}_1$  into  $O_1$  by a map  $\mathfrak{X}$  of the form (3.25) with  $k=\pm 1$  and considering  $g_{\mathfrak{X}} := \mathfrak{X} \circ g \circ \mathfrak{X}^{-1}$  in the general form (3.16) with  $r_1 = 0$ , we obtain, as before, that  $\mu^{-1}$  has the form (3.17) for some  $k \in \mathbb{R}^*$ ,  $t_0 \in \mathbb{R}$  and that  $a_v(z) = \lambda_0 z + i\beta(v)$ , where  $\lambda_0 > 0$  and  $\beta(v)$  is a function satisfying for every  $\lambda > 0$  and  $\gamma \in \mathbb{R}$  the following condition:

$$\beta \left(\lambda \mu^{-1}(v) + \gamma\right) - \lambda \beta \left(\mu^{-1}(v)\right) + \lambda \ln \lambda \left(\lambda_0 \mu^{-1}(v) - v\right) + \lambda_0 \beta \equiv \text{const.}$$

This gives that  $g_{\mathfrak{X}}$  extends to an automorphism of  $\mathcal{D}$  as a map of the form (3.21). It now follows that the cyclic group  $\Gamma$  generated by  $\hat{F}$  acts freely properly discontinuously on  $\mathcal{D}$ , which leads, as before, to to holomorphic equivalence between M and the non-hyperbolic manifold  $\mathcal{D}/\Gamma$ . This contradiction shows that in in fact exactly one of  $V_j$ , j=1,2, is a subset of D, and hence M is holomorphically equivalent to a G-invariant domain in  $\mathcal{D}$ .

The theorem is proved.

## References

[1] Alekseevsky, A. V. and Alekseevsky, D. V., Riemann G-manifolds with one-dimensional orbit space, Ann. Global Anal. Geom. 11(1993), 197–211.

- [2] Cartan, E., Sur la géométrie pseudo-conforme des hypersurfaces de deux variables complexes, I Ann. Math. Pura Appl. 11(1932), 17–90 (or Oeuvres Completes II, 2, 1231–1304), II Ann. Scuola Norm. Sup. Pisa 1(1932), 333–354 (or Oeuvres Completes III, 2, 1217–1238).
- [3] Gifford, J. A., Isaev, A. V. and Krantz, S. G., On the dimensions of the automorphism groups of hyperbolic Reinhardt domains, *Illinois J. Math.* 44(2000), 602–618.
- [4] Isaev, A. V., Hyperbolic n-dimensional manifolds with automorphism group of dimension  $n^2$ , preprint, available from Mathematics ArXiv at http://front.math.ucdavis.edu/math.CV/0502152
- [5] Isaev, A. V., Hyperbolic manifolds of dimension n with automorphism group of dimension  $n^2 1$ , J. Geom. Analysis 15(2005), 239–259.
- [6] Isaev, A. V., Analogues of Rossi's map and E. Cartan's classification of homogeneous strongly pseudoconvex 3-dimensional hypersurfaces, preprint, available from Mathematics ArXiv at http://front.math.ucdavis.edu/math.CV/0504353
- [7] Isaev, A. V. and Krantz, S. G., On the automorphism groups of hyperbolic manifolds, *J. Reine Angew. Math.* 534(2001), 187–194.
- [8] Isaev, A. V. and Kruzhilin, N. G., Effective actions of  $SU_n$  on complex *n*-dimensional manifolds, *Illinois J. Math.* 48(2004), 37–57.
- [9] Kaup, W., Reelle Transformationsgruppen und invariante Metriken auf komplexen Räumen, *Invent. Math.* 3(1967), 43–70.
- [10] Kim, K.-T. and Verdiani, L., Complex n-dimensional manifolds with a real  $n^2$ -dimensional automorphism group, J. Geom. Analysis 14 (2004), 701–713.

- [11] Kobayashi, S., Hyperbolic Manifolds and Holomorphic Mappings, Marcel Dekker, New York 1970.
- [12] Rossi, H., Attaching analytic spaces to an analytic space along a pseudoconcave boundary, Proc. Conf. Complex Analysis, Minneapolis, 1964, Springer-Verlag, 1965, 242–256.

Department of Mathematics The Australian National University Canberra, ACT 0200 AUSTRALIA

E-mail: alexander.isaev@maths.anu.edu.au