# Hyperbolic 2-Dimensional Manifolds with 3-Dimensional Automorphism Groups I<sup>\*†</sup>

A. V. Isaev

Let M be a Kobayashi-hyperbolic 2-dimensional complex manifold and Aut(M) the group of holomorphic automorphisms of M. We showed earlier that if dim Aut(M) = 3, then Aut(M)-orbits are closed submanifolds in M of (real) codimension 1 or 2. In this paper we classify all connected Kobayashi-hyperbolic 2-dimensional manifolds with 3-dimensional automorphism groups in the case when every orbit has codimension 1. A classification in the case when codimension 2 orbits are present will appear elsewhere.

## 0 Introduction

If M is a connected *n*-dimensional Kobayashi-hyperbolic complex manifold, then the group  $\operatorname{Aut}(M)$  of holomorphic automorphisms of M is a (real) Lie group in the compact-open topology, of dimension d(M) not exceeding  $n^2+2n$ , with the maximal value occurring only for manifolds holomorphically equivalent to the unit ball  $B^n \subset \mathbb{C}^n$  [11], [9]. We are interested in describing hyperbolic manifolds with lower values of d(M).

In [7], [4] we completely classified manifolds with  $n^2 \leq d(M) < n^2 + 2n$ (partial classifications for  $d(M) = n^2$  were also obtained in [3] and [10]). Note that for  $d(M) = n^2$  the manifold M may not be homogeneous, which makes this case substantially more difficult than the case  $d(M) > n^2$ , where homogeneity always takes place [9]. A further decrease in d(M) almost immediately leads to unclassifiable cases. Indeed, no reasonable classification exists for n = 2, d(M) = 2, in which case  $d(M) = n^2 - 2$  (observe, for example, that the automorphism group of a generic Reinhardt domain in  $\mathbb{C}^2$  is 2-dimensional). While it is possible that there is some classification for  $n \geq 3$ ,  $d(M) = n^2 - 2$ , as well as for particular pairs n, d(M) with  $d(M) < n^2 - 2$  (see [3] for the special case of Reinhardt domains), the case

<sup>\*</sup>Mathematics Subject Classification: 32Q45, 32M05, 32M10.

<sup>&</sup>lt;sup>†</sup>**Keywords and Phrases:** Kobayashi-hyperbolic manifolds, holomorphic automorphism groups.

 $d(M) = n^2 - 1$  is probably the only remaining candidate to investigate for the existence of an explicit classification for every  $n \ge 2$ . It turns out that all hyperbolic manifolds with  $n \ge 2$ ,  $d(M) = n^2 - 1$  indeed can be explicitly described. The case  $n \ge 3$  was treated in [5]. In this paper we deal with the far more difficult case n = 2.

For simplicity we will call connected 2-dimensional hyperbolic manifolds with 3-dimensional automorphism group (2,3)-manifolds. As we will see below, the classification of such manifolds is the richest among all previously obtained classifications. We will work with the group  $G(M) := \operatorname{Aut}(M)^c$ the connected identity component of Aut(M). Since the G(M)-action on M is proper [9], for every  $p \in M$  its isotropy subgroup  $I_p := \{f \in G(M) :$ f(p) = p is compact in G(M) and the orbit  $O(p) := \{f(p) : f \in G(M)\}$  is a connected closed submanifold in M. It follows from Proposition 2.1 of [5] (see Proposition 1.1 below) that for every  $p \in M$  the orbit O(p) has (real) codimension 1 or 2 in M (in the latter case O(p) is either a complex curve or a totally real submanifold). In this paper we obtain a complete classification of (2,3)-manifolds M under the additional assumption that every G(M)-orbit is of codimension 1, and hence is either strongly pseudoconvex or Levi-flat (see Theorems 2.1 and 3.2). Due to the substantial length of this article we decided to publish our classification of (2,3)-manifolds with codimension 2 orbits elsewhere. We note, however, that the codimension 1 case presented here contains the vast majority of all possible types of (2,3)-manifolds.

The paper is organized as follows. In Section 1 (see Proposition 1.1) we list some basic properties of G(M)-orbits. The most important properties for the purposes of this paper are: every orbit has codimension 1 or 2, and every Levi-flat codimension 1 orbit is foliated by complex curves holomorphically equivalent to the unit disk  $\Delta$  (that we will almost always realize as the right half-plane  $\mathcal{P} := \{z \in \mathbb{C} : \operatorname{Re} z > 0\}$ ). The last fact will be extensively used in Section 3.

In Section 2 we deal with the case when every orbit is strongly pseudoconvex and classify all (2,3)-manifolds with this property in Theorem 2.1. An important ingredient in the proof of Theorem 2.1 is E. Cartan's classification of 3-dimensional homogeneous strongly pseudoconvex CR-manifolds [2], together with the explicit determination of all covers of the non simplyconnected hypersurfaces on Cartan's list [6].

In Section 3 we first describe Levi-flat orbits together with all possible actions of G(M) (see Proposition 3.1). This description is used to classify (2,3)-manifolds with Levi-flat orbits in Theorem 3.2.

### **1** Initial Classification of Orbits

In this section we will list some initial facts about G(M)-orbits that follow from the results of [5]. For  $p \in M$  let  $L_p := \{d_p f : f \in I_p\}$  be the linear isotropy subgroup of p, where  $d_p f$  is the differential of a map f at p. The group  $L_p$  is a compact subgroup of  $GL(T_p(M), \mathbb{C})$  isomorphic to  $I_p$  by means of the isotropy representation

$$I_p \to L_p, \quad f \mapsto d_p f.$$

Proposition 2.1 of [5] implies the following

**Proposition 1.1** Let M be a (2,3)-manifold. Fix  $p \in M$  and let  $V_p := T_p(O(p))$ . Then the following holds:

(i) The orbit O(p) is either a closed real hypersurface, or a closed complex curve, or a closed totally real 2-dimensional submanifold of M.

(ii) If O(p) is a real Levi-flat hypersurface, it is foliated by complex curves holomorphically equivalent to  $\Delta$ , and there exist coordinates on  $T_p(M)$  such with respect to the orthogonal decomposition  $T_p(M) = (V_p \cap iV_p)^{\perp} \oplus (V_p \cap iV_p)$ we have  $L_p \subset \{\pm id\} \times L'_p$ , where  $L'_p$  is a finite subgroup of  $U_1$ .

(iii) If O(p) is a complex curve, it is holomorphically equivalent to  $\Delta$ . Furthermore, there exist holomorphic coordinates (z, w) on  $T_p(M)$  in which  $V_p = \{z = 0\}$  and the identity component  $L_p^c$  of  $L_p$  is given by either the matrices

$$\left(\begin{array}{cc}a^{\frac{k_1}{k_2}} & 0\\0 & a\end{array}\right),$$

for some  $k_1, k_2 \in \mathbb{Z}$ ,  $(k_1, k_2) = 1$ ,  $k_2 \neq 0$ , or the matrices

$$\left(\begin{array}{cc}a&0\\0&1\end{array}\right),$$

where |a| = 1.

(iv) If O(p) is totally real, then  $T_p(M) = V_p \oplus iV_p$ , and there are coordinates on  $V_p$  such that every transformation from  $L_p^c$  has the form:  $v_1 + iv_2 \mapsto$  $Av_1 + iAv_2, v_1, v_2 \in V_p$ , where  $A \in SO_2(\mathbb{R})$ .

Below we only consider (2,3)-manifolds without codimension 2 orbits.

#### 2 The Case of Strongly Pseudoconvex Orbits

In this section we will give a complete classification of (2,3)-manifolds M for which every G(M)-orbit is a strongly pseudoconvex real hypersurface in M. We will start with a large number of examples of such manifolds.

(1) Fix  $b \in \mathbb{R}$  such that  $|b| \ge 1$ ,  $b \ne 1$ , and choose  $0 \le s < t \le \infty$ . Define

$$R_{b,s,t} := \left\{ (z,w) \in \mathbb{C}^2 : s \left( \operatorname{Re} z \right)^b < \operatorname{Re} w < t \left( \operatorname{Re} z \right)^b, \operatorname{Re} z > 0 \right\}.$$

All these domains are obviously hyperbolic and the group  $\operatorname{Aut}(R_{b,s,t}) = G(R_{b,s,t})$  consists of all the maps

$$\begin{array}{lcl} z & \mapsto & \lambda z + i\beta, \\ w & \mapsto & \lambda^b w + i\gamma, \end{array} \tag{2.1}$$

where  $\lambda > 0$  and  $\beta, \gamma \in \mathbb{R}$ . The  $G(R_{b,s,t})$ -orbits are the following pairwise CR-equivalent hypersurfaces:

$$O_{\alpha}^{R_b} := \left\{ (z, w) \in \mathbb{C}^2 : \operatorname{Re} w = \alpha \left( \operatorname{Re} z \right)^b, \operatorname{Re} z > 0 \right\}, \quad s < \alpha < t,$$

and we set

$$\tau_b := O_1^{R_b}.\tag{2.2}$$

(2) For  $0 \le s < t \le \infty$  with either s > 0 or  $t < \infty$  define

$$U_{s,t} := \left\{ (z, w) \in \mathbb{C}^2 : \operatorname{Re} w \cdot \ln \left( s \operatorname{Re} w \right) < \operatorname{Re} z < \operatorname{Re} w \cdot \ln \left( t \operatorname{Re} w \right), \operatorname{Re} w > 0 \right\}$$

All these domains are clearly hyperbolic and the group  $\operatorname{Aut}(U_{s,t}) = G(U_{s,t})$  consists of all the maps

$$\begin{array}{rcl}
z & \mapsto & \lambda z + (\lambda \ln \lambda)w + i\beta, \\
w & \mapsto & \lambda w + i\gamma,
\end{array}$$
(2.3)

where  $\lambda > 0$  and  $\beta, \gamma \in \mathbb{R}$ . The  $G(U_{s,t})$ -orbits are the following pairwise CR-equivalent hypersurfaces:

$$O_{\alpha}^{U} := \left\{ (z, w) \in \mathbb{C}^{2} : \operatorname{Re} z = \operatorname{Re} w \cdot \ln \left( \alpha \operatorname{Re} w \right), \operatorname{Re} w > 0 \right\}, \quad s < \alpha < t,$$

and we set

$$\xi := O_1^U. \tag{2.4}$$

(3) For  $0 \le s < t < \infty$  define

$$\mathfrak{S}_{s,t} := \left\{ (z, w) \in \mathbb{C}^2 : s < (\operatorname{Re} z)^2 + (\operatorname{Re} w)^2 < t \right\}.$$

All these domains are clearly hyperbolic and the group  $\operatorname{Aut}(\mathfrak{S}_{s,t})$  consists of all maps of the form

$$\begin{pmatrix} z \\ w \end{pmatrix} \mapsto A \begin{pmatrix} z \\ w \end{pmatrix} + i \begin{pmatrix} \beta \\ \gamma \end{pmatrix}, \qquad (2.5)$$

where  $A \in O_2(\mathbb{R})$  and  $\beta, \gamma \in \mathbb{R}$ . The subgroup  $G(\mathfrak{S}_{s,t})$  is given by the condition  $A \in SO_2(\mathbb{R})$ . The  $G(\mathfrak{S}_{s,t})$ -orbits are the following pairwise CR-equivalent hypersurfaces:

$$O_{\alpha}^{\mathfrak{S}} := \left\{ (z, w) \in \mathbb{C}^2 : (\operatorname{Re} z)^2 + (\operatorname{Re} w)^2 = \alpha \right\}, \quad s < \alpha < t,$$

and we set

$$\chi := O_1^{\mathfrak{S}}.\tag{2.6}$$

(4) For more details on this example see [6]. Let  $\Phi_{\chi}^{(\infty)} : \mathbb{C}^2 \to \mathbb{C}^2$  be the following map:

$$z \mapsto \exp(\operatorname{Re} z)\cos(\operatorname{Im} z) + i\operatorname{Re} w, w \mapsto \exp(\operatorname{Re} z)\sin(\operatorname{Im} z) + i\operatorname{Im} w.$$

It is easy to see that  $\Phi_{\chi}^{(\infty)}$  is an infinitely-sheeted covering map onto  $\mathbb{C}^2 \setminus \{\operatorname{Re} z = 0, \operatorname{Re} w = 0\}$ . Introduce on the domain of  $\Phi_{\chi}^{(\infty)}$  a complex structure defined by the condition that the map  $\Phi_{\chi}^{(\infty)}$  is holomorphic (the pull-back complex structure under  $\Phi_{\chi}^{(\infty)}$ ), and denote the resulting manifold by  $M_{\chi}^{(\infty)}$ .

Next, for an integer  $n \geq 2$  consider the map  $\Phi_{\chi}^{(n)}$  from  $\mathbb{C}^2 \setminus \{\operatorname{Re} z = 0, \operatorname{Re} w = 0\}$  onto itself defined as follows:

$$z \mapsto \operatorname{Re}\left(\left(\operatorname{Re} z + i\operatorname{Re} w\right)^{n}\right) + i\operatorname{Im} z,$$
  
$$w \mapsto \operatorname{Im}\left(\left(\operatorname{Re} z + i\operatorname{Re} w\right)^{n}\right) + i\operatorname{Im} w.$$

Denote by  $M_{\chi}^{(n)}$  the domain of  $\Phi_{\chi}^{(n)}$  with the pull-back complex structure under  $\Phi_{\chi}^{(n)}$ . For  $0 \le s < t < \infty$  define

$$\begin{split} \mathfrak{S}_{s,t}^{(\infty)} &:= \left\{ (z,w) \in M_{\chi}^{(\infty)} : (\ln s)/2 < \operatorname{Re} z < (\ln t)/2 \right\}, \\ \mathfrak{S}_{s,t}^{(n)} &:= \left\{ (z,w) \in M_{\chi}^{(n)} : s^{1/n} < (\operatorname{Re} z)^2 + (\operatorname{Re} w)^2 < t^{1/n} \right\}, \quad n \ge 2. \end{split}$$

The domains  $\mathfrak{S}_{s,t}^{(\infty)}$  and  $\mathfrak{S}_{s,t}^{(n)}$  are respectively an infinite- and *n*-sheeted cover of the domain  $\mathfrak{S}_{s,t}$  introduced in (3) and hence are hyperbolic.

The group Aut  $\left(\mathfrak{S}_{s,t}^{(\infty)}\right)$  is generated by  $G\left(\mathfrak{S}_{s,t}^{(\infty)}\right)$  that consists of the maps

$$\begin{array}{lll} z & \mapsto & z + i\beta, \\ w & \mapsto & w + a, \end{array} \tag{2.7}$$

where  $\beta \in \mathbb{R}$ ,  $a \in \mathbb{C}$ , and the map

$$\begin{array}{l} z \quad \mapsto \quad \overline{z}, \\ w \quad \mapsto \quad \overline{w}. \end{array} \tag{2.8}$$

The  $G\left(\mathfrak{S}_{s,t}^{(\infty)}\right)$ -orbits are the following pairwise CR-equivalent hypersurfaces:

$$O_{\alpha}^{\mathfrak{S}^{(\infty)}} := \left\{ (z, w) \in M_{\chi}^{(\infty)} : \operatorname{Re} z = \alpha \right\}, \quad (\ln s)/2 < \alpha < (\ln t)/2,$$

and we set

$$\chi^{(\infty)} := O_0^{\mathfrak{S}^{(\infty)}}.$$
 (2.9)

The group Aut  $\left(\mathfrak{S}_{s,t}^{(n)}\right)$  for  $n \geq 2$  is generated by  $G\left(\mathfrak{S}_{s,t}^{(n)}\right)$  that consists of the maps

$$z \mapsto \cos \varphi \cdot \operatorname{Re} z + \sin \varphi \cdot \operatorname{Re} w + i\left(\cos(n\varphi) \cdot \operatorname{Im} z + \sin(n\varphi) \cdot \operatorname{Im} w + \beta\right),$$

$$w \mapsto -\sin \varphi \cdot \operatorname{Re} z + \cos \varphi \cdot \operatorname{Re} w + i\left(-\sin(n\varphi) \cdot \operatorname{Im} z + \cos(n\varphi) \cdot \operatorname{Im} w + \gamma\right),$$

$$(2.10)$$

where  $\varphi, \beta, \gamma \in \mathbb{R}$ , and the map

$$\begin{array}{rcl}
z & \mapsto & z, \\
w & \mapsto & -w.
\end{array}$$
(2.11)

The  $G\left(\mathfrak{S}_{s,t}^{(n)}\right)$ -orbits are the following pairwise CR-equivalent hypersurfaces:

$$O_{\alpha}^{\mathfrak{S}^{(n)}} := \left\{ (z, w) \in M_{\chi}^{(n)} : (\operatorname{Re} z)^2 + (\operatorname{Re} w)^2 = \alpha \right\}, \quad s^{1/n} < \alpha < t^{1/n},$$

and we set

$$\chi^{(n)} := O_1^{\mathfrak{S}^{(n)}}.$$
 (2.12)

(5) Fix b > 0 and for  $0 < t < \infty$ ,  $e^{-2\pi b}t < s < t$  consider

$$V_{b,s,t} := \left\{ (z, w) \in \mathbb{C}^2 : se^{b\varphi} < r < te^{b\varphi} \right\},\$$

where  $(r, \varphi)$  denote the polar coordinates in the (Re *z*, Re *w*)-plane with  $\varphi$  varying from  $-\infty$  to  $\infty$  (thus, the boundary of  $V_{b,t,s} \cap \mathbb{R}^2$  consists of two spirals accumulating to the origin and infinity). All these domains are hyperbolic and Aut $(V_{b,s,t}) = G(V_{b,s,t})$  consists of all maps of the form

$$\begin{pmatrix} z \\ w \end{pmatrix} \mapsto e^{b\psi} \begin{pmatrix} \cos\psi & \sin\psi \\ -\sin\psi & \cos\psi \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix} + i \begin{pmatrix} \beta \\ \gamma \end{pmatrix}, \quad (2.13)$$

where  $\psi, \beta, \gamma \in \mathbb{R}$ . The  $G(V_{b,s,t})$ -orbits are the following pairwise CR-equivalent hypersurfaces:

$$O_{\alpha}^{V_b} := \left\{ (z, w) \in \mathbb{C}^2 : r = \alpha e^{b\varphi} \right\}, \quad s < \alpha < t,$$

and we set

$$\rho_b := O_1^{V_b}. \tag{2.14}$$

(6) For  $1 \leq s < t < \infty$  define

$$E_{s,t} := \left\{ (z:w:\zeta) \in \mathbb{CP}^2 : s|z^2 + w^2 + \zeta^2| < |z|^2 + |w|^2 + |\zeta|^2 < t|z^2 + w^2 + \zeta^2| \right\},$$

The domain  $E_{s,t}$  is hyperbolic for each t since it is covered in a 2-to-1 fashion by the manifold

$$\Big\{(z, w, \zeta) \in \mathbb{C}^3 : s < |z|^2 + |w|^2 + |\zeta|^2 < t, \ z^2 + w^2 + \zeta^2 = 1\Big\},\$$

A. V. Isaev

which is clearly hyperbolic (the covering map is  $(z, w, \zeta) \mapsto (z : w : \zeta)$ ). The group  $\operatorname{Aut}(E_{s,t}) = G(E_{s,t})$  is given by

$$\begin{pmatrix} z \\ w \\ \zeta \end{pmatrix} \mapsto A \begin{pmatrix} z \\ w \\ \zeta \end{pmatrix}, \qquad (2.15)$$

where  $A \in SO_3(\mathbb{R})$ . The orbits of the action of the group  $G(E_{s,t})$  on  $E_{s,t}$  are the following pairwise CR non-equivalent hypersurfaces:

$$\mu_{\alpha} := \{ (z : w : \zeta) \in \mathbb{CP}^2 : |z|^2 + |w|^2 + |\zeta|^2 = \alpha |z^2 + w^2 + \zeta^2| \}, \\ s < \alpha < t.$$
(2.16)

(7) For more details on this example see [6]. Let  $\mathcal{Q}_+$  be the variety in  $\mathbb{C}^3$  given by

$$z_1^2 + z_2^2 + z_3^2 = 1$$

Consider the map  $\Phi_{\mu} : \mathbb{C}^2 \setminus \{0\} \to \mathcal{Q}_+$  defined by the formulas

$$z_{1} = -i(z^{2} + w^{2}) + i\frac{z\overline{w} - w\overline{z}}{|z|^{2} + |w|^{2}},$$

$$z_{2} = z^{2} - w^{2} - \frac{z\overline{w} + w\overline{z}}{|z|^{2} + |w|^{2}},$$

$$z_{3} = 2zw + \frac{|z|^{2} - |w|^{2}}{|z|^{2} + |w|^{2}}.$$

This map was first introduced in [12]. It is straightforward to verify that  $\Phi_{\mu}$  is a 2-to-1 covering map onto  $\mathcal{Q}_{+} \setminus \mathbb{R}^{3}$ . Introduce on the domain of  $\Phi_{\mu}$  the pull-back complex structure under  $\Phi_{\mu}$  and denote the resulting complex manifold by  $M_{\mu}^{(4)}$ .

For  $1 \leq s < t < \infty$  define

$$\begin{aligned} E_{s,t}^{(2)} &:= \left\{ (z_1, z_2, z_3) \in \mathbb{C}^3 : s < |z_1|^2 + |z_2|^2 + |z_3|^2 < t \right\} \cap \mathcal{Q}_+, \\ E_{s,t}^{(4)} &:= \left\{ (z, w) \in M_{\mu}^{(4)} : \sqrt{(s-1)/2} < |z|^2 + |w|^2 < \sqrt{(t-1)/2} \right\}. \end{aligned}$$

These domains are respectively 2- and 4-sheeted covers of the domain  $E_{s,t}$  introduced in (6) and hence are hyperbolic.

The group  $\operatorname{Aut}\left(E_{s,t}^{(2)}\right)$  consists of the maps

$$\begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix} \mapsto A \begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix}, \qquad (2.17)$$

where  $A \in O_3(\mathbb{R})$ , and  $G\left(E_{s,t}^{(2)}\right)$  is given by the condition  $A \in SO_3(\mathbb{R})$ . The  $G\left(E_{s,t}^{(2)}\right)$ -orbits are the following pairwise non-equivalent hypersurfaces:

$$\mu_{\alpha}^{(2)} := \left\{ (z_1, z_2, z_3) \in \mathbb{C}^3 : |z_1|^2 + |z_2|^2 + |z_3|^2 = \alpha \right\} \cap \mathcal{Q}_+, \ s < \alpha < t.$$
(2.18)

The group Aut  $\left(E_{s,t}^{(4)}\right)$  is generated by  $G\left(E_{s,t}^{(4)}\right)$  that consists of the maps

$$\left(\begin{array}{c}z\\w\end{array}\right)\mapsto A\left(\begin{array}{c}z\\w\end{array}\right),\tag{2.19}$$

where  $A \in SU_2$ , and the map

$$z \mapsto i \frac{z(|z|^2 + |w|^2) - \overline{w}}{\sqrt{1 + (|z|^2 + |w|^2)^2}},$$

$$w \mapsto i \frac{w(|z|^2 + |w|^2) + \overline{z}}{\sqrt{1 + (|z|^2 + |w|^2)^2}}.$$
(2.20)

The  $G\left(E_{s,t}^{(4)}\right)$ -orbits are the following pairwise CR non-equivalent hypersurfaces:

$$\mu_{\alpha}^{(4)} := \left\{ (z, w) \in M_{\mu}^{(4)} : |z|^2 + |w|^2 = \sqrt{(\alpha - 1)/2} \right\}, \quad s < \alpha < t.$$
 (2.21)

(8) For  $-1 \le s < t \le 1$  define

$$\Omega_{s,t} := \left\{ (z,w) \in \mathbb{C}^2 : s|z^2 + w^2 - 1| < |z|^2 + |w|^2 - 1 < t|z^2 + w^2 - 1| \right\}.$$

Clearly,  $\Omega_{s,t}$  is bounded if t < 1. Further,  $\Omega_{s,1}$  is hyperbolic since it is contained in the hyperbolic domain

$$\left\{ (z,w) \in \mathbb{C}^2 : z, w \notin (-\infty, -1] \cup [1,\infty) \right\}.$$

A. V. Isaev

The group  $\operatorname{Aut}(\Omega_{s,t})$  for every t consists of the maps

$$\begin{pmatrix} z \\ w \end{pmatrix} \mapsto \frac{\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}}{c_1 z + c_2 w + d}, \qquad (2.22)$$

where

$$Q := \begin{pmatrix} a_{11} & a_{12} & b_1 \\ a_{21} & a_{22} & b_2 \\ c_1 & c_2 & d \end{pmatrix} \in SO_{2,1}(\mathbb{R}).$$
(2.23)

The group  $\operatorname{Aut}(\Omega_{s,t})$  has two connected components (that correspond to the connected components of  $SO_{2,1}(\mathbb{R})$ ), and its identity component  $G(\Omega_{s,t})$  is given by the condition  $a_{11}a_{22} - a_{12}a_{21} > 0$ . The orbits of  $G(\Omega_{s,t})$  on  $\Omega_{s,t}$  are the following pairwise CR non-equivalent hypersurfaces:

$$\nu_{\alpha} := \{ (z, w) \in \mathbb{C}^2 : |z|^2 + |w|^2 - 1 = \alpha |z^2 + w^2 - 1| \} \setminus \{ (x, u) \in \mathbb{R}^2 : x^2 + u^2 = 1 \}, \quad s < \alpha < t.$$

$$(2.24)$$

(9) For  $1 \le s < t \le \infty$  define

$$D_{s,t} := \left\{ (z,w) \in \mathbb{C}^2 : s|1+z^2-w^2| < 1+|z|^2-|w|^2 < t|1+z^2-w^2|, \\ \operatorname{Im}\left(z(1+\overline{w})\right) > 0 \right\},$$

where  $D_{s,\infty}$  is assumed not to include the complex curve

$$\{(z,w) \in \mathbb{C}^2 : w^2 = 1 + z^2, \operatorname{Im}(z(1+\overline{w})) > 0\}$$

All these domains lie in the hyperbolic domain

$$\left\{(z,w)\in\mathbb{C}^2:\operatorname{Im} z>0,\,w\not\in(-\infty,-1]\cup[1,\infty)\right\},\$$

hence they are hyperbolic. For every matrix  $Q \in SO_{2,1}(\mathbb{R})^c$  as in (2.23) consider the map

$$\begin{pmatrix} z \\ w \end{pmatrix} \mapsto \frac{\begin{pmatrix} a_{22} & b_2 \\ c_2 & d \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix} + \begin{pmatrix} a_{21} \\ c_1 \end{pmatrix}}{a_{12}z + b_1w + a_{11}}.$$
 (2.25)

The group  $\operatorname{Aut}(D_{s,t}) = G(D_{s,t})$  consists of all such maps. The orbits of  $G(D_{s,t})$  on  $D_{s,t}$  are the following pairwise CR non-equivalent hypersurfaces:

$$\eta_{\alpha} := \begin{cases} (z, w) \in \mathbb{C}^2 : 1 + |z|^2 - |w|^2 = \alpha |1 + z^2 - w^2|, \\ \operatorname{Im}(z(1 + \overline{w})) > 0 \end{cases}, \quad s < \alpha < t. \end{cases}$$
(2.26)

(10) For more details on this example see [6]. Let  $\mathcal{Q}_{-}$  be the variety in  $\mathbb{C}^{3}$  given by

$$z_1^2 + z_2^2 - z_3^2 = 1,$$

and let

$$\Sigma := \left\{ (z, w) \in \mathbb{C}^2 : |z|^2 - |w|^2 > 0 \right\}.$$

For every integer  $n \ge 2$  consider the map  $\Phi^{(n)}$  from  $\Sigma$  to  $\mathcal{Q}_-$  defined as follows:  $z\overline{w} + w\overline{z}$ 

$$z_{1} = -i(z^{n} + z^{n-2}w^{2}) - i\frac{zw + wz}{|z|^{2} - |w|^{2}},$$

$$z_{2} = z^{n} - z^{n-2}w^{2} + \frac{z\overline{w} - w\overline{z}}{|z|^{2} - |w|^{2}},$$

$$z_{3} = -2iz^{n-1}w - i\frac{|z|^{2} + |w|^{2}}{|z|^{2} - |w|^{2}}.$$

The above maps were introduced in [6]. Further, set

$$\begin{split} \Omega^{(n)}_{\nu} &:= \left\{ (z,w) \in \mathbb{C}^2 : 0 < |z|^n - |z|^{n-2} |w|^2 < 1 \right\}, \\ \Omega^{(n)}_{\eta} &:= \left\{ (z,w) \in \mathbb{C}^2 : |z|^n - |z|^{n-2} |w|^2 > 1 \right\}. \end{split}$$

Clearly,  $\Omega_{\nu}^{(n)} \subset \Sigma$  and  $\Omega_{\eta}^{(n)} \subset \Sigma$  for all  $n \geq 2$ . Let  $\Phi_{\nu}^{(n)}$  and  $\Phi_{\eta}^{(n)}$  be the restrictions of  $\Phi^{(n)}$  to  $\Omega_{\nu}^{(n)}$  and  $\Omega_{\eta}^{(n)}$ , respectively. A straightforward calculation shows that  $\Phi_{\nu}^{(n)}$  and  $\Phi_{\eta}^{(n)}$  are *n*-to-1 covering maps onto

$$\Omega_{\nu} := \left\{ (z_1, z_2, z_3) \in \mathbb{C}^3 : -1 < |z_1|^2 + |z_2|^2 - |z_3|^2 < 1, \\ \operatorname{Im} z_3 < 0 \right\} \cap \mathcal{Q}_{-}$$

and

$$\Omega_{\eta} := \left\{ (z_1, z_2, z_3) \in \mathbb{C}^3 : |z_1|^2 + |z_2|^2 - |z_3|^2 > 1, \\ \operatorname{Im}(z_2(\overline{z_1} + \overline{z_3})) > 0 \right\} \cap \mathcal{Q}_-,$$

respectively. We now introduce on  $\Omega_{\nu}^{(n)}$ ,  $\Omega_{\eta}^{(n)}$  the pull-back complex structures under the maps  $\Phi_{\nu}^{(n)}$ ,  $\Phi_{\eta}^{(n)}$ , respectively, and denote the resulting complex manifolds by  $M_{\nu}^{(n)}$ ,  $M_{\eta}^{(n)}$ .

Further, let  $\Lambda : \mathbb{C} \times \Delta \to \Sigma$  be the following covering map:

$$\begin{array}{rccc} z & \mapsto & e^z, \\ w & \mapsto & e^z w, \end{array}$$

where  $z \in \mathbb{C}$ ,  $w \in \Delta$  and  $\Delta$  is the unit disk. Define

$$U_{\nu} := \{(z, w) \in \mathbb{C}^2 : |w| < 1, \exp(2\operatorname{Re} z)(1 - |w|^2) < 1\}, U_{\eta} := \{(z, w) \in \mathbb{C}^2 : |w| < 1, \exp(2\operatorname{Re} z)(1 - |w|^2) > 1\}.$$

Denote by  $\Lambda_{\nu}$ ,  $\Lambda_{\eta}$  the restrictions of  $\Lambda$  to  $U_{\nu}$ ,  $U_{\eta}$ , respectively. Clearly,  $U_{\nu}$  covers  $M_{\nu}^{(2)}$  by means of  $\Lambda_{\nu}$ , and  $U_{\eta}$  covers  $M_{\eta}^{(2)}$  by means of  $\Lambda_{\eta}$ . We now introduce on  $U_{\nu}$ ,  $U_{\eta}$  the pull-back complex structures under the maps  $\Lambda_{\nu}$ ,  $\Lambda_{\eta}$ , respectively, and denote the resulting complex manifolds by  $M_{\nu}^{(\infty)}$ ,  $M_{\eta}^{(\infty)}$ .

For  $-1 \le s < t \le 1$  we now define

$$\begin{split} \Omega_{s,t}^{(n)} &:= \left\{ (z,w) \in M_{\nu}^{(n)} : \quad \sqrt{(s+1)/2} < |z|^n - |z|^{n-2}|w|^2 < \\ &\qquad \sqrt{(t+1)/2} \right\}, \quad n \ge 2, \\ \Omega_{s,t}^{(\infty)} &:= \left\{ (z,w) \in M_{\nu}^{(\infty)} : \quad \sqrt{(s+1)/2} < \exp\left(\operatorname{Re} z\right) (1-|w|^2) < \\ &\qquad \sqrt{(t+1)/2} \right\}. \end{split}$$

The domain  $\Omega_{s,t}^{(n)}$ ,  $n \geq 2$ , is an *n*-sheeted cover of the domain  $\Omega_{s,t}$  introduced in (8) and the domain  $\Omega_{s,t}^{(\infty)}$  is its infinitely-sheeted cover. In particular, each of  $\Omega_{s,t}^{(n)}$ ,  $\Omega_{s,t}^{(\infty)}$  is hyperbolic.

The group Aut  $\left(\Omega_{s,t}^{(n)}\right)$  is generated by  $G\left(\Omega_{s,t}^{(n)}\right)$  that consists of all maps of the form

$$z \mapsto z \sqrt[n]{(a+bw/z)^2},$$

$$w \mapsto z \frac{\overline{b} + \overline{a}w/z}{a+bw/z} \sqrt[n]{(a+bw/z)^2},$$
(2.27)

where  $\sqrt[n]{}$  is any branch of the *n*th root and  $|a|^2 - |b|^2 = 1$ , and the map

$$z \mapsto \overline{z} \left( \sqrt[n]{\frac{\left(1 + z^{n-1}w(1 - |w|^2/|z|^2)\right)^2}{1 - |z|^{2n}(1 - |w|^2/|z|^2)^2}} \right)',$$

$$w \mapsto -\frac{\overline{w}/\overline{z} + z^n(1 - |w|^2/|z|^2)}{1 + z^{n-1}w(1 - |w|^2/|z|^2)} \times$$

$$\overline{z} \left( \sqrt[n]{\frac{\left(1 + z^{n-1}w(1 - |w|^2/|z|^2)\right)^2}{1 - |z|^{2n}(1 - |w|^2/|z|^2)^2}} \right)',$$
(2.28)

for some branch  $\binom{n}{\sqrt{}}$  of  $\frac{n}{\sqrt{}}$ . The  $G\left(\Omega_{s,t}^{(n)}\right)$ -orbits are the following pairwise CR non-equivalent hypersurfaces:

$$\nu_{\alpha}^{(n)} := \left\{ (z, w) \in M_{\nu}^{(n)} : |z|^{n} - |z|^{n-2}|w|^{2} = \sqrt{(\alpha+1)/2} \right\}, \qquad (2.29)$$
$$s < \alpha < t.$$

The group Aut  $\left(\Omega_{s,t}^{(\infty)}\right)$  is generated by  $G\left(\Omega_{s,t}^{(\infty)}\right)$  that consists of the maps of the form  $z \mapsto z + \ln(a + bw)$ 

$$w \mapsto \frac{\overline{b} + \overline{a}w}{a + bw}, \qquad (2.30)$$

where ln is any branch of the logarithm and  $|a|^2 - |b|^2 = 1$ , and the map

$$z \mapsto \overline{z} + \ln' \left( -\frac{1 + e^{2z}w(1 - |w|^2)}{\sqrt{1 - \exp(4\operatorname{Re} z)(1 - |w|^2)^2}} \right),$$

$$w \mapsto -\frac{\overline{w} + e^{2z}(1 - |w|^2)}{1 + e^{2z}w(1 - |w|^2)},$$
(2.31)

for some branch  $\ln'$  of the logarithm. The  $G\left(\Omega_{s,t}^{(\infty)}\right)$ -orbits are the following

pairwise CR non-equivalent hypersurfaces:

$$\nu_{\alpha}^{(\infty)} := \left\{ (z, w) \in M_{\nu}^{(\infty)} : \exp\left(2\operatorname{Re} z\right) \left(1 - |w|^2\right) = \sqrt{(\alpha + 1)/2} \right\}, \quad (2.32)$$
$$s < \alpha < t.$$

Next, for  $1 \leq s < t \leq \infty$  we define

$$\begin{split} D_{s,t}^{(2)} &:= \left\{ (z_1, z_2, z_3) \in \mathbb{C}^3 : \ s < |z_1|^2 + |z_2|^2 - |z_3|^2 < t, \\ &\operatorname{Im}(z_2(\overline{z_1} + \overline{z_3})) > 0 \right\} \cap \mathcal{Q}_-, \\ D_{s,t}^{(2n)} &:= \left\{ (z, w) \in M_{\eta}^{(n)} : \quad \sqrt{(s+1)/2} < |z|^n - |z|^{n-2}|w|^2 < \\ &\sqrt{(t+1)/2} \right\}, \quad n \ge 2, \\ D_{s,t}^{(\infty)} &:= \left\{ (z, w) \in M_{\eta}^{(\infty)} : \quad \sqrt{(s+1)/2} < \exp\left(\operatorname{Re} z\right) (1 - |w|^2) < \\ &\sqrt{(t+1)/2} \right\}. \end{split}$$

The domain  $D_{s,t}^{(2n)}$ ,  $n \geq 1$ , is an 2*n*-sheeted cover of the domain  $D_{s,t}$  introduced in (9) and the domain  $D_{s,t}^{(\infty)}$  is its infinitely-sheeted cover. To obtain an *n*-sheeted cover of  $D_{s,t}$  for odd  $n \geq 3$ , the domain  $D_{s,t}^{(4n)}$  must be factored by the action of the cyclic group of four elements generated by the following automorphism of  $M_{\eta}^{(2n)}$ :

$$\begin{aligned} z &\mapsto i z^2 \overline{z} \left( \sqrt[n]{\frac{1 - |w|^2 / |z|^2 + z^{-2n} \overline{w} / \overline{z}}{\sqrt{|z|^{4n} (1 - |w|^2 / |z|^2)^2 - 1}}} \right)', \\ w &\mapsto i \frac{1 + z^{2n-1} w (1 - |w|^2 / |z|^2)}{\overline{w} / \overline{z} + z^{2n} (1 - |w|^2 / |z|^2)} \times \\ z^2 \overline{z} \left( \sqrt[n]{\frac{1 - |w|^2 / |z|^2 + z^{-2n} \overline{w} / \overline{z}}{\sqrt{|z|^{4n} (1 - |w|^2 / |z|^2)^2 - 1}}} \right)', \end{aligned}$$

for some branch  $\binom{n}{\sqrt{}}$  of  $\sqrt[n]{}$ . Let  $\Pi_n$  denote the corresponding factorization map. Then  $D_{s,t}^{(n)} := \Pi_n \left( D_{s,t}^{(4n)} \right)$  is an *n*-sheeted cover of  $D_{s,t}$ . Clearly, each of  $D_{s,t}^{(n)}$ ,  $D_{s,t}^{(\infty)}$  is hyperbolic. The group  $G\left(D_{s,t}^{(2)}\right)$  consists of all maps of the form (2.17) with  $A \in SO_{2,1}(\mathbb{R})^c$ , and the full group  $\operatorname{Aut}\left(D_{s,t}^{(2)}\right)$  is generated by  $G\left(D_{s,t}^{(2)}\right)$  and the map

$$\begin{array}{l} z_1 & \mapsto & -z_1, \\ z_2 & \mapsto & -z_2, \\ z_3 & \mapsto & -z_3. \end{array}$$

$$(2.33)$$

The  $G\left(D_{s,t}^{(2)}\right)$ -orbits are the following pairwise non-equivalent hypersurfaces:  $\eta_{\alpha}^{(2)} := \left\{ (z_1, z_2, z_3) \in \Omega_{\eta} : |z_1|^2 + |z_2|^2 - |z_3|^2 = \sqrt{(\alpha+1)/2}, \right\}, \quad (2.34)$   $s < \alpha < t,$ 

For  $n \geq 2$  the group  $G\left(D_{s,t}^{(2n)}\right)$  consists of all maps of the form (2.27), and the full group  $\operatorname{Aut}\left(D_{s,t}^{(2n)}\right)$  is generated by  $G\left(D_{s,t}^{(2n)}\right)$  and the map

$$z \mapsto z^{2}\overline{z} \left( \sqrt[n]{\frac{\left(1 - |w|^{2}/|z|^{2} + z^{-n}\overline{w}/\overline{z}\right)^{2}}{|z|^{2n}(1 - |w|^{2}/|z|^{2})^{2} - 1}} \right)',$$

$$w \mapsto \frac{1 + z^{n-1}w(1 - |w|^{2}/|z|^{2})}{\overline{w}/\overline{z} + z^{n}(1 - |w|^{2}/|z|^{2})} \times$$

$$z^{2}\overline{z} \left( \sqrt[n]{\frac{\left(1 - |w|^{2}/|z|^{2} + z^{-n}\overline{w}/\overline{z}\right)^{2}}{|z|^{2n}(1 - |w|^{2}/|z|^{2})^{2} - 1}} \right)',$$
(2.35)

for some branch  $\binom{n}{\sqrt{}}$  of  $\sqrt[n]{}$ . The  $G\left(D_{s,t}^{(2n)}\right)$ -orbits are the following pairwise CR non-equivalent hypersurfaces:

$$\eta_{\alpha}^{(2n)} := \left\{ (z, w) \in M_{\eta}^{(n)} : |z|^{n} - |z|^{n-2}|w|^{2} = \sqrt{(\alpha+1)/2} \right\}, \qquad (2.36)$$
$$s < \alpha < t, n \ge 2.$$

Next, the group Aut  $(D_{s,t}^{(n)})$  for odd  $n \ge 3$  is connected and consists of all lifts from the domain  $D_{1,\infty}$  to  $D_{1,\infty}^{(n)} = \prod_n (M_{\eta}^{(2n)})$  of maps of the form

(2.25). The  $G\left(D_{s,t}^{(n)}\right)$ -orbits are the following pairwise CR non-equivalent hypersurfaces:

$$\eta_{\alpha}^{(n)} := \Pi_n \left( \eta_{\alpha}^{(4n)} \right). \tag{2.37}$$

Finally, the group  $G\left(D_{s,t}^{(\infty)}\right)$  consists of all maps of the form (2.30), and the full group Aut  $\left(D_{s,t}^{(\infty)}\right)$  is generated by  $G\left(D_{s,t}^{(\infty)}\right)$  and the map

$$z \mapsto 2z + \overline{z} + \ln' \left( i \frac{1 - |w|^2 + e^{-2z} \overline{w}}{\sqrt{\exp(4\text{Re}\,z)(1 - |w|^2)^2 - 1}} \right),$$

$$w \mapsto \frac{1 + e^{2z} w(1 - |w|^2)}{\overline{w} + e^{2z}(1 - |w|^2)}.$$
(2.38)

The  $G\left(D_{s,t}^{(\infty)}\right)$ -orbits are the following pairwise CR non-equivalent hypersurfaces:

$$\eta_{\alpha}^{(\infty)} := \left\{ (z, w) \in M_{\eta}^{(\infty)} : \exp\left(2\operatorname{Re} z\right) (1 - |w|^2) = \sqrt{(\alpha + 1)/2} \right\}, \quad (2.39)$$

$$s < \alpha < t.$$

We are now ready to formulate the main result of this section.

**THEOREM 2.1** Let M be a (2,3)-manifold. Assume that the G(M)-orbit of every point in M is a strongly pseudoconvex real hypersurface. Then M

is holomorphically equivalent to one of the following manifolds:

$$\begin{array}{ll} (i) & R_{b,s,t}, \ b \in \mathbb{R}, \ |b| \geq 1, \ b \neq 1, \ for \ 0 \leq s < t \leq \infty; \\ (ii) & U_{s,t}, \ for \ 0 \leq s < t \leq \infty, \ with \ either \ s > 0 \ or \ t < \infty; \\ (iii) & \mathfrak{S}_{s,t}, \ 0 \leq s < t < \infty; \\ (iv) & \mathfrak{S}_{s,t}^{(\infty)}, \ 0 \leq s < t < \infty; \\ (v) & \mathfrak{S}_{s,t}^{(n)}, \ 0 \leq s < t < \infty, \ n \geq 2; \\ (vi) & V_{b,s,t}, \ b > 0, \ 0 < t < \infty, \ e^{-2\pi b}t < s < t; \\ (vii) & E_{s,t}, \ 1 \leq s < t < \infty; \\ (viii) & E_{s,t}^{(4)}, \ 1 \leq s < t < \infty; \\ (ix) & E_{s,t}^{(2)}, \ 1 \leq s < t < \infty; \\ (x) & \Omega_{s,t}, \ -1 \leq s < t \leq 1; \\ (xi) & \Omega_{s,t}^{(\infty)}, \ -1 \leq s < t \leq 1; \\ (xii) & \Omega_{s,t}^{(n)}, \ -1 \leq s < t \leq 1; \\ (xii) & \Omega_{s,t}^{(n)}, \ -1 \leq s < t \leq \infty; \\ (xiii) & D_{s,t}, \ 1 \leq s < t \leq \infty; \\ (xiv) & D_{s,t}^{(\infty)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s,t}^{(n)}, \ 1 \leq s < t \leq \infty; \\ (xv) & D_{s$$

**Proof:** In [2] E. Cartan classified all homogeneous 3-dimensional strongly pseudoconvex CR-manifolds. Since the G(M)-orbit of every point in M is such a manifold, every G(M)-orbit is equivalent to a manifold on Cartan's list. We reproduce Cartan's classification below together with the corresponding groups of CR-automorphisms. Note that all possible covers of the

hypersurfaces  $\chi$ ,  $\mu_{\alpha}$ ,  $\nu_{\alpha}$  and  $\eta_{\alpha}$  appear below as explicitly calculated in [6].

(a) 
$$S^3$$
;  
(b)  $\mathcal{L}_m := S^3/\mathbb{Z}_m, m \in \mathbb{N}, m \ge 2$ ;  
(c)  $\sigma := \{(z,w) \in \mathbb{C}^2 : \operatorname{Re} w = |z|^2\}$ ;  
(d)  $\varepsilon_b := \{(z,w) \in \mathbb{C}^2 : \operatorname{Re} z = \exp(\operatorname{Re} w)\}$ ;  
(e)  $\omega := \{(z,w) \in \mathbb{C}^2 : |w| = \exp(|z|^2)\}$ ;  
(f)  $\delta := \{(z,w) \in \mathbb{C}^2 : |w| = \exp(|z|^2)\}$ ;  
(g)  $\tau_b, b \in \mathbb{R}, |b| \ge 1, b \ne 1 \text{ (see } (2.2));$   
(h)  $\xi (\operatorname{see } (2.4));$   
(j)  $\chi (\operatorname{see } (2.6));$   
(j')  $\chi^{(\infty)} (\operatorname{see } (2.9));$   
(j')  $\chi^{(n)}, n \ge 2 (\operatorname{see } (2.12));$   
(k)  $\rho_b, b > 0 (\operatorname{see } (2.14));$   
(l)  $\mu_{\alpha}, \alpha > 1 (\operatorname{see } (2.16));$   
(l')  $\mu^{(4)}_{\alpha}, \alpha > 1 (\operatorname{see } (2.21));$   
(l')  $\mu^{(2)}_{\alpha}, \alpha > 1 (\operatorname{see } (2.21));$   
(m')  $\nu^{(\infty)}_{\alpha}, -1 < \alpha < 1, (\operatorname{see } (2.32));$   
(m')  $\nu^{(\infty)}_{\alpha}, -1 < \alpha < 1, n \ge 2 (\operatorname{see } (2.29));$   
(n)  $\eta_{\alpha}, \alpha > 1, (\operatorname{see } (2.39));$   
(n')  $\eta^{(\infty)}_{\alpha}, \alpha > 1, (\operatorname{see } (2.39));$   
(n')  $\eta^{(\infty)}_{\alpha}, \alpha > 1, n \ge 2 (\operatorname{see } (2.34), (2.36), (2.37)).$ 

The above hypersurfaces are pairwise CR non-equivalent. The corresponding groups of CR-automorphisms are as follows:

 $\underline{(a)\operatorname{Aut}_{CR}(S^3)}:$ 

$$\begin{pmatrix} z \\ w \end{pmatrix} \mapsto \frac{\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}}{c_1 z + c_2 w + d},$$

with  $Q \in SU_{2,1}$ , where

$$Q := \begin{pmatrix} a_{11} & a_{12} & b_1 \\ a_{21} & a_{22} & b_2 \\ c_1 & c_2 & d \end{pmatrix};$$

(b) Aut<sub>CR</sub>(
$$\mathcal{L}_m$$
),  $m \ge 2$ :  

$$\begin{bmatrix} \begin{pmatrix} z \\ w \end{bmatrix} \mapsto \begin{bmatrix} U \begin{pmatrix} z \\ w \end{bmatrix} \end{bmatrix},$$

where  $U \in U_2$  and  $[(z, w)] \in \mathcal{L}_m$  denotes the equivalence class of  $(z, w) \in S^3$ under the action of  $\mathbb{Z}_m$  embedded in  $U_2$  as a subgroup of scalar matrices;

$$\underbrace{(c) \operatorname{Aut}_{CR}(\sigma)}_{w \mapsto \lambda} : \qquad z \mapsto \lambda e^{i\varphi} z + a, \\ w \mapsto \lambda^2 w + 2\lambda e^{i\varphi} \overline{a} z + |a|^2 + i\gamma,$$
(2.40)

where  $\lambda \in \mathbb{R}^*, \varphi, \gamma \in \mathbb{R}, a \in \mathbb{C};$ 

where  $\lambda, \beta, \mu, \kappa, \varphi \in \mathbb{R}, \, \lambda \kappa + \mu \beta = 1;$ 

(e)  $\operatorname{Aut}_{CR}(\omega)$  :

$$z \mapsto \frac{\lambda z + i\beta}{i\mu z + \kappa},$$

$$w \mapsto w - 2\ln(i\mu z + \kappa) + i\gamma,$$
(2.42)

where  $\lambda, \beta, \mu, \kappa, \gamma \in \mathbb{R}, \, \lambda \kappa + \mu \beta = 1;$ 

 $\underbrace{(\mathbf{f})\operatorname{Aut}_{CR}(\delta)}_{z \mapsto e^{i\varphi}z + a,} \\ w \mapsto e^{i\psi}\exp\left(2e^{i\varphi}\overline{a}z + |a|^2\right)w,$ 

where  $\varphi, \psi \in \mathbb{R}, a \in \mathbb{C};$ (g)  $\operatorname{Aut}_{CR}(\tau_b)$  : see (2.1); (h)  $\operatorname{Aut}_{CR}(\xi)$  : see (2.3); (j)  $\operatorname{Aut}_{CR}(\chi)$  : see (2.5); (j')  $\operatorname{Aut}_{CR}(\chi^{(\infty)})$  : see (2.7), (2.8); (j")  $\operatorname{Aut}_{CR}(\chi^{(n)}), n \ge 2$  : see (2.10), (2.11); (k)  $\operatorname{Aut}_{CR}(\rho_b)$  : see (2.13); (l)  $\operatorname{Aut}_{CR}(\mu_{\alpha})$  : see (2.15); (l') Aut<sub>CR</sub>  $\left(\mu_{\alpha}^{(4)}\right)$  : see (2.19), (2.20); (l")  $\operatorname{Aut}_{CR}\left(\mu_{\alpha}^{(2)}\right)$  : see (2.17); (m)  $\operatorname{Aut}_{CR}(\nu_{\alpha})$  : see (2.22); (m') Aut<sub>CR</sub>  $\left(\nu_{\alpha}^{(\infty)}\right)$  : see (2.30), (2.31); (m") Aut<sub>*CR*</sub>  $\left(\nu_{\alpha}^{(n)}\right)$   $n \ge 2$  : see (2.27), (2.28); (n)  $\operatorname{Aut}_{CR}(\eta_{\alpha})$  : see (2.25);  $(\mathbf{n}')\operatorname{Aut}_{CR}\left(\eta_{\alpha}^{(\infty)}\right) : \operatorname{see} (2.30), (2.38);$ (n")  $\operatorname{Aut}_{CR}\left(\eta_{\alpha}^{(2)}\right)$  : see (2.17) with  $A \in SO_{2,1}(\mathbb{R})^c$ , (2.33);

(n") Aut<sub>CR</sub> 
$$\left(\eta_{\alpha}^{(2n)}\right) n \ge 2$$
: see (2.27), (2.35);

 $\frac{(n^{"})\operatorname{Aut}_{CR}\left(\eta_{\alpha}^{(n)}\right), n \geq 3 \text{ is odd}: \text{this group is connected and consists of}}{\text{all lifts from the domain } D_{1,\infty} \text{ to } D_{1,\infty}^{(n)}} \text{ of maps of the form (2.25) (see (10)).}$ 

We will now show that the presence of an orbit of a particular kind in M determines the group G(M) as a Lie group. Fix  $p \in M$  and suppose that O(p) is CR-equivalent to  $\mathfrak{m}$ , where  $\mathfrak{m}$  is one of the model hypersurfaces (a)–(n") listed above. In this case we say that  $\mathfrak{m}$  is the model for O(p). Since G(M) acts properly and effectively on O(p), the CR-equivalence induces an isomorphism between G(M) and a closed connected 3-dimensional subgroup  $R_{\mathfrak{m}}$  of  $\operatorname{Aut}_{CR}(\mathfrak{m})$  that acts transitively on  $\mathfrak{m}$  (we will often say that the CR-equivalence transforms G(M) into  $R_{\mathfrak{m}}$ ). The subgroup  $R_{\mathfrak{m}}$  a priori depends on the choice of CR-equivalence between O(p) and  $\mathfrak{m}$ , but, as we will see below, this dependence is insignificant.

We will now list all possible groups  $R_{\mathfrak{m}}$  for each model (a)-(n"). In the following lemma  $\mathcal{P}$  denotes the right half-plane  $\{z \in \mathbb{C} : \operatorname{Re} z > 0\}$ .

#### Lemma 2.2 We have

- (A)  $R_{\mathfrak{m}} = Aut_{CR}(\mathfrak{m})^c$ , if  $\mathfrak{m}$  is one of (g)-(n");
- (B)  $R_{S^3}$  is conjugate in  $Aut_{CR}(S^3)$  to  $SU_2$ ,
- (C)  $R_{\mathcal{L}_m} = SU_2/(SU_2 \cap \mathbb{Z}_m), \ m \ge 2;$
- (D)  $R_{\sigma}$  is the Heisenberg group, that is, it consists of all elements of  $Aut_{CR}(\sigma)$  with  $\lambda = 1$ ,  $\varphi = 0$  in formula (2.40);
- (E)  $R_{\varepsilon_b}$  is either the subgroup of  $Aut_{CR}(\varepsilon_b)$  corresponding to a subgroup of  $Aut(\mathcal{P})$ , conjugate in  $Aut(\mathcal{P})$  to the subgroup  $\mathcal{T}$  given by

$$z \mapsto \lambda z + i\beta, \tag{2.43}$$

where  $\lambda > 0, \beta \in \mathbb{R}$ , or, for  $b \in \mathbb{Q}$ , is the subgroup  $V_b$  given by  $\varphi = 0$  in formula (2.41);

- (F)  $R_{\omega}$  is either the subgroup of  $Aut_{CR}(\omega)$  corresponding to a subgroup of  $Aut(\mathcal{P})$  conjugate in  $Aut(\mathcal{P})$  to the subgroup  $\mathcal{T}$  specified in (E), or is the subgroup  $V_{\infty}$  given by  $\gamma = 0$  in formula (2.42);
- (G)  $R_{\delta}$  coincides with the subgroup of  $Aut_{CR}(\delta)$  given by  $\varphi = 0$ .

**Proof:** Statement (A) is clear since in cases (g)-(n") we have dimAut<sub>CR</sub>( $\mathfrak{m}$ ) = d(M) = 3. Further, statements (B)–(D), (G) are obtained as in Lemma 3.4 of [5].

To prove statements (E) and (F) note that every codimension 1 subgroup of  $\operatorname{Aut}(\mathcal{P})$  is conjugate in  $\operatorname{Aut}(\mathcal{P})$  to the subgroup  $\mathcal{T}$  defined in (2.43).

The lemma is proved.

Lemma 2.2 implies, in particular, that if for some point  $p \in M$  the model for O(p) is  $S^3$ , then M admits an effective action of  $SU_2$ . Therefore, Mis holomorphically equivalent to one of the manifolds listed in [8]. However, none of the hyperbolic manifolds with 3-dimensional automorphism group on the list has a spherical orbit. Hence  $S^3$  is never the model of a G(M)-orbit.

We now observe that for each  $\mathfrak{m}$  the group  $\operatorname{Aut}_{CR}(\mathfrak{m})$  acts by holomorphic transformations on a certain complex manifold  $M_{\mathfrak{m}}$  containing  $\mathfrak{m}$ , and every  $R_{\mathfrak{m}}$ -orbit in  $M_{\mathfrak{m}}$  either is CR-equivalent to  $\mathfrak{m}$  (cases (b)–(k)), or belongs to the same family to which  $\mathfrak{m}$  belongs and the orbits are pairwise CR nonequivalent (cases (l)–(n")). The manifolds  $M_{\mathfrak{m}}$  are as follows:

- (b)  $M_{\mathcal{L}_m} = \mathbb{C}^2 \setminus \{0\} / \mathbb{Z}_m, \ m \ge 2;$
- (c)  $M_{\sigma} = \mathbb{C}^2;$
- (d)  $M_{\varepsilon_b} = \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0, w \neq 0\};$
- (e)  $M_{\omega} = \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0\};$
- (f)  $M_{\delta} = \{(z, w) \in \mathbb{C}^2 : w \neq 0\};$
- (g)  $M_{\tau_b} = \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0, \operatorname{Re} w > 0\};$
- (h)  $M_{\xi} = \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} w > 0\};$
- (j)  $M_{\chi} = \mathbb{C}^2 \setminus \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z = 0, \operatorname{Re} w = 0\};$
- (j')  $M_{\chi^{(\infty)}} = M_{\chi}^{(\infty)}$  (see (4));
- (j")  $M_{\chi^{(n)}} = M_{\chi}^{(n)}, n \ge 2$  (see (4));

$$\begin{split} &(\mathbf{k}) \ M_{\rho_{0}} = \mathbb{C}^{2} \setminus \{(z,w) \in \mathbb{C}^{2} : \operatorname{Re} z = 0, \operatorname{Re} w = 0\}; \\ &(\mathbf{l}) \ M_{\mu_{\alpha}} = \bigcup_{\alpha>1} \mu_{\alpha} = \mathbb{CP}^{2} \setminus \mathbb{RP}^{2}; \\ &(\mathbf{l}') \ M_{\mu_{\alpha}^{(4)}} = \bigcup_{\alpha>1} \mu_{\alpha}^{(4)} = M_{\mu}^{(4)} \ (\operatorname{see} \ (\mathbf{7})); \\ &(\mathbf{l}'') \ M_{\mu_{\alpha}^{(2)}} = \bigcup_{\alpha>1} \mu_{\alpha}^{(2)} = \mathcal{Q}_{+} \setminus \mathbb{R}^{3} \ (\operatorname{see} \ (\mathbf{7})); \\ &(\mathbf{m}) \ M_{\nu_{\alpha}} = \bigcup_{-1<\alpha<1} \nu_{\alpha} = \Omega_{-1,1} \ (\operatorname{see} \ (\mathbf{8})); \\ &(\mathbf{m}') \ M_{\nu_{\alpha}^{(\infty)}} = \bigcup_{-1<\alpha<1} \nu_{\alpha}^{(\infty)} = M_{\nu}^{(\infty)} = \Omega_{-1,1}^{(\infty)} \ (\operatorname{see} \ (\mathbf{10})); \\ &(\mathbf{m}'') \ M_{\nu_{\alpha}^{(n)}} = \bigcup_{-1<\alpha<1} \nu_{\alpha}^{(n)} = M_{\nu}^{(n)} = \Omega_{-1,1}^{(n)}, \ n \geq 2 \ (\operatorname{see} \ (\mathbf{10})); \\ &(\mathbf{n}) \ M_{\eta_{\alpha}} = \bigcup_{\alpha>1} \eta_{\alpha} = D_{1,\infty} \ (\operatorname{see} \ (\mathbf{10})); \\ &(\mathbf{n}) \ M_{\eta_{\alpha}^{(\infty)}} = \bigcup_{\alpha>1} \eta_{\alpha}^{(\infty)} = M_{\eta}^{(\infty)} = D_{1,\infty}^{(\infty)} \ (\operatorname{see} \ (\mathbf{10})); \\ &(\mathbf{n}'') \ M_{\eta_{\alpha}^{(2)}} = \bigcup_{\alpha>1} \eta_{\alpha}^{(2)} = \Omega_{\eta} \ (\operatorname{see} \ (\mathbf{10})); \\ &(\mathbf{n}'') \ M_{\eta_{\alpha}^{(2n)}} = \bigcup_{\alpha>1} \eta_{\alpha}^{(2n)} = M_{\eta}^{(n)} = D_{1,\infty}^{(2n)}, \ n \geq 2 \ (\operatorname{see} \ (\mathbf{10})); \\ &(\mathbf{n}'') \ M_{\eta_{\alpha}^{(2n)}} = \bigcup_{\alpha>1} \eta_{\alpha}^{(2n)} = M_{\eta}^{(n)} = D_{1,\infty}^{(2n)}, \ n \geq 2 \ (\operatorname{see} \ (\mathbf{10})); \\ &(\mathbf{n}'') \ M_{\eta_{\alpha}^{(2n)}} = \bigcup_{\alpha>1} \eta_{\alpha}^{(2n)} = M_{\eta}^{(n)} = D_{1,\infty}^{(2n)}, \ n \geq 2 \ (\operatorname{see} \ (\mathbf{10})); \\ &(\mathbf{n}'') \ M_{\eta_{\alpha}^{(2n)}} = \bigcup_{\alpha>1} \eta_{\alpha}^{(n)} = D_{1,\infty}^{(n)}, \ n \geq 3 \ \operatorname{is odd} \ (\operatorname{see} \ (\mathbf{10})). \end{split}$$

In cases (b)–(k) every two  $R_{\mathfrak{m}}\text{-}\mathrm{orbits}$  are  $CR\text{-}\mathrm{equivalent}$  (and equivalent

to  $\mathfrak{m}$ ) by means of automorphisms of  $M_{\mathfrak{m}}$  of the simple forms specified below:

(b) 
$$[(z,w)] \mapsto [(az,aw)], a > 0;$$
  
(c)  $z \mapsto z, w \mapsto w + a, a \in \mathbb{R};$   
(d)  $z \mapsto az, w \mapsto w, a > 0;$   
(e) as in (d);  
(f)  $z \mapsto z, w \mapsto aw, a > 0;$   
(g) as in (f); (2.44)  
(h)  $z \mapsto az, w \mapsto aw, a > 0;$   
(j) as in (h);  
(j')  $z \mapsto z + a, w \mapsto w, a \in \mathbb{R};$   
(j'') as in (h);  
(k) as in (h).

We will now show how the models can be glued together to form hyperbolic manifolds. The gluing procedure described below is similar to that used in [4], [5]. The procedure comprises the following steps:

(I). Start with a real hypersurface orbit O(p) with model  $\mathfrak{m}$  and consider a real-analytic CR-isomorphism  $f: O(p) \to \mathfrak{m}$ . Clearly, f satisfies

$$f(gq) = \varphi(g)f(q), \qquad (2.45)$$

for all  $g \in G(M)$  and  $q \in O(p)$ , where  $\varphi : G(M) \to R_{\mathfrak{m}}$  is an isomorphism.

(II). Observe that f can be extended to a biholomorphic map from a G(M)invariant connected neighborhood of O(p) in M onto an  $R_{\mathfrak{m}}$ -invariant neighborhood of  $\mathfrak{m}$  in  $M_{\mathfrak{m}}$ . First of all, extend f to some neighborhood U of O(p)to a biholomorphic map onto a neighborhood of  $\mathfrak{m}$  in  $M_{\mathfrak{m}}$ . Fix  $s \in U$  and  $s_0 \in$  O(s). Choose  $h_0 \in G(M)$  such that  $s_0 = h_0 s$  and define  $f(s_0) := \varphi(h_0) f(s)$ . To see that f is well-defined at  $s_0$ , suppose that for some  $h_1 \in G(M)$ ,  $h_1 \neq h_0$ , we have  $s_0 = h_1 s$ , and show that  $\varphi(h)$  fixes f(s), where  $h := h_1^{-1} h_0$ . Indeed, for every  $g \in G(M)$  identity (2.45) holds for  $q \in U_g$ , where  $U_g$  is the connected component of  $g^{-1}(U) \cap U$  containing O(p). Since  $h \in I_s$ , we have  $s \in U_h$  and the application of (2.45) to h and s yields that  $\varphi(h)$  fixes f(s), as required. Thus, f extends to  $U' := \bigcup_{q \in U} O(q)$ . The extended map satisfies (2.45) for all  $g \in G(M)$  and  $q \in U'$ .

(III). Consider a maximal G(M)-invariant domain  $D \subset M$  from which there exists a biholomorphic map f onto an  $R_{\mathfrak{m}}$ -invariant domain in  $M_{\mathfrak{m}}$  satisfying (2.45) for all  $q \in G(M)$  and  $q \in D$ . The existence of such a domain is guaranteed by the previous step. Assume that  $D \neq M$  and consider  $x \in \partial D$ . Let  $\mathfrak{m}_1$  be the model for O(x) and let  $f_1: O(x) \to \mathfrak{m}_1$  be a real-analytic CR-isomorphism satisfying (2.45) for all  $g \in G(M), q \in O(x)$  and some isomorphism  $\varphi_1: G(M) \to R_{\mathfrak{m}_1}$  in place of  $\varphi$ . As in (II), extend  $f_1$  to a biholomorphic map from a connected G(M)-invariant neighborhood V of O(x)onto an  $R_{\mathfrak{m}_1}$ -invariant neighborhood of  $\mathfrak{m}_1$  in  $M_{\mathfrak{m}_1}$ . The extended map satisfies (2.45) for all  $g \in G(M)$ ,  $q \in V$  and  $\varphi_1$  in place of  $\varphi$ . Consider  $s \in V \cap D$ . The maps f and  $f_1$  take O(s) onto an  $R_{\mathfrak{m}}$ -orbit  $\mathfrak{m}'$  in  $M_{\mathfrak{m}}$  and an  $R_{\mathfrak{m}_1}$ -orbit  $\mathfrak{m}'_1$  in  $M_{\mathfrak{m}_1}$ , respectively. Then  $F := f_1 \circ f^{-1}$  establishes a *CR*-isomorphism between  $\mathfrak{m}'$  and  $\mathfrak{m}'_1$ . Therefore,  $\mathfrak{m}_1$  lies in  $M_{\mathfrak{m}}$ , that is, we have  $M_{\mathfrak{m}} = M_{\mathfrak{m}_1}$ . Moreover, F is either an element of  $\operatorname{Aut}_{CR}(\mathfrak{m}')$  (if  $\mathfrak{m}' = \mathfrak{m}'_1$ ), or is a composition of an element of  $\operatorname{Aut}_{CR}(\mathfrak{m}')$  and a non-trivial map from list (2.44) that takes  $\mathfrak{m}'$  onto  $\mathfrak{m}'_1$  (if  $\mathfrak{m}' \neq \mathfrak{m}'_1$ ). Therefore, F extends to a holomorphic automorphism of  $M_{\mathfrak{m}}$ .

(IV). Since O(x) is strongly pseudoconvex and closed in M, for V small enough we have  $V = V_1 \cup V_2 \cup O(x)$ , where  $V_j$  are open connected nonintersecting sets. For each  $j, D \cap V_j$  is a union of G(M)-orbits and therefore is mapped by f onto a union W of  $R_{\mathfrak{m}}$ -orbits in  $M_{\mathfrak{m}}$ . If W is disconnected, so is D which contradicts the definition of D. Thus,  $D \cap V_j$  is connected for j = 1, 2, and, if V is sufficiently small, then each  $V_j$  is either a subset of D or disjoint from it. Suppose first that there is only one j for which  $D \cap V_j \neq \emptyset$ . In this case  $D \cap V$  is connected and  $V \setminus (D \cup O(x)) \neq \emptyset$ . Setting now

$$\tilde{f} := \begin{cases} f & \text{on } D, \\ F^{-1} \circ f_1 & \text{on } V, \end{cases}$$

$$(2.46)$$

we obtain a biholomorphic extension of f to  $D \cup V$ . By construction,  $\hat{f}$  satisfies (2.45) for  $g \in G(M)$  and  $q \in D \cup V$ . Since  $D \cup V$  is strictly larger than D, we obtain a contradiction with the maximality of D. Thus, in this case D = M, and hence M is holomorphically equivalent to an  $R_{\mathfrak{m}}$ -invariant domain in  $M_{\mathfrak{m}}$ .

Suppose now that  $V_j \,\subset D$  for j = 1, 2. Applying formula (2.46) we can extend  $f|_{V_1}$  and  $f|_{V_2}$  to biholomorphic maps  $\hat{f}_1$ ,  $\hat{f}_2$ , respectively, from a neighborhood of O(x) into  $M_{\mathfrak{m}}$ ; each of these maps satisfies (2.45). Let  $\hat{\mathfrak{m}}_j := \hat{f}_j(O(x)), j = 1, 2$ . Clearly,  $\hat{\mathfrak{m}}_1 \neq \hat{\mathfrak{m}}_2$ . The map  $\hat{F} := \hat{f}_2 \circ \hat{f}_1^{-1}$  is a CR-isomorphism from  $\hat{\mathfrak{m}}_1$  onto  $\hat{\mathfrak{m}}_2$ . In this case  $M_{\mathfrak{m}}$  is one of the manifolds occurring in (b)–(k), and  $\hat{F}$  is a composition of an element of  $\operatorname{Aut}_{CR}(\hat{\mathfrak{m}}_1)$  and a non-trivial map of the corresponding form on list (2.44) that takes  $\hat{\mathfrak{m}}_1$  onto  $\hat{\mathfrak{m}}_2$ . Let  $\Gamma$  denote the cyclic group of automorphisms of  $M_{\mathfrak{m}}$  generated by  $\hat{F}$ . It follows from the explicit form of  $\hat{F}$  and  $M_{\mathfrak{m}}$  in each of the cases (b)–(k) that  $\Gamma$  acts freely properly discontinuously on  $M_{\mathfrak{m}}$ . Let  $M_{\Gamma}$  be the complex manifold obtained by factoring  $M_{\mathfrak{m}}$  by the action of  $\Gamma$  and let  $\Pi_{\Gamma} : M_{\mathfrak{m}} \to M_{\Gamma}$ be the factorization map. Then the map  $\Pi_{\Gamma} \circ \mathcal{F}$ , where

$$\mathcal{F} := \begin{cases} f & \text{on } D, \\ \hat{f}_1 & \text{on } O(x), \end{cases}$$
(2.47)

establishes holomorphic equivalence between M and  $M_{\Gamma}$ . Observe, however, that in cases (b)–(f), (h)–(k) the manifold  $M_{\Gamma}$  is not hyperbolic, since it is covered by  $M_{\mathfrak{m}}$  and none of the manifolds  $M_{\mathfrak{m}}$  in these cases is hyperbolic. Further, it is straightforward to see that in the remaining case (g) we have  $d(M_{\Gamma}) \neq 3$  (in fact, in this case either  $d(M_{\Gamma}) = 2$  or  $d(M_{\Gamma}) = 4$ ). These contradictions show that in all cases (b)–(k) exactly one of  $V_j$ , j = 1, 2, is a subset of D, and hence M is holomorphically equivalent to an  $R_{\mathfrak{m}}$ -invariant domain in  $M_{\mathfrak{m}}$ .

(V). In each of cases (b)-(n") the determination of  $R_{\mathfrak{m}}$ -invariant domains is straightforward; they are as follows:

(b)  $\{(z,w) \in \mathbb{C}^2 : s < |z|^2 + |w|^2 < t\} / \mathbb{Z}_m, \ 0 \le s < t < \infty, \ m \ge 2;$ (c)  $\{(z,w) \in \mathbb{C}^2 : s + |z|^2 < \operatorname{Re} w < t + |z|^2\}, \ -\infty < s < t < \infty;$ 

- (d)  $\mathfrak{R}_{b,s,t} := \left\{ (z,w) \in \mathbb{C}^2 : s|w|^b < \operatorname{Re} z < t|w|^b, \ w \neq 0 \right\}, \text{ for } 0 \le s < t \le \infty, \text{ with either } s > 0 \text{ or } t < \infty;$
- (e)  $\mathfrak{R}_{s,t} := \{(z,w) \in \mathbb{C}^2 : s \exp(\operatorname{Re} w) < \operatorname{Re} z < t \exp(\operatorname{Re} w)\}, \text{ for } 0 \le s < t \le \infty, \text{ with either } s > 0 \text{ or } t < \infty;$
- (f)  $\{(z,w) \in \mathbb{C}^2 : s \exp(|z|^2) < |w| < t \exp(|z|^2)\}, 0 < s < t \le \infty;$

(g) 
$$R_{b,s,t}$$
, with  $0 \le s < t \le \infty$ ;

- (h)  $U_{s,t}$ , for  $0 \le s < t \le \infty$ , with either s > 0 or  $t < \infty$ ;
- (j)  $\mathfrak{S}_{s,t}, 0 \leq s < t < \infty;$
- (j')  $\mathfrak{S}_{s,t}^{(\infty)}, 0 \le s < t < \infty;$
- (j")  $\mathfrak{S}_{s,t}^{(n)}, 0 \le s < t < \infty, n \ge 2;$
- (k)  $V_{b,s,t}$ ,  $0 < t < \infty$ ,  $e^{-2\pi b}t < s < t$ ;
- (l)  $E_{s,t}, 1 \le s < t < \infty;$
- (l')  $E_{s,t}^{(4)}, 1 \le s < t < \infty;$
- (l")  $E_{s,t}^{(2)}, 1 \le s < t < \infty;$
- (m)  $\Omega_{s,t}, -1 \le s < t \le 1;$
- (m')  $\Omega_{s,t}^{(\infty)}, -1 \le s < t \le 1;$
- (m")  $\Omega_{s,t}^{(n)}, -1 \le s < t \le 1, n \ge 2;$
- (n)  $D_{s,t}, 1 \leq s < t \leq \infty;$
- (n')  $D_{s,t}^{(\infty)}, 1 \le s < t \le \infty;$
- (n")  $D_{s,t}^{(2)}, 1 \le s < t \le \infty;$

A. V. Isaev

(n") 
$$D_{s,t}^{(2n)}, 1 \le s < t \le \infty, n \ge 2;$$

(n")  $D_{s,t}^{(n)}$ ,  $1 \leq s < t \leq \infty$ ,  $n \geq 3$ , n is odd.

This concludes our gluing procedure. Note that in each of cases (d) and (e) we have two non-isomorphic possibilities for  $R_{\mathfrak{m}}$ . Each of the possibilities leads to the same set of  $R_{\mathfrak{m}}$ -invariant domains.

To finish the proof of the theorem it now remains to observe that the automorphism groups of all  $R_{\mathfrak{m}}$ -invariant domains that appear in cases (b)–(f) have dimension at least 4 rather than 3. The remaining domains form list (i)–(xv) as stated in the theorem.

The proof of the theorem is complete.

## 3 Levi-Flat orbits

In this section we deal with Levi-flat orbits. First, we will classify such orbits. In order to formulate our orbit classification result, we introduce the following groups of transformations.

For every  $b \in \mathbb{R}$  let  $G_b$  be the group of all maps of the form (2.1),  $G'_0$  be the group of all maps of the form

$$\begin{array}{rccc} z & \mapsto & \lambda z + i\beta, \\ w & \mapsto & e^{i\varphi}w, \end{array} \tag{3.1}$$

where  $\lambda > 0$ ,  $\beta, \varphi \in \mathbb{R}$ , and G the group of all maps of the form (2.3).

We will now prove the following proposition. Observe that the proposition applies to an arbitrary (2,3)-manifold (possibly containing codimension 2 orbits).

**Proposition 3.1** Let M be a (2,3)-manifold. Assume that for a point  $p \in M$  its orbit O(p) is Levi-flat. Then O(p) is equivalent to one of the following hypersurfaces:

$$O_1 := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0, \operatorname{Re} w = 0\}, O_2 := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0, |w| = 1\},\$$

by means of a real-analytic CR-map. The CR-equivalence can be chosen so that it transforms G(M) into either the group  $G_b$  for some  $b \in \mathbb{R}$  or the group G in the first case, and into the group  $G'_0$  in the second case. **Proof:** Recall that the hypersurface O(p) is foliated by complex manifolds holomorphically equivalent to  $\mathcal{P}$  (see (ii) of Proposition 1.1). Denote by  $\mathfrak{g}$ the Lie algebra of vector fields on O(p) arising from the action of G(M). The algebra  $\mathfrak{g}$  is isomorphic to the Lie algebra of G(M). For  $q \in O(p)$  we consider the leaf  $M_q$  of the foliation passing through q and the subspace  $\mathfrak{l}_q \subset \mathfrak{g}$  of all vector fields tangent to  $M_q$  at q. Since vector fields in  $\mathfrak{l}_q$  remain tangent to  $M_q$  at each point in  $M_q$ , the subspace  $\mathfrak{l}_q$  is in fact a Lie subalgebra of  $\mathfrak{g}$ . It follows from the definition of  $\mathfrak{l}_q$  that dim  $\mathfrak{l}_q = 2$ .

Denote by  $H_q$  the (possibly non-closed) connected subgroup of G(M)with Lie algebra  $l_q$ . It is straightforward to verify that the group  $H_q$  acts on  $M_q$  by holomorphic transformations. If some element  $g \in H_q$  acts trivially on  $M_q$ , then  $g \in I_q$ . If for every non-identical element of  $L_q$  its projection to  $L'_{q}$  is non-identical (see (ii) of Proposition 1.1), then every non-identical element of  $I_q$  acts non-trivially on  $M_q$  and thus g = id; if  $L_q$  contains a non-identical element with an identical projection to  $L'_{q}$  and  $g \neq id$ , then  $g = g_q$ , where  $g_q$  denotes the element of  $I_q$  corresponding to the non-trivial element in  $\mathbb{Z}_2$  (see (ii) of Proposition 1.1). Thus, either  $H_q$  or  $H_q/\mathbb{Z}_2$  acts effectively on  $M_q$  (the former case occurs if  $g_q \notin H_q$ , the latter if  $g_q \in H_q$ ). Since dim  $H_q = 2$ , we obtain that either  $H_q$  or  $H_q/\mathbb{Z}_2$  is isomorphic to a 2-dimensional (possibly non-closed) subgroup of  $\operatorname{Aut}(\mathcal{P})$ . As we noted in the proof of Lemma 2.2, every codimension 1 subgroup of  $Aut(\mathcal{P})$  is conjugate in  $\operatorname{Aut}(\mathcal{P})$  to the subgroup  $\mathcal{T}$  (see (2.43)). The Lie algebra of this subgroup is isomorphic to the 2-dimensional Lie algebra  $\mathfrak{h}$  given by two generators X and Y satisfying [X, Y] = X. Therefore,  $\mathfrak{l}_q$  is isomorphic to  $\mathfrak{h}$  for every  $q \in O(p)$ .

It is not difficult to determine all 3-dimensional Lie algebras containing a subalgebra isomorphic to  $\mathfrak{h}$ . Every such algebra has generators X, Y, Z that satisfy one of the following sets of relations:

Suppose first that  $\mathfrak{g}$  is given by relations (R1). In this case  $\mathfrak{g}$  is isomorphic to the Lie algebra of the simply-connected Lie group  $G_b$ . Indeed, the Lie algebra of  $G_b$  is isomorphic to the Lie algebra of vector fields on  $\mathbb{C}^2$  with the generators

$$X_1 := i \partial/\partial z,$$
  

$$Y_1 := z \partial/\partial z + bw \partial/\partial w,$$
  

$$Z_1 := i \partial/\partial w,$$

that clearly satisfy (R1).

Assume first that  $b \neq 0$ . In this case the center of  $G_b$  is trivial, and hence  $G_b$  is the only (up to isomorphism) connected Lie group whose Lie algebra is given by relations (R1). Therefore, G(M) is isomorphic to  $G_b$ . Assume further that  $b \neq 1$ . In this case, it is straightforward to observe that every subalgebra of  $\mathfrak{g}$  isomorphic to  $\mathfrak{h}$  is generated by either X and  $Y + \nu Z$ , or by Z and  $\nu X + Y$  for some  $\nu \in \mathbb{R}$ . The connected subgroup of  $G_b$  with Lie algebra generated by  $X_1$  and  $Y_1 + \nu Z_1$  is conjugate in  $G_b$  to the closed subgroup  $G_b^1$  given by  $\gamma = 0$ ; similarly, the connected subgroup of with Lie algebra generated by  $Z_1$  and  $\nu X_1 + Y_1$  is conjugate to the closed subgroup  $G_b^2$  given by  $\beta = 0$  (see (2.1)). Moreover, the conjugating element can be chosen to have the form

$$\begin{array}{lll} z & \mapsto & z \\ w & \mapsto & w + i\gamma, \quad \gamma \in \mathbb{R}, \end{array} \tag{3.3}$$

in the first case, and the form

$$\begin{array}{rcl}
z & \mapsto & z + i\beta, & \gamma \in \mathbb{R}, \\
w & \mapsto & w,
\end{array}$$
(3.4)

in the second case.

Thus we have shown that (upon identification of G(M) and  $G_b$ ), the subgroup  $H_q$  for every  $q \in O(p)$  is conjugate to either  $G_b^1$  or to  $G_b^2$  by a map of either the form (3.3) or the form (3.4), respectively. In particular,  $H_q$  is isomorphic to  $\mathcal{T}$  and hence does not have subgroups isomorphic to  $\mathbb{Z}_2$ . Therefore,  $H_q$  acts effectively on  $M_q$ . Since the subgroups  $H_q$  are conjugate to each other, it follows that either  $H_q$  is conjugate to  $G_b^1$  for every q, or  $H_q$ is conjugate to  $G_b^2$  for every q. Suppose, for example, that the former holds. Since the normalizer of  $G_b^1$  in  $G_b$  coincides with  $G_b^1$ , for every  $q_1, q_2 \in O(p)$ there exists a unique  $g \in G(M)$  of the form (3.3) such that  $gM_{q_1} = M_{q_2}$ . Fix  $q_0 \in O(p)$  and let  $f : M_{q_0} \to \mathcal{P}$  be a holomorphic equivalence that transforms  $H_{q_0}|_{M_{q_0}}$  into the group  $\mathcal{T}$  and such that  $\hat{X}_1|_{M_{q_0}}$  and  $\hat{Y}_1|_{M_{q_0}}$  are transformed into the vector fields  $i\partial/\partial z|_{\mathcal{P}}$  and  $z\partial/\partial z|_{\mathcal{P}}$ , respectively, where  $X_1, Y_1$  are the vector fields on O(p) corresponding to  $X_1, Y_1$ . For every  $q \in O(p)$  find the unique map g of the form (3.3) such that  $gM_{q_0} = M_q$ and define  $F(q) := (f(g^{-1}(q)), i\gamma) \in \mathbb{C}^2$ , with  $\gamma$  corresponding to g as in formula (3.3). Clearly, F is a real-analytic CR-isomorphism between O(p)and  $O_1$  that transforms G(M) into  $G_b$ . The case when  $H_q$  is conjugate to  $G_b^2$  for every  $q \in O(p)$  is treated similarly; arguing as above we construct a real-analytic *CR*-isomorphism between O(p) and

$$O'_{1} := \{(z, w) \in \mathbb{C}^{2} : \operatorname{Re} z = 0, \operatorname{Re} w > 0\}, \qquad (3.5)$$

that transforms G(M) into  $G_b$ . Further, interchanging the variables turns  $O'_1$  into  $O_1$  and  $G_b$  into  $G_{1/b}$ .

Suppose now that b = 1. In this case, in addition to the subalgebras analogous to those arising for  $b \neq 1$ , a subalgebra of  $\mathfrak{g}$  isomorphic to  $\mathfrak{h}$  can also be generated by  $X + \eta Z$  and  $Y + \nu Z$  for some  $\eta, \nu \in \mathbb{R}, \eta \neq 0$ . The connected subgroup of  $G_1$  corresponding to this subalgebra is conjugate in  $G_1$  to the subgroup  $G_{1,\eta}$  of all maps of the form (2.1) with b = 1,  $\gamma = \beta \eta$ . Moreover, the conjugating element can be chosen to have the form (3.3). Thus, upon identification of G(M) and  $G_1$ , the subgroup  $H_q$  for every  $q \in$ O(p) is conjugate to either  $G_1^1$  or to  $G_1^2$ , or to  $G_{1,\eta}$  for some  $\eta \neq 0$  (all these subgroups are closed). In particular,  $H_q$  is isomorphic to  $\mathcal{T}$  and hence acts effectively on  $M_q$ . Since the subgroups  $H_q$  are conjugate to each other, it follows that either  $H_q$  is conjugate to  $G_1^1$  for every q, or  $H_q$  is conjugate to  $G_1^2$  for every q, or  $H_q$  is conjugate to  $G_{1,\eta}$  for every q and a fixed  $\eta$ . The first two cases are treated as for  $b \neq 1$ . Suppose that  $H_q$  is conjugate to  $G_{1,\eta}$  for every  $q \in O(p)$ . Since the normalizer of  $G_{1,\eta}$  in  $G_1$  coincides with  $G_{1,\eta}$ , for every  $q_1, q_2 \in O(p)$  there exists a unique  $g \in G(M)$  of the form (3.3) such that  $gM_{q_1} = M_{q_2}$ . Fix  $q_0 \in O(p)$  and let  $f : M_{q_0} \to N_{\eta}$ , with  $N_{\eta} := \{(z, w) \in \mathbb{C}^2 : w = \eta z, \operatorname{Re} z > 0\}, \text{ be a holomorphic equivalence that}$ transforms  $H_{q_0}|_{M_{q_0}}$  into the group  $G_{1,\eta}|_{N_{\eta}}$  and such that  $X_1|_{M_{q_0}}$  and  $Y_1|_{M_{q_0}}$  are transformed into the vector fields  $i\partial/\partial z + i\eta\partial/\partial w|_{N_{\eta}}$  and  $z\partial/\partial z + w\partial/\partial w|_{N_{\eta}}$ , respectively. For every  $q \in O(p)$  find the unique map g of the form (3.3) such that  $gM_{q_0} = M_q$  and define  $F(q) := (f(g^{-1}(q)), i\gamma)$ , with  $\gamma$  corresponding to q as in formula (3.3). Clearly, F is a real-analytic CR-isomorphism between O(p) and

$$O_{1,\eta} := \left\{ (z, w) \in \mathbb{C}^2 : \operatorname{Re} w = \eta \operatorname{Re} z, \operatorname{Re} z > 0 \right\},\$$

that transforms G(M) into  $G_1$ . Finally, the transformation

$$\begin{array}{rccc} z & \mapsto & z, \\ w & \mapsto & w - \eta z \end{array}$$

turns  $O_{1,\eta}$  into  $O_1$  and preserves the group  $G_1$ .

Suppose now that b = 0. In this case there are exactly two (up to isomorphism) connected Lie groups with Lie algebra  $\mathfrak{g}$ :  $G_0$  and  $G'_0$ . It is straightforward to see that every subalgebra of  $\mathfrak{g}$  isomorphic to  $\mathfrak{h}$  is generated by X and  $Y + \nu Z$  for some  $\nu \in \mathbb{R}$ . Clearly, the connected subgroup of  $G_0$  with Lie algebra generated by  $X_1$  and  $Y_1 + \nu Z_1$  coincides with the closed normal subgroup  $G_{0,\nu}$  given by  $\lambda = e^t$ ,  $\gamma = \nu t$ ,  $t \in \mathbb{R}$  (see (2.1)). It then follows that if G(M) is isomorphic to  $G_0$ , there exists  $\nu \in \mathbb{R}$ , such that, identifying G(M) and  $G_0$ , we have  $H_q = G_{0,\nu}$  for every  $q \in O(p)$ . Further, let us realize the Lie algebra of  $G'_0$  as the Lie algebra generated by the following vector fields on  $\mathbb{C}^2$ :

$$\begin{array}{rcl} X_1' &:=& i \,\partial/\partial z, \\ Y_1' &:=& z \,\partial/\partial z, \\ Z_1' &:=& i w \,\partial/\partial w, \end{array}$$

which clearly satisfy (R1) of (3.2). Therefore, the connected subgroup of  $G'_0$ with Lie algebra generated by  $X'_1$  and  $Y'_1 + \nu Z'_1$  coincides with the closed normal subgroup  $G'_{0,\nu}$  of  $G'_0$  given by  $\lambda = e^t$ ,  $\varphi = \nu t$ ,  $t \in \mathbb{R}$  (see (3.1)). It then follows that if G(M) is isomorphic to  $G'_0$ , there exists  $\nu \in \mathbb{R}$ , such that, identifying G(M) and  $G'_0$ , we have  $H_q = G'_{0,\nu}$  for every  $q \in O(p)$ .

Thus, if b = 0, every subgroup  $H_q$  is normal, closed, isomorphic to  $\mathcal{T}$ (hence acts effectively on  $M_q$ ). In particular, all these subgroups coincide for  $q \in O(p)$ . Denote by H the coinciding subgroups  $H_q$ . The group H acts properly on O(p), and the orbits of this action are the leaves  $M_q$  of the foliation on O(p). Further, we have  $G(M) = H \times L$ , where L is either the subgroup given by  $\lambda = 1, \beta = 0$  in formula (2.1) where we set b = 0, or to the subgroup given by  $\lambda = 1, \beta = 0$  in formula (3.1), and hence isomorphic to either  $\mathbb{R}$  or  $S^1$ . For every  $q \in O(p)$  let  $S_q := \{g \in L : gM_q = M_q\}$ . Since  $M_q$  is closed,  $S_q$  is a closed subgroup of L. Clearly, for every  $g \in S_q$  there is  $h \in H$  such that  $hg \in I_q$ . The elements g and h lie in the projections of  $I_q$  to L and H, respectively. Since H is isomorphic to  $\mathcal{T}$ , it does not have non-trivial finite subgroups, hence the projection of  $I_q$  to H is trivial, and therefore  $S_q = I_q$ . Since all isotropy subgroups are contained in the Abelian subgroup L and are conjugate to each other in G(M), they are in fact identical. The effectiveness of the action of G(M) on M now implies that all isotropy subgroups are trivial and hence every  $S_q$  is trivial as well.

Thus, we have shown that for every  $q_1, q_2 \in O(p)$  there is a unique  $g \in L$ , such that  $gM_{q_1} = M_{q_2}$ . Fix  $q_0 \in O(p)$  and let  $f : M_{q_0} \to \mathcal{P}$  be a holomorphic equivalence that transforms  $H_{q_0}|_{M_{q_0}}$  into the group  $\mathcal{T}$ . For every  $q \in O(p)$  find the unique map  $g \in L$  such that  $gM_{q_0} = M_q$  and define

 $F(q) := (f(g^{-1}(q)), u)$ , where u corresponds to g as follows (see formula (2.1) with b = 0 and formula (3.1)):  $u = i\gamma$  if  $L \simeq \mathbb{R}$  and  $u = e^{i\varphi}$  if  $L \simeq S^1$ . Clearly, if  $G(M) \simeq G_0$ , the map F takes O(p) onto  $O_1$  and transforms G(M) into  $G_0$ , and if  $G(M) \simeq G'_0$ , it takes O(p) onto  $O_2$  and transforms G(M) into  $G'_0$ .

Suppose next that  $\mathfrak{g}$  is given by relations (R2) (see (3.2)). In this case  $\mathfrak{g}$  is isomorphic to the Lie algebra of the simply-connected Lie group G. Indeed, the Lie algebra of G is isomorphic to the Lie algebra of holomorphic vector fields on  $\mathbb{C}^2$  with the following generators:

$$\begin{aligned} X_2 &:= i \partial/\partial z, \\ Y_2 &:= (z+w) \partial/\partial z + w \partial/\partial w, \\ Z_2 &:= i \partial/\partial w, \end{aligned}$$

that clearly satisfy (R2). It is straightforward to observe that the center of G is trivial, and hence G is the only (up to isomorphism) connected Lie group whose Lie algebra is given by relations (R2). Therefore, G(M) is isomorphic to G. In this case every subalgebra of  $\mathfrak{g}$  isomorphic to  $\mathfrak{h}$  is generated by Xand  $Y + \nu Z$  for some  $\nu \in \mathbb{R}$ . The connected subgroup of G with Lie algebra generated by  $X_2$  and  $Y_2 + \nu Z_2$  is conjugate in G to the closed subgroup Qgiven by  $\gamma = 0$  (see (2.3)). Moreover, the conjugating element can be chosen to have the form (3.3).

Thus – upon identification of G(M) and G – the subgroup  $H_q$  for every  $q \in O(p)$  is conjugate to Q by a map of the form (3.3). Further, since the normalizer of Q in G coincides with Q, we proceed as in the case of relations (R1) with  $b \neq 0, 1$  and obtain that there exists a real-analytic CR-isomorphism between O(p) and  $O_1$  that transforms G(M) into G.

Suppose finally that  $\mathfrak{g}$  is given by relations (R3) (see (3.2)). We will show that this is in fact impossible. In this case  $\mathfrak{g}$  is isomorphic to the algebra  $\mathfrak{su}_{1,1}$ . All connected Lie groups with Lie algebra isomorphic to  $\mathfrak{su}_{1,1}$ are described as follows: the simply-connected group (the universal cover of  $SU_{1,1}$ ) is the group  $V_{\infty}$ , and any non simply-connected group is isomorphic to  $V_n$  with  $n \in \mathbb{N}$ , where  $V_{\infty}$  and  $V_n$  are the Lie groups defined in Lemma 2.2. Clearly, the set  $\mathcal{C} := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0\}$  is  $V_{\infty}$ -invariant, and the set  $\mathcal{C}' := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0, w \neq 0\}$  is  $V_n$ -invariant for  $n \in \mathbb{N}$ . Consider in  $V_j$  three one-parameter subgroups  $H_X$ ,  $H_Y$ ,  $H_Z$ 

for  $j = \infty$ :  $z \mapsto z - \frac{i}{2}\beta, \quad w \mapsto w,$   $z \mapsto \lambda z, \qquad w \mapsto w + \ln_0 \lambda,$  $z \mapsto \frac{z}{i\mu z + 1}, \quad w \mapsto w - 2\ln_0(i\mu z + 1),$ 

for  $j = n \in \mathbb{N}$ :

where  $\lambda > 0$ ,  $\beta, \mu \in \mathbb{R}$ ,  $t^{2/n} = \exp(2/n \ln_0 t)$  for  $t \in \mathbb{C} \setminus (-\infty, 0]$ , and  $\ln_0$  is the branch of the logarithm in  $\mathbb{C} \setminus (-\infty, 0]$  defined by the condition  $\ln_0 1 = 0$ . The vector fields on  $\mathbb{C}^2$  corresponding to these subgroups generate the Lie algebras of  $V_j$  for  $j \in \{1, 2, ..., \infty\}$  and are as follows:

for 
$$j = \infty$$
:  
 $X_3 := -\frac{i}{2} \partial/\partial z$ ,  
 $Y_3 := z \partial/\partial z + \partial/\partial w$ ,  
 $Z_3 := -iz^2 \partial/\partial z - 2iz \partial/\partial w$ ,

for  $j = n \in \mathbb{N}$ :  $X_{3} := -\frac{i}{2} \partial/\partial z,$   $Y_{3} := z \partial/\partial z + \frac{1}{n} \partial/\partial w,$   $Z_{3} := -iz^{2} \partial/\partial z - \frac{2izw}{n} \partial/\partial w.$ (3.6)

One can verify that these vector fields indeed satisfy relations (R3).

Next, it is straightforward to observe that any subalgebra of  $\mathfrak{g}$  isomorphic to  $\mathfrak{h}$  is generated by either  $X + \eta Y - \eta^2/2Z$ ,  $Y - \eta Z$ , with  $\eta \in \mathbb{R}$ , or by Y, Z.

For every  $j \in \{1, 2, ..., \infty\}$  the connected subgroups of  $V_j$  corresponding to the subalgebras generated by  $X_3 + \eta Y_3 - \eta^2/2 Z_3$ ,  $Y_3 - \eta Z_3$ , with  $\eta \in \mathbb{R}$ , or by  $Y_3$ ,  $Z_3$  are isomorphic to  $\mathcal{T}$  (hence  $H_q$  acts effectively on  $M_q$  for every q) and are all conjugate to each other in  $V_j$  by elements of the subgroup  $W_j$  that we will describe below. Let first  $j = \infty$ . Transform  $\mathcal{C}$  into  $\{(z, w) : |z| < 1\}$  by means of the map  $\Phi_\infty$  defined as

$$z \mapsto \frac{z-1}{z+1},$$

$$w \mapsto w-2\ln_0\left((z+1)/2\right),$$
(3.7)

and let  $W_{\infty}$  be the subgroup of  $V_{\infty}$  that transforms under  $\Phi_{\infty}$  into the group of maps

$$\begin{array}{rccc} z & \mapsto & e^{it}z, \\ w & \mapsto & w+it \end{array}$$

where  $t \in \mathbb{R}$ . Let now j = n. Transform  $\mathcal{C}'$  into  $\{(z, w) : |z| < 1, w \neq 0\}$  by means of the map  $\Phi_n$  defined as

$$z \mapsto \frac{z-1}{z+1},$$

$$w \mapsto \left(\frac{2}{z+1}\right)^{2/n} w,$$
(3.8)

and let  $W_n$  be the subgroup of  $V_n$  that transforms under  $\Phi_n$  into the group of maps

$$\begin{array}{rccc} z & \mapsto & e^{it}z, \\ w & \mapsto & e^{it/n}w, \end{array}$$

where  $0 \leq t < 2\pi n$ .

Observe that – upon identification of G(M) and  $V_j$  for a particular value of j – for every  $q \in O(p)$  every element  $g \in G(M)$  can be written in the form g = g'g'' with  $g' \in W_j$ ,  $g'' \in H_q$ . In particular, for every  $q \in O(p)$  there exists a neighborhood of  $M_q$  in O(p) that consists of leaves  $M_{q'}$  such that the groups  $H_{q'}$  corresponding to the leaves are pairwise distinct. Further, for every two points  $q_1, q_2 \in O(p)$  there exists  $g \in W_j$  such that  $gM_{q_1} = M_{q_2}$ . If for some  $q \in O(p)$  there is a non-trivial  $g \in W_j$  such that  $gM_q = M_q$ , then  $gH_qg^{-1} = H_q$  and hence  $\Phi_j \circ g \circ \Phi_j^{-1}$  has the form

for 
$$j = \infty$$
:  
 $z \mapsto z,$   
 $w \mapsto w + 2\pi i k_0, \quad k_0 \in \mathbb{Z} \setminus \{0\},$ 

for 
$$j = n \in \mathbb{N}$$
:  
 $z \mapsto z,$   
 $w \mapsto e^{2\pi i k_0/n} w, \quad k_0 \in \mathbb{N}, \, k_0 \le n-1$ 

Find  $h \in H_q$  such that  $gh \in I_q$ . However, every element of  $I_q$  has finite order (see (ii) of Proposition 1.1). If  $G(M) = V_\infty$ , then gh is clearly of infinite order. Assume that  $G(M) = V_n$  for some  $n \in \mathbb{N}$ . In this case ghcan only be of finite order if h = id. Hence  $g \in I_q$  and, since this argument can be applied to any point in  $M_q$ , we obtain that g fixes every point in  $M_q$ . Hence  $g = g_q$ , where, as before,  $g_q$  denotes the element of  $I_q$  corresponding to the non-trivial element in  $\mathbb{Z}_2$  (see (ii) of Proposition 1.1). Then for a pair of points  $q_1, q_2$  sufficiently close to q (and hence such that  $H_{q_1} \neq H_{q_2}$ ) we obtain that  $H_{q_1} = g_q H_{q_2} g_q^{-1} = H_{q_2}$  which is impossible. The above contradictions show that in fact  $gM_q \neq M_q$  for every  $q \in O(p)$  and every non-trivial  $g \in W_j$ .

Suppose that  $G(M) = V_{\infty}$ . Fix  $q_0 \in O(p)$  and let  $f : M_{q_0} \to \mathcal{P}$  be a holomorphic equivalence that transforms  $H_{q_0}|_{M_{q_0}}$  into the group  $\mathcal{T}$  and such that  $\hat{X}_3|_{M_{q_0}}$  and  $\hat{Y}_3|_{M_{q_0}}$  are transformed into the vector fields  $i\partial/\partial z|_{\mathcal{P}}$  and  $z\partial/\partial z|_{\mathcal{P}}$ , respectively, where  $\hat{X}_3$ ,  $\hat{Y}_3$  are the vector fields on O(p) corresponding to  $X_3$ ,  $Y_3$ . For every  $q \in O(p)$  find the unique map  $g \in W_{\infty}$  such that  $gM_{q_0} = M_q$  and define  $F_{\infty}(q) := \left(f(g^{-1}(q)), it\right) \in \mathbb{C}^2$ , where t is the parameter value corresponding to g. Clearly,  $F_{\infty}$  is a real-analytic CR-isomorphism between O(p) and  $O_1$  that transforms  $\hat{X}_3 - 1/2 \hat{Z}_3$  into  $i\partial/\partial w|_{O_1}$ , where  $\hat{Z}_3$ is the vector field on O(p) corresponding to  $Z_3$ .

Denote by X, Y, Z the vector fields on  $O_1$  into which  $F_{\infty}$  transforms  $X_3$ ,  $\hat{Y}_3, \hat{Z}_3$ , respectively. Since  $F_{\infty}$  is real-analytic, it extends to a biholomorphic map from a neighborhood of O(p) in M onto a neighborhood of  $O_1$  in  $\mathbb{C}^2$ . Clearly,  $\hat{X}_3, \hat{Y}_3, \hat{Z}_3$  extend to holomorphic vector fields on all of M and hence  $\tilde{X}, \tilde{Y}, \tilde{Z}$  extend to holomorphic vector fields defined in a neighborhood of  $O_1$ . Since  $\tilde{X}$  and  $\tilde{Y}$  are tangent to  $O_1$  and since their restrictions to  $\mathcal{P} \times \{0\} \subset O_1$ 

36

are  $i\partial/\partial z$  and  $z\partial/\partial z$ , respectively, they have the form

$$\tilde{X} = (i + \rho(z, w))\partial/\partial z + \sigma(z, w)\partial/\partial w, 
\tilde{Y} = (z + \mu(z, w))\partial/\partial z + \nu(z, w)\partial/\partial w,$$
(3.9)

where  $\rho,\sigma,\mu,\nu$  are functions holomorphic near  $O_1$  such that

$$\rho(z,0) \equiv \sigma(z,0) \equiv \mu(z,0) \equiv \nu(z,0), \qquad (3.10)$$

and

$$\operatorname{Re}\sigma(z,iv) \equiv \operatorname{Re}\nu(z,iv) \equiv 0. \tag{3.11}$$

Since  $[X_3 - 1/2 Z_3, X_3] = -1/2 Y_3$  and  $[X_3 - 1/2 Z_3, Y_3] = 2X_3 - (X_3 - 1/2 Z_3)$ , we obtain

$$\begin{bmatrix} \mathcal{Z}, \tilde{X} \end{bmatrix} = -\frac{1}{2} \tilde{Y},$$

$$\begin{bmatrix} \mathcal{Z}, \tilde{Y} \end{bmatrix} = 2\tilde{X} - \mathcal{Z},$$
(3.12)

where  $\mathcal{Z} := i\partial/\partial w$ .

Conditions (3.12) together with (3.9) and (3.10) uniquely determine the functions  $\rho, \sigma, \mu, \nu$  as follows:

$$\rho = \frac{i}{4} \Big( (z+2)e^w - (z-2)e^{-w} \Big) - i,$$
  

$$\sigma = -\frac{i}{4} \Big( e^w + e^{-w} - 2 \Big),$$
  

$$\mu = \frac{1}{2} \Big( (z+2)e^w + (z-2)e^{-w} \Big) - z,$$
  

$$\nu = -\frac{1}{2} \Big( e^w - e^{-w} \Big).$$

The functions  $\sigma$ ,  $\nu$  automatically satisfy conditions (3.11).

Integrating  $\tilde{X}, \tilde{Y}, \mathcal{Z}$  we respectively obtain the following one-parameter

subgroups of holomorphic transformations of  $O_1$ :

$$\begin{array}{rcl} z & \mapsto & ze^{-w}\left(e^w + \frac{i}{4}(e^w - 1)t\right)\left(1 + \frac{i}{4}(e^w - 1)t\right) + \\ & & \frac{i}{2}\left[t(e^w + e^{-w}) + t^2\frac{i}{4}(e^w - 1)(e^{-w} + 1)\right], \\ H_{\tilde{X}}: \\ w & \mapsto & \ln^w\left(\frac{e^w + \frac{i}{4}(e^w - 1)t}{1 + \frac{i}{4}(e^w - 1)t}\right), \end{array}$$

$$z \mapsto \frac{1}{4} \Big( z \left[ (2 + e^w + e^{-w}) e^t + (2 - e^w - e^{-w}) e^{-t} \right] + 2 (e^w - e^{-w}) (e^t - e^{-t}) \Big),$$
  

$$H_{\tilde{Y}}:$$

$$w \mapsto \ln^{w} \frac{e^{w} + 1 + (e^{w} - 1)e}{e^{w} + 1 - (e^{w} - 1)e}$$

$$\begin{array}{cccc} z & \mapsto & z, \\ H_{\mathcal{Z}}: & & \\ & w & \mapsto & w + it, \end{array}$$

where  $t \in \mathbb{R}$  and  $\ln^w := \ln'_0 + 2\pi i k$  for  $2\pi k < \operatorname{Im} w < 2\pi (k+1)$ , and  $\ln^{2\pi i k} \equiv 2\pi i k$ ,  $k \in \mathbb{Z}$ , with  $\ln'_0$  being the branch of the logarithm defined in  $\mathbb{C} \setminus [0, \infty)$  by the condition  $\ln'_0(-1) = \pi i$ .

Denote by  $\tilde{V}_{\infty}$  the Lie group of CR-transformations of  $O_1$  into which  $F_{\infty}$  transforms  $G(M)|_{O(p)}$ . Clearly,  $\tilde{V}_{\infty}$  is isomorphic to  $V_{\infty}$  and  $H_{\tilde{X}}$ ,  $H_{\tilde{Y}}$ ,  $H_{\mathcal{Z}}$  are one-parameter subgroups of  $\tilde{V}_{\infty}$ . Observe that the action of  $\tilde{V}_{\infty}$  on  $O_1$  does not extend to an action on any neighborhood of  $O_1$  in  $\mathbb{C}^2$ ; more precisely, for every point  $p_0 = (z_0, w_0) \in \mathbb{C}^2$  with  $\operatorname{Re} w_0 \neq 0$  there exists an element of  $\tilde{V}_{\infty}$  that has a singularity at  $p_0$ .

There exist a neighborhood  $\mathcal{W}$  of O(p) in M such that O(q) is of codimension 1 in M for every  $q \in \mathcal{W}$  (in the presence of a codimension 1 orbit there may be at most two codimension 2 orbits in M – see [1]). Choosing  $\mathcal{W}$  sufficiently small, we can assume that  $F_{\infty}$  extends to  $\mathcal{W}$  to a biholomorphic map onto a neighborhood  $\mathcal{W}'$  of  $O_1$  in  $\mathbb{C}^2$ . Suppose first that for every  $q \in \mathcal{W}$  the orbit O(q) is Levi-flat. Let  $\hat{q} := F_{\infty}^{-1}(1,0)$  and  $q_m := F_{\infty}^{-1}(1,\epsilon_m)$  with  $\epsilon_m \to 0$  as  $m \to \infty$  (we assume that m is sufficiently large to ensure that  $(1,\epsilon_m) \in \mathcal{W}'$ ). Let  $M_{q_m}^{O(q_m)}$  be the leaf of the foliation on  $O(q_m)$  passing through  $q_m$ . It now follows that for every m there exists  $g_m \in W_{\infty}$ , such that  $g_m H_{\hat{q}} g_m^{-1}$  is the connected subgroup of G(M) corresponding to the subalgebra of vector fields on  $O(q_m)$  tangent to  $M_{q_m}^{O(q_m)}$ . Clearly, we can choose the sequence  $\{g_m\}$  so that  $g_m \to \text{id}$  as  $m \to \infty$ . At the same time, conjugating the one-parameter subgroups  $H_{\tilde{X}}$  and  $H_{\tilde{Y}}$  by the element of  $H_{\mathcal{Z}}$  corresponding to the conjugated subgroups, we see that these vector fields at the points  $(1, \epsilon_m)$  span all of  $\mathbb{C}^2$  rather than a complex line in  $\mathbb{C}^2$  (the span coincides with that of the pair of vectors  $(3e^{\epsilon_m + it_m}, 1 - e^{\epsilon_m + it_m})$  and  $(-1, 1 - e^{\epsilon_m + it_m})$  which are linearly independent for large m).

Thus, for some  $q \in \mathcal{W}$  the orbit O(q) is strongly pseudoconvex. Since G(M) is isomorphic to  $V_{\infty}$ , the orbit O(q) is CR-equivalent to the hypersurface  $\omega$  (see Lemma 2.2). Suppose first that O(q) is CR-equivalent to  $\omega$ and let  $\Lambda$  be the equivalence map. Then  $\Lambda$  induces an automorphism of  $V_{\infty}$ (recall that G(M) is identified with  $V_{\infty}$ ), and hence an isomorphism  $\varphi_{\Lambda}$  between the Lie algebra  $\hat{\mathfrak{v}}_{\infty}$  generated by the vector fields  $\hat{X}_1$ ,  $\hat{Y}_1$ ,  $\hat{Z}_3$  on O(q)and the Lie algebra  $\mathfrak{v}_{\infty}$  of  $V_{\infty}$ . Every automorphism of  $\mathfrak{v}_{\infty}$  is either an inner automorphism corresponding to the conjugation by an element of  $V_{\infty}$ , or is a composition of such an inner automorphism and the automorphism given by

$$X_3 \mapsto -X_3, \quad Y_3 \mapsto Y_3, \quad Z_3 \mapsto -Z_3.$$

Hence  $\Lambda$  can be chosen so that  $\varphi_{\Lambda}(\hat{X}_3 - 1/2 \hat{Z}_3) = \pm (X_3 - 1/2 Z_3)$ . Arguing as in the proof of Theorem 2.1, we see that  $\Lambda$  can be extended to a holomorphic equivalence between a neighborhood of O(q) and the domain  $\Re_{s,t}$  for some 0 < s < t. Then the map  $F_{\infty} \circ \Lambda^{-1}$  transforms the vector field  $X_3 - 1/2 Z_3$ on an open subset of  $\Re_{s,t}$  into the vector field  $\pm i\partial/\partial w$ . It is not difficult to show that the general form of a locally defined biholomorphic map with this property is the composition  $\Gamma_{\infty} \circ \Psi_{\infty} \circ \Phi_{\infty}$ , where  $\Phi_{\infty}$  is defined in (3.7),  $\Psi_{\infty}$ is given by

and  $\Gamma_{\infty}$  is of the form

$$\begin{array}{rccc} z & \mapsto & r(z), \\ w & \mapsto & \pm(w + \ln(Cz)), \end{array}$$

for some branch of the logarithm, holomorphic function r and  $C \in \mathbb{C}$ . A straightforward calculation shows, however, that the composition  $\Gamma_{\infty} \circ \Psi_{\infty} \circ \Phi_{\infty}$  does not transform any of the vector fields from  $\mathfrak{v}_{\infty}$  into any of the vector fields  $\tilde{X}, \tilde{Y}$  on any open subset of  $\mathfrak{R}_{s,t}$ .

Suppose that  $G(M) = V_n$  for  $n \in \mathbb{N}$ . Fix  $q_0 \in O(p)$  and let  $f: M_{q_0} \to \mathcal{P}$ be a holomorphic equivalence that transforms  $H_{q_0}|_{M_{q_0}}$  into the group  $\mathcal{T}$  and such that  $\hat{X}_3|_{M_{q_0}}$  and  $\hat{Y}_3|_{M_{q_0}}$  are transformed into the vector fields  $i\partial/\partial z$  and  $z\partial/\partial z$ , respectively, where  $\hat{X}_3$ ,  $\hat{Y}_3$  are the vector fields on O(p) corresponding to  $X_3$ ,  $Y_3$ . For every  $q \in O(p)$  find the unique map  $g \in W_n$  such that  $gM_{q_0} =$  $M_q$  and define  $F_n(q) := \left(f(g^{-1}(q)), e^{it/n}\right) \in \mathbb{C}^2$ , where t is the parameter value corresponding to g. Clearly,  $F_n$  is a real-analytic CR-isomorphism between O(p) and  $O_2$  that transforms  $\hat{X}_3 - 1/2 \hat{Z}_3$  into  $iw/n \partial/\partial w|_{O_2}$ , where  $\hat{Z}_3$  is the vector field on O(p) corresponding to  $Z_3$ .

Denote by  $\tilde{X}$ ,  $\tilde{Y}$ ,  $\tilde{Z}$  the vector fields on  $O_2$  into which  $F_n$  transforms  $\hat{X}_3$ ,  $\hat{Y}_3$ ,  $\hat{Z}_3$ , respectively. Since  $F_n$  is real-analytic, it extends to a biholomorphic map from a neighborhood of O(p) in M onto a neighborhood of  $O_2$  in  $\mathbb{C}^2$ . Clearly,  $\hat{X}_3$ ,  $\hat{Y}_3$ ,  $\hat{Z}_3$  extend to holomorphic vector fields on all of M and hence  $\tilde{X}$ ,  $\tilde{Y}$ ,  $\tilde{Z}$  extend to holomorphic vector fields defined in a neighborhood of  $O_2$ . Since  $\tilde{X}$  and  $\tilde{Y}$  are tangent to  $O_2$  and since their restrictions to  $\mathcal{P} \times \{1\} \subset O_2$ are  $i\partial/\partial z$  and  $z\partial/\partial z$ , respectively, they have the form (3.9), where  $\rho, \sigma, \mu, \nu$ are functions holomorphic near  $O_2$  such that

$$\rho(z,1) \equiv \sigma(z,1) \equiv \mu(z,1) \equiv \nu(z,1), \qquad (3.13)$$

and

$$\sigma \overline{w} + \overline{\sigma} w \equiv \nu \overline{w} + \overline{\nu} w \equiv 0, \quad \text{for } |w| = 1.$$
(3.14)

As before, we obtain

$$\begin{bmatrix} \frac{iw}{n} \partial/\partial w, \tilde{X} \end{bmatrix} = -\frac{1}{2} \tilde{Y},$$

$$\begin{bmatrix} \frac{iw}{n} \partial/\partial w, \tilde{Y} \end{bmatrix} = 2\tilde{X} - \frac{iw}{n} \partial/\partial w.$$
(3.15)

Conditions (3.15) together with (3.9) and (3.13) uniquely determine the functions  $\rho, \sigma, \mu, \nu$  as follows:

$$\rho = \frac{i}{4} \left( (z+2)w^n - (z-2)w^{-n} \right) - i,$$
  

$$\sigma = -\frac{i}{4n} \left( w^{n+1} + w^{1-n} - 2w \right),$$
  

$$\mu = \frac{1}{2} \left( (z+2)w^n + (z-2)w^{-n} \right) - z,$$
  

$$\nu = -\frac{1}{2n} \left( w^{n+1} - w^{1-n} \right).$$

The functions  $\sigma$ ,  $\nu$  automatically satisfy conditions (3.14).

Integrating  $\tilde{X}$ ,  $\tilde{Y}$ ,  $\mathcal{Z}$  we respectively obtain the following one-parameter subgroups of holomorphic transformations of  $O_2$ :

$$\begin{aligned} z &\mapsto zw^{-n} \left( w^n + \frac{i}{4}(w^n - 1)t \right) \left( 1 + \frac{i}{4}(w^n - 1)t \right) + \\ &\qquad \frac{i}{2} \left[ t(w^n + w^{-n}) + t^2 \frac{i}{4}(w^n - 1)(w^{-n} + 1) \right], \\ H_{\tilde{X}}: \\ w &\mapsto \left( \frac{w^n + \frac{i}{4}(w^n - 1)t}{1 + \frac{i}{4}(w^n - 1)t} \right)^{1/n}, \end{aligned}$$

$$\begin{aligned} z &\mapsto \frac{1}{4} \Big( z \left[ \left( 2 + w^n + w^{-n} \right) e^t + \left( 2 - w^n - w^{-n} \right) e^{-t} \right] + \\ & 2 \left( w^n - w^{-n} \right) \left( e^t - e^{-t} \right) \Big), \\ H_{\tilde{Y}}: \\ w &\mapsto \left( \frac{w^n + 1 + (w^n - 1)e^{-t}}{w^n + 1 - (w^n - 1)e^{-t}} \right)^{1/n}, \end{aligned}$$

where  $t \in \mathbb{R}$ ,  $\tau^{1/n} = \exp(1/n \operatorname{\tilde{ln}}^w \tau)$ , where  $\operatorname{\tilde{ln}}^w := \ln'_0 + 2\pi i k$  for  $2\pi i k/n < \arg w < 2\pi i (k+1)/n$ , and  $\operatorname{\tilde{ln}}^{2\pi i k/n} \equiv 2\pi i k$ ,  $0 \le k \le n-1$ , with  $\ln'_0$  defined as above.

Denote by  $\tilde{V}_n$  the Lie group of CR-transformations of  $O_2$  into which  $F_n$  transforms  $G(M)|_{O(p)}$ . Clearly,  $\tilde{V}_n$  is isomorphic to  $V_n$  and  $H_{\tilde{X}}$ ,  $H_{\tilde{Y}}$ ,  $H_{\mathcal{Z}}$  are one-parameter subgroups of  $\tilde{V}_n$ . Observe that the action of  $\tilde{V}_n$  on  $O_2$  does not extend to an action on any neighborhood of  $O_2$  in  $\mathbb{C}^2$  since for every point  $p_0 = (z_0, w_0) \in \mathbb{C}^2$  with  $|w_0| \neq 1$  there exists an element of  $\tilde{V}_n$  that has a singularity at  $p_0$ .

As before, there exist a neighborhood  $\mathcal{W}$  of O(p) in M such that O(q) is a real hypersurface in M for every  $q \in \mathcal{W}$ . Choosing  $\mathcal{W}$  sufficiently small, we can assume that  $F_n$  extends to  $\mathcal{W}$  to a biholomorphic map onto a neighborhood  $\mathcal{W}'$  of  $O_2$  in  $\mathbb{C}^2$ . Suppose first that for every  $q \in \mathcal{W}$  the orbit O(q) is Levi-flat. Let  $\hat{q} := F_n^{-1}(1,1)$  and  $q_m := F_n^{-1}(1,\epsilon_m)$  with  $\epsilon_m \to 1$  as  $m \to \infty$ . Let  $M_{q_m}^{O(q_m)}$  be the leaf of the foliation on  $O(q_m)$  passing through  $q_m$ . It now follows that for every m there exists  $g_m \in W_n$ , such that  $g_m H_{\hat{q}} g_m^{-1}$  is the connected subgroup of G(M) corresponding to the subalgebra of vector fields on  $O(q_m)$  tangent to  $M_{q_m}^{O(q_m)}$ . The sequence  $\{g_m\}$  can be chosen so that  $g_m \to \text{id as } m \to \infty$ .

At the same time, conjugating the one-parameter subgroups  $H_{\tilde{X}}$  and  $H_{\tilde{Y}}$ by the element of  $H_{\mathcal{Z}}$  corresponding to  $g_m$  with  $t = t_m$  and computing the vector fields corresponding to the conjugated subgroups, we see that these vector fields at the point  $(1, \epsilon_m)$  span all of  $\mathbb{C}^2$  rather than a complex line in  $\mathbb{C}^2$  (the span coincides with that of the pair of vectors  $(3\epsilon_m^n e^{it_m}, \epsilon_m/n - \epsilon_m^{n+1}e^{it_m}/n)$  and  $(-1, \epsilon_m/n - \epsilon_m^{n+1}e^{it_m}/n)$  which are linearly independent for large m).

Thus, for some  $q \in \mathcal{W}$  the orbit O(q) is strongly pseudoconvex. Since G(M) is isomorphic to  $V_n$ , the orbit O(q) is CR-equivalent to a hypersurface  $\varepsilon_{n/k}$ , where  $k \in \mathbb{N}$  is such that (n, k) = 1 (see Lemma 2.2). Let  $\Lambda$  be the equivalence map. Then  $\Lambda$  induces an isomorphism  $V_n \to V_{n/k}$ , and hence an isomorphism  $\varphi_{\Lambda}$  between the respective Lie algebras  $\hat{\mathfrak{v}}_n$  and  $\mathfrak{v}_{n/k}$ . Here  $\hat{\mathfrak{v}}_n$  is generated by the vector fields  $\hat{X}_3, \hat{Y}_3, \hat{Z}_3$  on O(q) and  $\mathfrak{v}_{n/k}$  is generated by the vector fields  $X_{3,n/k}, Z_{3,n/k}$  defined as in (3.6) where n is replaced

with n/k. Every automorphism of  $\mathfrak{v}_{n/k}$  is either an inner automorphism corresponding to the conjugation by an element of  $V_{n/k}$ , or is a composition of such an inner automorphism and the automorphism given by

$$X_{3,n/k} \mapsto -X_{3,n/k}, \quad Y_{3,n/k} \mapsto Y_{3,n/k}, \quad Z_{3,n/k} \mapsto -Z_{3,n/k}.$$

Hence  $\Lambda$  can be chosen so that  $\varphi_{\Lambda}(\hat{X}_3 - 1/2\hat{Z}_3) = \pm (X_{3,n/k} - 1/2Z_{3,n/k})$ . Arguing as in the proof of Theorem 2.1, we see that a neighborhood of O(q) is holomorphically equivalent to the domain  $\Re_{n/k,s,t}$  for some 0 < s < t. Then the map  $F_n \circ \Lambda^{-1}$  transforms the vector field  $X_{3,n/k} - 1/2Z_{3,n/k}$  on an open subset of  $\Re_{n/k,s,t}$  into the vector field  $\pm iw\partial/\partial w$ . It is not difficult to show that the general form of a locally defined map with this property is the composition  $\Gamma_n \circ \Psi_n \circ \Phi_n \circ \Pi_k$ , where  $\Phi_n$  is defined in (3.8),  $\Psi_n$  is given by

$$\begin{array}{rccc} z & \mapsto & zw^{-n}, \\ w & \mapsto & w, \end{array}$$

 $\Gamma_n$  is of the form

$$\begin{array}{rccc} z & \mapsto & r(z), \\ w & \mapsto & \pm (Cz)^{1/n} w, \end{array}$$

for some branch of  $\sqrt[n]{}$ , holomorphic function r and  $C \in \mathbb{R}$ , and  $\Pi_k$  is given by

It is straightforward to verify, however, that the composition  $\Gamma_n \circ \Psi_n \circ \Phi_n \circ \Pi_k$ does not transform any of the vector fields from  $\mathfrak{v}_{n/k}$  into any of the vector fields  $\tilde{X}$ ,  $\tilde{Y}$  on any open subset of  $\mathfrak{R}_{n/k,s,t}$ .

Thus,  $\mathfrak{g}$  cannot be given by relations (R3), which completes the proof of the proposition.

We will now prove the following theorem that finalizes the classification of (2,3)-manifolds in the case when every orbit is a real hypersurface.

**THEOREM 3.2** Let M be a (2,3)-manifold. Assume that the G(M)-orbit of every point in M has codimension 1 and that at least one orbit is Levi-flat.

Then M is holomorphically equivalent to one of the following domains:

(i)  $S + i\mathbb{R}^2$ , where S is an angle with vertex at the origin of size less than  $\pi$  in the (Re z, Re w)-plane;

(ii) 
$$\begin{cases} (z,w) \in \mathbb{C}^2 : s (\operatorname{Re} z)^b < \operatorname{Re} w < t (\operatorname{Re} z)^b, \ \operatorname{Re} z > 0 \end{cases}, \text{ where } |b| \ge 1, \\ b \ne 1, -\infty \le s < 0 < t \le \infty \text{ and at least one of } s, t \text{ is finite;} \end{cases}$$

(iii) 
$$\left\{ (z,w) \in \mathbb{C}^2 : \operatorname{Re} w > r_{s,t}(z) |\operatorname{Re} z|^b \right\}$$
, where  $b > 1$ ,

$$r_{s,t}(z) := \begin{cases} s & \text{if } \operatorname{Re} z > 0, \\ t & \text{if } \operatorname{Re} z < 0, \\ 0 & \text{if } \operatorname{Re} z = 0, \end{cases}$$

and  $s, t \in \mathbb{R}$  are such that either  $s \cdot t < 0$ , or  $s \ge 0$ ,  $t \ge 0$ ,  $s^2 + t^2 > 0$ , and in the latter case  $s \ne t$  for b = 2;

(iv)  $\left\{ (z,w) \in \mathbb{C}^2 : 0 < \operatorname{Re} w < r(z) |\operatorname{Re} z|^b \right\}$ , where  $b \leq -1$ , r(z) as in (iii), and  $s \geq 0, t \geq 0, s^2 + t^2 > 0$ ;

(v) 
$$\{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z < R_{s,t}(w)\}, \text{ where }$$

$$R_{s,t}(w) := \begin{cases} Rew \cdot \ln(sRew) & \text{if } Rew > 0\\ Rew \cdot \ln(-tRew) & \text{if } Rew < 0\\ 0 & \text{if } Rew = 0 \end{cases}$$

with s, t > 0.

**Proof:** We will use the notation introduced earlier in this section. Observe that the set  $\mathfrak{L} := \{p \in M : O(p) \text{ is Levi-flat}\}$  is closed in M. Hence, if  $\mathfrak{L}$  is also open, then every orbit in M is Levi-flat. Let  $p \in \mathfrak{L}$  and suppose first that there exists a CR-isomorphism  $f : O(p) \to O_1$  that transforms G(M) into the group  $G_0$ . Since  $G_0$  acts on all of  $\mathcal{C}$ , arguing as on step (II) of the orbit gluing procedure introduced in the proof of Theorem 2.1, we extend f to a biholomorphic map between a G(M)-invariant neighborhood U of O(p) and a  $G_0$ -invariant neighborhood of  $O_1$  in  $\mathcal{C}$  that satisfies (2.45) for all  $g \in G(M)$ and  $q \in U$ , where  $\varphi : G(M) \to G_0$  is an isomorphism. Since every  $G_0$ -orbit in  $\mathcal{C}$  has the form

$$\mathfrak{b}_r := \{ (z, w) \in \mathcal{C} : \operatorname{Re} w = r \},\$$

for  $r \in \mathbb{R}$ , and hence is Levi-flat, the set  $\mathfrak{L}$  is open, and thus every orbit in M is Levi-flat. Since the group  $G_0$  is not isomorphic to any of the groups  $G_b$ 

for  $b \in \mathbb{R}^*$ , G and  $G'_0$ , it follows that every orbit in M is CR-equivalent to  $O_1$  by means of a map that transforms G(M) into  $G_0$ .

We will now further utilize the orbit gluing procedure from the proof of Theorem 2.1. Consider a maximal G(M)-invariant domain  $D \subset M$  from which there exists a biholomorphic map f onto a  $G_0$ -invariant domain in  $\mathcal{C}$ satisfying (2.45) for all  $g \in G(M), q \in D$  and some isomorphism  $\varphi: G(M) \to G(M)$  $G_0$ . Assume that  $D \neq M$  and consider  $x \in \partial D$ . Let  $f_1: O(x) \to O_1$  be a real-analytic CR-isomorphism satisfying (2.45) for all  $g \in G(M), q \in O(x)$ and an isomorphism  $\varphi_1: G(M) \to G_0$  in place of  $\varphi$ . As before, extend  $f_1$ to a biholomorphic map from a connected G(M)-invariant neighborhood V of O(x) onto a  $G_0$ -invariant neighborhood of  $O_1$  in  $\mathcal{C}$ . The extended map satisfies (2.45) for all  $g \in G(M)$ ,  $q \in V$  and  $\varphi_1$  in place of  $\varphi$ . Consider  $s \in V \cap D$ . The maps f and  $f_1$  take O(s) onto  $G_0$ -orbits  $\mathfrak{b}_{r_1}$  and  $\mathfrak{b}_{r_2}$ , respectively. Then  $F := f_1 \circ f^{-1}$  establishes a *CR*-isomorphism between  $\mathfrak{b}_{r_1}$ and  $\mathfrak{b}_{r_2}$ . We will now show that F extends to an automorphism of C. Clearly, F has the form  $F = \nu \circ g$ , where  $\nu$  is a real translation in the w-variable, and  $g \in \operatorname{Aut}_{CR}(\mathfrak{b}_{r_1})$ . Each of the maps f and  $f_1$  transforms the group G(M)into the group  $G_0$ , and therefore g lies in the normalizer of  $G_0$  in  $\operatorname{Aut}_{CR}(\mathfrak{b}_{r_1})$ . The general form of an element of  $\operatorname{Aut}_{CR}(\mathfrak{b}_{r_1})$  is

$$(z, r_1 + iv) \mapsto (a_v(z), r_1 + i\mu(v)),$$
 (3.16)

where  $v \in \mathbb{R}$ ,  $a_v \in \operatorname{Aut}(\mathcal{P})$  for every v, and  $\mu$  is a diffeomorphism of  $\mathbb{R}$ . Considering g in this form, we obtain that  $a_v \in \mathcal{T}$  for every v and that  $a_{v_1}aa_{v_1}^{-1} = a_{v_2}aa_{v_2}^{-1}$  for all  $a \in \mathcal{T}$  and all  $v_1, v_2 \in \mathbb{R}$ . Therefore,  $a_{v_1}a_{v_2}^{-1}$  lies in the center of  $\mathcal{T}$ , which is trivial. Hence we obtain that  $a_{v_1} = a_{v_2}$  for all  $v_1, v_2$ . In addition, there exists  $k \in \mathbb{R}^*$  such that  $\mu^{-1}(v) + c \equiv \mu^{-1}(v + kc)$ , for  $c \in \mathbb{R}$ . Differentiating this identity with respect to c at 0 gives

$$\mu^{-1}(v) = v/k + t_0 \tag{3.17}$$

for some  $t_0 \in \mathbb{R}$ . Therefore, F extends to a holomorphic automorphism of C as the following map:

$$\begin{array}{lll} z & \mapsto & \lambda z + i\beta, \\ w & \mapsto & kw + \sigma - ikt_0, \end{array} \tag{3.18}$$

where  $\lambda > 0, \beta, \sigma \in \mathbb{R}$ .

Since  $O_1$  splits  $\mathcal{C}$ , for V small enough we have  $V = V_1 \cup V_2 \cup O(x)$ , where  $V_j$  are open connected non-intersecting sets. For each  $j, D \cap V_j$  is connected,

and, if V is sufficiently small, each  $V_j$  is either a subset of D or disjoint from it. Suppose first that there is only one j for which  $D \cap V_j \neq \emptyset$ . In this case  $D \cap V$  is connected and  $V \setminus (D \cup O(x)) \neq \emptyset$ . Then the map  $\tilde{f}$ defined by formula (2.46) gives a biholomorphic extension of f to  $D \cup V$ . By construction,  $\tilde{f}$  satisfies (2.45) for  $g \in G(M)$  and  $q \in D \cup V$ . Since  $D \cup V$  is strictly larger than D, we obtain a contradiction with the maximality of D. Thus, in this case D = M, and hence M is holomorphically equivalent to a  $G_0$ -invariant domain in C.

Suppose now that  $V_j \,\subset D$  for j = 1, 2. Applying formula (2.46) we can extend  $f|_{V_1}$  and  $f|_{V_2}$  to biholomorphic maps  $\hat{f}_1$ ,  $\hat{f}_2$ , respectively, from a neighborhood of O(x) into  $\mathcal{C}$ ; each of these maps satisfies (2.45). Let  $\hat{f}_j(O(x)) = \mathfrak{b}_{\rho_j}, j = 1, 2$ . Clearly,  $\rho_1 \neq \rho_2$ . The map  $\hat{F} := \hat{f}_2 \circ \hat{f}_1^{-1}$  is a CRisomorphism from  $\mathfrak{b}_{\rho_1}$  onto  $\mathfrak{b}_{\rho_2}$ . It follows, as before, that  $\hat{F}$  has the form (3.18) with  $\sigma \neq \rho_1(1-k)$ . Let  $\Gamma$  denote the cyclic group of automorphisms of  $\mathcal{C}$  generated by  $\hat{F}$ . This group acts freely properly discontinuously on  $\mathcal{C}$ . Let  $\mathcal{C}_{\Gamma}$  be the complex manifold obtained by factoring  $\mathcal{C}$  by the action of  $\Gamma$ and let  $\Pi_{\Gamma} : \mathcal{C} \to \mathcal{C}_{\Gamma}$  be the factorization map. Then the map  $\Pi_{\Gamma} \circ \mathcal{F}$ , where  $\mathcal{F}$  is given by formula (2.47) establishes holomorphic equivalence between Mand  $\mathcal{C}_{\Gamma}$ . Observe, however, that  $\mathcal{C}_{\Gamma}$  is not hyperbolic, since it is covered by  $\mathcal{C}$ which is not hyperbolic. This contradiction shows that in fact exactly one of  $V_j, j = 1, 2$ , is a subset of D, and hence M is holomorphically equivalent to a  $G_0$ -invariant domain in  $\mathcal{C}$ , that is, to a domain of the form

$$\left\{ (z, w) \in \mathbb{C}^2 : \operatorname{Re} z > 0, \, r < \operatorname{Re} w < R \right\},\$$

for some  $0 \le r < R \le \infty$ . Thus, M is equivalent to  $\Delta^2$  which is impossible, since  $d(\Delta^2) = 6$ .

Next, if for  $p \in \mathfrak{L}$  there exists a CR-isomorphism between O(p) and  $O_2$  that transforms G(M) into the group  $G'_0$ , a similar argument gives that M is holomorphically equivalent to the product of  $\Delta \times A$ , where A is either an annulus or a punctured disk. This is impossible either since  $d(\Delta \times A) = 4$ .

Let now  $p \in \mathfrak{L}$  and suppose that there exists a CR-isomorphism  $f : O(p) \to O_1$  that transforms G(M) into the group  $G_1$ . The group  $G_1$  acts on

$$\mathcal{D} := \mathbb{C}^2 \setminus \left\{ (z, w) \in \mathbb{C}^2 : \operatorname{Re} z = \operatorname{Re} w = 0 \right\},\$$

and, as as before, we can extend f to a biholomorphic map between a G(M)invariant neighborhood U of O(p) and a  $G_1$ -invariant neighborhood of  $O_1$  in

 $\mathcal{D}$  that satisfies (2.45) for all  $g \in G(M)$  and  $q \in U$ , where  $\varphi : G(M) \to G_1$  is an isomorphism. A  $G_1$ -orbit in  $\mathcal{D}$  is either of the form

$$\left\{ (z, w) \in \mathbb{C}^2 : \operatorname{Re} w = r \operatorname{Re} z, \operatorname{Re} z > 0 \right\},\$$

or of the form

$$\left\{ (z,w) \in \mathbb{C}^2 : \operatorname{Re} w = r \operatorname{Re} z, \operatorname{Re} z < 0 \right\},\$$

for  $r \in \mathbb{R}$ , or coincides with either  $O'_1$  (see (3.5)), or

$$O_1'' := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z = 0, \operatorname{Re} w < 0\}, \qquad (3.19)$$

and hence is Levi-flat. Therefore the set  $\mathfrak{L}$  is open, and thus every orbit in M is Levi-flat. Since the group  $G_1$  is not isomorphic to any of the groups  $G_b$  for  $b \in \mathbb{R}^*$ ,  $b \neq 1$ , and G, it follows that every orbit in M is CR-equivalent to  $O_1$  by means of a map that transforms G(M) into  $G_1$ .

As before, consider a maximal G(M)-invariant domain  $D \subset M$ ,  $x \in \partial D$ , and a CR-isomorphism  $f_1 : O(x) \to O_1$ . Extend  $f_1$  to a biholomorphic map from a connected G(M)-invariant neighborhood V of O(x) onto a  $G_1$ invariant neighborhood of  $O_1$  in  $\mathcal{D}$  and consider  $s \in V \cap D$ . The maps f and  $f_1$  take O(s) onto some  $G_1$ -orbits  $o_1$ ,  $o_2$ , respectively. Clearly,  $F := f_1 \circ f^{-1}$ has the form  $F = \nu \circ g$ , where  $\nu$  is a map of one of the forms

with  $\sigma \in \mathbb{R}$ , and  $g \in \operatorname{Aut}_{CR}(o_1)$ . Any two  $G_1$ -orbits are equivalent by means of a map of one of the forms (3.20).

As before, g lies in the normalizer of  $G_1$  in  $\operatorname{Aut}_{CR}(o_1)$ . Let  $\mathfrak{X}$  be a map of the form (3.20) that transforms  $o_1$  into  $O_1$  and  $g_{\mathfrak{X}} := \mathfrak{X} \circ g \circ \mathfrak{X}^{-1}$ . Considering  $g_{\mathfrak{X}}$  in the general form (3.16) with  $r_1 = 0$  we see that for every  $\lambda > 0$ ,  $\beta, \gamma \in \mathbb{R}$ , the composition  $a_{\lambda v+\gamma} \circ a^{\lambda,\beta} \circ a_v^{-1}$ , where  $a^{\lambda,\beta}(z) := \lambda z + i\beta$ , belongs to  $\mathcal{T}$  and is independent of v. This implies that  $a_v(z) = \lambda_0 z + i(C_1 v + C_2)$  for some  $\lambda_0 > 0, C_1, C_2 \in \mathbb{R}$ . Also, for every  $\lambda > 0, \gamma \in \mathbb{R}$  there exist  $\lambda_1 > 0, \gamma_1 \in \mathbb{R}$ such that  $\mu(\lambda \mu^{-1}v + \gamma) = \lambda_1 v + \gamma_1$ , which gives that  $\mu^{-1}$ , as before, has the form (3.17) for some  $k \in \mathbb{R}^*$ ,  $t_0 \in \mathbb{R}$ . It then follows that  $g_{\mathfrak{X}}$  extends to a holomorphic automorphism of  $\mathcal{D}$  as follows:

$$\begin{array}{rcl}
z & \mapsto & \lambda_0 z + C_1 w + i C_2, \\
w & \mapsto & k w - i k t_0,
\end{array}$$
(3.21)

and thus F extends to an automorphism of  $\mathcal{D}$  as well.

As before, choose a neighborhood V of O(x) such that  $V = V_1 \cup V_2 \cup O(x)$ and suppose first that there is only one j for which  $D \cap V_j \neq \emptyset$ . Arguing as in the case of the group  $G_0$ , we now see that M is holomorphically equivalent to a  $G_1$ -invariant domain in  $\mathcal{D}$ . Every such a domain clearly has the form  $S + i\mathbb{R}^2$ , where S is an angle with vertex at the origin in the (Re z, Re w)plane. Such a domain is only hyperbolic if the size of S is less than  $\pi$ , and we obtain (i) of the theorem.

Suppose now that  $V_j \subset D$  for j = 1, 2. As before, applying formula (2.46) we can extend  $f|_{V_1}$  and  $f|_{V_2}$  to biholomorphic maps  $\hat{f}_1, \hat{f}_2$ , respectively, from a neighborhood of O(x) into  $\mathcal{D}$ , and let  $\hat{f}_j(O(x)) = \hat{o}_j, j = 1, 2$ . Clearly, the  $G_1$ -orbits  $\hat{o}_1$  and  $\hat{o}_2$  are distinct. The map  $\hat{F} := \hat{f}_2 \circ \hat{f}_1^{-1}$  is a CRisomorphism from  $\hat{o}_1$  onto  $\hat{o}_2$ , and it follows that  $\hat{F}$  has the same form as the map F introduced above, where  $\nu$  is non-trivial. Then the cyclic group  $\Gamma$  generated by  $\hat{F}$  acts freely properly discontinuously on  $\mathcal{D}$ , which gives, as before, that M is biholomorphically equivalent to  $\mathcal{D}_{\Gamma} := \mathcal{D}/\Gamma$ . However,  $\mathcal{D}_{\Gamma}$ is not hyperbolic since it is covered by the non-hyperbolic domain  $\mathcal{D}$ , and thus exactly one of  $V_j, j = 1, 2$ , is a subset of D; hence M is holomorphically equivalent to a  $G_1$ -invariant domain in  $\mathcal{D}$ , as specified above.

Let now  $p \in \mathfrak{L}$  and suppose that there exists a CR-isomorphism  $f : O(p) \to O_1$  that transforms G(M) into the group  $G_b$  for some  $b \in \mathbb{R}^*, b \neq 1$ . The group  $G_b$  acts on  $\mathcal{D}$ , and, as as before, we can consider an extension of f. Every  $G_b$ -orbit in  $\mathcal{D}$  is either strongly pseudoconvex and has one of the forms

$$\left\{ (z,w) \in \mathbb{C}^2 : \operatorname{Re} w = r (\operatorname{Re} z)^b, \operatorname{Re} z > 0 \right\},\$$
$$\left\{ (z,w) \in \mathbb{C}^2 : \operatorname{Re} w = r (-\operatorname{Re} z)^b, \operatorname{Re} z < 0 \right\},\$$

for  $r \in \mathbb{R}^*$ , or coincides with one of  $O_1, O'_1, O''_1$  (see (3.5), (3.19)), and

$$O_1''' := \{(z, w) \in \mathbb{C}^2 : \operatorname{Re} z < 0, \operatorname{Re} w = 0\}.$$
(3.22)

It then follows that every Levi-flat orbit in M has a G(M)-invariant neighborhood in which every other orbit is strongly pseudoconvex. Among the groups  $G_c$  for  $c \in \mathbb{R}^*$ ,  $c \neq 1$ , and G, the only group isomorphic to  $G_b$  is  $G_{1/b}$ . Thus, it follows that every Levi-flat orbit in M is CR-equivalent to  $O_1$  by means of a map that transforms G(M) into either  $G_b$  or  $G_{1/b}$ . In the latter case interchanging the variables we obtain a map that takes the Levi-flat orbit into  $O'_1$  and transforms G(M) into  $G_b$ . Next, it follows from Lemma 2.2 that every strongly pseudoconvex orbit is CR-equivalent to  $\tau_b$  (see (2.2)) by means of a CR-map that transforms G(M) into  $G_b$ .

As before, consider a maximal G(M)-invariant domain  $D \subset M$ ,  $x \in \partial D$ and a CR-map  $f_1 : O(x) \to O$ , where O is either  $O_1$  or  $O'_1$  or  $\tau_b$ . As before, extend  $f_1$  to a biholomorphic map from a connected G(M)-invariant neighborhood V of O(x) onto a  $G_b$ -invariant neighborhood of O in  $\mathcal{D}$ , and consider  $s \in V \cap D$  such that O(s) is strongly pseudoconvex. The maps f and  $f_1$  take O(s) onto some strongly pseudoconvex  $G_b$ -orbits. Then  $F := f_1 \circ f^{-1}$ establishes a CR-isomorphism between the orbits, and thus has the form  $F = \nu \circ g$ , where  $\nu$  is a map of the form

with  $k \in \mathbb{R}^*$ , and  $g \in G_b$ . Therefore, F extends to an automorphism of  $\mathcal{D}$ .

As before, choose a neighborhood V of O(x) such that  $V = V_1 \cup V_2 \cup O(x)$ . If there is only one j for which  $D \cap V_j \neq \emptyset$ , then, as before, we obtain that M is holomorphically equivalent to a  $G_b$ -invariant hyperbolic domain in  $\mathcal{D}$ . This leads to (ii), (iii), (iv) of the theorem (note that interchanging the variables transforms  $G_b$ -invariant domains into  $G_{1/b}$ -invariant domains).

Suppose now that  $V_j \subset D$  for j = 1, 2. As before, consider the *CR*isomorphism  $\hat{F} : \hat{o}_1 \to \hat{o}_2$ . If O(x) is strongly pseudoconvex, then  $\hat{F} = \nu \circ g$ , where  $\nu$  is a non-trivial map of the form (3.23), and  $g \in G_b$ . Suppose now that O(x) is Levi-flat. Then  $\hat{F} = \nu \circ g$ , where  $\nu$  is a non-trivial map of one of the forms

and  $g \in \operatorname{Aut}_{CR}(\hat{o}_1)$ . In this case, either g lies in the normalizer of  $G_b$  in  $\operatorname{Aut}_{CR}(\hat{o}_1)$ , or  $gG_bg^{-1} = G_{1/b}$  and  $gG_{1/b}g^{-1} = G_b$ . Transforming  $\hat{o}_1$  into  $O_1$  by a map  $\mathfrak{X}$  of one of the forms (3.24) and arguing as in the case of the group  $G_1$ , we obtain that  $g_{\mathfrak{X}} := \mathfrak{X} \circ g \circ \mathfrak{X}^{-1}$  extends to a holomorphic automorphism of  $\mathcal{D}$  as a map of the form (3.21) with  $C_1 = 0$ .

It now follows that the cyclic group  $\Gamma$  generated by  $\tilde{F}$  acts freely properly discontinuously on  $\mathcal{D}$ , which leads, as before, to holomorphic equivalence between M and the non-hyperbolic manifold  $\mathcal{D}/\Gamma$ . This contradiction shows that in fact exactly one of  $V_j$ , j = 1, 2, is a subset of D, and hence M is holomorphically equivalent to a  $G_b$ -invariant domain in  $\mathcal{D}$ . Let now  $p \in \mathfrak{L}$  and suppose that there exists a CR-isomorphism  $f : O(p) \to O_1$  that transforms G(M) into the group G. The group G acts on  $\mathcal{D}$ , and we can consider an extension of f. Every G-orbit in  $\mathcal{D}$  is either strongly pseudoconvex and has one of the forms

$$\left\{ (z,w) \in \mathbb{C}^2 : \operatorname{Re} z = \operatorname{Re} w \ln (r \operatorname{Re} w), \operatorname{Re} w > 0 \right\},$$
$$\left\{ (z,w) \in \mathbb{C}^2 : \operatorname{Re} z = \operatorname{Re} w \ln (-r \operatorname{Re} w), \operatorname{Re} w < 0 \right\},$$

for r > 0, or coincides with one of  $O_1$ ,  $O_1'''$  (see (3.22)).

It then follows that every Levi-flat orbit in M has a G(M)-invariant neighborhood in which every other orbit is strongly pseudoconvex, that every Levi-flat orbit in M is CR-equivalent to  $O_1$  by means of a map that transforms G(M) into G and that every strongly pseudoconvex orbit is CRequivalent to  $\xi$  by means of a map that transforms G(M) into G.

Consider a maximal G(M)-invariant domain  $D \subset M$ ,  $x \in \partial D$  and a CR-map  $f_1: O(x) \to O$ , where O is either  $O_1$  or  $\xi$ . As before, extend  $f_1$  to a biholomorphic map from a connected G(M)-invariant neighborhood V of O(x) onto a G-invariant neighborhood of O in  $\mathcal{D}$ , and consider  $s \in V \cap D$  such that O(s) is strongly pseudoconvex. The maps f and  $f_1$  take O(s) onto some strongly pseudoconvex G-orbits. Then  $F := f_1 \circ f^{-1}$  establishes a CR-isomorphism between the orbits, and thus has the form  $F = \nu \circ g$ , where  $\nu$  is a map of the form

with  $k \in \mathbb{R}^*$ , and  $g \in G$ . Therefore, F extends to an automorphism of  $\mathcal{D}$ .

As before, choose a neighborhood V of O(x) such that  $V = V_1 \cup V_2 \cup O(x)$ . If there is only one j for which  $D \cap V_j \neq \emptyset$ , then, as before, we obtain that M is holomorphically equivalent to a G-invariant hyperbolic domain in  $\mathcal{D}$ . This leads to (v) of the theorem.

Suppose now that  $V_j \,\subset\, D$  for j = 1, 2. As before, consider the *CR*isomorphism  $\hat{F} : \hat{o}_1 \to \hat{o}_2$ . If O(x) is strongly pseudoconvex, then  $\hat{F} = \nu \circ g$ , where  $\nu$  is a non-trivial map of the form (3.25), and  $g \in G$ . Suppose now that O(x) is Levi-flat. Then  $\hat{F} = \nu \circ g$ , where  $\nu$  is map (3.25) with k = -1, and  $g \in \operatorname{Aut}_{CR}(\hat{o}_1)$ . The element g lies in the normalizer of G in  $\operatorname{Aut}_{CR}(\hat{o}_1)$ . Transforming  $\hat{o}_1$  into  $O_1$  by a map  $\mathfrak{X}$  of the form (3.25) with  $k = \pm 1$  and considering  $g_{\mathfrak{X}} := \mathfrak{X} \circ g \circ \mathfrak{X}^{-1}$  in the general form (3.16) with  $r_1 = 0$ , we obtain, as before, that  $\mu^{-1}$  has the form (3.17) for some  $k \in \mathbb{R}^*$ ,  $t_0 \in \mathbb{R}$  and that  $a_v(z) = \lambda_0 z + i\beta(v)$ , where  $\lambda_0 > 0$  and  $\beta(v)$  is a function satisfying for every  $\lambda > 0$  and  $\gamma \in \mathbb{R}$  the following condition:

$$\beta \left( \lambda \mu^{-1}(v) + \gamma \right) - \lambda \beta \left( \mu^{-1}(v) \right) + \lambda \ln \lambda \left( \lambda_0 \mu^{-1}(v) - v \right) + \lambda_0 \beta \equiv \text{const.}$$

This gives that  $g_{\mathfrak{X}}$  extends to an automorphism of  $\mathcal{D}$  as a map of the form (3.21). It now follows that the cyclic group  $\Gamma$  generated by  $\hat{F}$  acts freely properly discontinuously on  $\mathcal{D}$ , which leads, as before, to to holomorphic equivalence between M and the non-hyperbolic manifold  $\mathcal{D}/\Gamma$ . This contradiction shows that in in fact exactly one of  $V_j$ , j = 1, 2, is a subset of D, and hence M is holomorphically equivalent to a G-invariant domain in  $\mathcal{D}$ .

The theorem is proved.

# References

- [1] Alekseevsky, A. V. and Alekseevsky, D. V., Riemann *G*-manifolds with one-dimensional orbit space, *Ann. Global Anal. Geom.* 11(1993), 197–211.
- [2] Cartan, E., Sur la géométrie pseudo-conforme des hypersurfaces de l'espace de deux variables complexes, I Ann. Math. Pura Appl. 11(1933), 17–90 (or Oeuvres Completes II, 2, 1231–1304), II Ann. Scuola Norm. Sup. Pisa 1(1933), 333–354 (or Oeuvres Completes III, 2, 1217–1238).
- [3] Gifford, J. A., Isaev, A. V. and Krantz, S. G., On the dimensions of the automorphism groups of hyperbolic Reinhardt domains, *Illinois J. Math.* 44(2000), 602–618.
- [4] Isaev, A. V., Hyperbolic *n*-dimensional manifolds with automorphism group of dimension  $n^2$ , to appear in *Geom. Funct. Anal. (GAFA)*, available from Mathematics ArXiv at http://front.math.ucdavis.edu/math.CV/0502152
- [5] Isaev, A. V., Hyperbolic manifolds of dimension n with automorphism group of dimension  $n^2 1$ , J. Geom. Analysis 15(2005), 239–259.

- [6] Isaev, A. V., Analogues of Rossi's map and E. Cartan's classification of homogeneous strongly pseudoconvex 3-dimensional hypersurfaces, J. Lie Theory 16(2006), 407–426.
- [7] Isaev, A. V. and Krantz, S. G., On the automorphism groups of hyperbolic manifolds, *J. Reine Angew. Math.* 534(2001), 187–194.
- [8] Isaev, A. V. and Kruzhilin, N. G., Effective actions of  $SU_n$  on complex *n*-dimensional manifolds, *Illinois J. Math.* 48(2004), 37–57.
- [9] Kaup, W., Reelle Transformationsgruppen und invariante Metriken auf komplexen Räumen, *Invent. Math.* 3(1967), 43–70.
- [10] Kim, K.-T. and Verdiani, L., Complex *n*-dimensional manifolds with a real  $n^2$ -dimensional automorphism group, J. Geom. Analysis 14 (2004), 701–713.
- [11] Kobayashi, S., *Hyperbolic Manifolds and Holomorphic Mappings*, Marcel Dekker, New York 1970.
- [12] Rossi, H., Attaching analytic spaces to an analytic space along a pseudoconcave boundary, Proc. Conf. Complex Analysis, Minneapolis, 1964, Springer-Verlag, 1965, 242–256.

Department of Mathematics The Australian National University Canberra, ACT 0200 AUSTRALIA E-mail: alexander.isaev@maths.anu.edu.au