

**A DEGREE THEORETIC APPROACH TO  
THE SOLVABILITY OF SYMMETRIC WORD  
EQUATIONS IN POSITIVE DEFINITE LETTERS**

SCOTT N. ARMSTRONG AND CHRISTOPHER J. HILLAR

ABSTRACT. Let  $S(X, B)$  be a symmetric (“palindromic”) word in two letters  $X$  and  $B$ . A theorem due to Hillar and Johnson states that for each pair of positive definite matrices  $B$  and  $P$ , there is a positive definite solution  $X$  to the word equation  $S(X, B) = P$ . They also conjectured that these solutions are unique. In this paper, we resolve this conjecture (negatively). Furthermore, we prove that, generically, the number of solutions is odd (and thus finite) in the real case. Our approach utilizes the theory of Brouwer degree and also provides a second proof of existence of such solutions in the real case.

1. INTRODUCTION

In this paper, we consider a natural matrix generalization to the elementary scalar equation

$$bx^s = p,$$

in which  $b > 0$ ,  $p \geq 0$ ,  $s \in \mathbb{Z}_+$  and  $x$  is a nonnegative real indeterminate. One difficulty with an extension is dealing with matrix noncommutativity, while another is determining what should be meant by the words “real” and “nonnegative.” Fortunately for us, the latter concerns have already long been addressed: the natural matrix interpretation of the reals are the Hermitian matrices, while nonnegative (resp. positive) numbers correspond to those complex Hermitian matrices with all nonnegative (resp. positive) eigenvalues, the so-called *positive semidefinite* (resp. *positive definite*) matrices. The issue of noncommutativity, however, is of a more subtle nature, and we first introduce some notation before addressing it.

Fix a positive integer  $k$ , and let  $W = W(X, B_1, \dots, B_k)$  be a word in the letters  $X$  and  $B_1, \dots, B_k$ . The *reversal* of  $W$  is the word written in reverse order, and it is denoted by  $W^*$ . A word is *symmetric* if it is identical to its reversal (in other contexts, the name “palindromic” is also used). As we shall soon see (see Sections 2 and 3), formulating our generalization requires restriction to a special class of words. For the purposes of this work, an *interlaced* word  $W = W(X, B_1, \dots, B_k)$  in the *interlacing* letter  $X$  is a juxtaposition of powers of letters that alternate in  $X$ . More precisely, an interlaced word is an expression of the form,

$$(1) \quad W = B_{i_1}^{q_1} \prod_{j=1}^m X^{p_j} B_{i_{j+1}}^{q_{j+1}},$$

---

1991 *Mathematics Subject Classification.* 15A24, 15A57; 15A18, 15A90.

*Key words and phrases.* matrix equation, positive definite matrix, word, nonunique solutions.

The work of the second author is supported under a National Science Foundation Postdoctoral Fellowship.

in which the exponents  $p_j > 0$ ,  $q_j \geq 0$  are nonnegative integers and  $\{i_1, \dots, i_{m+1}\} \subseteq \{1, \dots, k\}$ . (Here, of course, we consider the zeroth power of a letter to be the empty word, the identity element of the monoid). For example, the word  $B_1 X B_3^7 X^2 B_2^3 X^5$  is interlaced, whereas the word  $X B_1 B_2 X B_2 B_1 X$  is not. The integer  $s = p_1 + \dots + p_m$  is called the *degree* of the interlaced word  $W$ .

The interlacing letter  $X$  is distinguished, and is to be viewed as an indeterminate  $n \times n$  positive semidefinite matrix, while the letters  $B_1, \dots, B_k$  correspond to fixed  $n \times n$  positive definite matrices. For convenience, the letters  $X$  and  $B_i$  will also represent the substituted matrices (the context will make the distinction clear). When  $k = 1$ , the set of interlaced words is simply the set of all words in two letters containing at least one  $X$ . For notational simplicity, when  $k$  is understood, we write  $W(X, B_i)$  in place of  $W(X, B_1, \dots, B_k)$ .

Returning to our motivating example, notice that there is a unique nonnegative solution to the equation  $bx^s = p$  for every pair of positive  $b$  and nonnegative  $p$ ; we would like to generalize this observation. Our introductory remarks prepare us to make the following definition.

**Definition 1.1.** A *symmetric word equation* is an equation,  $S(X, B_i) = P$ , in which  $S(X, B_i)$  is an interlaced symmetric word. If the  $B_i$  are positive definite and  $P$  is positive semidefinite, then any positive semidefinite matrix  $X$  for which the equation holds is called a *solution* to the symmetric word equation.

A symmetric word equation will be called *solvable* if there exists a solution for every positive definite  $n \times n$  matrices  $B_i$  and  $n \times n$  positive semidefinite  $P$ . Moreover, if each such  $B_i$  and  $P$  give rise to a unique solution, the equation will be called *uniquely solvable*.

**Theorem 1.2** (Hillar and Johnson). *Every symmetric word equation is solvable. Moreover, if the parameters  $P$  and  $B_i$  are real, then there is a real solution.*

Theorem 1.2 first appeared in [9, Theorem 7.1]. Its proof was accomplished using fixed point methods. The authors left open the problem of uniqueness.

**Conjecture 1.3.** *Every symmetric word equation is uniquely solvable.*

Uniquely solvable equations are ubiquitous (see Section 5, where we produce a large family of them). Recently, Lawson and Lim [14] have verified the conjecture in the case that the degree of  $S(X, B_i)$  is not greater than five. Their approach utilizes the Riemannian metric on the set of positive definite matrices and Banach's fixed point theorem.

In this paper, we resolve Conjecture 1.3 negatively in the case  $n \geq 3$ .

**Theorem 1.4.** *There are symmetric word equations of degree 6 which have multiple real  $3 \times 3$  positive definite solutions.*

Conjecture 1.3 remains open in the case of  $2 \times 2$  matrices.

Theorem 1.4 shows that the result of Lawson and Lim is optimal. Although uniqueness fails in general, our approach allows us to verify that these equations are still well-behaved in the following sense.

**Theorem 1.5.** *Fix an interlaced symmetric word  $S$  and real positive definite matrices  $B_1, \dots, B_k$  and  $P$ . There is a bounded open subset  $U$  of real positive definite*

matrices such that all real solutions  $X$  of the equation  $f(X) = S(X, B_i) = P$  lie in  $U$ . Moreover, identifying the real symmetric matrices with  $\mathbb{R}^m$  we have that

$$\deg(f, U, P) = 1.$$

Here,  $\deg(f, U, P)$  is the Brouwer degree of  $f$  at  $P$  with respect to  $U$ ; in a vague sense, it gives a topological measure of the number of solutions to equations  $f(X) = P$ . See Example 2.5 for a (non-interlaced) symmetric word equation with an unbounded set of solutions. Theorem 1.5 implies a special case of Theorem 1.2, giving a second proof of existence in the real case.

**Corollary 1.6.** *Every symmetric word equation in real positive definite letters has a real positive definite solution.*

*Proof.* The result follows from Theorem 1.5 and Lemma 7.1.  $\square$

**Corollary 1.7.** *For almost every real positive semidefinite matrix  $P$ , the symmetric word equation*

$$S(X, B_i) = P$$

*has an odd (and thus finite) number of real solutions  $X$ .*

*Proof.* By Theorems 1.5 and 7.3, at any regular value  $P$  of the map  $X \mapsto S(X, B_i)$  the equation  $S(X, B_i) = P$  has an odd number of solutions  $X$ . By Sard's theorem, the set of regular values is a set of full measure, completing the proof.  $\square$

The proof of Theorem 1.5 is the content of Sections 8, 9 and 10. The arguments in the proof often employ the reductions found in Section 6. Some consequences of Theorem 1.5 are explored in Section 11, including a proof of Theorem 1.4. In Sections 2 and 3 we explain why we restrict our attention to interlaced symmetric words, and Sections 4 and 5 are devoted to applications and a special class of uniquely solvable words, respectively. In Section 7 we review the theory of Brouwer degree.

## 2. A COLLECTION OF EXAMPLES

The simplest instance of a symmetric word equation arises in the following example ([13, p. 413] and [13, p. 433]); it is the most straightforward generalization of the scalar case.

**Example 2.1.** *Let  $P$  be any positive semidefinite matrix and let  $S(X)$  be the word  $X^m$ , for a positive integer  $m$ . Then, there is a unique positive semidefinite solution to the equation  $S(X) = P$ . In fact, writing  $P = UDU^*$  for a unitary matrix  $U$  and a nonnegative diagonal matrix  $D$ , we have  $X = UD^{1/m}U^*$ .  $\square$*

As another example, we examine a special case of the *algebraic Riccati equation*, which is encountered frequently in control theory [19].

**Example 2.2.** *The equation,  $XBX = P$ , has a unique positive semidefinite solution given positive definite  $B$  and positive semidefinite  $P$ . Moreover, it is given by*

$$X = B^{-1/2}(B^{1/2}PB^{1/2})^{1/2}B^{-1/2}.$$

*This fact can be deduced from the proof of Proposition 5.2, in which a large class of word equations are shown to be uniquely solvable. When  $P$  is invertible, this solution can also be expressed as*

$$X = P^{1/2}(P^{-1/2}B^{-1}P^{-1/2})^{1/2}P^{1/2},$$

even though this expression appears quite different.  $\square$

As promised, we will now explain why it makes sense to restrict our attention to interlaced symmetric words. A first obstacle in generalizing the scalar case is that most words do not evaluate to positive semidefinite matrices upon substitution. One simple example is the word  $XB$ , which does not even have to be Hermitian when  $X$  and  $B$  are positive definite. Similarly, the unique matrix solution  $X$  of the equation  $XB = P$  is not in general positive semidefinite; and examples are easily generated. It turns out that the right class of words to consider are the symmetric ones, and this is evidenced by the following discussion.

Recall that two  $n \times n$  matrices  $X$  and  $Y$  are said to be *congruent* if there is an invertible  $n \times n$  matrix  $Z$  such that  $Y = Z^*XZ$  (here,  $C^*$  denotes the *conjugate transpose* of a complex matrix  $C$ ); and that congruence on Hermitian matrices preserves inertia (the ordered triple consisting of the number of positive, negative, and zero eigenvalues) and, thus, positive definiteness [12, p. 223]. A symmetric word evaluated at positive definite matrices is inductively congruent to the “center,” positive definite matrix. We conclude that

**Lemma 2.3.** *A symmetric word evaluated at positive definite matrices is positive definite.*

A more careful examination (or a simple continuity argument) also proves the following.

**Lemma 2.4.** *A symmetric word evaluated at positive semidefinite matrices is positive semidefinite.*

Conversely, it may be shown that symmetric words are the only words that are positive definite for all positive definite substitutions (see Section 3 for a proof). In light of these facts, restricting our consideration to symmetric words seems appropriate.

Next, we discuss the difficulties that arise when considering non-interlaced symmetric words. As the following examples demonstrate, both uniqueness and existence may fail even when  $k = n = 2$  and  $s = 3$ .

**Example 2.5.** *Let  $S(X, B_1, B_2) = XB_1B_2XB_2B_1X$  and set*

$$B_1 = \begin{bmatrix} 3 & -1 \\ -1 & 1 \end{bmatrix}, \quad B_2 = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}, \quad \text{and} \quad P = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

*Then, as is easily verified, the equation  $S(X, B_1, B_2) = P$  has symmetric solutions*

$$X = \begin{bmatrix} 0 & 0 \\ 0 & x \end{bmatrix} \quad \text{and} \quad X = \begin{bmatrix} x/5 & -x \\ -x & 5x \end{bmatrix},$$

*in which  $x$  is an arbitrary real number. In particular, there are infinitely many positive semidefinite solutions (in two distinct solution classes). Notice also that the kernel of a solution  $X$  and that of  $P$  can be different. For interlaced words, this situation cannot occur (see Lemma 6.1).  $\square$*

**Example 2.6.** *Let  $S$  and  $B_1, B_2$  be as in the previous example, but instead set*

$$P = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

Then, there are no positive semidefinite solutions to  $S(X, B_1, B_2) = P$ . To verify this, suppose that

$$X = \begin{bmatrix} e & f \\ g & h \end{bmatrix}$$

is a complex solution to  $S(X, B_1, B_2) = P$ . Computing the ideal generated by the 4 consequent polynomial equations (using Maple or Macaulay 2 to find the reduced Gröbner basis), we find that it is the entire ring  $\mathbb{C}[e, f, g, h]$ . In particular, there are no matrix solutions over  $\mathbb{C}$  to the given equation, much less positive semidefinite ones.  $\square$

### 3. RELATIONS BETWEEN POSITIVE DEFINITE WORDS

In this section, we explain why we restrict our attention to symmetric words. Specifically, we prove that a word  $W(A, B)$  in two letters  $A$  and  $B$  is positive definite for all positive definite substitutions if and only if the word is symmetric.

We begin by illustrating some of the subtlety of the problem. Let  $B$  and  $P$  be positive definite matrices. In Example (2.2) we saw that

$$P^{1/2} \left( P^{-1/2} B^{-1} P^{-1/2} \right)^{1/2} P^{1/2} = B^{-1/2} \left( B^{1/2} P B^{1/2} \right)^{1/2} B^{-1/2},$$

even though both expressions are quite different. In fact, both sides of the above equality are the unique solution  $X$  to the symmetric word equation,

$$S(X, B) = X B X = P.$$

Fortunately, such behavior does not occur with words, as the following discussion illustrates.

Let  $\mathcal{W}$  be the set of words in two letters  $A$  and  $B$ , and fix  $a, b$  to be two  $n \times n$  complex matrices. Consider the *evaluation homomorphism*  $Eval_{a,b} : \mathcal{W} \rightarrow \mathbb{M}_n(\mathbb{C})$  which sends a word  $W(A, B)$  to the matrix  $W(a, b)$  produced by substituting the matrices  $a$  and  $b$  for the letters  $A$  and  $B$ , respectively. By convention, the empty word is sent to the identity matrix by this map. We describe a pair of positive definite  $a$  and  $b$  for which this function is injective.

**Lemma 3.1.** *The map  $Eval_{a,b}$  is injective when*

$$a = \begin{bmatrix} 3 & 1 \\ 1 & 1 \end{bmatrix}, \quad b = \begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix}.$$

*Proof.* Let  $a, b$  be the matrices in the statement of the lemma, and let  $W_1$  and  $W_2$  be two words for which  $W_1(a, b) = W_2(a, b)$ ; we must show that  $W_1$  and  $W_2$  are the same word. If either  $W_1$  or  $W_2$  is the empty word, then the claim is clear (take a determinant). Furthermore, since  $a$  and  $b$  are invertible, we may suppose that  $W_1 = AU$  and  $W_2 = BV$  for some words  $U$  and  $V$ .

Let  $x$  and  $y$  be indeterminates. Given a word  $W$ , we set

$$\begin{bmatrix} W^x x + W^y y \\ W_x x + W_y y \end{bmatrix} = W(a, b) \begin{bmatrix} x \\ y \end{bmatrix},$$

for natural numbers  $W^x, W^y, W_x, W_y$ . Notice that by our choice of  $a$  and  $b$ , we cannot have both  $W^x$  and  $W^y$  equal to zero. A direct computation shows that  $(AU)^x - (AU)_x = 2U^x$  and that  $(BV)^x - (BV)_x = -2V_x$ . By assumption, these two numbers are equal so that  $U^x + V_x = 0$ . Since these two quantities are nonnegative

integers, it follows that  $U^x = V_x = 0$ . Similarly, the equality  $(AU)^y - (AU)_y = (BV)^y - (BV)_y$  implies that  $U^y = V_y = 0$ . This contradiction finishes the proof.  $\square$

We now list some corollaries of this lemma.

**Corollary 3.2.** *The following are equivalent for a word  $W$ .*

- (1)  $W$  is positive definite for all substitutions of positive definite  $A$  and  $B$
- (2)  $W$  is Hermitian for all substitutions of positive definite  $A$  and  $B$
- (3)  $W$  is Hermitian for all  $2 \times 2$  substitutions of positive definite  $A$  and  $B$
- (4)  $W$  is symmetric (“palindromic”)

*In particular, if a word is Hermitian for all  $2 \times 2$  substitutions of positive definite  $A$  and  $B$ , then the word is necessarily positive definite for all such substitutions.*

*Proof.* (1)  $\Rightarrow$  (2)  $\Rightarrow$  (3) is clear. If  $W(A, B)$  is always Hermitian for  $2 \times 2$  positive definite  $A$  and  $B$ , then  $W(A, B)^* = W(A, B)$  for all such  $A$  and  $B$ . But then Lemma 3.2 says that  $W^*$  and  $W$  must be identical as words. It follows that  $W$  is symmetric. This proves (3)  $\Rightarrow$  (4). Finally, if  $W$  is symmetric, Lemma 2.3 says that  $W$  will always be positive definite for any positive definite  $A$  and  $B$ . This completes the proof.  $\square$

#### 4. APPLICATIONS

Symmetric word equations were first encountered while studying a trace conjecture [7] involving words in two letters  $A$  and  $B$  (see also [8, 11]).

**Conjecture 4.1.** *A word in two letters  $A$  and  $B$  has positive trace for every pair of real positive definite  $A$  and  $B$  if and only if the word is symmetric or a product (juxtaposition) of 2 symmetric words.*

It turns out that for each solvable symmetric word equation, one can identify an infinite class of words that admit real positive definite matrices  $A$  and  $B$  giving those words a negative trace. The following is a brief description of this application. Consider the word  $W = BAB AAB$ , which is not symmetric nor a product of two symmetric words. In light of Conjecture 4.1, we would like to verify that there exist real positive definite matrices  $A$  and  $B$  giving  $W$  a negative trace. This is surprisingly difficult, as the methods in [7] show. Resulting  $A$  and  $B$  that exhibit a negative trace are, for example,

$$A_1 = \begin{bmatrix} 1 & 20 & 210 \\ 20 & 402 & 4240 \\ 210 & 4240 & 44903 \end{bmatrix} \quad \text{and} \quad B_1 = \begin{bmatrix} 36501 & -3820 & 190 \\ -3820 & 401 & -20 \\ 190 & -20 & 1 \end{bmatrix}.$$

Consider now the following extension. Let  $T$  be the word given by  $T = S_1 S_2$ , in which  $S_1$  and  $S_2$  are symmetric words in the letters  $A$  and  $B$ . If the simultaneous word equations

$$\begin{aligned} S_1(A, B) &= B_1, \\ S_2(A, B) &= A_1 \end{aligned}$$

may be solved for positive definite  $A$  and  $B$  given positive definite  $A_1$  and  $B_1$ , then the word  $TTT^*$  can have negative trace. Specializing to the case that  $S_2$  is the word  $A$ , we have the following.

**Corollary 4.2.** *Let  $S = S(A, B)$  be any symmetric word with at least one  $B$ . Then the word  $SASAAS$  admits real positive definite matrices  $A$  and  $B$  giving it negative trace.*

*Proof.* The matrix  $B_1A_1B_1A_1A_1B_1$  has negative trace. Using Corollary 1.6, the equation  $S(A_1, X) = B_1$  has a real positive definite solution  $X = B_2$ . The two matrices  $B = B_2$  and  $A = A_1$  are then the desired witnesses.  $\square$

We should remark that Conjecture 4.1, while interesting in its own right, arises from a long-standing problem in statistical physics. In [1], while studying partition functions of quantum mechanical systems, a conjecture was made regarding a positivity property of traces of matrices. If this property holds, explicit error bounds in a sequence of Padé approximants follow. Recently, in [15], and as previously communicated to us [7], the conjecture of [1] was reformulated as a question about the traces of certain sums of words in two positive definite matrices.

**Conjecture 4.3** (Bessis-Moussa-Villani). *The polynomial  $p(t) = \text{Tr}[(A + tB)^m]$  has all positive coefficients whenever  $A$  and  $B$  are  $n \times n$  positive definite matrices.*

The coefficient of  $t^k$  in  $p(t)$  is the trace of  $H_{m,k}(A, B)$ , the sum of all words of length  $m$  in  $A$  and  $B$ , in which  $k$   $B$ 's appear. Since its introduction in [1], many partial results and substantial computational experimentation have been given [3, 4, 7, 10, 18], all in favor of the conjecture's validity. However, despite much work, very little is known about the problem, and it has remained unresolved except in very special cases. Until recently, even the case  $m = 6$  and  $n = 3$  was unknown. In this case, all coefficients, except  $\text{Tr}[H_{6,3}(A, B)]$  were known to be positive [7]. The remaining coefficient  $\text{Tr}[H_{6,3}(A, B)]$  can be shown to be positive, but the proof requires notably different methods [10]. The difficulty is that some summands of  $H_{6,3}(A, B)$  can have negative trace, precisely the types of words such as  $BABAAB$  considered above.

A recent advance on the conjecture has been the derivation of a pair of equations satisfied by  $A$  and  $B$  with Euclidean norm 1 that minimizes a coefficient  $\text{Tr}[H_{m,k}(A, B)]$ :

$$\begin{cases} AH_{m-1,k}(A, B) & = A^2\text{Tr}[AH_{m-1,k}(A, B)] \\ BH_{m-1,k-1}(A, B) & = B^2\text{Tr}[BH_{m-1,k-1}(A, B)]. \end{cases}$$

Using these equations, one can show that if Conjecture 4.3 holds for some  $m_0$ , then it holds for all  $m < m_0$ .

## 5. A CLASS OF UNIQUELY SOLVABLE EQUATIONS

In this section, we describe a class of words that are uniquely solvable with solutions that can be explicitly constructed. These words generalize those found in Examples 2.1 and 2.2.

**Definition 5.1.** A symmetric word is called *totally symmetric* if it can be expressed as a composition of maps of the form

- (1)  $\pi_{m, B_i}(W) = (WB_i)^m W$ ,  $m$  a positive integer
- (2)  $\varphi_m(W) = W^m$ ,  $m$  a positive integer
- (3)  $\mathcal{C}_{B_i}(W) = B_iWB_i$

applied to the letter  $X$ .

For example, the word  $W = B_1X^2B_2X^2B_2X^2B_1$  may be expressed as the composition,  $\mathcal{C}_{B_1} \circ \pi_{2, B_2} \circ \varphi_2(X)$ . The utility of this definition becomes clear from the following proposition.

**Proposition 5.2.** *For every totally symmetric word  $S(X, B_i)$  and every positive definite  $B_i$  and positive semidefinite  $P$ , the equation  $S(X, B_i) = P$  has a unique positive semidefinite solution  $X$ .*

*Proof.* We induct on the number of compositions involved in the word  $S$ ; the base case  $S = X$  being trivial. If  $S = \varphi_m(W)$  for some word  $W$ , then  $W = P^{1/m}$  is a smaller totally symmetric word equation and any solution  $X$  to  $S(X, B_i) = P$  satisfies it. A similar statement holds when  $S = \mathcal{C}_{B_i}(W)$  (using Lemma 2.4), leaving us to deal with  $\pi_{m, B_i}$ .

Without loss of generality, we prove the result for the equation  $(XB)^m X = P$ . Assume that  $B$  and  $P$  are given and that  $X$  is a solution to  $(XB)^m X = P$ . Set  $Y = B^{1/2} X B^{1/2}$ , so that  $X = B^{-1/2} Y B^{-1/2}$ . Then,

$$P = (B^{-1/2} Y B^{1/2})^m B^{-1/2} Y B^{-1/2} = B^{-1/2} Y^{m+1} B^{-1/2}.$$

Therefore,  $Y^{m+1} = B^{1/2} P B^{1/2}$ , from which it follows that  $Y$  is uniquely determined as  $(B^{1/2} P B^{1/2})^{1/(m+1)}$ . Hence,  $X$  must be the positive semidefinite matrix  $B^{-1/2} (B^{1/2} P B^{1/2})^{1/(m+1)} B^{-1/2}$ . Finally, substituting this  $X$  into the original equation does verify that it is a solution. This completes the proof.  $\square$

The shortest symmetric word equation without a known (closed-form) solution, as above, is  $XBX^3BX = P$  (although it is uniquely solvable [14]). An exploration of which equations give rise to such explicit solutions is the focus of future work.

## 6. REDUCTIONS

The purpose of this section is to make some reductions that simplify the problem. Given the nature of Theorem 1.2 and Conjecture 1.3, we begin by noticing that we may assume our interlaced symmetric words are of the following form:

$$(2) \quad S = X^{p_1} B_1 X^{p_2} B_2 \cdots B_2 X^{p_2} B_1 X^{p_1},$$

in which the exponents  $p_j$  are positive. This simplification is accomplished by observing first, that powers of positive definite matrices are positive definite; and second, that congruences of positive semidefinite  $P$  are positive semidefinite (Lemma 2.4).

We next establish that it suffices to verify our claims when  $P$  is invertible. We begin with a useful lemma.

**Lemma 6.1.** *Let  $p_1, \dots, p_k > 0$  and let  $B_1, \dots, B_{k-1}$  be positive definite matrices. Then, for any positive semidefinite matrix  $X$ , we have*

$$\ker X = \ker X^{p_k} B_{k-1} \cdots B_2 X^{p_2} B_1 X^{p_1}.$$

*Proof.* Set  $X = UDU^*$  for a unitary matrix  $U$  and  $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ , in which  $\lambda_1 \geq \dots \geq \lambda_n \geq 0$ . Let  $Y = X^{p_k} B_{k-1} \cdots B_2 X^{p_2} B_1 X^{p_1}$ , and notice that  $\ker U^* X U = \ker U^* Y U$  if and only if  $\ker X = \ker Y$ . Thus, it suffices to argue that

$$\ker D = \ker D^{p_k} B_{k-1} \cdots B_2 D^{p_2} B_1 D^{p_1},$$

whenever the  $B_i$  are positive definite matrices.

Let  $m$  be the largest integer such that  $\lambda_m \neq 0$ , and for each  $i$ , let  $\tilde{B}_i$  denote the  $m \times m$  leading principal submatrix of  $B_i$ , which will be positive definite (see, for

instance, [12, p. 472]). Additionally, set  $\tilde{D} = \text{diag}(\lambda_1, \dots, \lambda_m)$ . A straightforward block matrix multiplication then gives us that

$$(3) \quad D^{p_k} B_{k-1} \cdots B_2 D^{p_2} B_1 D^{p_1} = \begin{bmatrix} \tilde{D}^{p_k} \tilde{B}_{k-1} \cdots \tilde{B}_2 \tilde{D}^{p_2} \tilde{B}_1 \tilde{D}^{p_1} & 0 \\ 0 & 0 \end{bmatrix}.$$

Since the leading principal  $m \times m$  matrix in this direct sum is invertible, the claim follows.  $\square$

Using this lemma, we can prove the following reduction.

**Theorem 6.2.** *If a symmetric word equation has a solution for every positive definite  $B_i$  and  $P$ , then the symmetric word equation has a solution for every positive definite  $B_i$  and positive semidefinite  $P$ .*

*Proof.* Performing a uniform unitary similarity, we may prove the theorem with the supposition that  $P$  is of the form,

$$\begin{bmatrix} \tilde{P} & 0 \\ 0 & 0 \end{bmatrix},$$

for a positive diagonal matrix  $\tilde{P}$  of rank  $m$ . Lemma 6.1 implies that any positive semidefinite solution  $X$  to the symmetric word equation  $S(X, B_i) = P$  has the same block form as  $P$ . As in the lemma, let  $\tilde{B}_i$  denote the  $m \times m$  leading principal (positive definite) submatrix of each  $B_i$ .

From these observations, it follows that positive semidefinite solutions  $X$  to the equation  $S(X, B_i) = P$  correspond in a one-to-one manner with positive definite solutions  $\tilde{X}$  to the equation  $S(\tilde{X}, \tilde{B}_i) = \tilde{P}$ . This completes the proof.  $\square$

The proof above also shows that the question of uniqueness found in Conjecture 1.3 may also be simplified.

**Theorem 6.3.** *If a symmetric word equation has a unique solution for all positive definite matrices  $B_i$  and  $P$ , then the symmetric word equation has a unique solution for all positive definite  $B_i$  and each positive semidefinite  $P$ .*

We close this section with an interesting interpretation of unique solvability.

**Proposition 6.4.** *Fix positive definite matrices  $B_i$  in the unit ball and an interlaced symmetric word  $S(X, B_i)$  whose equations are uniquely solvable. Then, the mapping  $X \mapsto S(X, B_i)$  from the set of positive semidefinite matrices in the (closed) unit ball to its image is a homeomorphism.*

*Proof.* The assumptions imply that our map is bijective. Since the set of positive semidefinite matrices in the unit ball is compact, it follows that its inverse is also continuous.  $\square$

## 7. BROUWER MAPPING DEGREE

In this section, we give a brief overview of degree theory and some of its main implications. The bulk of this discussion is material taken from [5, 16, 21]. First we introduce some notation. Let  $U$  be a bounded open subset of  $\mathbb{R}^m$ . We denote the set of  $r$ -times differentiable functions from  $U$  (resp.  $\bar{U}$ ) to  $\mathbb{R}^m$  by  $C^r(U, \mathbb{R}^m)$  (resp.  $C^r(\bar{U}, \mathbb{R}^m)$ ) (when  $r = 0$ ,  $C^r(U, \mathbb{R}^m)$  is the set of continuous functions). The

identity function  $\mathbb{1}$  satisfies  $\mathbb{1}(\mathbf{x}) = \mathbf{x}$ . If  $f \in C^1(U, \mathbb{R}^m)$ , then the *Jacobi matrix* of  $f$  at a point  $\mathbf{x} \in U$  is

$$J_f(\mathbf{x}) = \left[ \frac{\partial f_j}{\partial x_i}(\mathbf{x}) \right]_{1 \leq i, j \leq m}$$

and the *Jacobi determinant* (or simply *Jacobian*) of  $f$  at  $\mathbf{x}$  is

$$\det J_f(\mathbf{x}).$$

The set of *regular values* of  $f$  is

$$\text{RV}(f) = \{\mathbf{y} \in \mathbb{R}^m : \forall \mathbf{x} \in f^{-1}(\mathbf{y}), J_f(\mathbf{x}) \neq 0\}$$

and for  $\mathbf{y} \in \mathbb{R}^m$ , we set

$$D_{\mathbf{y}}^r(\overline{U}, \mathbb{R}^m) = \{f \in C^r(\overline{U}, \mathbb{R}^m) : \mathbf{y} \notin f(\partial U)\}.$$

A function  $\text{deg} : D_{\mathbf{y}}^0(\overline{U}, \mathbb{R}^m) \rightarrow \mathbb{R}$  which assigns to each  $f \in D_{\mathbf{y}}^0(\overline{U}, \mathbb{R}^m)$  and  $\mathbf{y} \in \mathbb{R}^m$  a real number  $\text{deg}(f, U, \mathbf{y})$  will be called a *degree* if it satisfies the following conditions:

- (1)  $\text{deg}(f, U, \mathbf{y}) = \text{deg}(f - \mathbf{y}, U, 0)$  (*translation invariance*).
- (2)  $\text{deg}(\mathbb{1}, U, \mathbf{y}) = 1$  if  $\mathbf{y} \in U$  (*normalization*).
- (3) If  $U_1$  and  $U_2$  are open, disjoint subsets of  $U$  such that  $\mathbf{y} \notin f(\overline{U} \setminus (U_1 \cup U_2))$ , then  $\text{deg}(f, U, \mathbf{y}) = \text{deg}(f, U_1, \mathbf{y}) + \text{deg}(f, U_2, \mathbf{y})$  (*additivity*).
- (4) If  $H(t) = tf + (1-t)g \in D_{\mathbf{y}}^0(\overline{U}, \mathbb{R}^m)$  for all  $t \in [0, 1]$ , then  $\text{deg}(f, U, \mathbf{y}) = \text{deg}(g, U, \mathbf{y})$  (*homotopy invariance*).

Motivationally, one should think of a degree map as somehow “counting” the number of solutions to  $f(\mathbf{x}) = \mathbf{y}$ . Condition (1) reflects that the solutions to  $f(\mathbf{x}) = \mathbf{y}$  are the same as those of  $f(\mathbf{x}) - \mathbf{y} = 0$ , and since any multiple of a degree will satisfy (1) and (3), condition (2) is a normalization. Additionally, (3) is natural since it requires  $\text{deg}$  to be additive with respect to components. The following lemma gives a method to show the existence of solutions to  $f(\mathbf{x}) = \mathbf{y}$  by calculating a degree.

**Lemma 7.1.** *Suppose that  $f \in D_{\mathbf{y}}^0(\overline{U}, \mathbb{R}^m)$ . If a degree satisfies  $\text{deg}(f, U, \mathbf{y}) \neq 0$ , then  $\mathbf{y} \in f(U)$ .*

*Proof.* Using property (3) above with  $U_1 = U$  and  $U_2 = \emptyset$ , we must have that  $\text{deg}(f, \emptyset, \mathbf{y}) = 0$ . Again using (3) with  $U_1 = U_2 = \emptyset$ , it follows that if  $\mathbf{y} \notin f(\overline{U})$  then  $\text{deg}(f, U, \mathbf{y}) = 0$ . The contrapositive is now what we want.  $\square$

Of course, we need a theorem guaranteeing that a degree even exists.

**Theorem 7.2.** *There is a unique degree  $\text{deg}$ . Moreover,  $\text{deg}(\cdot, U, \mathbf{y}) : D_{\mathbf{y}}^0(\overline{U}, \mathbb{R}^m) \rightarrow \mathbb{Z}$ .*

When functions are differentiable, the degree can be calculated explicitly in terms of Jacobians at solutions to the equation  $f(\mathbf{x}) = \mathbf{y}$ .

**Theorem 7.3.** *Suppose that  $f \in D_{\mathbf{y}}^1(\overline{U}, \mathbb{R}^m)$  and  $\mathbf{y} \in \text{RV}$ . Then the degree of  $f$  at  $\mathbf{y}$  with respect to  $U$  is given by*

$$\text{deg}(f, U, \mathbf{y}) = \sum_{\mathbf{x} \in f^{-1}(\mathbf{y})} \text{sgn} \det J_f(\mathbf{x}),$$

where this sum is finite and we adopt the convention that  $\sum_{\mathbf{x} \in \emptyset} = 0$ .

The final property of Brouwer degree that we will need is a stronger form of homotopy invariance than that provided by Property (4). We say that a function  $H : \overline{U} \times [0, 1] \rightarrow \mathbb{R}^m$  is a  $C^0$  homotopy between  $f, g \in C^r(\overline{U}, \mathbb{R}^m)$  if  $H$  is continuous on  $\overline{U} \times [0, 1]$  and if  $H(x, 0) = f(x)$  and  $H(x, 1) = g(x)$  for all  $x \in \overline{U}$ .

**Theorem 7.4.** *Suppose  $H$  is a  $C^0$  homotopy between  $f, g \in D_{\mathbf{y}}^0(\overline{U}, \mathbb{R}^m)$ . Set  $h_t(x) = H(x, t)$  and suppose that for each  $t \in [0, 1]$ ,  $h_t \in D_{\mathbf{y}}^0(\overline{U}, \mathbb{R}^m)$ . Then  $\deg(f, U, \mathbf{y}) = \deg(g, U, \mathbf{y})$ .*

## 8. ESTIMATES OF SOLUTIONS

This section is devoted to estimating the norms of positive definite solutions of symmetric word equations. In particular, we show that the set of positive definite solutions to a fixed symmetric word equation  $S(X, B_i) = P$  is bounded. Our estimate is the first step in the proof of Theorem 1.5. In what follows, we will be using the spectral norm [12, p. 295] on the set of  $n \times n$  matrices, so that for positive semidefinite  $A$ , the norm of  $A$  is just the largest eigenvalue of  $A$ .

**Lemma 8.1.** *Fix an interlaced symmetric word  $S(X, B_i)$  and a number  $\alpha \geq 1$ . Then there exists a constant  $C = C_{S, \alpha}$  depending only on  $S$  and  $\alpha$  such that for all positive definite matrices  $B_i$  with  $\|B_i\| \leq 1$  and  $\|B_i^{-1}\| \leq \alpha$  and all positive semidefinite matrices  $P$  with  $\|P\| \leq 1$  we have the estimate*

$$(4) \quad \|X\| \leq C$$

for any solution  $X$  of the word equation  $S(X, B_i) = P$ .

*Proof.* We proceed by way of contradiction. If the statement is false, then for each positive integer  $j$  there exist positive semidefinite matrices  $X_j, P_j$  and positive definite matrices  $B_{i,j}$  such that  $S(X_j, B_{i,j}) = P_j$ , where  $\|B_{i,j}\| \leq 1$ ,  $\|B_{i,j}^{-1}\| \leq \alpha$ ,  $\|P_j\| \leq 1$ , and  $\|X_j\| \geq j$ . By taking a subsequence, if necessary, we may assume that there are positive semi-definite matrices  $B_i, P$  and  $X$  such that  $B_{i,j} \rightarrow B_i$ ,  $P_j \rightarrow P$ , and  $\|X_j\|^{-1} X_j \rightarrow X$  as  $j \rightarrow \infty$ . It is clear that

$$(5) \quad \|X\| = 1.$$

Since  $\|B_{i,j}^{-1}\|$  is bounded uniformly in  $j$ , we have that  $B_i$  is positive definite. Let  $s$  be the degree of  $S$ . Keeping in mind that  $\|X_j\| \geq j$  and letting  $j \rightarrow \infty$  in the equation

$$S(\|X_j\|^{-1} X_j, B_{i,j}) = \|X_j\|^{-s} P_j$$

yields

$$S(X, B_i) = 0.$$

Finally, an application of Lemma 6.1 gives  $X = 0$ , which contradicts (5) and finishes the proof.  $\square$

Lemma 8.1 allows us to estimate  $\|X\|$  in terms of the norms of the  $B_i$  and the norm of the word  $S(X, B_i)$ .

**Proposition 8.2.** *Fix an interlaced symmetric word  $S(X, B_1, \dots, B_k)$  of the form (2) with degree  $s$  and a number  $\alpha \geq 1$ . There exists a constant  $C = C_{S, \alpha}$  depending only on  $S$  and  $\alpha$  such that for all positive definite matrices  $B_i$  with  $\|B_i\| \|B_i^{-1}\| \leq \alpha$  and any positive semidefinite  $X$  we have*

$$(6) \quad \|X\| \leq C \|B_1\|^{-\frac{2}{s}} \cdots \|B_k\|^{-\frac{2}{s}} \|S(X, B_i)\|^{\frac{1}{s}}.$$

*Proof.* Let  $C = C_{S,\alpha}$  be the constant in Lemma 8.1. By Lemma 6.1, if  $S(X, B_i) = 0$ , then  $X = 0$ , and the bound is trivial. Otherwise, set  $\tilde{B}_i = \|B_i\|^{-1}B_i$ ,  $P = S(X, B_i)$ ,  $\tilde{P} = \|P\|^{-1}P$ , and  $\tilde{X} = \|B_1\|^{\frac{2}{s}} \cdots \|B_k\|^{\frac{2}{s}} \|P\|^{-\frac{1}{s}} X$ . Noticing that  $\|\tilde{B}_i\| = \|B_i\| \|B_i^{-1}\| \leq \alpha$  and also that  $S(\tilde{X}, \tilde{B}_i) = \tilde{P}$ , we may apply Lemma 8.1 to get that

$$(7) \quad \|\tilde{X}\| \leq C.$$

Substituting  $\tilde{X} = \|B_1\|^{\frac{2}{s}} \cdots \|B_k\|^{\frac{2}{s}} \|P\|^{-\frac{1}{s}} X$  into (7) and rearranging produces (6).  $\square$

## 9. CALCULATION OF JACOBI MATRICES

From here on, we consider only real  $n \times n$  matrices. For the purposes of this section, we will identify  $\mathbb{M}_n = \mathbb{M}_n(\mathbb{R})$  with  $\mathbb{R}^d$ , where  $d = n^2$ , by means of the *vec* operator. If  $A = [a_{ij}] \in \mathbb{M}_n$  then  $\text{vec } A$  is the column vector obtained by stacking the columns of  $A$  below one another:

$$\text{vec } A = [a_{11} \cdots a_{n1} \ a_{12} \cdots \cdots a_{nn}]^T.$$

Recall that the *Kronecker product* of two  $n \times n$  matrices  $A$  and  $B$  is the matrix

$$A \otimes B = \begin{bmatrix} a_{11}B & \cdots & a_{1n}B \\ \vdots & \ddots & \vdots \\ a_{n1}B & \cdots & a_{nn}B \end{bmatrix} \in \mathbb{M}_d.$$

The following lemma can be found in [17, page 30]. We reproduce it here for the reader's convenience.

**Lemma 9.1.** *If  $A, B, X \in \mathbb{M}_n$ , then*

$$\text{vec}(AXB) = (B^T \otimes A)\text{vec } X.$$

*Proof.* For a given matrix  $Q$ , let  $Q_k$  denote the  $k$ th column of  $Q$ . Let  $B = [b_{ij}]$ . Then

$$\begin{aligned} (AXB)_k &= AXB_k \\ &= A \left( \sum_{i=1}^n b_{ik} X_i \right) = [ \ b_{1k}A \ \cdots \ b_{nk}A \ ] \text{vec } X. \end{aligned}$$

Therefore,

$$\text{vec}(AXB) = \begin{bmatrix} b_{11}A & \cdots & b_{n1}A \\ \vdots & \ddots & \vdots \\ b_{1n}A & \cdots & b_{nn}A \end{bmatrix} \text{vec } X = (B^T \otimes A)\text{vec } X. \quad \square$$

Suppose that  $Y(X) \in \mathbb{M}_n$  is a function of the matrix variable  $X \in \mathbb{M}_n$ . Following [17], we define the *derivative*  $\frac{dY}{dX}$  of  $Y$  with respect to  $X$  to be the Jacobi matrix of  $\text{vec } Y$  with respect to  $\text{vec } X$ . That is, if  $[y_1, \dots, y_d]^T = \text{vec } Y$  and  $[x_1, \dots, x_d]^T = \text{vec } X$ , then

$$\frac{dY}{dX} = \left[ \frac{\partial y_i}{\partial x_j} \right].$$

Notice that it follows from Lemma 9.1 that

$$(8) \quad \frac{d(AXB)}{dX} = B^T \otimes A.$$

Using (8), we may derive a matrix calculus version of the product rule (see [17] for more on matrix calculus).

**Proposition 9.2.** *Let  $Y(X) \in \mathbb{M}_n$  and  $Z(X) \in \mathbb{M}_n$  be functions of the matrix variable  $X \in \mathbb{M}_n$ . Then*

$$(9) \quad \frac{d(YZ)}{dX} = (Z^T \otimes I) \frac{dY}{dX} + (I \otimes Y) \frac{dZ}{dX}.$$

Motivated by Theorem 7.3, we want to calculate the derivative  $\frac{dW}{dX}$  of a word  $W = W(X, B_1, \dots, B_k)$ . To accomplish this, we first introduce some notation. Let  $W$  have degree  $s \geq 1$ . Enumerate the occurrences of  $X$  in  $W(X, B_i)$  from left to right, and for each  $j \in \{1, \dots, s\}$  let  $W_j^L(X, B_i)$  be the portion of  $W(X, B_i)$  that appears to the left of the  $j$ th occurrence of  $X$ . For instance, if

$$W(X, B_1, B_2) = B_2^3 X B_1^2 B_2 X B_2 B_1 X^2 B_2 X,$$

then  $W_4^L(X, B_1, B_2) = B_2^3 X B_1^2 B_2 X B_2 B_1 X$ . We adopt the convention that  $W_1^L = I$  if  $X$  is the first letter of the word. In a similar way we define  $W_j^R(X, B_i)$  to be the portion of  $W(X, B_i)$  that appears to the right of the  $j$ th appearance of  $X$ . Notice that

$$W(X, B_i) = W_j^L(X, B_i) X W_j^R(X, B_i)$$

for any  $j = 1, \dots, s$ .

**Proposition 9.3.** *Let  $W = W(X, B_i)$  be a word of degree  $s$ , and  $B_i \in \mathbb{M}_n$ . Then*

$$(10) \quad \frac{dW}{dX} = \sum_{j=1}^s W_j^R(X, B_i)^T \otimes W_j^L(X, B_i).$$

*Proof.* We proceed by induction on the length of  $W$ . For the words  $X$  and  $BX$  ( $B = B_{k_1}, \dots, B_{k_N}$ ), (10) is a special case of (8). Now suppose that (10) holds for a fixed word  $W = W(X, B_i)$  of degree  $s$ . Pick  $B \in \{B_1, \dots, B_k\}$  and set

$$\overline{W}(X, B_i) = W(X, B_i)B.$$

Then (8) and (9) imply that

$$\begin{aligned} \frac{d\overline{W}}{dX} &= (B^T \otimes I) \frac{dW}{dX} \\ &= (B^T \otimes I) \sum_{j=1}^s W_j^R(X, B_i)^T \otimes W_j^L(X, B_i) \\ &= \sum_{j=1}^s (W_j^R(X, B_i)B)^T \otimes W_j^L(X, B_i) \\ &= \sum_{j=1}^s \overline{W}_j^R(X, B_i)^T \otimes \overline{W}_j^L(X, B_i), \end{aligned}$$

so formula (10) holds for  $\overline{W}$ .

Now set  $\widetilde{W}(X, B_i) = W(X, B_i)X$ . Appealing again to (8) and (9) we compute

$$\begin{aligned} \frac{d\widetilde{W}}{dX} &= (X^T \otimes I) \frac{dW}{dX} + (I \otimes W(X, B_i)) \\ &= (X^T \otimes I) \sum_{j=1}^s W_j^R(X, B_i)^T \otimes W_j^L(X, B_i) + (I \otimes \widetilde{W}_{s+1}^L) \\ &= \sum_{j=1}^{s+1} \widetilde{W}_j^R(X, B_i)^T \otimes \widetilde{W}_j^L(X, B_i), \end{aligned}$$

and so (10) holds for  $\widetilde{W}$ . This completes the induction and the proof.  $\square$

We next write down expression (10) for some explicit interlaced symmetric words. We begin with the most basic symmetric word.

**Example 9.4.** *For a positive integer  $s$ , the Jacobi matrix of  $\text{vec } X^s$  with respect to  $\text{vec } X$  is given by*

$$(11) \quad \frac{dX^s}{dX} = \sum_{j=1}^s (X^{s-j})^T \otimes X^{j-1}.$$

*In particular, since the Kronecker product of two positive (semi)definite matrices is also positive (semi)definite (see [13, p. 245], page 245),  $\frac{dX^s}{dX}$  is positive (semi)definite whenever  $X$  is positive (semi)definite.*

**Example 9.5.** *Consider the symmetric word  $S$  in two letters given by*

$$S(X, B) = XBX^2B^3X^2BX.$$

*If  $B$  is positive definite and  $X$  is symmetric, then*

$$\begin{aligned} \frac{dS}{dX} &= XBX^2B^3X^2B \otimes I + XBX^2B^3X \otimes XB + XBX^2B^3X \otimes XBX \\ &\quad + XBX \otimes XBX^2B^3 + XB \otimes XBX^2B^3X + I \otimes XBX^2B^3X^2B. \end{aligned}$$

## 10. THE BROUWER DEGREE OF SYMMETRIC WORD EQUATIONS

Our proof of Theorem 1.5 consists of two main steps. In the first, we calculate the degree of the simple map  $\varphi_s(X) = X^s$  and show that it is 1. And in the second, we create a homotopy from the function  $f(X) = S(X, B_i)$  to  $\varphi_s(X)$  and apply Theorem 7.4. Before initiating our proof, we need to identify the set of real positive definite matrices with an *open* set in Euclidean space. To this end, we identify the set  $\text{Sym}_n$  of real symmetric matrices with  $\mathbb{R}^m$ , in which  $m = \frac{1}{2}n(n+1)$ , by identifying a real symmetric matrix  $A = [a_{ij}]$  with the point

$$\mu(A) = (a_{11}, \dots, a_{n1}, a_{22}, \dots, a_{n2}, \dots, a_{nn}).$$

More precisely, if  $A \in \mathbb{M}_n$  then we define  $\mu(A) = (y_1, \dots, y_m)$ , where

$$y_{\frac{1}{2}(2n-j)(j-1)+i} = a_{ij}, \quad 1 \leq j \leq i \leq n.$$

It follows that  $\mu|_{\text{Sym}_n}$  is a linear isomorphism from  $\text{Sym}_n$  onto  $\mathbb{R}^m$ . We denote by  $\nu$  the inverse of  $\mu|_{\text{Sym}_n}$ . Let

$$\mathcal{O} = \{\mu(X) \mid X \text{ is positive definite}\}.$$

The set of positive definite matrices is therefore identified with the open subset  $\mathcal{O} \subset \mathbb{R}^m$ , and the set of positive semidefinite matrices is identified with the set  $\overline{\mathcal{O}}$ .

Define a function  $\mathcal{P}_s : \mathbb{R}^m \rightarrow \mathbb{R}^m$  by

$$\mathcal{P}_s = \mu \circ \varphi_s \circ \nu.$$

Since  $\varphi_s$  maps  $\text{Sym}_n$  into itself, it follows that  $\mathcal{P}_s(\mu(X)) = \mu(X^s)$  for every symmetric matrix  $X$ . We intend to show that  $\det J_{\mathcal{P}_s}(\mu(X)) > 0$  when  $X$  is positive definite. To accomplish this, we need the following lemma. For a matrix  $H$ , we denote the set of eigenvalues of  $H$  by  $\sigma(H)$ .

**Lemma 10.1.** *Let  $f : \mathbb{R}^N \rightarrow \mathbb{R}^N$  be a  $C^1$  map and  $V \subseteq \mathbb{R}^N$  be a linear subspace of  $\mathbb{R}^N$  such that  $f(V) \subseteq V$ . Let  $\pi : \mathbb{R}^M \rightarrow V$  be a linear isomorphism, and let  $\tilde{f} : \mathbb{R}^M \rightarrow \mathbb{R}^M$  be given by  $\tilde{f} = \pi^{-1} \circ f \circ \pi$ . Then if  $\mathbf{x} \in V$ , we have*

$$\sigma(J_{\tilde{f}}(\pi^{-1}(\mathbf{x}))) \subseteq \sigma(J_f(\mathbf{x})).$$

*In particular, if  $\mathbf{x} \in V$  and  $J_f(\mathbf{x})$  is positive definite, then  $J_{\tilde{f}}(\pi^{-1}(\mathbf{x}))$  has positive eigenvalues.*

*Proof.* Let  $\{e_1, \dots, e_N\}$  be the standard basis for  $\mathbb{R}^N$ . By choosing a linear change of variables  $u : \mathbb{R}^N \rightarrow \mathbb{R}^N$  such that  $u(V) = \text{span}\{e_1, \dots, e_M\}$  and considering the  $C^1$  map  $g = u \circ f \circ u^{-1}$ , we may reduce to the case that  $V = \text{span}\{e_1, \dots, e_M\}$ . We may likewise assume that  $\pi(z_1, \dots, z_M) = (z_1, \dots, z_M, 0, \dots, 0)$ .

Write  $f = (f_1, \dots, f_N)$ . If  $j \leq M < k$ , we have

$$f_k(\mathbf{x} + te_j) = 0 \text{ for all } t \in \mathbb{R}$$

since  $f(V) \subseteq V$ . Therefore,

$$(12) \quad \frac{\partial f_k}{\partial x_j}(\mathbf{x}) = 0 \text{ for all } j \leq M < k.$$

It follows that  $J_f(\mathbf{x})$  has the block form

$$J_f(\mathbf{x}) = \begin{bmatrix} J_0 & * \\ 0 & * \end{bmatrix},$$

in which  $J_0$  is the  $M \times M$  leading principle submatrix of  $J_f(\mathbf{x})$ . In particular, this implies that  $\sigma(J_0) \subset \sigma(J_f(\mathbf{x}))$ . Since  $\pi$  has such a simple form, it is straightforward to verify that

$$\tilde{f}(x_1, \dots, x_m) = (f_1(x_1, \dots, x_m, 0, \dots, 0), \dots, f_m(x_1, \dots, x_m, 0, \dots, 0)),$$

from which it follows that  $J_{\tilde{f}}(\pi^{-1}(\mathbf{x})) = J_0$ . Thus

$$\sigma(J_{\tilde{f}}(\pi^{-1}(\mathbf{x}))) = \sigma(J_0) \subset \sigma(J_f(\mathbf{x})).$$

□

**Lemma 10.2.** *At any positive definite matrix  $X$ , the Jacobi matrix  $J_{\mathcal{P}_s}(\mu(X))$  of the map  $\mathcal{P}_s$  has positive eigenvalues. In particular,  $\det J_{\mathcal{P}_s}(\mu(X)) > 0$ .*

*Proof.* Let  $V = \{\text{vec } X \mid X \in \text{Sym}_n\}$  be the linear subspace of  $\mathbb{R}^d$  identified with  $\text{Sym}_n$ . The function  $\tilde{\mathcal{P}}_s : \mathbb{R}^d \rightarrow \mathbb{R}^d$  defined by

$$\tilde{\mathcal{P}}_s(\text{vec } X) = \text{vec } X^s$$

maps  $V$  into itself. Let  $\pi = \text{vec} \circ \nu$ , and notice that  $\pi : \mathbb{R}^m \rightarrow V$  is a linear isomorphism and that  $\mathcal{P}_s = \pi^{-1} \circ \tilde{\mathcal{P}}_s \circ \pi$ . By Example 9.4, if  $X$  is a positive definite

matrix then  $J_{\overline{\mathcal{P}}_s}(\text{vec } X) = dX^s/dX$  is also positive definite. Thus we may apply Lemma 10.1 to conclude that the Jacobi matrix  $J_{\mathcal{P}_s}(\pi^{-1}(\text{vec } X)) = J_{\mathcal{P}_s}(\mu(X))$  has positive eigenvalues at any positive definite  $X$ .  $\square$

**Proposition 10.3.** *Let  $s$  be a positive integer,  $P$  a positive definite matrix, and  $\mathcal{V} \subset \mathcal{O}$  a bounded open set containing  $\mu(P^{1/s})$ . Let  $g$  be the function  $\mathcal{P}_s$  restricted to  $\overline{\mathcal{O}}$ . Then*

$$\deg(g, \mathcal{V}, \mu(P)) = 1.$$

*Proof.* Lemma 10.2 implies that  $\mu(P)$  is a regular value for  $g$ . Using Theorem 7.3 and Lemma 10.2 we calculate

$$\deg(g, \mathcal{V}, \mu(P)) = \sum_{\mathbf{x} \in g^{-1}(\mu(P))} \text{sgn } \det J_g(\mathbf{x}) = \text{sgn } \det J_g(\mu(P^{1/s})) = 1.$$

$\square$

We are now ready to calculate the Brouwer degree of a symmetric word equation.

*Proof of Theorem 1.5.* From the discussion in Section 6 we may assume that our interlaced symmetric word  $S(X, B_1, \dots, B_k)$  is of the form (2). Fix positive definite matrices  $B_1, \dots, B_k$ , a positive definite matrix  $P$  and set  $f(X) = S(X, B_i)$ . Also set  $\tilde{f} = \mu \circ f \circ \nu$ . We will show that there is a bounded, open subset  $\mathcal{V} \subset \mathcal{O}$  such that for all bounded, open  $\mathcal{U} \subset \mathcal{O}$  with  $\mathcal{V} \subset \mathcal{U}$  we have

$$(13) \quad \deg(\tilde{f}, \mathcal{U}, \mu(P)) = 1.$$

By Proposition 8.2, there exists a constant  $K$  independent of  $t$  such that any positive definite solution  $X$  of the equation  $S(X, tB_i + (1-t)I) = P$  has  $\|X\| < K$ . Indeed, if

$$\alpha = \max_{1 \leq i \leq k, 0 \leq t \leq 1} \|tB_i + (1-t)I\| \cdot \|(tB_i + (1-t)I)^{-1}\| < \infty,$$

and

$$\beta = \min_{1 \leq i \leq k, 0 \leq t \leq 1} \|tB_i + (1-t)I\| > 0$$

then

$$\|X\| \leq C_{S, \alpha} \beta^{-\frac{2k}{s}} \|P\|^{\frac{1}{s}} < \infty.$$

Let  $V = V_K$  be the open set of positive definite matrices with norm less than  $K$ . For each  $t \in [0, 1]$ , let  $f_t$  be the function from the positive semidefinite matrices into itself given by  $f_t(X) = S(X, tB_i + (1-t)I)$ . From our choice of  $K$ , it follows that  $f_t(X) \neq P$  when  $X$  is positive definite with  $\|X\| = K$ . Moreover, if  $X$  is singular, then taking a determinant shows that  $f_t(X) \neq P$ . Thus  $P \notin f_t(\partial V)$  for each  $t \in [0, 1]$ .

Let  $\mathcal{V} = \mu(V)$  and  $\tilde{f}_t = \mu \circ f_t \circ \nu$ . Then if  $\mathcal{V} \subset \mathcal{U} \subset \mathcal{O}$  we have  $\mu(P) \notin \tilde{f}_t(\partial \mathcal{U})$  for  $t \in [0, 1]$ . Since  $(\mathbf{x}, t) \mapsto \tilde{f}_t(\mathbf{x})$  is continuous, Theorem 7.4 implies that

$$\deg(\tilde{f}_0, \mathcal{U}, \mu(P)) = \deg(\tilde{f}_1, \mathcal{U}, \mu(P)).$$

Since  $\tilde{f}_0 = \mathcal{P}_s$  and  $\tilde{f}_1 = \tilde{f}$ , (13) now follows from Proposition 10.3.  $\square$

11. PROOF OF THEOREM 1.4

A corollary of Theorem 1.5 will allow us to prove Theorem 1.4.

**Corollary 11.1.** *Fix an interlaced symmetric word  $S$  and let  $\tilde{f} = \tilde{f}_S$  be as in the proof of Theorem 1.5. Suppose there is a positive definite matrix  $X_0$  such that*

$$\det J_{\tilde{f}}(\mu(X_0)) < 0.$$

*Then the symmetric word equation*

$$S(X, B_i) = S(X_0, B_i)$$

*has at least two real solutions  $X$ .*

*Proof.* Set  $P = S(X_0, B_i)$ . If  $\mu(P)$  is a regular value of  $\tilde{f}$ , then Theorems 1.5 and 7.3 imply that there must be at least two solutions  $X_1$  and  $X_2$  of  $S(X, B_i) = P$  such that

$$\det J_{\tilde{f}}(\mu(X_i)) > 0, \quad i = 1, 2.$$

If  $\mu(P)$  is not a regular value of  $\tilde{f}$ , then there exists a positive definite matrix  $X_1$  such that  $S(X_1, B_i) = P$  and

$$J_{\tilde{f}}(\mu(X_1)) = 0.$$

Since  $J_{\tilde{f}}(\mu(X_1)) \neq J_{\tilde{f}}(\mu(X_0))$ , we have that  $X_0 \neq X_1$ . □

Let  $S$  and  $\tilde{f}$  be as in Corollary 11.1. We will now outline a method for obtaining the smaller Jacobian matrix  $J_{\tilde{f}}(\mu(X))$  from the larger Jacobian matrix  $dS/dX$ . To simplify the bookkeeping of indices, define

$$\alpha(i, j) = n(j - 1) + i, \quad i, j = 1, \dots, n$$

and

$$\beta(k, l) = \frac{1}{2}(2n - l)(l - 1) + k, \quad 1 \leq l \leq k \leq n.$$

Thus if  $X = [x_{ij}] \in \mathbb{M}_n$  then the  $\alpha(i, j)$ th entry of  $\text{vec } X$  is equal to  $x_{ij}$ ,  $i, j = 1, \dots, n$ . Likewise, the  $\beta(k, l)$ th entry of  $\mu(X)$  is  $x_{kl}$ ,  $1 \leq l \leq k \leq n$ .

The Jacobi matrix  $J_{\tilde{f}}$  of the map

$$\begin{aligned} \tilde{f} &= \mu \circ (X \mapsto S(X, B_i)) \circ \nu \\ &= (\mu \circ \text{vec}^{-1}) \circ (\text{vec} \circ (X \mapsto S(X, B_i)) \circ \text{vec}^{-1}) \circ (\text{vec} \circ \nu). \end{aligned}$$

is given by

$$J_{\tilde{f}}(\mu(X)) = M(dS/dX)N,$$

in which  $M \in \mathbb{M}_{m \times d}$  is the matrix representation of  $\mu \circ \text{vec}^{-1}$  and  $N \in \mathbb{M}_{d \times m}$  is the matrix representation of  $\text{vec} \circ \nu$ . It is easy to see that if  $1 \leq i, j \leq n$  and  $1 \leq l \leq k \leq n$  then the  $(\alpha(i, j), \beta(k, l))$  entry of  $N$  is

$$\begin{cases} 1 & \text{if } i = k, j = l \text{ or } i = l, j = k \\ 0 & \text{otherwise} \end{cases}$$

and the  $(\beta(k, l), \alpha(i, j))$  entry of  $M$  is

$$\begin{cases} 1 & \text{if } i = k, j = l \\ 0 & \text{otherwise.} \end{cases}$$

We are now ready to prove our main result.

*Proof of Theorem 1.4.* Let  $A_1$  and  $B_1$  be as in Section 4, and let  $S$  be the symmetric word

$$S(X, B) = XBX^2B^3X^2BX.$$

Let  $f(X) = S(X, A_1)$  and  $\tilde{f} = \mu \circ f \circ \nu$ . Using Maple<sup>1</sup>, we calculate

$$\det J_{\tilde{f}}(B_1) = -633705909477329213831177437148144640 < 0.$$

By Corollary 11.1, it follows that the symmetric word equation

$$S(X, A_1) = S(B_1, A_1)$$

has at least two distinct real positive definite solutions  $X$ . □

As a final remark, we note that there are many other words which can be proved to have multiple solutions using the techniques found in the proof of Theorem 1.4. We list a few of them below:

$$\begin{aligned} XBX^kBX, & \quad 9 \leq k \leq 20 \\ XBXB^2X^kB^2XBX, & \quad 2 \leq k \leq 16 \\ XBX^kB^3X^kBX, & \quad 2 \leq k \leq 15 \\ XB^2XBX^kBXB^2X, & \quad 6 \leq k \leq 40. \end{aligned}$$

## 12. ACKNOWLEDGMENTS

We would like to thank Bill Helton for many interesting and useful conversations about this problem.

## REFERENCES

- [1] D. Bessis, P. Moussa and M. Villani, *Monotonic converging variational approximations to the functional integrals in quantum statistical mechanics*, J. Math. Phys. **16**, 2318–2325 (1975).
- [2] R. Bhatia, *Matrix analysis*, Springer, New York, 1996.
- [3] M. Drmota, W. Schachermayer and J. Teichmann, *A hyper-geometric approach to the BMV-conjecture*, Monatshefte für Mathematik, to appear.
- [4] M. Fannes and D. Petz, *Perturbation of Wigner matrices and a conjecture*, Proc. Amer. Math. Soc. **131** (2003), 1981–1988.
- [5] I. Fonseca and W. Gangbo, *Degree theory in analysis and applications*, Oxford University Press, New York, 1995.
- [6] C. Hillar, *Advances on the Bessis-Moussa-Villani trace conjecture*, preprint.
- [7] C. Hillar and C. R. Johnson, *Eigenvalues of words in two positive definite letters*, SIAM J. Matrix Anal. Appl., **23** (2002), 916–928.
- [8] C. Hillar and C. R. Johnson, *Positive eigenvalues of generalized words in two Hermitian positive definite matrices*. Novel approaches to hard discrete optimization (Waterloo, ON, 2001), 111-122, Fields Inst. Commun., 37, Amer. Math. Soc., Providence, 2003.
- [9] C. Hillar and C. R. Johnson, *Symmetric word equations in two positive definite letters*, Proc. Amer. Math. Soc., **132** (2004), 945-953.
- [10] C. Hillar and C. R. Johnson, *On the positivity of the coefficients of a certain polynomial defined by two positive definite matrices*, J. Stat. Phys., **118** (2005), 781–789.
- [11] C. Hillar, C. R. Johnson and I. M. Spitkovsky, *Positive eigenvalues and two-letter generalized words*, Electronic Journal of Linear Algebra, **9** (2002), 21–26.
- [12] R. Horn and C. R. Johnson, *Matrix analysis*, Cambridge University Press, New York, 1985.
- [13] R. Horn and C. R. Johnson, *Topics in matrix analysis*, Cambridge University Press, New York, 1991.

---

<sup>1</sup>Code that performs this calculation is available at <http://math.berkeley.edu/~sarm>.

- [14] J. Lawson and Y. Lim, *private communication*, 2005.
- [15] E. H. Lieb, R. Seiringer, *Equivalent forms of the Bessis-Moussa-Villani conjecture*, J. Stat. Phys., **115** (2004), 185-190.
- [16] N. Lloyd, *Degree theory*, Cambridge University Press, London, 1978.
- [17] J. R. Magnus and M. Neudecker, *Matrix differential calculus with applications in statistics and econometrics*, John Wiley, New York, 1999.
- [18] Nathan Miller,  *$3 \times 3$  cases of the Bessis-Moussa-Villani conjecture*, Princeton University Senior Thesis, 2004.
- [19] Y. E. Nesterov and M. J. Todd, *Self-scaled barriers and interior-point methods for convex programming*, Mathematics of Operations Research **22** (1997), 1-42.
- [20] P. Sebastiani, *On the derivatives of matrix powers*, SIAM J. Matrix Anal. Appl., **17** (1996), 640-648.
- [21] G. Teschl, *Nonlinear Functional Analysis*, lecture notes available at <http://www.mat.univie.ac.at/~gerald/ftp/book-nlfa/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720.  
*E-mail address:* `sarm@math.berkeley.edu`

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843.  
*E-mail address:* `chillar@math.tamu.edu`