### NON-MARKOV RANDOM FIELDS W ITH LINEAR REGRESSIONS - A TOEPLITZ OPERATORS APPROACH

#### WOJCIECH MATYSIAK AND PAWEL J.SZABLOWSKI

Abstract. The aim of the paper is to analyze square integrable random sequences  $X = (X_k)_{k2Z}$  satisfying condition

$$E [X_{k} ::::; X_{k-2}; X_{k-1}; X_{k+2}; :::] = b_{j} X_{k-j} + X_{k+j}$$

with  $b_j \ge R$  and  $n \ge N$  [f+1 g. The question of existence of such sequences for all n 2 N including n = +1 is exam ined and som e conditions guaranteeing existence are provided. In order to give these conditions we analyze general problem of existence of processes de ned by regression coe cients. The problem is closely related to one considered by K ingm an and W illiam s.

O ne of the results presented in the paper is that one sided regressions of X are also linear:

$$E (X_{k} j ::: ; X_{k-2} ; X_{k-1}) = \int_{jX_{k-j}}^{X^{n}} x_{k-j}$$

for some  $\frac{1}{2}$  R and with the same n as before.

### 1. Introduction

W Bryc in 2001 (and later other authors) considered  $X = (X_k)_{k2Z}$  { a square integrable random sequence satisfying the following conditions in posed on their rst two conditionalmom ents:

(1) 
$$E[X_k F_{6k}] = a(X_{k-1} + X_{k+1}) + b;$$
  
(2)  $E[X_k^2 F_{6k}] = A[X_{k-1}^2 + X_{k+1}^2 + BX_{k-1}X_{k+1} + C + D(X_{k-1} + X_{k+1});$ 

where  $F_{6k} := (X_j: j \in k)$ . In the series of papers [3], [2], [6], [7], the consequences of these conditions were thoroughly analyzed and as a result, an almost complete characterization of possible values of parameters assuring existence of X and almost complete characterization of nite dimensional distributions of X were given. The crucial step in the analysis was to observe that (1) in plies that one sided regressions are linear:

$$E[X_k ::::X_{k-2};X_{k-1}] = X_{k-1}:$$

From symmetry, also  $E[X_k X_{k+1}; X_{k+2}; :::] = X_{k+1}$ . The next step was to realize that this is a part of a more general pattern: from (1) and (2) together, it

<sup>2000</sup> M athem atics Subject C lassi cation. Prim ary: 60G 07; Secondary: 60G 10, 60G 12.

K ey words and phrases. D iscrete tim e stochastic processes, stationary processes, harnesses, Toeplitz operators.

follows that

$$E [Q_n (X_{k+1}) j:::;X_k _ 1;X_k] = {}^n Q_n (X_k);$$
  

$$E [Q_n (X_k) jX_{k+1};X_{k+2};:::] = {}^n Q_n (X_{k+1});$$

where Q<sub>n</sub> are so called q-H erm ite polynom ials de ned by the recurrence

This enabled to read the properties of the random sequence X from the properties of the qH em ite polynom ials. In particular, one could identify the one dimensional distributions of X as measures that orthogonalize qH em ite polynom ials (for q 2 R); also, by (3) the M arkov property of X was easily established.

The paper is mainly devoted to processes that satisfy generalized condition (1). Namely, throughout this paper X will denote the discrete, norm alized, stochastic process satisfying the following condition:

(4) 
$$E [X_k F_{6k}] = b_j (X_{k-j} + X_{k+j})_{j=1}$$

for all k 2 Z: Here and further it will be assumed that the series on the right hand side of the above condition converges almost surely. We will also assume that covariance matrix of any subset  $fX_i : i2 J$  Zg is non-singular. As an important digression we will discuss the general problem of existence of processes de ned by regression coe cients, the problem discussed by W iliams [9] and K ingm an [5].

We formulate conditions under which processes de ned by (4) exist and are  $\rm L_2$ -stationary. Finally we shall show that

(5) 
$$E[X_{k} F_{k-1}] = \int_{j=1}^{X} X_{k-j}$$
:

for some coe cients  $_1$ ;  $_2$ ; ::: related to  $b_1$ ;  $b_2$ ; :::.

Furthermore, it will turn out that if there exists m 2 N such that  $b_j = 0$  for j > m, then j = 0 for j > m.

The investigation of the processes satisfying condition (4) involves many di erent tools known in analysis, including Toeplitz operators, i.e. operators de ned by matrices with constant diagonals.

The paper is organized as follows. In section 2 we consider general problem of mutual relationship between covariances and linear regressions coe cients. The statement of the problem and its analysis resemble that done by W illiams [9] and K ingm an [5]. However, we believe that our approach is more readable and understandable. The results of this general analysis conned to speci c situation de ned by condition (4) supply condition for the existence of processes given by (4). We also formulate some conditions under which an  $L^2$ -process can be decomposed into union of mutually uncorrelated  $L^2$ -processes. In section 3 we give conditions for the existence of  $L^2$ -processes satisfying (4). In section 4 main results of the paper are proved. They concern the form of one-sided regressions in both nite and in nite case.

### 2. Covariances and linear regressions

Let  $Y = fY_jg_{j2J}$ , J Z, be a square integrable zero-mean random vector consisting of linearly independent random variables. Linear independence means that there do not exist  $c_j$ ,  $j \ge J$ , (not all zero), such that

$$P \stackrel{0}{=} X \\ p \stackrel{1}{=} c_{j}Y_{j} = 0^{A} = 1:$$

Equivalently, if we denote  $R := [C \text{ ov } (Y_i; Y_j)]_{i;j2J}$  then linear independence m eans that any m ain submatrix of R is non-singular.

De ne

$$\begin{array}{rcl} H_{6j} & \rightleftharpoons & \operatorname{span} fY n Y_{j}g; & j 2 J; \\ H & \rightleftharpoons & \operatorname{span} Y; \end{array}$$

where closure is in  $L^2$ . Obviously H and H<sub>6k</sub> (k 2 Z) are closed subspaces of  $L^2$  and therefore H ilbert spaces with inner product hS;Ti = Cov(S;T).

By Proj(Y\_j)Y\_{\textit{6} j}) we denote the orthogonal projection of Y\_j on H\_{\textit{6} j} and assume that

(6) 
$$P \operatorname{roj}(Y_{j}) \stackrel{L^{2}}{=} X_{a_{j;i}Y_{i}}$$

for som e complex  $a_{j;i}$ . (Notice that if # J < 1 then form ulas for  $P \operatorname{roj}(Y_j)_{f \in j}$ ) are the well-known linear regression form ulas.) Let us introduce matrix A with (i; j)-th entry  $(i; j \ge J)$  de ned as

Let us also de ne

$$v_{j} \coloneqq E Y_{j}^{2} \qquad a_{j;i}Cov(Y_{j};Y_{i}); j 2 J;$$

$$J_{J_{j}} = diag(v_{j}; j 2 J):$$

Sincefork€j

$$C \text{ ov } (Y_{j};Y_{k}) = C \text{ ov } \operatorname{\mathbb{P}} \operatorname{roj}(Y_{j}) _{f \in j});Y_{k}];$$

wehave

$$v_j = E[Y_j Proj(Y_j)Y_{ij})]^2$$
:

Therefore  $v_j > 0$  for all j 2 J. Indeed, if  $v_j = 0$  for some j 2 J, then  $Y_j = P \operatorname{roj}(Y_j)_{f \in j}$  P-almost surely { contradiction with the assumption of linear independence. (Again, if # J < 1 then the explicit formula for  $v_j$  in terms of covariances of Y is wellknown.)

Lemma 2.1. If Y is a random vector with A, R and de ned as in above, then

(7) A R = :

Proof. See [5] or [6].

Corollary 2.2.

(1) R is invertible if and only if A is invertible.

(2) R is invertible and has a block diagonal structure if and only if A is invertible and has a block diagonal structure.

Proof. The rst assertion is trivial. To obtain the second, observe that if R is block diagonal, then so is its inverse. Since  $A = R^{-1}$  and is diagonal, A must be block diagonal. The inverse implication follows from the analogous reasoning.

2.1. C om plete correlation. Now consider a random vector Y such that A is symmetric and invertible. By the symmetry we have AR = = RA, so  ${}^{1}R$  and R  ${}^{1}$  are left-hand side and right-hand side reciprocals of A (see [4]). Since A is invertible,  ${}^{1}R = R$   ${}^{1}$ , that is

(8) 
$$v_{i}^{1}(R)_{j;i} = v_{j}^{1}(R)_{i;j}; i; j \ge J:$$

However, it does not imply that  $v_i = v_j$  for all i; j 2 J.

Therefore let us introduce a relation 1 on the set J: i and j from J are in relation 1 (shortly: il j) if there exists 2 N,  $k_1$ ;:::; $k_s 2 J$  such that

 $k_1 = i;k_s = j \text{ and } (R)_{k_m ;k_{m+1}} \in 0 \quad 8 \text{ m} = 1; :::;s \quad 1:$ 

Having the relation introduced, we can state the following lemma with straightforward proof.

Lem m a 2.3. Assume that A is symmetric and invertible. If il j then  $v_i = v_j$ .

Obviously 1 is the equivalence relation, so it divides J into (disjoint) equivalence classes [ $_1$ ]; if i; j 2 [k]<sub>1</sub> for some k 2 J then il j and if i 2 [k]<sub>1</sub> and j 2 [m]<sub>1</sub>, where [k]<sub>1</sub>  $\in$  [m]<sub>1</sub>, then Y<sub>i</sub> and Y<sub>j</sub> are uncorrelated. Thus the random vector Y is divided into mutually uncorrelated disjoint subsets of random variables.

Remark 2.1. This fact is rejected in the structure of the matrix R. Namely, it can be reordered (in fact the coordinates  $(Y_j)_{j2J}$  can be reordered) in such a way, that R would have block diagonal structure. Each such diagonal block would consist of covariances of random variables with indices from one of the equivalence classes [i]. By C orollary 2.2, matrix A would also have block diagonal structure.

M oreover disjoint subsets of indices  $K_{k}^{\circ}$ ; k 2 J for which  $J = \int_{k^2 Z}^{k} K_{k}^{\circ}$  and (A)<sub>i;j</sub> =

0;8i2  $K_k^{\circ}$ ;j2  $K_m^{\circ}$ ;k  $\notin$  m are the same as previously de ned. That is  $K_k = K_k^{\circ}$  under proper permutation of J: Thus relationship 1 can be de ned with the help of entries of matrix A :

De nition 2.1. Random vector  $Y = fY_jg_{j2J}$ , J Z, is completely correlated if  $[j]_1 = J$  for all j 2 J.

In other words, Y is completely correlated if it cannot be decomposed into disjoint subsets of mutually uncorrelated random variables.

C orollary 2.4. If Y is a completely correlated random vector such that A is sym – metric and invertible, then there exists a positive constant v such that = v I.

In connection with the problem s considered in this section, the following questions seem to be open:

(1) How to characterize those matrices R that result in symmetric matrices A?

(2) How to characterize matrices R that result in matrices of the form v I?

# 3. Existence and L<sup>2</sup>-stationarity of non-M arkov random fields with linear regressions

From now on, we analyze sequences  $X = (X_k)_{k2Z}$  of square integrable, linearly independent and standardized random variables, satisfying (4) for all k 2 Z. For a given real sequence  $(b_n)_{n2N}$ , let B denote a doubly in nite symmetric matrix with (i; j)-th entry (i; j 2 Z) dened as

(9) (B)<sub>*i*;*j*</sub> := 
$$b_{ji \ jj}$$
; if if *i j*;  
1; if *i i j*.

If we denote  $R := [C \text{ ov } (X_i; X_j)]_{i; j \ge Z}$  then by Lem m a 2.1

 $BR = diag(v_k; k 2 Z);$ 

with  $v_k \coloneqq E X_k E [X_k F_{\epsilon_k}]^2$ .

Lem m a 3.1. Let X satisfy (4) for a sequence  $(b_n)_{n \ge N}$ . If

(10)  $GCD (fn 2 N : b_n \in 0g) = 1$ 

then there exists v > 0 such that  $v_k$  = v for all  $k \ 2 \ Z$  .

Proof. First of all notice that for each in nite set I one can nd its nite subset  $I_E$  I such that GCD ( $I_E$ ) = GCD (I). We see that in order to have complete correlation of the set Y it is enough to be able to nd for each pair of indices (i; j) (for simplicity assume that i < j) a sequence of natural numbers  $fk_1;k_2;\ldots;k_m$  g such that the numbers  $b_{ji}$   $k_1;j;b_{jk_1}$   $k_2;j;\ldots;b_{jk_m-1}$   $k_m$   $j;b_{jk_m}$   $j_j$  are nonzero and moreover ji  $k_1j$   $j_k$   $k_2j$   $\ldots$   $j_k$   $j_j = j$  i. But this can be done since ji  $k_1j;j_k$   $k_2j;\ldots;j_m$   $j_j = 1$ 

C orollary 3.2. If  $(b_n)_{n \ge N}$  is a sequence satisfying (10) and such that B is symmetric, invertible and positive de nite, then there exists a square integrable sequence  $X = (X_k)_{k \ge Z}$  satisfying (4). Additionally, if B<sup>-1</sup> is a Toeplitz matrix, then X is  $L^2$ -stationary.

Proof. Since G aussian processes have linear regression we can reduce the problem of existence of the process X to the existence of G aussian processes satisfying (6). On the other hand for the existence of G aussian process it is enough to show that its covariance m atrix is positive de nite. We deduce (by (7)) that B = is equal to inverse of R : And conversely R = is equal to inverse of B : . Now if inverse of B is Toeplitz and positive de nite, then so is R. As remarked earlier, the fact that R is positive de nite, in plies that G aussian process with R as its covariance m atrix exists. M oreover if R is Toeplitz it m eans that X is  $L_2$  stationary.

3.1. Toeplitz m atrices and their sym bols. To each Toeplitz m atrix  $A = \begin{bmatrix} i & j \end{bmatrix}_{i,j2Z}$  one relates a function (in fact a distribution)

A (!) = 
$$\begin{bmatrix} X \\ k^{2} Z \end{bmatrix}_{k^{2} Z} e^{-jk!}$$

This function is called symbol of Toeplitz matrix A (see e.g. [1]). Many properties of Toeplitz matrices are relected in the properties of their symbols. Generally one can relate symbols to a wide class of Toeplitz matrices, e.g. assuming only that

which results in the fact that the sym bolof Toeplitz m atrix with such entries would, in general, be a periodic distribution (see e.g. [8]). On the other hand, having a periodic distribution we can consider sequence of its Fourier coe cients and construct a Toeplitz m atrix with these coe cients as its entries. Thus we have m utual 1 1 correspondence between subclass of in nite Toeplitz m atrices with entries satisfying (11) and the space of periodic distributions. Since product of two distributions is not always well de ned, the product of two Toeplitz m atrices m ay not be Toeplitz. In order to avoid unnecessary com plications we will con ne ourselves to in nite Toeplitz m atrices will be denoted C . We have the following simple Lemma, which we state for com pleteness, since all its assertions are known, but within di erent contexts.

Lemma 3.3.

- (1) Linear combination of Toeplitz matrices belonging to C is Toeplitz matrix to C . Moreover, symbol of the combination is the combination of symbols.
- (2) P roduct of two m atrices from C also belongs to C and sym bolof a product is product of two sym bols.
- (3) An in nite symmetric matrix  $A = \lim_{ji j j i, j \ge Z} belonging to C is positive de nite if and only if sequence <math>(i)_{i=0}$  is positive de nite, or equivalently, its symbol is a positive function.

Proof. First assertion is trivial. To prove the second, let  $A = \begin{bmatrix} i & j \end{bmatrix}_{i,j2Z}$ ;  $B = \begin{bmatrix} i & j \end{bmatrix}_{i,j2Z}$ ;  $B = \begin{bmatrix} i & j \end{bmatrix}_{i,j2Z}$  be two in nite Toeplitz matrices and  $AB = D = \begin{bmatrix} d_{ij} \end{bmatrix}_{i,j2Z}$ . We have

$$d_{ij} = \underset{\substack{k \ge Z}{k \ge Z}}{\overset{\Lambda}{}} i + k + j:$$

If m = i k then for xed i; k = i m : W e have

$$d_{ij} = \begin{bmatrix} x^{i} & & \\ & m & i & m & j \end{bmatrix}$$

Thus  $d_{ij}$  depends on (i j); proving that D is Toeplitz matrix. Hence we can write x

$$d_k = \prod_{\substack{m \ k \ m \ 2 \ Z}} m k m :$$

Now, calculating the Fourier transform D (!) of the sequence  $(d_k)_{k2Z}$  and using the well known properties of the Fourier transform and convolution we see that

where A  $(!\ )$  and B  $(!\ )$  are Fourier transform s of matrices respectively A  $% (!\ )$  and B .

To prove the third assertion, notice that for any nite nonzero sequence of complex numbers  $(x_i)_{i=1}^N$ , we have

$$X^{N}$$
  
ji jj $x_{i}x_{j} > 0$ ;  
i;j= 1

which m eans positive de niteness of the sequence ( $_{i}$ ). On the other hand, it is also well known (Herglotz theorem) that for a positive de nite sequence there exists a nonnegative function (in fact a positive m easure) such that the sequence is a sequence of Fourier coe cients of the function.

C om bining assertions of C orollary 3.2 and Lemma 3.3 we get the following Theorem .

Theorem 3.4. Assume that  $(b_n)_{n \ge N}$  is a sequence such that (10) holds. If symbol of Toeplitz matrix B de ned by (P),

(12) B (!) = 1 
$$2 b_n \cos(n!)$$
  
 $n=1$ 

is continuous and positive for ! 2 [ ; ), then there exists a square integrable and  $L^2$ -stationary random eld X =  $(X_k)_{k2Z}$  satisfying (4) for the sequence  $(b_n)_{n2N}$ , and having spectral density of the form =B (!) for some .

Proof. From Lemma 3.3 it follows that the process X exists if and only if the coe cients fb<sub>i</sub>g are such that B (!) > 0, ! 2< ; ): Since distribution B is positive, we deduce that sequence 1;  $b_1$ ;  $b_2$ ;::: is positive de nite. Moreover since B is continuous we deduce that  $b_i$ ! 0 as i! 1 and also that symbol of B<sup>-1</sup> is of a form  $\frac{1}{B(!)}$  and is also continuous: Hence the process X is L<sub>2</sub> stationary and moreover its covariance matrix  $R = r_{ji \ jj}_{i,j2Z}$  has the property that  $r_{jnj}$ ! 0 as n! 1. Besides since  $R = B^{-1}$  we deduce that matrix R is de ned by the function =B (!): On the other hand this function is, by the de nition, a spectral density of the process X :

Remark 3.1. The form of the spectral density discussed above generalizes the one obtained by Bryc in [3].

3.1.1. Bounded and positive symbols. W hat conditions should sequence  $(b_n)_{n 2 N}$  satisfy for function B de ned in (12) to be bounded and positive? One has to distinguish two cases:

(1)  $9 N > 0 : 8 n N b_{h} = 0$ , (2)  $8N 1 9 n N : b_{h} \neq 0$ :

In the rst case, B is a trigonom etric polynom ialand conditions for its positivity are given by the well known Fejer { R iesz T heorem that states that trigonom etric polynom ial of order N is positive if and only if there exists a polynom ial C (z) =  $P_{\substack{i=0\\j=0}}^{N} c_i z^i$  with real coe cients, such that

B (!) = C 
$$e^{i!}^{2}$$
:

M oreover,

$$X^{N}$$
  $c_{i}^{2} = 1;$   $b_{k} = \int_{i=0}^{N} X^{k} c_{k+i} c_{i};$   
 $b_{k-i} = 0$ 

and C can be chosen in such a way that it has all zeros inside the unit circle.

In the second case, from the Szego Theorem it follows that B is positive if and only if  $$\rm Z$$ 

and then

B (!) = C 
$$e^{i!}$$
 <sup>2</sup>;

where

C (z) = exp 
$$\frac{1}{2}c_0 + \sum_{i=1}^{X^i} c_i z^i$$
 :

In this case, the relation between coe cients (c  $_n$ ) and (b $_n$ ) is rather complicated. Notice how ever that if only coe cients (b  $_n$ ) satisfy

(13) 
$$X \quad p_n j < \frac{1}{2};$$

then B is positive. Besides, when, as in [6],  $b_n = 0$  for n 2, (13) reduces to  $b_1 \leq 1=2$ , (com pare [6]).

### 4. Linearity of one-sided regressions

4.1. F in ite case.

Theorem 4.1. Let  $X = (X_k)_{k \ge Z}$  be  $L^2$ -stationary sequence of random variables such that there exist n 2 N and real numbers  $b_1$ ;:::; $b_n$  with  $b_n \notin 0$ 

(14) 
$$E[X_k F_{6k}] = b_j (X_{k-j} + X_{k+j}) P-a.s.;$$
  
 $j=1$ 

Then there exist  $_1$ ; ...;  $_n$  2 R such that

(15) 
$$E[X_{k} F_{k-1}] = \begin{cases} X^{n} \\ jX_{k-j} \end{cases}$$

Lem m a 4.2. For all k 2 Z; m 2 N conditional expectations E [X  $_k$  F  $_k$  1;  $_{k+m}$ ] and E [X  $_{k+m}$  1 F  $_k$  1;  $_{k+m}$ ] are linear combinations of random variables

X<sub>k</sub> n;:::;X<sub>k</sub> 1;X<sub>k+m</sub> ;:::;X<sub>k+m+n</sub> 1;

with some real coe cients that do not depend on k (coe cients at X  $_k$   $_n$  and X  $_{k+m\,+\,n}$   $_1$  are nonzero).

P roof. We proceed by induction on m . For m = 1 the assertion is true for all k 2 Z by (14). Now assume the assertion is true for all m M (M 1):

(16) 
$$E [X_{k} F_{k}]_{; k+m} = X^{n} X_{k} j + j_{m} X_{k+m} 1+j 8 k 2 Z;$$
  
 $E [X_{k+m}] F_{k}]_{; k+m} = X^{n} X_{k} j + J_{jm} X_{k+m} 1+j 8 k 2 Z;$   
 $j=1 X^{n} X_{k} j + J_{jm} X_{k+m} 1+j 8 k 2 Z;$   
 $j=1 j=1 j=1$ 

 $(j_m; -j_m; j_m; -j_m 2 R and, in particular, n_m - n_m \in 0)$ . Then

$$E [X_{k} F_{k-1}; k+M+1] = E E [X_{k} F_{k-1}; k+M] F_{k-1}; k+M+1] = = X^{n} jM X_{k-j} + 1M E [X_{k+M} F_{k-1}; k+M+1] + j= 1 j= 2$$

$$E [X_{k+M} \quad F_{k-1}; \ _{k+M+1}] = E E X_{(k+1)+M-1} \quad F_{(k+1)-1}; \ _{(k+1)+M} \quad F_{k-1}; \ _{k+M+1}] = \\ = \sum_{1,M} E [X_{k} \quad F_{k-1}; \ _{k+M+1}] + \sum_{j=2}^{N} \sum_{j=1}^{N} X_{k+1-j} + \sum_{j=1}^{N} X_{k+M+j};$$

If  $_{1,M} = 0 = \sim_{1,M}$  then the assertion follows immediately from the inductive assumptions. If  $_{1,M} \in 0$  or  $\sim_{1,M} \in 0$  then we obtained a system of linear equations with unknowns  $x \coloneqq E[X_k \ F_{k-1}; \ _{k+M+1}]$  and  $y \coloneqq E[X_{k+M} \ F_{k-1}; \ _{k+M+1}]$ . The system has only one solution since otherwise 1  $_{1,M} \sim_{1,M} = 0$  and assumed linear independence of random variables would imply  $_{n,M} = \overset{\circ}{}_{n,M} = 0$  -a contradiction. It is easy to check that coe cients at X  $_{k-n}$  and X  $_{k+M+n}$  in x and y are nonzero.

Proof of Theorem 4.1. By Lemma 4.2, (16) holds and multiplying both of its sides respectively by  $X_{k-n}$ ;:::; $X_{k-1}$  and  $X_{k+m}$ ;:::; $X_{k+m+n-1}$ , and taking expectations, we obtain the following system of 2n linear equations with 2n unknowns  $j_{\pi}$  and  $j_{\pi}$  (j = 1;:::;n):

Here and in the sequel

$$r_k \coloneqq Cov(X_0; X_k) = E(X_0X_k); k 2 Z:$$

Since the main matrix of (17) is the covariance matrix of random vector

$$[X_0; :::; X_n_1; X_{m+n}; :::; X_{m+2n-1}]^T;$$

its determ inant is nonzero. Observe that

$$X^{n} = X^{n}$$

$$r_{1} = \sum_{j=1}^{j+1} r_{j1 \ jj}; \ l = 1; \dots; n;$$

$$0 = X^{n} = \sum_{j+1}^{N^{n}} r_{j1+1 \ jj}; \ l = 1; \dots; n:$$

In particular, the last n equations in ply that  $_{j;1} = 0$  for j = 1; ...; n.

By the Levy D ownward Theorem applied to (16) with m  $\ ! \ 1$  , we deduce that lim it

$$L \coloneqq \lim_{\substack{m \mid 1 \\ j=1}} X^n$$

exists alm ost surely and

(18) 
$$E [X_{k}]_{m=1}^{n} F_{k-1; k+m} = X^{n}_{j;1} X_{k-j} + L:$$

and

But

$$E jL j = E \lim_{\substack{m \mid 1 \\ j=1}} X_{k+m} \sum_{\substack{m \mid 1 \atop j=1}} X_{k+m} \sum_{\substack{m \mid$$

so L = 0 a.s. Taking the conditional expectation of both sides of (18) with respect to F  $_{\rm k-1}$  yields (15).

Proposition 4.3. There is a 1 1 relationship between  $\infty e$  cients  $b^0 s$  and  $^0 s$  mentioned in Theorem 4.1:

(19a) 
$$\frac{1}{b_{h}} = \frac{1 + \frac{2}{1} + \cdots + \frac{2}{n}}{n};$$
$$\frac{b_{h-1}}{b_{h}} = \frac{1 + \frac{2}{1} + \cdots + \frac{2}{n}}{n};$$
$$\frac{b_{h-1}}{b_{h}} = \frac{1 + \frac{2}{1} + \cdots + \frac{2}{n}}{n};$$
$$\frac{b_{h}}{b_{h}} = \frac{n + 2 + 1 + n + 1 + 2 + n}{n};$$

and (20a)

$$b_{1} = \frac{n \cdot 1 - 1 \cdot n}{1 + \frac{2}{1} + \frac{1}{1} + \frac{2}{1} + \frac{2}{1}};$$
  

$$b_{n-1} = \frac{1 - 1 \cdot 2 \cdot \frac{2}{1} + \frac{1}{1} + \frac{1}{1} + \frac{1}{1}}{1 + \frac{2}{1} + \frac{2}{1} + \frac{2}{1}};$$
  

$$b_{n} = \frac{n}{1 + \frac{2}{1} + \frac{2}{1} + \frac{2}{1} + \frac{2}{1}};$$

Proof. The fact that coe cients b  $_{\rm j}$  de ne $_{\rm j}$  follows from Theorem 4.1. To obtain converse relationship, we multiply both sides of (15) by X  $_{\rm k}$   $_{\rm m}$ , m = 1;2;::; and then take expectation to get

(21) 
$$\mathbf{r}_{m} = \begin{bmatrix} X^{n} & & \\ & j \mathbf{r}_{m} & j ; \\ & j=1 \end{bmatrix}$$

analogously, from (14) we get

(22) 
$$r_{m} = \sum_{j=1}^{X^{n}} b_{j} (r_{m} + r_{m+j}):$$

Characteristic polynom ial of (21) is

$$q_n(x) \coloneqq x^n \qquad \qquad x^{n} \qquad \qquad y^{n-j}$$

and (monic) characteristic polynom ial of (22) is

$$p_{2n}(\mathbf{x}) \coloneqq b_n^{-1} \cdots b_j x^{n-j} + x^{n+j} x^n ;$$

$$r_{m} = \sum_{j=1}^{\hat{X}^{n}} c_{j} (m) x_{j}^{m};$$

where  $c_j$  is a polynom ial (in variable m) of degree equal to the multiplicity of the root  $x_j$ . Observe that if  $p_{2n}(x) = 0$  for some  $x \notin 0$  then  $p_{2n} \times x^{-1} = 0$ . Observe also that  $jx_j \notin 1$ , j = 1;:::;2n. This is so since if y was one of the roots of  $p_{2n}$  with  $jy_j = 1$  then we would have

$$y^{m} = \sum_{j=1}^{X^{n}} b_{j} y^{m} j + y^{m+j};$$

which implies  $1 = {P \atop j=1}^{n} b_j y^{j} + y^{j}$ , and, consequently

$$1 = 2 \sum_{j=1}^{X^n} b_j \cos j ;$$

where  $y = \exp(i) - \operatorname{contradiction} w$  ith assumption 1  $2^{P_{j=1}} b_j \cos j > 0$  for all . Hence all the roots of  $p_{2n}$  are either inside or outside unit circle. Thus if  $j_{x_j} j < 1$  then exists i 2 f1;:::;2ng n fjg such that  $x_i = x_j^{-1}$  and  $j_{x_i} j > 1$ . Hence we have exactly (counting multiplicities) n roots of  $p_{2n}$  with absolute values sm aller than 1. Since  $r_n$  tends to 0,

$$r_m = \sum_{j=1}^{X^{\mathbf{P}}} c_j (m) x_j^m;$$

where 1 **e** n is a number of distinct roots  $x_j$  of polynom  $ialp_{2n}$  with  $jx_j j < 1$ . Since  $(r_m)_m$  satis es (21),  $q_n(x_j) = 0$  for j = 1; ...; **e**. Since

 $n^{1}x^{n}q_{n}x^{1}$ 

is a monic polynom ial (in variable x) of degree n with roots that are inverses of the roots of polynom ial  $q_h$ , we deduce that

(23) 
$$p_{2n}(x) = \int_{n}^{1} x^{n} q_{n}(x) dx^{-1} q_{n}(x)$$

W e use the above equation to get relationship between  $\cos$  cients b<sup>0</sup>s and <sup>0</sup>s: N amely,  $\cos$  paring the appropriate  $\cos$  cients at both sides of (23) yields (20) and (19).

Corollary 4.4. For allk 2 Z and m 2 N

$$E [X_{k} F_{k}r] = X_{k}^{T} r [1;0;:::;0]^{T};$$
where  $X_{k} = [X_{k};:::;X_{k}n+1]^{T}$  and  $= [i;j]_{i;j=1}^{n} 2 R^{n}$  with
$$\begin{cases} 8 \\ \ge i & \text{if } j = 1; \\ i;j = i+1 \text{ and } i < n; \\ 0 & \text{otherw ise:} \end{cases}$$

P roof. M in icking the proof of Lem m a 4.2, one can deduce that there exist constants  $_{1,r}$ ; ...;  $_{n,r}$  (and they do not depend on k) such that

(24) 
$$E [X_{k} F_{k} r] = \int_{j \neq 1}^{X^{n}} X_{k} r_{j+1}:$$

Sinœ

$$\mathbb{E} [X_k F_{k-r}] = \mathbb{E} \mathbb{E} [X_k F_{k-r+1}] F_{k-r}];$$

and by the linear independence assumption, applying (24) in the above equation and comparing the respective coe cients we get

Thus

j;r = j+1;r 1 + 1;r 1 j; j = 1;:::;n 1; n;r = 1;r 1 n:

Denen 1 vectors  $\mathbf{b}_r \coloneqq [1;r;\ldots;n;r]^T$  for r 2 N and  $\mathbf{b}_0 \coloneqq [1;0;\ldots;0]^T$ . Consequently

$$b_r = b_{r,1} r 2 N;$$

and the Corollary follows.

42. In nite case.

Theorem 4.5. Let  $X = (X_k)_{k22}$  be a square integrable random eld satisfying (4) for a sequence  $(b_n)_n$  constituting Toeplitz matrix with continuous and positive sym bol.

If  $_{i}jr_{i}j < 1$  then there exist a sequence (  $_{j})_{k2N}$  such that

(25) 
$$E [X_k F_{k-1}] = {}_{j}X_{k-j} P \text{-a.s.}$$

Lem m a 4.6. If  $X = (X_k)_{k2Z}$  satis es 4) then for all k 2 Z; m 2 N conditional expectations E  $[X_k F_{k-1}; k+m]$  and E  $[X_{k+m-1} F_{k-1}; k+m]$  are linear combinations of random variables

with some real coe cients that do not depend on k. In particular,

(26) 
$$E[X_k F_{k-1}; k+m] = \int_{jm}^{M} X_{k-j} + \int_{jm}^{M} X_{k+m+j-1}$$

Proof. See nite case.

Let  $l^1 \underset{P}{=} l^1(Z)$  denote Banach space of complex sequences  $x = (x_k)_{k2Z}$  with  $kxk_1 \coloneqq_{k2Z} \dot{y}_k j < 1$  and let B  $l^1$  denote Banach space of bounded linear operators acting on  $l^1$  (with usual operator norm).

Lem m a 4.7. Suppose that  $(b_n)_{n \ge N} \ge l^1$ ,  $(T_n)_{n \ge N} \ge B l^1$ . Assume also that there exist  $b_1 \ge l^1$  and invertible operator  $T_1 \ge B l^1$  such that  $b_n \ge b_1$  in  $l^1$  and  $T_n \ge T_1$  in  $B l^1$ . If sequence  $(x_n)_{n \ge N}$  satisfies

$$T_n x_n = b_n;$$

then  $x_n ! x_1 \coloneqq T_1^{-1}b_1$ .

Proof. De ne  $B_n := T_1^{-1}T_n$  for n 2 N. Since

$$kB_{n} Ik_{l^{1} l^{1} l^{1}} = kT_{1}^{1} (T_{n} T_{1})k_{l^{1} l^{1} l^{1}} kT_{1}^{1}k_{l^{1} l^{1} l^{1}}kT_{n} T_{1}k_{l^{1} l^{1} l^{1}} ! 0;$$

(I stands for the identity operator), B<sub>n</sub> tends to I in B  $l^1$ . Therefore there exists N 2 N such that kI  $B_n k_{l^1 ! \ l^1} < 1$  for all n > N. From now on we will consider only n > N. For such n the inverse  $B_n^{-1}$  exists and

(27) 
$$B_n^{1} = \sum_{k=0}^{X^{1}} (I B_n)^{k}; k B_n^{1} k_{l^{1}! l^{1}} \frac{1}{1 k I B_n k_{l^{1}! l^{1}}};$$

Notice that

$$\begin{array}{rcl} kB_{n}x_{n} & B_{n}x_{1} & k_{l^{1}} = kT_{1}^{-1}T_{n}x_{n} & B_{n}T_{1}^{-1}b_{l} & k_{l^{1}} = \\ & = kT_{1}^{-1}T_{n}x_{n} & T_{1}^{-1}b_{l} + T_{1}^{-1}b_{l} + B_{n}T_{1}^{-1}b_{l} & k_{l^{1}} \\ & & kT_{1}^{-1}k_{l^{1}+l^{1}} & kb_{n} & b_{l} & k_{l^{1}} + kT_{1}^{-1} & B_{n}T_{1}^{-1}k_{l^{1}+l^{1}} & kb_{l} & k_{l^{1}} \end{array}$$

Sinœ

$$\Gamma_{1}^{1} B_{n}T_{1}^{1} = T_{1}^{1} T_{1}^{1}T_{n}T_{1}^{1} = T_{1}^{1} I T_{n}T_{1}^{1}$$

and

Proof of Theorem 4.5. By Lemma 4.6, (26) holds and multiplying both of its sides respectively by

and

and taking expectations, we obtain the following (in nite) system of linear equations with unknowns  $_{jm}$  and  $_{jm}$  (j 2 N):

Here and in the sequel

0

 $r_k \coloneqq Cov(X_0; X_k) = E(X_0X_k); k 2 Z:$ 

Linear system (28) can be written in the matrix form

 $A_m x_m = b_m$ ;

where

$$\mathbf{x}_{m} = (:::; _{2,m} ; _{1,m} ; _{1,m} ; _{2,m} ;:::)^{T} ; \mathbf{b}_{m} = (:::; \mathbf{r}_{2} ; \mathbf{r}_{1} ; \mathbf{r}_{m} ; \mathbf{r}_{p+1} ; :::)^{T} ; \mathbf{A}_{m} = \begin{pmatrix} \mathbf{A}_{m}^{(1)} & \mathbf{A}_{m}^{(2)} \\ \mathbf{A}_{m}^{(3)} & \mathbf{A}^{(4)} \end{pmatrix}$$

0

1

and

$$A^{(1)} = \begin{bmatrix} B \\ 0 \end{bmatrix} \qquad \begin{array}{c} 1 & 1 & r & r_{2} & C_{A} \\ 1 & r & 1 & r_{1} & A \\ 0 \end{array} ; A^{(2)} = \begin{bmatrix} B \\ 0 \\ r_{m+2} & r_{m+3} & r_{m+2} \\ r_{m+2} & r_{m+3} \\ 0 \end{array} ; A^{(3)}_{m+2} = \begin{bmatrix} B \\ 0 \\ r_{m+2} & r_{m+3} \\ r_{m+1} & r_{m+2} \\ r_{m+2} & r_{m+1} \\ r_{m+3} & r_{m+2} & C_{A} \\ r_{m+4} & r_{m+3} & r_{m+3} \\ r_{m+4} & r_{m+3} & r_{m+3} \\ \end{array} ; A^{(4)} = \begin{bmatrix} B \\ 0 \\ 0 \\ r_{2} & r_{1} & 1 \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1} \\ r_{1} & r_{1} \\ r_{2} & r_{1} \\ r_{1} & r_{1}$$

1

(observe that in nite submatrices A  $^{(1)}$  and A  $^{(4)}$  do not depend on m ). Since  ${}_{j}\ jr_{j}\ j<1$  ,

$$b_m ! b_1 := (:::;r_2;r_1;0;0;:::):$$

Since norm of an in nite matrix A = (A  $_{i,j})_{i_{j}j\,2\,Z}$  , treated as element of B  $\,l^1\,$  , is

$$kA k_{l^{1}! l^{1}} = \sup_{j^{2} Z \atop i^{2} Z} A_{i;j}j;$$

we see that  $A_m$  tends in  $B l^1$  to  $A_1$  de ned as

$$A_1 = \begin{array}{c} A^{(1)} & 0 \\ 0 & A^{(4)} \end{array}$$
:

Inverse A  $_1$  <sup>1</sup> exists and is of the form

$$A_{1}^{1} = \begin{array}{c} A_{(1)}^{(1)} & 1 & 0 \\ 0 & A_{(4)}^{(4)} & 1 \end{array};$$

By Lemma 4.7,

$$x_m ! x_1 \coloneqq A_1 b_1$$

in  $l^1$ .

By the Levy D ownward Theorem applied to (26) with m  $!\,$  1 , we deduce that lim it of  $$x^{\!i}$$ 

alm ost surely and in  $L^1$ . If we denote

$$\mathbf{x}_{1} = (:::; 2;1; 1;1;1;1;2;1;:::)^{T}$$

then  $j_{i1} = 0, j = 1;2; :::, and$ 

Since

and

when m tends to in nity, we get that the right hand side of (26) tends in  $L^1$  to

Hence

(29) 
$$E X_{k} K_{k} F_{k 1; k+m} = j_{1} X_{k j} P-a.s.$$

$$m = 1 j = 1$$

Taking the conditional expectation of both sides of (29) with respect to F  $_{\rm k-1}$  yields (25).

5. Convergence of 
$$P_{n-1}$$
  $\dot{\mathbf{y}}_n$  j

From Lusin-Den joy theorem (see [10]) it follows that if trigonom etric series

is absolutely convergent on a set of positive measure, then  $p_j(j_{a_j}j_{a_j}+j_{a_j}j_{a_j}) < 1$ : On the other hand, we have two theorems on absolute convergence: Bernstein's theorem stating that if a function f 2 Lip ;  $> \frac{1}{2}$ ; then Fourier series of f converges absolutely and a theorem of Zygm und stating that if f is of bounded variation and 2 Lip ; > 0; then Fourier series of f converges absolutely.

Thus applying these results to spectral density of process  ${\tt X}$  i.e. function

f (!) = 1 + 2 
$$r_j \cos j!$$
;

applying relationship f (!) =  $\frac{\text{const}}{B(!)}$ ; where B (!) is a symbol of Toeplitz matrix B we can utter the following

Proposition 5.1. If symbol of matrix B i.e. function B (!) is of Lipschitz class > 0 and either > 1=2 or additionally B (!) is of bounded variation, then

## jr<sub>j</sub>j< 1 :

i 0

Х

A cknow ledgem ent. The authors thank W Bryc, R Latala and JW esolow ski for helpful rem arks and conversations. In particular, J.W esolow ski suggested the signi cant im provem ent of the proof of Lem m a 42.

### References

- 1. A lbrecht Bottcher and Sergei M. Grudsky, Toeplitz matrices, asymptotic linear algebra, and functional analysis, Birkhauser Verlag, Basel, 2000. MR MR 1772773 (2001g:47043)
- 2. W lodzim ierz Bryc, Stationary Markov chains with linear regressions, Stochastic Process. Appl.93 (2001), no.2, 339{348. M R M R 1828779 (2002d:60058)
- 3. W lodzim ierz Bryc, Stationary random elds with linear regressions, Ann. Probab.29 (2001), no.1, 504(519. M R M R 1825162 (2002d:60014)
- 4. Richard G.Cooke, In nite matrices and sequence spaces, Dover Publications Inc., New York, 1965. M R M R 0193472 (33 # 1692)
- 5. J. F. C. K ingm an, Random variables with unsymmetrical linear regressions, M ath. Proc. Cambridge Philos. Soc. 98 (1985), no. 2, 355{365. MR M R 795900 (87b:60021)
- 6. W ojciech M atysiak and Pawel J. Szablowski, A few remarks on Bryc's paper on random elds with linear regressions, Ann. Probab. 30 (2002), no. 3, 1486 (1491. M R M R 1920274 (2003e:60111)
- 7. \_\_\_\_\_, Bryc's random elds: the existence and distributions analysis, Preprint, 2003.
- 8. Laurent Schwartz, Metody matematyczne w zyce, Panstwowe Wydawnictwo Naukowe (PWN), Warsaw, 1984. MR MR 786468 (86m :00002)
- 9. David W illiam s, Som e basic theorem s on harnesses, Stochastic analysis (a tribute to the m em ory of Rollo Davidson) (D.G. K endall and E.F. Harding, eds.), W iley, London, 1973, pp. 349(363. M R M R 0362565 (50 # 15005)
- 10. A . Zygm und, Trigonom etric series. Vol. I, II, third ed., C am bridge M athem atical Library, C am bridge U niversity P ress, C am bridge, 2002, W ith a foreword by R obert A .Fe erm an. M R M R 1963498 (2004h:01041)

Faculty of M athematics and Information Science, W arsaw University of Technology, pl. Politechniki 1, 00-661 W arszawa, Poland

E-m ailaddress: matysiak@mini.pw.edu.pl

Faculty of M athematics and Information Science, W arsaw University of Technology,pl.Politechniki1,00-661 W arszawa,Poland

E-m ailaddress: pszablowski@elka.pw.edu.pl