UNIVERSAL FORMULAS FOR CHARACTERISTIC CLASSES ON THE HILBERT SCHEMES OF POINTS ON SURFACES

SAMUEL BOISSIÈRE AND MARC A. NIEPER-WISSKIRCHEN

ABSTRACT. This article can be seen as a sequel to the first author's article "Chern classes of the tangent bundle on the Hilbert scheme of points on the affine plane", where he calculates the total Chern class of the Hilbert schemes of points on the affine plane by proving a result on the existence of certain universal formulas expressing characteristic classes on the Hilbert schemes in term of Nakajima's creation operators.

The purpose of this work is (at least) two-fold. First of all, we clarify the notion of "universality" of certain formulas about the cohomology of the Hilbert schemes by defining a universal algebra of creation operators. This helps us to reformulate and extend a lot of the first author's previous results in a very precise manner.

Secondly, we are able to extend the previously found results by showing how to calculate any characteristic class of the Hilbert scheme of points on the affine plane in terms of the creation operators. In particular, we have included the calculation of the total Segre class and the square root of the Todd class.

Using this methods, we have also found a way to calculate any characteristic class of any tautological sheaf on the Hilbert scheme of points on the affine plane. This in fact gives another complete description of the ring structure of the cohomology spaces of the Hilbert schemes of points on the affine plane.

Contents

1. Introduction	2
2. Notation	3
2.1. Completed graded super-rings	3
2.2. Partitions	4
2.3. The symmetric algebra	4
2.4. Complex manifolds	5
3. Characteristic classes	5
3.1. Marked complex manifolds	5
3.2. Characteristic classes	6
3.3. Multiplicative classes	7
4. The Hilbert schemes of points on surfaces	9
4.1. Nakajima's creation operators	9
4.2. Hilbert schemes of marked complex surfaces	11
5. Universal formulas	13
6. The Hilbert scheme of the affine plane	16
Appendix A. Lagrange Inversion Theorem	21
Appendix B. Two algebraic identities	22
Beferences	24

Date: August 26, 2018.

²⁰⁰⁰ Mathematics Subject Classification. Primary 14C05; Secondary 14C17.

Key words and phrases. Hilbert schemes of points, characteristic classes, universal formulas.

1. INTRODUCTION

The Hilbert scheme of n points on a complex surface X (see [13] or [3] for the non-algebraic case) parametrises the zero-dimensional subspaces of length n on X. It is denoted in this text by $X^{[n]}$. By a result of Fogarty ([5]), it is smooth of dimension 2n. The vector space structure of its rational cohomology ring has been calculated by Göttsche ([6]), i.e. he calculated the Betti numbers. His formulas can be a given a particularly nice form if one considers all Hilbert schemes at once, i.e. if one studies Hilb $X := \prod_{n\geq 0} X^{[n]}$. Nakajima ([12]) and Grojnowski ([7]) were led by this result to the construction of certain operators $q_n(\alpha)$ and $p_n(\alpha)$ on the cohomology $\mathbf{H}_{\mathbf{Q}}X$ of Hilb X that are part of a vertex algebra structure (see [9]). Here α is a cohomology class on the surface X and n is a positive integer. The upshot of this construction is that every cohomology class in $\mathbf{H}_{\mathbf{Q}}X$ can be given as an application of a polynomial in the $q_n(\alpha)$ on a certain element $|\mathbf{0}\rangle$ in $\mathbf{H}_{\mathbf{Q}}X$, called the vacuum. In the case of the affine plane, i.e. $X = \mathbf{C}^2$, this leads to the isomorphism $\mathbf{H}_{\mathbf{Q}}\mathbf{C}^2 \simeq \mathbf{Q}[[p_1, p_2, \ldots]]$, the power series ring in infinitely many variables, as vector spaces (not rings!).

Each sheaf \mathcal{F} on X gives rise to a sheaf $\mathcal{F}^{[n]}$ on each $X^{[n]}$, called a tautological sheaf. One can ask what the total Chern classes of these sheaves in terms of polynomials in the $q_n(\alpha)$ applied to the vacuum are. Lehn gives in [9] a closed formula for the case that \mathcal{F} is an invertible sheaf.

Another natural sheaf on the Hilbert schemes is given by the tangent sheaf. The first author has started to attack the problem of expressing the total Chern classes of these tangent sheaves in terms of the $q_n(\alpha)$ in [2]. It turns out that these expressions do not really depend on the surface X (or the invertible sheaf \mathcal{F}) but only on some algebraic invariants of X and \mathcal{F} (namely more or less their Chern classes). Thus one can say there are *universal formulas* governing the Chern classes of the tautological and the tangent sheaves. Unfortunately the coefficients in these universal formulas are not known in a closed form for the case of the tangent sheaves of the Hilbert schemes on an arbitrary surface (or the tautological sheaves of a sheaf that is not of rank one). However, when one specialises to $X = \mathbb{C}^2$, the affine plane, there is a closed formula for the total Chern class of the tangent sheaves on the Hilbert schemes over X (expressed as elements in the ring $\mathbb{Q}[[p_1, p_2, \ldots]])$, which is the main result of [2].

This article builts upon these results. We clarify what is meant by universality of certain formulas by introducing a universal algebra of operators together with an evaluation map for each surface X that specialises these operators to the operators $q_n(\alpha)$. Equipped with this machinery, we easily rediscover the results of [2] that state how certain formulas have to look like.

We improve the methods that were used in [2] to calculate the total Chern classes of the tangent sheaves $\Theta_{X^{[n]}}$ of the Hilbert schemes of points on the affine plane to be able to calculate all multiplicative classes of $\Theta_{X^{[n]}}$, i.e. the total Segre class or Todd class. Given a multiplicative class $\gamma = \prod_{i=1}^{\infty} f(\lambda_i)$ where the λ_i are the (universal) Chern roots and $f \in 1 + xR[[x]]$ is a power series with coefficients in a ring R, one of our main results states that

$$\sum_{n\geq 0} \gamma(\Theta_{(\mathbf{C}^2)^{[n]}}) = \exp\left(\sum_{k\geq 1} g_{2k+1} \cdot \frac{p_{2k+1}}{2k+1}\right) \in R[[p_1, p_2, \dots]],$$

where the power series $g = \sum_{n \ge 1} g_n t^n$ is defined by $\frac{\partial g}{\partial t} \left(\frac{x}{f(x)f(-x)} \right) = f(x)f(-x)$. As every characteristic class appears in a sufficiently general multiplicative class, one can use this result to calculate effectively all characteristic classes of the sheaves $\Theta_{(\mathbf{C}^2)^{[n]}}$. As mentioned before, Lehn has calculated the total Chern classes of the tautological sheaves in the rank one case. We are able to extend his result to arbitrary multiplicative classes in the case of the affine plane (where there are only the tautological sheaves of the structure sheaf). Our second main result states that

$$\sum_{n\geq 0} \gamma(\mathcal{O}^{[n]}) = \exp\left(\sum_{n\geq 1} g_n \cdot \frac{p_n}{n}\right) \in R[[p_1, p_2, \dots]],$$

where this the power series $g = \sum_{n\geq 1} g_n t^n$ is defined by $\frac{\partial g}{\partial t} \left(\frac{x}{f(-x)}\right) = f(-x)$. As explained in the last section of this article, one can use this result to effectively calculate the cup-product in $\mathbf{H}_R \mathbf{C}^2 \simeq R[[p_1, p_2, \ldots]]$.

It is remarkable that these formulas for the case of the affine plane do also give partial results for the characteristic classes of the tautological sheaves and the tangent sheaves of the Hilbert schemes of points on any surface by virtue of the universality and uniqueness of the involved formulas. This is explained in the main text.

Let us outline the structure of this article. The first of the sections that follow is dedicated to the introduction of some notations and notions that are used throughout the text.

The third section gives some general results on (multiplicative) characteristic classes. As we want to discuss characteristic classes of the tangent sheaf of a complex manifold and characteristic classes of bundles on it uniformly, we introduce the notion of a marked complex manifold (X, α) , where X is a complex manifold and α is a cohomology class on X. In the applications, α will often by the Chern character of a sheaf on X, from which all other characteristic classes can be built. We introduce the ring of all (virtual) marked complex manifolds.

In the fourth section, we come to the Hilbert scheme of points on a surface. We give the definition and the construction of one half of Nakajima's operators (the half that interests us). It ends with defining the notion of a Hilbert scheme of a marked complex surface and explaining that the Hilbert scheme operation is like an exponential mapping from the abelian group of (virtual) marked complex surfaces to the ring of all (virtual) marked complex manifolds.

The fifth section is devoted to the universality of certain formulas, which has already been stated in [2]. Here, we add the remark that the universal expressions for multiplicative classes evaluated on the Hilbert schemes are unique, which makes it possible to get results for the general case from specific results on the Hilbert scheme of points on the affine plane.

The last section contains the two main results in that have been stated above. An algebraic identity that is used in the proof of these two theorems (and that makes it possible to generalise the result of [2] where hypergeometric methods were used instead), can be found in the appendix together with a certain form of the Lagrange Inversion Theorem, that is used to give our results a nice form.

2. NOTATION

2.1. Completed graded super-rings. By a *(unital)* super-ring R we understand a $\mathbb{Z}/(2)$ -graded unital ring that is commutative in the graded sense. In particular R comes with a decomposition $R = R_0 \oplus R_1$ such that in particular R_0 is a subring and R_1 is a module over R_0 . The elements in R_0 are the *even elements* (or *elements* of *even parity*) and the elements in R_1 are the *odd* ones.

Let R be such a super-ring. A grading on R is a decomposition $R = \bigoplus_{n \ge 0} R^n$ of R into subgroups such that R^n for n even consists entirely of even elements and R^n for n odd consists entirely of odd elements. We require further that $R^m \cdot R^n \subset$ R^{m+n} . A super-ring with a grading is a graded super-ring. The elements in R^n are the homogeneous elements of degree n.

A completed graded super-ring \hat{R} is a super-ring \hat{R} that comes with an inclusion $R \subset \hat{R}$ of a graded super-ring R such that \hat{R} is the *I*-adic completion of R with respect to the ideal $I := \bigoplus_{n>0} R^n$.

An ordinary ring R becomes trivially a completed graded super-ring by choosing the grading $R = R^0$. For example, we may view **Q** as a completed graded superring. Every graded super-ring can be made into a completed graded super-ring by passing to its completion with respect to the ideal generated by the elements of positive degree.

All of the rings that occur in this article will be completed graded super-ring. In order to simplify notions, we will sloppily call these completed graded super-rings just *rings*. It frequently happens that a construction, e.g. the tensor product, does not yield a completed graded super-ring but just a graded super-ring. In these cases, it is to be understood that one has to implicitely pass to the completion. In other words, "tensor product" will always mean "completed tensor product", etc.

For the rest of this paper, R will be a ring, i.e. a completed graded super-ring.

Let us mention that there are the obvious notions of modules and algebras over completed graded rings. We will make heavy use of them. One example will be the cohomology ring $H^*(X, R)$ of a complex manifold X. It is an R-algebra (in the above sense) with the grading $\bigoplus_{p+q>0} H^p(X, R^q)$.

Given an *R*-module M and a natural number $l \in \mathbf{N}_0$, we denote by M[-l] the *R*-module whose grading is given by

$$(M[-l])^n = \begin{cases} M^{n-l} & \text{for } n \ge l\\ 0 & \text{otherwise.} \end{cases}$$

2.2. **Partitions.** By **P**, we denote the set of partitions of the positive natural numbers, i.e. non-increasing finite sequences $(\lambda_1, \ldots, \lambda_n)$ of positive natural numbers. We view the non-zero natural numbers **N** as a subset of **P** by considering a number $n \in \mathbf{N}$ as the trivial partition (n) of itself.

Given a partition $\lambda = (\lambda_1, \ldots, \lambda_r) \in \mathbf{P}$, we call

$$|\lambda| := r$$

the length of λ and

$$\|\lambda\| := \lambda_1 + \dots + \lambda_r$$

the weight of λ , i.e. a partition λ of weight n is a partition of n.

2.3. The symmetric algebra. Let M be an R-module. We denote by $S_R M$ its (completed) symmetric algebra (symmetric in the super-sense, of course), i.e. $S_R M$ is the free R-algebra (in our category of completed graded commutative unital super-R-algebras) generated by M. It carries a natural decomposition

$$S_R M = \prod_{k=0}^{\infty} S_R^k M$$

with $S_R^0 M = R$ and $S_R^1 M = M$. Given an element $q \in S_R M$, we can write

$$q = \sum_{k=0}^{\infty} q_k$$

with each $q_k \in S_R^k M$. For each $n \in R$, we set

$$q(n) := \sum_{k=0}^{\infty} n^k q_k.$$

2.4. Complex manifolds. By \emptyset we denote the complex manifold without any point. By \ast we denote the zero-dimensional complex manifold with exactly one point.

For each complex manifold X, let Θ_X be its tangent sheaf.

Given any coherent sheaf \mathcal{F} on X, its Chern character in $H^{2*}(X, \mathbf{Q}) \subset H^*(X, R)$ is denoted by ch(\mathcal{F}). It decomposes as

$$\operatorname{ch}(\mathcal{F}) = \sum_{n=0}^{\infty} \operatorname{ch}_n(\mathcal{F})$$

with $\operatorname{ch}_n(\mathcal{F})$ being of degree 2n.

Given a product $X \times Y$ of two complex manifolds X and Y, we usually identify the cohomology space $H^*(X \times Y, R)$ with the tensor product $H^*(X, R) \otimes_R H^*(Y, R)$ given by Künneth's theorem.

3. Characteristic classes

3.1. Marked complex manifolds.

Definition 1. A marking α (over R) on a complex manifold X is an element $\alpha \in H^*(X, R)$.

A marked complex manifold (X, α) is a complex manifold X together with a marking α on X.

A morphism $f : (X, \alpha) \to (Y, \beta)$ between marked complex manifolds (X, α) and (Y, β) is a holomorphic map $f : X \to Y$ such that $f^*(\beta) = \alpha$.

Given a marking α on a complex manifold X, we write

$$\alpha = \sum_{n=0}^{\infty} \alpha_n$$

with $\alpha_n \in \bigoplus_{p+q=n} H^p(X, \mathbb{R}^q)$ being the component of α of (total) degree n.

Definition 2. Let (X, α) and (Y, β) be two marked complex manifolds. Their sum $(X, \alpha) + (Y, \beta)$ is given by

 $(X \amalg Y, i_*\alpha + j_*\beta),$

where $i: X \to X \amalg Y$ and $j: Y \to X \amalg Y$ are the natural inclusion maps. Their *product* $(X, \alpha) \cdot (Y, \beta)$ is given by

$$(X \times Y, p^*\alpha + q^*\beta),$$

where $p: X \times Y \to X$ and $q: X \times Y \to Y$ are the natural projection maps.

Remark 1. The neutral element with respect to the sum is the marked complex manifold $(\emptyset, 0)$. The neutral element with respect to the product is the marked complex manifold (*, 0). We thus may speak of the *commutative semiring of marked complex manifolds*.

Remark 2. By formally adding additive inverses, we may turn this semiring into the "ring" of (virtual) marked complex manifolds. The word "ring" is in quotation marks as this "ring" is a proper class and not a set. By tensoring this ring with any other ring S, we arrive at the S-algebra of marked complex manifolds, i.e. we can study formal S-linear combinations of marked complex manifolds. All the constructions for marked complex manifolds that will follow also extend to this formal combinations. 3.2. Characteristic classes. We define the ring

$$U_R := R[p_0, p_1, p_2, \dots, b_0, b_{\frac{1}{2}}, b_1, \dots]$$

with the grading defined by deg $p_i = 2i$ and deg $b_i = 2i$. (Note that this ring is in fact a power series ring as we have to implicitly complete it.)

Definition 3. The *R*-algebra U_R is the ring of characteristic classes of marked complex manifolds over *R*.

The reason for giving this name to the ring is the following: Given a marked complex manifold (X, α) , there is a unique homomorphism

$$U_R \to H^*(X, R), \quad \gamma \mapsto \gamma|_{(X,\alpha)}$$

of R-algebras with

$$p_i|_{(X,\alpha)} = \operatorname{ch}_i(\Theta_X)$$
 and $b_i|_{(X,\alpha)} = \alpha_{2i}$.

We set

$$\operatorname{ch} := \sum_{i=0}^{\infty} p_i$$
 and $b := \sum_{j=0}^{\infty} b_{j/2}.$

Then $\operatorname{ch}_{(X,\alpha)} = \operatorname{ch}(\Theta_X)$ and $b|_{(X,\alpha)} = \alpha$.

Definition 4. The element $ch \in U_R$ is the *universal Chern character*, the element $b \in U_R$ is the *universal marking*.

Lemma 1. Let $\gamma, \gamma' \in U_R$ with $\gamma|_{(X,\alpha)} = \gamma'|_{(X,\alpha)}$ for all marked complex manifolds (X, α) . Then $\gamma = \gamma'$.

Proof. This result is well-known for complex manifolds without markings, you may consult [8].

It is enough to consider the case that γ and γ' are sums of homogeneous elements of degree $n \in \mathbf{N}$ or less. The general case then follows by passing to the limit. Let us also assume for a moment that γ and γ' do not contain p_0 and b_0 .

Consider the (compact) complex manifold

$$X := \mathbf{G}_k(\mathbf{C}^{n+k}) \times \mathbf{C}^r / \mathbf{Z}^{2r},$$

which is the product of the Graßmannian of k-dimensional subspaces in \mathbf{C}^{n+k} and an r-dimensional torus with $r, k \gg 0$. Recall that the rational cohomology of the Graßmannian is freely generated up to degree n by the components ch_1, ch_2, \ldots, ch_k of the Chern character of its tangent bundle. Further, we have $H^p(\mathbf{C}^r/\mathbf{Z}^{2r}, R) = \Lambda_R^p(\mathbf{C}^r)$, so we find cohomology classes $\alpha_1, \alpha_2, \ldots, \alpha_n$ in the cohomology of the torus that are algebraic independent up to degree n, provided r is big enough. These classes define a marking α on X.

By these freeness results, we find that $\gamma|_{(X,\alpha)} = \gamma'|_{(X,\alpha)}$ implies $\gamma = \gamma'$.

It remains to consider the more general case with p_0 and b_0 appearing in γ . For this, consider γ as a polynomial $\gamma(p_0, b_0)$ in p_0 and b_0 . Let (X, α) be any marked complex manifold. From this, we construct for each pair $r, s \in \mathbb{Z}$ a new marked complex manifold $(X \times \mathbb{C}^r / \mathbb{Z}^{2r}, \alpha + s)$ with $\alpha + s$ being the image of $\alpha + s \in H^*(X, R)$ under the inclusion map

$$H^*(X,R) \to H^*(X \times \mathbf{C}^r / \mathbf{Z}^{2r}, R) \simeq H^*(X,R) \otimes_R H^*(\mathbf{C}^r / \mathbf{Z}^{2r}, R), \quad v \mapsto v \otimes 1.$$

By virtue of this embedding, we consider $H^*(X, R)$ as a subspace of the cohomology ring $H^*(X \times \mathbf{C}^r / \mathbf{Z}^{2r}, R)$. Then we can formulate:

$$\gamma|_{(X \times \mathbf{C}^r / \mathbf{Z}^{2r}, \alpha + s)} = \gamma(p_0 + r, b_0 + s)|_{(X, \alpha)}.$$

The same equation holds true for γ' in place of γ . Thus, we have

$$\gamma(p_0 + r, b_0 + s)|_{(X,\alpha)} = \gamma'(p_0 + r, b_0 + s)|_{(X,\alpha)}.$$

Expand both sides in powers of $p_0 + r$ and $b_0 + s$, e.g.

$$\gamma(p_0 + r, b_0 + s)|_{(X,\alpha)} = \sum_{l,m \ge 0} \gamma_{l,m} (p_0 + r)^l (b_0 + s)^m|_{(X,\alpha)}$$

with certain coefficients $\gamma_{l,m} \in U_R$ that do not contain p_0 and b_0 . We get a another expansion with γ replaced by γ' . By comparing coefficients, it follows that

$$\gamma_{l,m}|_{(X,\alpha)} = \gamma'_{l,m}|_{(X,\alpha)}$$

for all $l, m \ge 0$. By the previous result on the case where p_0 and b_0 do not appear, we get $\gamma_{l,m} = \gamma'_{l,m}$ and therefore $\gamma = \gamma'$.

Remark 3. Each characteristic class $\gamma \in U_R$ is "additive" in the following sense: It is

$$\gamma|_{(\emptyset,0)} = 0$$

and

$$\gamma|_{(X,\alpha)+(Y,\beta)} = i_*\gamma|_{(X,\alpha)} + j_*\gamma|_{(Y,\beta)}$$

in $H^*(X \amalg Y, R)$ for all marked manifolds (X, α) and (Y, β) .

3.3. **Multiplicative classes.** On the other hand, not every class is multiplicative in the following sense:

Definition 5. A characteristic class $\gamma \in U_R$ is *multiplicative* if

$$\gamma|_{(\{*\},0)} = 1$$

and

$$\gamma|_{(X,\alpha)\cdot(Y,\beta)} = p^*\gamma|_{(X,\alpha)} \otimes q^*\gamma|_{(Y,\beta)}$$

in $H^*(X \times Y, R) \simeq H^*(X, R) \otimes_R H^*(Y, R)$ for all marked manifolds (X, α) and (Y, β) . Here as before, p and q denote the two projections from $X \times Y$ onto its two factors.

To describe all multiplicative classes, we have to consider the ring

$$R[\lambda_1, \lambda_2, \dots]$$

with deg $\lambda_i = 2$. The symmetric group

$$\mathfrak{S} = \lim_{n \in \mathbf{N}_0} \mathfrak{S}_n$$

acts naturally on this R-algebra by permuting the variables λ_i . We define

$$\Lambda_R := R[\lambda_1, \lambda_2, \dots]^{\mathfrak{S}},$$

i.e. Λ_R is the *R*-algebra of \mathfrak{S} -invariants of the polynomial ring.

Definition 6. The *R*-algebra Λ_R is the (completed) ring of symmetric functions over *R*.

Example 1. The *n*-th elementary symmetric function in the λ_i is an element in Λ_R . It is denoted by c_n , i.e.

$$\sum_{i=0}^{\infty} c_i = \prod_{i=1}^{\infty} (1+\lambda_i)$$

in Λ_R with deg $c_i = 2i$.

Remark 4. By the theorem of elementary symmetric functions, the homomorphism

$$R[c_1, c_2, \dots] \to \Lambda_R$$

of *R*-algebras that map the (formal) variable c_i of degree 2i to the *n*-th symmetric function c_i is an isomorphism.

Consider the (completed) tensor product $\Lambda_R \otimes_R \Lambda_R$. To distinguish between the first and the second factor, let us decorate the variables coming from the second factor with a prime, e.g. we write λ'_i . Thus we have

$$\Lambda_R \otimes_R \Lambda_R = R[\lambda_1, \lambda_2, \dots, \lambda_1', \lambda_2', \dots]^{\mathfrak{S} \times \mathfrak{S}}.$$

Similarly, everything else arising from the second factor, e.g. any elementary symmetric function, is primed.

Remark 5. There is a unique homomorphism

$$\Lambda_R \otimes_R \Lambda_R \to U_R$$

of *R*-algebras mapping $\sum_{i=1}^{\infty} \lambda_i^n$ to p_n and $\sum_{i=1}^{\infty} \lambda_i'^n$ to b_n for all $n \in \mathbf{N}$. This homomorphism is a monomorphism. By virtue of this homomorphism, we view $\Lambda_R \otimes_R \Lambda_R$ as a subring of U_R .

We make use of this identification in the following theorem:

Theorem 1. Let $f, f' \in 1 + xR[[x]]$ be two power series in R with $f_0 = f'_0 = 1$ Then

$$\prod_{i=1}^{\infty} f(\lambda_i) \cdot \prod_{i=1}^{\infty} f'(\lambda_i')$$

is a multiplicative characteristic class in U_R .

On the other hand, every multiplicative class in uniquely given by such a construction.

Proof. This result is classical for complex manifolds without marking. See again [8]. To see that $\gamma = \prod_{i=1}^{\infty} f(\lambda_i) \cdot \prod_{i=1}^{\infty} f'(\lambda'_i)$ is multiplicative, we write

$$\gamma = \exp\left(\sum_{n \ge 1} (a_n p_n + a'_n b_n)\right)$$

with $\log f = \sum_{n \ge 1} a_n x^n$ and $\log f' = \sum_{n \ge 1} a'_n x^n$. Then one calculates $\gamma|_{(*,0)} = \exp(0) = 1$ and

$$\gamma|_{(X,\alpha)+(X',\alpha')} = \exp\left(\sum_{n\geq 1} \left((a_n p_n + a'_n b_n)|_{(X,\alpha)} + (a_n p_n + a'_n b_n)|_{(X',\alpha')} \right) \right)$$
$$= \gamma|_{(X,\alpha)} \cdot \gamma|_{(X',\alpha')}.$$

It remains to prove the implication in the other direction. So let $\gamma \in U_R$ be a multiplicative class. From $\gamma|_{(*,0)} = 1$ it follows, that the logarithm $\log \gamma \in U_R$ exists. Let

$$m: H^*(X^n, R) \simeq H^*(X, R)^{\otimes n} \to H^*(X, R)$$

be the multiplication map given the cup-product. As γ is multiplicative, we have

$$m\left(\log \gamma|_{(X,\alpha)^n}\right) = n\left(\log \gamma|_{(X,\alpha)}\right).$$

Any power series (with trivial constant term) in the p_i and b_i that fulfills this equality in place of $\log \gamma$, has to be linear in the p_i and b_i . Thus $\log \gamma$ is an *R*-linear combination of the p_i and b_i , which proves the other direction.

Example 2. The total Chern classes $c := 1 + c_1 + c_2 + \ldots$ and $c' := 1 + c'_1 + c'_2 + \ldots$ are multiplicative classes.

4. The Hilbert schemes of points on surfaces

4.1. Nakajima's creation operators. Let X be a complex surface. For each $n \in \mathbf{N}_0$ we denote by $X^{[n]}$ the Douady space (see [3]) of the zero-dimensional subspaces of X of length n, i.e. holomorphic maps from a complex space S to $X^{[n]}$ parametrise subspaces Z of $S \times X$ such that the restriction $p|_Z : Z \to X^{[n]}$ of the projection map $S \times X \to S$ is flat and finite of degree n.

In case X is an algebraic surface, the Douady spaces are algebraic as well. In fact, they are the corresponding Hilbert schemes of points on X viewed as a scheme. By abuse of notion, we shall not distinguish between the two notions:

Definition 7. The space $X^{[n]}$ is the Hilbert scheme of n points on X.

Example 3. It is $X^{[0]} = *$, the complex manifold consisting of one point, and $X^{[1]} = X$.

For each point $\xi \in X^{[n]}$, we denote by $\operatorname{supp} \xi = \operatorname{supp} \mathcal{O}_{\xi}$ its *support* which is a finite subset of X whose cardinality lies between 1 and n. Given two zerodimensional subspaces ξ and ξ' of X with $\operatorname{supp} \xi \subset \operatorname{supp} \xi'$, we denote by $\mathcal{I}_{\xi,\xi'}$ the kernel of the natural epimorphism $\mathcal{O}_{\xi'} \to \mathcal{O}_{\xi}$.

The identity $X^{[n]} \to X^{[n]}$ corresponds to a subspace Ξ^n of $X^{[n]} \times X$ such that the restriction $p|_{\Xi^n} : \Xi^n \to X^{[n]}$ of the projection $p : X^{[n]} \times X \to X^{[n]}$ is flat and finite of degree n. We denote its structure sheaf by \mathcal{O}_n and use the same symbol to denote the corresponding quotient sheaf of $\mathcal{O}_{X^{[n]} \times X}$ on $X^{[n]} \times X$.

Definition 8. The space Ξ^n is the universal family over $X^{[n]}$.

We generalise this subspace as follows: Let $n, l \in \mathbf{N}_0$ be two natural numbers. By $\Xi^{n,l}$ we denote the reduced subspace of $X^{[l+n]} \times X^{[l]} \times X$ whose support is the closure of the subset

$$\{(\xi',\xi,x)\in X^{[l+n]}\times X^{[l]}\times X|\operatorname{supp}\mathcal{I}_{\xi,\xi'}\subset\{x\}\}$$

We denote the projections of $X^{[l+n]}\times X^{[l]}\times X$ uniformly by $p,\,q$ and r as given in the diagram

$$\begin{array}{cccc} X^{[l+n]} & \xleftarrow{p} & X^{[l+n]} \times X^{[l]} \times X & \xrightarrow{r} & X \\ & & q \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\$$

The restriction $p|_{\Xi^{n,l}}:\Xi^{n,l}\to X^{[n+l]}$ of p is proper. Thus a push-forward operator

$$p_*(\cdot \cap [\Xi^{n,l}]) : H^*(X^{[l+n]} \times X^{[l]} \times X, R) \to H^*(X^{[l+n]}, R)$$

is well-defined. We use this map to construct Nakajima's creation operators in the sequel.

In order to do this, it is convenient to consider all Hilbert schemes of points of X at once. For this, we define

$$\operatorname{Hilb} X := \prod_{l=0}^{\infty} X^{[l]}.$$

The total cohomology space (up to some shifts to make it completed graded) is given by

$$\mathbf{H}_{R}X := \prod_{l=0}^{\infty} H^{*}(X^{[l]}, R)[-2l].$$

There are (at least) two distinguished elements in this cohomology space: One of them is the unit $|1\rangle$ of the (unshifted) cohomology ring, which is given by the infinite sum of the units in each $H^*(X^{[n]}, R)$, and the other one is the so-called *vacuum* $|0\rangle$ that is the image of $1 \in H^*(X^{[0]}, R) = H^*(*, R) = R$ in $\mathbf{H}_R X$.

Elements in the subspace $H^*(X^{[l]}, R)[-2l]$ of $H^*(\text{Hilb } X, R)$, $l \in \mathbf{N}_0$, are called elements of weight l.

Given $n \in \mathbf{N}$, let

$$q_n: H^*(X, R) \to \operatorname{End} \mathbf{H}_R X$$

be the operator-valued linear map such that for each $\alpha \in H^*(X, R)$ the restriction of $q_n(\alpha) \in \operatorname{End} \mathbf{H}_R X$ to $H^*(X^{[l]}, R)$ is given by

$$q_n(\alpha): H^*(X^{[l]}, R) \to H^*(X^{[l+n]}, R), \quad \beta \mapsto p_*((r^* \alpha \cup q^* \beta) \cap [\Xi^{n,l}]).$$

Given $n_1, \ldots, n_r \in \mathbf{N}$, we set

$$q_{n_1} \circ \cdots \circ q_{n_r} : H^*(X, R)^{\otimes r} \to \operatorname{End} \mathbf{H}_R X,$$
$$\alpha_1 \otimes \cdots \otimes \alpha_{n_r} \mapsto q_{n_1}(\alpha_1) \circ \cdots \circ q_{n_r}(\alpha_r),$$

and for a partition $\lambda = (\lambda_1, \dots, \lambda_r)$ of length r, we set

$$q_{\lambda}: H^*(X, R) \to \operatorname{End} \mathbf{H}_R X,$$

 $\alpha \mapsto (q_{n_1} \circ \cdots \circ q_{n_r})(\delta_* \alpha).$

where $\delta: X \to X^r$ is the diagonal map, inducing a push-forward map

$$\delta_* : H^*(X, R) \to H^*(X^r, R) \simeq H^*(X, R)^{\otimes r}$$

on cohomology.

Definition 9. The operators $q_n(\alpha)$ and $q_\lambda(\alpha)$ for $\alpha \in H^*(X, R)$, $n \in \mathbb{N}$ and $\lambda \in \mathbb{P}$ are Nakajima's creation operators over X (and R).

Remark 6. For homogeneous α , the operator $q_{\lambda}(\alpha)$ is an operator of degree

$$\deg \alpha + 4\|\lambda\| + 2|\lambda| - 4$$

(and weight $2\|\lambda\|$).

Set

$$\mathbf{N}H^*(X,R) := \prod_{n=1}^{\infty} H^*(X,R)[-2n].$$

The image of $\alpha \in H^*(X, R)$ in $\mathbf{N}H^*(X, R)$ under the inclusion map corresponding to a positive natural number n is denoted by $q_n(\alpha)$, which is not to be confused with the *operator* $q_n(\alpha) \in \text{End } \mathbf{H}_R X$. Finally consider the symmetric algebra

$$N_R X := S_R(\mathbf{N}H^*(X, R)).$$

(The "N" stands for Nakajima.)

Theorem 2. The *R*-linear map

$$N_R X \to \mathbf{H}_R X, \quad q_{n_1}(\alpha_1) \dots q_{n_r}(\alpha_r) \mapsto (q_{n_1}(\alpha_1) \circ \dots \circ q_{n_r}(\alpha_r)) |\mathbf{0}\rangle$$

is an isomorphism of R-modules.

In particular, any two creation operators commute (in the super-sense, of course).

Proof. The reader may find the original proof in [12].

Example 4. It is

$$|\mathbf{1}\rangle = \exp(q_1(1))|\mathbf{0}\rangle.$$

The multiplication morphism $(N_R X)^{\otimes n} \to N_R X$ of the *R*-algebra $N_R X$ induces via the isomorphism in Nakajima's theorem a "multiplication map"

$$\mu: (\mathbf{H}_R X)^{\otimes n} \to \mathbf{H}_R X,$$

which is *not* the cup-product.

 $\langle q_m$

We can express the Poincaré pairing on the cohomology of the Hilbert schemes of points on a compact surface X in terms of the creation operators as follows:

Proposition 1. Given $\alpha_1, \ldots, \alpha_r, \beta_1, \ldots, \beta_s \in H^*(X, R)$ and natural numbers $m_1, \ldots, m_r, n_1, \ldots, n_s \in \mathbf{N}$ we do have

$$= \sum_{\substack{\sigma:\{1,\dots,s\}\\ \to \{1,\dots,r\}}} \pm \prod_{i=1}^{s} m_i \cdot \delta_{m_i,n_{\sigma(i)}} \cdot \langle \alpha_i, \beta_{\sigma(i)} \rangle$$

where the brackets denote the Poincaré pairing on $\mathbf{H}_R X$ and $H^*(X, R)$, respectively, and the sum runs over all bijections. The sign is the Koszul sign arising from reordering the α_i and β_i .

In particular, the pairing vanishes if $r \neq s$ or each index of a creation operator does not occur with the same multiplicity on both sides.

Proof. There are adjoint operators to the creation operators acting on $\mathbf{H}_R X$, called *annihilation operators*. Nakajima calculated commutators, which are fundamental for the study of the cohomology of the Hilbert schemes, between the creation and annihilation operators. From these commutation relations, one can deduce the above result. The way to do this can be found in [10] (see in particular the first paragraph in section 6).

In the sequel we need a simple fact following from this description of the Poincaré pairing. This fact is stated in the following lemma where we also assume that X is compact. Let further $\alpha_1, \ldots, \alpha_n \in H^*(X, R)$ be linearly independent cohomology classes on the surface. We choose classes β_1, \ldots, β_n such that $\langle \alpha_i, \beta_j \rangle = \delta_{ij}$ for $i, j \in$ $\{1, \ldots, n\}$. For each $r \geq 1$ we set $\check{\beta}_i^r := p^*\beta$ where $p: X^r \to X$ is the projection onto the first factor. Recall the push-forward map $\delta_*: H^*(X, R) \to H^*(X^r, R)$. We set $\alpha_i^r := \delta_*(\alpha_i)$. One verifies that $\langle \check{\beta}_i^s, \alpha_j^r \rangle \neq 0$ if i = j and s = r and zero otherwise. For each partition $\lambda = (n_1, \ldots, n_r)$, we finally set

$$\check{q}_{\lambda}(\beta_i) := (q_{n_1} \circ \cdots \circ q_{n_r})(\beta_i^r)$$

for $i \in \{1, ..., n\}$.

Lemma 2. Given partitions $\lambda_1, \ldots, \lambda_r, \mu_1, \ldots, \mu_s$ and maps $\rho : \{1, \ldots, r\} \rightarrow \{1, \ldots, n\}$ and $\sigma : \{1, \ldots, s\} \rightarrow \{1, \ldots, n\}$, the expression

$$\langle \check{q}_{\lambda_1}(\beta_{\rho(1)}) \dots \check{q}_{\lambda_r}(\beta_{\rho(r)}) | \mathbf{0} \rangle, q_{\mu_1}(\alpha_{\sigma(1)}) \dots q_{\mu_r}(\alpha_{\sigma(r)}) | \mathbf{0} \rangle \rangle$$

is non-zero if and only there is a bijection $\tau : \{1, \ldots, r\} \to \{1, \ldots, s\}$ such that $\lambda_i = \mu_{\tau(i)}$ and $\rho(i) = \sigma(\tau(i))$.

Proof. This follows from the previous proposition and the orthogonality of the α_i and β_i .

4.2. Hilbert schemes of marked complex surfaces. For each marking $\alpha \in H^*(X, R)$ on a complex surface X and each $n \in \mathbf{N}_0$, we set

$$\alpha^{[n]} := p_*(\operatorname{ch}(\mathcal{O}_{\Xi^n}) \cup q^*(\operatorname{td}(X) \cup \alpha)),$$

which is a marking on $X^{[n]}$, which depends *R*-linearly on α . The corresponding marked complex manifold is denoted by

$$(X, \alpha)^{[n]} := (X^{[n]}, \alpha^{[n]}).$$

Definition 10. The marked complex manifold $(X, \alpha)^{[n]}$ is the Hilbert scheme of n points on (X, α) .

Again, it is a good idea to consider all these spaces at once, so we set

$$\operatorname{Hilb}(X,\alpha) := \sum_{l=0}^{\infty} (X,\alpha)^{[n]}.$$

The function $Hilb(\cdot)$ works like an exponential. By this, we mean the following:

Proposition 2. Let (X, α) and (Y, β) be two marked complex surfaces. Then

$$\operatorname{Hilb}(\emptyset, 0) = (*, 0)$$

and

$$\operatorname{Hilb}((X, \alpha) + (Y, \beta)) = \operatorname{Hilb}(X, \alpha) \cdot \operatorname{Hilb}(Y, \beta).$$

Proof. From the fact that a zero-dimensional subspace of $X \amalg Y$ of length n is given by a pair of a zero-dimensional subspace of X of length p and a zero-dimensional subspace of Y of length q with p + q = n, one proves that

$$(X \amalg Y)^{[n]} = \prod_{p+q=n} X^{[p]} \times Y^{[q]},$$

see e.g. [4]. Furthermore, the universal family over $(X \amalg Y)^{[n]}$ restricted to the component $X^{[p]} \times Y^{[q]}$ is given by

$$(\Xi^p \times Y^{[q]}) \amalg (X^{[p]} \times \Upsilon^q) \subset (X^{[p]} \times X \times Y^{[q]}) \amalg (X^{[p]} \times Y^{[q]} \times Y),$$

where Ξ^p is the universal family over $X^{[p]}$ and Υ^q is the universal family over $Y^{[q]}$. Thus by definition of $(\alpha + \beta)^{[n]}$, its restriction onto the component $X^{[p]} \times Y^{[q]}$ is given by

$$\alpha^{[p]} \boxtimes 1 + 1 \boxtimes \beta^{[q]} \in H^*(X^{[p]} \times Y^{[q]}, R).$$

Summing all up, this yields the claimed formula.

Let $n \in \mathbf{N}_0$. Given a marked complex surface (X, α) , we may consider the *n*-fold sum $n(X, \alpha)$, which is again a marked surface. The previous proposition tells us that

$$\operatorname{Hilb}(n(X,\alpha)) = \operatorname{Hilb}(X,\alpha)^n.$$

In particular, we have

$$\mathbf{H}_R(n(X,\alpha)) = (\mathbf{H}_R(X,\alpha))^{\otimes n}.$$

Recall the *R*-linear map $\mu : (\mathbf{H}_R X)^{\otimes n} \to \mathbf{H}_R X$ given by Nakajima's description of the cohomology space in terms of the creation operators applied to the vacuum. This give us thus an *R*-linear map

$$\mu: \mathbf{H}_R(n(X,\alpha)) \to \mathbf{H}_R(X,\alpha).$$

We shall need this map in the next section.

5. Universal formulas

Consider the **Q**-algebra

$$A := \mathbf{Q}[K, e, \alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4]$$

with deg K = 2, deg e = 4 and deg $\alpha_i = i$. Let $A^{\geq 5}$ be the ideal generated by the elements of degree five or more in A. We set

$$V_R := R \otimes_{\mathbf{Q}} (A/A^{\geq 5}).$$

The element $K \in V_R$ is called the *universal canonical class*, the element $e \in V_R$ is called the *universal Euler class*.

Definition 11. The *R*-algebra V_R is the universal cohomology ring for marked complex surfaces over *R*.

This is due to the following: Given a marked complex surface (X, α) , there is a unique homomorphism

$$V_R \to H^*(X, R), \quad \gamma \mapsto \gamma|_{(X,\alpha)}$$

of R-algebras with

$$K|_{(X,\alpha)} = -c_1(\Theta_X), \quad e|_{(X,\alpha)} = c_2(\Theta_X), \quad \text{and} \quad \alpha_i|_{(X,\alpha)} = \alpha_i.$$

Consider a **Q**-algebra S. A (virtual) marked compact complex surface (X, α) in the S-algebra of marked complex surfaces is called *versal* if the homomorphism

$$V_R \to H^*(X, R), \quad \mapsto \gamma|_{(X,\alpha)}$$

is a monomorphism. There are versal marked surfaces. They can be constructed by taking a sufficiently general linear combination of copies of \mathbf{P}^2 , $\mathbf{P}^1 \times \mathbf{P}^1$, $\mathbf{C}^2/\mathbf{Z}^4$ and $\mathbf{P}^1 \times \mathbf{C}/\mathbf{Z}^2$ with sufficiently general markings.

Lemma 3. Let $\gamma, \gamma' \in V_R$ with $\gamma|_{(X,\alpha)} = \gamma'|_{(X,\alpha)}$ for all marked complex surfaces (X, α) . Then $\gamma = \gamma'$.

Proof. This follows by evaluating γ and γ' on a versal marked surface.

Consider the R-algebra

$$\mathbf{P}V_R := \prod_{\lambda \in \mathbf{P}} V_R[-2\|\lambda\|].$$

The canonical injection from $V \to \mathbf{P}V_R$ corresponding to the factor indexed by a partition $\lambda \in \mathbf{P}$ is denoted by

$$V_R \to \mathbf{P} V_R, \quad v \mapsto q_\lambda(v).$$

An element in $\mathbf{P}V_R$ is called an element of weight $l, l \in \mathbf{N}_0$, if it is a (finite) sum of elements of the form $q_{\lambda}(v)$ with $\|\lambda\| = l$.

We set

$$Q_R := S_R(\mathbf{P}V_R).$$

Note that $\mathbf{P}V_R$ becomes a subspace of Q_R .

Definition 12. The *R*-algebra Q_R is the universal creation operator algebra of the Hilbert schemes of points of marked surfaces over *R*.

The algebra Q_R is universal in the following sense: Given a marked complex surface (X, α) , there is a unique homomorphism of *R*-algebras

$$Q_R \to \operatorname{End}(\mathbf{H}_R X), \quad q \mapsto q|_{(X,\alpha)}$$

mapping $q_{\lambda}(v)$ for $v \in V_R$ and $\lambda \in \mathbf{P}$ to $q_{\lambda}(v|_{(X,\alpha)})$.

Lemma 4. Let $q, q' \in Q_R$ with $q|_{(X,\alpha)} = q'|_{(X,\alpha)}$ for all marked complex surfaces (X, α) . Then q = q'.

Proof. We may assume that X is a versal surface. Then $(\alpha_0, \ldots, \alpha_4, 1, K, e) \in V_R$ maps to a linearly independent system on $H^*(X, R)$. We choose a "dual system" $(\beta_0, \ldots, \beta_7)$ as in lemma 2 to this system. Applying this lemma enables us to extract the components of q and q' that have to occur with the same coefficients. From this, we conclude q = q'.

Recall the definition of q(n) given in the section about notations.

Lemma 5. Let (X, α) be a marked complex surface and $n \in \mathbf{N}_0$ a natural number. Then

$$\mu \circ q|_{n(X,\alpha)} = q(n)|_{(X,\alpha)} \circ \mu$$

as operators from $(\mathbf{H}_R X)^{\otimes n}$ to $\mathbf{H}_R X$ for all $q \in Q_R$.

Proof. By linearity, we may assume that

$$q = q_{\lambda_1}(v_1) \dots q_{\lambda_r}(v_r)$$

for partitions $\lambda_1, \ldots, \lambda_r \in \mathbf{P}$ and elements $v_1, \ldots, v_r \in V_R$. Then

$$\mu(q_{\lambda_1}(v_1)\dots q_{\lambda_r}(v_r)|_{n(X,\alpha)}|\mathbf{0}\rangle) = \mu(q_{\lambda_1}(n\cdot\beta_1)\dots q_{\lambda_r}(n\cdot\beta_r)|\mathbf{0}\rangle),$$

with $\beta_i := v_i|_{(X,\alpha)}$ and where

$$n \cdot \beta_i := \sum_{j=1}^n (i_j)_* \beta \quad \in H^*(n(X,\alpha),R) \simeq H^*(X,\alpha)^{\oplus n}$$

where $i_j : H^*(X, R) \to H^*(X, R)^{\oplus n}$ is the injection map into the *j*-th summand. Of course, $|\mathbf{0}\rangle$ is the vacuum vector in $\mathbf{H}_{n(X,\alpha)}$.

The operator μ "commutes" with the q_{λ} in the sense that

$$\mu \circ q_{\lambda_i}(n \cdot \beta_i) = n \cdot q_{\lambda_i}(\beta_i) \circ \mu$$

(This follows from

$$q_l(n \cdot \beta_i)(\epsilon_1 \otimes \cdots \otimes \epsilon_n) = \sum_{i=1}^n \epsilon_1 \otimes \cdots \otimes \epsilon_{i-1} \otimes q_l(\beta_i)(\epsilon_i) \otimes \epsilon_{i+1} \otimes \cdots \otimes \epsilon_n$$

for $l \in \mathbf{N}$ and $\epsilon_1 \otimes \cdots \otimes \epsilon_n \in \mathbf{H}_R(n \cdot (X, \alpha)) \simeq (\mathbf{H}_R X)^{\otimes n}$.) This gives

$$\mu(q_{\lambda_1}(n \cdot \beta_1) \dots q_{\lambda_r}(n \cdot \beta_r) |\mathbf{0}\rangle) = n^r(q_{\lambda_1}(\beta_1) \dots q_{\lambda_r}(\beta_r))(\mu(|\mathbf{0}\rangle))$$
$$= n^r(q_{\lambda_1}(v_1) \dots q_{\lambda_r}(v_r))|_{(X,\alpha)}(\mu(|\mathbf{0}\rangle)).$$

Together with the fact that acting on $|0\rangle$ is faithful, this yields

$$\mu \circ q_{\lambda_1}(v_1) \dots q_{\lambda_r}(v_r)|_{n(X,\alpha)} = n^r (q_{\lambda_1}(v_1) \dots q_{\lambda_r}(v_r))|_{(X,\alpha)} \circ \mu.$$

Lemma 6. Let $q \in Q_R$ with

 $q_{(X,\alpha)+(X',\alpha')} = q|_{(X,\alpha)} \otimes \mathrm{id} + \mathrm{id} \otimes q|_{(X',\alpha')}$

in End $\mathbf{H}_R((X, \alpha) + (X', \alpha')) = \text{End}(\mathbf{H}_R X \otimes \mathbf{H}_R X')$ for two marked complex surfaces. Then $q \in \mathbf{P}V_R \subset Q_R$.

Proof. For each natural number $n \in \mathbf{N}$ and each complex surface (X, α) , we have

$$nq|_{(X,\alpha)} \circ \mu = \mu \circ q|_{n(X,\alpha)} = q(n)|_{(X,\alpha)} \circ \mu$$

as maps from $(\mathbf{H}_R X)^{\otimes n} \to \mathbf{H}_R(X)$, where the first equality is due to the assumption and the second due to the previous lemma. It follows that $nq|_{(X,\alpha)} = q(n)|_{(X,\alpha)}$ and thus nq = q(n) for all $n \in \mathbf{N}$. This means that q(n) is linear in n, i.e. $q \in \mathbf{P}V_R$. \Box **Theorem 3.** For each characteristic class γ of marked complex manifolds, there exists a unique class $q(\gamma) \in Q_R$ such that

$$q(\gamma)|_{(X,\alpha)}|\mathbf{1}\rangle = \gamma|_{\mathrm{Hilb}(X,\alpha)}$$

for all marked complex surfaces (X, α) .

Proof. In [2], it is stated that each polynomial π in the Chern classes of tautological sheaves $u^{[\cdot]}$ of the Hilbert schemes of points on a surfaces can be expressed by a "universal formula". In our language, this means that there exists a $q \in Q_R$ such that

$$\sum_{n \ge 0} \pi(u^{[n]}) = q|_{(X, ch(u))}|\mathbf{0}|$$

for all complex surfaces X and each "sheaf" $u \in K(X)$. (In fact, the results in [2] are only stated for algebraic surfaces, however, the result generalises easily.)

We may generalise this result even further: The theorem in [2] remains true if we substitute ch(u) by an arbitrary class $\alpha \in H^*(X, R)$ (however, one has to start to keep track of the "Koszul signs" arising), i.e. for each polynomial π in the cohomology classes $\alpha^{[\cdot]}$, there exists a $q \in Q_R$ such that

$$\sum_{n\geq 0} \pi(\alpha^{[n]}) = q|_{(X,\alpha)} |\mathbf{0}\rangle$$

for each marked complex surface (X, α) :

An analogous statement holds true for polynomials in the Chern classes of the tangent sheaves of the Hilbert schemes of points on a surface, which is also proven in [2]. Thus, everything holds true for "mixed" polynomials in the $\alpha^{[\cdot]}$ and the Chern classes of the tangent sheaves, and thus, via completion, for γ .

This proves that there exists a $q(\gamma)$ with

$$q(\gamma)|_{(X,\alpha)}|\mathbf{1}\rangle = \gamma_{\mathrm{Hilb}(X,\alpha)}.$$

(Recall that $|\mathbf{1}\rangle = \exp(q_1(1))|\mathbf{0}\rangle$.) The uniqueness of $q(\gamma)$ follows from lemma 4. \Box

Corollary 1. Consider the universal Chern character $ch \in U_R$ and the universal marking $\alpha \in U_R$. It is $q(ch) \in \mathbf{P}V_R \subset Q_R$ and $q(\alpha) \in \mathbf{P}V_R \subset Q_R$.

Proof. We apply $q(ch)|_{(X,\alpha)+(X',\alpha')}$ to the unit, which gives:

$$\begin{aligned} q(\mathrm{ch})|_{(X,\alpha)+(X',\alpha')}|\mathbf{1}\rangle &= \mathrm{ch}(\mathrm{Hilb}((X,\alpha)+(X',\alpha'))) \\ &= \mathrm{ch}(\mathrm{Hilb}(X,\alpha)\cdot\mathrm{Hilb}(X',\alpha')) \\ &= \mathrm{ch}(\mathrm{Hilb}(X,\alpha))\otimes|\mathbf{1}\rangle + |\mathbf{1}\rangle\otimes\mathrm{ch}(\mathrm{Hilb}(X',\alpha')) \\ &= \left(q(\mathrm{ch})|_{(X,\alpha)}\otimes\mathrm{id} + \mathrm{id}\otimes q(\mathrm{ch})|_{(X',\alpha')}\right)|\mathbf{1}\rangle. \end{aligned}$$

This proves the statement for q(ch) by the previous lemma.

The proof of the statement for $q(\alpha)$ is virtually the same, so we shall omit it. \Box

Corollary 2. For each multiplicative characteristic class γ of marked complex manifolds, there exists a unique class $\log q(\gamma) \in \mathbf{P}V_R$ such that

$$\exp(\log q(\gamma)) = q(\gamma)$$

in Q_R .

Proof. It is

$$q(\gamma)|_{(\emptyset,0)}|\mathbf{1}\rangle = \gamma|_{\mathrm{Hilb}(\emptyset,0)} = \gamma|_{(*,0)} = 1,$$

thus $q(\gamma) \in 1 + (\mathbf{P}V_R) \cdot Q_R \subset Q_R$ Therefore, there exists a $\log q(\gamma) \in Q_R$ with $\exp(\log q(\gamma)) = q(\gamma)$.

It remains to show that $\log q(\gamma) \in \mathbf{P}V_R \subset Q_R$. This follows from the previous lemma since

$$\log q(\gamma)|_{(X,\alpha)+(X',\alpha')} = \log q(\gamma)|_{(X,\alpha)} \otimes \mathrm{id} + \mathrm{id} \otimes \log q(\gamma)|_{(X',\alpha')}$$

This in turn is due to the fact that

$$q(\gamma)|_{(X,\alpha)+(X',\alpha')}|\mathbf{1}\rangle = \gamma|_{\mathrm{Hilb}((X,\alpha)+(X',\alpha'))}$$

= $\gamma|_{\mathrm{Hilb}(X,\alpha)\cdot\mathrm{Hilb}(X',\alpha')}$
= $\gamma|_{\mathrm{Hilb}(X,\alpha)} \otimes \gamma|_{\mathrm{Hilb}(X',\alpha')}$
= $q(\gamma)|_{(X,\alpha)}|\mathbf{1}\rangle \otimes q(\gamma)|_{(X',\alpha')}|\mathbf{1}\rangle$
= $(q(\gamma)|_{(X,\alpha)} \otimes q(\gamma)|_{(X',\alpha')})|\mathbf{1}\rangle$

for marked complex surfaces (X, α) and (X', α') .

Let $L: V_R \to V_R$ be the homomorphism of *R*-algebras that maps *K* to *K*, *e* to e, α_{2i} to $\frac{(\alpha_2)^i}{i!}$ for $i \in \{0, 1, 2\}$ and α_1 and α_3 to zero. In particular

$$L(\alpha) = \exp(\alpha_2)$$

with $\alpha = \sum_{i=0}^{4} \alpha_i$. This homomorphism induces homomorphisms of $\mathbf{P}V_R$ and Q_R , which will also be denoted by L.

Proposition 3. Using the homomorphism L, we can formulate the following partial result on $q(c') \in Q_R$, namely

$$L(\log q(c')) = \sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{m} q_m (1+\alpha_2) - q_1(1).$$

Proof. It is

$$L(\log q(c'))|_{(X,\alpha)}|\mathbf{1}\rangle = \log q(c')|_{(X,\exp\delta)}|\mathbf{1}\rangle = \sum_{n\geq 0} c'|_{(X,\alpha)^{[n]}}$$

with $\delta := \exp(\alpha_2)$. We may assume that α_2 is algebraic, i.e. $\alpha_2 = c_1(\mathcal{L})$ for a invertible sheaf \mathcal{L} (from which $\delta = \operatorname{ch}(\mathcal{L})$ follows). Then the proposition follows from Lehn's theorem

$$\sum_{n\geq 0} c(\mathcal{L}^{[n]}) = \exp\left(\sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{m} q_m(c_1(\mathcal{L}))\right) |\mathbf{0}\rangle.$$

given in [9] as $c(\mathcal{L}^{[n]}) = c'|(\alpha^{[n]}).$

Corollary 3. Given a marked complex surface (X, α) with $\alpha = \exp(\delta)$ for a class $\delta \in H^2(X, R)$, we have

$$c'|_{\mathrm{Hilb}(X,\alpha)} = \exp\left(\sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{m} q_m (1+\delta)\right) |\mathbf{0}\rangle.$$

6. The Hilbert scheme of the Affine plane

Let $\gamma \in V_R$ be a multiplicative characteristic class, i.e. $\gamma = \prod_{i=1}^{\infty} f(\lambda_i)$ for a uniquely determined power series $f \in 1 + xR[[x]]$.

We define a power series $g \in tR[[t]]$ by

$$\frac{\partial g}{\partial t}\left(\frac{x}{f(x)f(-x)}\right) = f(x)f(-x),$$

i.e. $t\frac{\partial g(t)}{\partial t}$ is the compositional inverse of the power series $\frac{x}{f(x)f(-x)}$. In particular, g is an odd power series.

Let $A: V_R \to V_R$ be the homomorphism of R-algebras that maps K and e to zero and α to 1. The R-linear map $\mathbf{P}V_R \to \mathbf{P}V_R$ that maps $q_{\lambda}(v)$ to $q_{\lambda}(A(v))$ for $|\lambda| = 1$ and $q_{\lambda}(v)$ to zero for $|\lambda| > 1$ will also be denoted by A. This map in turn induces a homomorphism of R-algebra $A: Q_R \to Q_R$.

Theorem 4. Using the homomorphism A, the following partial result on $q(\gamma) \in Q_R$ holds, namely

$$A(\log q(\gamma)) = \sum_{k\geq 1}^{\infty} \left(\frac{1}{k!} \left. \frac{\partial^k}{\partial t^k} \right|_{t=0} g \right) \cdot q_k(1) - q_1(1).$$

Proof. The proof follows the strategy also used in [2]. First of all note that

$$A(q(\gamma))|_{X,\alpha}|\mathbf{1}\rangle = q(\gamma)|_{(\mathbf{C}^2,0)}|\mathbf{1}\rangle = \sum_{n\geq 0} \gamma|_{(\mathbf{C}^2,0)^{[n]}}.$$

Thus the proof of the theorem reduces to calculating multiplicative classes on the Hilbert schemes of points on the affine plane. By corollary 2, the right hand side can be written as

$$\sum_{n\geq 0} \gamma|_{(\mathbf{C}^2,0)^{[n]}} = \exp\left(\sum_{k\geq 1} g_k q_k(1)\right) |\mathbf{0}\rangle$$

for certain $g_k \in R$. The reader may wonder why no $q_{\lambda}(1)$ with $|\lambda| \geq 2$ appears on the right hand side, but these $q_{\lambda}(1)$ act trivially on $\mathbf{H}_{\mathbf{C}^2}$ as $\delta_* : H^*(\mathbf{C}^2, R) \to H^*((\mathbf{C}^2)^r, R)$ is the zero map for $r \geq 2$.

Thus it remains to calculate the g_k that appear above, in fact to prove that $g = \sum_{k\geq 1} g_k t^k$. Restricting the previous formula to the classes of (maximal) degree n-1 on each Hilbert $(\mathbf{C}^2)^{[n]}$ scheme yields

$$\sum_{k\geq 1} g_k q_k(1) |\mathbf{0}\rangle = \sum_{n\geq 0} (\gamma_{n-1})|_{(\mathbf{C}^2, 0)^{[n]}}$$

where γ_{n-1} is the component of γ of degree n-1. As explained in [2], this follows from the fact that $H^r((\mathbf{C}^2)^{[n]}, R) = 0$ for $r \ge n$.

Looking in [1], we find the following formula in the proof of theorem 5.1:

$$c_{n-1}|_{(\mathbf{C}^{2},0)^{[n]}} = \sum_{\|\lambda\|=n} \frac{1}{h(\lambda)} \left(\frac{1}{(n-1)!} \frac{\partial^{n-1}}{\partial x^{n-1}} \bigg|_{x=0} \prod_{w \in D(\lambda)} ((1+h(w)x)(1-h(w)x)) \right) \\ \cdot \chi^{\lambda}_{(n)} \cdot z^{-1}_{(n)} \cdot q_{n}(1) |\mathbf{0}\rangle_{2}$$

where the symbols $D(\lambda)$, $h(\lambda)$, h(x), χ^{λ}_{μ} and z_{μ} are defined in [1] (or [2]). (This formula follows from theorem 4.2 in [2].)

Slightly generalising this formula by replacing the Chern class c_{n-1} by γ_{n-1} , which amounts to replace the power series 1 + x by the more general f, yields

$$\gamma_{n-1}|_{(\mathbf{C}^{2},0)^{[n]}} = \sum_{\|\lambda\|=n} \frac{1}{h(\lambda)} \left(\frac{1}{(n-1)!} \frac{\partial^{n-1}}{\partial x^{n-1}} \bigg|_{x=0} \prod_{w \in D(\lambda)} (f(h(w)x)f(-h(w)x)) \right) \cdot \chi_{(n)}^{\lambda} \cdot z_{(n)}^{-1} \cdot q_{n}(1) |\mathbf{0}\rangle.$$

(Already in [2] is has been stated that similar formulas hold true for more than just the total Chern class.)

We proceed as in [2], noting that

$$\chi_{(n)}^{\lambda} = \begin{cases} (-1)^{s} & \text{for } \lambda = (n - s, \underbrace{1, \dots, 1}_{s-\text{times}}) \text{ and } 0 \le s < n \\ 0 & \text{otherwise} \end{cases}$$

and that $z_{(n)} = n$, $\{h(w) : w \in D(\lambda)\} = \{1, \ldots, s, 1, \ldots, n - s - 1, n\}$ (with multiplicities) and $h(\lambda) = s! \cdot (n - s - 1)!n$ for a partition λ of n the form $\lambda = (n - s, 1, \ldots, 1), 0 \le s < n$. This simplifies the above expression for $\gamma_{n-1}|_{(\mathbf{C}^2, 0)}$ to

$$\begin{split} \gamma_{n-1}|_{(\mathbf{C}^{2},0)^{[n]}} &= \frac{q_{n}(1)}{n} |\mathbf{0}\rangle \cdot \sum_{s=0}^{n-1} \frac{(-1)^{s}}{n \cdot s!(n-1-s)!} \\ &\cdot \left(\frac{1}{(n-1)!} \frac{\partial^{n-1}}{\partial x^{n-1}} \bigg|_{x=0} \prod_{\substack{k \in \{1,\dots,s,\\1,\dots,n-1-s,n\}}} (f(k \cdot x) \cdot f(-k \cdot x)) \right) \\ &= \frac{q_{n}(1)}{n} |\mathbf{0}\rangle \cdot \frac{1}{(n-1)!} \frac{\partial^{n-1}}{\partial x^{n-1}} \bigg|_{x=0} f(n \cdot x) \cdot f(-n \cdot x) \\ &\cdot \sum_{s=0}^{n-1} \frac{(-1)^{s}}{n \cdot s!(n-1-s)!} \prod_{k=-(n-s)}^{s} (f(k \cdot x)f(-k \cdot x)). \end{split}$$

By lemma 8 (to be found in the appendix) this gives

$$\gamma_{n-1}|_{(\mathbf{C}^{2},0)^{[n]}} = \frac{q_{n}(1)}{n} |\mathbf{0}\rangle \cdot \frac{(-1)^{n-1}}{n!} \frac{\partial^{n-1}}{\partial x^{n-1}} \bigg|_{x=0} (f(x) \cdot f(-x))^{n}$$

As $f(x) \cdot f(-x)$ is an even power series, we can leave out the sign $(-1)^{n-1}$. Together with the Langrange Inversion Theorem, see proposition 4 in the appendix, this yields

$$\sum_{n\geq 0} \gamma_{n-1}|_{(\mathbf{C}^2,0)^{[n]}} = \sum_{k\geq 1} \left(\left. \frac{1}{k!} \frac{\partial^k}{\partial t^k} \right|_{t=0} g \right) \cdot q_k(1) |\mathbf{0}\rangle,$$

which is all that remained to show.

Corollary 4. Consider the marked complex surface $(\mathbf{C}^2, 0)$. We have

$$\gamma|_{\mathrm{Hilb}(\mathbf{C}^{2},0)} = \exp\left(\sum_{k\geq 1}^{\infty} \left(\frac{1}{k!} \left.\frac{\partial^{k}}{\partial t^{k}}\right|_{t=0} g(t)\right) \cdot q_{k}(1)\right) |\mathbf{0}\rangle.$$

Example 5. Let $\gamma = c$ be the total Chern class. Then f = 1 + x and thus

$$g = \frac{1 - \sqrt{1 + 4t^2}}{2t} + \arcsin(2t) = \sum_{n=0}^{\infty} (-1)^n \frac{1}{(n+1) \cdot (2n+1)} \cdot \binom{2n}{n} \cdot t^{2n+1}.$$

By the previous theorem, this gives

$$A(\log q(c)) = \sum_{n \ge 0} \frac{(-1)^n}{n+1} \cdot \binom{2n}{n} \cdot \frac{q_{2n+1}(1)}{2n+1} - q_1(1).$$

This formula is one of the main results in [1].

Example 6. Let $\gamma = s = c^{-1}$ be the total Segre class, i.e. $f = (1+x)^{-1}$. Thus $\frac{\partial g}{\partial t}(x \cdot (1-x^2)) = (1-x^2)^{-1}.$ By the Lagrange Inversion Theorem, we have

$$g = \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \cdot \binom{3n}{n} \cdot t^{2n+1},$$

and therefore

$$A(\log q(s)) = \sum_{n \ge 0} \frac{1}{2n+1} \cdot \binom{3n}{n} \cdot \frac{q_{2n+1}(1)}{2n+1} - q_1(1).$$

This example is new.

Example 7. Let $\gamma = \sqrt{\text{td}}$ be the square root of the Todd class, i.e. $f = \sqrt{\frac{x}{1 - \exp(-x)}}$. Then

$$g = 2\operatorname{Shi}\left(\frac{t}{2}\right) = \sum_{n=0}^{\infty} \frac{1}{4^n \cdot (2n+1) \cdot (2n+1)!} \cdot t^{2n+1},$$

where $\operatorname{Shi}(y) := \int_0^y \frac{\sinh \eta}{\eta} d\eta$ is the hyperbolic sine integral. Therefore

$$A(\log q(\sqrt{\mathrm{td}})) = \sum_{n \ge 0} \frac{1}{4^n \cdot (2n+1)!} \cdot \frac{q_{2n+1}(1)}{2n+1} - q_1(1)$$

This example is new. (The calculation for td instead of \sqrt{td} leads to a much more complicated result.)

Remark 7. As every product of Chern classes occurs in a suitable multiplicative class (e.g. $\gamma = \prod_{i,j=1}^{\infty} (1 + \lambda_i t_j) \in U_R$ with $R = \mathbf{Q}[t_1, t_2, \ldots]$), we can use the above corollary to calculate any characteristic class of the Hilbert schemes of points on the affine plane in terms of Nakajima's creation operators.

We turn now from the tangent bundle to the multiplicative classes built up from the b_i : Let $\gamma \in V_R$ be a multiplicative characteristic class of the form $\gamma = \prod_{i=1}^{\infty} f(\lambda'_i)$ for a (uniquely determined) power series $f \in 1 + xR[[x]]$.

We define a power series $g \in tR[[t]]$ by

$$\frac{\partial g}{\partial t}\left(\frac{x}{f(-x)}\right) = f(-x),$$

i.e. $-t\frac{\partial g}{\partial t}$ is the compositional inverse of the power series $-\frac{x}{f(x)}$.

Theorem 5. Using the homomorphism A, the following partial result on $q(\gamma) \in Q_R$ holds, namely

$$A(\log q(\gamma)) = \sum_{k\geq 1}^{\infty} \left(\frac{1}{k!} \left. \frac{\partial^k}{\partial t^k} \right|_{t=0} g(t) \right) \cdot q_k(1) - q_1(1).$$

Proof. We have

$$A(q(\gamma))|_{(X,\alpha)}|\mathbf{1}\rangle = q(\gamma)|_{(\mathbf{C}^2,1)}|\mathbf{1}\rangle = \sum_{n\geq 0} \gamma|_{(\mathbf{C}^2,1)^{[n]}}.$$

Thus again, everything boils down to calculating universal classes on the Hilbert schemes of points on the affine plane. As in the proof of the previous theorem, we have to show that

$$\sum_{k\geq 1} g_k q_k(1) |\mathbf{0}\rangle = \sum_{n\geq 0} \gamma_{n-1}|_{(\mathbf{C}^2, 1)^{[n]}}$$

with $g = \sum_{k\geq 1} g_k t^k$. The right hand side of the equation is in fact a sum of characteristic classes of the tautological bundles $\mathcal{O}^{[n]}$ as $ch(\mathcal{O}^{[n]}) = 1^{[n]}$ (note that $ch(\mathcal{O}) = 1$). To be more precise, it is

$$\gamma_{n-1}|_{(\mathbf{C}^2,1)^{[n]}} = \frac{1}{(n-1)!} \frac{\partial^{n-1}}{\partial x^{n-1}} \bigg|_{x=0} \prod_{i=1}^n f(\epsilon_i^{[n]} \cdot x),$$

where $\epsilon_1^{[n]}, \ldots, \epsilon_n^{[n]}$ are the Chern roots of $\mathcal{O}^{[n]}$.

From here, we continue as in the proof of the previous theorem. We just have to replace the tangent bundle by the tautological bundle. By [1], we have

$$\gamma_{n-1}|_{(\mathbf{C}^2,1)^{[n]}} = \sum_{\|\lambda\|=n} \frac{1}{h(\lambda)} \left(\frac{1}{(n-1)!} \frac{\partial^{n-1}}{\partial x^{n-1}} \bigg|_{x=0} \prod_{w \in D(\lambda)} f(h(w)x) \right) \cdot \chi^{\lambda}_{(n)} \cdot z^{-1}_{(n)} \cdot q_n(1) |\mathbf{0}\rangle,$$

where the symbols have the same meaning as in the proof of the previous theorem. (Again, we have replaced the total Chern class by an arbitrary multiplicative class.) Doing the same calculations as above, one eventually arrives at

$$\gamma_{n-1}|_{(\mathbf{C}^{2},1)^{[n]}} = \frac{q_{n}(1)}{n}|\mathbf{0}\rangle \cdot \frac{(-1)^{n-1}}{n!} \frac{\partial^{n-1}}{\partial x^{n-1}}\Big|_{x=0} (f(x))^{n}.$$

Again by the Langrange Inversion Theorem this yields

$$\sum_{n\geq 0} \gamma_{n-1}|_{(\mathbf{C}^2,1)^{[n]}} = \sum_{k\geq 1} \left(\left. \frac{1}{k!} \frac{\partial^k}{\partial t^k} \right|_{t=0} g \right) \cdot q_k(1) |\mathbf{0}\rangle,$$

which again is all that remained to show.

Corollary 5. Consider the marked complex surface $(\mathbf{C}^2, 1)$. We have

$$\gamma|_{\mathrm{Hilb}(\mathbf{C}^2,1)} = \exp\left(\sum_{k\geq 1}^{\infty} \left(\frac{1}{k!} \left. \frac{\partial^k}{\partial t^k} \right|_{t=0} g\right) \cdot q_k(1) \right) |\mathbf{0}\rangle.$$

Remark 8. From this result, one can deduce a formula for

$$\gamma|_{\mathrm{Hilb}(\mathbf{C}^2,r)} = \sum_{n \ge 0} \gamma|_{(X^{[n]},r^{[n]})}$$

for each $r \in \mathbf{Z}$, as $\gamma(X^{[n]}, r^{[n]}) = \gamma(X^{[n]}, 1^{[n]})^r$. From this one can conclude that this formula holds for all $r \in \mathbb{R}$.

We do this in the following example:

Example 8. Let $r \in \mathbf{Z}$ and let $\gamma = c'^r$, i.e. $f = (1+x)^r$. Then

$$\frac{\partial g}{\partial t}(x \cdot (1-x)^{-r}) = (1-x)^r.$$

By the Lagrange Inversion Theorem,

$$g = \sum_{n \ge 0} \frac{(-1)^{n-1}}{n^2} \cdot \binom{rn}{n-1} t^n.$$

By the theorem, this gives

$$A(\log q((c')^r)) = \sum_{n \ge 0} \frac{(-1)^{n-1}}{n} \cdot \binom{rn}{n-1} \cdot \frac{q_n(1)}{n} - q_1(1)$$

and by the corollary,

$$c'|_{(\mathbf{C}^2,r)} = \gamma|_{(\mathbf{C}^2,1)} = \exp\left(\sum_{n\geq 1} \frac{(-1)^{n-1}}{n} \cdot \binom{rn}{n-1} \cdot \frac{q_n(1)}{n}\right) |\mathbf{0}\rangle,$$

which is an instance of Lehn's formula for r = 1, i.e. corollary 3 applied to $\delta = 0$. . The case $r \ge 2$ appears to be new.

Remark 9. One can use the previous corollary to obtain a complete description of the ring structure of the cohomology ring of the Hilbert schemes of points on the affine plane. (This has be studied successfully before, see [11].)

To understand this, consider the power series

$$g = \sum_{k \ge 1} (\rho_k + \rho'_k) \cdot t^k \in tR[t]$$

with $R := \mathbf{Q}[\rho_1, \rho_2, \dots, \rho'_1, \rho'_2, \dots]$. There is a unique power series $f \in 1 + xR[[x]]$ such that

$$\frac{\partial g}{\partial t}\left(\frac{x}{f(-x)}\right) = f(-x).$$

Let $\gamma := \prod_{i=1}^{\infty} f(\lambda'_i)$ be the multiplicative class defined by f. Then

$$\gamma|_{\mathrm{Hilb}\,(X,\alpha)} = \exp\left(\sum_{k\geq 1} (\rho_k + \rho'_k) \cdot q_k(1)\right) |\mathbf{0}\rangle,$$

i.e. $\gamma|_{(X,\alpha)}$ can be seen as a *universal cohomology class in* $\mathbf{H}_{\mathbf{C}^2}$ as every class in $\mathbf{H}_{\mathbf{C}^2}$ can be written as a sum of polynomials in the $q_k(1)$ acting on the vacuum $|\mathbf{0}\rangle$.

Now, γ^2 is another multiplicative class, which is defined by f^2 and which applied to Hilb (X, α) gives

$$\gamma^{2}|_{\mathrm{Hilb}\,(X,\alpha)} = \exp\left(\sum_{k\geq 1} (\rho_{k} + \rho_{k}') \cdot q_{k}(1)\right) |\mathbf{0}\rangle \cdot \exp\left(\sum_{k\geq 1} (\rho_{k} + \rho_{k}') \cdot q_{k}(1)\right) |\mathbf{0}\rangle,$$

which includes all cup-products of all polynomials in the $q_k(1)$ acting on the vacuum.

Now let $h \in tR[[t]]$ be the unique power series with

$$\frac{\partial h}{\partial t}\left(\frac{x}{f^2(-x)}\right) = f^2(-x).$$

Then

$$\left(\exp\left(\sum_{k\geq 1}(\rho_k+\rho'_k)\cdot q_k(1)\right)|\mathbf{0}\rangle\right)^2 = \exp\left(\sum_{k\geq 1}\left(\frac{1}{k!}\frac{\partial^k}{\partial t^k}\Big|_{t=0}h(t)\right)\cdot q_k(1)\right)|\mathbf{0}\rangle.$$

Thus, we can effectively calculate all powers of the universal cohomology class, which encodes completely the ring structure.

APPENDIX A. LAGRANGE INVERSION THEOREM

For the convenience of the reader, we state the Langrange Inversion Theorem (also called Lagrange expansion) in the form in which it is needed in the main text. For a proof of the general statement see e.g. [14].

Let $f \in 1 + xR[[x]]$ be a power series and define $g \in tR[[t]]$ such that

$$\frac{\partial g}{\partial t}\left(\frac{x}{f}\right) = f,$$

i.e. such that $t\frac{\partial q}{\partial t}$ is the compositional inverse of $\frac{x}{t}$.

Proposition 4. The power series g is given by

$$g = \sum_{n=1}^{\infty} \frac{t^n}{n \cdot n!} \left(\frac{\partial^{n-1}}{\partial x^{n-1}} \bigg|_{x=0} f^n \right).$$

Appendix B. Two algebraic identities

The following identity holds true in the domain of the rationals for all $m \in \mathbf{N}_0$ and $0 \le p \le m$:

Lemma 7. It is

$$\sum_{s=0}^{m} \frac{(-1)^s s^p}{s!(m-s)!} = \begin{cases} 0 & \text{for } p < m\\ (-1)^m & \text{for } p = m. \end{cases}$$

Proof. Observe that

$$\begin{split} \sum_{s=0}^{m} \frac{(-1)^{s} s^{p}}{s!(m-s)!} &= \left. \sum_{s=0}^{m} \frac{(-x)^{s} s^{p}}{s!(m-s)!} \right|_{x=1} \\ &= \left. \left(x \frac{d}{dx} \right)^{p} \sum_{s=0}^{m} \frac{(-x)^{s}}{s!(m-s)!} \right|_{x=1} \\ &= \frac{1}{m!} \left(x \frac{d}{dx} \right)^{p} (1-x)^{m} \Big|_{x=1}, \end{split}$$

where x is a formal variable. As

$$\left(x\frac{d}{dx}\right)^p (1-x)^m$$

is divisable as a polynomial in x by (1 - x) for p < m and

$$\left(x\frac{d}{dx}\right)^m (1-x)^m = m! \cdot x^n + f$$

where f is a polynomial divisable by 1 - x, the lemma follows.

Let $f \in 1 + x\mathbf{R}[[x]]$ any formal power series with constant coefficient one. For each $n \in \mathbf{N}_0$ define the formal power series

$$P_n := \sum_{s=0}^n \frac{(-1)^s}{s!(n-s)!} \cdot \prod_{k=-(n-s)}^s f(k \cdot x)$$

in 1 + xR[[x]].

Lemma 8. The power series P_n has leading term $(-1)^n \left(\frac{1}{n!} \frac{d}{dx^n}\Big|_{x=0} f^{n+1}\right) \cdot x^n$, i.e.

$$P_n = (-1)^n \cdot \left(\left. \frac{d}{dx^n} \right|_{x=0} f^{n+1} \right) \cdot x^n + O(x^{n+1}).$$

Proof. First of all, we consider the logarithm

$$\log f = \sum_{i=1}^{\infty} F_i x^i$$

with certain coefficients $F_i \in R$ as a formal power series in xR[[x]]. This enables us to write

$$\prod_{k=-(n-s)}^{s} f(k \cdot x) = \exp\left(\sum_{k=-(n-s)}^{s} (\log f)(k \cdot x)\right)$$
$$= \exp\left(\sum_{i=1}^{\infty} \sum_{k=-(n-s)}^{s} F_i k^i x^i\right)$$
$$= \exp\left(\sum_{i=1}^{\infty} F_i x^i \left(\sum_{k=1}^{s} k^i + \sum_{k=1}^{n-s} (-k)^i\right)\right)$$

Recall the classical summation formula involving the Bernoulli numbers. These show, that terms like $\sum_{k=1}^{s} k^i$ can be expressed as polynomials in s of degree i + 1, in fact

$$\sum_{k=1}^{s} k^{i} = \frac{1}{i+1} \left(s^{i+1} + \frac{1}{2}(i+1)s^{i} + \dots \right),$$

where " \dots " denotes term of lower degree in s. Similarly, one has

$$\sum_{k=1}^{n-s} (-k)^{i} = \frac{(-1)^{i}}{i+1} \left((n-s)^{i+1} + \frac{1}{2}(i+1)(n-s)^{i} + \dots \right)$$
$$= \frac{1}{i+1} \left(-s^{i+1} + \frac{1}{2}(i+1)(2n+1)s^{i} + \dots \right).$$

Summing up yields

$$\sum_{k=1}^{s} k^{i} + \sum_{k=1}^{n-s} (-k)^{i} = (n+1)s^{i} + \dots$$

Thus, we have

$$\prod_{k=-(n-s)}^{s} f(k \cdot x) = \exp\left(\sum_{i=1}^{\infty} ((n+1)F_i x^i (s^i + \dots))\right) = f^{n+1}(s \cdot x) + g(s, x)$$

for a power series $g \in R[s][[x]]$, in which we collect all the "..."-terms. In each homogeneous component of g the exponent of s is strictly less than the exponent of x. Plugging this into our definition of P_n , we arrive at

$$P_n = \sum_{s=0}^n \frac{(-1)^s}{s!(n-s)!} f^{n+1}(s \cdot x) + xg(s \cdot x).$$

In order to complete the proof, we have to differentiate this expression up to *n*-times with respect to x. So, let $0 \le p \le n$. Then

$$\frac{1}{p!}\frac{d}{dx^p}\Big|_{x=0} P_n = \sum_{s=0}^n \frac{(-1)^s}{s!(n-s)!} \frac{1}{p!}\frac{d}{dx^p}\Big|_{x=0} \left(f^{n+1}(s\cdot x) + xg(s\cdot x)\right).$$

The term $\frac{1}{p!} \frac{d}{dx^p} \Big|_{x=0} f^{n+1}(s \cdot x)$ is of degree p (or less) in s, the term $\frac{1}{p!} \frac{d}{dx^p} \Big|_{x=0} xg(s \cdot x)$ is of degree strictly less than p in s. Therefore, in view of the previous lemma, we have

$$\frac{1}{p!} \frac{d}{dx^p} \Big|_{x=0} P_n = \begin{cases} (-1)^n \frac{1}{n!} \frac{d}{dx^n} \Big|_{x=0} f^{n+1} & \text{for } p = n \\ 0 & \text{for } p < n. \end{cases}$$

References

- 1. Samuel Boissière, On the McKay correspondences for the Hilbert scheme of points on the affine plane, To appear in Math. Ann.; see arXiv:math.AG/0410458.
- <u>_____</u>, Chern classes of the tangent bundle on the Hilbert scheme of points on the affine plane, J. Algebraic Geom. 14 (2005), 761–787.
- Mark Andrea A. de Cataldo and Luca Migliorini, The Douady space of a complex surface, Adv. Math. 151 (2000), no. 2, 283–312. MR MR1758249 (2002a:32008)
- Geir Ellingsrud, Lothar Göttsche, and Manfred Lehn, On the cobordism class of the Hilbert scheme of a surface, J. Algebraic Geom. 10 (2001), no. 1, 81–100. MR MR1795551 (2001k:14005)
- John Fogarty, Algebraic families on an algebraic surface, Amer. J. Math 90 (1968), 511–521. MR MR0237496 (38 #5778)
- Lothar Göttsche, The Betti numbers of the Hilbert scheme of points on a smooth projective surface, Math. Ann. 286 (1990), no. 1-3, 193–207. MR MR1032930 (91h:14007)
- I. Grojnowski, Instantons and affine algebras. I. The Hilbert scheme and vertex operators, Math. Res. Lett. 3 (1996), no. 2, 275–291. MR MR1386846 (97f:14041)
- 8. F. Hirzebruch, Topological methods in algebraic geometry, Third enlarged edition. New appendix and translation from the second German edition by R. L. E. Schwarzenberger, with an additional section by A. Borel. Die Grundlehren der Mathematischen Wissenschaften, Band 131, Springer-Verlag New York, Inc., New York, 1966. MR MR0202713 (34 #2573)
- Manfred Lehn, Chern classes of tautological sheaves on Hilbert schemes of points on surfaces, Invent. Math. 136 (1999), no. 1, 157–207. MR MR1681097 (2000h:14003)
- <u>_____</u>, Lectures on Hilbert schemes, Algebraic structures and moduli spaces, CRM Proc. Lecture Notes, vol. 38, Amer. Math. Soc., Providence, RI, 2004, pp. 1–30. MR MR2095898 (2005g:14010)
- Manfred Lehn and Christoph Sorger, Symmetric groups and the cup product on the cohomology of Hilbert schemes, Duke Math. J. 110 (2001), no. 2, 345–357. MR MR1865244 (2002i:14004)
- Hiraku Nakajima, Heisenberg algebra and Hilbert schemes of points on projective surfaces, Ann. of Math. (2) 145 (1997), no. 2, 379–388. MR MR1441880 (98h:14006)
- Lectures on Hilbert schemes of points on surfaces, University Lecture Series, vol. 18, American Mathematical Society, Providence, RI, 1999. MR MR1711344 (2001b:14007)
- 14. E. T. Whittaker and G. N. Watson, A course of modern analysis, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1996, An introduction to the general theory of infinite processes and of analytic functions; with an account of the principal transcendental functions, Reprint of the fourth (1927) edition. MR MR1424469 (97k:01072)

INSTITUT FÜR MATHEMATIK, JOHANNES-GUTENBERG UNIVERSITÄT, 55099 MAINZ, GERMANY *E-mail address*: boissiere@mathematik.uni-mainz.de, nieper@mathematik.uni-mainz.de