On Conjugates for Integer Compositions and Set Partitions

DAVID CALLAN

Department of Statistics University of Wisconsin-Madison Medical Science Center 1300 University Ave Madison, WI 53706-1532

callan@stat.wisc.edu

Aug 19, 2005

Abstract

There is a familiar conjugate for integer partitions: transpose the Ferrers diagram, and an analogous conjugate for integer compositions. Here we propose a conjugate for set partitions and exhibit statistics interchanged by the conjugate, both for compositions and set partitions.

0 The Conjugate of an Integer Partition A partition of n is a weakly decreasing list of positive integers, called its parts, whose sum is n. The Ferrers diagram of a partition $a_1 \ge a_2 \ge \ldots \ge a_k \ge 1$ is the k-row left-justified array of dots with a_i dots in the i-th row. The conjugate, obtained by transposing the Ferrers diagram, is a well known involution on partitions of n that interchanges the largest part and the number of parts.

1 A Conjugate for Set Partitions The partitions of an *n*-element set, say $[n] = \{1, 2, ..., n\}$, into nonempty blocks are counted by the Bell numbers, A000110 in OEIS. A singleton is a block containing just 1 element and an *adjacency* is an occurrence of two consecutive elements of [n] in the same block. Consecutive is used here in the cyclic sense so that n and 1 are also considered to be consecutive (the standard sense is considered in [2]). We say *i* initiates an adjacency if *i* and $i + 1 \mod n$ are in the same block and analogously for terminating an adjacency.

Theorem. There is a bijection ϕ on partitions of [n] that interchanges number of singletons and number of adjacencies.

To prove this, we need to consider partitions on arbitrary subsets rather than just initial segments of the positive integers. (I thank Robin Chapman [1] for pointing out that my original proof was incorrect.) We always write a partition with elements increasing within each block and blocks arranged in increasing order of their first (smallest) elements. Thus, for example, with a dash separating blocks, the partition $\pi = 3512 - 4810 - 7$ has support $\supp(\pi) = \{3, 4, 5, 7, 8, 10, 12\}$. The successor of an element of the support is the next largest element except that the successor of max(support) is min(support) and for a one-element support $\{a\}$, the successor of a is a itself. A successive pair in $supp(\pi)$ is a pair $(a, b) \subseteq supp(\pi)$ with b the successor of a and $b \neq a$. Thus (5,7) and (12,3) are successive pairs in the preceding example. The notions of adjacency initiator and terminator generalize in the obvious way.

Consider the operation SeparateIS on such partitions defined by SeparateIS(π) = (ρ , (I, S)) where I is the set of adjacency initiators of π , S is the set of singleton elements of π , and ρ is the partition obtained by suppressing the elements of $I \cup S$ in π . Thus, for $\pi = 3512$ -4810-7, we have $I = \{8, 12\}$, $S = \{7\}$ and $\rho = 35$ -410. Also, for a one-block partition $\pi = a_1 a_2 \dots a_k$, SeparateIS(π) = (ϵ , ($\{a_1, \dots, a_k\}, \emptyset$)) where ϵ denotes the empty partition, unless k = 1 in which case it is (ϵ , ($\{a_1\}, \{a_1\}$)). Clearly, for ρ a partition and A, B finite sets of positive integers, the pair ($\rho, (A, B$)) lies in the range of SeparateIS iff (i) $\rho = \epsilon$, A = Band both are singletons, or (ii) $\operatorname{supp}(\rho)$, A, B are disjoint and for no successive pair (a, b) in $\operatorname{supp}(\rho) \cup A \cup B$ is $a \in A$ and $b \in B$ (except for case (i), the successor of an adjacency initiator cannot be a singleton).

Analogously, define SeparateST with S the set of singleton elements and T the set of adjacency terminators. Note that the condition for $(\rho, (A, B))$ to lie in range(SeparateST) is precisely the same as for it to lie in range(SeparateIS). Both are injective and so we may define their respective inverses CombineIS and CombineST and we will make use of the crucial property that these inverses have identical domains. For example, CombineST is defined on $(310-47-12, ({11}, {1,2}))$ and yields the partition 1212-310-47-11.

Now we can define the desired bijection. Given a partition π on [n], form a sequence $(\rho_1, (I_1, S_1)), (\rho_2, (I_2, S_2)), \ldots, (\rho_k, (I_k, S_k))$ by $(\rho_1, (I_1, S_1)) = \text{SeparateIS}(\pi), (\rho_2, (I_2, S_2)) = \text{SeparateIS}(\rho_1), \ldots, (\rho_k, (I_k, S_k)) = \text{SeparateIS}(\rho_{k-1})$ stopping when ρ_k has no adjacency initiators and no singletons (as must eventually occur). For example, with n = 12 and $\pi = 1 - 2 - 3 11 12 - 47 10 - 59 6 8$, the results are laid out in the following table

j	$ ho_j$	I_j	S_j
1	312-4710-59-68	{11}	$\{1, 2\}$
2	3 - 4710 - 59 - 68	{12}	Ø
3	4710-59-68	Ø	{3}
4	47 - 59 - 68	{10}	Ø

and ρ_4 has no adjacency initiators (hence, no adjacencies) and no singletons. Next, form a sequence of partitions $\tau_k, \tau_{k-1}, \ldots, \tau_1, \tau_0$ by reversing the procedure but using CombineST rather than CombineIS. More precisely, set $\tau_k = \rho_k$, $\tau_{k-1} = \text{CombineST}$ on $(\tau_k, (I_k, S_k))$, $\tau_{k-2} =$ CombineST on $(\tau_{k-1}, (I_{k-1}, S_{k-1}))$, $\ldots, \tau_0 = \text{CombineST}$ on $(\tau_1, (I_1, S_1))$. Note that for j = $k, k - 1, \ldots, 1$ in turn, CombineST is defined on $(\tau_j, (I_j, S_j))$ because (i) $\text{supp}(\tau_j) = \text{supp}(\rho_j)$, (ii) CombineIS is certainly defined on $(\rho_j, (I_j, S_j))$, and (iii) CombineST, CombineIS have the same domain. The example yields

j	$ au_j$	I_j	S_j
4	$4\ 7\ -\ 5\ 9\ -\ 6\ 8$	{10}	Ø
3	47 - 59 - 68 - 10	Ø	{3}
2	$3\ 10\ -\ 4\ 7\ -\ 5\ 9\ -\ 6\ 8$	$\{12\}$	Ø
1	310-47-59-68-12	{11}	$\{1, 2\}$
0	1212 - 310 - 47 - 59 - 68 - 11		

Now $\phi : \pi \to \tau_0$ is the desired bijection: ϕ is clearly reversible and sends adjacency initiators to singletons and singletons to adjacency terminators, and hence interchanges #adjacencies and # singletons.

We can use ϕ as follows to construct an *involution*—our proposed conjugate—that interchanges # singletons and # adjacencies. Given a partition of [n], if # singletons = # adjacencies, apply the identity map; if # singletons < # adjacencies, apply ϕ ; if # singletons > # adjacencies, apply ϕ^{-1} .

The number of k-block partitions of [n] containing no adjacencies is considered in a recent Monthly problem proposed by Donald Knuth [3]. Also, Knuth [4, 7.2.1.5., Ex. 20] mentions a different conjugate for set partitions on $[n]: \pi \to n + 1 - \pi$ (elementwise).

2 The Conjugate of an Integer Composition

A composition of n is a list of positive integers—its parts—whose sum is n. There is a bijection from compositions of n to subsets of [n-1] via partial sums: $(c_i)_{i=1}^k \mapsto \{\sum_{j=1}^k c_j\}_{i=1}^{k-1}$, and a further bijection from subsets S of [n-1] to lattice paths of n-1 unit steps North (N) or East (E): the *i*th step is N if $i \in S$ and E otherwise. The conjugate of a composition is defined by: pass to lattice path, flip the path in the 45° line, and pass back. For example, with n = 8 and k = 4,

$$\begin{array}{c} \text{composition} \\ (2,1,2,3) \rightarrow \{2,3,5\} \end{array} \xrightarrow{\text{path}} \\ \begin{array}{c} \text{flip} \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ \end{array} \xrightarrow{\text{path}} \\ \begin{array}{c} \text{flip} \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ \end{array} \xrightarrow{\text{path}} \\ \begin{array}{c} \text{subset} \\ \text{subset} \end{array} \xrightarrow{\text{conjugate}} \\ \begin{array}{c} \text{composition} \\ \text{composition} \end{array} \xrightarrow{\text{conjugate}} \\ \begin{array}{c} \text{conjugate} \end{array} \xrightarrow{\text{conjugate}} \end{array} \xrightarrow{\text{conjugate}} \\ \begin{array}{c} \text{conjugate} \end{array} \xrightarrow{\text{conjugate}} \xrightarrow{\text{conjugate}} \end{array} \xrightarrow{\text{conjugate}} \end{array} \xrightarrow{\text{conjugate}} \xrightarrow{\text{conjugate}} \end{array} \xrightarrow{\text{conjugate}} \end{array} \xrightarrow{\text{conjugate}} \xrightarrow{\text{conjugate}} \xrightarrow{\text{conjugate}} \end{array} \xrightarrow{\text{conjugate}} \xrightarrow{$$

There is a neat graphical construction for the lattice path of a composition using a kind of

shifted Ferrers diagram [5]. Represent a part a_i as a row of a_i dots. Stack the rows so each starts where its predecessor ends. Then join up the dots with E and N steps.



If the compositions of n of a given length are listed in lex (dictionary) order, then so are the corresponding subsets, and the length of the conjugate is n+1- length of the original. It follows that if the compositions of n are sorted, primarily by length and secondarily by lex order (n = 4 is shown),

(4), (1 3), (2 2), (3 1), (1 1 2), (1 2 1), (2 1 1), (1 1 1 1)

then the conjugate of the *i*th composition from the left is the *i*th composition from the right.

There are two statistics on compositions of n that are interchanged by conjugation, and these statistics again involve "singletons" (parts = 1) and adjacent parts. To define them, observe that each part has two neighbors except the end parts which have only one or, in the case of a one-part composition, none ("wraparound" neighbors are not allowed here). Say a part ≥ 2 is big; a part = 1 is small. Now define

 μ = sum of the big parts,

 ν = sum of the small parts + total number of neighbors of the big parts.

For example, in the composition (3, 1, 1, 4, 2), the big parts 3, 4, 2 have 1, 2, 1 neighbors respectively; so $\mu = 3 + 4 + 2 = 9$ and $\nu = (1 + 1) + (1 + 2 + 1) = 6$.

Theorem. Conjugation interchanges the statistics μ and ν on compositions of n except when n = 1.

Proof Carefully translate μ and ν to the corresponding lattice paths. Using the Iverson notation that [statement] = 1 if statement is true, = 0 if it is false, we find

$$\mu = \# Es + \# ENs + [path ends E], and$$
$$\nu = \# Ns + \# ENs + [path starts N]$$

It is then routine to check that μ on the flipped path agrees with ν on the original.

Finally, we remark that the genesis of the statistics μ and ν was the following graphical construction of the conjugate that involves "local" rather than "global" flipping. Suppose given

a composition, say $(4, 2, 1, 2, 1^3, 3)$, where consecutive 1s have been collected so that 1^3 is short for 1, 1, 1. Represent it as a list of vertical strips (for parts > 1) and horizontal strips (for the 1s):



The vertical strips thus consist of 2 or more squares and may occur consecutively, but no two horizontal strips are consecutive. Insert an (initially) empty horizontal strip between each pair of consecutive vertical strips so that horizontal strips H and vertical strips V alternate (the first strip may be either an H or a V):



For each horizontal strip H, transfer one square from each of its neighboring vertical strips to H (there will be two such neighbors in general, but possibly just one or even none in the case of the all-1s composition):



Since each vertical strip originally contained ≥ 2 squares this will always be possible, though some vertical strips may afterward be empty, in which case just erase them. Finally, rotate all strips 90° so that Vs become Hs and vice versa:



The result is the conjugate composition: $(1^3, 2, 3, 5, 1^2)$.

References

[1] Robin Chapman, personal communication.

- [2] Winston Yang, Bell numbers and k-trees, Disc. Math. 156 (1996), 247–252.
- [3] Donald Knuth, Problem 11151, Amer. Math. Monthly 112, April 2005, p. 367.
- [4] Donald Knuth, The Art of Computer Programming, Volume 4, Fascicle 3: Generating All Combinations and Partitions, Addison-Wesley Professional, 2005. Preliminary version.
- [5] P. A. MacMahon, Combinatory Analysis, Vol. 1 Cambridge University Press, 1915; reprinted 2 vols. in 1, Chelsea Publishing Company, New York, 1984, p. 153.