# ENDOMORPHISM ALGEBRAS OF HYPERELLIPTIC JACOBIANS AND FINITE PROJECTIVE LINES

ARSEN ELKIN AND YURI G. ZARHIN

#### 1. Statement of results

Let K be a field with  $\operatorname{char}(K) \neq 2$ . Let us fix an algebraic closure  $K_a$  of K. Let us put  $\operatorname{Gal}(K) := \operatorname{Aut}(K_a/K)$ . If X is an abelian variety of positive dimension over  $K_a$  then we write  $\operatorname{End}(X)$  for the ring of all its  $K_a$ -endomorphisms and  $\operatorname{End}^0(X)$ for the corresponding (semisimple finite-dimensional) Q-algebra  $\operatorname{End}(X) \otimes \mathbb{Q}$ . We write  $\operatorname{End}_K(X)$  for the ring of all K-endomorphisms of X and  $\operatorname{End}^0_K(X)$  for the corresponding (semisimple finite-dimensional) Q-algebra  $\operatorname{End}_K(X) \otimes \mathbb{Q}$ . The absolute Galois group  $\operatorname{Gal}(K)$  of K acts on  $\operatorname{End}(X)$  (and therefore on  $\operatorname{End}^0(X)$ ) by ring (resp. algebra) automorphisms and

$$\operatorname{End}_{K}(X) = \operatorname{End}(X)^{\operatorname{Gal}(K)}, \operatorname{End}_{K}^{0}(X) = \operatorname{End}^{0}(X)^{\operatorname{Gal}(K)},$$

since every endomorphism of X is defined over a finite separable extension of K.

If n is a positive integer that is not divisible by char(K) then we write  $X_n$  for the kernel of multiplication by n in  $X(K_a)$ . It is well-known [17] that  $X_n$  is a free  $\mathbb{Z}/n\mathbb{Z}$ -module of rank  $2\dim(X)$ . In particular, if  $n = \ell$  is a prime then  $X_{\ell}$  is an  $\mathbb{F}_{\ell}$ -vector space of dimension  $2\dim(X)$ .

If X is defined over K then  $X_n$  is a Galois submodule in  $X(K_a)$ . It is known that all points of  $X_n$  are defined over a finite separable extension of K. We write  $\bar{\rho}_{n,X,K}$ :  $\operatorname{Gal}(K) \to \operatorname{Aut}_{\mathbb{Z}/n\mathbb{Z}}(X_n)$  for the corresponding homomorphism defining the structure of the Galois module on  $X_n$ ,

$$\tilde{G}_{n,X,K} \subset \operatorname{Aut}_{\mathbb{Z}/n\mathbb{Z}}(X_n)$$

for its image  $\bar{\rho}_{n,X,K}(\operatorname{Gal}(K))$  and  $K(X_n)$  for the field of definition of all points of  $X_n$ . Clearly,  $K(X_n)$  is a finite Galois extension of K with Galois group  $\operatorname{Gal}(K(X_n)/K) = \tilde{G}_{n,X,K}$ . If  $n = \ell$  then we get a natural faithful linear representation

$$G_{\ell,X,K} \subset \operatorname{Aut}_{\mathbb{F}_{\ell}}(X_{\ell})$$

of  $\tilde{G}_{\ell,X,K}$  in the  $\mathbb{F}_{\ell}$ -vector space  $X_{\ell}$ . Recall [24] that all endomorphisms of X are defined over  $K(X_4)$ ; this gives rise to the natural homomorphism

$$\kappa_{X,4}: \tilde{G}_{4,X,K} \to \operatorname{Aut}(\operatorname{End}^0(X))$$

and  $\operatorname{End}_{K}^{0}(X)$  coincides with the subalgebra  $\operatorname{End}^{0}(X)^{\tilde{G}_{4,X,K}}$  of  $\tilde{G}_{4,X,K}$ -invariants [33, Sect. 1].

Let  $f(x) \in K[x]$  be a polynomial of degree  $n \geq 3$  without multiple roots. Let  $\mathfrak{R}_f \subset K_a$  be the *n*-element set of roots of f. Then  $K(\mathfrak{R}_f)$  is the splitting field of f and  $\operatorname{Gal}(f) := \operatorname{Aut}(K(\mathfrak{R}_f)/K)$  is the Galois group of f (over K). One may view  $\operatorname{Gal}(f)$  as a group of permutations of  $\mathfrak{R}_f$ ; it is transitive if and only if f(x) is irreducible.

Let us consider the hyperelliptic curve  $C_f : y^2 = f(x)$  and its jacobian  $J(C_f)$ . It is well-known that  $J(C_f)$  is a  $\left[\frac{n-1}{2}\right]$ -dimensional abelian variety defined over K. The aim of this paper is to study  $\operatorname{End}^0(J(C_f))$ , assuming that n = q+1 where q is a power of a prime p and  $\operatorname{Gal}(f) = \operatorname{PSL}_2(\mathbb{F}_q)$  acts via fractional-linear transformations on  $\mathfrak{R}_f$  identified with the projective line  $\mathbb{P}^1(\mathbb{F}_q)$ . It follows from results of [27, 30, 32] that for every q in all characteristics do exist K and f with  $\operatorname{End}^0(J(C_f)) = \mathbb{Q}$ . On the other hand, it is known that  $\operatorname{End}^0(J(C_f)) = \mathbb{Q}$  [27, 28, 34] if p = 2 and  $q \ge 8$ . (It is also true for q = 4 if one assumes that  $\operatorname{char}(K) \neq 3$  [32].) However, if q = 5 then there are examples where  $\operatorname{End}^0(J(C_f))$  is a (real) quadratic field (even in characteristic zero) [2, 8, 26, 5].

Our main result is the following statement.

**Theorem 1.1.** Let us assume that  $\operatorname{char}(K) = 0$ . Suppose that n = q + 1 where  $q \geq 5$  is a prime power that is congruent to  $\pm 3$  modulo 8. Suppose that f(x) is irreducible and  $\operatorname{Gal}(f) \cong \operatorname{PSL}_2(\mathbb{F}_q)$ . Then one of the following two conditions holds:

- (i)  $\operatorname{End}^{0}(J(C_{f})) = \mathbb{Q}$  or a quadratic field. In particular,  $J(C_{f})$  is an absolutely simple abelian variety.
- (ii) q is congruent to 3 modulo 8 and  $J(C_f)$  is  $K_a$ -isogenous to a self-product of an elliptic curve with complex multiplication by  $\mathbb{Q}(\sqrt{-q})$ .

**Remark 1.2.** It follows from results of [11] (see also [23], [21]) that if the case (ii) of Theorem 1.1 holds then  $J(C_f)$  is isomorphic over  $K_a$  to a product of mutually isogenous elliptic curves with complex multiplication by  $\mathbb{Q}(\sqrt{-q})$ .

The paper is organized as follows. Section 2 contains auxiliary results about endomorphism algebras of abelian varieties. In Section 3 we prove the main result. In Section 4 we discuss additional conditions that guarantee the absolute simplicity of  $J(C_f)$  when q is congruent to 3 modulo 8. Section 5 contains examples.

### 2. Endomorphism algebras of Abelian varieties

**Remark 2.1.** Recall [6] (see also [29, p. 199]) that a surjective homomorphism of finite groups  $\pi : \mathcal{G}_1 \to \mathcal{G}$  is called a *minimal cover* if no proper subgroup of  $\mathcal{G}_1$  maps onto  $\mathcal{G}$ . If H is a normal subgroup of  $\mathcal{G}_1$  that lies in ker $(\pi)$  then the induced surjection  $\mathcal{G}_1/H \to \mathcal{G}$  is also a minimal cover.

- (i) If a surjection  $\mathcal{G}_2 \twoheadrightarrow \mathcal{G}_1$  is also a minimal cover then one may easily check that the composition  $\mathcal{G}_2 \to \mathcal{G}$  is surjective and a minimal cover.
- (ii) Clearly, if  $\mathcal{G}$  is simple then every proper normal subgroup in  $\mathcal{G}'$  lies in ker $(\pi)$ .
- (iii) If  $\mathcal{G}$  is perfect then its universal central extension is a minimal cover [25].
- (iv) If  $\mathcal{G}_1 \twoheadrightarrow \mathcal{G}$  is an arbitrary surjective homomorphism of finite groups then there always exists a subgroup  $\mathcal{H} \subset \mathcal{G}'$  such that  $\mathcal{H} \to \mathcal{G}$  is surjective and a minimal cover. Clearly, if  $\mathcal{G}$  is perfect then  $\mathcal{G}'$  is also perfect.

The field inclusion  $K(X_2) \subset K(X_4)$  induces a natural surjection [33, Sect. 1]

$$\tau_{2,X}: G_{4,X,K} \to G_{2,X,K}.$$

**Definition 2.2.** We say that K is 2-balanced with respect to X if  $\tau_{2,X}$  is a minimal cover.

**Remark 2.3.** Clearly, there always exists a subgroup  $H \subset \tilde{G}_{4,X,K}$  such that  $\mathcal{H} \to \tilde{G}_{2,X,K}$  is surjective and a minimal cover. Let us put  $L = K(X_4)^H$ . Clearly,

$$K \subset L \subset K(X_4), \ L \bigcap K(X_2) = K$$

and L is a maximal overfield of K that enjoys these properties. It is also clear that

$$K(X_2) \subset L(X_2), \ L(X_4) = K(X_4), \ H = G_{4,X,L}, G_{2,X,L} = G_{2,X,K}$$

and L is 2-balanced with respect to X.

The following assertion (and its proof) is (are) inspired by Theorem 1.6 of [33] (and its proof).

**Theorem 2.4.** Suppose that  $E := \operatorname{End}_{K}^{0}(X)$  is a field that contains the center C of  $\operatorname{End}^{0}(X)$ . Let  $C_{X,K}$  be the centralizer of  $\operatorname{End}_{K}^{0}(X)$  in  $\operatorname{End}^{0}(X)$ . Then:

(i)  $C_{X,K}$  is a central simple E-subalgebra in  $\operatorname{End}^0(X)$ . In addition, the centralizer of  $C_{X,K}$  in  $\operatorname{End}^0(X)$  coincides with  $E = \operatorname{End}^0_K(X)$  and

$$\dim_E(C_{X,K}) = \frac{\dim_C(\operatorname{End}^0(X))}{[E:C]^2}.$$

(ii) Assume that K is 2-balanced with respect to X and G
<sub>2,X,K</sub> is a non-abelian simple group. If End<sup>0</sup>(X) ≠ E (i.e., not all endomorphisms of X are defined over K) then there exist a finite perfect group Π ∈ C<sup>\*</sup><sub>X,K</sub> and a surjective homomorphism Π → G
<sub>X,K,2</sub> that is a minimal cover. In addition, the induced homomorphism

$$\mathbb{Q}[\Pi] \to C_{X,K}$$

is surjective, i.e.,  $C_{X,K}$  is isomorphic to a direct summand of the group algebra  $\mathbb{Q}[\Pi]$ .

*Proof.* Since E is a field, C is a subfield of E and therefore  $\text{End}^{0}(X)$  is a central simple C-algebra. Now the assertion (i) follows from Theorem 2 of Sect. 10.2 in Chapter VIII of [3].

Now let us prove the assertion (ii). Since

Recall that there is the homomorphism

$$\kappa_{X,4}: G_{4,X,K} \to \operatorname{Aut}(\operatorname{End}^0(X))$$

such that

$$\operatorname{End}^{0}(X)^{G_{4,X,K}} = \operatorname{End}^{0}_{K}(X) = E \supset C.$$

This implies that

$$\kappa_{X,4}(\tilde{G}_{4,X,K}) \subset \operatorname{Aut}_E(\operatorname{End}^0(X)) \subset \operatorname{Aut}_C(\operatorname{End}^0(X))$$

and we get a homomorphism

$$\kappa_E : \tilde{G}_{4,X,K} \to \operatorname{Aut}_E(C_{X,K})$$

such that

(1) 
$$C_{X,K}^{\tilde{G}_{4,X,K}} = E$$

Assume that  $E = C_{X,K}$ , i.e., E coincides with its own centralizer in  $\text{End}^0(X)$ . It follows from the Skolem-Noether theorem that  $\text{Aut}_C(\text{End}^0(X)) = \text{End}^0(X)^*$ . This implies that the group

$$\operatorname{Aut}_{E}(\operatorname{End}^{0}(X)) = C^{*}_{X,K}/C^{*} = E^{*}/C^{*}$$

is commutative. It follows that  $\kappa_{X,4}(\tilde{G}_{4,X,K})$  is commutative. Since  $\tilde{G}_{4,X,K}$  is perfect,  $\kappa_{X,4}(\tilde{G}_{4,X,K})$  is perfect commutative and therefore trivial, i.e.,  $\operatorname{End}^0(X) = \operatorname{End}^0_K(X)$ .

Assume that  $E \neq C_{X,K}$ . This means that the group  $\Gamma := \kappa_E(\tilde{G}_{4,X,K})$  is not  $\{1\}$ , i.e.,  $\ker(\kappa_E) \neq \tilde{G}_{4,X,K}$ . Clearly,  $\Gamma$  is a finite perfect subgroup of  $\operatorname{Aut}_E(C_{X,K})$ .

The minimality of  $\tau_{2,X}$  and the simplicity of  $\tilde{G}_{2,X,K}$  imply the existence of a minimal cover

$$\Gamma \to G_{2,X,K},$$

thanks to Remark 2.1.

Since  $C_{X,K}$  is a central simple *E*-algebra, all its automorphisms are inner, i.e., Aut<sub>*E*</sub>( $C_{X,K}$ ) =  $C_{X,K}^*/E^*$ . Let  $\Delta \to \Gamma$  be the universal central extension of  $\Gamma$ . It is well-known that  $\Delta$  is a finite perfect group. The universality property implies that  $\Delta \to \Gamma$  is a minimal cover and the inclusion map  $\Gamma \subset C_{X,K}^*/E^*$  lifts (uniquely) to a homomorphism  $\pi : \Delta \to C_{X,K}^*$ . Clearly, ker( $\pi$ ) lies in the kernel of  $\Delta \to \Gamma$  and we get a minimal cover

$$r(\Delta) \cong \Delta / \ker(\pi) \twoheadrightarrow \Gamma_{\pi}$$

thanks to Remark 2.1. Taking the compositions of minimal covers  $\pi(\Delta) \twoheadrightarrow \Gamma$  and  $\Gamma \twoheadrightarrow \tilde{G}_{2,X,K}$ , we obtain a minimal cover  $\pi(\Delta) \twoheadrightarrow \tilde{G}_{2,X,K}$ . If we put

$$\Pi := \pi(\Delta) \subset C^*_{X,K}.$$

then we get a minimal cover

$$\Pi \twoheadrightarrow \tilde{G}_{2,X,K}$$

The equality (1) means that the centralizer of  $\pi(\Delta) = \Pi$  in  $C_{X,K}$  coincides with E. It follows that if  $\mathbb{Q}[\Pi]$  is the group  $\mathbb{Q}$ -algebra of  $\Pi$  then the inclusion  $\Pi \subset C^*_{X,K}$  induces the  $\mathbb{Q}$ -algebra homomorphism  $\omega : \mathbb{Q}[\Pi] \to C_{X,K}$  such that the centralizer of its image in  $C_{X,K}$  coincides with E.

We claim that  $\omega(\mathbb{Q}[\Pi]) = C_{X,K}$  and therefore  $C_{X,K}$  is isomorphic to a direct summand of  $\mathbb{Q}[\Pi]$ . This claim follows easily from the next lemma that was proven in [33, Lemma 1.7]

**Lemma 2.5.** Let F be a field of characteristic zero, T a semisimple finite-dimensional F-algebra, S a finite-dimensional central simple F-algebra,  $\beta : T \to S$  an F-algebra homomorphism that sends 1 to 1. Suppose that the centralizer of the image  $\beta(T)$  in S coincides with the center F. Then  $\beta$  is surjective, i. e.  $\beta(T) = S$ .

**Theorem 2.6.** Suppose that  $\operatorname{End}^{0}(X)$  is a simple  $\mathbb{Q}$ -algebra,  $\tilde{G}_{2,X,K}$  is a simple non-abelian group, whose order is not a divisor of  $2\dim(X)$  and  $\operatorname{End}_{\tilde{G}_{2,X,K}}(X_{2}) \cong \mathbb{P}$ 

 $\mathbb{F}_4$ . Let us put  $E := \operatorname{End}_K^0(X)$  and let  $C_{X,K}$  be the centralizer of E in  $\operatorname{End}^0(X)$ .

Then the center C of  $\operatorname{End}^0(X)$  is either  $\mathbb{Q}$  or a quadratic field. In addition, there exists a finite separable extension L/K such that  $\tilde{G}_{2,X,L} = \tilde{G}_{2,X,K}$ , the map  $\tau_{2,X} : \tilde{G}_{4,X,L} \twoheadrightarrow \tilde{G}_{2,X,L}$  is surjective and a minimal cover, the  $\mathbb{Q}$ -algebra  $\operatorname{End}^0_L(X)$ is either  $\mathbb{Q}$  or a quadratic field and  $C \subset \operatorname{End}^0_L(X)$ . In particular, C is either  $\mathbb{Q}$  or a quadratic field. If  $\operatorname{End}^0(X) \neq \operatorname{End}^0_L(X)$  then there exist a finite perfect group  $\Pi \subset C^*_{X,L}$  and a surjective homomorphism  $\Pi \to \tilde{G}_{X,K,2}$  that is a minimal cover. In addition, the induced homomorphism

$$\mathbb{Q}[\Pi] \to C_{X,L}$$

is surjective, i.e.,  $C_{X,L}$  is isomorphic to a direct summand of the group algebra  $\mathbb{Q}[\Pi]$ .

*Proof.* Choose a field L as in Remark 2.3. Then  $\tilde{G}_{2,X,L} = \tilde{G}_{2,X,K}$ , the map

$$\tau_{2,X}: \tilde{G}_{4,X,L} \to \tilde{G}_{2,X,L} = \tilde{G}_{2,X,K}$$

is surjective and a minimal cover. We have

$$\operatorname{End}_{L}(X) \otimes \mathbb{Z}/2\mathbb{Z} \hookrightarrow \operatorname{End}_{\operatorname{Gal}(L)}(X_{2}) = \operatorname{End}_{\tilde{G}_{2,X,L}}(X_{2}) = \mathbb{F}_{4}$$

It follows that the rank of the free  $\mathbb{Z}$ -module  $\operatorname{End}_L(X)$  is 1 or 2; Lemma 1.3 of [33] implies that  $\operatorname{End}_L(X)$  has no zero divisors. This implies that  $\operatorname{End}_L^0(X) = \operatorname{End}_L(X) \otimes \mathbb{Q}$  is a division algebra of  $\mathbb{Q}$ -dimension 1 or 2. This means that  $E = \operatorname{End}_L^0(X)$  is either  $\mathbb{Q}$  or a quadratic field.

Recall that the center C of  $\operatorname{End}^{0}(X)$  is a number field, whose degree  $[C : \mathbb{Q}]$ divides  $2\dim(X)$ . The group  $\tilde{G}_{4,X,L}$  acts via automorphisms on C and

$$C^{G_{4,X,L}} = C \bigcap \operatorname{End}_{L}^{0}(X)$$

is either  $\mathbb{Q}$  or a quadratic field. Since  $\tilde{G}_{2,X,L} = \tilde{G}_{2,X,K}$  has no normal subgroups of index dividing  $2\dim(X)$ , the same is true for  $\tilde{G}_{4,X,L}$  and therefore  $\tilde{G}_{4,X,L}$  acts on C trivially, i.e.,  $C \subset \operatorname{End}_{L}^{0}(X)$ . In order to finish the proof, one has only to apply Theorem 2.4 ( to L instead of K).

**Theorem 2.7.** Let us assume that  $\operatorname{char}(K) = 0$ . Suppose that  $\operatorname{End}^0(X)$  is a simple  $\mathbb{Q}$ -algebra with the center C. Suppose that  $\tilde{G}_{2,X,K}$  is a simple non-abelian group, whose order is not a divisor of  $2\dim(X)$  and  $\operatorname{End}_{\tilde{G}_{2,X,K}}(X_2) \cong \mathbb{F}_4$ . Assume, in addition that  $\tilde{G}_{2,X,K}$  is a known simple group that is not one of Lie type in characteristic 2 that is listed in the table in Theorem 3 of p. 316 in [12]. Suppose also that  $\dim(X)$  coincides with the smallest positive integers d such that  $\tilde{G}_{2,X,K}$  is isomorphic to a subgroup of  $\operatorname{PGL}(d, \mathbb{C})$ . Then:

- (i) The center C of  $\operatorname{End}^{0}(X)$  is either  $\mathbb{Q}$  or quadratic field.
- (ii) Either  $\operatorname{End}^{0}(X) = C$  or the following conditions hold:
  - (1) There exist a finite perfect group  $\Pi \subset \operatorname{End}^0(X)^*$  and a surjective homomorphism  $\Pi \to \tilde{G}_{X,K,2}$  that is a minimal cover and a central extension. In addition, the induced homomorphism

$$\mathbb{Q}[\Pi] \to \mathrm{End}^0(X)$$

is surjective, i.e.,  $\operatorname{End}^0(X)$  is isomorphic to a direct summand of the group algebra  $\mathbb{Q}[\Pi]$ .

- (2) If  $C = \mathbb{Q}$  then either  $\operatorname{End}^{0}(X) = \mathbb{Q}$  or  $\dim(X)$  is even and X is isogenous over  $K_{a}$  to a self-product of an abelian surface Y such that  $\operatorname{End}^{0}(Y)$  is an indefinite quaternion  $\mathbb{Q}$ -algebra.
- (3) If  $C \neq \mathbb{Q}$  then C is an imaginary quadratic field and X is isogenous over  $K_a$  to a self-product of an elliptic curve with complex multiplication by C

*Proof.* Using Theorem 2.6 and replacing if necessary K by its suitable extension, we may assume that K is 2-balanced with respect to X, the algebra  $E := \operatorname{End}_{K}^{0}(X)$  is either  $\mathbb{Q}$  or a quadratic field,  $C \subset E$  and the following conditions hold.

Either  $\operatorname{End}^0(X) = E$  or there exist a finite perfect group  $\Pi \subset C^*_{X,K}$  and a surjective homomorphism  $\Pi \to \tilde{G}_{X,K,2}$  that is a minimal cover and such that the induced homomorphism

$$\mathbb{Q}[\Pi] \to C_{X,K}$$

is surjective, i.e.,  $C_{X,K}$  is isomorphic to a direct summand of the group algebra  $\mathbb{Q}[\Pi]$ . (Here as above  $C_{X,K}$  is the centralizer of E in  $\mathrm{End}^0(X)$ 

Assume that  $\operatorname{End}^0(X) \neq E$ . We are going to prove that  $\Pi \to \tilde{G}_{X,K,2}$  is a central extension, using results of Feit-Tits and Kleidman-Liebeck [6, 12]. Without loss of generality we may assume that there is a field embedding  $K_a \hookrightarrow \mathbb{C}$  and consider X as complex abelian variety. Let  $t_X$  be the Lie algebra of X that is a dim(X)-dimensional complex vector space. By functoriality, this gives us the embeddings

$$j : \operatorname{End}^{0}(X) \hookrightarrow \operatorname{End}_{\mathbb{C}}(t_{X}) \cong \operatorname{M}_{\dim(X)}(\mathbb{C}), \ j : \operatorname{End}^{0}(X)^{*} \hookrightarrow \operatorname{Aut}_{\mathbb{C}}(t_{X}) \cong \operatorname{GL}(\dim(X), \mathbb{C})$$

Clearly, only central elements of  $\Pi$  go to scalars under j. It follow that there exists a central subgroup Z in  $\Pi$  such that j(Z) consists of scalars and  $\Pi/Z \hookrightarrow$  PGL(dim $(X), \mathbb{Z}$ ). The simplicity of  $\tilde{G}_{X,K,2}$  implies that Z lies in the kernel of  $\Pi \to \tilde{G}_{X,K,2}$  and the induced map  $\Pi/Z \to \tilde{G}_{X,K,2}$  is also a minimal cover. It follows from Theorem on p. 1002 of [6] and Theorem 3 on p. 316 of [12] that  $\Pi/Z \to \tilde{G}_{X,K,2}$  is a central extension of  $\tilde{G}_{X,K,2}$ . Since  $\Pi$  is a central extension of  $\Pi/Z$ , it follows [25] that  $\Pi$  is a central extension of  $\tilde{G}_{X,K,2}$ .

Now notice that  $t_X$  carries a natural structure of  $E \otimes_{\mathbb{Q}} \mathbb{C}$ -module. Assume that  $E \neq \mathbb{Q}$ , i.e., E is a quadratic field. Let  $\sigma, \tau : E \hookrightarrow \mathbb{C}$  be the two different embeddings of E into  $\mathbb{C}$ . Then

$$E \otimes_{\mathbb{O}} \mathbb{C} = \mathbb{C}_{\sigma} \oplus \mathbb{C}_{\tau}$$

with

$$\mathbb{C}_{\sigma} = E \otimes_{E,\sigma} \mathbb{C} = \mathbb{C}, \ \mathbb{C}_{\tau} = E \otimes_{E,\tau} \mathbb{C} = \mathbb{C}$$

and  $t_X$  splits into a direct sum

$$t_X = \mathbb{C}_{\sigma} t_X \oplus \mathbb{C}_{\tau} t_X.$$

Suppose that both  $\mathbb{C}_{\sigma}t_X$  and  $\mathbb{C}_{\tau}t_X$  do not vanish. Then the  $\mathbb{C}$ -dimension  $d_{\sigma}$  of non-zero  $\mathbb{C}_{\sigma}t_X$  is strictly less than dim(X). Clearly,  $\mathbb{C}_{\sigma}t_X$  is  $C_{X,K}$ -stable and we get a nontrivial homomorphism

$$C_{X,K} \to \operatorname{End}_{\mathbb{C}}(\mathbb{C}_{\sigma}t_X) \cong \operatorname{M}_{d_{\sigma}}(\mathbb{C})$$

that must be an embedding in light of the simplicity of  $C_{X,K}$ . This gives us an embedding

$$C^*_{X,K} \to \operatorname{Aut}_{\mathbb{C}}(\mathbb{C}_{\sigma}t_X) \cong \operatorname{GL}(d_{\sigma},\mathbb{C}).$$

One may easily check that all the elements of  $\Pi$  that go to scalars in  $\operatorname{Aut}_{\mathbb{C}}(\mathbb{C}_{\sigma}t_X)$ constitute a central subgroup  $Z_{\sigma}$  that lies in the kernel of  $\Pi \to \tilde{G}_{X,K,2}$ . This gives us a central extension  $\Pi/Z_{\sigma} \twoheadrightarrow \tilde{G}_{X,K,2}$  that is a minimal cover and an embedding  $\Pi/Z_{\sigma} \hookrightarrow \operatorname{Aut}_{\mathbb{C}}(\mathbb{C}_{\sigma}t_X) \cong \operatorname{GL}(d_{\sigma},\mathbb{C})$ . Since  $d_{\sigma} < \dim(X)$ , Theorem on p. 1002 of [6] and Theorem 3 on p. 316 of [12] provide us with a contradiction. It follows that either  $\mathbb{C}_{\sigma}t_X$  or  $\mathbb{C}_{\tau}t_X$  does vanish. We may assume that  $\mathbb{C}_{\tau}t_X = 0$ . This means that each  $e \in E$  acts on  $t_X$  as multiplication by complex number  $\sigma(e)$ , i.e., j(E) consists of scalars. Recall that the exponential map identifies  $X(\mathbb{C})$  with the complex torus  $t_X/\Lambda$  where  $\Lambda$  is a discrete lattice of rank  $2\dim(X)$ . In addition,  $\Lambda$  is  $j(\operatorname{End}_K(X))$ -stable where  $\operatorname{End}_K(X)$  is an order in the quadratic field  $\operatorname{End}_K^0(X)$ . Now the discreteness of  $\Lambda$  implies that E cannot be real and therefore is an imaginary quadratic field. It follows easily that X is isogenous over  $\mathbb{C}$  to a self-product of an elliptic curve with complex multiplication by E. In particular, E = C and  $C_{X,K} = \operatorname{End}^0(X)$ .

Now let us assume that  $E = \mathbb{Q}$ . Then  $C_{X,K} = \operatorname{End}^0(X)$ . Let Y be an absolutely simple abelian variety such that X is isogenous to a self-product  $Y^r$  for some positive integer r with  $r \mid \dim(X)$ . Then  $\operatorname{End}^0(X) \cong \operatorname{M}_r(\operatorname{End}^0(Y))$ . In particular, the center of the division algebra  $\operatorname{End}^0(Y)$  is  $\mathbb{Q}$ . It follows from Albert's classification [17] that  $\operatorname{End}^0(Y)$  is either  $\mathbb{Q}$  or a quaternion  $\mathbb{Q}$ -algebra.

If  $\operatorname{End}^{0}(Y) = \mathbb{Q}$  then  $\operatorname{End}^{0}(X) \cong \operatorname{M}_{r}(\mathbb{Q})$  and  $\Pi/Z \hookrightarrow \operatorname{PGL}(r, \mathbb{Q}) \subset \operatorname{PGL}(r, \mathbb{C})$ . It follows that  $r = \dim(X)$ , i.e. Y is an elliptic curve without complex multiplication.

Suppose that  $\operatorname{End}^0(Y)$  is a quaternion  $\mathbb{Q}$ -algebra. Since  $\dim(Y) = \dim(X)/r$  and we live in characteristic zero, 2r divides  $\dim(X)$ . Clearly,

$$\operatorname{End}^{0}(Y) \subset \operatorname{End}^{0}(Y) \otimes_{\mathbb{Q}} \mathbb{C} \cong \operatorname{M}_{2}(\mathbb{C})$$

and therefore

$$\operatorname{End}^{0}(X) \cong \operatorname{M}_{r}(\operatorname{End}^{0}(Y)) \hookrightarrow \operatorname{M}_{2r}(\mathbb{C}).$$

This implies that  $\Pi \hookrightarrow \operatorname{GL}(2r, \mathbb{C})$ . It follows that  $2r = \dim(X)$ , i.e.,  $\dim(Y) = 2$ . It follows from the classification of endomorphism algebras of abelian surfaces [18, Sect. 6] that  $\operatorname{End}^0(Y)$  is an *indefinite* quaternion  $\mathbb{Q}$ -algebra.  $\Box$ 

**Theorem 2.8.** Let us assume that  $\operatorname{char}(K) = 0$ . Suppose that  $\operatorname{End}^0(X)$  is a simple  $\mathbb{Q}$ -algebra. Suppose that  $d := \dim(X) = (q-1)/2$  where  $q \ge 5$  is an odd prime power. Suppose that  $\tilde{G}_{2,X,K} \cong \operatorname{PSL}_2(\mathbb{F}_q)$  and  $\operatorname{End}_{\tilde{G}_{2,X,K}}(X_2) \cong \mathbb{F}_4$ . Then one of the following two conditions holds:

- End<sup>0</sup>(X) =  $\mathbb{Q}$  or a quadratic field. In particular, X is an absolutely simple abelian variety.
- q is congruent to 3 modulo 4 and X is  $K_a$ -isogenous to a self-product of an elliptic curve with complex multiplication by  $\mathbb{Q}(\sqrt{-q})$ .

*Proof.* It is well-known that  $\operatorname{SL}_2(\mathbb{F}_q)$  is the universal central extension of  $\operatorname{PSL}_2(\mathbb{F}_q)$ and therefore every projective representation of  $\operatorname{PSL}_2(\mathbb{F}_q)$  lifts to a linear representation of  $\operatorname{SL}_2(\mathbb{F}_q)$ . The well-known list of irreducible representations of  $\operatorname{SL}_2(\mathbb{F}_q)$ over complex numbers tells us that the smallest degree of a nontrivial representation of  $\operatorname{SL}_2(\mathbb{F}_q)$  is (q-1)/2 = d. This implies that we are in position to apply Theorem 2.7. We need to rule out the following possibilities:

- (1) dim(X) is even and X is isogenous over  $K_a$  to a self-product of an abelian surface Y such that  $\operatorname{End}^0(Y)$  is an indefinite quaternion  $\mathbb{Q}$ -algebra. In particular,  $\operatorname{End}^0(X)$  is a  $d^2$ -dimensional central simple  $\mathbb{Q}$ -algebra.
- (2) q is congruent to 1 modulo 4 and X is isogenous over  $K_a$  to a self-product of an elliptic curve with complex multiplication. In particular,  $\operatorname{End}^0(X)$  is a  $d^2$ -dimensional central simple algebra over the imaginary quadratic field C unramified at  $\infty$ .

By Theorem 2.7, there exist a finite perfect group  $\Pi$  and a minimal central cover  $\Pi \to \mathrm{PSL}_2(\mathbb{F}_q)$  such that  $\mathrm{End}^0(X)$  is a quotient of the group algebra  $\mathbb{Q}[\Pi]$ . It follows easily that  $\Pi = \mathrm{PSL}_2(\mathbb{F}_q)$  or  $\mathrm{SL}_2(\mathbb{F}_q)$ , so we may always view  $\mathrm{End}^0(X)$ 

as a simple quotient (direct summand) D of  $\mathbb{Q}[\operatorname{SL}_2(\mathbb{F}_q)]$  that corresponds to an irreducible (complex) character of degree d as in Lemma 24.7 of [4]. These quotients D are described explicitly in [10, 7]. In particular, the center of H is an imaginary quadratic field  $\mathbb{Q}(\sqrt{-q})$  if q is congruent to 3 modulo 4. If q is congruent to 1 modulo 4 but not a square then the center of H is a totally real quadratic field  $\mathbb{Q}(\sqrt{q})$ . If q is a square then the center of H is ramified at infinity. Now the result follows easily.

#### 3. Hyperelliptic Jacobians

Suppose that  $f(x) \in K[x]$  is a polynomial of degree  $n \geq 5$  without multiple roots. Let  $\mathfrak{R}_f \subset K_a$  be the set of roots of f. Clearly,  $\mathfrak{R}_f$  consists of n elements. Let  $K(\mathfrak{R}_f) \subset K_a$  be the splitting field of f. Clearly,  $K(\mathfrak{R}_f)/K$  is a Galois extension and we write  $\operatorname{Gal}(f)$  for its Galois group  $\operatorname{Gal}(K(\mathfrak{R}_f)/K)$ . By definition,  $\operatorname{Gal}(K(\mathfrak{R}_f)/K)$  permutes elements of  $\mathfrak{R}_f$ ; further we identify  $\operatorname{Gal}(f)$  with the corresponding subgroup of  $\operatorname{Perm}(\mathfrak{R}_f)$  where  $\operatorname{Perm}(\mathfrak{R}_f)$  is the group of permutations of  $\mathfrak{R}_f$ .

We write  $\mathbb{F}_{2}^{\mathfrak{R}_{f}}$  for the *n*-dimensional  $\mathbb{F}_{2}$ -vector space of maps  $h: \mathfrak{R}_{f} \to \mathbb{F}_{2}$ . The space  $\mathbb{F}_{2}^{\mathfrak{R}_{f}}$  is provided with a natural action of  $\operatorname{Perm}(\mathfrak{R}_{f})$  defined as follows. Each  $s \in \operatorname{Perm}(\mathfrak{R}_{f})$  sends a map  $h: \mathfrak{R}_{f} \to \mathbb{F}_{2}$  to  $sh: \alpha \mapsto h(s^{-1}(\alpha))$ . The permutation module  $\mathbb{F}_{2}^{\mathfrak{R}_{f}}$  contains the  $\operatorname{Perm}(\mathfrak{R}_{f})$ -stable hyperplane

$$(\mathbb{F}_2^{\mathfrak{R}_f})^0 = \{h : \mathfrak{R}_f \to \mathbb{F}_2 \mid \sum_{\alpha \in \mathfrak{R}_f} h(\alpha) = 0\}$$

and the Perm $(\mathfrak{R}_f)$ -invariant line  $\mathbb{F}_2 \cdot 1_{\mathfrak{R}_f}$  where  $1_{\mathfrak{R}_f}$  is the constant function 1. Clearly,  $(\mathbb{F}_2^{\mathfrak{R}_f})^0$  contains  $\mathbb{F}_2 \cdot 1_{\mathfrak{R}_f}$  if and only if n is even.

If n is even then let us define the  $\operatorname{Gal}(f)$ -module  $Q_{\mathfrak{R}_f} := (\mathbb{F}_2^{\mathfrak{R}_f})^0/(\mathbb{F}_2 \cdot 1_{\mathfrak{R}_f})$ . If n is odd then let us put  $Q_{\mathfrak{R}_f} := (\mathbb{F}_2^{\mathfrak{R}_f})^0$ . If  $n \neq 4$  the natural representation of  $\operatorname{Gal}(f)$  is faithful, because in this case the natural homomorphism  $\operatorname{Perm}(\mathfrak{R}_f) \to \operatorname{Aut}_{\mathbb{F}_2}(Q_{\mathfrak{R}_f})$ is injective.

The canonical surjection  $\operatorname{Gal}(K) \twoheadrightarrow \operatorname{Gal}(K(\mathfrak{R}_f)/K) = \operatorname{Gal}(f)$  provides  $Q_{\mathfrak{R}_f}$ with a natural structure of  $\operatorname{Gal}(K)$ -module. It is well-known that the  $\operatorname{Gal}(K)$ modules  $J(C_f)_2$  and  $Q_{\mathfrak{R}_f}$  are isomorphic (see for instance [19, 20, 28]). It follows easily that  $K(J(C_f)_2) = K(\mathfrak{R}_f)$  and  $\tilde{G}_{2,J(C_f),K} = \operatorname{Gal}(f)$ .

Let us put  $X = J(C_f)$  and  $G := \tilde{G}_{2,X,K}$ . Then  $G \cong \text{Gal}(f)$ , and the *G*-modules  $X_2$  and  $Q_{\mathfrak{R}_f}$  are isomorphic. We freely interchange these two modules throughout this section.

**Example 3.1.** Suppose that n = q + 1 where  $q \ge 5$  is a power of an odd prime p. Suppose that  $\operatorname{Gal}(f) = \operatorname{PSL}_2(\mathbb{F}_q)$ . Assume that that f(x) is *irreducible*, i.e.,  $\operatorname{Gal}(f) = \operatorname{PSL}_2(\mathbb{F}_q)$  acts transitively on the (q + 1)-element set  $\mathfrak{R}_f$ . If  $\beta \in \mathfrak{R}_f$  then its stabilizer  $\operatorname{Gal}(f)_\beta$  is a subgroup of index q + 1 and therefore contains a Sylow p-subgroup of  $\operatorname{PSL}_2(\mathbb{F}_q)$ . It follows from the classification of subgroups of  $\operatorname{PSL}_2(\mathbb{F}_q)$  [25, Theorem 6.25 on page 412] and explicit description of its Sylow p-subgroup and their normalizers [9, p. 191–192] that that  $\operatorname{Gal}(f)_\beta$  is conjugate to the (Borel) subgroup of upper-triangular matrices and therefore the  $\operatorname{PSL}_2(\mathbb{F}_q)$ -set  $\mathfrak{R}_f$  is isomorphic to the projective line  $\mathbb{P}^1(\mathbb{F}_q)$  with the standard action of  $\operatorname{PSL}_2(\mathbb{F}_q)$  (by fractional-linear transformations), which is well-known to be doubly transitive.

Assume, in addition that q is congruent to  $\pm 3 \mod 8$ . Then it is known [16] that

$$\operatorname{End}_{\operatorname{Gal}(f)}(Q_{\mathfrak{R}_f}) = \mathbb{F}_4.$$

**Theorem 3.2.** Suppose that  $\operatorname{char}(K) \neq 2$  and n = q + 1 where  $q \geq 5$  is a prime power that is congruent to  $\pm 3$  modulo 8. Suppose that  $\operatorname{Gal}(f) = \operatorname{PSL}_2(\mathbb{F}_q)$  acts doubly transitively on  $\mathfrak{R}_f$  (where  $\mathfrak{R}_f$  is identified with the projective line  $\mathbb{P}^1(\mathbb{F}_q)$ ). Then  $\operatorname{End}^0(J(C_f))$  is a simple Q-algebra, i.e.  $J(C_f)$  is either absolutely simple or isogenous to a power of an absolutely simple abelian variety.

*Proof.* See [33, Theorem 3.10].

*Proof of Theorem 1.1.* The result follows from Theorem 2.8 combined with Example 3.1 and Theorem 3.2.  $\Box$ 

### 4. CRITERIA FOR ABSOLUTE SIMPLICITY

Sometimes, it is possible to rule out the second outcome of Theorem 1.1. First, recall Goursat's lemma [13, p. 75]:

**Lemma 4.1.** Let  $G_1$  and  $G_2$  be finite group, and H a subgroup of  $G_1 \times G_2$  such that the restrictions  $p_1 : H \to G_1$  and  $p_2 : H \to G_2$  of the projection maps are surjective. Let  $H_1$  and  $H_2$  be the normal subgroups of  $G_1$  and  $G_2$ , respectively, such the groups  $H_1 \times \{1\}$  and  $\{1\} \times H_2$  are kernels of  $p_2$  and  $p_1$ , respectively. Then there exists an isomorphism  $\gamma : G_1/H_1 \cong G_1/H_2$  such that H coincides with the preimage in  $G_1 \times G_2$  of the graph of  $\gamma$  in  $G_1/H_1 \times G_2/H_2$ .

**Example 4.2.** Let  $G_1$  be a finite simple group and  $G_2$  be a finite group which does not admit  $G_1$  as a quotient. If H is a subgroup of  $G_1 \times G_2$  which satisfies the conditions of Goursat's lemma, then  $H = G_1 \times G_2$ .

Indeed, since  $G_1$  is simple,  $H_1 = \{1\}$  or  $G_1$ . We have  $H_1 \neq \{1\}$ , since otherwise  $G_1/H_1 \cong G_1$  and no quotient of  $G_2$  is isomorphic to  $G_1$ . Therefore,  $H_1 = G_1$ ,  $G_1/H_1 \cong G_2/H_2 = \{1\}$ , and  $H_2 = G_2$ . Since  $G_1/H_1 \times G_2/H_2$  is a trivial group, the graph of  $\gamma$  coincides with  $G_1/H_1 \times G_2/H_2$ , and its preimage H coincides with  $G_1 \times G_2$ .

**Theorem 4.3.** Let K be a field of characteristic zero. Suppose that  $f(x) \in K[x]$  is a polynomial of degree  $n \ge 5$  without multiple roots. Let us consider the hyperelliptic curve  $C_f : y^2 = f(x)$  and its jacobian  $J(C_f)$ . Suppose that  $h(x) \in K[x]$  is an irreducible cubic polynomial and let us consider the elliptic curve  $Y : y^2 = h(x)$ .

Let us assume that f(x) and h(x) enjoy the following properties:

- (1)  $\operatorname{Gal}(K(\mathfrak{R}_f)/K) = \operatorname{PSL}_2(\mathbb{F}_q)$  for some odd prime power  $q \equiv 3 \mod 8$  with n = q + 1, and  $\operatorname{Gal}(K(\mathfrak{R}_f)/K)$  acts doubly transitively on  $\mathfrak{R}_f$  (where  $\mathfrak{R}_f$  is identified with the projective line  $\mathbb{P}^1(\mathbb{F}_q)$ );
- (2)  $\operatorname{Gal}(K(\mathfrak{R}_h)/K) = SS_3.$

Then  $\operatorname{Hom}(J(C_f), Y) = 0$  and  $\operatorname{Hom}(Y, J(C_f)) = 0$ . In particular,  $J(C_f)$  is not  $K_a$ -isogenous to a self-product of Y.

*Proof.* First, we prove that  $K(\mathfrak{R}_f)$  and  $K(\mathfrak{R}_h)$  are linearly disjoint over K. Let us put  $G_1 := \operatorname{Gal}(K(\mathfrak{R}_f)/K), G_2 := \operatorname{Gal}(K(\mathfrak{R}_h)/K),$  and  $H := \operatorname{Gal}(K(\mathfrak{R}_f, \mathfrak{R}_h)/K),$  the Galois group of the compositum of  $K(\mathfrak{R}_f)$  and  $K(\mathfrak{R}_h)$  over K. By Theorem 1.14 of [13], H can be considered to be a subgroup of  $G_1 \times G_2$ , where the Galois restriction maps coincide with restrictions of projection maps  $p_i : G_1 \times G_2 \to G_i$ ,

with i = 1, 2, to H. It follows from Example 4.2 that  $H \cong G_1 \times G_2$ , and  $K(\mathfrak{R}_f)$ and  $K(\mathfrak{R}_h)$  are linearly disjoint over K. The equalities  $\operatorname{Hom}(J(C_f), Y) = 0$  and  $\operatorname{Hom}(Y, J(C_f)) = 0$  follow from the definitions (s) and (p3) and Theorem 2.5 of [31]. Since for any positive integer r we have  $\operatorname{Hom}(J(C_f), Y^r) = \prod_{i=1}^r \operatorname{Hom}(J(C_f), Y) = 0$ , we conclude that  $\operatorname{Hom}(J(C_f), Y^r) = 0$ .

We write j for the classical modular function [14, Ch. 3, Sect. 3].

**Corollary 4.4.** Let p be a prime such that  $p \equiv 3 \mod 8$  and  $3 \le p \le 1051$ . Let  $q \ge 11$  be an odd power of p. (In particular,  $q \equiv p \equiv 3 \mod 8$ .) Let us put

$$\omega := \frac{-1 + \sqrt{-p}}{2}, \ \alpha := j(\omega) \in \mathbb{C}, \ K := \mathbb{Q}(j(\omega)) \subset \mathbb{C}.$$

Suppose that  $f(x) \in K[x]$  is an irreducible polynomial of degree q + 1 such that  $\operatorname{Gal}(f/K) = \operatorname{PSL}_2(\mathbb{F}_q)$  acts doubly transitively on  $\mathfrak{R}_f$  (where  $\mathfrak{R}_f$  is identified with the projective line  $\mathbb{P}^1(\mathbb{F}_q)$ ).

Then  $J(C_f)$  is an absolutely simple abelian variety, and  $\operatorname{End}^0(J(C_f)) = \mathbb{Q}$  or a quadratic field.

*Proof.* Clearly,  $\mathbb{Q}(\sqrt{-p}) = \mathbb{Q}(\sqrt{-q})$ . Since  $p \equiv 3 \mod 8$ , the ring of integers  $\emptyset$  in  $\mathbb{Q}(\sqrt{-p})$  coincides with  $\mathbb{Z} + \mathbb{Z}\omega$ . If p > 3 let us consider the polynomial

$$h_p(x) := x^3 - \frac{27\alpha}{4(\alpha - 1728)}x - \frac{27\alpha}{4(\alpha - 1728)} \in K[x].$$

If p = 3 then

$$\omega = \frac{-1 + \sqrt{-3}}{2}, \ \alpha = j(\omega) = 0, \ K = \mathbb{Q}$$

and we put

$$h_3(x) := x^3 - 2 \in \mathbb{Q}[x] = K[x].$$

Using MAGMA computational package [1], we verified that if p satisfies the conditions of the theorem then  $h_p(x)$  is irreducible over K and its Galois group  $\operatorname{Gal}(h_p/K) \cong SS_3$ . The elliptic curve  $Y : y^2 = h_p(x)$  is defined over K and its j-invariant coincides with  $\alpha = j(\omega)$ , i.e.,  $Y(\mathbb{C}) = \mathbb{C}/(\mathbb{Z} + \mathbb{Z}\omega)$ . Hence Y admits complex multiplication by  $\mathbb{Z} + \mathbb{Z}\omega = \emptyset$ . By Theorem 4.3,  $J(C_f)$  is not isogenous to a self-product of Y. Now the result follows from Theorem 1.1.

**Remark 4.5.** If p = 3, 11, 19, 43, 67 or 163 then  $j(\omega) \in \mathbb{Z}$  [22], so  $K = \mathbb{Q}(\alpha) = \mathbb{Q}(j(\omega)) = \mathbb{Q}$ .

**Remark 4.6.** It would be interesting to find out if  $\operatorname{Gal}(h_p/K) \cong SS_3$  for all p with  $p \equiv 3 \mod 8$ . If this is the case then one may drop the restriction  $p \leq 1051$  in the statement of Corollary 4.4.

### 5. Examples

**Example 5.1.** Let S be a transcendental over  $\mathbb{Q}$ ,  $T = 2^8 3^5 / (11S^2 + 1)$ , and put

$$f_{11,S}(x) := (x^3 - 66x - 308)^4 - 9T(11x^5 - 44x^4 - 1573x^3 + 1892x^2 + 57358x + 103763) - 3T^2(x - 11).$$

According to Table 10 of the Appendix in [15],  $\operatorname{Gal}(f_{11,S}/\mathbb{Q}(S)) = \operatorname{PSL}_2(\mathbb{F}_{11})$ . It can be verified using MAGMA [1] that when s = m/n for any nonzero integers  $-5 \leq m, n \leq 5$ , then  $\operatorname{Gal}(f_{11,s}/\mathbb{Q}) = \operatorname{PSL}_2(\mathbb{F}_{11})$ . Define the hyperelliptic curve

$$C_{11,s}: y^2 = f_{11,s}(x)$$

over  $\mathbb{Q}(j((-1+\sqrt{-11})/2)) = \mathbb{Q}$  for any one of these s. Then by Corollary 4.4 the 5-dimensional abelian variety  $J(C_{11,s})$  is absolutely simple.

s	$f_{11,s}(x)$
-5	$\begin{array}{r} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -370808064/23x^5-1289570832/23x^4+5531810240/23x^3\\ +56937133440/23x^2+4018853846400/529x+4650105431488/529\end{array}$
-5/2	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -501830208/31x^5-1729935504/31x^4+7748414080/31x^3\\ +76389540480/31x^2+6968010969600/961x+7873029928192/961\end{array}$
-5/3	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -1156940928/71x^5-3931758864/71x^4+18831433280/71x^3\\ +173651575680/71x^2+33695317737600/5041x\\ +36730846917952/5041\end{array}$
-5/4	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -1594524096/97x^5-5315899248/97x^4+27717247360/97x^3\\ +234848972160/97x^2+55633666214400/9409x\\ +58251794030848/9409 \end{array}$
-4	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -951939648/59x^5-3305084112/59x^4+14295594368/59x^3\\ +145929472128/59x^2+26218129108992/3481x\\ +30197814439936/3481\end{array}$
-4/3	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -3033885888/185x^5-10167437808/185x^4+51830793344/185x^3\\ +449149117824/185x^2+209588584132608/34225x\\ +221614446456832/34225\end{array}$

s	$f_{11,s}(x)$
-4/5	$\begin{array}{c} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -1130006592/67x^5-3557269584/67x^4+23239604608/67x^3\\ +157290173568/67x^2+15890216497152/4489x\\ +16829869010944/4489\end{array}$
-3	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -404034048/25x^5-1397780208/25x^4+6153235264/25x^3\\ +61719318144/25x^2+4619555772288/625x+5269380605632/625 \end{array}$
-3/2	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -1682911296/103x^5-5685690384/103x^4+27966944896/103x^3\\ +251137535616/103x^2+68417021833728/10609x\\ +73580830369024/10609\end{array}$
-3/4	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -1950011712/115x^5-6063968592/115x^4+41382960256/115x^3\\ +268178587776/115x^2+39951908510208/13225x\\ +44813546981632/13225\end{array}$
-3/5	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -537584256/31x^5-1586919312/31x^4+12861242944/31x^3\\ +70239844224/31x^2+737335045248/961x+2380957877056/961\end{array}$
-2	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -81183168/5x^5-278050608/5x^4+1284466304/5x^3\\ +12279129984/5x^2+174855579648/25x+194281618432/25\end{array}$
-2/3	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -908715456/53x^5-2754640944/53x^4+20504206976/53x^3\\ +121872871296/53x^2+5409113745408/2809x\\ +7911924849664/2809\end{array}$
-2/5	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -421616448/23x^5-1086337296/23x^4+12797409152/23x^3\\ +48198091392/23x^2-2880749180928/529x+2494254889984/529\end{array}$
-1	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -16612992x^5-54104688x^4+310712512x^3+2391092352x^2\\ +4956865152x+5044849216 \end{array}$
-1/2	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -88710336/5x^5-247941936/5x^4+2360851328/5x^3\\ +10984457088/5x^2-41887170048/25x+64716475648/25\end{array}$

## HYPERELLIPTIC JACOBIANS

S	$f_{11,s}(x)$
-1/3	$ \begin{aligned} &x^{12} - 264  x^{10} - 1232  x^9 + 26136  x^8 + 243936  x^7 - 580800  x^6 \\ &- 94355712/5  x^5 - 225360432/5  x^4 + 3168140096/5  x^3 \\ &+ 10013452416/5  x^2 - 227206388352/25  x + 217926334912/25 \end{aligned} $
-1/4	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -19749312x^5-41559408x^4+759206272x^3+1851645312x^2\\ -15393289728x+19418603776 \end{array}$
-1/5	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -20376576x^5-39050352x^4+848905024x^3+1743755904x^2\\ -20185928832x+30242044096 \end{array}$
1	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -16612992x^5-54104688x^4+310712512x^3+2391092352x^2\\ +4956865152x+5044849216 \end{array}$
1/2	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\-88710336/5x^5-247941936/5x^4+2360851328/5x^3\\+10984457088/5x^2-41887170048/25x+64716475648/25\end{array}$
1/3	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -94355712/5x^5-225360432/5x^4+3168140096/5x^3\\ +10013452416/5x^2-227206388352/25x+217926334912/25\end{array}$
1/4	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -19749312x^5-41559408x^4+759206272x^3\\ +1851645312x^2-15393289728x+19418603776 \end{array}$
1/5	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -20376576x^5-39050352x^4+848905024x^3\\ +1743755904x^2-20185928832x+30242044096 \end{array}$
2	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\-81183168/5x^5-278050608/5x^4+1284466304/5x^3\\+12279129984/5x^2+174855579648/25x+194281618432/25\end{array}$
2/3	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -908715456/53x^5-2754640944/53x^4+20504206976/53x^3\\ +121872871296/53x^2+5409113745408/2809x\\ +7911924849664/2809\end{array}$
2/5	$x^{12} - 264 x^{10} - 1232 x^9 + 26136 x^8 + 243936 x^7 - 580800 x^6$

s	$f_{11,s}(x)$
	$-421616448/23 x^5 - 1086337296/23 x^4 + 12797409152/23 x^3 \\ +48198091392/23 x^2 - 2880749180928/529 x + 2494254889984/529$
3	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -404034048/25x^5-1397780208/25x^4+6153235264/25x^3\\ +61719318144/25x^2+4619555772288/625x+5269380605632/625 \end{array}$
3/2	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -1682911296/103x^5-5685690384/103x^4+27966944896/103x^3\\ +251137535616/103x^2+68417021833728/10609x\\ +73580830369024/10609 \end{array}$
3/4	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -1950011712/115x^5-6063968592/115x^4+41382960256/115x^3\\ +268178587776/115x^2+39951908510208/13225x\\ +44813546981632/13225\end{array}$
3/5	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -537584256/31x^5-1586919312/31x^4+12861242944/31x^3\\ +70239844224/31x^2+737335045248/961x+2380957877056/961\end{array}$
4	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -951939648/59x^5-3305084112/59x^4+14295594368/59x^3\\ +145929472128/59x^2+26218129108992/3481x\\ +30197814439936/3481\end{array}$
4/3	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -3033885888/185x^5-10167437808/185x^4+51830793344/185x^3\\ +449149117824/185x^2+209588584132608/34225x\\ +221614446456832/34225\end{array}$
4/5	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -1130006592/67x^5-3557269584/67x^4+23239604608/67x^3\\ +157290173568/67x^2+15890216497152/4489x\\ +16829869010944/4489\end{array}$
5	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -370808064/23x^5-1289570832/23x^4+5531810240/23x^3\\ +56937133440/23x^2+4018853846400/529x+4650105431488/529\end{array}$
5/2	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -501830208/31x^5-1729935504/31x^4+7748414080/31x^3\\ +76389540480/31x^2+6968010969600/961x+7873029928192/961\end{array}$

HYPERELLIPTIC JACOBIANS

s	$f_{11,s}(x)$
5/3	$\begin{array}{l} x^{12}-264x^{10}-1232x^9+26136x^8+243936x^7-580800x^6\\ -1156940928/71x^5-3931758864/71x^4+18831433280/71x^3\\ +173651575680/71x^2+33695317737600/5041x\\ +36730846917952/5041 \end{array}$
5/4	$\begin{array}{l} x^{12}-264x^{10}-1232x^{9}+26136x^{8}+243936x^{7}-580800x^{6}\\ -1594524096/97x^{5}-5315899248/97x^{4}+27717247360/97x^{3}\\ +234848972160/97x^{2}+55633666214400/9409x\\ +58251794030848/9409\end{array}$

Example 5.2. If we define

$$f_{13,S}(x) := (x^2 + 36)(x^3 - x^2 + 35x - 27)^4 - 4T(7x^2 - 2x + 247)(x^2 + 39)^6/27$$

with  $T = 1/(39S^2 + 1)$  then again [15] we have  $\operatorname{Gal}(f_{13,S}/\mathbb{Q}(S)) = \operatorname{PSL}_2(\mathbb{F}_{13})$ . Similarly, when s = m/n for any nonzero integers  $-5 \le m, n \le 5$ , then  $\operatorname{Gal}(f_{13,s}/\mathbb{Q}) = \operatorname{PSL}_2(\mathbb{F}_{13})$ . If we define

$$C_{13,s}: y^2 = f_{13,s}(x)$$

over  $\mathbb{Q}$ , then by Theorem 2.7 the 6-dimensional abelian variety  $J(C_{13,s})$  is absolutely simple.

TABLE 2. Some polynomials over  $\mathbb{Q}$  with Galois group  $PSL_2(\mathbb{F}_{13})$ 

s	$f_{13,s}(x)$
-5	$6581/6588x^{14} - 13175/3294x^{13} + 1197131/6588x^{12} - 123695/183x^{11}$
	$+9748765/732x^{10}-5569187/122x^9+120857113/244x^8$
	$-94078582/61x^7+2328530789/244x^6-3195234315/122x^5$
	$+ 20264560875/244x^4 - 11171912661/61x^3 + 42022670157/244x^2$
	-11972359197/122 x - 27521811477/244
-5/2	$26321/26433  x^{14} - 105700/26433  x^{13} + 4780646/26433  x^{12}$
,	$-1984580/2937  x^{11} + 3529565/267  x^{10} - 44670184/979  x^9$
	$+478725208/979  x^8 - 1508830648/979  x^7 + 9104797949/979  x^6$
	$-25609631580/979x^5 + 75819375450/979x^4 - 178819338204/979x^3$
	$+98304582987/979  x^2 - 92948708304/979  x - 496309720932/979$
-5/3	$731/738x^{14} - 1475/369x^{13} + 132431/738x^{12} - 27690/41x^{11}$
,	$+1070615/82 x^{10} - 1869361/41 x^9 + 39240539/82 x^8$
	$-63116092/41x^7+729660217/82x^6-1070384445/41x^5$
	$+ 5590660725/82x^4 - 7455579066/41x^3 - 1500464979/82x^2$
	-3676323591/41  x - 95001153831/82
-5/4	$26309/26757 x^{14} - 106900/26757 x^{13} + 4749134/26757 x^{12}$
3/1	$-2006420/2973 x^{11} + 38145835/2973 x^{10} - 45136936/991 x^9$

l	s	$f_{13,s}(x)$
		$\begin{array}{l} +459912232/991x^8-1523123992/991x^7+8267497121/991x^6\\ -25800659820/991x^5+54863903250/991x^4-179094280716/991x^3\\ -180839807577/991x^2-81628047216/991x-2041199621028/991 \end{array}$
	-4	$\begin{array}{l} 16847/16875x^{14}-67492/16875x^{13}+612742/3375x^{12}\\ -1267292/1875x^{11}+24936457/1875x^{10}-5705648/125x^9\\ +61765964/125x^8-192758696/125x^7+1187183959/125x^6\\ -3273066108/125x^5+51248611674/625x^4-114408633924/625x^3\\ +19840740777/125x^2-60958859256/625x-116802534384/625 \end{array}$
	-4/3	$\begin{array}{l} 1871/1899x^{14}-7588/1899x^{13}+338078/1899x^{12}-142428/211x^{11}\\ +2720393/211x^{10}-9613136/211x^9+98762612/211x^8\\ -324440792/211x^7+1792573081/211x^6-5497560900/211x^5\\ +12426098958/211x^4-38197309644/211x^3-28964185497/211x^2\\ -17803917288/211x-382243044816/211 \end{array}$
	-4/5	$\begin{array}{l} 16823/17523x^{14}-69892/17523x^{13}+3000686/17523x^{12}\\ -1310972/1947x^{11}+2143427/177x^{10}-29461744/649x^{9}\\ +271203868/649x^{8}-992380168/649x^{7}+4261318139/649x^{6}\\ -16747387020/649x^{5}+9337667274/649x^{4}-114958518948/649x^{3}\\ -459085077243/649x^{2}-38317537080/649x-3206582334576/649 \end{array}$
	-3	$\begin{array}{l} 2369/2376x^{14}-4751/1188x^{13}+430547/2376x^{12}-44603/66x^{11}\\ +318227/24x^{10}-2008019/44x^9+43257721/88x^8-16957967/11x^7\\ +827107073/88x^6-1151557875/44x^5+7015835619/88x^4\\ -4022805033/22x^3+11406236049/88x^2-4234578237/44x\\ -30506384133/88 \end{array}$
	-3/2	$\begin{array}{l} 9473/9585x^{14}-38308/9585x^{13}+342862/1917x^{12}-719108/1065x^{11}\\ +13832143/1065x^{10}-3236168/71x^9+33665528/71x^8\\ -109245656/71x^7+619820617/71x^6-1852044012/71x^5\\ +22824474426/355x^4-64433616156/355x^3-4832230689/71x^2\\ -31046460624/355x-508248011556/355\end{array}$
	-3/4	$\begin{array}{l} 9461/9909x^{14}-39508/9909x^{13}+1682798/9909x^{12}-740948/1101x^{11}\\ +13152763/1101x^{10}-16647592/367x^9+149514664/367x^8\\ -560521624/367x^7+2261802257/367x^6-9451248300/367x^5\\ +1869002226/367x^4-64708558668/367x^3-303305544009/367x^2\\ -19725799536/367x-2053137911652/367\end{array}$
	-3/5	$\frac{2363/2538 x^{14} - 5051/1269 x^{13} + 414791/2538 x^{12} - 94666/141 x^{11}}{+ 3160807/282 x^{10} - 2124707/47 x^9 + 720239/2 x^8 - 71405204/47 x^7} \\ + 408456659/94 x^6 - 1199314935/47 x^5 - 3461900481/94 x^4$

s	$f_{13,s}(x)$
	$-8114345694/47  x^3 - 128165959233/94  x^2 - 1404412965/47  x \\ -802951334181/94$
-2	$\begin{array}{l} 4211/4239x^{14}-16948/4239x^{13}+763958/4239x^{12}-318188/471x^{11}\\ +6191653/471x^{10}-7161232/157x^9+76031644/157x^8\\ -241841704/157x^7+1431648647/157x^6-4103271900/157x^5\\ +11502435906/157x^4-28619342388/157x^3+7354401561/157x^2\\ -14532173496/157x-125756252352/157 \end{array}$
-2/3	$\begin{array}{l} 467/495x^{14}-1972/495x^{13}+16510/99x^{12}-36972/55x^{11}\\ +57967/5x^{10}-498160/11x^9+4232644/11x^8-16758040/11x^7\\ +58229873/11x^6-282041604/11x^5-822626298/55x^4\\ -9600879132/55x^3-11916123921/11x^2-2328355368/55x\\ -385227617472/55\end{array}$
-2/5	$\begin{array}{l} 4187/4887x^{14}-19348/4887x^{13}+700934/4887x^{12}-361868/543x^{11}\\ +4832893/543x^{10}-8094736/181x^9+38405692/181x^8\\ -270428392/181x^7-242953009/181x^6-4485328380/181x^5\\ -30408508494/181x^4-29169227412/181x^3-550934379567/181x^2\\ +8109148680/181x-3215536052544/181\end{array}$
-1	$\begin{array}{l} 263/270x^{14}-539/135x^{13}+9451/54x^{12}-10114/15x^{11}\\ +376363/30x^{10}-45487x^9+891605/2x^8-1533844x^7\\ +15279043/2x^6-25943931x^5+391472991/10x^4-896502438/5x^3\\ -780396201/2x^2-365687757/5x-31998670461/10 \end{array}$
-1/2	$\begin{array}{l} 1049/1161x^{14}-4612/1161x^{13}+181142/1161x^{12}-86372/129x^{11}\\ +1335607/129x^{10}-1936168/43x^9+13128856/43x^8\\ -64927096/43x^7+96255653/43x^6-1085514300/43x^5\\ -3672976086/43x^4-7240755132/43x^3-85394021661/43x^2\\ -95336784/43x-514217156868/43 \end{array}$
-1/3	$\begin{array}{l} 29/36x^{14}-71/18x^{13}+4667/36x^{12}-663x^{11}+29237/4x^{10}\\ -88777/2x^9+440843/4x^8-1476722x^7-21051641/4x^6\\ -48546225/2x^5-1033701903/4x^4-153235719x^3\\ -16808682033/4x^2+192566889/2x-96493440159/4 \end{array}$
-1/4	$\begin{array}{l} 1037/1485x^{14}-5812/1485x^{13}+29926/297x^{12}-108212/165x^{11}\\+59657/15x^{10}-480584/11x^9-1136824/11x^8-15844088/11x^7\\-148209035/11x^6-255308508/11x^5-24628448286/55x^4\\-7515697644/55x^3-72907682445/11x^2+11225324304/55x\\-2059107056964/55\end{array}$

s	$f_{13,s}(x)$
-1/5	$\begin{array}{c} 257/432x^{14}-839/216x^{13}+31499/432x^{12}-7787/12x^{11}\\ +36673/48x^{10}-344123/8x^9-4948463/16x^8-2810639/2x^7\\ -342255199/16x^6-177476715/8x^5-10086263109/16x^4\\ -482619033/4x^3-143474176287/16x^2+2464477515/8x\\ -804443620509/16\end{array}$
1	$\begin{array}{l} 263/270x^{14}-539/135x^{13}+9451/54x^{12}-10114/15x^{11}\\ +376363/30x^{10}-45487x^9+891605/2x^8-1533844x^7\\ +15279043/2x^6-25943931x^5+391472991/10x^4-896502438/5x^3\\ -780396201/2x^2-365687757/5x-31998670461/10 \end{array}$
1/2	$\begin{array}{l} 1049/1161x^{14}-4612/1161x^{13}+181142/1161x^{12}-86372/129x^{11}\\ +1335607/129x^{10}-1936168/43x^9+13128856/43x^8\\ -64927096/43x^7+96255653/43x^6-1085514300/43x^5\\ -3672976086/43x^4-7240755132/43x^3-85394021661/43x^2\\ -95336784/43x-514217156868/43 \end{array}$
1/3	$\begin{array}{l} 29/36x^{14}-71/18x^{13}+4667/36x^{12}-663x^{11}+29237/4x^{10}\\ -88777/2x^9+440843/4x^8-1476722x^7-21051641/4x^6\\ -48546225/2x^5-1033701903/4x^4-153235719x^3\\ -16808682033/4x^2+192566889/2x-96493440159/4 \end{array}$
1/4	$\begin{array}{l} 1037/1485x^{14}-5812/1485x^{13}+29926/297x^{12}-108212/165x^{11}\\ +59657/15x^{10}-480584/11x^9-1136824/11x^8-15844088/11x^7\\ -148209035/11x^6-255308508/11x^5-24628448286/55x^4\\ -7515697644/55x^3-72907682445/11x^2+11225324304/55x\\ -2059107056964/55 \end{array}$
1/5	$\begin{array}{l} 257/432x^{14}-839/216x^{13}+31499/432x^{12}-7787/12x^{11}\\ +36673/48x^{10}-344123/8x^9-4948463/16x^8-2810639/2x^7\\ -342255199/16x^6-177476715/8x^5-10086263109/16x^4\\ -482619033/4x^3-143474176287/16x^2+2464477515/8x\\ -804443620509/16 \end{array}$
2	$\begin{array}{l} 4211/4239x^{14}-16948/4239x^{13}+763958/4239x^{12}-318188/471x^{11}\\ +6191653/471x^{10}-7161232/157x^9+76031644/157x^8\\ -241841704/157x^7+1431648647/157x^6-4103271900/157x^5\\ +11502435906/157x^4-28619342388/157x^3+7354401561/157x^2\\ -14532173496/157x-125756252352/157 \end{array}$
2/3	$\begin{array}{l} 467/495x^{14}-1972/495x^{13}+16510/99x^{12}-36972/55x^{11}\\ +57967/5x^{10}-498160/11x^9+4232644/11x^8-16758040/11x^7\\ +58229873/11x^6-282041604/11x^5-822626298/55x^4 \end{array}$

s	$f_{13,s}(x)$
	$-9600879132/55x^3-11916123921/11x^2$
	-2328355368/55  x - 385227617472/55
2/5	$\begin{array}{l} 4187/4887x^{14}-19348/4887x^{13}+700934/4887x^{12}-361868/543x^{11}\\ +4832893/543x^{10}-8094736/181x^9+38405692/181x^8\\ -270428392/181x^7-242953009/181x^6-4485328380/181x^5\\ -30408508494/181x^4-29169227412/181x^3-550934379567/181x^2\\ +8109148680/181x-3215536052544/181\end{array}$
3	$\begin{array}{l} 2369/2376x^{14}-4751/1188x^{13}+430547/2376x^{12}-44603/66x^{11}\\ +318227/24x^{10}-2008019/44x^9+43257721/88x^8-16957967/11x^7\\ +827107073/88x^6-1151557875/44x^5+7015835619/88x^4\\ -4022805033/22x^3+11406236049/88x^2\\ -4234578237/44x-30506384133/88 \end{array}$
3/2	$\begin{array}{l} 9473/9585x^{14}-38308/9585x^{13}+342862/1917x^{12}\\ -719108/1065x^{11}+13832143/1065x^{10}-3236168/71x^9\\ +33665528/71x^8-109245656/71x^7+619820617/71x^6\\ -1852044012/71x^5+22824474426/355x^4-64433616156/355x^3\\ -4832230689/71x^2-31046460624/355x-508248011556/355\end{array}$
3/4	$\begin{array}{l} 9461/9909x^{14}-39508/9909x^{13}+1682798/9909x^{12}-740948/1101x^{11}\\ +13152763/1101x^{10}-16647592/367x^9+149514664/367x^8\\ -560521624/367x^7+2261802257/367x^6-9451248300/367x^5\\ +1869002226/367x^4-64708558668/367x^3-303305544009/367x^2\\ -19725799536/367x-2053137911652/367\end{array}$
3/5	$\begin{array}{l} 2363/2538x^{14}-5051/1269x^{13}+414791/2538x^{12}\\ -94666/141x^{11}+3160807/282x^{10}-2124707/47x^9+720239/2x^8\\ -71405204/47x^7+408456659/94x^6-1199314935/47x^5\\ -3461900481/94x^4-8114345694/47x^3-128165959233/94x^2\\ -1404412965/47x-802951334181/94 \end{array}$
4	$\begin{array}{l} 16847/16875x^{14}-67492/16875x^{13}+612742/3375x^{12}\\ -1267292/1875x^{11}+24936457/1875x^{10}-5705648/125x^9\\ +61765964/125x^8-192758696/125x^7+1187183959/125x^6\\ -3273066108/125x^5+51248611674/625x^4-114408633924/625x^3\\ +19840740777/125x^2-60958859256/625x-116802534384/625 \end{array}$
4/3	$\begin{array}{l} 1871/1899x^{14}-7588/1899x^{13}+338078/1899x^{12}-142428/211x^{11}\\ +2720393/211x^{10}-9613136/211x^9+98762612/211x^8\\ -324440792/211x^7+1792573081/211x^6-5497560900/211x^5\\ +12426098958/211x^4-38197309644/211x^3-28964185497/211x^2\\ \end{array}$

s	$f_{13,s}(x)$
	-17803917288/211  x - 382243044816/211
4/5	$\begin{array}{l} 16823/17523x^{14}-69892/17523x^{13}+3000686/17523x^{12}\\ -1310972/1947x^{11}+2143427/177x^{10}-29461744/649x^{9}\\ +271203868/649x^{8}-992380168/649x^{7}+4261318139/649x^{6}\\ -16747387020/649x^{5}+9337667274/649x^{4}-114958518948/649x^{3}\\ -459085077243/649x^{2}-38317537080/649x-3206582334576/649 \end{array}$
5	$\begin{array}{l} 6581/6588x^{14}-13175/3294x^{13}+1197131/6588x^{12}-123695/183x^{11}\\ +9748765/732x^{10}-5569187/122x^9+120857113/244x^8\\ -94078582/61x^7+2328530789/244x^6-3195234315/122x^5\\ +20264560875/244x^4-11171912661/61x^3+42022670157/244x^2\\ -11972359197/122x-27521811477/244 \end{array}$
5/2	$\begin{array}{l} 26321/26433x^{14}-105700/26433x^{13}+4780646/26433x^{12}\\ -1984580/2937x^{11}+3529565/267x^{10}-44670184/979x^9\\ +478725208/979x^8-1508830648/979x^7+9104797949/979x^6\\ -25609631580/979x^5+75819375450/979x^4-178819338204/979x^3\\ +98304582987/979x^2-92948708304/979x-496309720932/979 \end{array}$
5/3	$\begin{array}{l} 731/738x^{14}-1475/369x^{13}+132431/738x^{12}-27690/41x^{11}\\ +1070615/82x^{10}-1869361/41x^9+39240539/82x^8-63116092/41x^7\\ +729660217/82x^6-1070384445/41x^5+5590660725/82x^4\\ -7455579066/41x^3-1500464979/82x^2-3676323591/41x\\ -95001153831/82 \end{array}$
5/4	$\begin{array}{l} 26309/26757x^{14}-106900/26757x^{13}+4749134/26757x^{12}\\ -2006420/2973x^{11}+38145835/2973x^{10}-45136936/991x^9\\ +459912232/991x^8-1523123992/991x^7+8267497121/991x^6\\ -25800659820/991x^5+54863903250/991x^4-179094280716/991x^3\\ -180839807577/991x^2-81628047216/991x-2041199621028/991 \end{array}$

#### References

- W. Bosma, J. Cannon, C. Playoust, The Magma Algebra System I: The User Language. J. Symb. Comp. 24 (1997), 235–265; http://magma.maths.usyd.edu.au/magma/.
- [2] A. Brumer, The Rank of  $J_0(N)$ , Astérique **228** (1995), 41–68.
- [3] N. Bourbaki, Algébre, Chapitre VIII, Hermann, Paris, 1958.
- [4] L. Dornhoff, Group Representation Theory, Part A. Marcel Dekker, Inc., New York, 1972.
- [5] A. Elkin, Hyperelliptic jacobians with real multiplication. J. Number Theory, to appear; http://front.math.ucdavis.edu/math.AG/0403553.
- [6] W. Feit, J. Tits, Projective representations of minimum degree of group extensions. Canad. J. Math. 30 (1978), 1092–1102.
- [7] W. Feit, The computations of some Schur indices. Israel J. Math. 46 (1983), 274–300.
- [8] K. Hashimoto, On Brumer's family of RM-curves of genus two, Tohoku Math. J. 52 (2000), 475–488.
- [9] B. Huppert, Endliche Gruppen I. Springer-Verlag, Berlin Heidelberg New York, 1967.
- [10] G. Janusz, Simple components of  $\mathbb{Q}[SL(2,q)]$ . Commun. Algebra 1 (1974), 1–22.
- [11] T. Katsura, On the structure of singular abelian varieties. Proc. Japan Acad. 51 (1975), no. 4, 224–228.

- [12] P. B. Kleidman, M. W. Liebeck, On a theorem of Feit and Tits. Proc. Amer. Math. Soc. 107 (1989), 315–322.
- [13] S. Lang, Algebra, rev. 3rd ed., Springer-Verlag, New York, 2002.
- [14] S. Lang, Elliptic functions, 2nd edition, Springer-Verlag, New York, 1987.
- [15] G. Malle, B. H. Matzat, Inverse Galois Theory, Springer-Verlag, Berlin, 1999.
- [16] B. Mortimer, The modular permutation representations of the known doubly transitive groups. Proc. London Math. Soc. (3) 41 (1980), 1–20.
- [17] D. Mumford, Abelian varieties, 2nd edn, Oxford University Press, 1974.
- [18] F. Oort, Endomorphism algebras of abelian varieties. In: Algebraic Geometry and Commutative Algebra in Honor of M. Nagata (Ed. H. Hijikata et al), Kinokuniya Cy, Tokyo 1988; Vol. II, pp. 469 - 502.
- B. Poonen, E. Schaefer, Explicit descent for Jacobians of cyclic covers of the projective line. J. reine angew. Math. 488 (1997), 141–188.
- [20] E. Schaefer, Computing a Selmer group of a Jacobian using functions on the curve. Math. Ann. 310 (1998), 447–471.
- [21] C. Schoen, Produkte Abelscher Varietäten und Moduln über Ordnungen. J. Reine Angew. Math. 429 (1992), 115–123.
- [22] J.-P. Serre, Complex Multiplication. In: Algebraic Number Theory (J. Cassels, A. Frölich, eds.), Chapter XIII, pp. 292–296. Academic Press, 1967.
- [23] T. Shioda, N. Mitani, Singular abelian surfaces and binary quadratic forms. In: Classification of Algebraic and Compact Complex Manifolds, Springer Lect. Notes in Math. 412 (1974), 259–287.
- [24] A. Silverberg, Fields of definition for homomorphisms of abelian varieties. J. Pure Appl. Algebra 77 (1992), 253–262.
- [25] M. Suzuki, Group theory. I. Springer-Verlag, Berlin-New York, 1982.
- [26] J. Wilson, Explicit moduli for curves of genus 2 with real multiplication by  $\mathbb{Q}(\sqrt{5})$ , Acta Arith. **93** (2000), 121–138.
- [27] Yu. G. Zarhin, Hyperelliptic jacobians without complex multiplication. Math. Res. Letters 7 (2000), 123–132.
- [28] Yu. G. Zarhin, Hyperelliptic jacobians and modular representations. In: Moduli of abelian varieties (eds. C. Faber, G. van der Geer and F. Oort). Progress in Math., vol. 195 (2001), Birkhäuser, pp. 473–490.
- [29] Yu. G. Zarhin, Hyperelliptic Jacobians without Complex Multiplication, Doubly Transitive Permutation Groups and Projective Representations. In: Algebraic Number Theory and Algebraic Geometry (Parshin Festschrift), Contemp. Math. **300** (2002), 195–210.
- [30] Yu. G. Zarhin, Very simple 2-adic representations and hyperelliptic jacobians. Moscow Math. J. 2 (2002), issue 2, 403-431.
- [31] Yu. G. Zarhin, Homomorphisms of hyperelliptic Jacobians. In: Number Theory, Algebra and Algebraic Geometry (Shafarevich Festschrift), Tr. Mat. Inst. Steklova 241 (2003), 90–104; Proc. Steklov Inst. Math. 241 2003, 79–92.
- [32] Yu. G. Zarhin, Non-supersingular hyperelliptic jacobians. Bull. Soc. Math. France 132 (2004), 617–634
- [33] Yu. G. Zarhin, Homomorphisms of abelian varieties. In: Proceedings of the European school "Algebraic Geometry and Information Theory" and of the conference "Arithmetic, Geometry and Coding Theory-9", Séminaires et Congres 11, to appear.
- [34] Yu. G. Zarhin, Hyperelliptic jacobians without complex multiplication and Steinberg representations in positive characteristic, http://front.math.ucdavis.edu/math.NT/0301177.

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA

E-mail address: elkin@math.psu.edu

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA

E-mail address: zarhin@math.psu.edu