

ABELIANIZATION CONJECTURES FOR SOME ARITHMETIC SQUARE COMPLEX GROUPS

DIEGO RATTAGGI

ABSTRACT. We extend a conjecture of Kimberley-Robertson on the abelianizations of certain square complex groups.

1. INTRODUCTION

Throughout this paper, let p, l be any pair of distinct odd prime numbers,

$$r_{p,l} := \gcd\left(\frac{p-1}{4}, \frac{l-1}{4}, 6\right) \in \{1, 2, 3, 6\},$$

and $q \in \{p, l\}$. We first recall the definition of the group $\Gamma_{p,l}$ from [2, 3, 4, 5]. Let \mathbb{Q}_q be the field of q -adic numbers. We fix elements $c_p, d_p \in \mathbb{Q}_p$ and $c_l, d_l \in \mathbb{Q}_l$ such that

$$c_p^2 + d_p^2 + 1 = 0 \in \mathbb{Q}_p \quad \text{and} \quad c_l^2 + d_l^2 + 1 = 0 \in \mathbb{Q}_l.$$

Note that we can take $d_q = 0$, if $q \equiv 1 \pmod{4}$.

Let $\mathbb{H}(\mathbb{Q})^*$ be the multiplicative group of invertible rational Hamilton quaternions, i.e. the set

$$\{x_0 + x_1i + x_2j + x_3k : x_0, x_1, x_2, x_3 \in \mathbb{Q}\} \setminus \{0\}$$

equipped with the multiplication induced by the rules $i^2 = j^2 = k^2 = -1$ and $ij = k = -ji$. If $x = x_0 + x_1i + x_2j + x_3k$, we define as usual the conjugate $\bar{x} := x_0 - x_1i - x_2j - x_3k$, and the norm $|x|^2 := x\bar{x} = x_0^2 + x_1^2 + x_2^2 + x_3^2$.

Let ψ_q be the homomorphism of groups $\mathbb{H}(\mathbb{Q})^* \rightarrow \mathrm{PGL}_2(\mathbb{Q}_q)$ defined by

$$\psi_q(x_0 + x_1i + x_2j + x_3k) := \left[\begin{array}{cc} x_0 + x_1c_q + x_3d_q & -x_1d_q + x_2 + x_3c_q \\ -x_1d_q - x_2 + x_3c_q & x_0 - x_1c_q - x_3d_q \end{array} \right]$$

and let the homomorphism

$$\psi_{p,l} : \mathbb{H}(\mathbb{Q})^* \rightarrow \mathrm{PGL}_2(\mathbb{Q}_p) \times \mathrm{PGL}_2(\mathbb{Q}_l)$$

be given by $\psi_{p,l}(x) := (\psi_p(x), \psi_l(x))$. Observe that it satisfies $\psi_{p,l}(-x) = \psi_{p,l}(x)$ and $\psi_{p,l}(x)^{-1} = \psi_{p,l}(\bar{x})$.

Let $\mathbb{H}(\mathbb{Z})$ be the set of integer Hamilton quaternions and X_q the subset of quaternions

$$X_q := \{x = x_0 + x_1i + x_2j + x_3k \in \mathbb{H}(\mathbb{Z}) ; \quad |x|^2 = q ; \\ x_0 \text{ odd, if } q \equiv 1 \pmod{4} ; x_1 \text{ even, if } q \equiv 3 \pmod{4}\}$$

Date: November 11, 2018.

Supported by the Swiss National Science Foundation, No. PP002-68627.

of cardinality $2(q+1)$.

Finally, let $Q_{p,l}$ be the subgroup of $\mathbb{H}(\mathbb{Q})^*$ generated by $(X_p \cup X_l) \subset \mathbb{H}(\mathbb{Z})$ and let $\Gamma_{p,l} < \mathrm{PGL}_2(\mathbb{Q}_p) \times \mathrm{PGL}_2(\mathbb{Q}_l)$ be its image $\psi_{p,l}(Q_{p,l})$, which is a finitely presented, torsion-free linear group.

The starting point for this work was the following conjecture of Kimberley and Robertson for the abelianization $\Gamma_{p,l}^{ab} := \Gamma_{p,l}/[\Gamma_{p,l}, \Gamma_{p,l}]$ of the group $\Gamma_{p,l}$ in the case $p, l \equiv 1 \pmod{4}$. We use the notation $\mathbb{Z}_n := \mathbb{Z}/n\mathbb{Z}$ and $\mathbb{Z}_n^m := \mathbb{Z}/n\mathbb{Z} \times \dots \times \mathbb{Z}/n\mathbb{Z}$ (m times).

Conjecture 1. (*Kimberley-Robertson* [1, Section 6]) *If $p, l \equiv 1 \pmod{4}$, then*

$$\Gamma_{p,l}^{ab} \cong \begin{cases} \mathbb{Z}_2 \times \mathbb{Z}_4^3, & \text{if } r_{p,l} = 1 \\ \mathbb{Z}_2^3 \times \mathbb{Z}_8^2, & \text{if } r_{p,l} = 2 \\ \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_4^3, & \text{if } r_{p,l} = 3 \\ \mathbb{Z}_2^3 \times \mathbb{Z}_3 \times \mathbb{Z}_8^2, & \text{if } r_{p,l} = 6. \end{cases}$$

In Section 2, we will give an equivalent formulation of this conjecture and a new conjecture relating the abelianization of $\Gamma_{p,l}$ to the number $t_{p,l}$ of certain pairs of commuting quaternions, defined as

$$t_{p,l} := |\{(x, y) \in Y_p \times Y_l : xy = yx\}|,$$

where Y_q is any subset of X_q of cardinality $(q+1)/2$ such that $x \in Y_q$ implies $x_0 > 0$ and $\bar{x} \notin Y_q$. Note that the definition of $t_{p,l}$ does not depend on the choice of elements in Y_p and Y_l , and that $\psi_{p,l}(Y_p \cup Y_l)$ is a generating set of $\Gamma_{p,l}$ of cardinality $(p+1)/2 + (l+1)/2$.

The case $p, l \equiv 3 \pmod{4}$ is treated in Section 3 and the remaining (mixed) case in Section 4. The final section is devoted to conjectures on the abelianizations of some subgroups of $\Gamma_{p,l}$.

The Conjectures 2, 3, 4, 5, 7, 9, 10, 11 and 12 have been stated in the authors Ph.D. thesis ([2, Chapter 3]). We have checked Conjectures 2, 4, 5, 7, 9, 10, 11 and 12 for more than 100 different pairs (p, l) which are explicitly listed in [2, Table 3.13].

2. THE CASE $p, l \equiv 1 \pmod{4}$

In this section, we restrict to the ‘‘classical’’ case $p, l \equiv 1 \pmod{4}$. The following conjecture is equivalent to Conjecture 1.

Conjecture 2. *Let $p, l \equiv 1 \pmod{4}$.*

If $p, l \equiv 1 \pmod{8}$, then

$$\Gamma_{p,l}^{ab} \cong \begin{cases} \mathbb{Z}_2^3 \times \mathbb{Z}_3 \times \mathbb{Z}_8^2, & \text{if } p, l \equiv 1 \pmod{3} \\ \mathbb{Z}_2^3 \times \mathbb{Z}_8^2, & \text{else.} \end{cases}$$

If $p \equiv 5 \pmod{8}$ or $l \equiv 5 \pmod{8}$, then

$$\Gamma_{p,l}^{ab} \cong \begin{cases} \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_4^3, & \text{if } p, l \equiv 1 \pmod{3} \\ \mathbb{Z}_2 \times \mathbb{Z}_4^3, & \text{else.} \end{cases}$$

Proof of the equivalence of Conjecture 1 and Conjecture 2. If $r_{p,l} = 6$, then $(p-1)/4 = 6s$ and $(l-1)/4 = 6t$ for some $s, t \in \mathbb{N}$, i.e. $p = 24s + 1$ and $l = 24t + 1$. It follows $p, l \equiv 1 \pmod{8}$ and $p, l \equiv 1 \pmod{3}$.

If $r_{p,l} = 3$, then $(p-1)/4 = 3s$ and $(l-1)/4 = 3t$, where s or t is odd (otherwise $r_{p,l}$ would be 6). Consequently, we have $p = 12s + 1$ and $l = 12t + 1$, in particular $p, l \equiv 1 \pmod{3}$. If s is odd, then $p \equiv 5 \pmod{8}$. If t is odd, then $l \equiv 5 \pmod{8}$.

If $r_{p,l} = 2$, then $(p-1)/4 = 2s$ and $(l-1)/4 = 2t$, i.e. $p = 8s + 1$ and $l = 8t + 1$, hence $p, l \equiv 1 \pmod{8}$. Moreover, $s \not\equiv 0 \pmod{3}$ or $t \not\equiv 0 \pmod{3}$ (otherwise $r_{p,l}$ would be 6). In the first case, we have $p \not\equiv 1 \pmod{3}$, in the second case $l \not\equiv 1 \pmod{3}$.

If $r_{p,l} = 1$, then $(p-1)/4 = 2s - 1$ or $(l-1)/4 = 2t - 1$ (otherwise $r_{p,l}$ would be even), hence $p = 8s - 3$ or $l = 8t - 3$, i.e. $p \equiv 5 \pmod{8}$ or $l \equiv 5 \pmod{8}$. Moreover: $(p-1)/4 = 3s + 1$ or $(p-1)/4 = 3s + 2$ or $(l-1)/4 = 3t + 1$ or $(l-1)/4 = 3t + 2$ for some $s, t \in \mathbb{N}_0$ (otherwise $r_{p,l}$ would be a multiple of 3), hence $p = 12s + 5$ or $p = 12s + 9$ or $l = 12t + 5$ or $l = 12t + 9$, in particular $p \not\equiv 1 \pmod{3}$ or $l \not\equiv 1 \pmod{3}$. \square

The equivalence of the two conjectures above is also expressed in Table 1.

$r_{p,l}$	$l \equiv 1$	5	9	13	17	21 (mod 24)
$p \equiv 1$	6	1	2	3	2	1
5	1	1	1	1	1	1
9	2	1	2	1	2	1
13	3	1	1	3	1	1
17	2	1	2	1	2	1
21	1	1	1	1	1	1

TABLE 1. $r_{p,l}$ for p, l taken modulo 24

The structure of $\Gamma_{p,l}^{ab}$ also seems to depend only on the number $t_{p,l}$ defined in Section 1. Observe that

$$3 \leq t_{p,l} \leq \min \left\{ \frac{p+1}{2}, \frac{l+1}{2} \right\},$$

if $p, l \equiv 1 \pmod{4}$.

Conjecture 3. *Let $p, l \equiv 1 \pmod{4}$. Then*

$$t_{p,l} \equiv \begin{cases} 3 \pmod{12}, & \text{if } r_{p,l} = 1 \\ 9 \pmod{12}, & \text{if } r_{p,l} = 2 \\ 7 \pmod{12}, & \text{if } r_{p,l} = 3 \\ 1 \pmod{12}, & \text{if } r_{p,l} = 6. \end{cases}$$

We have checked Conjecture 3 for all pairs of distinct prime numbers $p, l < 1000$ such that $p, l \equiv 1 \pmod{4}$. The following values for $t_{p,l}$ appear in this range:

$$t_{p,l} \in \begin{cases} \{3, 15, 27, 39, 51, 63, 75, 87, 99\}, & \text{if } r_{p,l} = 1 \\ \{9, 21, 33, 45, 57, 69, 81, 93, 105, 117, 129, 153\}, & \text{if } r_{p,l} = 2 \\ \{7, 19, 31, 43, 55, 67, 79, 91, 103, 115, 127, 151\}, & \text{if } r_{p,l} = 3 \\ \{37, 49, 61, 73, 85, 97, 109, 121, 133\}, & \text{if } r_{p,l} = 6. \end{cases}$$

See Table 2 for the frequencies of the values of $t_{p,l}$, where $p, l \equiv 1 \pmod{4}$ are prime numbers such that $p < l < 1000$.

$t_{p,l}$	3	15	27	39	51	63	75	
#	1242	449	143	56	34	17	7	
	87	99						
	5	2						1955
$t_{p,l}$	9	21	33	45	57	69	81	
#	178	158	84	57	40	21	8	
	93	105	117	129	141	153		
	9	12	5	2		1		575
$t_{p,l}$	7	19	31	43	55	67	79	
#	236	130	79	42	18	8	12	
	91	103	115	127	139	151		
	6	1	4	2		1		539
$t_{p,l}$	1	13	25	37	49	61	73	
#				26	15	15	16	
	85	97	109	121	133			
	7	4	3	2	3			91
								3160

TABLE 2. $t_{p,l}$ and its frequency, $p < l < 1000$

Combining Conjecture 3 with Conjecture 1, we get another conjecture:

Conjecture 4. *Let $p, l \equiv 1 \pmod{4}$. Then*

$$\Gamma_{p,l}^{ab} \cong \begin{cases} \mathbb{Z}_2 \times \mathbb{Z}_4^3, & \text{if } t_{p,l} \equiv 3 \pmod{12} \\ \mathbb{Z}_2^3 \times \mathbb{Z}_8^2, & \text{if } t_{p,l} \equiv 9 \pmod{12} \\ \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_4^3, & \text{if } t_{p,l} \equiv 7 \pmod{12} \\ \mathbb{Z}_2^3 \times \mathbb{Z}_3 \times \mathbb{Z}_8^2, & \text{if } t_{p,l} \equiv 1 \pmod{12}. \end{cases}$$

3. THE CASE $p, l \equiv 3 \pmod{4}$

If $p, l \equiv 3 \pmod{4}$, we have a conjecture similar to Conjecture 2.

Conjecture 5. *Let $p, l \equiv 3 \pmod{4}$.
If $p \pmod{8} = l \pmod{8}$, then*

$$\Gamma_{p,l}^{ab} \cong \begin{cases} \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_8^2, & \text{if } p, l \equiv 1 \pmod{3} \quad (=: \text{case (B1)}) \\ \mathbb{Z}_2 \times \mathbb{Z}_8^2, & \text{else} \quad (=: \text{case (B2)}). \end{cases}$$

If $p \pmod{8} \neq l \pmod{8}$, then

$$\Gamma_{p,l}^{ab} \cong \begin{cases} \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_4^2, & \text{if } p, l \equiv 1 \pmod{3} \quad (=: \text{case (B3)}) \\ \mathbb{Z}_2 \times \mathbb{Z}_4^2, & \text{else} \quad (=: \text{case (B4)}). \end{cases}$$

The four cases (B1)–(B4) defined in the conjecture above can also be expressed taking p and l modulo 24, see Table 3.

	$l \equiv 3$	7	11	15	19	23 (mod 24)
$p \equiv 3$	(B2)	(B4)	(B2)	(B4)	(B2)	(B4)
7	(B4)	(B1)	(B4)	(B2)	(B3)	(B2)
11	(B2)	(B4)	(B2)	(B4)	(B2)	(B4)
15	(B4)	(B2)	(B4)	(B2)	(B4)	(B2)
19	(B2)	(B3)	(B2)	(B4)	(B1)	(B4)
23	(B4)	(B2)	(B4)	(B2)	(B4)	(B2)

TABLE 3. Cases (B1)–(B4) for p, l taken modulo 24

The connection to $t_{p,l}$ is not as nice as in Section 2. We get the following values for $t_{p,l}$, if $p, l \equiv 3 \pmod{4}$ are distinct prime numbers less than 1000.

$$t_{p,l} \in \begin{cases} (\{4, 6, \dots, 104\} \cup \{110, 114, 122, 124, 132\}) \setminus \{84, 88\}, & \text{in case (B1)} \\ \{0, 2, \dots, 78\} \cup \{84, 100, 110\}, & \text{in case (B2)} \\ \{0\}, & \text{in case (B3)} \\ \{0\}, & \text{in case (B4)}. \end{cases}$$

In general, i.e. without the restriction $p, l < 1000$, it is easy to see that $t_{p,l}$ is always even. Moreover, it follows from [4, Section 5] that $t_{p,l} = 0$ in the cases (B3), (B4), and $t_{p,l} > 0$ in case (B1). The computations of $t_{p,l}$ combined with Conjecture 5 lead to the following conjecture:

Conjecture 6. *Let $p, l \equiv 3 \pmod{4}$.*

- (1) *If $t_{p,l} = 0$, then $\Gamma_{p,l}^{ab} \cong \mathbb{Z}_2 \times \mathbb{Z}_8^2$ or $\Gamma_{p,l}^{ab} \cong \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_4^2$ or $\Gamma_{p,l}^{ab} \cong \mathbb{Z}_2 \times \mathbb{Z}_4^2$.*
- (2) *If $t_{p,l} = 2$, then $\Gamma_{p,l}^{ab} \cong \mathbb{Z}_2 \times \mathbb{Z}_8^2$.*
- (3) *If $t_{p,l} \geq 4$, then $\Gamma_{p,l}^{ab} \cong \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_8^2$ or $\Gamma_{p,l}^{ab} \cong \mathbb{Z}_2 \times \mathbb{Z}_8^2$.*
- (4) *If $\Gamma_{p,l}^{ab} \cong \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_4^2$ or $\Gamma_{p,l}^{ab} \cong \mathbb{Z}_2 \times \mathbb{Z}_4^2$, then $t_{p,l} = 0$.*
- (5) *If $\Gamma_{p,l}^{ab} \cong \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_8^2$, then $t_{p,l} \geq 4$.*

4. THE CASE $p \equiv 3 \pmod{4}$, $l \equiv 1 \pmod{4}$

The remaining case is $p \pmod{4} \neq l \pmod{4}$. Since $\Gamma_{p,l} \cong \Gamma_{l,p}$, we can restrict to $p \equiv 3 \pmod{4}$, $l \equiv 1 \pmod{4}$.

Conjecture 7. *Let $p \equiv 3 \pmod{4}$, $l \equiv 1 \pmod{4}$.*

If $l \equiv 1 \pmod{8}$, then

$$\Gamma_{p,l}^{ab} \cong \begin{cases} \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_8^2, & \text{if } p, l \equiv 1 \pmod{3} & (= : \text{case (C1)}) \\ \mathbb{Z}_2 \times \mathbb{Z}_8^2, & \text{else} & (= : \text{case (C2)}). \end{cases}$$

If $l \equiv 5 \pmod{8}$, then

$$\Gamma_{p,l}^{ab} \cong \begin{cases} \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_4^2, & \text{if } p, l \equiv 1 \pmod{3} & (= : \text{case (C3)}) \\ \mathbb{Z}_2 \times \mathbb{Z}_4^2, & \text{else} & (= : \text{case (C4)}). \end{cases}$$

Observe that the four conjectured possibilities for $\Gamma_{p,l}^{ab}$ are exactly the same as in Conjecture 5.

See Table 4 for the cases (C1)–(C4) expressed by p and l taken modulo 24.

	$l \equiv 1$	5	9	13	17	21 (mod 24)
$p \equiv 3$	(C2)	(C4)	(C2)	(C4)	(C2)	(C4)
7	(C1)	(C4)	(C2)	(C3)	(C2)	(C4)
11	(C2)	(C4)	(C2)	(C4)	(C2)	(C4)
15	(C2)	(C4)	(C2)	(C4)	(C2)	(C4)
19	(C1)	(C4)	(C2)	(C3)	(C2)	(C4)
23	(C2)	(C4)	(C2)	(C4)	(C2)	(C4)

TABLE 4. Cases (C1)–(C4) for p, l taken modulo 24

The behaviour of $t_{p,l}$ seems to be very similar as in Section 3. We get the following values for $t_{p,l}$, if $p \equiv 3 \pmod{4}$, $l \equiv 1 \pmod{4}$ are prime numbers less than 1000.

$$t_{p,l} \in \begin{cases} (\{4, 6, \dots, 48\} \cup \{58\}) \setminus \{40\}, & \text{in case (C1)} \\ \{0, 2, \dots, 54\} \cup \{60\}, & \text{in case (C2)} \\ \{0\}, & \text{in case (C3)} \\ \{0\}, & \text{in case (C4)}. \end{cases}$$

Conjecture 8. *Conjecture 6 also holds if $p \equiv 3 \pmod{4}$, $l \equiv 1 \pmod{4}$.*

5. MORE CONJECTURES

In this section, we give conjectures for the abelianization of the commutator subgroup $[\Gamma_{p,l}, \Gamma_{p,l}]$ of $\Gamma_{p,l}$ and for a certain subgroup $\Lambda_{p,l}$ of $\Gamma_{p,l}$ of index 4 defined below.

Conjecture 9. *Let $p, l \equiv 1 \pmod{4}$.*

If $p, l \equiv 1 \pmod{8}$, then

$$[\Gamma_{p,l}, \Gamma_{p,l}]^{ab} \cong \begin{cases} \mathbb{Z}_2^2 \times \mathbb{Z}_{16}^2 \times \mathbb{Z}_{64}, & \text{if } p, l \equiv 1 \pmod{3} \\ \mathbb{Z}_3 \times \mathbb{Z}_{16}^2 \times \mathbb{Z}_{64}, & \text{else.} \end{cases}$$

If $p \equiv 5 \pmod{8}$ or $l \equiv 5 \pmod{8}$, then

$$[\Gamma_{p,l}, \Gamma_{p,l}]^{ab} \cong \begin{cases} \mathbb{Z}_2^2 \times \mathbb{Z}_{16}^3, & \text{if } p, l \equiv 1 \pmod{3} \\ \mathbb{Z}_3 \times \mathbb{Z}_{16}^3, & \text{else.} \end{cases}$$

Conjecture 10. *Let $p, l \equiv 3 \pmod{4}$.*

If $p \pmod{8} = l \pmod{8}$, then

$$[\Gamma_{p,l}, \Gamma_{p,l}]^{ab} \cong \begin{cases} \mathbb{Z}_2^2 \times \mathbb{Z}_8^2 \times \mathbb{Z}_{64}, & \text{if } p, l \equiv 1 \pmod{3} \\ \mathbb{Z}_8^2 \times \mathbb{Z}_{64}, & \text{if } p = 3 \text{ or } l = 3 \\ \mathbb{Z}_3 \times \mathbb{Z}_8^2 \times \mathbb{Z}_{64}, & \text{else.} \end{cases}$$

If $p \pmod{8} \neq l \pmod{8}$, then

$$[\Gamma_{p,l}, \Gamma_{p,l}]^{ab} \cong \begin{cases} \mathbb{Z}_2^2 \times \mathbb{Z}_8^2 \times \mathbb{Z}_{16}, & \text{if } p, l \equiv 1 \pmod{3} \\ \mathbb{Z}_8^2 \times \mathbb{Z}_{16}, & \text{if } p = 3 \text{ or } l = 3 \\ \mathbb{Z}_3 \times \mathbb{Z}_8^2 \times \mathbb{Z}_{16}, & \text{else.} \end{cases}$$

The groups appearing in Conjecture 11 are again the same as in Conjecture 10:

Conjecture 11. *Let $p \equiv 3 \pmod{4}$ and $l \equiv 1 \pmod{4}$.*

If $l \equiv 1 \pmod{8}$, then

$$[\Gamma_{p,l}, \Gamma_{p,l}]^{ab} \cong \begin{cases} \mathbb{Z}_2^2 \times \mathbb{Z}_8^2 \times \mathbb{Z}_{64}, & \text{if } p, l \equiv 1 \pmod{3} \\ \mathbb{Z}_8^2 \times \mathbb{Z}_{64}, & \text{if } p = 3 \\ \mathbb{Z}_3 \times \mathbb{Z}_8^2 \times \mathbb{Z}_{64}, & \text{else.} \end{cases}$$

If $l \equiv 5 \pmod{8}$, then

$$[\Gamma_{p,l}, \Gamma_{p,l}]^{ab} \cong \begin{cases} \mathbb{Z}_2^2 \times \mathbb{Z}_8^2 \times \mathbb{Z}_{16}, & \text{if } p, l \equiv 1 \pmod{3} \\ \mathbb{Z}_8^2 \times \mathbb{Z}_{16}, & \text{if } p = 3 \\ \mathbb{Z}_3 \times \mathbb{Z}_8^2 \times \mathbb{Z}_{16}, & \text{else.} \end{cases}$$

Let $\Lambda_{p,l}$ be the following subgroup of $\Gamma_{p,l}$.

$$\Lambda_{p,l} := \psi_{p,l}(\{x = x_0 + x_1i + x_2j + x_3k \in \mathbb{H}(\mathbb{Z}); x_0 \text{ odd}; |x|^2 = p^s l^t, s, t \in 2\mathbb{N}_0\}).$$

Observe that $\Lambda_{p,l}$ is the kernel of the surjective homomorphism $\Gamma_{p,l} \rightarrow \mathbb{Z}_2 \times \mathbb{Z}_2$ determined by

$$\psi_{p,l}(x) \mapsto \begin{cases} (1 + 2\mathbb{Z}, 0 + 2\mathbb{Z}), & \text{if } |x|^2 = p \\ (0 + 2\mathbb{Z}, 1 + 2\mathbb{Z}), & \text{if } |x|^2 = l, \end{cases}$$

in particular $\Lambda_{p,l}$ is a normal subgroup of $\Gamma_{p,l}$ of index 4. It seems that the abelianization of $\Lambda_{p,l}$ does not depend on p and l , if $p, l > 3$.

Conjecture 12. *Let p, l be any pair of distinct odd prime numbers. Then*

$$\Lambda_{p,l}^{ab} \cong \begin{cases} \mathbb{Z}_2 \times \mathbb{Z}_8^2, & \text{if } p = 3 \text{ or } l = 3 \\ \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_8^2, & \text{else.} \end{cases}$$

Since the conjectured abelianizations of the groups $\Gamma_{p,l}$, $[\Gamma_{p,l}, \Gamma_{p,l}]$ and $\Lambda_{p,l}$ are never 2-generated, we also get the following conjecture:

Conjecture 13. *Let p, l be any pair of distinct odd prime numbers. Then the groups $\Gamma_{p,l}$, $[\Gamma_{p,l}, \Gamma_{p,l}]$ and $\Lambda_{p,l}$ are not 2-generated.*

REFERENCES

- [1] Kimberley, Jason S.; Robertson, Guyan, *Groups acting on products of trees, tiling systems and analytic K-theory*, New York J. Math. **8**(2002), 111–131 (electronic).
- [2] Rattaggi, Diego, *Computations in groups acting on a product of trees: normal subgroup structures and quaternion lattices*, Ph.D. thesis, ETH Zürich, 2004.
- [3] Rattaggi, Diego, *Anti-tori in square complex groups*, to appear in *Geom. Dedicata*. A preprint is available at arXiv:math.GR/0411547.
- [4] Rattaggi, Diego, *On infinite groups generated by two quaternions*, Preprint 2005, see arXiv:math.GR/0502512.
- [5] Rattaggi, Diego; Robertson, Guyan, *Abelian subgroup structure of square complex groups and arithmetic of quaternions*, J. Algebra **286**(2005), no. 1, 57–68.

UNIVERSITÉ DE GENÈVE, SECTION DE MATHÉMATIQUES, 2–4 RUE DU LIÈVRE, CP 64, CH–1211 GENÈVE 4, SWITZERLAND

E-mail address: rattaggi@math.unige.ch