# Error bounds for semi-Galerkin approximations of nonhomogeneous incompressible fluids

P. Braz e Silva\*†and M. A. Rojas-Medar<sup>‡§</sup>

Math Subject Classification: 35Q30, 76M22, 65M15, 65M60 Keywords: Semi-Galerkin, Navier-Stokes equations, nonhomogeneous fluids

#### Abstract

We consider spectral semi-Galerkin approximations for the strong solutions of the nonhomogeneous Navier-Stokes equations. We derive an optimal uniform in time error bound in the  $H^1$  norm for approximations of the velocity. We also derive an error estimate for approximations of the density in some spaces  $L^r$ .

### 1 Introduction

Let  $\Omega \subseteq \mathbb{R}^n$ , n=2 or 3, a  $C^{1,1}$ -regular bounded domain, and T>0. We are interested in the initial boundary value problem

$$\begin{cases}
\rho(\mathbf{u}_{t} + \mathbf{u} \cdot \nabla \mathbf{u}) - \Delta \mathbf{u} + \nabla p = \rho \mathbf{f} & \text{in } \Omega \times [0, T), \\
\operatorname{div} \mathbf{u} = 0 & \text{in } \Omega \times (0, T), \\
\rho_{t} + \mathbf{u} \cdot \nabla \rho = 0 & \text{in } \Omega \times (0, T), \\
\mathbf{u} = 0 & \text{on } \partial \Omega \times [0, T), \\
\mathbf{u}(x, 0) = \mathbf{u}_{0}(x), \forall x \in \Omega, \\
\rho(x, 0) = \rho_{0}(x), \forall x \in \Omega.
\end{cases}$$
(1.1)

These are the equations of motion for nonhomogeneous incompressible fluids. The unknowns are the velocity  $\mathbf{u}(x,t) \in \mathbb{R}^n$  of the fluid, its density  $\rho(x,t) \in \mathbb{R}$  and the hydrostatic pressure  $p(x,t) \in \mathbb{R}$ . The functions  $\mathbf{u}_0(x)$  and  $\rho_0(x)$  are respectively the initial velocity and initial density. The function  $\mathbf{f}(x,t)$  is the density by unit of mass of the external force acting on the fluid. Here, without loss of generality to our aim, the viscosity is considered to be one.

<sup>\*</sup>Departamento de Matemática - Universidade Federal de Pernambuco CEP 50740-540 Recife, PE, Brazil (pablo@dmat.ufpe.br)

<sup>&</sup>lt;sup>‡</sup>Instituto de Matemática, Estatística e Computação Científica - UNICAMP, Cx. Postal 6065, CEP 13083-970, Campinas, SP, Brazil (marko@ime.unicamp.br)

 $<sup>^\</sup>dagger P.$  Braz e Silva was supported for this work by FAPESP/Brazil, Grant #02/13270-1 and is currently supported in part by CAPES/MECD-DGU Brazil/Spain, Grant #2137-05-4.

 $<sup>\</sup>S$  M. Rojas-Medar is partially supported by CNPq-Brazil, Grant #301354/03-0, CAPES/MECD-DGU Brazil/Spain, grant #2137-05-4 and project BFM2003-06446-CO-01, Spain.

Different methods have been used to study the existence, uniqueness and regularity of solutions for problem (1.1)(see [2], [3], [5], [8] [9], [10], [11], [12], [13], [14], [17], [18]). Here, we are interested in the spectral Semi-Galerkin method applied to system (1.1), the word spectral being used to indicate that the eigenfunctions of the associated Stokes operator are used as a basis of approximations. It is very important to derive error bounds for Galerin methods, due to the wide application of these methods in numerical experiments. Even the case of spectral Galerkin method may be used as a preparation and guide for the more practical finite element Galerkin method.

A systematic development of error bounds for the spectral Galerkin method applied to the classical Navier-Stokes equations was given in [15]. These error bounds are local in time in the sense that they depend on functions which grow exponentially with time. As observed in [6], this is the best one may expect without any assumptions about the stability of the solution being approximated. Optimal uniform in time error estimates for the velocity in the Dirichlet norm were also derived in [6], assuming uniform boundedness in time of the  $L^2$ -norm of the gradient of the velocity and exponential stability in the Dirichlet norm of the solution.

For the variable density case, error bounds were first obtained in [16], where local in time error bounds were derived. Moreover, a result of uniform in time error estimates in the  $L^2$  norm was also stated. This last result, however, is not optimal. Moreover, it requires the assumption  $\mathbf{u} \in L^{\infty}(0,T,\mathbf{H}^3(\Omega))$ . As pointed out in [7], this assumption is pretty restrictive, since it requires a global compatibility condition on the initial data even for the classical Navier-Stokes equations. Error bounds were also derived in [4] without explicitly assuming stability, but requiring exponential decay of the external force field. This hypothesis though is very restrictive as well, since gravitational forces do not satisfy it.

Here we derive error bounds assuming the solution  $(\mathbf{u}, \rho)$  to be  $p_0$ -conditionally asymptotically stable, a notion to be defined in section 3. The number  $p_0$  is required to satisfy  $6 \leq p_0 \leq \infty$ , and is related with the regularity of allowed perturbations of the density equation. In [6], a similar notion has been used to treat the classical Navier-Stokes equations (see also [16]). Here, we adapt it in the proper way to be used in the variable density case. With this assumption, we obtain an uniform in time optimal error bound in the Dirichlet norm for the velocity. An error bound depending on time for the density in some spaces  $L^r$  is also derived.

In section 2 we state some preliminary useful results. In section 3 we describe the approximation method, the stability notion to be used, and state the main result. Section 4 is dedicated to deriving a priori estimates. Finally, we present the proof of the main result in section 5.

To simplify the notation, we denote by C a generic finite positive constant depending only on  $\Omega$  and the other fixed parameters of the problem. As usual, it may have different values in different expressions. When necessary, we emphasize that the constants may have different values using the notation  $C_1$ ,  $C_2$ , and so on.

### 2 Preliminaries

Throughout this work, we consider the usual Sobolev spaces

$$W^{m,q}(D) = \left\{ f \in L^q(D), : ||\partial^{\alpha} f||_{L^q(D)} < +\infty, |\alpha| \le m \right\},\,$$

for a multi-index  $\alpha$ , a nonnegative integer m and  $1 \leq q \leq +\infty$ , where the domain is  $D = \Omega$  or  $D = \Omega \times (0,T)$ ,  $0 < T \leq +\infty$ , depending on the context. We write  $H^m(D) := W^{m,2}(D)$  and denote by  $H_0^m(D)$  the closure of  $C_0^\infty(D)$  in  $H^m(D)$ . If B is a Banach space, we denote by  $L^q([0,T];B)$  the Banach space of B-valued functions defined on the interval [0,T] that are  $L^q$ -integrable in Bochner's sense. Spaces of  $\mathbb{R}^n$  valued functions, as well as their elements, are represented by bold face letters. We write

$$\mathbf{C}_{0,\sigma}^{\infty}(\Omega) := \{ \mathbf{v} \in \mathbf{C}_0^{\infty}(\Omega) / \text{ div } \mathbf{v} = 0 \text{ in } \Omega \},$$

and denote by **H** and **V** the closure of  $\mathbf{C}_{0,\sigma}^{\infty}(\Omega)$  in  $\mathbf{L}^{2}(\Omega)$  and  $\mathbf{H}^{1}(\Omega)$  respectively.

Throughout this work, the orthogonal projection from  $\mathbf{L}^2(\Omega)$  onto  $\mathbf{H}$  is written as P. Thus, the well known Stokes operator is  $-P\Delta$ , defined on  $\mathbf{V} \cap \mathbf{H}^2(\Omega)$ . Its eigenfunctions and eigenvalues are denoted by  $\mathbf{w}^k$  and  $\lambda_k$  respectively. The usual  $L^2(\Omega)$  inner product and norm are respectively indicated by  $(\cdot, \cdot)$ , and  $\|\cdot\|$ .

It is well known that  $\{\mathbf{w}^k(x)\}_{k=1}^{\infty}$  form an complete orthogonal system in the spaces  $\mathbf{H}$ ,  $\mathbf{V}$  and  $\mathbf{V} \cap \mathbf{H}^2(\Omega)$  equipped with the usual inner products  $(\mathbf{u}, \mathbf{v}), (\nabla \mathbf{u}, \nabla \mathbf{v})$  and  $(P\Delta \mathbf{u}, P\Delta \mathbf{v})$  respectively.

For each  $k \in \mathbb{N}$ , we denote by  $P_k$  the orthogonal projection from  $\mathbf{L}^2(\Omega)$  onto  $\mathbf{V}_k := \operatorname{span}[\mathbf{w}^1, ..., \mathbf{w}^k]$ . For all  $\mathbf{f}, \mathbf{g} \in \mathbf{L}^2(\Omega)$  and  $k, m \in \mathbb{N}$ , it holds

- (i)  $(P_k \mathbf{f}, \mathbf{g}) = (\mathbf{f}, P_k \mathbf{g}),$
- (ii)  $(P\mathbf{f}, \mathbf{g}) = (\mathbf{f}, P\mathbf{g}),$
- (iii)  $((P_m P_k)\mathbf{f}, \mathbf{g}) = (\mathbf{f}, (P_m P_k)\mathbf{g}),$
- (iv)  $((P P_k)\mathbf{f}, \mathbf{g}) = (\mathbf{f}, (P P_k)\mathbf{g}).$

The following bounds are useful to our ends.

Lemma 2.1 (Rautmann[15]). If  $v \in V$ , then

$$\|\boldsymbol{v} - P_k \boldsymbol{v}\|^2 \le \frac{1}{\lambda_{k+1}} \|\nabla \boldsymbol{v}\|^2.$$

Moreover, if  $\mathbf{v} \in \mathbf{V} \cap \mathbf{H}^2(\Omega)$ , then

$$\|\boldsymbol{v} - P_k \boldsymbol{v}\|^2 \le \frac{1}{\lambda_{k+1}^2} \|P \triangle \boldsymbol{v}\|^2,$$

$$\|\nabla \boldsymbol{v} - \nabla P_k \boldsymbol{v}\|^2 \le \frac{1}{\lambda_{k+1}} \|P \triangle \boldsymbol{v}\|^2.$$

Remark 2.2. From Lemma 2.1, it follows that if  $\mathbf{f} \in \mathbf{H}^1(\Omega)$ , then

$$\|(I - P_k)P\mathbf{f}\|^2 \le \frac{1}{\lambda_{k+1}} \|\nabla P\mathbf{f}\|^2.$$

Moreover, since  $P: \mathbf{H}^1(\Omega) \to \mathbf{H}^1(\Omega)$  is a continuous operator[19], we have

$$\|\nabla P\mathbf{f}\|^2 \le C\|\mathbf{f}\|_{\mathbf{H}^1}^2.$$

Thus, for all  $\mathbf{f} \in \mathbf{H}^1(\Omega)$ , one has

$$\|(I - P_k)P\mathbf{f}\| \le \frac{C}{\lambda_{k+1}}\|\mathbf{f}\|_{\mathbf{H}^1}^2$$

Since  $PP_k = P_k P = P_k$ , one equivalently obtains

$$||P\mathbf{f} - P_k\mathbf{f}||^2 \le \frac{C}{\lambda_{k+1}} ||\mathbf{f}||_{\mathbf{H}^1}^2.$$

Moreover, the above relations also hold if one replaces P by any  $P_m$ , m > k. Analogously, one may check that

$$||(I - P_k)P\mathbf{f}||^2 \le \frac{C}{\lambda_{k+1}^2} ||\mathbf{f}||_{\mathbf{H}^2}^2$$

for all  $\mathbf{f} \in \mathbf{H}^2(\Omega)$ . We also have, as an easy consequence of the  $L^2$ -orthogonality of the functions  $\{\mathbf{w}^k\}_{k=1}^{\infty}$ , the following: Let  $m > k, m, k \in \mathbb{N}$ ,  $\mathbf{f} \in \mathbf{L}^2(\Omega)$  and  $\mathbf{v}^m \in \mathbf{V}_m$ ,  $\mathbf{v}^k \in \mathbf{V}_k$ . Then

$$((P_m - P_k)\mathbf{f}, \mathbf{v}^m - \mathbf{v}^k) = (\mathbf{f}, (I - P_k)\mathbf{v}^m).$$

# 3 Stability concept and main result

We consider problem (1.1) for all time  $t \ge 0$ , and suppose the data given therein satisfy

$$\mathbf{u}_0 \in \mathbf{V} \cap \mathbf{H}^2(\Omega),\tag{3.1}$$

$$\sup_{t \ge 0} \|\mathbf{f}\|_{\mathbf{H}^1} < \infty \; ; \; \sup_{t \ge 0} \|\mathbf{f}_t\| < \infty, \tag{3.2}$$

$$\rho_0 \in C^1(\bar{\Omega}) \; ; \; 0 < \alpha \le \rho_0 \le \beta, \tag{3.3}$$

where  $\alpha$  and  $\beta$  are constants. We also suppose that there exists M > 0 such that the solution  $(\mathbf{u}, \rho)$  of (1.1) satisfies

$$\|\nabla \mathbf{u}(t)\| \le M \; ; \; \forall \, t \ge 0. \tag{3.4}$$

If n=2, then conditions (3.1) and (3.2) imply that (3.4) holds. If n=3, then inequality (3.4) holds for  ${\bf f}$  and  ${\bf u}_0$  sufficiently small (see [5]). For a given  $p_0$ ,  $6 \le p_0 \le \infty$ , we also assume  $({\bf u},\rho)$  to be  $p_0$ -conditionally asymptotically stable

(see [6, 16] for similar notions of stability). To define this notion of stability, we first define perturbations of system (1.1). The pair  $(\boldsymbol{\xi}(x,t),\eta(x,t))$ , defined for  $t \geq t_0 \geq 0$ , is called a perturbation of  $(\mathbf{u},\rho)$  at time  $t_0$  if  $(\widehat{\mathbf{u}} := \mathbf{u} + \boldsymbol{\xi}, \widehat{\rho} := \rho + \eta)$  is a solution of (1.1), with  $\boldsymbol{\xi}|_{\partial\Omega} = 0$ . Therefore, setting  $\boldsymbol{\xi}_0 := \boldsymbol{\xi}(\cdot,t_0), \, \eta_0 := \eta(\cdot,t_0),$  the pair  $(\boldsymbol{\xi},\widehat{\rho})$  is a solution of the initial boundary value problem

$$\begin{cases}
\widehat{\rho}\boldsymbol{\xi}_{t} + \widehat{\rho}(\mathbf{u} \cdot \nabla)\boldsymbol{\xi} + \widehat{\rho}(\boldsymbol{\xi} \cdot \nabla)\mathbf{u} + \widehat{\rho}(\boldsymbol{\xi} \cdot \nabla)\boldsymbol{\xi} + \nabla q = \\
= \Delta \boldsymbol{\xi} + (\rho - \widehat{\rho})(\mathbf{u}_{t} + (\mathbf{u} \cdot \nabla)\mathbf{u} - \mathbf{f}) & \text{in } \Omega \times (t_{0}, \infty), \\
\widehat{\rho}_{t} + ((\boldsymbol{\xi} + \mathbf{u}) \cdot \nabla)\widehat{\rho} = 0 & \text{in } \Omega \times (t_{0}, \infty), \\
\nabla \cdot \boldsymbol{\xi} = 0 & \text{in } \Omega \times (t_{0}, \infty), \\
\boldsymbol{\xi}|_{\partial\Omega} = 0, \\
\boldsymbol{\xi}(x, t_{0}) = \boldsymbol{\xi}_{0}(x), \ \forall x \in \Omega, \\
\widehat{\rho}(x, t_{0}) = \rho(x, t_{0}) + \eta_{0}(x), \ \forall x \in \Omega.
\end{cases}$$
(3.5)

Now, for a given  $p_0$ ,  $6 \le p_0 \le \infty$ , we define the concept of  $p_0$ -conditional asymptotic stability.

**Definition 3.1.** The pair  $(\mathbf{u}, \rho)$  is said to be  $p_0$ -conditionally asymptotically stable if for all  $t_0 \geq 0$  there exist positive numbers  $A, B, \delta, M_1, M_2$  and a continuous decreasing function  $F: [0, \infty) \to \mathbb{R}^+, F(0) = 1$ ,  $\lim_{t \to \infty} F(t) = 0$  such that, for all  $\boldsymbol{\xi}_0 \in V \cap H^2(\Omega), \ \eta_0 \in L^{\infty}(\Omega) \cap W^{1,p_0}(\Omega)$ , satisfying  $\|\nabla \boldsymbol{\xi}_0\| < \delta$ ,  $\|P\Delta \boldsymbol{\xi}_0\| < A, \|\eta_0\|_{L^{\infty}} < B$ , problem (3.5) is uniquely solvable with

$$\boldsymbol{\xi} \in L^{2}_{loc}([t_{0}, \infty); \mathbf{V} \cap \mathbf{H}^{2}(\Omega)),$$
  
$$\boldsymbol{\xi}_{t} \in L^{2}_{loc}([t_{0}, \infty); \mathbf{H}^{1}(\Omega)),$$
  
$$\eta \in L^{\infty}([t_{0}, \infty); L^{\infty}(\Omega) \cap W^{1, p_{0}}(\Omega)).$$

Moreover,

$$\|\nabla \eta(\cdot, t)\|_{L^{p_0}} \le M_1, \, \forall \, t \ge t_0,$$
 (3.6)

$$\|\nabla \xi(\cdot, t)\| \le M_2 \|\nabla \xi_0\| F(t - t_0), \ \forall \ t \ge t_0. \tag{3.7}$$

Remark 3.2. We use a general function F(t) in Definition 3.1 just to stress out that the results here do not require an exponential decay rate.

The solution of problem (1.1) can be obtained through a spectral semi-Galerkin approximation, that is, a spectral Galerkin approximation

$$\mathbf{u}^{n}(x,t) = \sum_{k=1}^{n} C_{kn}(t)\mathbf{w}^{k}(x)$$

for the velocity **u**, uniquely determined by

$$(\rho^{n}\mathbf{u}_{t}^{n}, \boldsymbol{\phi}^{n}) + (\rho^{n}\mathbf{u}^{n} \cdot \nabla \mathbf{u}^{n}, \boldsymbol{\phi}^{n}) + (\nabla \mathbf{u}^{n}, \boldsymbol{\phi}^{n}) = (\rho^{n}\mathbf{f}, \boldsymbol{\phi}^{n}), \ t \ge 0, \ (3.8)$$
$$(\mathbf{u}^{n}(x, 0) - \mathbf{u}_{0}(x), \boldsymbol{\phi}^{n}) = 0, \tag{3.9}$$

for all  $\phi^n$  of the form  $\phi^n(x) = \sum_{k=1}^n d_k \mathbf{w}^k(x)$ , and an infinite dimensional approximation  $\rho^n$  for the density, solution of

$$\rho_t^n + \mathbf{u}^n \cdot \nabla \rho^n = 0,$$
  

$$\rho^n(0) = \rho_0.$$
(3.10)

It can be proved that  $(\mathbf{u}^n, \rho^n)$  converges in an appropriate sense to  $(\mathbf{u}, \rho)$ , solution of (1.1). Our main result is

**Theorem 3.3.** Suppose  $(u, \rho)$  to be  $p_0$ -conditionally asymptotically stable, for some  $p_0$ ,  $6 \le p_0 \le \infty$ . Then, there exist constants  $N \in \mathbb{N}$  and  $C \ge 0$  such that if  $n \ge N$  then, for all  $t \ge 0$ ,

$$\|\nabla \boldsymbol{u}(\cdot,t) - \nabla \boldsymbol{u}^{n}(\cdot,t)\| \le \frac{C}{(\lambda_{n+1})^{\frac{1}{2}}}.$$
(3.11)

Moreover, if  $6 \le p_0 < \infty$ , then

$$\|\rho(\cdot,t) - \rho^n(\cdot,t)\|_{L^r} \le \frac{Ct}{(\lambda_{n+1})^{\frac{1}{2}}}, \quad 2 \le r \le \frac{6p_0}{6+p_0},$$
 (3.12)

and if  $p_0 = \infty$ , then

$$\|\rho(\cdot,t) - \rho^n(\cdot,t)\|_{L^r} \le \frac{Ct}{(\lambda_{n+1})^{\frac{1}{2}}}, \ 2 \le r \le 6.$$
 (3.13)

The constants N, C, depend only on the domain, on the norms of the data in (3.1), (3.2) and on the constants introduced in (3.4) and definition 3.1.

## 4 A priori estimates

We first state a general simple result that will be used later on. A proof is given in the Appendix.

**Lemma 4.1.** Let h(t) be an integrable nonnegative function. Suppose there exist nonnegative constants  $a_1$ ,  $a_2$  satisfying

$$\int_{t_0}^t h(\tau)d\tau \le a_1(t - t_0) + a_2,$$

for all t,  $t_0$  with  $0 \le t_0 \le t$ . Then,

$$\sup_{t\geq 0} \left(e^{-t} \int_0^t e^{\tau} h(\tau) d\tau\right) < \infty.$$

For **u**, solution of (1.1), and the perturbations  $\boldsymbol{\xi}$ , we have:

**Lemma 4.2.** Given  $\epsilon > 0$ , the bounds

$$\frac{1}{2}\frac{d}{dt}\|\nabla u(\cdot,t)\|^{2} + \frac{\epsilon}{8}\|P\Delta u(\cdot,t)\|^{2} + \frac{1}{4}\|\rho^{\frac{1}{2}}u_{t}(\cdot,t)\|^{2} \leq$$
(4.1)

$$C\left(\|\mathbf{f}(\cdot,t)\|^2 + \|\nabla \mathbf{u}(\cdot,t)\|^6\right),$$

$$\|\boldsymbol{\xi}_{t}(\cdot,t)\|^{2} + \frac{1}{2}\frac{d}{dt}\|\nabla\boldsymbol{\xi}(\cdot,t)\|^{2} + \frac{\epsilon}{42}\|P\Delta\boldsymbol{\xi}(\cdot,t)\|^{2} \leq$$
(4.2)

$$C(\|\nabla \xi(\cdot,t)\|^2\|P\Delta u(\cdot,t)\|^2 + \|\nabla \xi(\cdot,t)\|^6 + \|u_t(\cdot,t)\|^2 + \|P\Delta u(\cdot,t)\|^2),$$

hold for all  $t \geq 0$ .

*Proof.* Inequality (4.1) was proved in [18]. Inequality (4.2) can be proved in a completely analogous way.

Corollary 4.3. For all  $t \geq t_0$ , we have

$$\int_{t_0}^{t} \|P\Delta u(\cdot, \tau)\|^2 d\tau \le C + C(t - t_0), \tag{4.3}$$

$$\int_{t_0}^{t} \|\boldsymbol{u}_t(\cdot,\tau)\|^2 d\tau \le C + C(t-t_0), \tag{4.4}$$

$$\int_{t_0}^{t} \|P\Delta \xi(\cdot, \tau)\|^2 d\tau \le C + C(t - t_0), \tag{4.5}$$

$$\int_{t_0}^{t} \|\boldsymbol{\xi}_t(\cdot, \tau)\|^2 d\tau \le C + C(t - t_0). \tag{4.6}$$

Moreover, combining inequalities (4.3), (4.4), (4.5) and (4.6) with lemma 4.1, one gets

$$\sup_{t \ge 0} e^{-t} \int_0^t e^{\tau} \| \mathbf{u}_t(\cdot, \tau) \|^2 d\tau < \infty, \tag{4.7}$$

$$\sup_{t\geq 0} e^{-t} \int_0^t e^{\tau} \|P\Delta \mathbf{u}(\cdot,\tau)\|^2 d\tau < \infty, \tag{4.8}$$

$$\sup_{t>0} e^{-t} \int_0^t e^{\tau} \|\boldsymbol{\xi}_t(\cdot, \tau)\|^2 d\tau < \infty, \tag{4.9}$$

$$\sup_{t\geq 0} e^{-t} \int_0^t e^{\tau} \|P\Delta \boldsymbol{\xi}(\cdot, \tau)\|^2 d\tau < \infty. \tag{4.10}$$

The following lemma states some bounds for  ${\bf u}$  which are very important for our later arguments.

#### Lemma 4.4. We have

$$\sup_{t\geq 0} \|\boldsymbol{u}_t(\cdot,t)\| < \infty, \tag{4.11}$$

$$\sup_{t>0} \|P\Delta \mathbf{u}(\cdot, t)\| < \infty, \tag{4.12}$$

$$\sup_{t\geq 0} e^{-t} \int_0^t e^{\tau} \|\nabla \mathbf{u}_t(\cdot, \tau)\|^2 d\tau < \infty. \tag{4.13}$$

*Proof.* We begin by proving (4.12), supposing (4.11) holds. Setting  $\mathbf{v} = -P\Delta\mathbf{u}$  in the weak formulation of problem (1.1), we get

$$-(\rho \mathbf{u}_t, P\Delta \mathbf{u}) - (\rho \mathbf{u} \cdot \nabla \mathbf{u}, P\Delta \mathbf{u}) + ||P\Delta \mathbf{u}||^2 = -(\rho \mathbf{f}, P\Delta \mathbf{u}).$$

Thus,

$$\begin{split} \|P\Delta\mathbf{u}\| & \leq \|\rho\mathbf{u}_t\| + \|\rho\mathbf{u} \cdot \nabla\mathbf{u}\| + \|\rho\mathbf{f}\| \\ & \leq \beta\|\mathbf{u}_t\| + \beta\|\mathbf{u}\|_{L^6} \|\nabla\mathbf{u}\|_{L^3} + \beta\|\mathbf{f}\| \\ & \leq \beta\|\mathbf{u}_t\| + C\beta\|P\Delta\mathbf{u}\|^{\frac{1}{2}} \|\nabla\mathbf{u}\|^{\frac{3}{2}} + \beta\|\mathbf{f}\| \\ & \leq \beta\left(\|\mathbf{u}_t\| + \|\mathbf{f}\|\right) + \frac{(C\beta)^2}{2} \|\nabla\mathbf{u}\|^3 + \frac{1}{2} \|P\Delta\mathbf{u}\| \\ & \leq 2\beta\|\mathbf{u}_t\| + 2\beta\|\mathbf{f}\| + (C\beta)^2 M^3 + \frac{1}{2} \|P\Delta\mathbf{u}\|. \end{split}$$

Therefore, by (4.11) and (3.2), we have

$$\sup_{t\geq 0} \|P\Delta \mathbf{u}(\cdot,t)\| \leq 2\beta \sup_{t\geq 0} \|\mathbf{u}_t(\cdot,t)\| + 2\beta \sup_{t\geq 0} \|\mathbf{f}(\cdot,t)\| + (C\beta)^2 M^3 < \infty,$$

which proves (4.12). To prove (4.11) and (4.13), differentiate the weak formulation of problem (1.1), and set  $\mathbf{v} = \mathbf{u}_t$  to get

$$(\rho_t \mathbf{u}_t, \mathbf{u}_t) + (\rho \mathbf{u}_{tt}, \mathbf{u}_t) + ((\rho(\mathbf{u} \cdot \nabla)\mathbf{u})_t, \mathbf{u}_t) + (\nabla \mathbf{u}_t, \nabla \mathbf{u}_t) = ((\rho \mathbf{f})_t, \mathbf{u}_t).$$

Therefore,

$$\frac{1}{2} \frac{d}{dt} \| \sqrt{\rho} \mathbf{u}_t \|^2 + \| \nabla \mathbf{u}_t \|^2 = -\frac{1}{2} \int_{\Omega} \rho \mathbf{u} \cdot \nabla (\mathbf{u}_t \cdot \mathbf{u}_t) \, dx 
- ((\rho (\mathbf{u} \cdot \nabla) \mathbf{u})_t, \mathbf{u}_t) + ((\rho \mathbf{f})_t, \mathbf{u}_t). \quad (4.14)$$

Now, bound each term on the right hand side of (4.14) as follows:

$$\begin{vmatrix}
-\frac{1}{2} \int_{\Omega} \rho \mathbf{u} \cdot \nabla (\mathbf{u}_{t} \cdot \mathbf{u}_{t}) dx \\
\leq \|\rho\|_{\infty} \|\mathbf{u}\|_{L^{4}} \|\mathbf{u}_{t}\|_{L^{4}} \|\nabla \mathbf{u}_{t}\| \\
\leq C\beta \|\nabla \mathbf{u}\| \|\nabla \mathbf{u}_{t}\|_{4} \|\mathbf{u}_{t}\|_{4}^{\frac{1}{4}} \|\nabla \mathbf{u}_{t}\|_{4}^{\frac{3}{4}} \\
= C\beta \|\nabla \mathbf{u}\| \|\mathbf{u}_{t}\|_{4}^{\frac{1}{4}} \|\nabla \mathbf{u}_{t}\|_{4}^{\frac{7}{4}} \\
\leq C_{\epsilon} (C\beta \|\nabla \mathbf{u}\|)^{8} \|\mathbf{u}_{t}\|^{2} + \epsilon \|\nabla \mathbf{u}_{t}\|^{2},$$

$$|(\rho \mathbf{f}_t, \mathbf{u}_t)| \le \frac{\beta^2}{2} ||\mathbf{f}_t||^2 + \frac{1}{2} ||\mathbf{u}_t||^2,$$

$$|(\rho_{t}\mathbf{f},\mathbf{u}_{t})| = \left| \int_{\Omega} \rho_{t}\mathbf{f} \cdot \mathbf{u}_{t} dx \right| = \left| -\int_{\Omega} \operatorname{div}(\rho \mathbf{u})\mathbf{f} \cdot \mathbf{u}_{t} dx \right| = \left| -\int_{\Omega} \rho \mathbf{u} \cdot \nabla(\mathbf{f} \cdot \mathbf{u}_{t}) dx \right|$$

$$\leq \|\rho\|_{L^{\infty}} \|\mathbf{u}\| \|\nabla(\mathbf{f} \cdot \mathbf{u}_{t})\| \leq C_{\epsilon} (C\beta \|\nabla \mathbf{u}\| \|\nabla \mathbf{f}\|)^{2} + \epsilon \|\nabla \mathbf{u}_{t}\|^{2}$$

$$\leq C_{\epsilon} (C\beta M \|\nabla \mathbf{f}\|)^{2} + \epsilon \|\nabla \mathbf{u}_{t}\|^{2},$$

$$|(\rho(\mathbf{u}_t \cdot \nabla)\mathbf{u}, \mathbf{u}_t)| \leq \|\rho\|_{L^{\infty}} \|\mathbf{u}_t\|_{L^4}^2 \|\nabla \mathbf{u}\| \leq C\beta \|\nabla \mathbf{u}\| \|\mathbf{u}_t\|^{\frac{1}{2}} \|\nabla \mathbf{u}_t\|^{\frac{3}{2}}$$
$$\leq C_{\epsilon} (C\beta \|\nabla \mathbf{u}\| \|\mathbf{u}_t\|^{\frac{1}{2}})^4 + \epsilon \|\nabla \mathbf{u}_t\|^2 \leq C_{\epsilon} C^4 \beta^4 M^4 \|\mathbf{u}_t\|^2 + \epsilon \|\nabla \mathbf{u}_t\|^2$$

$$|(\rho(\mathbf{u}\cdot\nabla)\mathbf{u}_t,\mathbf{u}_t)| \leq \|\rho\|_{L^{\infty}}\|\mathbf{u}_t\|^{\frac{1}{4}}\|\nabla\mathbf{u}\|\|\nabla\mathbf{u}_t\|^{\frac{7}{4}} \leq C_{\epsilon}(C\beta M)^8\|\mathbf{u}_t\|^2 + \epsilon\|\nabla\mathbf{u}_t\|^2,$$

$$|(\rho_t(\mathbf{u} \cdot \nabla)\mathbf{u}, \mathbf{u}_t)| = |\int_{\Omega} \rho_t(\mathbf{u} \cdot \nabla)\mathbf{u} \cdot \mathbf{u}_t| = |-\int_{\Omega} \operatorname{div}(\rho \mathbf{u})(\mathbf{u} \cdot \nabla)\mathbf{u} \cdot \mathbf{u}_t|$$

$$\leq CC_{\epsilon} \beta^2 M^4 ||P\Delta \mathbf{u}||^2 + 4\epsilon ||\nabla \mathbf{u}_t||^2.$$

Therefore, from equation (4.14) one obtains

$$\frac{1}{2} \frac{d}{dt} \| \sqrt{\rho} \mathbf{u}_t \|^2 + \| \nabla \mathbf{u}_t \|^2 \le C_{\epsilon} (C\beta M)^8 \| \mathbf{u}_t \|^2 + \frac{\beta^2}{2} \| \mathbf{f}_t \|^2 + C_{\epsilon} (C\beta M \| \nabla \mathbf{f} \|)^2 
+ C_{\epsilon} C^4 \beta^4 M^4 \| \mathbf{u}_t \|^2 + C C_{\epsilon} \beta^2 M^4 \| P \Delta \mathbf{u} \|^2 + \left( 8\epsilon + \frac{1}{2} \right) \| \nabla \mathbf{u}_t \|^2.$$
(4.15)

Now, choose  $\epsilon < \frac{1}{16}$  to conclude

$$\frac{d}{dt} \|\sqrt{\rho} \mathbf{u}_t(\cdot, t)\|^2 + \widetilde{C} \|\nabla \mathbf{u}_t(\cdot, t)\|^2 \le C + C \|P\Delta \mathbf{u}(\cdot, t)\|^2 + C \|\mathbf{u}_t(\cdot, t)\|^2,$$
(4.16)

where  $\widetilde{C} > 0$  is an absolute constant, and the constant C depends only on  $\Omega$ ,  $\|\rho\|_{L^{\infty}}$ ,  $\|\nabla \mathbf{f}\|$ ,  $\|\mathbf{f}_t\|$ ,  $\sup_{t\geq 0} \|\nabla \mathbf{u}\|$ . Now, multiplying inequality (4.16) by  $e^t$  and integrating over [0,t], one gets

$$e^{t} \|\sqrt{\rho} \mathbf{u}_{t}(\cdot, t)\|^{2} + \widetilde{C} \int_{0}^{t} e^{\tau} \|\nabla \mathbf{u}_{t}(\cdot, \tau)\|^{2} d\tau \leq \|(\sqrt{\rho} \mathbf{u}_{t})(\cdot, 0)\|^{2} + C \int_{0}^{t} e^{\tau} \|P\Delta \mathbf{u}(\cdot, \tau)\|^{2} d\tau + C \int_{0}^{t} e^{\tau} d\tau + C \int_{0}^{t} e^{\tau} \|\mathbf{u}_{t}(\cdot, \tau)\|^{2} d\tau.$$

Hence,

$$\|\sqrt{\rho}\mathbf{u}_{t}(\cdot,t)\|^{2} + \widetilde{C}e^{-t} \int_{0}^{t} e^{\tau} \|\nabla\mathbf{u}_{t}(\cdot,\tau)\|^{2} d\tau \leq e^{-t} \beta \|\mathbf{u}_{t}(\cdot,0)\|^{2} + Ce^{-t} \int_{0}^{t} e^{\tau} \|P\Delta\mathbf{u}(\cdot,\tau)\|^{2} d\tau + Ce^{-t} \int_{0}^{t} e^{\tau} d\tau + Ce^{-t} \int_{0}^{t} e^{\tau} \|\mathbf{u}_{t}(\cdot,\tau)\|^{2} d\tau.$$

Using inequalities (4.7) and (4.8), we get the desired result.

Corollary 4.5. For all  $t_0, t \in \mathbb{R}$ ,  $0 \le t_0 \le t$ , one has

$$\int_{t_0}^{t} \|\nabla u_t(\cdot, \tau)\|^2 d\tau \le C(t - t_0) + C. \tag{4.17}$$

*Proof.* Integrating inequality (4.16) from  $t_0$  to t, one gets

$$\|\sqrt{\rho}\mathbf{u}_{t}(\cdot,t)\|^{2} + \widetilde{C} \int_{t_{0}}^{t} \|\nabla\mathbf{u}_{t}(\cdot,\tau)\|^{2} d\tau \leq \|\sqrt{\rho}\mathbf{u}_{t}(\cdot,t_{0})\|^{2} + C \int_{t_{0}}^{t} d\tau + C \int_{t_{0}}^{t} \|P\Delta\mathbf{u}(\cdot,\tau)\|^{2} d\tau + C \int_{t_{0}}^{t} \|\mathbf{u}_{t}(\cdot,\tau)\|^{2} d\tau \leq \|\sqrt{\rho}\mathbf{u}_{t}(\cdot,t_{0})\|^{2} + C(t-t_{0}) + C(t-t_{0}) \left(\sup_{t\geq 0} \|P\Delta\mathbf{u}(\cdot,t)\|^{2}\right) + C(t-t_{0}) \left(\sup_{t\geq 0} \|\mathbf{u}_{t}(\cdot,t)\|^{2}\right).$$

Using inequalities (4.11) and (4.12), one obtains the bound (4.17).

A priori estimates for the solution  $\boldsymbol{\xi}$  of problem (3.5), similar to those in Lemma 4.4 for  $\mathbf{u}$ , also hold. Indeed, if  $\|\nabla \boldsymbol{\xi}_0\| < \delta$ , where  $\delta$  is the number referred to in Definition 3.1, then it follows by (3.7) that  $\|\nabla \boldsymbol{\xi}(\cdot,t)\| \leq \delta M_2$ . Therefore, the function  $\widehat{\mathbf{u}} = \mathbf{u} + \boldsymbol{\xi}$  is a solution of the nonhomogeneous Navier-Stokes equations satisfying  $\|\nabla \widehat{\mathbf{u}}\| \leq M + \delta M_2$ . Moreover, if  $\|P\Delta \boldsymbol{\xi}(\cdot,t_0)\|$  is bounded then  $\|P\Delta \widehat{\mathbf{u}}(\cdot,t_0)\|$  is also bounded. In this case, analogously to the proof of Lemma 4.4, one can bound  $\|P\Delta \widehat{\mathbf{u}}(\cdot,t)\|$ , for  $t \geq t_0$ . This bound implies that  $\|P\Delta \boldsymbol{\xi}(\cdot,t)\|$  is bounded, for  $t \geq t_0$ . Summarizing,

**Lemma 4.6.** For perturbations  $\boldsymbol{\xi}$  satisfying  $\|\nabla \boldsymbol{\xi}_0\| < \delta$  and  $\|P\Delta \boldsymbol{\xi}_0\| \leq C_0$ ,  $C_0 > 0$ , we have  $\|P\Delta \boldsymbol{\xi}(\cdot,t)\| \leq C$ , for all  $t \geq t_0$ . The constant C depends on  $\|P\Delta \boldsymbol{\xi}_0\|$ ,  $C_0$ ,  $\Omega$ , the initial data of problem (1.1) and on the norms and constants appearing in (3.6) and (3.7).

It also holds

**Lemma 4.7.** The function  $\xi$  satisfies

$$\int_{t_0}^t \|\nabla \xi_t(\cdot, \tau)\|^2 d\tau \le C(t - t_0) + C, \tag{4.18}$$

for all  $t_0$ , t,  $0 \le t_0 \le t$ .

*Proof.* Note that

$$\int_{t_0}^t \|\nabla \boldsymbol{\xi}_t(\cdot, \tau)\|^2 d\tau \le C \left( \int_{t_0}^t \|\nabla \widehat{\mathbf{u}}_t(\cdot, \tau)\|^2 d\tau + \int_{t_0}^t \|\nabla \mathbf{u}_t(\cdot, \tau)\|^2 d\tau \right).$$

The second term on the right hand side of this inequality is bounded (Corollary 4.5). Therefore, it only remains to bound  $\int_{t_0}^t \|\nabla \widehat{\mathbf{u}}_t(\cdot,\tau)\|^2 d\tau$ . This bound follows analogously to the bound for  $\mathbf{u}$ .

Now, let  $\mathbf{u} = \sum_{k=1}^{\infty} C_k(t) \mathbf{w}^k(x)$  be the expression of  $\mathbf{u}$ , the solution of (1.1), in

terms of the eigenfunctions of the Stokes operator. Let  $\mathbf{v}^n := \sum_{k=1}^n C_k(t) \mathbf{w}^k(x)$ 

be its n-th partial sum, and define  $\mathbf{e}^n := \mathbf{u} - \mathbf{v}^n$  and  $\boldsymbol{\psi}^n := \mathbf{u}^n - \mathbf{v}^n$ . We begin by bounding  $\mathbf{e}^n$ .

Lemma 4.8. The bounds

$$\|\nabla \mathbf{e}^{n}(\cdot,t)\|^{2} \leq \frac{C}{\lambda_{n+1}} \tag{4.19}$$

and

$$\|\mathbf{e}^{n}(\cdot,t)\|^{2} \leq \frac{C}{\lambda_{n+1}^{2}}$$
 (4.20)

hold for all  $t \geq 0$ .

Proof. We bound

$$\|\nabla \mathbf{e}^{n}(\cdot,t)\|^{2} = \|\nabla \sum_{k=n+1}^{\infty} C_{k}(t)\mathbf{w}^{k}(\cdot)\|^{2} \leq \frac{1}{\lambda_{n+1}} \|P\Delta \sum_{k=n+1}^{\infty} C_{k}(t)\mathbf{w}^{k}(\cdot)\|^{2}$$

$$\leq \frac{C}{\lambda_{n+1}} \|P\Delta \mathbf{u}(\cdot,t)\|^{2} \leq \frac{C}{\lambda_{n+1}}, \tag{4.21}$$

as desired. Moreover,

$$\|\mathbf{e}^{n}(\cdot,t)\|^{2} = \|\sum_{k=n+1}^{\infty} C_{k}(t)\mathbf{w}^{k}(\cdot)\|^{2} \leq \frac{1}{\lambda_{n+1}} \|\nabla \sum_{k=n+1}^{\infty} C_{k}(t)\mathbf{w}^{k}(\cdot)\|^{2}$$

$$\leq \frac{1}{\lambda_{n+1}} \|\nabla \mathbf{e}^{n}(\cdot,t)\|^{2} \leq \frac{C}{\lambda_{n+1}^{2}}.$$
(4.22)

Now, we prove that a suitable bound for  $\|\nabla \psi^n(\cdot,t)\|$  implies in a bound for  $\|P\Delta \psi^n(\cdot,t)\|$ .

**Lemma 4.9.** If for some constant K > 0 the inequality  $\|\nabla \psi^n(\cdot,t)\|^2 \leq \frac{K}{\lambda_{n+1}}$  holds on an interval  $[0,t^*]$ , then there exists C > 0, independent of n, such that

$$||P\Delta\psi^n(\cdot,t)||^2 \le C,\tag{4.23}$$

for all  $t \in [0, t^*]$ .

**Proof:** Since  $\psi^n = \mathbf{u}^n - \mathbf{v}^n$  and  $\sup_{t \geq 0} \|P\Delta \mathbf{v}^n(\cdot,t)\| \leq \sup_{t \geq 0} \|P\Delta \mathbf{u}(\cdot,t)\| < \infty$ , one only needs to bound  $\|P\Delta \mathbf{u}^n\|$ . To this end, note that

$$\|\nabla \psi^n\|^2 = \|\nabla \mathbf{u}^n - \nabla \mathbf{v}^n\|^2 \le \frac{K}{\lambda_{n+1}}.$$

Therefore,

$$\|\nabla \mathbf{u}^n\| \le \left(\frac{K}{\lambda_{n+1}}\right)^{\frac{1}{2}} + \|\nabla \mathbf{v}^n\| \le \left(\frac{K}{\lambda_1}\right)^{\frac{1}{2}} + M.$$

We also have, for  $\mathbf{u}^n$ , the bounds

$$\int_{t_0}^t \|P\Delta \mathbf{u}^n(\cdot,\tau)\|^2 d\tau \le C + C(t-t_0),$$
$$\int_{t_0}^t \|\mathbf{u}_t^n(\cdot,\tau)\|^2 d\tau \le C + C(t-t_0),$$

which are analogous to the bounds (4.3) and (4.4) for  $\mathbf{u}$ , and can be proved by analogous arguments. Therefore, by Lemma 4.1,

$$\sup_{t\geq 0} e^{-t} \int_0^t e^{\tau} \|\mathbf{u}_t^n(\cdot, \tau)\|^2 d\tau \leq C,$$
  
$$\sup_{t\geq 0} e^{-t} \int_0^t e^{\tau} \|P\Delta \mathbf{u}^n(\cdot, \tau)\|^2 d\tau \leq C.$$

Using these inequalities, one can show that

$$\frac{d}{dt} \|\sqrt{\rho} \mathbf{u}_{t}^{n}\|^{2} + \widetilde{C} \|\nabla \mathbf{u}_{t}^{n}\|^{2} \le C + C \|P\Delta \mathbf{u}^{n}\|^{2} + C \|\mathbf{u}_{t}^{n}\|^{2}, \tag{4.24}$$

for all  $t \in [0, t^*]$ . At this point, we need to restrict the time interval, since the constant C depends also on  $\sup_{t>0} \|\nabla \mathbf{u}^n\|$ , and we can assure this term to be

bounded, uniformly with respect to n, only in the interval  $[0, t^*]$ . Using inequality (4.24), it follows that

$$\|\mathbf{u}_t^n(\cdot,t)\| \le C. \tag{4.25}$$

Finally, inequality (4.25) allows one to prove

$$||P\Delta \mathbf{u}^n(\cdot, t)|| < C, \tag{4.26}$$

for all  $t \in [0, t^*]$ . We do not give the details of the proof, since it is completely analogous to the proof of Lemma 4.4.

Now we bound, for later use, the term  $\nabla P_n(\psi^n - \xi) = \nabla \psi^n - \nabla P_n \xi$ . First, note that  $\mathbf{v}^n$  satisfies

$$(\rho \mathbf{u}_t, \boldsymbol{\phi}^n) + (\nabla \mathbf{v}^n, \nabla \boldsymbol{\phi}^n) + (\rho \mathbf{u} \cdot \nabla \mathbf{u}, \boldsymbol{\phi}^n) = (\rho \mathbf{f}, \boldsymbol{\phi}^n), \tag{4.27}$$

for all  $\phi^n$  of the form  $\phi^n(x) = \sum_{k=1}^n d_k \mathbf{w}^k(x)$ . Subtracting equation (3.8) from equation (4.27), we get

$$(\rho^{n}\boldsymbol{\psi}_{t}^{n},\boldsymbol{\phi}^{n}) + (\nabla\boldsymbol{\psi}^{n},\nabla\boldsymbol{\phi}^{n}) = (\rho^{n}\mathbf{e}_{t}^{n},\boldsymbol{\phi}^{n}) + ((\rho-\rho^{n})(\mathbf{u}_{t}+\mathbf{u}\cdot\nabla\mathbf{u}-\mathbf{f}),\boldsymbol{\phi}^{n}) + (\rho^{n}\mathbf{u}\cdot\nabla\mathbf{u},\boldsymbol{\phi}^{n}) - (\rho^{n}\mathbf{u}^{n}\cdot\nabla\mathbf{u}^{n},\boldsymbol{\phi}^{n}).$$
(4.28)

On the other hand, taking the inner product of the fist equation in (3.5) with  $\phi^n$  and integrating by parts, one gets

$$(\widehat{\rho}\boldsymbol{\xi}_{t},\boldsymbol{\phi}^{n}) + (\widehat{\rho}\mathbf{u} \cdot \nabla\boldsymbol{\xi},\boldsymbol{\phi}^{n}) + (\widehat{\rho}\boldsymbol{\xi} \cdot \nabla\mathbf{u},\boldsymbol{\phi}^{n}) + (\widehat{\rho}\boldsymbol{\xi} \cdot \nabla\boldsymbol{\xi},\boldsymbol{\phi}^{n}) + (\nabla\boldsymbol{\xi},\nabla\boldsymbol{\phi}^{n}) = ((\rho - \rho^{n})(\mathbf{u}_{t} + \mathbf{u} \cdot \nabla\mathbf{u} - \mathbf{f}),\boldsymbol{\phi}^{n}). \quad (4.29)$$

Subtract equation (4.29) from equation (4.28) to get

$$(\rho^{n}\boldsymbol{\theta}_{t},\boldsymbol{\phi}^{n}) + (\nabla\boldsymbol{\theta},\nabla\boldsymbol{\phi}^{n}) = ((\widehat{\rho}-\rho^{n})(\mathbf{u}_{t}+\mathbf{u}\cdot\nabla\mathbf{u}-\mathbf{f}),\boldsymbol{\phi}^{n}) + ((\widehat{\rho}-\rho^{n})\boldsymbol{\xi}_{t},\boldsymbol{\phi}^{n}) + (\rho^{n}\mathbf{e}_{t}^{n},\boldsymbol{\phi}^{n}) + (\rho^{n}\mathbf{u}\cdot\nabla\mathbf{u},\boldsymbol{\phi}^{n}) - (\rho^{n}\mathbf{u}^{n}\cdot\nabla\mathbf{u}^{n},\boldsymbol{\phi}^{n}) + (\widehat{\rho}\mathbf{u}\cdot\nabla\boldsymbol{\xi},\boldsymbol{\phi}^{n}) + (\widehat{\rho}\boldsymbol{\xi}\cdot\nabla\mathbf{u},\boldsymbol{\phi}^{n}) + (\widehat{\rho}\boldsymbol{\xi}\cdot\nabla\boldsymbol{\xi},\boldsymbol{\phi}^{n}), \quad (4.30)$$

where  $\theta := \psi^n - \xi$ . Now, since  $P_n \theta = P_n (\psi^n - \xi) = \psi^n - P_n \xi$ , one has

$$(\rho^{n}\mathbf{u}\cdot\nabla\psi^{n},\phi^{n}) = (\rho^{n}\mathbf{u}\cdot\nabla P_{n}\boldsymbol{\theta},\phi^{n}) + (\rho^{n}\mathbf{u}\cdot\nabla P_{n}\boldsymbol{\xi},\phi^{n}),$$

$$(\rho^{n}\psi^{n}\cdot\nabla\mathbf{u},\phi^{n}) = (\rho^{n}P_{n}\boldsymbol{\theta}\cdot\nabla\mathbf{u},\phi^{n}) + (\rho^{n}P_{n}\boldsymbol{\xi}\cdot\nabla\mathbf{u},\phi^{n}),$$

$$(\rho^{n}\psi^{n}\cdot\nabla\psi^{n},\phi^{n}) = (\rho^{n}\psi^{n}\cdot\nabla P_{n}\boldsymbol{\theta},\phi^{n}) + (\rho^{n}\psi^{n}\cdot\nabla P_{n}\boldsymbol{\xi},\phi^{n}).$$

Therefore,

$$(\rho^{n}\mathbf{u}\cdot\nabla\mathbf{u},\phi^{n})-(\rho^{n}\mathbf{u}^{n}\cdot\nabla\mathbf{u}^{n},\phi^{n})=(\rho^{n}\psi^{n}\cdot\nabla\mathbf{e}^{n},\phi^{n})+(\rho^{n}\mathbf{e}^{n}\cdot\nabla\psi^{n},\phi^{n})$$

$$+(\rho^{n}\mathbf{u}\cdot\nabla\mathbf{e}^{n},\phi^{n})+(\rho^{n}\mathbf{e}^{n}\cdot\nabla\mathbf{v}^{n},\phi^{n})-(\rho^{n}\mathbf{u}\cdot\nabla P_{n}\boldsymbol{\theta},\phi^{n})-(\rho^{n}\mathbf{u}\cdot\nabla P_{n}\boldsymbol{\xi},\phi^{n})$$

$$-(\rho^{n}P_{n}\boldsymbol{\theta}\cdot\nabla\mathbf{u},\phi^{n})-(\rho^{n}P_{n}\boldsymbol{\xi}\cdot\nabla\mathbf{u},\phi^{n})-(\rho^{n}\psi^{n}\cdot\nabla P_{n}\boldsymbol{\theta},\phi^{n})-(\rho^{n}\psi^{n}\cdot\nabla P_{n}\boldsymbol{\xi},\phi^{n}).$$

Moreover, since  $\boldsymbol{\xi} = P_n \boldsymbol{\xi} + Q_n \boldsymbol{\xi}$ , one can show, after some computations, that

$$\begin{split} &(\rho^{n}\mathbf{u}\cdot\nabla\mathbf{u},\phi^{n})-(\rho^{n}\mathbf{u}^{n}\cdot\nabla\mathbf{u}^{n},\phi^{n})+(\widehat{\rho}\mathbf{u}\cdot\nabla\boldsymbol{\xi},\phi^{n})+(\widehat{\rho}\boldsymbol{\xi}\cdot\nabla\mathbf{u},\phi^{n})+(\widehat{\rho}\boldsymbol{\xi}\cdot\nabla\boldsymbol{\xi},\phi^{n})=\\ &(\rho^{n}\boldsymbol{\psi}^{n}\cdot\nabla\mathbf{e}^{n},\phi^{n})+(\rho^{n}\mathbf{e}^{n}\cdot\nabla\boldsymbol{\psi}^{n},\phi^{n})+(\rho^{n}\mathbf{u}\cdot\nabla\mathbf{e}^{n},\phi^{n})+(\rho^{n}\mathbf{e}^{n}\cdot\nabla\mathbf{v}^{n},\phi^{n})\\ &-(\rho^{n}\mathbf{u}\cdot\nabla P_{n}\boldsymbol{\theta},\phi^{n})-(\rho^{n}P_{n}\boldsymbol{\theta}\cdot\nabla\mathbf{u},\phi^{n})+((\widehat{\rho}-\rho^{n})\mathbf{u}\cdot\nabla P_{n}\boldsymbol{\xi},\phi^{n})\\ &+((\widehat{\rho}-\rho^{n})P_{n}\boldsymbol{\xi}\cdot\nabla\mathbf{u},\phi^{n})+(\widehat{\rho}\mathbf{u}\cdot\nabla Q_{n}\boldsymbol{\xi},\phi^{n})+(\widehat{\rho}Q_{n}\boldsymbol{\xi}\cdot\nabla\mathbf{u},\phi^{n})-(\rho^{n}\boldsymbol{\psi}^{n}\cdot\nabla P_{n}\boldsymbol{\theta},\phi^{n})\\ &-(\rho^{n}P_{n}\boldsymbol{\theta}\cdot\nabla P_{n}\boldsymbol{\xi},\phi^{n})+(\rho^{n}P_{n}\boldsymbol{\xi}\cdot\nabla Q_{n}\boldsymbol{\xi},\phi^{n})+(\rho^{n}Q_{n}\boldsymbol{\xi}\cdot\nabla P_{n}\boldsymbol{\xi},\phi^{n})\\ &+(\rho^{n}Q_{n}\boldsymbol{\xi}\cdot\nabla Q_{n}\boldsymbol{\xi},\phi^{n})+((\widehat{\rho}-\rho^{n})\boldsymbol{\xi}\cdot\nabla\boldsymbol{\xi},\phi^{n}). \end{split}$$

Applying this identity to (4.30), and taking  $\phi^n = P_n \theta_t$ , one obtains

$$\|\sqrt{\rho^{n}}\boldsymbol{\theta}_{t}\|^{2} + \frac{1}{2}\frac{d}{dt}\|\nabla P_{n}\boldsymbol{\theta}\|^{2} = ((\widehat{\rho} - \rho^{n})(\mathbf{u}_{t} + \mathbf{u} \cdot \nabla \mathbf{u} - \mathbf{f} + \boldsymbol{\xi}_{t} + \mathbf{u} \cdot \nabla P_{n}\boldsymbol{\xi}), P_{n}\boldsymbol{\theta}_{t})$$

$$+ ((\widehat{\rho} - \rho^{n})(P_{n}\boldsymbol{\xi} \cdot \nabla \mathbf{u} + \boldsymbol{\xi} \cdot \nabla \boldsymbol{\xi}), P_{n}\boldsymbol{\theta}_{t}) - (\rho^{n}\boldsymbol{\theta}_{t}, Q_{n}\boldsymbol{\xi}_{t})$$

$$+ (\rho^{n}\mathbf{e}_{t}^{n}, P_{n}\boldsymbol{\theta}_{t}) + (\rho^{n}\boldsymbol{\psi}^{n} \cdot \nabla \mathbf{e}^{n}, P_{n}\boldsymbol{\theta}_{t}) + (\rho^{n}\mathbf{e}^{n} \cdot \nabla \boldsymbol{\psi}^{n}, P_{n}\boldsymbol{\theta}_{t})$$

$$+ (\rho^{n}\mathbf{u} \cdot \nabla \mathbf{e}^{n}, P_{n}\boldsymbol{\theta}_{t}) + (\rho^{n}\mathbf{e}^{n} \cdot \nabla \mathbf{v}^{n}, P_{n}\boldsymbol{\theta}_{t}) - (\rho^{n}\mathbf{u} \cdot \nabla P_{n}\boldsymbol{\theta}, P_{n}\boldsymbol{\theta}_{t})$$

$$- (\rho^{n}P_{n}\boldsymbol{\theta} \cdot \nabla \mathbf{u}, P_{n}\boldsymbol{\theta}_{t}) + (\widehat{\rho}\mathbf{u} \cdot \nabla Q_{n}\boldsymbol{\xi}, P_{n}\boldsymbol{\theta}_{t}) + (\widehat{\rho}Q_{n}\boldsymbol{\xi} \cdot \nabla \mathbf{u}, P_{n}\boldsymbol{\theta}_{t})$$

$$- (\rho^{n}\boldsymbol{\psi}^{n} \cdot \nabla P_{n}\boldsymbol{\theta}, P_{n}\boldsymbol{\theta}_{t}) - (\rho^{n}P_{n}\boldsymbol{\theta} \cdot \nabla P_{n}\boldsymbol{\xi}, P_{n}\boldsymbol{\theta}_{t})$$

$$+ (\rho^{n}P_{n}\boldsymbol{\xi} \cdot \nabla Q_{n}\boldsymbol{\xi}, P_{n}\boldsymbol{\theta}_{t}) + (\rho^{n}Q_{n}\boldsymbol{\xi} \cdot \nabla P_{n}\boldsymbol{\xi}, P_{n}\boldsymbol{\theta}_{t})$$

$$+ (\rho^{n}Q_{n}\boldsymbol{\xi} \cdot \nabla Q_{n}\boldsymbol{\xi}, P_{n}\boldsymbol{\theta}_{t}) .$$

Now, we bound each term on the right hand side of identity (4.31). Given  $\epsilon > 0$ , we bound

$$\begin{split} |(\rho^{n}\boldsymbol{\theta}_{t},Q_{n}\boldsymbol{\xi}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}\|\nabla\boldsymbol{\xi}_{t}\|^{2}, \\ |(\rho^{n}\mathbf{e}_{t}^{n},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}\|\nabla\mathbf{u}_{t}\|^{2}, \\ |(\rho^{n}\boldsymbol{\psi}^{n}\cdot\nabla\mathbf{e}^{n},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}\|P\Delta\boldsymbol{\psi}^{n}\|^{2}, \\ |(\rho^{n}\mathbf{e}^{n}\cdot\nabla\boldsymbol{\psi}^{n},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}\|P\Delta\boldsymbol{\psi}^{n}\|^{2}, \\ |(\rho^{n}\mathbf{u}\cdot\nabla\mathbf{e}^{n},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}, \\ |(\rho^{n}\mathbf{e}^{n}\cdot\nabla\mathbf{v}^{n},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}, \\ |(\rho^{n}\mathbf{u}\cdot\nabla P_{n}\boldsymbol{\theta},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + C(\epsilon)\|\nabla P_{n}\boldsymbol{\theta}\|^{2}, \\ |(\rho^{n}P_{n}\boldsymbol{\theta}\cdot\nabla\mathbf{u},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + C(\epsilon)\|\nabla P_{n}\boldsymbol{\theta}\|^{2}, \\ |(\hat{\rho}\mathbf{u}\cdot\nabla Q_{n}\boldsymbol{\xi},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}\|P\Delta\boldsymbol{\xi}\|^{2}, \\ |(\hat{\rho}Q_{n}\boldsymbol{\xi}\cdot\nabla\mathbf{u},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}\|P\Delta\boldsymbol{\xi}\|^{2}, \\ |(\rho^{n}P_{n}\boldsymbol{\theta}\cdot\nabla P_{n}\boldsymbol{\xi},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + C(\epsilon)\|P\Delta\boldsymbol{\psi}^{n}\|^{2}\|\nabla P_{n}\boldsymbol{\theta}\|^{2}, \\ |(\rho^{n}P_{n}\boldsymbol{\theta}\cdot\nabla P_{n}\boldsymbol{\xi},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + C(\epsilon)\|P\Delta\boldsymbol{\xi}\|^{2}\|\nabla P_{n}\boldsymbol{\theta}\|^{2}, \\ |(\rho^{n}P_{n}\boldsymbol{\xi}\cdot\nabla Q_{n}\boldsymbol{\xi},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}\|P\Delta\boldsymbol{\xi}\|^{4}, \\ |(\rho^{n}Q_{n}\boldsymbol{\xi}\cdot\nabla P_{n}\boldsymbol{\xi},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}\|P\Delta\boldsymbol{\xi}\|^{4}, \\ |(\rho^{n}Q_{n}\boldsymbol{\xi}\cdot\nabla P_{n}\boldsymbol{\xi},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}\|P\Delta\boldsymbol{\xi}\|^{4}, \\ |(\rho^{n}Q_{n}\boldsymbol{\xi}\cdot\nabla P_{n}\boldsymbol{\xi},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}\|P\Delta\boldsymbol{\xi}\|^{4}, \\ |(\rho^{n}Q_{n}\boldsymbol{\xi}\cdot\nabla P_{n}\boldsymbol{\xi},P_{n}\boldsymbol{\theta}_{t})| & \leq \quad \epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}}\|P\Delta\boldsymbol{\xi}\|^{4}. \end{split}$$

It remains to bound  $|(\pi g^n, P_n \boldsymbol{\theta}_t)|$ , where  $\pi := \widehat{\rho} - \rho^n$  and  $\mathbf{g}^n := \mathbf{u}_t + \mathbf{u} \cdot \nabla \mathbf{u} - \mathbf{f} + \boldsymbol{\xi}_t + \mathbf{u} \cdot \nabla P_n \boldsymbol{\xi} + P_n \boldsymbol{\xi} \cdot \nabla \mathbf{u} + \boldsymbol{\xi} \cdot \nabla \boldsymbol{\xi}$ . We begin by bounding  $\mathbf{g}^n$ .

**Lemma 4.10.** For all  $p, 2 \le p \le 6$ , the bound

$$\|\mathbf{g}^{n}(\cdot,t)\|_{L^{p}}^{2} \leq C + C\|P\Delta\boldsymbol{\xi}(\cdot,t)\|^{2} + C\|P\Delta\boldsymbol{\xi}(\cdot,t)\|^{4} + C\|\nabla\boldsymbol{u}_{t}(\cdot,t)\|^{2} + C\|\nabla\boldsymbol{\xi}_{t}(\cdot,t)\|^{2}$$

$$(4.32)$$

holds for all  $t \geq 0$ .

*Proof.* Since  $2 \le p \le 6$ , we have

$$\begin{aligned} \|\mathbf{g}^{n}(\cdot,t)\|_{L^{p}}^{2} & \leq & C\left\{\|\mathbf{u}_{t}(\cdot,t)\|_{L^{p}}^{2} + \|\mathbf{u}\cdot\nabla\mathbf{u}(\cdot,t)\|_{L^{p}}^{2} + \|\mathbf{f}(\cdot,t)\|_{L^{p}}^{2} + \|\boldsymbol{\xi}_{t}(\cdot,t)\|_{L^{p}}^{2} \\ & + \|\mathbf{u}\cdot\nabla P_{n}\boldsymbol{\xi}(\cdot,t)\|_{L^{p}}^{2} + \|P_{n}\boldsymbol{\xi}\cdot\nabla\mathbf{u}(\cdot,t)\|_{L^{p}}^{2} + \|\boldsymbol{\xi}\cdot\nabla\boldsymbol{\xi}(\cdot,t)\|_{L^{p}}^{2}\right\} \\ & \leq & C\left\{\|\nabla\mathbf{u}_{t}(\cdot,t)\|^{2} + \|P\Delta\mathbf{u}(\cdot,t)\|^{4} + \|\nabla\mathbf{f}(\cdot,t)\|^{2} + \|\nabla\boldsymbol{\xi}_{t}(\cdot,t)\|^{2} \\ & + \|P\Delta\mathbf{u}(\cdot,t)\|^{2}\|P\Delta\boldsymbol{\xi}(\cdot,t)\|^{2} + \|P\Delta\boldsymbol{\xi}(\cdot,t)\|^{4}\right\} \\ & \leq & C + C\|P\Delta\boldsymbol{\xi}(\cdot,t)\|^{2} + C\|P\Delta\boldsymbol{\xi}(\cdot,t)\|^{4} + C\|\nabla\mathbf{u}_{t}(\cdot,t)\|^{2} + C\|\nabla\boldsymbol{\xi}_{t}(\cdot,t)\|^{2}. \end{aligned}$$

**Lemma 4.11.** If  $6 \le p_0 < \infty$ , then the bound

$$\|\pi(\cdot,t)\|_{L^{r}}^{2} \leq C\|\pi(\cdot,t_{0})\|_{L^{r}}^{2} + C(t-t_{0}) \int_{t_{0}}^{t} \|\nabla P_{n}\boldsymbol{\theta}(\cdot,\tau)\|^{2} d\tau + \frac{C}{\lambda_{n+1}} (t-t_{0})^{2} + \frac{C}{\lambda_{n+1}} (t-t_{0}) \int_{t_{0}}^{t} \|P\Delta\boldsymbol{\xi}(\cdot,\tau)\|^{2} d\tau$$
(4.33)

holds for all  $t \ge 0$  and all r,  $2 \le r \le \frac{6p_0}{6+p_0}$ . If  $p_0 = \infty$ , then the bound (4.33) is valid for all  $t \ge 0$  and all r,  $2 \le r \le 6$ .

*Proof.* First note that

$$\pi_t + \mathbf{u}^n \cdot \nabla \pi = (\mathbf{u}^n - \widehat{\mathbf{u}}) \cdot \nabla \widehat{\rho}.$$

Since  $\hat{\mathbf{u}} = \mathbf{u} + \boldsymbol{\xi}$ , we write the equation above as

$$\pi_t + \mathbf{u}^n \cdot \nabla \pi = (P_n \boldsymbol{\theta} - Q_n \boldsymbol{\xi} - \mathbf{e}^n) \cdot \nabla \widehat{\rho}. \tag{4.34}$$

Let r belonging to the suitable interval depending on the value of  $p_0$ . Multiply equation (4.34) by  $|\pi|^{r-1}$  and integrate to get

$$\frac{1}{r} \frac{d}{dt} \|\pi(\cdot, t)\|_{L^{r}}^{r} \leq \int_{\Omega} (P_{n} \boldsymbol{\theta} - Q_{n} \boldsymbol{\xi} - \mathbf{e}^{n}) \cdot \nabla \widehat{\rho} |\pi|^{r-1} dx$$

$$\leq \left( \int_{\Omega} |(P_{n} \boldsymbol{\theta} - Q_{n} \boldsymbol{\xi} - \mathbf{e}^{n}) \cdot \nabla \widehat{\rho}|^{r} dx \right)^{\frac{1}{r}} \left( \int_{\Omega} |\pi|^{r} dx \right)^{\frac{r-1}{r}}$$

$$\leq \|(P_{n} \boldsymbol{\theta} - Q_{n} \boldsymbol{\xi} - \mathbf{e}^{n}) \cdot \nabla \widehat{\rho}\|_{L^{r}} \|\pi\|_{L^{r}}^{r-1}.$$

Therefore,

$$\frac{d}{dt} \|\pi(\cdot,t)\|_{L^r} \leq \|(P_n\boldsymbol{\theta} - Q_n\boldsymbol{\xi} - \mathbf{e}^n) \cdot \nabla \widehat{\rho}\|_{L^r} \leq \|P_n\boldsymbol{\theta} - Q_n\boldsymbol{\xi} - \mathbf{e}^n\|_{L^p} \|\nabla \widehat{\rho}\|_{L^{p_0}},$$

where p is chosen as  $\frac{1}{p} = \frac{1}{r} - \frac{1}{p_0}$  if  $6 \le p_0 < \infty$ , and as p = r if  $p_0 = \infty$ . Note that in the case  $6 \le p_0 < \infty$ , this choice of p implies  $2 < \frac{2p_0}{p_0 - 2} \le p \le 6$ . In the case  $p_0 = \infty$ , we have  $p_0 = \infty$ .

$$\frac{d}{dt} \|\pi(\cdot,t)\|_{L^{r}} \leq C (\|\nabla P_{n}\boldsymbol{\theta}(\cdot,t)\| + \|\nabla Q_{n}\boldsymbol{\xi}(\cdot,t)\| + \|\nabla \mathbf{e}^{n}(\cdot,t)\|) 
\leq C \left(\|\nabla P_{n}\boldsymbol{\theta}(\cdot,t)\| + \frac{C}{(\lambda_{n+1})^{\frac{1}{2}}} + \frac{C}{(\lambda_{n+1})^{\frac{1}{2}}} \|P\Delta\boldsymbol{\xi}(\cdot,t)\|\right),$$

where, for the last inequality, we used Lemma 2.1 and inequality (4.19). Integrating this inequality from  $t_0$  to t,

$$\|\pi(\cdot,t)\|_{L^{r}}^{2} \leq C \left\{ \|\pi(\cdot,t_{0})\|_{L^{r}}^{2} + \left( \int_{t_{0}}^{t} \|\nabla P_{n}\boldsymbol{\theta}(\cdot,\tau)\|d\tau \right)^{2} + \left( \int_{t_{0}}^{t} \frac{1}{(\lambda_{n+1})^{\frac{1}{2}}}d\tau \right)^{2} + \left( \int_{t_{0}}^{t} \frac{1}{(\lambda_{n+1})^{\frac{1}{2}}} \|P\Delta\boldsymbol{\xi}(\cdot,\tau)\|d\tau \right)^{2} \right\}$$

$$\leq C \left\{ \|\pi(\cdot,t_{0})\|_{L^{r}}^{2} + (t-t_{0}) \int_{t_{0}}^{t} \|\nabla P_{n}\boldsymbol{\theta}(\cdot,\tau)\|^{2}d\tau + \frac{1}{\lambda_{n+1}}(t-t_{0})^{2} + \frac{1}{\lambda_{n+1}}(t-t_{0}) \int_{t_{0}}^{t} \|P\Delta\boldsymbol{\xi}(\cdot,\tau)\|^{2}d\tau \right\},$$

which is the desired bound.

Getting back to inequality (4.31), we have

$$\alpha \|\boldsymbol{\theta}_{t}\|^{2} + \frac{1}{2} \frac{d}{dt} \|\nabla P_{n} \boldsymbol{\theta}\|^{2} \leq 15\epsilon \|\boldsymbol{\theta}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}} \|\nabla \boldsymbol{\xi}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}} \|\nabla \mathbf{u}_{t}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}} \|P\Delta \boldsymbol{\psi}^{n}\|^{2}$$

$$+ \frac{C(\epsilon)}{\lambda_{n+1}} + \frac{C(\epsilon)}{\lambda_{n+1}} \|P\Delta \boldsymbol{\xi}\|^{2} + \frac{C(\epsilon)}{\lambda_{n+1}} \|P\Delta \boldsymbol{\xi}\|^{4} + C(\epsilon) \|\boldsymbol{\pi}\|_{L^{r}}^{2} \|\mathbf{g}^{n}\|_{L^{p}}^{2}$$

$$+ C(\epsilon) \|\nabla P_{n} \boldsymbol{\theta}\|^{2} \left\{1 + \|P\Delta \boldsymbol{\psi}^{n}\|^{2} + \|P\Delta \boldsymbol{\xi}\|^{2}\right\},$$

where, in the case  $6 \le p_0 < \infty$ , the inequality above holds for each  $r \in \left[3, \frac{6p_0}{6+p_0}\right]$ , with  $p \in \left[\frac{3p_0}{p_0-3}, 6\right]$  chosen such that  $\frac{1}{r} + \frac{1}{p} = \frac{1}{2}$ . In the case  $p_0 = \infty$ , it holds for all  $r \in [3,6]$ , with  $p \in [3,6]$  chosen such that  $\frac{1}{r} + \frac{1}{p} = \frac{1}{2}$ . Now fix  $\epsilon = \frac{1}{15} \left(\alpha - \frac{1}{2}\right)$ . Integrating the inequality from  $t_0$  to t, and using

Lemma 4.6, Lemma 4.9 and inequality (4.18), we get

$$\|\nabla P_{n}\boldsymbol{\theta}(\cdot,t)\|^{2} + \int_{t_{0}}^{t} \|\boldsymbol{\theta}_{t}(\cdot,\tau)\|^{2} d\tau \leq \|\nabla P_{n}\boldsymbol{\theta}(\cdot,t_{0})\|^{2} + \frac{C}{\lambda_{n+1}}(t-t_{0}) + \frac{C}{\lambda_{n+1}}\int_{t_{0}}^{t} \|\nabla \boldsymbol{\xi}_{t}(\cdot,\tau)\|^{2} d\tau + \frac{C}{\lambda_{n+1}}\int_{t_{0}}^{t} \|\nabla \mathbf{u}_{t}(\cdot,\tau) d\tau\|^{2} + C\int_{t_{0}}^{t} \|\pi(\cdot,\tau)\|_{L^{r}}^{2} \|\mathbf{g}^{n}(\cdot,\tau)\|_{L^{p}}^{2} d\tau + C\int_{t_{0}}^{t} \|\nabla P_{n}\boldsymbol{\theta}(\cdot,\tau)\|^{2} d\tau.$$
(4.35)

Adding inequalities (4.33) and (4.35), one obtains

$$\|\nabla P_{n}\boldsymbol{\theta}(\cdot,t)\|^{2} + \|\pi(\cdot,t)\|_{L^{r}}^{2} + \int_{t_{0}}^{t} \|\boldsymbol{\theta}_{t}(\cdot,\tau)\|^{2} d\tau \leq \|\nabla P_{n}\boldsymbol{\theta}(\cdot,t_{0})\|^{2} + C\|\pi(\cdot,t_{0})\|_{L^{r}}^{2}$$

$$+ \frac{C}{\lambda_{n+1}}(t-t_{0}) + \frac{C}{\lambda_{n+1}}(t-t_{0})^{2} + \frac{C}{\lambda_{n+1}}\int_{t_{0}}^{t} \|\nabla \boldsymbol{\xi}_{t}(\cdot,\tau)\|^{2} d\tau + \frac{C}{\lambda_{n+1}}\int_{t_{0}}^{t} \|\nabla \mathbf{u}_{t}(\cdot,\tau) d\tau\|^{2}$$

$$+ C\int_{t_{0}}^{t} \|\pi(\cdot,\tau)\|_{L^{r}}^{2} \|\mathbf{g}^{n}(\cdot,\tau)\|_{L^{p}}^{2} d\tau + C\int_{t_{0}}^{t} \|\nabla P_{n}\boldsymbol{\theta}(\cdot,\tau)\|^{2} d\tau + C(t-t_{0})\int_{t_{0}}^{t} \|\nabla P_{n}\boldsymbol{\theta}(\cdot,\tau)\|^{2} d\tau.$$

$$(4.36)$$

Fixing  $\bar{t} > t_0$ , and using Corollary 4.5 and Lemma 4.7, one concludes that

$$\|\nabla P_{n}\boldsymbol{\theta}(\cdot,t)\|^{2} + \|\pi(\cdot,t)\|_{L^{r}}^{2} + \int_{t_{0}}^{t} \|\boldsymbol{\theta}_{t}(\cdot,\tau)\|^{2} d\tau \leq \|\nabla P_{n}\boldsymbol{\theta}(\cdot,t_{0})\|^{2}$$

$$+ C\|\pi(\cdot,t_{0})\|_{L^{r}}^{2} + \frac{C}{\lambda_{n+1}} + \frac{C}{\lambda_{n+1}}(t-t_{0}) + \frac{C}{\lambda_{n+1}}(t-t_{0})^{2}$$

$$+ C\int_{t_{0}}^{t} \|\pi(\cdot,\tau)\|_{L^{r}}^{2} \|\mathbf{g}^{n}(\cdot,\tau)\|_{L^{p}}^{2} d\tau + C\int_{t_{0}}^{t} \{1 + (\bar{t}-t_{0})\} \|\nabla P_{n}\boldsymbol{\theta}(\cdot,\tau)\|^{2} d\tau,$$

$$(4.37)$$

for all  $t \in [t_0, \bar{t}]$ . Let  $\Lambda(t) := \|\nabla P_n \boldsymbol{\theta}(\cdot, t)\|^2 + \|\pi(\cdot, t)\|_{L^r}^2 + \int_{t_0}^t \|\boldsymbol{\theta}_t(\cdot, \tau)\|^2 d\tau$ . Thus, inequality (4.37) gives

$$\Lambda(t) \leq C\Lambda(t_0) + \frac{C}{\lambda_{n+1}} + \frac{C}{\lambda_{n+1}}(t - t_0) + \frac{C}{\lambda_{n+1}}(t - t_0)^2 + C\int_{t_0}^t \left\{1 + \bar{t} - t_0 + \|\mathbf{g}^n(\cdot, \tau)\|_{L^p}^2\right\} \Lambda(\tau) d\tau.$$

Applying a corollary of Gronwall's Lemma(see [1], page 90, corollary 6.2), one gets

$$\Lambda(t) \le \left( C\Lambda(t_0) + \frac{C}{\lambda_{n+1}} + \frac{C}{\lambda_{n+1}} (t - t_0) + \frac{C}{\lambda_{n+1}} (t - t_0)^2 \right) \exp\left\{ C \int_{t_0}^t (1 + \bar{t} - t_0 + \|g^n(\cdot, \tau)\|_{L^p}^2) d\tau \right\}.$$

We summarize the results in the following lemma.

**Lemma 4.12.** Suppose  $\|\nabla \psi^n(\cdot,t)\| \leq K/\lambda_{k+1}$  holds for a constant K > 0 and all t in a given interval  $0 \leq t_0 \leq t \leq \bar{t}$ . Let  $\boldsymbol{\xi}$  as in problem (3.5), and the

functions  $\pi$ ,  $\theta$ ,  $\mathbf{g}^n$  defined as before. If  $6 \leq p_0 < \infty$  then, for all  $t \in [t_0, \overline{t}]$ , one has

$$\|\nabla P_n \boldsymbol{\theta}(\cdot, t)\|^2 + \|\pi(\cdot, t)\|_{L^r}^2 + \int_{t_0}^t \|\boldsymbol{\theta}_t(\cdot, \tau)\|^2 d\tau \tag{4.38}$$

$$\leq C \left( \|\nabla P_n \boldsymbol{\theta}(\cdot, t_0)\|^2 + \|\pi(\cdot, t_0)\|_{L^r}^2 + \frac{1}{\lambda_{n+1}} + \frac{1}{\lambda_{n+1}} (t - t_0) + \frac{1}{\lambda_{n+1}} (t - t_0)^2 \right) e^{C \int_{t_0}^t a(\tau) d\tau},$$

for all  $r \in \left[3, \frac{6p_0}{6+p_0}\right]$ , and  $p \in \left[\frac{3p_0}{p_0-3}, 6\right]$  chosen such that  $\frac{1}{r} + \frac{1}{p} = \frac{1}{2}$ , where  $a(t) := 1 + \overline{t} - t_0 + \|\mathbf{g}^n(\cdot, t)\|_{L^p}^2$ . If  $p_0 = \infty$ , then the bound holds for all  $r \in [3, 6]$  and  $p \in [3, 6]$  such that  $\frac{1}{r} + \frac{1}{p} = \frac{1}{2}$ .

Remark 4.13. From inequalities (4.32), (4.17), (4.18), and Lemma 4.6, one can bound

$$\int_{t_0}^t a(\tau)d\tau \le \widetilde{C} + \widetilde{C}(t - t_0) + \widetilde{C}(\overline{t} - t_0)(t - t_0).$$

Therefore,

$$\exp\left\{\int_{t_0}^t a(\tau)d\tau\right\} \le \exp\left\{\widetilde{C} + \widetilde{C}(t - t_0) + \widetilde{C}(\overline{t} - t_0)(t - t_0)\right\}. \tag{4.39}$$

From now on, we fix the constants C and  $\widetilde{C}$  appearing in inequalities (4.38) and (4.39). We prove now that the bound for  $\|\nabla \psi^n\|$  required in lemma (4.9) and lemma (4.12) hold for n large enough.

**Proposition 4.14.** There exist K > 0 and  $N \in \mathbb{N}$  such that if  $n \geq N$ , then  $\|\nabla \psi^n(\cdot,t)\|^2 < \frac{K}{\lambda_{n+1}}$  for all  $t \geq 0$ .

Proof. Choose T such that  $M_2^2(F(T))^2 \leq \frac{1}{4}$ . Let  $K := 8C(1+T+T^2) \exp\{\widetilde{C} + \widetilde{C}T + \widetilde{C}T^2\}$  and let N to be large enough such that  $\frac{K}{\lambda_{n+1}} < \delta$  if  $n \geq N$ . Under these conditions, we have

$$\|\nabla \psi^n(\cdot, t)\|^2 < \frac{K}{\lambda_{n+1}},\tag{4.40}$$

for all  $t \ge 0$ . Indeed, suppose that inequality (4.40) does not hold. Thus, there exist  $n \ge N$  and  $t^* > 0$  such that

$$\|\nabla \psi^n(\cdot, t^*)\|^2 = \frac{K}{\lambda_{n+1}}.$$
 (4.41)

We may have either  $t^* \leq T$  or  $t^* > T$ . If  $t^* \leq T$ , consider  $t_0 = 0$ ,  $\xi = 0$ ,  $\eta = 0$ ,  $\bar{t} = t^*$ . In this case,  $\|\nabla P_n \boldsymbol{\theta}\| = \|\nabla \psi^n\|$ . Therefore, using Lemma 4.12, we have

$$\begin{split} \|\nabla \psi^n(\cdot,t^*)\|^2 + \|\pi(\cdot,t^*)\|_{L^r}^2 + \int_0^{t^*} \|\psi^n_t(\cdot,\tau)\|^2 d\tau & \leq & \left(\frac{C}{\lambda_{n+1}} + \frac{C}{\lambda_{n+1}}T + \frac{C}{\lambda_{n+1}}T^2\right) e^{\tilde{C} + \tilde{C}T + \tilde{C}T^2}, \\ & = & \frac{K}{8\lambda_{n+1}} < \frac{K}{\lambda_{n+1}}, \end{split}$$

which contradicts (4.41). If  $t^* > T$ , apply Lemma 4.12 with  $\bar{t} = t^*$ ,  $t_0 = t^* - T$  and  $\boldsymbol{\xi}(x,t)$ ,  $\eta(x,t)$  satisfying

$$\xi(x, t_0) = \psi^n(x, t_0), 
\eta(x, t_0) = \rho^n(x, t_0) - \rho(x, t_0),$$

to get

$$\begin{split} &\|\nabla \psi^{n}(\cdot,t^{*}) - \nabla P_{n}\xi(\cdot,t^{*})\|^{2} + \|\rho(\cdot,t^{*}) - \rho^{n}(\cdot,t^{*}) + \eta(\cdot,t^{*})\|_{L^{r}}^{2} \\ &+ \int_{t^{*}-T}^{t^{*}} \|\psi^{n}_{t}(\cdot,\tau) - \xi_{t}(\cdot,\tau)\|^{2} d\tau \leq \left(\frac{C}{\lambda_{n+1}} + \frac{C}{\lambda_{n+1}}T + \frac{C}{\lambda_{n+1}}T^{2}\right) e^{\tilde{C} + \tilde{C}T + \tilde{C}T^{2}} = \frac{K}{8\lambda_{n+1}}. \end{split}$$

Therefore,

$$\|\nabla \psi^{n}(\cdot, t^{*})\|^{2} \leq 2\left(\|\nabla \psi^{n}(\cdot, t^{*}) - \nabla P_{n} \boldsymbol{\xi}(\cdot, t^{*})\|^{2} + \|\nabla P_{n} \boldsymbol{\xi}(\cdot, t^{*})\|^{2}\right)$$

$$\leq 2\left(\frac{K}{8\lambda_{n+1}} + M_{2}^{2}\|\nabla \boldsymbol{\xi}(\cdot, t_{0})\|^{2}F(T)^{2}\right)$$

$$\leq 2\left(\frac{K}{8\lambda_{n+1}} + \frac{K}{4\lambda_{n+1}}\right) = \frac{3}{4}\frac{K}{\lambda_{n+1}} < \frac{K}{\lambda_{n+1}}$$

which again contradicts (4.41). This proves the proposition.

### 5 Proof of Theorem 3.3

Using the bounds (4.19) and (4.40), we directly get

$$\|\nabla \mathbf{u}(\cdot,t) - \nabla \mathbf{u}^n(\cdot,t)\|^2 \le 2(\|\nabla \psi^n(\cdot,t)\|^2 + \|\nabla \mathbf{e}^n(\cdot,t)\|^2) \le \frac{C}{\lambda_{n+1}},\tag{5.1}$$

which is the first bound stated in Theorem 3.3. In order to prove the bound (3.12) for the density, first note that

$$\rho_t + \mathbf{u} \cdot \nabla \rho = 0 \tag{5.2}$$

$$\rho_t^n + \mathbf{u}^n \cdot \nabla \rho^n = 0. \tag{5.3}$$

Subtracting equation (5.3) from equation (5.2), one gets

$$(\rho - \rho^n)_t + \mathbf{u}^n \cdot \nabla(\rho - \rho^n) = (\mathbf{u}^n - \mathbf{u}) \cdot \nabla \rho. \tag{5.4}$$

Now, if  $6 \le p_0 < \infty$ , let  $r \in \left[2, \frac{6p_0}{6+p_0}\right]$ . Choose  $p \in \left[\frac{2p_0}{p_0-2}, 6\right]$  such that  $\frac{1}{r} = \frac{1}{p} + \frac{1}{p_0}$ . Multiplying equation (5.4) by  $|\rho - \rho^n|^{r-1}$  and integrating over  $\Omega$ , we obtain

$$\frac{1}{r} \frac{d}{dt} \| \rho - \rho^n \|_{L^r}^r = \int_{\Omega} |\rho - \rho^n|^{r-1} (\mathbf{u}^n - \mathbf{u}) \cdot \nabla \rho dx$$

$$\leq \|\rho - \rho^n\|_{L^r}^{r-1} \| (\mathbf{u}^n - \mathbf{u}) \cdot \nabla \rho \|_{L^r}$$

$$\leq \|\rho - \rho^n\|_{L^r}^{r-1} \| \nabla \rho \|_{L^{p_0}} \| \mathbf{u}^n - \mathbf{u} \|_{L^p},$$

If  $p_0 = \infty$ , the bounds above hold for all  $r \in [2, 6]$  and p = r. Thus,

$$\frac{d}{dt} \|\rho(\cdot,t) - \rho^n(\cdot,t)\|_{L^r} \le C \|\mathbf{u}^n(\cdot,t) - \mathbf{u}(\cdot,t)\|_{L^r} \le C \|\nabla \mathbf{u}^n(\cdot,t) - \nabla \mathbf{u}(\cdot,t)\|. \tag{5.5}$$

Integrating inequality (5.5) from 0 to t and using (5.1), one gets

$$\|\rho(\cdot,t) - \rho^n(\cdot,t)\|_{L^r} \le \frac{C}{(\lambda_{n+1})^{\frac{1}{2}}} t + \|\rho(\cdot,0) - \rho^n(\cdot,0)\|_{L^r} = \frac{C}{(\lambda_{n+1})^{\frac{1}{2}}} t, \quad (5.6)$$

since  $\rho^n(x,0) = \rho_0(x)$ . This finishes the proof of Theorem 3.3.

### A Proof of Lemma 4.1

Suppose h(t) integrable, nonnegative and satisfying, for all  $t, t_0$  with  $0 \le t_0 \le t$ ,

$$\int_{t_0}^t h(\tau)d\tau \le a_1(t - t_0) + a_2,\tag{A.1}$$

where  $a_1$  and  $a_2$  are nonnegative constants. We first consider the case  $0 \le t \le 1$ . In this case,

$$e^{-t} \int_{t_0}^t e^{\tau} h(\tau) d\tau \le \int_{t_0}^t h(\tau) d\tau \le \int_0^1 h(\tau) d\tau \le a_1 + a_2.$$

Now, if t > 1, let  $n \in \mathbb{N}$  and  $r \in [0, 1)$  such that t = n + r. Then,

$$\int_{0}^{t} e^{\tau} h(\tau) d\tau = \int_{0}^{n+r} e^{\tau} h(\tau) d\tau = \sum_{j=1}^{n} \int_{j-1}^{j} e^{\tau} h(\tau) d\tau + \int_{n}^{n+r} e^{\tau} h(\tau) d\tau 
\leq \sum_{j=1}^{n} e^{j} \int_{j-1}^{j} h(\tau) d\tau + e^{n+r} \int_{n}^{n+r} h(\tau) d\tau 
\leq \sum_{j=1}^{n} e^{j} \int_{j-1}^{j} h(\tau) d\tau + e^{n+1} \int_{n}^{n+1} h(\tau) d\tau.$$

Inequality (A.1) implies

$$\int_{j-1}^{j} h(\tau)d\tau \le a_1 + a_2 \quad , \quad j = 1, \dots, n+1.$$

Therefore,

$$e^{-t} \int_0^t e^{\tau} h(\tau) d\tau = e^{-n-r} \int_0^{n+r} e^{\tau} h(\tau) d\tau = e^{-n-r} \sum_{j=1}^{n+1} e^j \int_{j-1}^j h(\tau) d\tau$$

$$\leq (a_1 + a_2) e^{-n-r} \sum_{j=1}^n e^j = (a_1 + a_2) e^{-n-r} \frac{e^{n+2} - e}{e - 1}$$

$$= (a_1 + a_2) \frac{e^{2-r} - e^{-n-r+1}}{e - 1} \leq (a_1 + a_2) \frac{e^2}{e - 1}.$$

Thus,

$$\sup_{t \ge 0} e^{-t} \int_0^t e^{\tau} h(\tau) d\tau \le (a_1 + a_2) \frac{e^2}{e - 1} < \infty,$$

which proves the lemma.

### References

- [1] H. Amann, Ordinary differential equations. an introduction to nonlinear analysis, de Gruyter Studies in Mathematics, vol. 13, Walter de Gruyter & Co., Berlin, 1990.
- [2] S. N. Antontsev, A. V. Kazhikhov, and V. N. Monakhov, Boundary value problems in mechanics of nonhomogeneous fluids, Studies in Mathematics and its Applications, vol. 22, North-Holland Publishing Co., Amsterdam, 1990.
- [3] J. L. Boldrini and M. A. Rojas-Medar, Global solutions to the equations for the motion of stratified incompressible fluids, Mat. Contemp. 3 (1992), 1–8.
- [4] \_\_\_\_\_, An error estimate uniform in time for spectral semi-Galerkin approximations of the nonhomogeneous Navier-Stokes equations, Numer. Funct. Anal. Optim. 15 (1994), no. 7-8, 755-778.
- [5] \_\_\_\_\_\_, Global strong solutions of the equations for the motion of nonhomogeneous incompressible fluids, Numerical methods in mechanics (Harlow) (C. Conca and G. N. Gatica, eds.), Pitman Res. Notes Math. Ser., no. 371, Longman, 1997, pp. 35–45.
- [6] J. G. Heywood, An error estimate uniform in time for spectral Galerkin approximations of the Navier-Stokes problem, Pacific J. Math 98 (1982), no. 2, 333–345.
- [7] J. G. Heywood and R. Rannacher, Finite element approximation of the nonstationary Navier-Stokes problem I: regularity of solutions and second order error estimates for spatial discretization, SIAM J. Numer. Anal. 19 (1982), no. 2, 275–311.
- [8] A. V. Kazhikov, Solvability of the initial-boundary value problem for the equations of motion of an inhomogeneous viscous incompressible fluid, Dokl. Akad. Nauk SSSR 216 (1974), 1008–1010.
- [9] J. U. Kim, Weak solutions of an initial-boundary value problems for an incompressible viscous fluid with nonnegative density, SIAM J. Math. Anal. 18 (1987), no. 1, 89–96.

- [10] O. A. Ladyzhenskaya and V. A. Solonnikov, The unique solvability of an initial-boundary value problem for viscous incompressible inhomogeneous fluids, Boundary value problems of mathematical physics, and related questions of the theory of functions 8, Zap. Naučn Sem. Leningrad. Otdel. Mat. Inst. Steklov (LOMI), vol. 52, 1975, pp. 52–109,218–219.
- [11] J.-L. Lions, On some problems connected with Navier-Stokes equations, Nonlinear evolution equations (Proc. Sympos., Univ. Wisconsin, Madison, Wis., 1977) (New York-London), Publ. Math. Res. Center Univ. Wisconsin, vol. 40, Academic Press, 1978, pp. 59–84.
- [12] \_\_\_\_\_\_, On some questions in boundary value problems of mathematical physics, Contemporary developments in continuum mechanics and partial differential equations (Proc. Internat. Sympos., Inst. Mat., Univ. Fed. Rio de Janeiro, Rio de Janeiro, 1977) (Amsterdam-New York), North-Holland Math. Stud., vol. 30, North-Holland, 1978, pp. 284–346.
- [13] P.-L. Lions, Mathematical topics in fluid mechanics. vol i: Incompressible models, Oxford Lecture Series in Mathematics and its Applications, vol. 3, The Clarendon Press, Oxford University Press, New York, 1996.
- [14] H. Okamoto, On the equation of nonstationary stratified fluid motion: uniqueness and existence of the solutions, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **30** (1984), no. 3, 615–643.
- [15] R. Rautmann, On the convergence rate of nonstationary Navier-Stokes approximations, Approximation methods for Navier-Stokes problems (Proc. Sympos., Univ. Paderborn, Paderborn, 1979) (Berlin), Lecture Notes in Math., vol. 771, Springer, 1980, pp. 425–449.
- [16] R. Salvi, Error estimates for the spectral Galerkin approximations of the solutions of Navier-Stokes type equations, Glasgow Math. J. **31** (1989), no. 2, 199–211.
- [17] \_\_\_\_\_, The equations of viscous incompressible nonhomogeneous fluid: on the existence and regularity, J. Austral. Math. Soc. Ser. B **33** (1991), no. 1, 94–110.
- [18] J. Simon, Nonhomogeneous viscous incompressible fluids: existence of velocity, density, and pressure, SIAM J. Math. Anal. 21 (1990), no. 5, 1093–1117.
- [19] W. von Wahl, *The equations of Navier-Stokes and abstract parabolic equations*, Aspects of Mathematics, vol. E8, Friedr. Vieweg & Sohn, Braunschweig, 1985.