

SESHADRI CONSTANTS VIA LELONG NUMBERS

THOMAS ECKL

ABSTRACT. One of Demailly's characterizations of Seshadri constants on ample line bundles works with Lelong numbers of certain positive singular hermitian metrics. In this note sections of multiples of the line bundle are used to produce such metrics and then to deduce another formula for Seshadri constants. It is applied to compute Seshadri constants on blown up products of curves, to disprove a conjectured characterization of maximal rationally connected quotients and to introduce a new approach to Nagata's conjecture.

In 1990 Demailly introduced Seshadri constants $\epsilon(L, x)$ for nef line bundles L on projective complex manifolds X [Dem92]:

$$\epsilon(L, x) := \inf_{C \ni x} \frac{L \cdot X}{\text{mult}_x C}$$

where the infimum is taken over all irreducible curves passing through x . They refine constants $\epsilon(L)$ appearing in Seshadri's ampleness criterion [Laz04, Thm.1.4.13] and quantify how much of the positivity of an ample line bundle can be localized at a given point.

These constants gained immediately a lot of interest in algebraic geometry; for example lower bounds on Seshadri constants were used to produce sections in adjoint bundles [Laz97]. It also turned out that explicit calculations of Seshadri constants are difficult in almost every concrete situation (see for example the work of Garcia [Gar05] on ruled surfaces) and it is not easier to give (interesting) upper and lower bounds for them. From their very definition it seems easier to determine upper bounds (by showing that a curve with appropriate intersection number and multiplicity exists) than lower bounds (via the non-existence of such curves) [EKL95, Bau99].

On the other hand Demailly gave two more equivalent definitions of Seshadri constants ([Dem92, Thm.6.4] or Prop. 6):

$$\epsilon(L, x) = \gamma(L, x) := \sup_{\gamma \in \mathbb{R}^+} \left\{ \begin{array}{l} \exists \text{ singular metric } h \text{ on } L : i\Theta_h \geq 0, \\ x \text{ isolated pole of } \Theta_h, \nu(\Theta_h, x) = \gamma \end{array} \right\},$$

where $\nu(\Theta_h, x)$ is the Lelong number of the curvature current $i\Theta_h$ in x , and

$$\epsilon(L, x) = \sigma(L, x) := \sup_{k \in \mathbb{N}} \frac{1}{k} s(kL, x)$$

where

$$s(kL, x) := \max_{s \in \mathbb{N}} \left\{ \begin{array}{l} \text{the sections in } H^0(X, kL) \text{ generate all } s\text{-jets in} \\ J_x^s kL = \mathcal{O}_X(kL)/\mathfrak{m}_x^{s+1} \end{array} \right\}.$$

1991 *Mathematics Subject Classification.* 32J25, 14J26, 14C20.

Key words and phrases. Seshadri constants, Lelong numbers, Nagata conjecture.

In the nef case we still have

$$\epsilon(L, x) \geq \gamma(L, x) \geq \sigma(L, x).$$

The definitions of $\sigma(L, x)$ and $\gamma(L, x)$ allow to give lower bounds for Seshadri constants by constructing sections of kL with special properties. For $\sigma(L, x)$ this is obvious whereas for $\gamma(L, x)$ we need a better understanding of singular hermitian metrics, their curvature currents and Lelong numbers. Let us first repeat the definitions (consult [Dem00] for further properties and examples):

A singular hermitian metric h on a holomorphic line bundle L is given in any trivialization $\theta : L|_{\Omega} \xrightarrow{\cong} \Omega \times \mathbb{C}$ by

$$\|\xi\|_h = |\theta(\xi)|e^{-\phi_h(x)}, \quad x \in \Omega, \quad \xi \in L_x,$$

where the function $\phi_h \in L^1_{\text{loc}}(\Omega)$ is called the *weight* of the metric h (w.r.t. the trivialization θ).

The curvature current Θ_h of L is given by the closed $(1, 1)$ -current $\Theta_h = \frac{i}{\pi} \partial \bar{\partial} \phi_h$. This current exists in the sense of distribution theory because of $\phi_h \in L^1_{\text{loc}}(\Omega)$ and is independent of the chosen trivialization.

The most important example for the rest of these notes is the possibly singular metric on L induced by non-zero holomorphic sections $\sigma_1, \dots, \sigma_N$ of L which is given in any trivialization θ by

$$\|\xi\|_h^2 = \frac{|\theta(\xi)|^2}{|\theta(\sigma_1(x))|^2 + \dots + |\theta(\sigma_N(x))|^2}.$$

Then the associated weight function is

$$\phi_h(x) = \log \left(\sum_{1 \leq j \leq N} |\theta(\sigma_j(x))|^2 \right)^{\frac{1}{2}}$$

which is a plurisubharmonic function, so Θ_h is a (closed) positive current. The order of logarithmic poles of a plurisubharmonic function in a point $x \in X$ is measured by the Lelong number

$$\nu(\phi, x) := \liminf_{z \rightarrow x} \frac{\phi(z)}{\log |z - x|}.$$

Lemma 1. *Let $\phi(z) = \log(\sum_{1 \leq j \leq N} |\theta(\sigma_j(x))|^2)^{\frac{1}{2}}$ be the weight of the hermitian metric on L induced by the holomorphic sections $\sigma_1, \dots, \sigma_N$ w.r.t. a trivialization θ on an open subset $\Omega \subset X$. Then for every point $x \in \Omega$:*

$$\nu(\phi, x) = \min_{1 \leq j \leq N} \{\text{ord}_x \sigma_j\}.$$

Proof. In dimension 1 this is clear by definition. In higher dimensions restriction to sufficiently general lines shows at least

$$\nu(\phi, x) \leq \min_{1 \leq j \leq N} \{\text{ord}_x \sigma_j\}.$$

To get the other inequality set $x := 0$ and add enough monomials z^α with $|\alpha| = \nu$ until we get

$$\phi(z) \leq \phi'(z) = \log \left(\sum_{|\alpha|=\nu} |\alpha! \cdot z^\alpha \cdot h_\alpha(z)|^2 \right)^{\frac{1}{2}}$$

for some holomorphic functions h_α defined around 0. These h_α are bounded in a neighborhood of 0 hence there exist constants $C, C' > 0$ such that

$$\phi'(z) \leq \log C + \log\left(\sum_{|\alpha|=\nu} |\alpha! \cdot z^\alpha|^2\right)^{\frac{1}{2}} \leq \log C' + \log(|z^\alpha|^{2\nu})^{\frac{1}{2}}.$$

But then

$$\liminf_{z \rightarrow x} \frac{\log(|z^\alpha|^{2\nu})^{\frac{1}{2}}}{\log |z|} = \nu.$$

□

Now we are able to calculate $\gamma(L, x)$ from multiplicities of divisors passing through x :

Theorem 1. *Let L be an ample line bundle on an n -dimensional projective complex manifold X . Then*

$$\epsilon(L, x) = \sup_{k; D_1, \dots, D_n \ni x} \left\{ \frac{\min_{i=1, \dots, n} \{\text{mult}_x D_i\}}{k} \right\}$$

where the supremum is taken over all divisors $D_1, \dots, D_n \in |kL|$ such that x is an isolated point of $D_1 \cap \dots \cap D_n$.

Proof. We know that for ample line bundles

$$\epsilon(L, x) = \sigma(L, x) = \sup_k \frac{1}{k} s(kL, x).$$

so for every $\delta > 0$ there is a $k \gg 0$ such that

$$\epsilon(L, x) - \delta \leq \frac{s}{k} := \frac{1}{k} s(kL, x) \leq \epsilon(L, x).$$

Consequently $H^0(X, kL)$ generates all s -jets and we can find holomorphic sections f_1, \dots, f_n whose s_k -jets are the monomials z_i^s , $i = 1, \dots, n$. The weight of the associated metric on L has an isolated pole of Lelong number $\frac{s}{k}$ in x and in the limit we get

$$\epsilon(L, x) \leq \sup_{k; D_1, \dots, D_n \ni x} \left\{ \frac{\min_{i=1, \dots, n} \{\text{mult}_x D_i\}}{k} \right\}.$$

On the other hand n holomorphic sections $\sigma_1, \dots, \sigma_n$ of kL (which are the 0-divisors of D_1, \dots, D_n define a metric h on kL . If we multiply ϕ_h by $\frac{1}{k}$ we get a metric on L with isolated pole in x hence the other inequality. □

We use Theorem 1 for computing Seshadri constants on products of two curves and some of their blow ups.

Proposition 2. *Let $X = C_1 \times C_2$ be the product of two smooth projective curves, $p_1 : X \rightarrow C_1$, $p_2 : X \rightarrow C_2$ the projections and \mathbf{a} a divisor of degree $a > 0$ on C_1 , \mathbf{b} a divisor of degree $b > 0$ on C_2 . Let $L = p_1^* \mathbf{a} + p_2^* \mathbf{b}$ be an ample divisor on X and x any point on X . Then*

$$\epsilon(L, x) = \min(a, b).$$

Proof. Let F_i be the numerical equivalence class of the fibers of p_i , $i = 1, 2$. Let E be the exceptional divisor of the blow up of X in x . Then

$$(aF_1 + bF_2 - mE)(F_1 - E) = b - m, \quad (aF_1 + bF_2 - mE)(F_2 - E) = a - m.$$

Since $F_1 - E$, $F_2 - E$ are the (numerical classes of the) strict transforms of the vertical resp. horizontal fiber through x a line bundle of numerical class $aF_1 + bF_2 - mE$ is ample only if $b - m > 0$, $a - m > 0$ by Seshadri's ampleness criterion. This implies

$$\epsilon(L, x) \leq \min(a, b).$$

On the other hand let $P_1 = p_1(x) \in C_1$ and $P_2 = p_2(x) \in C_2$. For $a' = ak$, $k \gg 0$, there exists an $A_2 \in |a'P_1|$ such that $A_2 - P_1$ is not effective. Similarly for $b' = bk$ there is a $B_2 \in |b'P_2|$ such that $B_2 - P_2$ is not effective. Setting $A_1 = a'P_1$, $B_1 = b'P_2$ we conclude that x is an isolated point of $(A_1 + B_2) \cap (A_2 + B_1)$ and

$$\frac{\min(\text{mult}_x(A_1 + B_2), \text{mult}_x(A_2 + B_1))}{k} = \frac{\min(ak, bk)}{k} = \min(a, b).$$

Now Seshadri constants are by definition numerical invariants. Since $A_1 + B_2$ and $A_2 + B_1$ belong to a line bundle with numerical class $k(aF_1 + bF_2)$ the theorem follows from Theorem 1. \square

Let $X = C_1 \times C_2$, \mathbf{a} , \mathbf{b} , a , b and L be as in the last proposition.

Proposition 3. *Let $\pi_n : X^{(n)} \rightarrow X$ be the blow up of X in n points x_1, \dots, x_n where no two of them lie on the same horizontal or vertical fiber. Then for*

$$L^{(n)} := \pi_n^* L - \sum_{i=1}^n m_i E_i, \quad 0 < m_i < \min(a, b), \quad \sum_{i=1}^n m_i \leq \max(a, b),$$

E_i the exceptional divisor over x_i we have

$$\epsilon(L^{(n)}, y) = \min(a, b)$$

for every point y not lying on the same horizontal or vertical fiber as one of the x_i .

Proof. Starting with Proposition 2 and using inductively the proposition together with Seshadri's ampleness criterion we conclude that $L^{(n)}$ is ample. As before we compute for the blow up $X^{(n+1)}$ of $X^{(n)}$ in y that (E_{n+1} the exceptional divisor over y)

$$(\pi_{n+1}^* L - \sum_{i=1}^n m_i E_i - m E_{n+1})(F_1 - E_{n+1}) = b - m$$

and analogously for $F_2 - E_{n+1}$. Consequently

$$\epsilon(L^{(n)}, y) \leq \min(a, b).$$

On the other hand we can suppose w.l.o.g. that $\min(a, b) = a$. Then we can choose integers $k_i, l_i \geq 0$, $i = 1, \dots, n$, such that

$$\sum_i k_i \leq a, \quad \sum_i l_i \leq b - a, \quad k_i + l_i \geq m_i.$$

Such partitions always exist if a, b and the m_i satisfy the assumptions of the proposition.

Let $P_i = p_1(x_i)$, $P = p_1(y)$, $Q_i = p_2(x_i)$, $Q = p_2(y)$. We distinguish two cases:

- (1) $a > \sum_i k_i$. Then there is q such that $|qaP - q \sum k_i P_i|$ is a base point free linear system on C_1 . Consequently we have a divisor $P'_1 + \dots + P'_{q(a - \sum k_i)}$ in this linear system such that all $P'_j \neq P$. Similarly, for $q \gg 0$ the linear

system $|qaQ - q \sum k_i Q_i|$ is base point free on C_2 and contains an element $Q'_1 + \dots + Q'_{q(a-\sum k_i)}$ with $Q'_j \neq Q$. Setting

$$\begin{aligned} A_1 &= qaP, \quad A_2 = q \sum k_i P_i + \sum P'_j, \\ B_1 &= q \sum (k_i + l_i) Q_i + \sum Q'_j + q(b-a - \sum l_i) Q', \\ B_2 &= qaQ + q \sum l_i Q_i + q(b-a - \sum l_i) Q' \end{aligned}$$

for some point $Q' \neq Q$ we can apply Theorem 1 on

$$\pi_n^* L' - \sum m_i E_i \equiv q(\pi_n^* L - \sum m_i E_i)$$

where $L' = \mathcal{O}(A_1 + B_1) = \mathcal{O}(A_2 + B_2)$ and conclude

$$\epsilon(L, y) \geq \frac{qa}{q} = a.$$

(2) $a = \sum_i k_i$. Then for all $q \in \mathbb{N}$ there exists a q' such that

$$|-(q'qa - q')P + q'q \sum k_i P_i| \quad \text{and} \quad |q'q \sum k_i Q_i - (q'qa - q')Q|$$

are base point free linear systems on C_1 and C_2 . Consequently we have divisors $P''_1 + \dots + P''_{q'}$ and $Q''_1 + \dots + Q''_{q'}$ in these linear systems such that $P''_j \neq P$ and $Q''_j \neq Q$. Setting

$$\begin{aligned} A_1 &= (q'qa - q')P + \sum P''_j, \quad A_2 = q'q \sum k_i P_i, \\ B_1 &= q'q \sum (k_i + l_i) Q_i + q'q(b - \sum (k_i + l_i)) Q'' \\ B_2 &= (q'qa - q')Q + q'q \sum l_i Q_i + \sum Q''_j + q'q(b - \sum (k_i + l_i)) Q'' \end{aligned}$$

for some point $Q'' \neq Q$ we can again apply Theorem 1 on

$$\pi_n^* L' - \sum m_i E_i \equiv q'q(\pi_n^* L - \sum m_i E_i)$$

where $L' = \mathcal{O}(A_1 + B_1) = \mathcal{O}(A_2 + B_2)$ and conclude

$$\epsilon(L, y) \geq \frac{q'qa - q'}{q'q} = a - \frac{1}{q}.$$

But since q can be chosen arbitrarily big this implies

$$\epsilon(L, y) \geq a.$$

□

Of course we can similarly calculate Seshadri constants of (some) ample line bundles on the the blown up variety when some of the x_i lie on the same horizontal or vertical fiber or on exceptional divisors. Let us instead use Prop. 3 to give a counter example to a conjecture concerning maximal rationally connected quotients:

Let X be an n -dimensional projective complex manifold and $T_{\mathcal{F}} \subset T_X$ the tangent sheaf of a foliation on X . By a result of Bogomolov and McQuillan ([BM01], see also [KTS05]) the existence of a curve C such that $T_{\mathcal{F}|_C}$ is an ample vector bundle implies that all the leaves of \mathcal{F} through points of C are rationally connected algebraic subvarieties.

There is a natural way of getting foliations \mathcal{F} and curves C such that $T_{\mathcal{F}|_C}$ is ample: Fix an ample line bundle A and consider sufficiently general complete intersection curves $C \in |k_1 A \cap \dots \cap k_{n-1} A|$, $k_i \gg 0$. By Mehta-Ramanathan's theorem the

restriction of the Harder-Narasimhan filtration of T_X on C is again the Harder-Narasimhan filtration of $T_{X|C}$. Furthermore the maximal ample subbundle contained in $T_{X|C}$ is a bundle occurring in this filtration. Hence it is the restriction of a subbundle $\mathcal{H} \subset T_X$ occurring in the Harder-Narasimhan filtration of T_X and simple calculations using stability properties show that \mathcal{H} is saturated and involutive. Consequently \mathcal{H} is the tangent sheaf of a foliation such that the restriction to sufficiently general complete intersection curves of type $k_1 A \cap \dots \cap k_{n-1} A$ is ample. On the other hand Campana [Cam81, Cam94] and Kollár, Miyaoka and Mori [KMM92] constructed the maximal rationally connected (MRC) quotient of a projective manifold X . It is tempting to

Conjecture 4. *Let A be an ample line bundle on X and C a sufficiently complete intersection curve of type A . Then the MRC quotient induces a foliation \mathcal{F} such that $T_{\mathcal{F}|C}$ is the maximal ample part of the Harder-Narasimhan filtration of $T_{\mathcal{F}|C}$.*

Unfortunately there exists a counter example to this conjecture already on surfaces: Let $X = C \times \mathbb{P}^1$ be the ruled product surface over an elliptic curve C with projections $p_1 : X \rightarrow C$, $p_2 : X \rightarrow \mathbb{P}^1$. Let x_1, x_2, x_3 be 3 points on X such that no two of them lie on the same horizontal or vertical fiber. Let $\pi : \widehat{X} \rightarrow X$ be the blow up of X in x_1, x_2, x_3 and let

$$L = p_1^* \mathbf{a} + p_2^* \mathbf{b}, \quad \deg_C \mathbf{a} = 3, \quad \deg_{\mathbb{P}^1} \mathbf{b} = 4.$$

L is an ample line bundle on X and it follows from Prop. 3 that

$$L' = \pi^* L - 2E_1 - 2E_2 - 2E_3$$

is ample, too. (Here the E_i are the exceptional divisors on \widehat{X} over x_i .) On the other hand the MRC quotient of \widehat{X} is just the blown-up ruling of X (that is a simple consequence of surface classification). The induced foliation \mathcal{F} has tangent sheaf

$$T_{\mathcal{F}} = p_2^* \mathcal{O}(2) - E_1 - E_2 - E_3$$

as a local computation around the blown up points show (see [Bru04]). We have

$$L' \cdot T_{\mathcal{F}} = 0$$

and hence no complete intersection curve $C \in |kL'|$ intersects $T_{\mathcal{F}}$ positively.

Finally we turn to Seshadri constants associated to sets of points.

Definition 5. *Let L be an ample line bundle on an irreducible projective variety X and let x_1, \dots, x_r be arbitrary points on X . Then we define*

$$\epsilon(L; x_1, \dots, x_r) = \max\{\epsilon \geq 0 : \pi^* L - \epsilon \cdot \sum_{i=1}^r E_i \text{ is nef}\}$$

where $\pi : \widetilde{X} \rightarrow X$ is the blow up of X in x_1, \dots, x_r and $E_i \subset \widetilde{X}$ is the exceptional divisor over x_i .

Proposition 6. *Let L be an ample line bundle on a surface X and $x_1, \dots, x_r \in X$ be arbitrary points. Then $\epsilon(L; x_1, \dots, x_r)$ equals*

$$\inf_C \frac{L \cdot C}{\sum_{i=1}^r \text{mult}_{x_i} C},$$

where the infimum is taken over all irreducible curves $C \subset X$,

$$\sup_k \frac{s(kL; x_1, \dots, x_r)}{k}$$

where $s(kL; x_1, \dots, x_r)$ is defined as the maximal s such that $H^0(X, kL) \rightarrow \bigoplus_{i=1}^r \mathcal{O}_{X, x_i} / \mathfrak{m}_{x_i}^{s+1}$ is surjective and

$$\sup_{k; D_1, D_2} \frac{\min_j \min_{i=1,2}(\text{mult}_{x_j} D_i)}{k}$$

where the supremum is taken over all pairs of divisors $D_1, D_2 \in |kL|$ such that x_1, \dots, x_r are isolated points in $D_1 \cap D_2$.

Proof. We slightly generalize the arguments in [Dem92, Thm.6.4]: The first equality is a direct consequence of Seshadri's ampleness criterion. Next suppose that there are two divisors $D_1, D_2 \in |kL|$ such that x_1, \dots, x_r are isolated points in $D_1 \cap D_2$. If f_{1j}, f_{2j} are local equations of D_1, D_2 w.r.t. a trivialization θ_j in a neighborhood Ω_j of x_j ,

$$|\xi|_h := |\theta_j(\xi)| e^{-\phi_j}$$

with weight function $\phi_j = \frac{1}{k} \log(|f_{1j}|^2 + |f_{2j}|^2)$ defines a positive singular hermitian metric on L with isolated poles at the x_j and Lelong numbers

$$\gamma_j := \nu(\phi_j, x_j) = \min\{\text{mult}_{x_j} D_1, \text{mult}_{x_j} D_2\}.$$

Set $c(L) = \frac{i}{\pi} \partial \bar{\partial} \phi$ on Ω_j and let C be an irreducible curve. Then

$$\begin{aligned} L.C &= \int_C c(L) \geq \sum_{j=1}^r \int_{C \cap \Omega_j} \frac{i}{\pi} \partial \bar{\partial} \phi \geq \sum_{j=1}^r \gamma_j \nu(C, x_j) \\ &= \sum_{j=1}^r \gamma_j \text{mult}_{x_j} C \geq \frac{\min_j \min_{i=1,2}(\text{mult}_{x_j} D_i)}{k} \sum_{j=1}^r \text{mult}_{x_j} C \end{aligned}$$

because the last integral is larger than the Lelong number of the integration current $[C]$ w.r.t. the weight ϕ_j and we may apply the comparison theorem with the ordinary Lelong number associated to the weight $\log|z-x|$ (see [Dem00, 2.B] for more details). Therefore

$$\epsilon(L; x_1, \dots, x_r) \geq \sup_{k; D_1, D_2} \frac{\min_j \min_{i=1,2}(\text{mult}_{x_j} D_i)}{k}.$$

Now fix a coordinate system (z_1, z_2) on Ω_j centered at x_j . For $s := s(kL; x_1, \dots, x_r)$ the sections in $H^0(X, kL)$ generate all combinations of s -jets at all points x_j and we can find two holomorphic sections f_1, f_2 whose s -jets are z_1^s, z_2^s at every x_j . We define a global positive singular metric by

$$|\xi|_h := |\theta_j(\xi)| e^{-\phi_j}$$

with weight function $\phi_j = \frac{1}{k} \log(|\theta_j(f_1)|^2 + |\theta_j(f_2)|^2)$. Then ϕ_j has isolated poles with Lelong number $\frac{s}{k}$ at x_j thus

$$\sup_{k; D_1, D_2} \frac{\min_j \min_{i=1,2}(\text{mult}_{x_j} D_i)}{k} \geq \frac{s}{k}.$$

Finally by using Kodaira's vanishing theorem for $\mathcal{O}(p\pi^*L - q\sum_{j=1}^r E_j)$ with

$$\frac{q}{p} < \epsilon(L; x_1, \dots, x_r)$$

we can show that

$$\sup_k \frac{s(kL; x_1, \dots, x_r)}{k} \geq \epsilon(L; x_1, \dots, x_r).$$

□

A celebrated conjecture of Nagata may be seen as a statement on Seshadri constants associated to sets of points in \mathbb{P}^2 : Let

$$\pi : \tilde{X} = \text{Bl}_r(\mathbb{P}^2) \rightarrow \mathbb{P}^2$$

be the blow up of the projective plane in a finite set $S \subset \mathbb{P}^2$ consisting of r very general points. Denote by H the pull back of a line to \tilde{X} , and let E_1, \dots, E_r be the exceptional divisors over the x_j . Nagata [Nag59] conjectured in effect that the \mathbb{R} -divisor

$$H - \sqrt{\frac{1}{r}} \sum_{j=1}^r E_j$$

is nef on \tilde{X} provided that $r \geq 9$.

It is well known that Nagata's conjecture is implied by another conjecture of Harbourne and Hirschowitz about spaces $\mathcal{L}_d(n^m)$ of plane curves of given degree d and multiplicity at least m at n general points [Mir99, CM01]. This conjecture tries to detect those of the spaces $\mathcal{L}_d(n^m)$ which do not have the expected dimension

$$\frac{d(d+3)}{2} - n \cdot \frac{m(m+1)}{2}.$$

In particular it implies that the $\mathcal{L}_d(n^m)$ do have expected dimension if $d \geq 3m$ [CM00].

We show now that it is not necessary to prove the Harbourne–Hirschowitz conjecture for every triple (d, n, m) to get Nagata's conjecture:

Theorem 7. *Let $r > 9$ be an integer and (d_i, m_i) a sequence of pairs of positive integers such that $\frac{d_i^2}{m_i^2 \cdot r} \xrightarrow{i \rightarrow \infty} 1$ and the space $\mathcal{L}_{d_i}(r^{m_i+1})$ has expected dimension ≥ 0 . Then Nagata's conjecture is true for r general points in \mathbb{P}^2 .*

The advantage of this approach to Nagata's conjecture is that recent attacks on the Harbourne–Hirschowitz conjecture (see especially [CM00]) work with recursions which unfortunately have some gaps. But it might be worth checking if around these gaps there are sequences of triples (r, d_i, m_i) as in the theorem.

Note furthermore that the assumptions of the theorem on the spaces $\mathcal{L}_{d_i}(r^{m_i+1})$ follow from the Harbourne–Hirschowitz conjecture since $\frac{d_i}{m_i} \rightarrow \sqrt{r}$ implies $d_i \geq 3m_i$ for i big enough if $r > 9$.

Proof of the Theorem. Note first that if we define $m(d, r)$ as the maximal m such that

$$\frac{d(d+3)}{2} - r \cdot \frac{m(m+1)}{2} \geq 0$$

we get for fixed r that

$$\lim_{d \rightarrow \infty} \frac{m(d, r)}{d} = \frac{1}{\sqrt{r}}.$$

Next we observe the following fact: Since $\mathcal{L}_{d_i}(r^1)$ has also the expected dimension (this is the Multiplicity One Theorem in [CM98]) the intermediate spaces $\mathcal{L}_{d_i}(r^{m_i})$ and

$$\mathcal{L}_{d_i}(k^{m_i}, (r-k)^{m_i+1}), \quad k = 0, \dots, r$$

(denoting the space of curves having multiplicity m_i in k points and multiplicity $m_i + 1$ in $r - k$ points) have expected dimension.

Now let $\pi : \tilde{X} = \text{Bl}_r(\mathbb{P}^2) \rightarrow \mathbb{P}^2$ be the blow up of \mathbb{P}^2 in r general points x_1, \dots, x_r . We choose m_i points $y_{1j}, \dots, y_{m_i j}$ on every exceptional divisor E_j over x_j and consider the space of curves $C \subset \tilde{X}$ passing through the points y_{kj_0} , $k = 1, \dots, m_i$ for some fixed j_0 such that

- (i) $\deg \pi(C) = d_i$ and
- (ii) $\pi(C)$ has multiplicity $m_i + 1$ in the points $x_1, \dots, x_{j_0-1}, x_{j_0+1}, \dots, x_r$.

The expected dimension of this space is

$$\begin{aligned} \dim \mathcal{L}_{d_i}(1^{m_i}, (r-1)^{m_i+1}) - m_i &> \dim \mathcal{L}_{d_i}(1^{m_i}, (r-1)^{m_i+1}) - m_i - 1 = \\ &= \mathcal{L}_{d_i}(r^{m_i+1}). \end{aligned}$$

Hence there exists a curve $C \subset \mathbb{P}^2$ in this space having multiplicity exactly m_i in x_{j_0} and whose tangent cone is given by the points $y_{1j_0}, \dots, y_{m_i j_0}$.

The r -tuples of pairwise disjoint points in \mathbb{P}^2 with one marked point is parametrized by the subvariety

$$\begin{array}{ccc} W \hookrightarrow & ((\mathbb{P}^2)^r - D) \times \mathbb{P}^2 & \xrightarrow{pr} \mathbb{P}^2 \\ & \downarrow pr & \\ & (\mathbb{P}^2)^r - D & \end{array}$$

where D collects all r -tuples in $(\mathbb{P}^2)^r$ containing one point (at least) twice. A fiber of W over an r -tuple in $(\mathbb{P}^2)^r - D$ is just the scheme of the r points in the tuple. Since the fibers of the other projection are Zariski-open and hence irreducible subsets of $(\mathbb{P}^2)^{r-1}$ the algebraic subset W is irreducible.

Consequently no Zariski-closed subset of W can dominate $(\mathbb{P}^2)^r - D$ and we can find a tuple (x_1, \dots, x_r) such that for every $j_0 = 1, \dots, r$ there are curves C as described above. Considering these curves as divisors of degree d_i and adding them gives a divisor D which has exactly multiplicity m_i in every point x_i and whose tangent cone in x_i is described by the y_{ij} , $j = 1, \dots, m_i$.

Choosing points y'_{ij} such that

$$\{y_{i1}, \dots, y_{im_i}\} \cap \{y'_{i1}, \dots, y'_{im_i}\} = \emptyset, \quad i = 1, \dots, r$$

we get in the same way another divisor D' such that $D \cap D'$ contains x_1, \dots, x_r as isolated points and $\text{mult}_{x_i} D = \text{mult}_{x_i} D' = m_i$. The two divisors D and D' may be used to derive from Prop. 6

$$\frac{m_i}{d_i} \leq \epsilon(H; x_1, \dots, x_r).$$

On the other hand computing the self intersection of $\pi^* H - \frac{1}{\sqrt{r}} \sum E_i$ shows that

$$\epsilon(H; x_1, \dots, x_r) \leq \frac{1}{\sqrt{r}}$$

hence the theorem. \square

REFERENCES

- [Bau99] Th. Bauer. Seshadri Constants on Algebraic Surfaces. *Math. Ann.*, 313:547–583, 1999.
- [BM01] F. Bogomolov and M. McQuillan. Rational curves on foliated varieties. IHES Preprint, 2001.
- [Bru04] M. Brunella. *Birational Geometry of Foliations*. IMPA Mathematical Publications, 2004.
- [Cam81] F. Campana. Coréduction algébrique d’un espace analytique faiblement kählerien compact. *Inv. Math.*, 63:187–223, 1981.
- [Cam94] F. Campana. Remarques sur les revêtements universels des variétés kähleriennes compacte. *Bull. SMF*, 122:255–284, 1994.
- [CM98] C. Ciliberto and R. Miranda. Degenerations of planar linear systems. *J. Reine Angew. Math.*, 501:191–220, 1998.
- [CM00] C. Ciliberto and R. Miranda. Linear systems of plane curves with base points of equal multiplicity. *Trans. AMS*, 352:4037–4050, 2000.
- [CM01] C. Ciliberto and R. Miranda. The Segre and Harbourne–Hirschowitz conjectures. In *Applications of algebraic geometry to coding theory, physics and computation (Eilat, 2001)*, volume 36 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 37–51. Kluwer Acad. Publ., Dordrecht, 2001.
- [Dem92] J.-P. Demailly. Singular hermitian metrics on positive line bundles. In *Complex Algebraic Varieties, Bayreuth 1990*, volume 1507 of *SLNM*, pages 87–104, 1992.
- [Dem00] J.-P. Demailly. Multiplier ideal sheaves and analytic methods in algebraic geometry. Lecture Notes, School on Vanishing theorems and effective results in Algebraic Geometry, ICTP Trieste, April 2000.
- [EKL95] L. Ein, O. Küchle, and R. Lazarsfeld. Local positivity of ample line bundles. *J. Diff. Geom.*, 42:193–219, 1995.
- [Gar05] L. F. Garcia. Seshadri constants on ruled surfaces: the rational and the elliptic cases. [math.AG/0503253](https://arxiv.org/abs/math/0503253), 2005.
- [KMM92] J. Kollár, Y. Miyaoka, and S. Mori. Rationally connected varieties. *J. Alg. Geom.*, 1:429–448, 1992.
- [KTS05] S. Kebekus, M. Toma, and L. Sola. Rationally connected foliations after Bogomolov and McQuillan. Preprint, [math.AG/0505222](https://arxiv.org/abs/math/0505222), 2005.
- [Laz97] R. Lazarsfeld. Lectures on Linear Series. In *Complex Algebraic Geometry (Park City, Utah, 1993)*, volume 3 of *IAS/Park City Math. Ser.*, pages 161–219, Providence, 1997. AMS.
- [Laz04] R. Lazarsfeld. *Positivity in Algebraic Geometry I*. Erg. Math. 48. Springer, New York, 2004.
- [Mir99] R. Miranda. Linear Systems of plane curves. *Notices AMS*, 46:192–201, 1999.
- [Nag59] M. Nagata. On the 14-th problem of Hilbert. *Amer.J.Math*, 81:766–772, 1959.

THOMAS ECKL, MATHEMATISCHES INSTITUT, UNIVERSITÄT ZU KÖLN, 50931 KÖLN, GERMANY
E-mail address: thomas.eckl@math.uni-koeln.de
URL: <http://www.mi.uni-koeln.de/~teckl>