INDUCTIVE LIMITS OF K-THEORETIC COMPLEXES WITH TORSION COEFFICIENTS

SØREN EILERS AND ANDREW S. TOMS

ABSTRACT. We present the first range result for the total K-theory of C^* -algebras. This invariant has been used successfully to classify certain separable, nuclear C^* -algebras of real rank zero. Our results complete the classification of the so-called AD algebras of real rank zero.

1. INTRODUCTION

A theorem which classifies the objects of a category up to some notion of equivalence via an invariant begs naturally the question of range for the classifying invariant. In the classification theory of C^* -algebras by K-theoretic invariants the fundamental range-of-invariant result is the theorem of Effros, Handelman, and Shen ([9]), which states that the ordered groups arising as K₀-groups of AF algebras are exactly the dimension groups studied first by Riesz ([22]) and Fuchs. ([17]). Post AF classification results for C^* -algebras have required invariants more complex than K₀ alone, yet it has typically been possible to pair such results with an Effros-Handelman-Shen type theorem establishing the range of the classifying invariant. Notable examples include [24] and [15].

The aim of the present paper is to give the first range-of-invariant result associated to the classification of certain C^* -algebras of real rank zero, completed by Dadarlat and Gong in [6]. To obtain complete invariants for this class of non-simple C^* -algebras, the ordered K_{*}group $K_0(-) \oplus K_1(-)$ must be augmented: the addition of ordered K-groups with torsion coefficients and certain natural homomorphisms between them is required. While the completeness of this invariant has been established for almost a decade in a number of cases ([11], [8], [6]), there have been no range results available until now.

The situation is complicated by the intricate nature of the augmented K-theory. Following [6], this invariant associates to each C^* -algebra \mathfrak{A}

²⁰⁰⁰ Mathematics Subject Classification. Primary 46L35, Secondary 46L80.

Key words and phrases. C*-algebras, K-theory with coefficients, classification.

a family of groups

 $\mathrm{K}_{0}(\mathfrak{A}), \mathrm{K}_{1}(\mathfrak{A}), \mathrm{K}_{0}(\mathfrak{A}; \mathbb{Z}/n), \mathrm{K}_{1}(\mathfrak{A}; \mathbb{Z}/n)$

with n ranging over $\{2, 3, ...\}$, as well as an order structure on

$$\mathrm{K}_{0}(\mathfrak{A}) \oplus \mathrm{K}_{1}(\mathfrak{A}) \oplus \bigoplus_{n \in \{2,3,4,\dots\}} [\mathrm{K}_{0}(\mathfrak{A}; \mathbb{Z}/n) \oplus \mathrm{K}_{1}(\mathfrak{A}; \mathbb{Z}/n)]$$

and families of group homomorphisms

$$\rho_n^i : \mathrm{K}_i(\mathfrak{A}) \longrightarrow \mathrm{K}_i(\mathfrak{A}; \mathbb{Z}/n)$$
$$\beta_n^i : \mathrm{K}_i(\mathfrak{A}; \mathbb{Z}/n) \longrightarrow \mathrm{K}_{i+1}(\mathfrak{A})$$
$$\kappa_{n,m}^i : \mathrm{K}_i(\mathfrak{A}; \mathbb{Z}/n) \longrightarrow \mathrm{K}_i(\mathfrak{A}; \mathbb{Z}/m).$$

Thus, an isomorphism of invariants amounts to a family

$$\phi_i: \mathrm{K}_i(\mathfrak{A}) \longrightarrow \mathrm{K}_i(\mathfrak{B}) \qquad \psi_i: \mathrm{K}_i(\mathfrak{A}; \mathbb{Z}/n) \longrightarrow \mathrm{K}_i(\mathfrak{B}; \mathbb{Z}/n)$$

which preserves the order structure and intertwines all morphisms ρ, β, κ .

To keep technicalities to a minimum while staying in a class where, by the counterexamples given in [7] and [5], the full force of such an invariant is really needed, we shall concentrate on the class of so-called AD algebras of real rank zero. Recall that an AD algebra is an inductive limit of finite direct sums of matrix algebras over elements of

$$\mathbf{D} := \{ \mathbb{C}, C(S^1), \mathbb{I}_2^{\sim}, \mathbb{I}_3^{\sim}, \mathbb{I}_4^{\sim}, \dots \},\$$

where \mathbb{I}_n^{\sim} is the dimension drop algebra

 $\{f \in C([0,1], \mathbf{M}_n(\mathbb{C})) \mid f(0), f(1) \in \mathbb{C}\mathbf{1}\}.$

Such C^* -algebras may be classified by a more manageable invariant of the form

$$\mathrm{K}_{0}(\mathfrak{A})\otimes\mathbb{Q}\longrightarrow\mathrm{K}_{0}(\mathfrak{A};\mathbb{Q}/\mathbb{Z})\longrightarrow\mathrm{K}_{1}(\mathfrak{A})$$

provided that they are of real rank zero, cf. [4]. Here, as we shall recall below, $K_0(\mathfrak{A}; \mathbb{Q}/\mathbb{Z})$ should be thought of as a kind of conglomerate of $K_0(\mathfrak{A}; \mathbb{Z}/n)$ for all *n*. If the torsion part of $K_1(\mathfrak{A})$ is annihilated by a fixed integer *n*, then the even more manageable invariant

$$\overline{K}_n(\mathfrak{A}) \qquad \mathrm{K}_0(\mathfrak{A}) \longrightarrow \mathrm{K}_0(\mathfrak{A}; \mathbb{Z}/n) \longrightarrow \mathrm{K}_1(\mathfrak{A})$$

suffices.

Our strategy is to establish a range result in the latter case first, then use this to derive the general result. The key technical element of the proof is a decomposition result for refinement monoids attributed to Tarski by Wehrung ([25]).

To illustrate our results we revisit examples of AD algebras originally considered by Dadarlat and Loring, which showed that such algebras could have isomorphic ordered K_{*}-groups without having isomorphic augmented K-theory. Using our main result, we parametrize the real rank zero AD algebras with K_* -groups as considered in [7], and show that there are uncountably many non-isomorphic such algebras.

2. Building blocks

In this section we introduce the notion of an *n*-coefficient complex. This type of object is meant to abstract the characteristics of certain augmented K-theoretic invariants for AD algebras of real rank zero — invariants which will be reviewed in detail in section 3. We begin with some preliminaries and notation.

A graded ordered group is a graded group $G_0 \oplus G_1$ in which the G_0 -component dominates the order in the sense that

$$\begin{array}{c} (x,y) \geq 0 \\ (x,y') \geq 0 \end{array} \right\} \Longrightarrow (x,y \pm y') \geq 0$$

For any group G, we denote by G[n] the subgroup of elements of G annihilated by $n \in \mathbb{N}$. When G is an ordered group, we denote by I(x) the order ideal containing x. Recall the notions of unperforated and weakly unperforated groups from [20].

Let G be an ordered abelian group, H an abelian group, and $f : G \to H$ a surjective group homomorphism. Say that $h \in H$ is positive if it is the image of a positive element in G. The important and obvious feature of the order on H thus defined (the so-called *quotient order*) is that every positive element in H lifts to a positive element in G ([20]).

Definition 2.1. Let $n \in \{2, 3, ...\}$. An *n*-coefficient complex \overline{G} is an exact sequence

$$\mathbf{G}_0 \xrightarrow{\rho} \mathbf{G}_n \xrightarrow{\beta} \mathbf{G}_1$$

of abelian groups which, setting

$$\mathbf{G}_* := \mathbf{G}_0 \oplus \mathbf{G}_1, \quad \mathbf{G}_{\overline{n}} := \mathbf{G}_0 \oplus \mathbf{G}_n,$$

has the following properties:

- (i) $n\mathbf{G}_n = 0$
- (*ii*) ker $\rho = n \mathbf{G}_0$, im $\beta = \mathbf{G}_1[n]$.
- (iii) G_* and $G_{\underline{P}}$ are graded ordered groups restricting to the same order on G_0
- (iv) G_* has the Riesz interpolation property.
- (v) $G_0 \oplus \rho(G_0)$ has the quotient order coming from $id_{G_0} \oplus \rho$
- (vi) $G_0 \oplus \beta(G_n)$ has the quotient order coming from $id_{G_0} \oplus \beta$
- (vii) G_0 is unperforated and G_* is weakly unperforated.

We say that an element (x, y, z) is positive in \overline{G} if and only if $gd \ni (x, y) \ge 0$ and $G_* \ni (x, z) \ge 0$.

A morphism $\overline{\theta} : \overline{G} \to \overline{H}$ of *n*-coefficient complexes is a positive ordered triple of linear maps $(\theta_0, \theta_n, \theta_1)$ such that

 $\theta_0: \mathbf{G}_0 \to \mathbf{H}_0, \ \theta_n: \mathbf{G}_n \to \mathbf{H}_n, \ \theta_1: \mathbf{G}_1 \to \mathbf{H}_1,$

and the maps commute with ρ or β as appropriate.

We conclude this section by introducing three types of n-coefficient complexes — our so-called building blocks.

Definition 2.2. The

(i) (\mathbb{C}, n) complex

$$\mathbb{Z} \xrightarrow{\rho} \mathbb{Z}/n \xrightarrow{\beta} 0$$
$$\rho: 1 \mapsto \overline{1}, \beta: \overline{1} \mapsto 0,$$

(ii) (\mathbb{I}_m^{\sim}, n) complex

$$\mathbb{Z} \xrightarrow{\rho} \mathbb{Z}/n \oplus \mathbb{Z}/m \xrightarrow{\beta} \mathbb{Z}/m$$
$$\rho: 1 \mapsto (\overline{1}, \overline{0}); \beta: (\overline{1}, \overline{0}) \mapsto 0, (\overline{0}, \overline{1}) \mapsto \overline{1}$$

and

(iii) $(C(S^1), n)$ complex

$$\mathbb{Z} \xrightarrow{\rho} \mathbb{Z}/n \xrightarrow{\beta} \mathbb{Z}$$
$$\rho: 1 \mapsto \overline{1}, \beta: \overline{1} \mapsto 0,$$

where G_{\square} and G_* have the strict order coming from the first direct summand, are n-coefficient complexes.

The motivation for these defining these objects, hinted at by their very names, will be made clear in the following section.

3. K-THEORY WITH COEFFICIENTS

In this section we collect a suite of known results which together prove that *n*-coefficient complexes appear as the K-theory of certain C^* -algebras.

The following definitions, originating in the work of Dadarlat, Gong, and the first named author (see [4]) are based on the observation (from [26, 2.3]) that the lattices of order ideals of $K_0(\mathfrak{A})$ and of ideals of \mathfrak{A} are naturally isomorphic for C^* -algebras with minimal ranks. **Definition 3.1.** Let \mathfrak{A} be a C^* -algebra of real rank zero and stable rank one. When I is an order ideal of $K_0(\mathfrak{A})$ we define

$$\lfloor I \rfloor = \iota_*(\mathcal{K}_1(\mathfrak{I})) \text{ and } \lceil I \rceil = \iota_*(\mathcal{K}_0(\mathfrak{I}; \mathbb{Z}/n)),$$

where \mathfrak{I} is the unique ideal of \mathfrak{A} with $I = \iota_*(K_0(\mathfrak{I}))$ and $\iota : \mathfrak{I} \hookrightarrow \mathfrak{A}$ is the inclusion map.

We now equip $K_*(\mathfrak{A}) = K_0(\mathfrak{A}) \oplus K_1(\mathfrak{A})$ and $K_{\underline{n}}(\mathfrak{A}) = K_0(\mathfrak{A}) \oplus K_0(\mathfrak{A}; \mathbb{Z}/n)$ with the orders given by

$$(x,y) \ge 0 \Longrightarrow \left\{ \begin{array}{c} x \ge 0\\ y \in \lfloor \mathbf{I}(x) \rfloor \end{array} \right\}$$

and

$$(x,z) \ge 0 \Longrightarrow \left\{ \begin{array}{c} x \ge 0 \\ z \in \lceil \mathbf{I}(x) \rceil \end{array} \right\},$$

respectively.

It is well known (cf. [14]) that the order thus defined on $K_*(\mathfrak{A})$ will coincide with the standard order on $K_*(\mathfrak{A})$ derived from the isomorphism

$$\mathrm{K}_*(\mathfrak{A}) \cong \mathrm{K}_0(\mathfrak{A} \otimes C(S^1))$$

In general, the order on $K_{\underline{m}}(\mathfrak{A})$ will <u>not</u> be the one similarly derived from the isomorphism

$$\mathrm{K}_{\overline{n}}(\mathfrak{A}) \cong \mathrm{K}_0(\mathfrak{A} \otimes \mathbb{I}_n^{\sim}).$$

But since, as seen in [4], these two order structures allow the same positive group isomorphisms for a large class of C^* -algebras including the AD algebras, the choice of order structure for the invariant has no influence on the associated classification results.

Proposition 3.2. Let \mathfrak{A} be a C^* -algebra of real rank zero and stable rank one. Assume that $K_*(\mathfrak{A})$ is weakly unperforated, and that $K_0(\mathfrak{A})$ is unperforated. For any $n \in \{2, 3, \ldots\}$,

$$\overline{K}_n(\mathfrak{A}): \qquad \mathrm{K}_0(\mathfrak{A}) \longrightarrow \mathrm{K}_0(\mathfrak{A}; \mathbb{Z}/n) \longrightarrow \mathrm{K}_1(\mathfrak{A})$$

is an n-coefficient complex.

Proof: We verify properties (i) - (vii) from Definition 2.1.

The purely algebraic properties (i) and (ii) hold true for any such sequence, cf. [23]. Furthermore, it is clear from our definition of the order on $K_*(\mathfrak{A})$ and $K_{\square}(\mathfrak{A})$ that they are graded order groups based on the same order on $K_0(\mathfrak{A})$. Since we have noted that we are in fact working with the standard order on $K_*(\mathfrak{A})$, [3] or [14] show that condition (iv) is met. Inspection of the diagram

(1)
$$\begin{array}{ccc} \mathrm{K}_{0}(\mathfrak{I}) & \xrightarrow{\rho} \mathrm{K}_{0}(\mathfrak{I}; \mathbb{Z}/n) & \xrightarrow{\beta} \mathrm{K}_{1}(\mathfrak{I}) \\ & & \downarrow & & \downarrow & & \downarrow \\ & & \downarrow & & \downarrow & & \downarrow \\ & & \mathrm{K}_{0}(\mathfrak{A}) & \xrightarrow{\rho} \mathrm{K}_{0}(\mathfrak{A}; \mathbb{Z}/n) & \xrightarrow{\beta} \mathrm{K}_{1}(\mathfrak{A}) \end{array}$$

when \mathfrak{I} is the ideal of \mathfrak{A} corresponding to the order ideal I(x) shows that if $G_0 \ni x \ge 0$, then $G_n \ni (x, \rho(x)) \ge 0$, and, similarly, that if $G_n \ni (x, y) \ge 0$, then $G_* \ni (x, \beta(y)) \ge 0$.

To prove property (v) we look again at the diagram (1). By assumption, $y \in K_0(\mathfrak{A}; \mathbb{Z}/n)$ is in the image of both maps with target $K_0(\mathfrak{A}; \mathbb{Z}/n)$, so an easy diagram chase gives the desired result whenever $\iota_* : K_1(\mathfrak{I}) \longrightarrow K_1(\mathfrak{A})$ is injective. But ι_* is always injective by [21]. Combining this fact with the observation of the preceding paragraph, we have property (v).

For (vi), we do a similar diagram chase. Finally, we have explicitly required the properties in (vii).

4. Decomposition Lemmas

In this section we establish some decomposition results in the spirit of Riesz for n-coefficient complexes. These lemmas will allow us to prove an Effros-Handelman-Shen-type result for these complexes, realising them as inductive limits of our building blocks.

We shall rely heavily on results in [15] pertaining to the family of ordered $K_*(-)$ -groups of AH algebras with real rank zero. These groups have the following property:

Definition 4.1. (Cf. Goodearl ([19, Lemma 8.1]) and Elliott ([15])) An ordered group G is said to be weakly unperforated if

- (i) whenever $mx \in G_+$ there exists $t \in tor(G)$ with $x + t \in G_+$ and mt = 0;
- (ii) whenever $y \in G_+$, $t \in tor(G)$, and $ny + t \in G_+$ for some $n \in \mathbb{N}$, then $y \pm t \in G_+$.

Note that property (ii) is automatic in our case since all torsion is localized in the odd part of a graded ordered group. Although we do not apply the next observation in the sequel, we nevertheless record it for possible future use: all of the results in this section hold true if the condition of unperforation in G₀ in Definition 2.1(*vii*) is relaxed to weak unperforation. **Lemma 4.2** (Elliott ([15, Corollary 6.6])). Let $G_* = G_0 \oplus G_1$ be a weakly unperforated graded ordered group with the Riesz decomposition property. If $s_1, \ldots, s_m \leq g$ where $g \in G_0^+$ and $s_1, \ldots, s_m \in G_1$, then $g = g_1 + \cdots + g_m$ with $g_1, \ldots, g_m \in G_0^+$ and $s_i \leq g_i$.

We say that a family H_1, \ldots, H_n of subgroups of a given groups G is *independent* if

$$\sum_{i=1}^{n} x_i = 0, x_i \in H_i \Longrightarrow x_1 = \dots = x_n = 0$$

Lemma 4.3 (Elliott ([15, Corollary 6.3])). Let $G_* = G_0 \oplus G_1$ be a weakly unperforated graded ordered group with the Riesz decomposition property. Suppose $x \leq \sum_{j=1}^{k} g_j$, where $x \in G_1$ and $g_j \in G_0^+$. Then, there exist an independent family H_j , $j \in \{1, \ldots, k\}$, of finitely generated subgroups of G_1 such that $H_j \leq g_j$ and a decomposition $x = \sum_{j=1}^{k} x_j$ such that $x_j \in H_j$.

Note that if x is as in Lemma 4.3 and has order m, then each x_j has order at most m by the independence of the x_j . Thus, by property (*ii*) of Definition 2.1, if the G_* of Lemma 4.3 is in fact G_* for some n-coefficient complex \overline{G} and x is in the image of β , then so too are the x_j .

Wehrung attributes the following observation to Tarski:

Lemma 4.4 (Cf. Wehrung ([25, Lemma 1.9])). Let G_0 be an ordered group with the Riesz interpolation property, and let $a, b \in G_0$ satisfy $a, b \ge 0$ and $a \le nb$, $n \in \mathbb{N}$. Then, there exist $b_0, \ldots, b_n \ge 0$ such that $b = \sum_{i=0}^n b_i$ and $a = \sum_{i=1}^n ib_i$.

Lemma 4.5. Let $n \in \{2, 3, ...\}$ and let \overline{G} be an n-coefficient complex. Let $(e, f, g) \in \overline{G}$ be a positive element and let there be given a decomposition $e = \sum_{j=1}^{k} e_j, e_j \in \mathcal{G}_0^+$. Then, there exist elements

$$g_1, g_2, \ldots, g_k \in \mathcal{G}_1$$
 and $f_1, f_2, \ldots, f_k \in \mathcal{G}_n$

such that

$$(e, f, g) = \sum_{j=1}^{k} (e_j, f_j, g_j)$$

and (e_j, f_j, g_j) is positive in \overline{G} for each $j \in \{1, \ldots, k\}$.

Proof: Since $\beta(f), g \leq \sum_{j=1}^{k} e_j$, there exist elements

$$g_1,\ldots,g_k,l_1,\ldots,l_k\in G_1$$

such that $\beta(f) = \sum_{j=1}^{k} l_j$ and $g = \sum_{j=1}^{k} g_j$ with $l_j, g_j \leq e_j$ (Lemma 4.3). As noted in the comment following that lemma, we may assume that $l_i \in \operatorname{im} \beta$, so by (vi) of Definition 2.1, the l_j have β -lifts \tilde{l}_j such that $\tilde{l}_j \leq e_j$. Thus, both f and $\sum_{j=1}^{k} \tilde{l}_j$ are majorised by e and have the same image under β . We conclude that the difference $f - \sum_{j=1}^{k} \tilde{l}_j$ is in the image of ρ , and is majorised by e. By property (v) of Definition 2.1, we may choose $c \in I(e)$ so that $f = \sum_{j=1}^{k} \tilde{l}_j + \rho(c), c \in I(e)$. Since $I(e) = I(e_1) + \cdots + I(e_k)$, there is a decomposition $c = c_1 + \cdots + c_k$, $c_j \in I(e_j)$. Put $f_j = \tilde{l}_j + \rho(c_j)$, so that $f = \sum_{j=1}^{k} f_j$. By construction, we have $f_j, g_j \leq e_j$, so that (e_j, f_j, g_j) is positive in the *n*-coefficient complex for each $j \in \{1, \ldots, k\}$.

Note that the lemma above holds even when one specifies the $g_j \leq e_j$ a priori.

In the following, we will use the term *refinement* of a collection of elements x_1, \ldots, x_s to denote a new collection of elements $\tilde{x}_1, \ldots, \tilde{x}_t$ with the property that $\{1, \ldots, t\}$ can be partitioned into s subsets, such that the sum of the elements corresponding to the indices in the *j*th subset is exactly x_j .

Lemma 4.6. Fix $n \in \{2, 3, ...\}$ and let \overline{G} be an n-coefficient complex. Let $(e_i, f_i, 0), i \in \{1, ..., k\}$, be positive in \overline{G} . Let there be given elements $x_1, ..., x_r \in G_0^+$ and $z_1, ..., z_r \in G_1$ such that $z_j \leq x_j$, and non-negative integers $\lambda_{ij}, \delta_{ij}, i \in \{1, ..., k\}, j \in \{1, ..., r\}$, such that

$$e_i = \sum_{j=1}^r \lambda_{ij} x_j, \quad \beta(f_i) = \sum_{j=1}^r \delta_{ij} z_j.$$

Then, there exist refinements $\tilde{x}_1, \ldots, \tilde{x}_s$ of x_1, \ldots, x_r and $\tilde{z}_1, \ldots, \tilde{z}_s$ of z_1, \ldots, z_r , and lifts $\tilde{y}_l \in G_n$ of the \tilde{z}_l with $\tilde{y}_l \leq \tilde{x}_l$, having the following property: there are non-negative integers γ_{il} , κ_{il} , and n_{il} , $i \in \{1, \ldots, k\}$ and $l \in \{1, \ldots, s\}$, such that

$$e_i = \sum_{l=1}^s \gamma_{il} \widetilde{x_l}$$

and

$$f_i = \sum_{l=1}^{s} \kappa_{il} \widetilde{y}_l + n_{il} \rho(\widetilde{x}_l)$$

Furthermore, $\gamma_{il} \neq 0$ whenever $n_{il} \neq 0$.

Proof: Following the proof of Lemma 4.5, we may assume that we have lifts y_i of each z_i , and positive elements $c_{ij} \in I(x_j)$ such that

 $f_i = \sum_{j=1}^r \delta_{ij} y_l + \rho(c_{ij})$ and $\delta_{ij} y_j + \rho(c_{ij}) \leq x_j, 1 \leq j \leq r, 1 \leq i \leq k$. Fix j. By Lemma 4.4 there is, for each $i \in \{1, \ldots, k\}$, a decomposition $x_j = x_{j,1}^i + \cdots + x_{j,k_i}^i, k_i \in \mathbb{N}$, such that c_{ij} is in the non-negative integral linear span of $\{x_{j,1}^i, \ldots, x_{j,k_i}^i\}$. Choose by the Riesz property in G_0 a decomposition $x_j = x_{j,1} + \cdots + x_{j,m_j}$, some $m_j \in \mathbb{N}$, which simultaneously refines all of the $x_j = x_{j,1}^i + \cdots + x_{j,k_i}^i$ decompositions, $1 \leq i \leq k$. Then, there exist non-negative integers $n_{j,1}^i, \ldots, n_{j,m_j}^i, 1 \leq i \leq k$, such that $c_{ij} = n_{j,1}^i x_{j,1} + \cdots + n_{j,m_j}^i x_{j,m_j}$. Since $\delta_{ij} y_j + \rho(c_{ij}) \leq x_j$ we have that $\delta_{ij} y_j \leq x_j$ (property (v) of Definition 2.1), whence $y_j \leq x_j = x_{j,1} + \cdots + x_{j,m_j}$ (Lemma 4.2). By Lemma 4.5 there is a decomposition $y_j = y_{j,1} + \cdots + y_{j,m_j}$ such that $y_{j,l} \leq x_{j,l}, 1 \leq l \leq m_j$. Thus,

$$\delta_{ij}y_j + \rho(c_{ij}) = \sum_{p=1}^{m_j} \delta_{ij}y_{j,p} + n^i_{j,p}\rho(x_{j,p}).$$

Define

$$\{\widetilde{x}_1,\ldots,\widetilde{x}_s\} := \bigcup_{j \le r} \{x_{j,1},\ldots,x_{j,m_j}\},\\ \{\widetilde{y}_1,\ldots,\widetilde{y}_s\} := \bigcup_{j \le r} \{y_{j,1},\ldots,y_{j,m_j}\},$$

and $\widetilde{z}_l := \beta(\widetilde{y}_l)$. The lemma follows.

5. Bounded Torsion

5.1. A Local Property. In this section we establish that n-coefficient complexes satisfy a local property such as the one whose importance was realized by Shen (cf. [9]) in the setting of classical dimension groups.

Lemma 5.1. Let $\overline{\mathcal{G}}$ be an n-coefficient complex. Let $\overline{G} = \bigoplus_{i=1}^{n} \overline{G_i}$ be a direct sum of n-coefficient complex (with the direct sum order structure), where each $\overline{G_i}$ is a (\mathbb{C}, n) , (\mathbb{I}_m^{\sim}, n) or $(C(S^1), n)$ complex. Let $\overline{\theta} : \overline{G} \to \overline{\mathcal{G}}$ be a morphism. Then, there exist an n-coefficient complex $\overline{H} = \bigoplus_{j=1}^{m} \overline{H_j}$ with each $\overline{H_j}$ a (\mathbb{C}, n) , (\mathbb{I}_m^{\sim}, n) or $(C(S^1), n)$ complex, and morphisms $\overline{\gamma} : \overline{G} \to \overline{H}$ and $\overline{\lambda} : \overline{H} \to \overline{\mathcal{G}}$ such that the diagram



commutes and ker $\overline{\gamma} = \ker \overline{\theta}$.

Proof: Suppose that the conclusion above is relaxed to read

$$\ker \overline{\gamma} = \ker \overline{\theta} \ (\mathrm{mod}\rho(\mathbf{G}_0)),$$

with all other things being equal. Then, the original conclusion of the lemma follows. Indeed, suppose that $a \in \ker \overline{\theta}$ is in the image of ρ , i.e., $a = \rho(q)$ for some $q \in G_0^+$. Then, $\theta_0(q) = n \cdot q'$ for some $q' \in \mathcal{G}_0^+$. Put $\overline{H}' = \overline{H} \oplus \overline{R}$, where \overline{R} is a (\mathbb{C}, n) complex, and extend $\overline{\lambda}$ to \overline{H}' by sending the positive generator of R_0 to q'. Apply the weakened conclusion above to find a direct sum of building block complexes \overline{H}'' and $\overline{\lambda}'' : \overline{H}' \to \overline{\mathcal{G}}$ such that the diagram



commutes. Note that $\ker(\overline{\gamma}'' \circ \overline{\gamma}) = \ker \overline{\theta} \pmod{\rho(G_0)}$. Furthermore, $a \in \ker(\overline{\gamma}'' \circ \overline{\gamma})$, since $(\gamma_0'' \circ \gamma_0)(q)$ must be $\operatorname{ord}(\rho(q))$ times the image of the positive generator of R_0 under γ_0'' . Repeating this procedure for each of the finitely many elements in $\ker \overline{\theta} \cap \rho(G_0)$ yields the conclusion of the lemma proper.

It remains to prove that the lemma holds if we only require that $\ker \overline{\gamma} = \ker \overline{\theta} \pmod{\rho(G_0)}$. Let $G_{0i} = \langle e_i \rangle \ (e_i \geq 0), \ G_{1i} = \langle g_i \rangle$, and choose $f_i \in G_{ni}$ (necessarily $\leq e_i$) such that $G_{ni} = \langle \rho(e_i) \rangle \oplus \langle f_i \rangle$. Note that if \overline{G}_i is a (\mathbb{C}, n) or $(C(S^1), n)$ complex, then we may (and do) take $f_i = 0$. In the case of a (I_m, n) complex, $\beta(f_i) = g_i$. Define $a_i := \theta_0(e_i), b_i := \theta_n(f_i)$ and $c_i := \theta_1(g_i)$. By the main Theorem in Section 5 of [15], there is a complex $\overline{H}' = \bigoplus_{l=1}^k \overline{H}'_l$ with each \overline{H}'_l a building block (and elements e'_i, f'_i and g'_l playing roles analogous to those of the e_i, f_i and g_i above), and maps

$$\gamma_0^{'}: \mathrm{G}_0 \to \mathrm{H}_0^{'}, \ \gamma_1^{'}: \mathrm{G}_1 \to \mathrm{H}_1^{'}$$

and

$$\lambda_{0}^{'}: \operatorname{H}_{0}^{'} \to \mathcal{G}_{0}, \ \lambda_{1}^{'}: \operatorname{H}_{1}^{'} \to \mathcal{G}_{1}$$

such that



commutes and ker $(\theta_0, \theta_1) = \text{ker}(\gamma'_0, \gamma'_1)$. To be fair, [15] only provides the H'_{0j} and H'_{1j} , but we may clearly associate a building block complex to any pair $(\text{H}'_{0j}, \text{H}'_{1j}) = (\mathbb{Z}, R), R \in \{0, \mathbb{Z}, \mathbb{Z}/2, \mathbb{Z}/3, \dots\}$. Of course, this association is token for now, as we have no positive maps γ'_n : $\text{G}_n \to \text{H}'_n$ and $\lambda'_n : \text{H}'_n \to \mathcal{G}_n$ to fill in the diagram above.

 $G_n \to H'_n$ and $\lambda'_n : H'_n \to \mathcal{G}_n$ to fill in the diagram above. We have $a_i = \sum_{l=1}^k \kappa_{il} \lambda'_0(e'_l), \mathbb{Z} \ni \kappa_{il} \ge 0$, and $\beta(b_i) = \sum_{l=1}^k \phi_{il} \lambda'_1(g'_l), \phi_{il} \in \mathbb{Z}$. By Lemma 4.6 there exist refinements

$$\{\widetilde{a_1}, \ldots, \widetilde{a_m}\}\ (\widetilde{a_j} \ge 0, 1 \le j \le m) \text{ and } \{\widetilde{c_1}, \ldots, \widetilde{c_m}\}\$$

of

$$\{\lambda_{0}^{'}(e_{1}^{'}),\ldots,\lambda_{0}^{'}(e_{k}^{'})\} \text{ and } \{\lambda_{1}^{'}(g_{1}^{'}),\ldots,\lambda_{1}^{'}(g_{k}^{'})\},\$$

respectively, and elements $b_1, \ldots, b_s \in \mathcal{G}_n$ such that for some integers $\zeta_{ij}, \xi_{ij}, \iota_{ij}$ and $n_{ij}, 1 \leq i \leq k, 1 \leq j \leq m$, we have

$$a_{i} = \sum_{j=1}^{m} \zeta_{ij} \widetilde{a}_{j},$$
$$b_{i} = \sum_{j=1}^{m} \xi_{ij} \widetilde{b}_{j} + n_{ij} \rho(\widetilde{a}_{j}),$$

and

$$c_i = \sum_{j=1}^m \zeta_{ij} \widetilde{c_j}.$$

Furthermore, $\widetilde{b_j}$ is a lift of $\widetilde{c_j}$ whenever $\widetilde{c_j}$ is in the image of β , and is zero otherwise. Note that the $\widetilde{c_j}$ can be and should be chosen to be in the image of β whenever it is a torsion element. By the main Theorem of Section 5, [15], there is an *n*-coefficient complex $\overline{H} = \bigoplus_{j=1}^m \overline{H_j}, \overline{H_j}$ a building block complex for each j, and there are maps

$$\gamma_0: \mathbf{G}_0 \to \mathbf{H}_0, \ \gamma_1: \mathbf{G}_1 \to \mathbf{H}_1$$

and

$$\lambda_0: \mathrm{H}_0 \to \mathcal{G}_0, \ \lambda_1: \mathrm{H}_1 \to \mathcal{G}_1$$

such that (with the dotted arrows representing desired but as yet undefined maps)



commutes, and γ_0 and γ_1 factor through H'_0 and H'_1 , respectively. Thus, ker $(\theta_0, \theta_1) = \operatorname{ker}(\gamma_0, \gamma_1)$. Let $\widetilde{e_j}$, $\widetilde{f_j}$, and $\widetilde{g_j}$ play roles in $\overline{H_j}$ analogous to the roles of the e_i , f_i , and g_i in $\overline{G_i}$. Then $\lambda_0(\widetilde{e_j}) = \widetilde{a_j}$ and $\lambda_1(\widetilde{g_j}) = \widetilde{c_j}$. For every pair (i, j), $1 \leq i \leq k$, $1 \leq j \leq m$, define a partial map $\gamma_n^{ij} : \operatorname{G}_{ni} \to \operatorname{H}_{nj}$ by

$$\gamma_n^{ij}(f_i) := \xi_{ij} \widetilde{f_j} + n_{ij} \rho(\widetilde{e_j}).$$

Let $\lambda_n : \mathrm{H}_n \to \mathcal{G}_n$ be defined by $\lambda(\widetilde{f}_j) := \widetilde{b}_j$, and put $\gamma_n = \bigoplus_{j=1}^m (\sum_{i=1}^k \gamma_n^{ij})$. These maps are positive and so complete the morphisms $\overline{\lambda}$ and $\overline{\gamma}$, establishing the desired weak version of the lemma. \Box

5.2. The Range Result.

Lemma 5.2. Fix $n \in \{2, 3, ...\}$, let $\overline{\mathcal{G}}$ be a n-coefficient complex, and consider a positive element $(e, f, g) \in \overline{\mathcal{G}}$. There exists a n-coefficient complex \overline{H} which is a finite direct sum of (\mathbb{C}, n) , (\mathbb{I}_m^{\sim}, n) and $(C(S^1), n)$ complexes, and a positive morphism $\overline{\theta} : \overline{H} \longrightarrow \overline{\mathcal{G}}$ such that

$$(e, f, g) \in \overline{\theta}(\overline{H}_+)$$

Proof: First consider the case when f = 0. We define $\overline{\theta}_g : \overline{H}_g \longrightarrow \overline{\mathcal{G}}$ by

$$1 \mapsto e; \overline{1} \mapsto \rho(e); 1 \mapsto g$$

on the $(C(S^1), n)$ -complex, and note that $(e, 0, g) = \overline{\theta}(1, \overline{0}, 1)$. Similarly, when g = 0, we define $\overline{\theta}_f : \overline{H}_f \longrightarrow \overline{\mathcal{G}}$ by

$$1\mapsto e; (\overline{1},\overline{0})\mapsto \rho(e), (\overline{0},\overline{1})\mapsto f; \overline{1}\mapsto \beta(f)$$

on the (\mathbb{I}_n^{\sim}, n) -complex, and note that $(e, f, 0) = \overline{\theta}(1, (\overline{0}, \overline{1}), \overline{0})$.

In the general case, we consider



with $\overline{\gamma}, \overline{\lambda}$ chosen by Lemma 5.1 above. By assumption,

$$x = \gamma_0((1,0)) = \gamma_0((0,1))$$

so that $(x, \gamma_n([(\overline{0}, \overline{0}), (\overline{0}, \overline{1})]), \gamma_1([1, \overline{0}]))$ is a positive preimage of (e, f, g).

Theorem 5.3. Fix $n \in \{2, 3, ...\}$, and let $\overline{\mathcal{G}}$ be an n-coefficient complex. Then, $\overline{\mathcal{G}} = \lim_{i \to \infty} \overline{G}_i$, where each \overline{G}_i is a finite direct sum of $(\mathbb{C}, n), (\mathbb{I}_m^{\sim}, n)$ and $(C(S^1), n)$ complexes.

Proof: Enumerate the positive elements of $\overline{\mathcal{G}}$ as (e_i, f_i, g_i) and apply Lemmas 5.2 and 5.1 alternately to get a diagram



where (e_i, f_i, g_i) is the image under $\overline{\theta}_i$ of a positive element and ker $\overline{\theta}_i =$ ker $\overline{\gamma}_i$. The maps will then induce an order isomorphism.

An inductive system of finite direct sums of building blocks is said to have *large denominators* if all connecting morphisms either are zero on K_1 or have the K_0 -component greater than or equal to 2.

Theorem 5.4. Let $n \in \{2, 3, ...\}$ and let \overline{G} be a complex. The following are equivalent

(i) \overline{G} is a n-coefficient complex;

- (ii) \overline{G} is an inductive limit of finite direct sums of (\mathbb{C}, n) , (\mathbb{I}_m^{\sim}, n) and $(C(S^1), n)$ complexes, and G_* has the Riesz property;
- (iii) \overline{G} is an inductive limit of finite direct sums of (\mathbb{C}, n) , (\mathbb{I}_m^{\sim}, n) and $(C(S^1), n)$ complexes, such that the inductive system has large denominators;
- (iv) $\overline{G} \cong \overline{K}_n(\mathfrak{A})$, where \mathfrak{A} is an AD algebra of real rank zero.

Proof: Note first that $(iv) \Longrightarrow (i)$ was seen in Proposition 3.2. Theorem 5.3 proves $(i) \Longrightarrow (ii)$, and since the property of large denominators involves only the groups in G_* , [16, 8.1] proves $(ii) \Longrightarrow (iii)$. By compressing an inductive system such as in (iii) if necessary, we may assume that each morphism among building blocks at level i and level i + 1 is either zero on K_1 or greater than or equal to M_i on K_0 , where M_i is the largest number for which there is an $(\mathbb{I}_{M_i}^{\sim}, n)$ complex among the building blocks at level i. Then by [11] the inductive system can be realized by direct sums of building blocks from the set $\{\mathbb{C}, C(S^1), \mathbb{I}_2^{\sim}, \mathbb{I}_3^{\sim}, \ldots\}$ and *-homomorphisms among them. Furthermore, [16, 8.1] shows how to arrange for real rank zero in the limit.

By construction, the inductive limit \mathfrak{A} of this C^* -inductive system is an AD algebra with the desired invariant

$$\overline{K}_n(\mathfrak{A}): \qquad \mathrm{K}_0(\mathfrak{A}) \longrightarrow \mathrm{K}_0(\mathfrak{A}; \mathbb{Z}/n) \longrightarrow \mathrm{K}_1(\mathfrak{A}).$$

However, since we have not used – or even defined – an ideal based order on the building blocks $C(S^1)$ and \mathbb{I}_m^{\sim} , cf. Definition 3.1, we need to verify that the order on $\overline{K}_n(\mathfrak{A})$ coincides with the order on \overline{G} . Since we have used the strict order on all the algebraic building blocks this would follow directly if we knew that all ideals of \mathfrak{A} arise as inductive limits or direct sums of subcollections of the building blocks in the system. And this in turn is a consequence of the minimal real rank of \mathfrak{A} , or directly by the construction yielding this property in [16, 8.1]. \Box

It is essential to note at this stage that the ordered complex $\overline{K}_n(\mathfrak{A})$ is not complete for real rank zero AD algebras unless we know that the torsion of K_1 is annihilated by the number n. Thus, it is only in this case — covered by [11] — that Theorem 5.4 gives a one-to-one correspondence between a class of C^* -algebras and a class of algebraic invariants.

In this case, when the equivalent statements above hold true, we may write the AD algebra as an inductive limit using only the building blocks $C(S^1)$ and \mathbb{I}_n^{\sim} , cf. [11]

6. The general case

In the following we shall briefly recall definitions from [4]. Let Δ denote the ordered set (\mathbb{N}, \leq) where

$$x \leq y \iff x \text{ divides } y.$$

Note that Δ is directed, so that we may construct inductive limits over Δ . We will denote these by

$$\lim_{\longrightarrow \Delta} (G_p, f_{q,p})$$

where $f_{q,p}: G_p \to G_q$ are the bonding maps. When a cofinal subset Δ' of Δ , is given, we may restrict attention to this, as

$$\lim_{\longrightarrow \Delta} G_n \cong \lim_{\longrightarrow \Delta'} G_n$$

We define graded group homomorphisms

$$\kappa_{mn,m} : \mathrm{K}_{0}(\mathfrak{A}; \mathbb{Z} \oplus \mathbb{Z}/m) \to \mathrm{K}_{0}(\mathfrak{A}; \mathbb{Z} \oplus \mathbb{Z}/mn)$$

by

$$\begin{bmatrix} \chi_{mn,n} & 0 \\ 0 & \kappa_{mn,m} \end{bmatrix},$$

where $\chi_{mn,n}$ is just multiplication by m between the relevant copies of $K_0(\mathfrak{A})$. The maps $\kappa_{mn,m}$ are positive, so we may define:

$$\begin{aligned} \mathrm{K}_{0}(\mathfrak{A};\mathbb{Q}\oplus\mathbb{Q}/\mathbb{Z}) &= \lim_{\longrightarrow\Delta} \left(\mathrm{K}_{0}(\mathfrak{A};\mathbb{Z}\oplus\mathbb{Z}/n),\kappa_{mn,n} \right); \\ \mathrm{K}_{0}(\mathfrak{A};\mathbb{Q}\oplus\mathbb{Q}/\mathbb{Z})^{+} &= \lim_{\longrightarrow\Delta} \left(\mathrm{K}_{0}(\mathfrak{A};\mathbb{Z}\oplus\mathbb{Z}/n)^{+},\kappa_{mn,n} \right). \end{aligned}$$

This gives the limit groups the structure of graded ordered groups. The even parts are naturally isomorphic to $K_0(\mathfrak{A}) \otimes \mathbb{Q}$ since

$$\lim_{d\to\Delta} (G, \chi_{mn,n}) \cong \lim_{d\to\Delta} (G \otimes \mathbb{Z}, \mathrm{id} \otimes \chi_{mn,n})$$
$$\cong G \otimes \left(\lim_{d\to\Delta} (\mathbb{Z}, \chi_{mn,n})\right) \cong G \otimes \mathbb{Q}$$

naturally. We shall invoke this isomorphism tacitly in section 7. The maps

$$\mathfrak{K}_{mn,m}: \mathrm{K}_{0}(\mathfrak{A}; \mathbb{Z} \oplus \mathbb{Z}/m) \to \mathrm{K}_{0}(\mathfrak{A}; \mathbb{Z} \oplus \mathbb{Z}/mn)$$

can be described explicitly when $\mathfrak{A} \in \{\mathbb{C}, C(S^1), \mathbb{I}_2^{\sim}, \mathbb{I}_3^{\sim}, \mathbb{I}_4^{\sim}, \dots\}$ ([10]). In this section we study exact sequences of abelian groups

$$\mathcal{G}_0 \longrightarrow \mathcal{G}_0 \otimes \mathbb{Q} \xrightarrow{\rho} \mathcal{G}_n \xrightarrow{\beta} \mathcal{G}_1$$

which are meant to represent the natural K-theoretic invariants for AD algebras of real rank zero having unbounded torsion in K_1 . We

begin by listing the properties that such an abstract sequence should have before one may even consider whether the sequence arises as the invariant

$$\overline{K}_{\mathbf{n}}: \qquad \qquad \mathrm{K}_{0}(\mathfrak{A}) \longrightarrow \mathrm{K}_{0}(\mathfrak{A}) \otimes \mathbb{Q} \xrightarrow{\rho} \mathrm{K}_{0}(\mathfrak{A}; \mathbb{Q}/\mathbb{Z}) \xrightarrow{\beta} \mathrm{K}_{1}(\mathfrak{A})$$

for some AD algebra \mathfrak{A} of real rank zero. We shall denote the invariant consisting of two graded ordered groups and two group homomorphisms as in (2) by $\underline{K}_{\mathbf{n}}(\mathfrak{A})$ and will see that the conditions in the definition below are sufficient to ensure that the sequence

$$\mathcal{G}_0 \longrightarrow \mathcal{G}_0 \otimes \mathbb{Q} \xrightarrow{\rho} \mathcal{G}_n \xrightarrow{\beta} \mathcal{G}_1$$

does indeed arise in such a manner.

The next definition should be compared with Definition 2.1.

Definition 6.1. An exact sequence

$$\mathcal{G}_0 \longrightarrow \mathcal{G}_0 \otimes \mathbb{Q} \xrightarrow{\rho} \mathcal{G}_n \xrightarrow{\beta} \mathcal{G}_1$$

(which we denote by $\overline{\mathcal{G}}$) of countably generated abelian groups is an **n**-coefficient complex if

- $(i_{\mathbf{n}}) \mathcal{G}_{\mathbf{n}}$ is pure torsion
- $(ii_{\mathbf{n}})$ im $\beta = \operatorname{tor} \mathcal{G}_{1}$, and every element $x \in \mathcal{G}_{1}$ of order l has a β -lift of order l
- (iii_n) $\mathcal{G}_* := \mathcal{G}_0 \oplus \mathcal{G}_1$ and $\mathcal{G}_{\underline{\mathbf{n}}} := \mathcal{G}_0 \otimes \mathbb{Q} \oplus \mathcal{G}_{\mathbf{n}}$ are graded ordered groups inducing the same order on \mathcal{G}_0
- $(iv_n) \mathcal{G}_*$ has the Riesz interpolation property.
- $(v_{\mathbf{n}}) \ \mathcal{G}_0 \otimes \mathbb{Q} \oplus \rho(\mathcal{G}_0)$ has the quotient order coming from $\mathrm{id}_{\mathcal{G}_0 \otimes \mathbb{Q}} \oplus \rho$
- $(vi_{\mathbf{n}}) \ \mathcal{G}_0 \oplus \beta(\mathcal{G}_{\mathbf{n}})$ has the quotient order coming from $id_{\mathcal{G}_0} \oplus \beta$

 $(vii_{\mathbf{n}}) \mathcal{G}_0$ is unperforated and \mathcal{G}_* is weakly unperforated.

Note that since \mathcal{G}_0 is torsion free, it is determined by $\mathcal{G}_0 \otimes \mathbb{Q}$ in this setup. In the proofs below we may hence concentrate our work on the rightmost three groups in the complex.

Proposition 6.2. Let

$$\overline{\mathcal{G}}: \qquad \mathcal{G}_0 \otimes \mathbb{Q} \stackrel{
ho}{\longrightarrow} \mathcal{G}_{\mathbf{n}} \stackrel{
ho}{\longrightarrow} \mathcal{G}_1$$

be an **n**-coefficient complex. Then, there exist a strictly increasing sequence of natural numbers $\langle n_i \rangle$ with $n_i | n_{i+1}$, a n_i -coefficient complex $\overline{G^i}$ for each i, and positive morphisms $\overline{\theta_i} : \overline{G^i} \to \overline{G^{i+1}}$ such that

$$\overline{\mathcal{G}} \simeq (\overline{G^i}, \overline{\theta_i}).$$

(2)

Proof: Let $\langle n_i \rangle$ be a strictly increasing sequence of natural numbers with the property that every natural number divides some n_i . By the main theorem of section 5 of [15], the graded ordered group $(\mathcal{G}_0 \otimes \mathbb{Q}, \mathcal{G}_1)$ in Definition 6.1 is the limit of an inductive sequence

$$((G_0^i, G_1^i), (\phi_0^i, \phi_1^i))$$

of graded ordered groups (G_0^i, G_1^i) . Furthermore, each (G_0^i, G_1^i) consists of the first and third groups of an n_i -coefficient complex which is a direct sum of (\mathbb{C}, n_i) , $(\mathbb{I}_m^{\sim}, n_i)$ and $(C(S^1), n_i)$ complexes. Let

$$\phi_0^{i\infty}: G_0^i \to \mathcal{G}_0, \quad \phi_1^{i\infty}: G_1^i \to \mathcal{G}_1,$$

be the canonical maps. Assume that the n_i have been chosen large enough for the inclusions $G_0^i \subseteq \rho^{-1}(\mathcal{G}_{\mathbf{n}}[n_i])$ and $\operatorname{tor}(G_1^i) \subseteq \beta(\mathcal{G}_{\mathbf{n}}[n_i])$ to hold. We then have

$$(\mathcal{G}_0 \otimes \mathbb{Q}, \mathcal{G}_1) = \lim_{i \to \infty} \left((\rho^{-1}(\mathcal{G}_{\mathbf{n}}[n_i]), \beta(\mathcal{G}_{\mathbf{n}}[n_i]) \cup \phi_1^{i\infty}(G_1^i)), \iota \right),$$

where ι is the inclusion map. Note that

$$(\rho^{-1}(\mathcal{G}_{\mathbf{n}}[n_i]), \beta(\mathcal{G}_{\mathbf{n}}[n_i]) \cup \phi_1^{i\infty}(G_1^i))$$

is a graded ordered group for each natural number i — it is an order hereditary subgroup of $(\mathcal{G}_0 \otimes \mathbb{Q}, \mathcal{G}_1)$. One can then verify that the complex

$$\rho^{-1}(\mathcal{G}_{\mathbf{n}}[n_i]) \xrightarrow{\rho} \mathcal{G}_{\mathbf{n}}[n_i] \xrightarrow{\beta} \beta(\mathcal{G}_{\mathbf{n}}[n_i]) \cup \phi_1^{i\infty}(G_1^i)$$

is an n_i -coefficient complex. (The only subtle point is the exactness of the sequence, which follows from the second half of property iii_n of Definition 6.1.) The limit of the inductive sequence

$$\left(\rho^{-1}(\mathcal{G}_{\mathbf{n}}[n_i]) \xrightarrow{\rho} \mathcal{G}_{\mathbf{n}}[n_i] \xrightarrow{\beta} \beta(\mathcal{G}_{\mathbf{n}}[n_i]) \cup \phi_1^{i\infty}(G_1^i), \kappa_{i,i+1}\right)$$

 $-\kappa_{i,i+1}$ is the inclusion map — is then

$$\mathcal{G}_0\otimes \mathbb{Q} \stackrel{
ho}{\longrightarrow} \mathcal{G}_{\mathbf{n}} \stackrel{
ho}{\longrightarrow} \mathcal{G}_1$$

by construction.

The following theorem is the generalised integer coefficient version of Theorem 5.3.

Theorem 6.3. Let $\overline{\mathcal{G}}$ be an **n**-coefficient complex. Then,

$$\overline{\mathcal{G}} = \lim_{i \to \infty} (\overline{H}_i, \overline{\gamma_i}),$$

where each \overline{H}_i is a direct sum of (\mathbb{C}, n_i) , $(\mathbb{I}_m^{\sim}, n_i)$ and $(C(S^1), n_i)$ complexes, some $n_i \in \mathbb{N}$.

Proof: Assume the inductive sequence decomposition of $\overline{\mathcal{G}}$ from Lemma 7.2. For brevity, write

$$\overline{H}^{n_i} = \rho^{-1}(\mathcal{G}_{\mathbf{n}}[n_i]) \xrightarrow{\rho} \mathcal{G}_{\mathbf{n}}[n_i] \xrightarrow{\beta} \beta(\mathcal{G}_{\mathbf{n}}[n_i]) \cup \phi_1^{i\infty}(G_1^i)$$

Put $\kappa_{l,m} = \kappa_{m-1,m} \circ \cdots \circ \kappa_{l,l+1}$. By Theorem 5.3, each \overline{H}^{n_i} is the limit of an inductive system $(\overline{H}_k^{n_i}, \overline{\theta}_{k,k+1}^{n_i})$, where each $\overline{H}_k^{n_i}$ is a direct sum of $(\mathbb{C}, n_i), (I_m^{\sim}, n_i)$ and (S^1, n_i) complexes.

It will suffice to define a sequence of positive morphisms

$$\gamma_{i,i+1}: \overline{H}_i^{n_i} \to \overline{H}_{i+1}^{n_{i+1}}$$

making the diagram



commute; by compressing the sequence for H^{n_i} one can ensure that every positive element in \mathcal{G} is the image under $\gamma_{i,\infty}$ of a positive element in $\overline{H}_i^{n_i}$ for some $i \in \mathbb{N}$. Let $\overline{\theta}_{1,\infty}^{n_1} : \overline{H}_1^{n_1} \to \overline{H}^{n_1}$ be the canonical morphism. Let M be a

Let $\overline{\theta}_{1,\infty}^{n_1} : \overline{H}_1^{n_1} \to \overline{H}^{n_1}$ be the canonical morphism. Let M be a minimal set of positive generators for $\overline{H}_1^{n_1}$. Find, by compressing the inductive sequence for \overline{H}^{n_2} if necessary, a set \widetilde{M} of positive pre-images via $\overline{\theta}_{1,\infty}^{n_2}$ of the elements of $\kappa_{1,2} \circ \overline{\theta}_{1,\infty}(M)$ in $\overline{H}_2^{n_2}$. Note for future reference that each element of $\kappa_{1,2} \circ \overline{\theta}_{1,\infty}(M)$ is divisible by n_{i+1}/n_i inside \overline{H}^{n_2} , so we may assume that $m(n_i/n_{i+1}) \in \overline{H}_2^{n_2}$ whenever $m \in \widetilde{M}$. Define $\gamma_{1,2}$ by sending an element $m \in M$ to the corresponding pre-image of $\kappa_{1,2} \circ \overline{\theta}_{1,\infty}(m)$ in \widetilde{M} .

The following theorem is the generalised integer coefficients version of Theorem 5.4.

Theorem 6.4. The following are equivalent:

- (i) $\overline{\mathcal{G}}$ is a **n**-coefficient complex;
- (ii) $\overline{\mathcal{G}}$ is an inductive limit of finite direct sums of (\mathbb{C}, n) , $(\mathbb{I}_{m}^{\sim}, n)$ and $(C(S^{1}), n)$ complexes, where n ranges over the natural numbers, and \mathcal{G}_{*} is a Riesz group;
- (iii) $\overline{\mathcal{G}}$ is an inductive limit of finite direct sums of (\mathbb{C}, n) , $(\mathbb{I}_{m}^{\sim}, n)$ and $(C(S^{1}), n)$ complexes where n ranges over the natural numbers, and such that the inductive system has large denominators;
- (iv) $\overline{\mathcal{G}} \cong \underline{K}_{\mathbf{n}}(\mathfrak{A})$, where \mathfrak{A} is an AD algebra of real rank zero.

Proof: The proof follows the proof of Theorem 5.4 with the exception that in $(iii) \Longrightarrow (iv)$, one needs to realize maps from $\overline{K}_{n_i}(\mathfrak{A}_i)$ to $\overline{K}_{n_{i+1}}(\mathfrak{A}_i)$ by a triple of maps of the form

$$(\chi_{n_{i+1},n_i} \circ f_*, \kappa_{n_{i+1},n_i} \circ f_*, f_*)$$

rather than directly by a *-homomorphism. However, as noted at the end of the proof of Theorem 6.4, the maps in question will have a K₀-component which is divisible by n_{i+1}/n_i , so this may be arranged as in the proof of Theorem 5.4.

7. The example of Dadarlat and Loring

It follows from the work of Bödigheimer ([1], [2]) that the unspliced short exact sequence

$$0 \longrightarrow \mathrm{K}_{0}(\mathfrak{A})/n \longrightarrow \mathrm{K}_{0}(\mathfrak{A}; \mathbb{Z}/n) \longrightarrow \mathrm{K}_{1}(\mathfrak{A})[n] \longrightarrow 0$$

will always split. This has been useful in the analysis of other aspects of this object ([12], [13]) but we have not been able to employ the fact in the proofs leading to Theorem 5.4 – since the splitting map is unnatural, it is difficult to use it when trying to establish the range of the invariant. By contrast, it is a useful result when trying to describe the amount of freedom one has in the choice of equipping \overline{G} as an *n*-coefficient complex when G_* is fixed, as we shall se below.

However, since Theorem 5.4 combines with the results mentioned above to prove that every *n*-coefficient complex will split when unspliced, it is perhaps worthwhile to note that this follows already from properties (i), (ii) and the fact (contained in (vii)) that G_0 is torsion free. One proves this by first establishing that im ρ is a pure subgroup of $G_{\overline{P}}$ and then appealing to [18]. **Remark 7.1.** For G_0 not necessarily torsion free the properties (i)–(vi), (viii)-(ix) would not imply splitness, as the example

 $\mathbb{Z}[\frac{1}{4}] \oplus \mathbb{Z}/2 \longrightarrow \mathbb{Z}/4 \longrightarrow \mathbb{Z}/2$

shows. When this is equipped with the strict order induced by the standard order on $\mathbb{Z}[\frac{1}{4}]$ it has all the properties of our 4-coefficient complexes except unperforation, but could not be the augmented K-theory of a C^{*}algebra. Thus to extend range results beyond the case considered above, on would have to impose an extra condition; for instance that im ρ was a pure subgroup of G_n .

Using the splitness, we may, up to isomorphism, for any *n*-coefficient complex write $G_{[n]} = R \oplus B$ such that $R = G_0/n$, $B = G_1[n]$ and

$$\rho(x) = (x + n\mathbf{G}_0, 0) \qquad \beta(r, b) = b$$

The properties (i)-(ix) simplify accordingly.

Remark 7.2. Note that graded ordered every dimension group with torsion G_* can be extended to an n-coefficient complex when G_0 is unperforated, simply by ordering $G_0 \oplus (R \oplus B)$ by

$$(x, (r, b)) \Longleftrightarrow \begin{cases} (x, (r, 0)) \ge 0\\ (x, b) \ge 0 \end{cases}$$

where the quotient order of $G_0 \oplus G_0$ and the order on G_* , respectively, are used to determine whether (x, (r, 0)) and (x, b) are positive.

Fix G_* . We have seen that up to isomorphism, every *n*-coefficient complex is of the form

$$G_0 \longrightarrow R \oplus B \longrightarrow G_1,$$

so determining how many *n*-coefficient complexes with this particular G_* are possible comes out to determining which order structures on $G_{\overline{III}}$ will satisfy properties (*iii*), (*v*), (*vi*), (*viii*) and (*ix*).

As an example this process, and an application of Theorem 5.4, let us return to the example of Dadarlat and Loring which originally established the need for ordered K-theory with coefficients. They considered the G_{*}-group given by

(3)
$$G_0 = \left\{ (x, y_i) \in \mathbb{Z}[\frac{1}{n+1}] \oplus \prod_{-\infty}^{\infty} \mathbb{Z} \mid y_i = x(n+1)^{|i|}, a.e.(i) \right\}$$

 $G_1 = \mathbb{Z}/n$

equipped with the standard order on G_0 and the strict order herefrom on G_* . In [7] examples were given to show, in effect, that there were two different ways to complete G_* to an *n*-coefficient complex.

For convenience, let n = 2. With the notation above we have

$$R = \left\{ (a, b_i) \in \mathbb{Z}/2 \oplus \prod_{-\infty}^{\infty} \mathbb{Z}/2 \mid b_i = a, a.e.(i) \right\}$$
$$B = \{c \in \mathbb{Z}/2\}$$

where

$$\rho(x, y_i) = (\overline{a}, \overline{y_i}) \text{ with } x = \frac{a}{3^i}.$$

In the proof below we need elements $\delta_j, \Delta_N \in (\mathbb{Z}/2)^{\mathbb{Z}}$ defined by

$$(\delta_j)_i = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases} \qquad (\Delta_N)_i = \begin{cases} 1 & |i| > N \\ 0 & |i| \le N \end{cases}$$

Proposition 7.3. Up to isomorphism every 2-coefficient complex completing G_* defined in (3) will order $G_{\overline{Pl}}$ by

$$((x, y_i), (a, b_i, c)) \ge 0 \iff \begin{cases} x \vdash a \\ y_i \vdash b_i + \epsilon_i c \\ x \vdash c \end{cases}$$

where $(\epsilon_i)_{i\in\mathbb{Z}}$ is a sequence in $(\mathbb{Z}/2)^{\mathbb{Z}}$.

Proof: It is easy to see that this defines a 2-coefficient complex. In the other direction, first note that since the order of $G_0 \oplus (R \oplus 0)$ is induced by the graded ordered group $G_0 \oplus G_0$ we have

$$((x, y_i), (a, b_i, 0)) \ge 0 \iff \begin{cases} x \vdash a \\ y_i \vdash b_i \end{cases}$$

Lift the element $((1, 3^{|i|}(1 - \delta_j)), 1) \in G_*$ to a positive element $((1, 3^{|i|}(1 - \delta_j)), (a, b_i, 1) \in G_{\overline{[n]}}$. If a = 1, note that also

$$((1,3^{|i|}(1-\delta_j)),(a+1,b_i+1-\delta_j,1+0))$$

is positive, so that we may without loss of generality assume that a = 0. Similarly, we may assume that $b_i = 0$ for all $i \neq j$.

We have hence seen that at least one of

$$((1,3^{|i|}(1-\delta_j)),(0,\delta_j,1)) \qquad ((1,3^{|i|}(1-\delta_j)),(0,0,1))$$

is positive in $G_{[n]}$. We will define ϵ_j accordingly such that

$$((1, 3^{|i|}(1 - \delta_j)), (0, \epsilon_j \delta_j, 1)) \ge 0.$$

If $((x, y_i), (a, b_i, 1)) \ge 0$ then x > 0 because $(x, 1) \ge 0$ in G_* . Further, if $y_i = 0$ then since

$$((1+x,3^{|i|}(1-\delta_j)+y_i),(a,b_i+\epsilon_i\delta_j,1+1)) \ge 0$$

we have that $b_j + \epsilon_j = 0$. We conclude that $x \vdash a, c$ and $y_j \vdash b_j + \epsilon_j c_j$.

In the other direction, assume that $x \vdash a, c$ and $y_j \vdash b_j + \epsilon_j c$. We already know that $((x, y_i), (a, b_i, c)) \geq 0$ when c = 0, so we can focus on the case c = 1. In this case we will have x > 0 and hence $y_i > 0$ for $|j| \geq N$ for some $N \in \mathbb{N}$. We lift $((x, y_i), 1)$ to some positive element $((x, y_i), (a', b'_i, 1))$. If $a \neq a'$, we note that $((x, y_i), (1, \Delta_N, 0)) \geq 0$ whence also

$$((x, y_i), (a'+1, b'_i + \Delta_N, 1)) \ge 0$$

so that we without loss of generality may assume that a = a' and hence that $b_i = b'_i$ for all but finitely many *i*. For the remaining *i*s, if $y_i > 0$, we can adjust to get $b_j = b'_j$ in a similar fashion. And if $y_i = 0$ then since we have

$$((x, y_i)(a + a', b_j + b'_j, 1 + 1)) = ((x, y_i)(0, b_j + b'_j, 0)) \ge 0$$

we get that $b_i = b'_i$.

Let "~" be the finest equivalence relation on $(\mathbb{Z}/2)^{\mathbb{Z}}$ such that

$$(\epsilon_j) \sim (\epsilon_{j+k}) \sim (1+\epsilon_j) \sim (\epsilon_j \Delta_N(j)) \sim (\epsilon_{(-1)^{r_{|j|}}j})$$

for any $k \in \mathbb{Z}$, $N \in \mathbb{N}$ and any $r_j \in (\mathbb{Z}/2)^{\mathbb{N} \cup \{0\}}$. Using methods from [7] one can prove that the augmentations associated to (ϵ_i) and (η_i) in $(\mathbb{Z}/2)^{\mathbb{Z}}$ are isomorphic precisely when $(\epsilon_i) \sim (\eta_i)$. The examples given in [7] correspond to $(\epsilon_i) = (0)$ and (η_i) given by 1 on positive entries and 0 on negative ones.

One sees easily that there are uncountably many nonisomorphic 2coefficient complexes in this case — even though we have only added one bit of information to $G_0 \oplus (R \oplus 0)$ the amount of freedom in choosing the order structure is immense.

References

- C.F. Bödigheimer, Splitting the Künneth sequence in K-theory, Math. Ann. 242 (1979), 159–171.
- [2] _____, Splitting the Künneth sequence in K-theory, II, Math. Ann. 251 (1980), 249–252.
- [3] L.G. Brown, The Riesz interpolation property for $K_0(A) \oplus K_1(A)$, preprint.
- [4] M. Dădărlat and S. Eilers, Compressing coefficients while preserving ideals in the K-theory for C*-algebras, K-Theory 14 (1998), 281–304.
- [5] _____, The Bockstein map is necessary, Canad. Math. Bull. 42 (1999), no. 3, 274–284. MR 2000d:46070

- [6] M. Dădărlat and G. Gong, A classification result for approximately homogeneous C^{*}-algebras of real rank zero, Geom. Funct. Anal. 7 (1997), no. 4, 646–711.
- [7] M. Dădărlat and T.A. Loring, Classifying C*-algebras via ordered, mod-p Ktheory, Math. Ann. 305 (1996), no. 4, 601–616.
- [8] M. Dădărlat and T.A. Loring, A universal multicoefficient theorem for the Kasparov groups, Duke Math. J. 84 (1996), no. 2, 355–377.
- [9] E.G. Effros, D.E. Handelman, and C.L. Shen, Dimension groups and their affine representations, Amer. J. Math. 102 (1980), no. 2, 385–407. MR 83g:46061
- [10] S. Eilers, *Invariants for AD algebras*, Ph.D. thesis, Copenhagen University, November 1995.
- [11] _____, A complete invariant for AD algebras with real rank zero and bounded torsion in K_1 , J. Funct. Anal. **139** (1996), 325–348.
- S. Eilers, Künneth splittings and classification of C*-algebras with finitely many ideals, Operator algebras and their applications (Waterloo, ON, 1994/1995), Fields Inst. Commun., vol. 13, Amer. Math. Soc., Providence, RI, 1997, pp. 81– 90.
- [13] _____, Approximate homogeneity of C*-algebras with finitely many ideals, Math. Proc. R. Ir. Acad. 101A (2001), no. 2, 149–162. MR 1 925 347
- [14] S. Eilers and G.A. Elliott, The Riesz property for the K_{*}-group of a C^{*}-algebra of minimal stable and real rank, C. R. Math. Acad. Sci. Soc. R. Can. 25 (2003), no. 4, 108–113. MR 2 013 159
- [15] G.A. Elliott, Dimension groups with torsion, Internat. J. Math. 1 (1990), no. 4, 361–380.
- [16] _____, On the classification of C*-algebras of real rank zero, J. Reine Angew. Math. 443 (1993), 179–219.
- [17] L. Fuchs, *Riesz groups*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. **19** (1965), 1–34.
- [18] _____, Infinite abelian groups I, Academic Press, New York, San Francisco, London, 1970.
- [19] K. R. Goodearl, K₀ of multiplier algebras of C^{*}-algebras with real rank zero, K-Theory **10** (1996), no. 5, 419–489.
- [20] K.R. Goodearl, Partially ordered abelian groups with interpolation, American Mathematical Society, Providence, R.I., 1986.
- [21] H. Lin and M. Rørdam, Extensions of inductive limits of circle algebras, J. London Math. Soc. 51 (1995), no. 2, 603–613.
- [22] F. Riesz, Sur quelques notions fondamentales dans la théorie générale des opérations linéaires, Ann. of Math. 41 (1940), 174–206. MR 1,147d
- [23] C. Schochet, Topological methods for C*-algebras IV: Mod p homology, Pacific J. Math. 114 (1984), 447–468.
- [24] J. Villadsen, The range of the elliott invariant of the simple ah-algebras with slow dimension growth, K-theory 15 (1998), no. 1, 1–12.
- [25] F. Wehrung, Injective positively ordered monoids. I, J. Pure Appl. Algebra 83 (1992), no. 1, 43–82. MR 93k:06023
- [26] S. Zhang, A Riesz decomposition property and ideal structure of multiplier algebras, J. Operator Theory 24 (1990), 209–225.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COPENHAGEN, UNIVER-SITETSPARKEN 5, 2100 KØBENHAVN, DANMARK *E-mail address*: eilers@math.ku.dk

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW BRUNSWICK, FREDERICTON, NEW BRUNSWICK, CANADA, E3B 5A3 *E-mail address*: atoms@unb.ca