

Deformed multiplication in the semigroup \mathcal{PT}_n

Galyna Tsyaputa

Abstract

Pairwise non isomorphic semigroups obtained from the semigroup \mathcal{PT}_n of all partial transformations by the deformed multiplication proposed by Ljapin are classified.

1 Introduction

Let X and Y be two nonempty sets, S be the set of maps from X to Y . Fix some $\alpha : Y \rightarrow X$ and define the multiplication of elements in S in the following way: $\phi \circ \psi = \phi \alpha \psi$ (the composition of the maps is from left to right). The action defined by this rule is associative. In his famous monograph [1, p.353] Ljapin set the problem of investigation of the properties of this semigroup depending on the restrictions on set S and map α .

Magill [2] studies this problem in the case when X and Y are topological spaces and the maps are continuous. In particular under the assumption that α be onto he describes the automorphisms of such semigroups and determines the isomorphism criterion.

A bit later Sullivan [3] proves that if $|Y| \leq |X|$ then Ljapin's semigroup is embedding into transformation semigroup on the set $X \cup \{a\}$, $a \notin X$.

For us the important case is if $X = Y$, T_X is a transformation semigroup on the set X , $\alpha \in T_X$. Symons [4] establishes the isomorphism criterion for such semigroups and investigates the properties of their automorphisms.

The latter problem may be generalized to arbitrary semigroup S : for a fixed $a \in S$ define the operation $*_a$ via $x *_a y = xay$. The obtained semigroup is denoted by $(S, *_a)$ and operation $*_a$ is called the multiplication deformed by element a (or just the deformed multiplication).

In the paper we classify with respect to isomorphism all semigroups which are obtained from the semigroup of all partial transformations of an n -element set by the deformed multiplication.

2 \mathcal{PT}_n with the deformed multiplication

Let \mathcal{PT}_n be the semigroup of all partial transformations of the set $N = \{1, 2, \dots, n\}$. For arbitrary partial transformation $a \in \mathcal{PT}_n$ denote by $\text{dom}(a)$ the domain of a , and by $\text{ran}(a)$ its image. The value $|\text{ran}(a)|$ is called the rank of a and is denoted by $\text{rank}(a)$. Denote

$Z(a) = N \setminus \text{dom}(a)$ the set of those elements on which transformation a is not defined, and denote z_a the cardinality of this set. The type of $a \in \mathcal{PT}_n$ we call the set $(\alpha_1, \alpha_2, \dots, \alpha_n)$, where α_k is the number of those elements $y \in N$, the full inverse image $a^{-1}(y)$ of which contains exactly k elements. It is obvious that $1 \cdot \alpha_1 + 2 \cdot \alpha_2 + \dots + n \cdot \alpha_n = |\text{dom}(a)|$, and the sum $\alpha_1 + \alpha_2 + \dots + \alpha_n$ is equal to the rank of a .

Definition 1. Element $x \in \mathcal{S}$ is called left (right) annihilator of semigroup \mathcal{S}^0 provided that $xs = 0$ ($sx = 0$), $s \in \mathcal{S}$.

Element which is both the left and the right annihilator is called the annihilator of semigroup \mathcal{S} .

Proposition 1. Transformations $x \in \mathcal{PT}_n$ such that $\text{ran}(x) \subset Z(a)$ are left annihilators of semigroup $(\mathcal{PT}_n, *_a)$. The number of left annihilators equals $(z_a + 1)^n$.

Transformations $y \in \mathcal{PT}_n$ such that $Z(y) \supset \text{ran}(a)$ are right annihilators of semigroup $(\mathcal{PT}_n, *_a)$. The number of right annihilators equals $(n + 1)^{n - \text{rank}(a)}$.

Transformations $c \in \mathcal{PT}_n$ such that $\text{ran}(c) \subset Z(a)$ and $Z(c) \supset \text{ran}(a)$ are annihilators of $(\mathcal{PT}_n, *_a)$, moreover, the number of annihilators equals $(z_a + 1)^{n - \text{rank}(a)}$.

Proof. Let $x \in \mathcal{PT}_n$ be left annihilator of $(\mathcal{PT}_n, *_a)$. This means that for arbitrary u from \mathcal{PT}_n $x *_a u = 0$ or, what is the same, $xa \cdot u = 0$. Therefore xa is left zero of semigroup \mathcal{PT}_n , that is, a nowhere defined map. The latter is possible if and only if $\text{ran}(x) \subset Z(a)$. Evidently, the condition $\text{ran}(x) \subset Z(a)$ is sufficient for partial transformation x be the left annihilator of $(\mathcal{PT}_n, *_a)$. Now it is clear, that the number of such transformations is equal to $(z_a + 1)^n$.

Let $y \in \mathcal{PT}_n$ be the right annihilator of semigroup $(\mathcal{PT}_n, *_a)$. Then for any v from \mathcal{PT}_n $v \cdot ay = 0$ and ay is right zero of semigroup \mathcal{PT}_n , that is, nowhere defined map. This is equivalent to $Z(y) \supset \text{ran}(a)$. Therefore right annihilators of semigroup $(\mathcal{PT}_n, *_a)$ are those partial transformations \mathcal{PT}_n which are defined only in elements $N \setminus \text{ran}(a)$. It is clear that the number of such transformations equals $(n + 1)^{n - \text{rank}(a)}$.

Now the statement about annihilators follows from the definition and the above arguments. \square

On semigroup $(\mathcal{PT}_n, *_a)$ define the equivalence relation \sim_a by the rule: $x \sim_a y$ if and only if $x *_a u = y *_a u$ for all $u \in \mathcal{PT}_n$. Analogously on $(\mathcal{PT}_n, *_b)$ define the relation \sim_b .

Lemma 1. For an arbitrary isomorphism $\varphi : (\mathcal{PT}_n, *_a) \rightarrow (\mathcal{PT}_n, *_b)$ $\varphi(x) \sim_b \varphi(y)$ if and only if $x \sim_a y$.

Proof. In fact, let $\varphi : (\mathcal{PT}_n, *_a) \rightarrow (\mathcal{PT}_n, *_b)$ be isomorphism and $x \sim_a y$. Then for all $u \in \mathcal{PT}_n$: $x *_a u = y *_a u$, therefore $\varphi(x) *_b \varphi(u) = \varphi(y) *_b \varphi(u)$. However $\varphi(u)$ runs over the whole set \mathcal{PT}_n , hence $\varphi(x) \sim_b \varphi(y)$. Since the inverse map $\varphi^{-1} : (\mathcal{PT}_n, *_b) \rightarrow (\mathcal{PT}_n, *_a)$ is also isomorphism, $\varphi(x) \sim_b \varphi(y)$ implies $x \sim_a y$. Therefore, $x \sim_a y$ if and only if $\varphi(x) \sim_b \varphi(y)$. \square

Lemma 2. $x \sim_a y$ if and only if $xa = ya$.

Proof. Obviously, the equality $xa = ya$ implies $x \sim_a y$. Now let $xa \neq ya$. Then there exists k in N , such that $(xa)(k) \neq (ya)(k)$. Chose element u in \mathcal{PT}_n which has different images in the points $(xa)(k)$ and $(ya)(k)$. Then $x *_a u = xau$ and $y *_a u = yau$ have different images in k . Hence $x *_a u \neq y *_a u$ and $x \not\sim_a y$. \square

Theorem 1. *Semigroups $(\mathcal{PT}_n, *_a)$ and $(\mathcal{PT}_n, *_b)$ are isomorphic if and only if partial transformations a and b have the same type.*

Proof. Necessity. Let $(\mathcal{PT}_n, *_a)$ and $(\mathcal{PT}_n, *_b)$ be isomorphic. By lemma 1 arbitrary isomorphism between $(\mathcal{PT}_n, *_a)$ and $(\mathcal{PT}_n, *_b)$ maps equivalence classes of the relation \sim_a into equivalence classes of the relation \sim_b . Therefore for equivalence relations \sim_a and \sim_b cardinalities and the number of equivalence classes must be equal. We show that by the cardinalities of the classes of the relation \sim_a the type $(\alpha_1, \alpha_2, \dots, \alpha_n)$ of transformation a can be found uniquely.

Denote ρ_a the partition of set $\{1, 2, \dots, n\}$ induced by $a \in \mathcal{PT}_n$ (that is, l and m belong to the same block of the partition ρ_a provided that $a(l) = a(m)$; $Z(a)$ forms a separate block). We count the cardinality of equivalence class $\overline{x_0} = \{x | xa = x_0a\}$ of the relation \sim_a for the fixed transformation $x_0 \in \mathcal{PT}_n$. Consider element

$$y := x_0a = \begin{pmatrix} i_1 & i_2 & \dots & i_p & i_{p+1} & \dots & i_n \\ y_1 & y_2 & \dots & y_p & \emptyset & \dots & \emptyset \end{pmatrix}.$$

Obviously y_i belongs to the image of a , $i = 1, \dots, p$. Denote $N_a(a_i)$ the block of the partition ρ_a which is defined by a_i in the image of a . Denote $n_a(a_i)$ the cardinality of this block. The equality $xa = y$ is equivalent to that for every i $(xa)(i) = y_i$, or, what is the same, $x(i) \in N_a(y_i)$. Hence $x(i)$ can be chosen in $n_a(y_i)$ ways, $i = 1, \dots, p$. For $i = p+1, \dots, n$ the meaning of $x(i)$ can be chosen in $z_a + 1$ ways. Since the images of x in different points are chosen independently, partial transformation x can be defined in

$$n_a(y_1)n_a(y_2) \dots n_a(y_p)(z_a + 1)^{n-p} \quad (1)$$

ways, and this gives the cardinality of class $\overline{x_0}$.

It is clear that $\overline{0}$ is the set of all left annihilators of the semigroup $(\mathcal{PT}_n, *_a)$. By proposition 1 the cardinality of class $\overline{0}$ equals $(z_a + 1)^n$. Hence the value z_a is defined by abstract property the semigroup $(\mathcal{PT}_n, *_a)$. Denote m the smallest cardinality of the blocks of the partition ρ_a . Then the cardinality of the equivalence class of the relation \sim_a is the least if in (1) all multipliers are equal to $\min(m, z_a + 1)$. Consider the corresponding cases.

Let $m > z_a + 1$. Then class $\overline{0}$ is the only equivalence class of the least possible cardinality. The next larger class contains $m(z_a + 1)^{n-1}$ transformations. Now count the number of different equivalence classes of the relation \sim_a which have the cardinality $m(z_a + 1)^{n-1}$. Since $m > z_a + 1$ and this is the least of the cardinalities of the blocks of the partition ρ_a , to make the product in (1) be equal to $m(z_a + 1)^{n-1}$ there should be $p = 1$ and $n_a(y_1) = m$. Element i_1 can be chosen in n ways. We know that $|\{t : n_a(t) = m\}| = \alpha_m$, so y_1 can have α_m different meanings. Therefore, there are $n \cdot \alpha_m$ different transformations

$y = x_0 a$, such that the corresponding class $\overline{x_0}$ contains $m(z_a + 1)^{n-1}$ elements. Therefore by the relation \sim_a we may define the number m and the meaning α_m of the first nonzero component of the type $(\alpha_1, \alpha_2, \dots, \alpha_n)$ of element a . Components α_l for $l > m$ can be defined recursively. Assume that $\alpha_1, \alpha_2, \dots, \alpha_l$ are found. For the relation \sim_a denote C the number of equivalence classes of the cardinality $l \cdot (z_a + 1)^{n-1}$. Then C is equal to the number of sets $(p; i_1, i_2, \dots, i_p)$, where some of i_1, i_2, \dots, i_p may coincide in general, such that

$$n_a(y_{i_1})n_a(y_{i_2}) \dots n_a(y_{i_p})(z_a + 1)^{n-p} = l \cdot (z_a + 1)^{n-1}. \quad (2)$$

Since $\alpha_1, \alpha_2, \dots, \alpha_{l-1}$ are known, we may find the number A of sets $(p; i_1, i_2, \dots, i_p)$ such that all multipliers in the left hand side of (2) are less than l . This value equals

$$\sum_{k=1}^n \sum_{\substack{(m_1, \dots, m_k) \\ m \leq m_1, \dots, m_k \leq l \\ m_1 \cdot m_2 \dots m_k = l \cdot (z_a + 1)^{n-1}}} \binom{n}{k} \prod_{j=1}^k \alpha_{m_j}.$$

The number of sets $(p; i_1, i_2, \dots, i_n)$ such that one of the multipliers in the left hand side of (2) equals l , is equal to $n \cdot \alpha_l$. Then α_l can be found from the equality $A + n \cdot \alpha_l = C$.

Now let $m \leq z_a + 1$. Then the cardinality of the equivalence class of the relation \sim_a is the least if in (1) all multipliers are equal to m , that is, this cardinality equals m^n . Count the number of different equivalence classes $\overline{x_0}$ of the relation \sim_a of the cardinality m^n . To make all multipliers in (1) equal to m , there should be $n_a(y_i) = m$ for all i . However $|\{t : n_a(t) = m\}| = \alpha_m$. So every y_i can be chosen in α_m ways. Since the meanings of y_i for different i are chosen independently, there are α_m^n different $y = x_0 a$, such that the corresponding class $\overline{x_0}$ contains m^n elements. Hence by the relation \sim_a we may define the number m and the meaning α_m of the first non zero component of the type $(\alpha_1, \alpha_2, \dots, \alpha_n)$ of a .

Components α_l for $l > m$ can be found recursively applying the same arguments as above. Some changes include the following: in (2) the right hand side must be substituted with $l \cdot m^{n-1}$, and the number of sets $(p; i_1, i_2, \dots, i_n)$, such that one of the multipliers in the left hand side of (2) equals l , and other $n - 1$ multipliers equal m is equal to $n \cdot \alpha_l \cdot \alpha_m^{n-1}$ (if $z_a + 1 \neq l$ then $p = n$). Therefore α_l can be found from the equality $A + n \cdot \alpha_l \cdot \alpha_m^{n-1} = C$.

Analogously we may find the type $(\beta_1, \beta_2, \dots, \beta_n)$ of b via the cardinalities of the equivalence classes of the relation \sim_b . Since for isomorphic semigroups $(\mathcal{PT}_n, *_a)$ and $(\mathcal{PT}_n, *_b)$ the number of the equivalence classes of the same cardinality coincide, and the values α_k and β_k , $k = 1, \dots, n$ are defined uniquely by the number of equivalence classes, for all k we have $\alpha_k = \beta_k$, that is, elements a and b have the same types.

Sufficiency. Let elements a and b have the type $(\alpha_1, \alpha_2, \dots, \alpha_n)$. Then there exist permutations π and τ in \mathcal{S}_n such that $b = \tau \alpha \pi$. The map $f : (\mathcal{PT}_n, *_a) \rightarrow (\mathcal{PT}_n, *_b)$ such that $f(x) = \pi^{-1} x \tau^{-1}$ defines isomorphism between $(\mathcal{PT}_n, *_a)$ and $(\mathcal{PT}_n, *_b)$. In fact f is

bijjective and

$$\begin{aligned} f(x *_a y) &= \pi^{-1} x *_a y \tau^{-1} = \pi^{-1} x \tau^{-1} \tau a \pi \pi^{-1} y \tau^{-1} = \\ &= \pi^{-1} x \tau^{-1} b \pi^{-1} y \tau^{-1} = f(x) *_b f(y). \end{aligned}$$

□

Corollary 1. Let $p(k)$ denote the number of ways in which one can split positive integer k into non ordered sum of the natural integers. Then there are $\sum_{k=0}^n p(k)$ pairwise non-isomorphic semigroups obtained from \mathcal{PT}_n by the deformed multiplication.

References

- [1] Ljapin Y.S., Semigroups, Moscow, Fizmatgiz, 1960 (Russian).
- [2] Magill Kenneth D., Semigroup structures for families of functions. II. Continuous functions. // J. Austral. Math. Soc. 7 (1967), 95-107.
- [3] Sullivan R.P., Generalized partial transformation semigroups. // J. Austral. Math. Soc. 19 (1975), part 4, 470-473
- [4] Symons J.S.V., On a generalization of the transformation semigroup. // J. Austral. Math. Soc. 19 (1975), 47-61
- [5] Artamonov V.A, Salij V.N., Skorniyakov L.A. and others, General Algebra, Moscow, Nauka, 1991, vol. 1 (Russian).

Department of Mechanics and Mathematics,
Kiev Taras Shevchenko University,
64, Volodymyrska st., 01033, Kiev, Ukraine,
e-mail: *gtsyaputa@univ.kiev.ua*

Given to the editorial board 30.09.2003