

Hochschild cohomology of truncated quiver algebras ^{*}

Yunge Xu ^a, Yang Han ^b and Wenfeng Jiang ^a

a. Faculty of Mathematics & Computer Science, Hubei University,
Wuhan 430062, P.R.China E-mail: xuy@hubu.edu.cn

b. Academy of Mathematics and Systems science, Chinese Academy of Sciences,
Beijing 100080, P.R. China E-mail: hany@iss.ac.cn

Abstract: For a truncated quiver algebra over a field of arbitrary characteristic, its Hochschild cohomology is calculated. Moreover, it is shown that its Hochschild cohomology algebra is finite-dimensional if and only if its global dimension is finite if and only if its quiver has no oriented cycles.

MSC(2000): 16E40, 16E10, 16G10

Introduction

Let k be a fixed field of arbitrary characteristic and A a finite-dimensional k -algebra. Let $A^e := A^{op} \otimes_k A$ be the *enveloping algebra* of A . The n -th *Hochschild cohomology group* of A is $H^n(A) := \text{Ext}_{A^e}^n(A, A)$ and the *Hochschild cohomology algebra* of A is $H^\bullet(A) := \prod_{n \geq 0} H^n(A)$ where the multiplication is induced by the Yoneda product for extensions [1].

Let $Q = (Q_0, Q_1)$ be a finite connected quiver where Q_0 (resp. Q_1) is the set of vertices (resp. arrows) in Q . Given a path p in Q , we denote by $o(p)$ and $t(p)$ the origin and the terminus of p respectively, and by $l(p)$ the length of p . For $0 \leq j \leq l(p)$, denote by ${}^j p$ the subpath of p with length j and origin $o(p)$. A path $a_1 \cdots a_n$ with $n \geq 1$ is called an *oriented cycle* of length n if $t(a_i) = o(a_{i+1})$ for $i = 1, \dots, n-1$ and $t(a_n) = o(a_1)$. An oriented cycle is called a *basic cycle* if it has no self intersection.

A *truncated quiver algebra* is an algebra kQ/k^nQ where $n \geq 2$ and k^nQ denotes the ideal of kQ generated by all paths of length $\geq n$ (cf. [2]). It is

^{*}Projects 10426014 and 10201004 supported by NSFC

further called a *truncated basic cycle algebra* if Q is a basic cycle. We refer to [3] for the theory of quivers and their representations.

Though Hochschild cohomology is theoretically computable for a concrete algebra via derived functors, actual calculation for a class of algebras is still very convenient, important and difficult. So far the Hochschild cohomology was calculated for hereditary algebras [4, 5], incidence algebras [6, 7], algebras with narrow quivers [4, 5], and so on. For the Hochschild cohomology of truncated quiver algebras partial results are known: Cibils calculated the Hochschild cohomology of radical square zero algebras [8]. In case the field k is of characteristic zero, Zhang and Locateli calculated the Hochschild cohomology of truncated basic cycle algebras and truncated quiver algebras respectively [9, 10, 11]. For truncated basic cycle algebras over a field k of arbitrary characteristic, Bardzell, Locateli and Marcos calculated their Hochschild cohomology [12]. In this paper, we complete the calculation of the Hochschild cohomology of truncated quiver algebras over a field of arbitrary characteristic. We shall employ Locateli's strategy and do more and deeper analysis.

In 1989, Happel asked the following question: Whether for a finite dimensional algebra A over an algebraically closed field k , $\text{gl.dim.}A < \infty$ if and only if the Hochschild cohomology algebra of A is finite-dimensional [5]? Though Happel's question has negative answer [13], it does have some positive answers: Avramov and Iyengar's result implies that Happel's question has positive answer for commutative algebras [14]. Cibils' result implies that Happel's question has positive answer for radical square zero algebras [8]. Moreover, Locateli proved that, in case the underlying field k is of characteristic 0, the Hochschild cohomology algebra of a truncated quiver algebra is finite-dimensional if and only if its quiver has no oriented cycle. In this paper, we show by constructing explicitly some Hochschild cocycles that for a truncated quiver algebra over a field of arbitrary characteristic, its Hochschild cohomology algebra is finite-dimensional if and only if its global dimension is finite if and only if its quiver has no oriented cycles.

1 Hochschild cohomology groups

Let $A = kQ/k^nQ$, $n \geq 2$, be a truncated quiver algebra. If S and T are sets of paths in Q we denote by $(S//T)$ the set $\{(p, q) \in S \times T \mid o(p) = o(q) \text{ and } t(p) = t(q)\}$. If S (resp. T) is the set of paths of length i (resp. j) then we denote $(S//T)$ by $(i//j)$. Given a set S , kS denotes the k -vector space with basis S .

If Q is a basic cycle, Bardzell, Locateli and Marcos calculated the Hochschild cohomology of A :

Theorem 1. ([12, p. 1619]) *Let $A = kQ/k^nQ$ with Q a basic cycle of length e and $n = me + r$, $0 \leq r \leq e - 1$. Then we have*

$$\dim_k H^0(A) = \begin{cases} m, & \text{if } r = 0; \\ m + e, & \text{if } r = 1; \\ m + 1, & \text{if } r \geq 2. \end{cases}$$

$$\dim_k H^1(A) = \begin{cases} m, & \text{if } r = 0, 1; \\ m + 1, & \text{if } r \geq 2. \end{cases}$$

and for $i \geq 1$,

$$\dim_k H^{2i}(A) = \dim_k H^{2i+1}(A) = \begin{cases} c_{n,e,i}, & \text{if } \text{char} k \nmid n \text{ or } e \nmid (ni - n + 1); \\ c_{n,e,i} + 1, & \text{if } \text{char} k | n \text{ and } e | (ni - n + 1). \end{cases}$$

where $c_{n,e,i} := |\{j | 0 \leq j \leq n - 2, j \equiv ri(\text{mode})\}|$.

Thanks to Theorem 1, it remains to consider the case that Q is not a basic cycle. It follows from [11, Lemma 2 and Lemma 6] that for all $i \geq 0$, the i -th Hochschild cohomology group of A is just the i -th cohomology group of the cochain complex

$$(M^\bullet, d^\bullet) : 0 \rightarrow M^0 \xrightarrow{d^1} M^1 \xrightarrow{d^2} \dots \xrightarrow{d^{i-1}} M^{i-1} \xrightarrow{d^i} M^i \xrightarrow{d^{i+1}} M^{i+1} \xrightarrow{d^{i+2}} \dots$$

where

$$\begin{aligned} M^{2i} &= k(0//ni) \oplus \dots \oplus k(n-1//ni), \\ M^{2i+1} &= k(0//ni+1) \oplus \dots \oplus k(n-1//ni+1), \\ d^{2i+1} &= \begin{bmatrix} 0 & & & & & \\ d_0^{2i+1} & 0 & & & & \\ & d_1^{2i+1} & \dots & & & \\ & & \ddots & & & \\ & & & \ddots & & \\ & & & & 0 & \\ & & & & d_{n-2}^{2i+1} & 0 \end{bmatrix}, \\ d^{2i} &= \begin{bmatrix} 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 \\ d_0^{2i} & 0 & \dots & 0 \end{bmatrix} \end{aligned}$$

with

$$d_j^{2i+1} : k(j//ni) \rightarrow k(j+1//ni+1), (p, q) \mapsto \sum_{a \in Q_1} (ap, aq) - \sum_{a \in Q_1} (pa, qa)$$

for $0 \leq j \leq n - 2$ and

$$d_0^{2i} : k(0//n(i-1)+1) \rightarrow k(n-1//ni), (o(p), p) \mapsto \sum_{l(sq)=n-1} (sq, spq).$$

Therefore we have

$$\text{Ker } d^{2i+1} = \left(\bigoplus_{j=0}^{n-2} \text{Ker } d_j^{2i+1} \right) \oplus k(n-1//ni), \quad \text{Im } d^{2i+1} = \bigoplus_{j=0}^{n-2} \text{Im } d_j^{2i+1},$$

$$\text{Ker } d^{2i} = \text{Ker } d_0^{2i} \oplus \bigoplus_{j=1}^{n-1} k(j//n(i-1)+1), \quad \text{Im } d^{2i} = \text{Im } d_0^{2i},$$

$$H^{2i+1}(A) = \text{Ker } d_0^{2i+2} \oplus \bigoplus_{j=1}^{n-1} (k(j//ni+1)/\text{Im } d_{j-1}^{2i+1})$$

and

$$H^{2i}(A) = (k(n-1//ni)/\text{Im } d_0^{2i}) \oplus \bigoplus_{j=0}^{n-2} \text{Ker } d_j^{2i+1}.$$

Now we recall some notations introduced by Cibils in [2]: For $1 \leq j \leq n - 2$ and $(p, q) \in (j//ni)$. We say that (p, q) *starts together* (resp. *ends together*) if the first (resp. last) arrow of p and q coincide. And we say that (p, q) *starts at a source* (resp. *ends at a sink*) if $o(p) = o(q)$ is a source (resp. if $t(p) = t(q)$ is a sink).

Suppose that (p, q) starts together, so $(p, q) = (ap', aq')$ for some $a \in Q_1$, and moreover suppose that (p, q) does not end at a sink. A *+movement* of (p, q) is a couple $(p'b, q'b)$ where $b \in t(p)Q_1$. If (p, q) does not start together, or if (p, q) end at a sink, then no *+movement* of (p, q) is defined, and we call (p, q) a *+extreme*.

In analogous way, if (p, q) ends together and (p, q) does not start at a source, then we define a *-movement* of (p, q) . If (p, q) does not end together, or starts at a source, we say (p, q) is a *-extreme*.

Consider the equivalence relation on $(j//ni)$ by declaring (p, q) *equivalent* to (p', q') if there exists a finite sequence of *+* or *-* movements transforming (p, q) into (p', q') . We call an equivalence class a *j-extreme* if all its *+extreme* elements end at a sink and all its *-extreme* elements start at a source.

For notational convenience we denote $\{0\text{-extremes}\} = \emptyset$ and $\{(n-1)\text{-extremes}\} = (n-1//ni)$. One can check Locateli's proofs step by step and obtain that the following proposition and two lemmas hold for the underlying field k of arbitrary characteristic.

Proposition 1. ([11, Proposition 3]) *If Q is not a basic cycle, then*

$$\dim_k \text{Ker } d^{2i+1} = \sum_{j=1}^{n-1} |\{j\text{-extremes}\}|$$

and

$$\dim_k \text{Im } d^{2i+1} = \sum_{j=0}^{n-2} (|(j//ni)| - |\{j\text{-extremes}\}|).$$

Let $x = \sum_{(p,q) \in (i//j)} x_{(p,q)}(p, q) \in k(i//j)$. The *support* of x is the set $\text{Supp}(x) = \{(p, q) | x_{(p,q)} \neq 0\}$.

Lemma 1. ([11, Lemma 7]) (Case $i = 1$.) *Let $(o(p), p) \in (0//1)$ and $(sq, spq) \in \text{Supp}(d_0^{2i}(o(p), p))$ with $l(sq) = n - 1$. Then (sq, spq) is not in the support of $d_0^{2i}(o(p'), p')$ for any other $p' \neq p$.*

Let p and q be two paths in Q . We denote by $p \cap q$ the set of the vertices that belong to both p and q . Given a cycle $p = a_1 \cdots a_j \cdots a_{n(i-1)+1}$ where $i \geq 2$ and $a_j \in Q_1$ for $1 \leq j \leq n(i-1) + 1$, the cycle $p \langle a_1 \cdots a_j \rangle := a_{j+1} \cdots a_{n(i-1)+1} a_1 \cdots a_j$ is called the *rotate* of p with origin $o(a_{j+1})$.

Lemma 2. ([11, Lemma 8]) (Case $i \geq 2$.) *Let $(o(p), p) \in (0//n(i-1) + 1)$ with $i \geq 2$ and $(sq, spq) \in \text{Supp}(d_0^{2i}(o(p), p))$ with $l(sq) = n - 1$. Let $(o(p'), p') \in (0//n(i-1) + 1)$ and $(o(p), p) \neq (o(p'), p')$. Then $(sq, spq) \in \text{Supp}(d_0^{2i}(o(p'), p'))$ if and only if $|sq \cap p| \geq 2$ and p' is a rotate of p with origin in some vertex in $sq \cap p$.*

Proposition 2. *If Q is not a basic cycle then the map $d_0^{2i} : k(0//n(i-1) + 1) \rightarrow k(n-1//ni)$ is injective for all $i \geq 1$.*

Proof. Case $i = 1$: Let $x = \sum_{(o(p), p) \in (0//1)} x_{(o(p), p)}(o(p), p) \in \text{Ker } d_0^2$ where $x_{(o(p), p)} \in k$. For any $(o(p), p) \in (0//1)$ and $(sq, spq) \in \text{Supp}(d_0^2(o(p), p))$, by Lemma 1, we have that (sq, spq) is not in the support of $d_0^2(o(p'), p')$ for any $p' \neq p$. Since $d_0^2(x) = 0$, we must have $x_{(o(p), p)} = 0$. So $x = 0$ and d_0^2 is injective.

Case $i \geq 2$: Define an equivalence relation on $(0//n(i-1) + 1)$ by saying that p is *equivalent* to q if p is a rotate of q . Denote by \mathcal{C} the set of equivalence classes. We can define a complete order in the set $(0//n(i-1) + 1)$ as follows: First give a complete order $<'$ to the set of classes \mathcal{C} . Next, for each $c \in \mathcal{C}$, choose $p = a_1 \cdots a_{n(i-1)+1} \in c$ and order the elements of c by the complete order $<''$ given by $p_1 := p <'' p_2 := p \langle a_1 \rangle <'' p_3 := p \langle a_1 a_2 \rangle <'' \cdots <'' p_{n(i-1)+1} := p \langle a_1 \cdots a_{n(i-1)} \rangle$. Then consider in $(0//n(i-1) + 1)$ the

cycle, $B_{\bar{p}} = (2)$ and $N_{\bar{p}}$ is in the form

$$\begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix}.$$

Hence

$$\text{rank} \begin{bmatrix} B_{\bar{p}} \\ N_{\bar{p}} \end{bmatrix} = 1 = e.$$

If $e > 2$, the rank of the matrix

$$B_{\bar{p}} = \begin{bmatrix} 1 & 1 & & & \\ & 1 & 1 & & \\ & & 1 & \cdots & \\ & & & \ddots & 1 \\ 1 & & & & 1 \end{bmatrix}_{e \times e}$$

is

$$\text{rank } B_{\bar{p}} = \begin{cases} e, & \text{if char } k \neq 2; \\ e - 1, & \text{if char } k = 2. \end{cases}$$

Since $\text{char } k \neq 2$, we have $\text{rank } B_{\bar{p}} = e$.

Each block $B_{\bar{p}}$, $\bar{p} \in \mathcal{C}$, is of full rank, so the map d_0^{2i} is injective.

(ii). $n > 2$ and $\text{char } k \nmid n$: We write $n = me + r$, with $0 \leq r < e$. It is not difficult to see that $({}^{n-1}q, q \cdot {}^{n-1}q)$ in $\mathcal{B}_{\bar{p}}$ appears in $d_0^{2i}(o(p), p)$ with coefficient $m + 1$ if $q = p_j$ with $1 \leq j \leq r$, or m if $q = p_j$ with $j > r$. Thus each matrix $B_{\bar{p}}$ is of the form

$$B_{\bar{p}} = \begin{bmatrix} m+1 & m & m & \cdots & m+1 \\ m+1 & m+1 & m & \cdots & m+1 \\ m+1 & m+1 & m+1 & \cdots & m+1 \\ \vdots & \vdots & \vdots & & \vdots \\ m+1 & m+1 & m+1 & \cdots & m+1 \\ m+1 & m+1 & m+1 & \cdots & m \\ m & m+1 & m+1 & \cdots & m \\ m & m & m+1 & \cdots & m \\ \vdots & \vdots & \vdots & & \vdots \\ m & m & m & \cdots & m \\ m & m & m & \cdots & m+1 \end{bmatrix}_{e \times e}$$

which is a cyclic matrix, where in each column we have r entries $(m+1)$ and $(e-r)$ entries m .

Note that if $(o(p), p) \in (0//n(i-1)+1)$ then one must have $ni \equiv n-1 \pmod{l(p)}$. So $\gcd(n, l(p)) = 1$ and thus $\gcd(e, r) = 1$. By [12, p. 1619], we have

$$\text{rank } B_{\bar{p}} = \begin{cases} e, & \text{if char } k \nmid n; \\ e-1, & \text{if char } k \mid n. \end{cases}$$

Since $\text{char } k \nmid n$, we have $\text{rank } B_{\bar{p}} = e$.

Each block $B_{\bar{p}}, \bar{p} \in \mathcal{C}$, is of full rank, so the map d_0^{2i} is injective.

(iii). $n > 2$ and $\text{char } k \mid n$, or $n = 2$ and $\text{char } k = 2$: Since Q is connected but not a basic cycle, without loss of generality, we may assume that Q contains an arrow b which does not appear in w and $t(b) = o(p)$. Of course it is possible that $o(b)$ belongs to the vertices of w . Moreover, we assume $w = a_1 a_2 \cdots a_e$ and $p = w^h$ for some $h \geq 1$ and $n = me + r$, with $0 \leq r < e$. Clearly, $(b \cdot {}^{n-2}p, bp \cdot {}^{n-2}p) \in \mathcal{N}_{\bar{p}}$ and ${}^{n-2}p = w^m \cdot {}^{r-2}p$.

(1). If $m = 0$ then $n - 2 = r - 2$, thus $n - 2 < e - 2$. Hence

$$\begin{aligned} (b \cdot {}^{n-2}p, bp \cdot {}^{n-2}p) &= (b \cdot a_1 \cdots a_{n-2}, b(a_1 \cdots a_e)a_1 \cdots a_{n-2}) \\ &= (ba_1 \cdot a_2 \cdots a_{n-2}, ba_1(a_2 \cdots a_e a_1)a_2 \cdots a_{n-2}) \\ &= \cdots \\ &= (ba_1 \cdots a_{n-2}, ba_1 \cdots a_{n-2}(a_{n-1} \cdots a_e a_1 \cdots a_{n-2})). \end{aligned}$$

So $(b \cdot {}^{n-2}p, bp \cdot {}^{n-2}p)$ appears only in $d_0^{2i}(o(p'), p')$ with coefficient 1 for $p' = p_j$ with $1 \leq j \leq n-1$, and with coefficient 0 otherwise. Thus the row corresponding to $(b \cdot {}^{n-2}p, bp \cdot {}^{n-2}p)$ in the matrix $N_{\bar{p}}$ is of the form

$$\underbrace{(1, 1, \dots, 1)}_{n-1}, 0, 0, \dots, 0)$$

whose row-sum is $n-1 \neq 0$. Therefore

$$\text{rank} \begin{bmatrix} B_{\bar{p}} \\ N_{\bar{p}} \end{bmatrix} = e.$$

(2). If $m \geq 1$ and $r \geq 2$ then $(b \cdot {}^{n-2}p, bp \cdot {}^{n-2}p)$ appears in $d_0^{2i}(o(p'), p')$ for $p' = p_j$ with coefficient $m+1$ if $1 \leq j < r$, or m if $r \leq j \leq e$. So the row of the matrix $N_{\bar{p}}$ corresponding to $(b \cdot {}^{n-2}p, bp \cdot {}^{n-2}p)$ is of the form

$$\underbrace{(m+1, m+1, \dots, m+1)}_{r-1}, \underbrace{(m, \dots, m)}_{e-r+1}$$

whose row-sum is $n - 1 \neq 0$. Thus

$$\text{rank} \begin{bmatrix} B_{\bar{p}} \\ N_{\bar{p}} \end{bmatrix} = e.$$

(3). If $m \geq 1$ and $r = 1$ then ${}^{n-2}p = w^{m-1}a_1a_2 \cdots a_{e-1}$. So the coefficient of $(b \cdot {}^{n-2}p, bp \cdot {}^{n-2}p)$ in $d_0^{2i}(o(p'), p')$ for $p' = p_j$ is m if $1 \leq j \leq e$. Thus the row corresponding to $(b \cdot {}^{n-2}p, bp \cdot {}^{n-2}p)$ in $N_{\bar{p}}$ is of the form

$$\underbrace{(m, m, \dots, m)}_e$$

whose row-sum is also $(n - 1)$ and hence nonzero. So

$$\text{rank} \begin{bmatrix} B_{\bar{p}} \\ N_{\bar{p}} \end{bmatrix} = e.$$

(4). If $m \geq 1$ and $r = 0$ then ${}^{n-2}p = w^{m-1}a_1a_2 \cdots a_{e-2}$. The row corresponding to $(b \cdot {}^{n-2}p, bp \cdot {}^{n-2}p)$ in $N_{\bar{p}}$ is of the form

$$\underbrace{(m, m, \dots, m, m - 1)}_{e-1}$$

whose row-sum is $n - 1 \neq 0$. Therefore

$$\text{rank} \begin{bmatrix} B_{\bar{p}} \\ N_{\bar{p}} \end{bmatrix} = e.$$

Anyway, each block $\begin{bmatrix} B_{\bar{p}} \\ N_{\bar{p}} \end{bmatrix}$, $\bar{p} \in \mathcal{C}$, is of full rank, so the map d_0^{2i} is injective. \square

By Proposition 2, we immediately have:

Proposition 3. *If Q is not a basic cycle then, for $i \geq 1$, we have*

$$\dim_k \text{Ker } d^{2i} = \sum_{j=1}^{n-1} |(j // n(i - 1) + 1)|$$

and

$$\dim_k \text{Im } d^{2i} = |(0 // n(i - 1) + 1)|.$$

Now we can complete the calculation of the Hochschild cohomology of truncated quiver algebras over a field of arbitrary characteristic as follows:

Theorem 2. *Let $A = kQ/k^nQ$ be a truncated algebra. If Q is not a basic cycle quiver then we have*

$$H^0(A) = Z(A),$$

$$\dim_k H^1(A) = \dim_k Z(A) - \sum_{j=0}^{n-1} |(j//0)| + \sum_{j=1}^{n-1} |(j//1)|$$

and for $i \geq 1$

$$\dim_k H^{2i+1}(A) = \sum_{j=1}^{n-1} |(j//ni + 1)| - \sum_{j=0}^{n-2} (|(j//ni)| - |\{j\text{-extremes}\}|)$$

$$\dim_k H^{2i}(A) = \sum_{j=1}^{n-1} |\{j\text{-extremes}\}| - |(0//n(i-1) + 1)|.$$

where $Z(A)$ denotes the center of the algebra A .

Proof. It is well-known that $H^0(A) = Z(A)$. Since $H^0(A) = \text{Ker } d^1$, we have

$$\dim_k \text{Im } d^1 = \dim_k M^0 - \dim_k \text{Ker } d^1 = \sum_{j=0}^{n-1} |(j//0)| - \dim_k Z(A).$$

By Proposition 3, we have

$$\dim_k H^1(A) = \dim_k \text{Ker } d^2 - \dim_k \text{Im } d^1 = \sum_{j=1}^{n-1} |(j//1)| - \sum_{j=0}^{n-1} |(j//0)| + \dim_k Z(A).$$

If $i \geq 1$ then by Proposition 1 and Proposition 3 we have

$$\begin{aligned} \dim_k H^{2i+1}(A) &= \dim_k \text{Ker } d^{2(i+1)} - \dim_k \text{Im } d^{2i+1} \\ &= \sum_{j=1}^{n-1} |(j//ni + 1)| - \sum_{j=0}^{n-2} (|(j//ni)| - |\{j\text{-extremes}\}|) \end{aligned}$$

and

$$\begin{aligned} \dim_k H^{2i}(A) &= \dim_k \text{Ker } d^{2i+1} - \dim_k \text{Im } d^{2i} \\ &= \sum_{j=1}^{n-1} |\{j\text{-extremes}\}| - |(0//n(i-1) + 1)|. \end{aligned}$$

□

2 Hochschild cohomolgy algebras.

The main result of this section is the following:

Theorem 3. *Let $A = kQ/k^nQ$ be a truncated quiver algebra. Then the following conditions are equivalent:*

- (1) *The algebra $H^*(A)$ is finite-dimensional;*
- (2) *$\text{gl.dim.}A < \infty$;*
- (3) *Q has no oriented cycles.*

Proof. (3) \Rightarrow (2) \Rightarrow (1): Clearly.

(1) \Rightarrow (3): If Q has an oriented cycle then we take the shortest oriented cycle $w = a_1a_2 \cdots a_e$ in Q . Thus the vertices of w must be different from each other.

Now we show that $(a_e, a_e(a_1 \cdots a_e)^{nr}) \in k(1//nre + 1) \setminus \text{Im } d_0^{2re+1}$, thus d_0^{2re+1} is not surjective, hence $H^{2re+1}(A) \neq 0$, for all $r \geq 0$:

Assume on the contrary $(a_e, a_e(a_1 \cdots a_e)^{nr}) \in \text{Im } d_0^{2re+1}$. Then there is $\sum_{i=1}^e x_i(v_i, (a_i \cdots a_e a_1 \cdots a_{i-1})^{nr}) + \sum_{j=1}^s y_j(u_j, b_{j1} \cdots b_{j(nr)}) \in k(0//nre)$ with

$$\begin{aligned}
& \{b_{j1}, \dots, b_{j(nr)}\} \not\subseteq \{a_1, \dots, a_e\} \text{ such that } (a_e, a_e(a_1 \cdots a_e)^{nr}) \\
&= d_0^{2re+1} \left(\sum_{i=1}^e x_i(v_i, (a_i \cdots a_e a_1 \cdots a_{i-1})^{nr}) + \sum_{j=1}^s y_j(u_j, b_{j1} \cdots b_{j(nr)}) \right) \\
&= \sum_{i=1}^e x_i \sum_{a \in Q_1} ((av_i, a(a_i \cdots a_e a_1 \cdots a_{i-1})^{nr}) - (v_i a, (a_i \cdots a_e a_1 \cdots a_{i-1})^{nr} a)) \\
&\quad + \sum_{j=1}^s y_j \sum_{a \in Q_1} ((au_j, ab_{j1} \cdots b_{j(nr)}) - (u_j a, b_{j1} \cdots b_{j(nr)} a)) \\
&= \sum_{i=1}^e x_{i+1} (a_i, a_i(a_{i+1} \cdots a_e a_1 \cdots a_i)^{nr}) \\
&\quad - \sum_{i=1}^e x_i (a_i, (a_i \cdots a_e a_1 \cdots a_{i-1})^{nr} a_i) \\
&\quad + \sum_{i=1}^e x_i \sum_{a \in Q_1 \setminus \{a_{i-1}\}} (av_i, a(a_i \cdots a_e a_1 \cdots a_{i-1})^{nr}) \\
&\quad - \sum_{i=1}^e x_i \sum_{a \in Q_1 \setminus \{a_i\}} (v_i a, (a_i \cdots a_e a_1 \cdots a_{i-1})^{nr} a) \\
&\quad + \sum_{j=1}^s y_j \sum_{a \in Q_1} ((au_j, ab_{j1} \cdots b_{j(nr)}) - (u_j a, b_{j1} \cdots b_{j(nr)} a))
\end{aligned}$$

where $x_{e+1} := x_1$.

Let W be the k -subspace of $k(1//nre + 1)$ generated by e elements $(a_i, a_i(a_{i+1} \cdots a_e a_1 \cdots a_i)^{nr})$, $1 \leq i \leq e$. Then $k(1//nre + 1) = W \oplus \overline{W}$ where \overline{W} is the complement of W in $k(1//nre + 1)$. According to this decomposition we have $(a_e, a_e(a_1 \cdots a_e)^{nr}) = \sum_{i=1}^e x_{i+1} (a_i, a_i(a_{i+1} \cdots a_e a_1 \cdots a_i)^{nr}) -$

$\sum_{i=1}^e x_i(a_i, (a_i \cdots a_e a_1 \cdots a_{i-1})^{nr} a_i)$. Comparing the coefficients on two sides we obtain

$$\begin{aligned} 1 &= x_1 - x_e \\ 0 &= x_2 - x_1 \\ 0 &= x_3 - x_2 \\ \dots &\dots \dots \\ 0 &= x_e - x_{e-1}. \end{aligned}$$

It is a contradiction. □

References

- [1] Mac Lane S., Homology, Grundlehren 114, Third corrected printing, Springer-Verlag, 1975.
- [2] Cibils C., Rigidity of truncated quiver algebras, Adv. Math., 1990, 79: 18-42.
- [3] Auslander M., Reiten I. and SmalØ S.O., Representation theory of artin algebras, Cambridge studies in advanced mathematics 36, Cambridge university press, Cambridge, 1995.
- [4] Cibils C., On the Hochschild cohomology of finite-dimensional algebras, Comm. Algebra, 1988, 16: 645-649.
- [5] Happel D., Hochschild cohomology of finite-dimensional algebras, Springer Lecture Notes in Math. 1404, 1989: 108-126.
- [6] Cibils C., Cohomology of incidence algebras and simplicial complexes, J. Pure Appl. Algebra, 1989, 56: 221-232.
- [7] Gerstenhaber M. and Schack S.P., Simplicial homology is Hochschild cohomology, J. Pure Appl. Algebra, 1983, 30: 143-156.
- [8] Cibils C., Hochschild cohomology algebra of radical square zero algebras, CMS Conf. Proc. 1998, 24: 93-101.
- [9] Zhang P., Hochschild cohomology of truncated algebras, Sci. China, Ser. A (in Chinese), 1994, 24: 1121-1125.
- [10] Zhang P., Hochschild cohomology of truncated basic cycle, Sci. China, Ser. A, 1997, 40: 1272-1278.
- [11] Locateli A.C., Hochschild cohomology of truncated quiver algebras, Comm. Algebra, 1999, 27: 645-664.
- [12] Bardzell M.J., Locateli A.C. and Marcos E.N., On the Hochschild cohomology of truncated cycle algebras, Comm. Algebra, 2000, 28: 1615-1639.
- [13] Buchweitz R.O., Green E.L., Madsen D. and Solberg Ø., Finite Hochschild cohomology without finite global dimension, Math. Res. Letters (in press).
- [14] Avramov L.L. and Iyengar S., Gaps in Hochschild cohomology imply smoothness for commutative algebras, Math. Res. Letters (in press).