# Murre's conjectures for certain product varieties

Kenichiro Kimura

#### Abstract

We consider Murre's conjectures on Chow groups for a fourfold which is a product of two curves and a surface. We give a result which concerns Conjecture D: the kernel of a certain projector is equal to the homologically trivial part of the Chow group. We also give a proof of Conjecture B for a product of two surfaces. MSC number: 14C25

#### Introduction 1

Let X be a smooth projective variety over  $\mathbb{C}$  of dimension d. Let  $\Delta \subset X \times X$  be the diagonal. There is a cohomology class  $cl(\Delta) \in H^{2d}(X \times X)$ .

In this paper we use Betti cohomology with rational coefficients. There is the Künneth decomposition

 $H^{2d}(X \times X) \simeq \bigoplus_{i=0}^{2d} H^{2d-i}(X) \otimes H^i(X).$ We write  $cl(\Delta) = \sum_{i=0}^{2d} \pi_i^{hom}$  according to this decomposition. Here  $\pi_i^{hom} \in H^{2d-i}(X) \otimes H^i(X)$ .  $H^{i}(X)$ . If the Künneth conjecture is true, then each  $\pi_{i}^{hom}$  is an algebraic cycle. Murre([Mu],[Mu2]) formulated the following conjecture. For an abelian group M, we

- write  $M_{\mathbb{Q}} = M \otimes \mathbb{Q}$ .
- (A) The  $\pi_i^{hom}$  lift to a set of orthogonal projectors  $\pi_i$  in  $CH^d(X \times X)_{\mathbb{Q}}$  which satisfy the equality

$$\sum_{i=0}^{2d} \pi_i = \Delta.$$

- (B) The correspondences  $\pi_0, \dots, \pi_{j-1}, \pi_{2j+1}, \dots, \pi_{2d}$  act as zero on  $CH^j(X)_{\mathbb{Q}}$ .
- (C) Let  $F^{\nu}CH^{j}(X) = Ker\pi_{2j} \cap Ker\pi_{2j-1} \cdots \cap Ker\pi_{2j-\nu+1}$ . Then the filtration  $F^{\cdot}$  is independent of the choice of  $\pi_i$ .

(D) 
$$F^1CH^j(X)_{\mathbb{Q}} = CH^j(X)_{hom,\mathbb{Q}}$$

It is shown by Jannsen(Ja) that this conjecture of Murre is equivalent to Beilinson's conjectures on the filtration on Chow groups.

There are not yet many evidences for this conjecture. For a projective smooth curve Cand a closed point p on C, set  $\pi_0 = p \times C$ ,  $\pi_2 = C \times p$  and  $\pi_1 = \Delta - \pi_0 - \pi_2$ . Then Conjectures (A), (B) and (D) are true for these projectors. For a projective smooth surface Murre([Mu]) constructed a set of projectors  $\pi_0, \dots, \pi_4$  for which Conjectures (A), (B) and (D) are true. About Conjecture (C) he proved that the filtration on Chow groups given by these projectors is a natural one in the following sense(Theorem 3 in [Mu]):

- $F^1(CH^1(S)_{\mathbb{Q}}) = Ker(\pi_2) = Pic^0(S)_{\mathbb{Q}}.$
- $F^1(CH^2(S)_{\mathbb{Q}}) = CH^2(S)_{hom,\mathbb{Q}}$ .  $F^2(CH^2(S)_{\mathbb{Q}}) = Ker(\pi_3) = Ker(alb:CH^2(S)_{hom,\mathbb{Q}} \rightarrow Alb(S)_{\mathbb{Q}})$ .

Conjecture (A) is also true for abelian varieties (Shermenev [Sh], Deninger-Murre [DM]), hypersurfaces (easy), certain class of threefolds (del Angel-Müller-Stach [deM],[deM2]), and some modular varieties (Gordon-Murre [GM], Gordon-Hanamura-Murre [GHM], [GHM2], Miller-Müller-Stach-Wortmann-Yang-Zuo [Pic]).

Note that if Conjecture (A) is true for varieties X and Y, then it is also true for  $X \times Y$ . One can put  $\pi_{iX \times Y} = \sum_{p+q=i} \pi_{p_X} \times \pi_{q_Y}$ .

In [Mu2] Murre proves that Conjectures (B) and (D) are true for a product of a curve and a surface for this product Chow-Künneth decomposition.

Recently Murre([KMP]) proved the validity of Conjecture (B) and some part of Conjecture (D) for a product of two surfaces. More precisely, Murre proved that Conjecture (D) is true for a product  $S_1 \times S_2$  of two smooth projective surfaces except the following part:

The projector  $\pi_{2S_1} \times \pi_{2S_2}$  act as zero on  $CH^2(S_1 \times S_2)_{hom,\mathbb{Q}}$ .

If this is true for the case of a self-product  $S_1 = S_2$  of a surface, then Bloch's conjecture  $(p_g = 0 \Rightarrow \text{albanese map is injective})$  for  $S_1$  is true. If one assumes that the Chow group of  $S_1$  is finite dimensional in the sense of Kimura([Ki]), then for an element  $z \in CH^2(S_1 \times S_1)_{hom,\mathbb{Q}}$  one has the equality

$$(\pi_2 \times \pi_2(z))^n = 0$$

where <sup>n</sup> means the power as a correspondence and n is determined by the second Betti number of  $S_1$ .

In this paper we consider Conjecture (D) for the case where X is a product of two curves and a surface  $C_1 \times C_2 \times S$ . In this case the most crucial part is to show that  $\pi_{1C_1} \times \pi_{1C_2} \times \pi_{2S}$ act as zero on  $CH^2(X)_{hom,\mathbb{Q}}$ . Here the projectors  $\pi_{1C_i}$  for i = 1 and 2 are defined as above and we refer the reader to [Mu] for the definition of the projector  $\pi_{2S}$ . Our original aim was to show that if the cohomology  $H^1(C_1) \otimes H^1(C_2) \otimes H^2(S)$  has no non-zero Hodge cycle, then  $\pi_{1C_1} \times \pi_{1C_2} \times \pi_{2S}$  kills all the codimension 2 cycles on X. We could not completely solve the problem, so instead we studied what kind of cycles are killed by  $\pi_{1C_1} \times \pi_{1C_2} \times \pi_{2S}$ . It seems that under certain assumptions on X, "generic" cycles are killed by this projector (Theorem 2.1). This is the main result of this paper.

We also give a proof of the essential part of Conjecture (B) for a product of two surfaces. Our proof is similar to that of Murre in that we make essential use of the properties of the Chow-Künneth projectors for surfaces constructed by Murre. However there are still some differences so we decided to include our proof here.

This paper is organized as follows. In Section two we prove our main result about Conjecture (D). Section three is devoted to a proof of Conjecture (B) for a product of two surfaces.

The author expresses his deep gratitude to Jacob Murre for his patience in reading an earlier version of this paper and for valuable comments and encouragement.

## 2 The main result.

Let  $C_1$  and  $C_2$  be smooth projective curves over  $\mathbb{C}$ . Let S be a smooth projective surface over  $\mathbb{C}$ . We assume that these varieties are sufficiently general so that they satisfy the following conditions:

- $NS(S) \otimes \mathbb{Q} = \mathbb{Q}H$  where H is a hyperplane section of S.
- The cohomology groups  $H^1(C_1) \otimes H^1(C_2)$ ,  $H^1(C_2) \otimes H^1(S)$  and  $H^1(C_1) \otimes H^1(S)$  have no non-zero Hodge cycle.

Let  $X = C_1 \times C_2 \times S$ .

Let Z be a closed irreducible subvariety of X of codimension 2. Consider the following conditions for Z.

- 1.  $pr_{12}(Z) \subset C_1 \times C_2$  has dimension  $\leq 1$ .
- 2.  $pr_3(Z) \subset S$  has dimension  $\leq 1$ .
- 3.  $pr_{12}: Z \to C_1 \times C_2$  and  $pr_3: Z \to S$  are surjective and Z is a Cartier divisor either of  $C_1 \times pr_{23}(Z)$  or of  $C_2 \times pr_{13}(Z)$ .

Here we denote by  $pr_*$  the various projections from  $X = C_1 \times C_2 \times S$ . About the condition 3, if we assume that the projection  $pr_{23} : Z \to pr_{23}(Z)$  is flat, then one can show that Z is a Cartier divisor on  $C_1 \times pr_{23}(Z)$  (Lemma 2.2).

**Theorem 2.1.** Let  $C_1$  and  $C_2$  be a projective smooth curves over  $\mathbb{C}$  and let S be a projective smooth surface over  $\mathbb{C}$ .

Let  $X = C_1 \times C_2 \times S$ . Assume that these varieties satisfy the following conditions:

- $NS(S) \otimes \mathbb{Q} = \mathbb{Q}H$  where H is a hyperplane section of S.
- The cohomology groups  $H^1(C_1) \otimes H^1(C_2)$ ,  $H^1(C_2) \otimes H^1(S)$  and  $H^1(C_1) \otimes H^1(S)$ have no non-zero Hodge cycle.

Assume that the surface Z satisfies one of the conditions 1,2 and 3 above. Then the Chow-Künneth projector  $\pi_{1C_1} \times \pi_{1C_2} \times \pi_{2S}$  kills Z in  $CH^2(X)_{\mathbb{Q}}$ .

*Proof.* Assume that the condition 1 holds for Z. Note that we have a factorization

$$\pi_{1C_1} \times \pi_{1C_2} \times \pi_{2S} = (\pi_{1C_1} \times \pi_{1C_2} \times id_S) \circ (id_{C_1 \times C_2} \times \pi_{2S})$$

and they commute. We write  $C = pr_{12}(Z) \subset C_1 \times C_2$ . Let  $\eta_C \stackrel{j}{\hookrightarrow} C$  be the generic point of C. We apply the projector  $id_{C_1 \times C_2} \times (\pi_2)_S$  on Z as a cycle on  $C \times S$ . We have the equality

$$(j \times id_S)^*(id_C \times \pi_{2S})(Z) = (\eta_C \times \pi_{2S})((j \times id_S)^*Z).$$

We write  $(j \times id_S)^* Z = Z_\eta$ .

Since  $Z_{\eta}$  is algebraically equivalent to a cycle  $\eta_C \times E$  on the surface  $\eta_C \times S$  where E is a divisor on S defined over the base field  $\mathbb{C}$ , we see that

$$(\eta_C \times \pi_{2S}) Z_\eta = (\eta_C \times \pi_{2S}) (\eta_C \times E).$$

Here we use that  $\pi_{2S}(Pic^0(S)_{\mathbb{Q}}) = 0$ . By taking the closure of this equality in  $C \times S$ , it follows that

$$(id_C \times \pi_{2S})(Z) = C \times \pi_{2S}(E) + \sum_t p_t \times S$$

where for each  $t p_t$  is a closed point on C. Applying  $id_C \times \pi_{2S}$  on both sides of the equality kills  $p_t \times S$  because by Conjecture (B) for  $S \pi_{2S}(S) = 0$ .

Then we apply  $\pi_{1C_1} \times \pi_{1C_2} \times id_S$  on both sides of the equality. Since the cohomology  $H^1(C_1) \otimes H^1(C_2)$  has no non-zero Hodge cycle, it follows that

$$\pi_{1C_1} \times \pi_{1C_2}(C) \in Pic^0(C_1 \times C_2).$$

Since  $Pic^0(C_1 \times C_2) = pr_1^* Pic^0(C_1) + pr_2^* Pic^0(C_2)$  by applying  $\pi_{1C_1} \times \pi_{1C_2}$  again on  $\pi_{1C_1} \times \pi_{1C_2}(C)$  we see that it is zero because  $\pi_{1C_i} CH^0(C_i) = 0$  for i = 1 and 2.

The proof is similar if we assume that the condition 2 holds for Z.

Next we assume that the condition 3 holds for Z. Assume that Z is a Cartier divisor on  $C_1 \times pr_{23}(Z)$ .

**Lemma 2.1.** The subvariety  $pr_{23}(Z) \subset C_2 \times S$  is an ample divisor.

*Proof.* By the assumptions on  $C_2$  and S, we see that

 $NS(C_2 \times S) \otimes \mathbb{Q} = \mathbb{Q}(pt \times S) \oplus \mathbb{Q}(C_2 \times H).$ 

We denote  $D_1 = pt \times S$  and  $D_2 = C_2 \times H$ . Write  $aD_1 + bD_2$  for the class of  $pr_{23}(Z)$  in  $NS(C_2 \times S) \otimes \mathbb{Q}$ . We see that  $a = (C_2 \times pt, pr_{23}(Z)) > 0$  and  $b = \frac{(pt \times H, pr_{23}(Z))}{(H,H)} > 0$ . Here (\*, \*) denotes intersection number. So it follows that  $pr_{23}(Z) - aD_1 - bD_2 \in Pic^0(C_2 \times S) \simeq Pic^0(C_2) \oplus Pic^0(S)$ . So there are divisors  $d_1 \in Pic(C_2)$  and  $d_2 \in Pic(S)$  such that  $pr_{23}(Z) = pr_2^*d_1 + pr_3^*d_2$  in  $Pic(C_2 \times S)$ . By Nakai's criterion  $d_2$  is an ample divisor on S and  $d_1$  is ample on  $C_2$ .

By Lemma 2.1 it follows that  $H^1(pr_{23}(Z)) \simeq H^1(C_2 \times S) \simeq H^1(C_2) \oplus H^1(S)$ . Since Z is a Cartier divisor on  $C_1 \times pr_{23}(Z)$ , we can consider its cohomology class

 $cl(Z) \in H^2(C_1 \times pr_{23}(Z)) \simeq H^2(C_1) \oplus H^1(C_1) \otimes H^1(pr_{23}(Z)) \oplus H^2(pr_{23}(Z))$  which is the class associated to the line bundle  $\mathcal{O}(Z)$ .

We write  $cl(Z) = c_1 + c_2 + c_3$  according to this decomposition.

Let  $f: \mathbf{S} \to pr_{23}(Z)$  be a resolution of singularity. Since Z is not contained in the singular locus of  $C_2 \times pr_{23}(Z)$  we can take the pullback  $(id \times f)^*Z$  in  $C_2 \times \mathbf{S}$  with the associated line bundle  $(id \times f)^*\mathcal{O}(Z)$ . This pullback is compatible with the pullback on cohomology. The class  $(id_{C_1} \times f)^*c_2 \in H^1(C_1) \otimes f^*(H^1(C_2) \oplus H^1(S))$  is a Hodge cycle and by the assumption the cohomology  $H^1(C_1) \otimes (H^1(C_2) \oplus H^1(S))$  has no non-zero Hodge cycle. So we see that  $(id_{C_1} \times f)^*c_2 = 0$ .

So there are divisors  $d_1 \in Pic(C_1)$  and  $d_2 \in Pic(\mathbf{S})$  such that in  $Pic(C_1 \times \mathbf{S})$  there is an equality

$$(id_{C_1} \times f)^* Z = d_1 \times \mathbf{S} + C_1 \times d_2.$$

Pushing down to  $C_1 \times pr_{23}(Z)$  by the map  $id_{C_1} \times f$  we have an equality

$$Z = d_1 \times pr_{23}(Z) + C_1 \times f_*(d_2)$$

in  $A_2(C_1 \times pr_{23}(Z))$ .

Once Z is of this form, one can see that Chow-Künneth projector  $\pi_{1C_1} \times \pi_{1C_2} \times \pi_{2S}$  kills Z in  $CH^2(X)$  because by Conjcture (B) for  $C_2 \times S$  ([Mu2])  $\pi_{1C_2} \times \pi_{2S}$  kills  $pr_{23}(Z)$  and  $\pi_{1C_1}$  kills  $C_1$ . **Lemma 2.2.** If the projection  $pr_{23} : Z \to pr_{23}(Z)$  is flat, then Z is a Cartier divisor on  $C_1 \times pr_{23}(Z)$ .

Proof. Let  $I_Z$  be the ideal sheaf of Z in  $C_1 \times pr_{23}(Z)$ . For any point  $x \in pr_{23}(Z)$ , Let  $\{z_i\}_i$  be the set of closed points on the fiber  $Z \times_{pr_{23}(Z)} \operatorname{Spec}(x)$ . The image of  $I_Z$  in the local ring  $\mathcal{O}_{C_1 \times_{\mathbb{C}} \operatorname{Spec}(x), z_i}$  is a principal ideal  $(f_i)$ . For each i take a local section  $\tilde{f}_i \in I_Z$  which has the image  $f_i$  in  $\mathcal{O}_{C_1 \times_{\mathbb{C}} \operatorname{Spec}(x), z_i}$ .

For a sufficiently small neighborhood U of x in  $pr_{23}(Z)$  we can consider a Cartier divisor D on  $C_1 \times U$  which is defined by the equation  $\tilde{f}_i$  in a neighborhood of  $z_i$ . Let K be the kernel of natural surjection  $\mathcal{O}_D \to \mathcal{O}_Z$ .

Let  $\phi_D$  be the function on the set of points on U defined by

$$\phi_D(y) = \dim_{\kappa(y)} \mathcal{O}_D \otimes_{\mathcal{O}_U} \kappa(y).$$

It is an upper semicontinuous function on U. So there is an neighborhood  $U' \subset U$  of x such that for any  $y \in U'$ , one has

$$\phi_D(y) \le \phi_D(x)$$

On the other hand,  $\dim_{\kappa(y)} \mathcal{O}_Z \otimes_{\mathcal{O}_U} \kappa(y)$  is a constant function since Z is flat over  $pr_{23}(Z)$ . Also note that  $\phi_D(y) \geq \dim_{\kappa(y)} \mathcal{O}_Z \otimes_{\mathcal{O}_U} \kappa(y)$  on U'. Since  $\phi_D(x) = \dim_{\kappa(x)} \mathcal{O}_Z \otimes_{\mathcal{O}_U} \kappa(x)$  it follows that

$$\phi_D(y) = \dim_{\kappa(y)} \mathcal{O}_Z \otimes_{\mathcal{O}_U} \kappa(y)$$

on U'. As  $\mathcal{O}_Z$  is a flat  $\mathcal{O}_U$  module, it follows that  $K \otimes_{\mathcal{O}_U} \kappa(y) = 0$  for any point  $y \in U'$ . Hence K = 0.

# 3 A proof of Conjecture (B) for a product of two surfaces.

In this section we give a proof of the essential part of Conjecture (B) for a product of two surfaces.

Let  $S_1$  and  $S_2$  be projective smooth surfaces over  $\mathbb{C}$  and let  $X = S_1 \times S_2$ . For each  $S_i$  there is a Chow-Künneth decomposition  $\pi_{0S_i}, \dots, \pi_{4S_i}$  of the diagonal constructed by Murre([Mu]). They have the following properties:

 $\pi_4, \pi_3$  and  $\pi_0$  act as 0 on  $CH^1(S_i)_{\mathbb{Q}}$ .  $F^1CH^1(S_i)_{\mathbb{Q}} = Ker(\pi_2) = CH^1(S_i)_{hom,\mathbb{Q}}$ .  $F^2CH^1(S_i)_{\mathbb{Q}} = Ker(\pi_1|_{F^1}) = 0.$ 

 $\pi_0 \text{ and } \pi_1 \text{ act as } 0 \text{ on } CH^2(S_i)_{\mathbb{Q}}. F^1CH^2(S_i)_{\mathbb{Q}} = Ker(\pi_4) = CH^2(S_i)_{hom,\mathbb{Q}}. F^2CH^2(S_i)_{\mathbb{Q}} = Ker(\pi_3|_{F^1}) = Ker(alb: CH^2(S_i)_{hom,\mathbb{Q}} \to Alb(S_i) \otimes \mathbb{Q}). F^3CH^2(S_i)_{\mathbb{Q}} = Ker(\pi_2|_{F^2}) = 0.$ 

There is a Chow-Künneth decomposition for X given by the product of those for  $S_i$ . Murre has proven Conjecture (B) for X. Here we give another proof of the essential part of his result.

**Theorem 3.1.** The Chow-Künneth projectors  $\pi_{3S_1} \times \pi_{3S_2}$  and  $\pi_{3S_1} \times \pi_{2S_2}$  act as zero on  $CH^2(X)_{\mathbb{Q}}$ .

*Proof.* Let Z be an element of  $CH^2(X)$ . Let  $\eta_i \xrightarrow{j_i} S_i$  be the generic point of  $S_i$  for i = 1, 2 and  $Z_{\eta_i}$  be the generic fiber of Z.

The case of  $\pi_{3S_1} \times \pi_{3S_2}$ .  $(id_{S_1} \times j_2)^* (\pi_{3S_1} \times id_{S_2})(Z) = \pi_3 \times \eta_2((id_{S_1} \times j_2)^*Z)$ . We Write  $\pi_3 \times \eta_2 = \pi_{3\eta_2}$  and  $(id_{S_1} \times j_2)^* Z = Z_{\eta_2}$ . For p = 1 and  $2 \text{ let } C_p \stackrel{i_p}{\hookrightarrow} S_p$  be a smooth hyperplane section defined over the base field  $\mathbb{C}$ . Then by Lemma 2.3 of [Mu],  $i_{p_*}: Jac(C_p) \to Alb(S_p)$ is a surjection. So it follows that  $i_{1*}: Jac(C_1)(\eta_2)_{\mathbb{Q}} \to Alb(S_1)(\eta_2)_{\mathbb{Q}}$  is also surjective. Let d be the degree of  $Z_{\eta_2}$  and let  $e_1$  be a closed point on  $S_1$  which is rational over the base field  $\mathbb{C}$ . Then  $Z_{\eta_2} - d(e_1) \in CH^2(S_{1\underline{\eta}_2})_{hom,\mathbb{Q}}$  and so there is a cycle  $D \in Pic^0C_1(\eta_2)_{\mathbb{Q}}$ such that  $alb(Z_{n_2} - d(e_1)) = i_{1*}(D)$ . Let  $\overline{D}$  be the closure of D in X. Since D is supported on  $C_1 \times \eta_2$ ,  $\overline{D}$  is supported on  $\overline{C_1 \times \eta_2} = C_1 \times S_2$ .

Since  $Ker\pi_3 = Ker(alb)$ , we have the equality

$$(id_{S_1} \times j_2)^* (\pi_{3S_1} \times id_{S_2}) (Z - d(e_1) \times S_2 - \bar{D}) = \pi_{3S_1\eta_2} (Z_{\eta_2} - d(e_1) - i_{1*}D) = 0.$$

So it follows that

 $(\pi_{3S_1} \times id_{S_2})(Z) = (\pi_{3S_1} \times id_{S_2})(\bar{D}) + d\pi_{3S_1}(e_1) \times S_2 + \sum_k D_k$ 

where for each k  $D_k$  is supported on  $S_1 \times Y_k$  for an irreducible curve  $Y_k$ . We apply the projector  $\pi_{3S_1} \times id_{S_2}$  again on both sides of the equality. We apply  $\pi_{3S_1} \times id_{S_2}$  on each  $D_k$  as a cycle on  $S_1 \times Y_k$ . Let  $\eta_Y \stackrel{\mathcal{Y}}{\hookrightarrow} Y_k$  be the generic point of  $Y_k$ . We have the equality  $(id_{S_1} \times j_Y)^* (\pi_{3S_1} \times id_{Y_k})(D_k) = (\pi_{3S_1} \times \eta_Y)((id_{S_1} \times j_Y)^* D_k).$ 

Since  $(id_{S_1} \times j_Y)^* D_k$  is a divisor on the surface  $S_1 \times \eta_Y$ , from Conjecture (B) for  $S_1$  it follows that

 $(\pi_{3S_1} \times \eta_Y)((id_{S_1} \times j_Y)^*D_k) = 0.$ 

By taking closure of this equality in  $S_1 \times Y_k$  we have the equality

$$(\pi_{3S_1} \times id_{Y_k})(D_k) = \sum_i S_1 \times p_i$$

where for each i  $p_i$  is a closed point on  $Y_k$ . Applying  $\pi_{3S_1} \times id_{S_2}$  again on both sides of the equality it follows that  $(\pi_{3S_1} \times id_{S_2})(S_1 \times p_i) = 0$  since by Conjecture (B) for  $S_1$  $\pi_{3S_1}(S_1) = 0.$ 

Next we apply  $id_{S_1} \times \pi_{3S_2}$  on both sides of the equality. By Conjecture (B) for  $S_2$  we see that

$$(id_{S_1} \times \pi_{3S_2})(d\pi_{3S_1}(e_1) \times S_2) = d\pi_{3S_1}(e_1) \times \pi_{3S_2}(S_2) = 0.$$

Since

$$(\pi_{3S_1} \times id_{S_2})(id_{S_1} \times \pi_{3S_2}) = (id_{S_1} \times \pi_{3S_2})(\pi_{3S_1} \times id_{S_2})$$

it follows that

$$(id_{S_1} \times \pi_{3S_2})(\pi_{3S_1} \times id_{S_2})(\bar{D}) = (\pi_{3S_1} \times id_{S_2})(id_{S_1} \times \pi_{3S_2})(\bar{D}) = (\pi_{3S_1} \times id_{S_2})(\sum_l p_l \times S_2)$$

for a set of closed points  $p_l$  on  $S_1$ . Here we apply  $id_{S_1} \times \pi_{3S_2}$  on  $\overline{D}$  as a cycle on  $C_1 \times S_2$ . So we are reduced to the case where each component of Z is of the form  $pt \times S_2$  for a closed point pt. We can see that the projector  $\pi_{3S_1} \times \pi_{3S_2}$  kills  $pt \times S_2$  bacause by Conjecture (B) for surfaces  $\pi_{3S_i}(S_i) = 0$  for i = 1 and 2.

**Remark.** Murre pointed out that there is a simpler argument than the one above. We use the equality

$$\pi_{3S_1} \times id_{S_2}(Z) = Z \circ {}^t \pi_{3S_1} = Z \circ \pi_{1S_1}$$

where  $\circ$  is composition as correspondences and <sup>t</sup> is transpose. By construction of  $\pi_1$  there is a curve C on  $S_1$  such that  $\pi_{1S_1}$  is supported on  $C \times S_1(\text{cf.}$  (ii) of Proposition 2.1 in [KMP]). So one can immediately conclude that  $\pi_{3S_1} \times id_{S_2}(Z)$  is supported on  $C \times S_2$ .

The case of  $\pi_{3S_1} \times \pi_{2S_2}$ . We use the factorization  $\pi_{3S_1} \times \pi_{2S_2} = (id_{S_1} \times \pi_{2S_2})(\pi_{3S_1} \times id_{S_2})$ . We have the equality

$$(\pi_{3S_1} \times id_{S_2})(Z) = (\pi_{3S_1} \times id_{S_2})(\bar{D}) + d\pi_{3S_1}(e_1) \times S_2 + \sum_k D_k$$

where for each  $k D_k$  is supported on  $S_1 \times Y_k$  for an irreducible curve  $Y_k$  and  $\overline{D}$  is supported on  $C_1 \times S_2$ . The  $D_k$  part can be treated as above. Then we apply  $id_{S_1} \times \pi_{2S_2}$  on both sides of the equality. By Conjecture (B) for  $S_2$  it follows that

 $(id_{S_1} \times \pi_{2S_2})(d\pi_{3S_1}(e_1) \times S_2) = d\pi_{3S_1}(e_1) \times \pi_{2S_2}(S_2) = 0.$ By using the equality

$$(\pi_{3S_1} \times id_{S_2})(id_{S_1} \times \pi_{2S_2}) = (id_{S_1} \times \pi_{2S_2})(\pi_{3S_1} \times id_{S_2})$$

we have

$$(id_{S_1} \times \pi_{2S_2})(\pi_{3S_1} \times id_{S_2})(\bar{D}) = (\pi_{3S_1} \times id_{S_2})(id_{S_1} \times \pi_{2S_2})(\bar{D}).$$

Let  $\eta_{C_1} \stackrel{j_{C_1}}{\hookrightarrow} C_1$  be the generic point of  $C_1$ . We apply  $id_{S_1} \times \pi_{2S_2}$  on  $\overline{D}$  as a cycle on  $C_1 \times S_2$ . Since the divisor  $(j_{C_1} \times id_{S_2})^*(\overline{D})$  on  $\eta_{C_1} \times S_2$  is algebraically equivalent to a divisor  $\eta_{C_1} \times E$  on  $\eta_{C_1} \times S_2$  where E is a divisor on  $S_2$  defined over the base field  $\mathbb{C}$ , it follows that

$$(j_{C_1} \times id_{S_2})^* (id_{S_1} \times \pi_{2S_2}) (\bar{D} - C_1 \times E) = (\eta_{C_1} \times \pi_{2S_2}) ((j_{C_1} \times id_{S_2})^* (\bar{D}) - \eta_{C_1} \times E) = 0.$$

So by taking the closure of equality in  $C_1 \times S_2$  it follows that

$$(id_{S_1} \times \pi_{2S_2})(\bar{D}) = (id_{S_1} \times \pi_{2S_2})(C_1 \times E) + \sum_k p_k \times S_2$$

for a set  $\{p_k\}$  of closed points on  $S_1$ . In this way we are reduced to the case where each component of Z is a product of two curves or is of the form  $pt \times S_2$  or  $S_1 \times pt$ . By Conjecture (B) for surfaces one can see that the projector  $\pi_{3S_1} \times \pi_{2S_2}$  kills the cycles of this form in  $CH^2(X)_{\mathbb{Q}}$ .

## References

- [deM] del Angel, P.L., Müller-Stach, S.: Motives of uniruled 3-folds. Compositio Math. **112** (1998), 1-16.
- [deM2] del Angel, P.L., Müller-Stach, S.: On Chow motives of 3-folds. Trans. Amer. Math. Soc. **352** (2000), 1623-1633.
- [DM] Deninger, C., Murre, J.P.: Motivic decomposition of abelian schemes and the Fourier transform. J. Reine Angew. Math. **422** (1991), 201-219.
- [GM] Gordon, B., Murre, J.P.: Chow motives of elliptic modular threefolds. J. reine angew. Math. **514** (1999), 145–164.

[GHM]	Gordon, B., Hanamura, M., Murre, J.P.: Relative Chow-Kunneth projectors for modular varieties. J. reine angew. Math. <b>558</b> (2003), 1–14.
[GHM2]	Gordon, B., Hanamura, M., Murre, J.P.: Absolute Chow-Kunneth projectors for modular varieties. J. reine angew. Math. <b>580</b> (2005), 139–155.
[Ja]	Jannsen, U.: Motivic sheaves and filtrations on Chow groups. Proc. Sympos. Pure Math. <b>55 Part 1</b> (1994), 245–302.
[Ki]	Kimura, S.: Chow groups are finite dimensional, in some sense. Math. Ann. <b>331</b> (2005), 173–201.
[KMP]	Kahn, B., Murre, J.P., Pedrini, C.: On the transcendental part of the motive of a surface. preprint, (2005).
[Mu]	Murre, J.P.: On the motive of an algebraic surface J. reine angew. Math. <b>409</b> (1990), 190-204.
[Mu2]	Murre, J.P.: On a conjectural filtration on the Chow groups of an algebraic variety, (I and II), Indag. Math. (N.S.) ${\bf 4}$ (1993), 177-201.
[Pic]	Miller, A., Müller-Stach, S., Wortmann, S., Yang, YH., Zuo, K.: Chow-Künneth decomposition for universal families over Picard modular surfaces preprint, (2005).
[Sh]	Shermenev, A. M. Motif of an Abelian variety. Funckcional. Anal. i Priložen. <b>8</b> (1974), 55–61.
Kenichiro Kimura Institute of Mathematics University of Tsukuba	

Tsukuba, Ibaraki 305-8571 Japan email: kimurak@math.tsukuba.ac.jp