

The moduli space of curves is rigid

Paul Hacking

November 8, 2018

1 Introduction

Kapranov has proposed the following informal statement [Kapranov97]. Given a smooth variety $X = X(0)$, consider the moduli space $X(1)$ of varieties obtained as deformations of $X(0)$, the moduli space $X(2)$ of deformations of $X(1)$, and so on. Then this process should stop after $n = \dim X$ steps, i.e., $X(n)$ should be rigid (no infinitesimal deformations). Roughly speaking, one thinks of $X(1)$ as H^1 of a sheaf of non-abelian groups on $X(0)$. Indeed, at least the tangent space to $X(1)$ at $[X]$ is identified with $H^1(T_X)$, where T_X is the tangent sheaf, the sheaf of first order infinitesimal automorphisms of X . Then one regards $X(m)$ as a kind of non-Abelian H^m , and the analogy with the usual definition of Abelian H^m suggests the statement above.

In particular, the moduli space of curves should be rigid. In this paper, we verify this in the following precise form: the moduli stack of stable curves of genus g with n marked points is rigid for each g and n .

On the other hand, moduli spaces of surfaces should have non-trivial deformations in general. I do not know any examples at present, but hope to address this question in the future. It seems plausible that there should be a non-trivial deformation of a moduli space of surfaces whose fibres parametrise “generalised surfaces” in some sense, for example non-commutative surfaces. From this point of view the result of this paper says that the concept of a curve cannot be deformed.

Let us also note that our result can be thought of as a version of Mostow rigidity for the mapping class group. Recall that the moduli space M_g of smooth complex curves of genus g is the quotient of the Teichmüller space T_g by the mapping class group Γ_g . The space T_g is a bounded domain in \mathbb{C}^{3g-3} , which is homeomorphic to a ball, and Γ_g acts discontinuously on T_g with finite stabilisers. We thus obtain M_g as a complex orbifold with orbifold fundamental group Γ_g . The space T_g admits a natural metric, the Weil–Petersson metric, which has negative holomorphic sectional curvatures. So,

roughly speaking, M_g looks like a quotient of a complex ball by a discrete group Γ of isometries, with finite volume. Mostow rigidity predicts that such a quotient is uniquely determined by the group Γ up to complex conjugation. (This is certainly true if Γ acts freely with compact quotient, see [Siu80]). In particular, it should have no infinitesimal deformations. Unfortunately I do not know a proof along these lines.

Acknowledgements: We thank M. Kapranov for suggesting the problem and subsequent encouragement, S. Keel for help with $\overline{\mathcal{M}}_{g,n}$, and M. Olsson for help with log structures. We also thank I. Dolgachev, G. Farkas, S. Grushchinsky, J. Kollár, S. Kovács, R. Lazarsfeld, A. Marian, S.P. Smith and T. Stafford for helpful discussions.

2 Statements

We work over an algebraically closed field k of characteristic zero. Let g and n be non-negative integers such that $2g - 2 + n > 0$. Let $\overline{\mathcal{M}}_{g,n}$ denote the moduli stack of stable curves of genus g with n marked points. The stack $\overline{\mathcal{M}}_{g,n}$ is a smooth proper Deligne–Mumford stack of dimension $3g - 3 + n$.

Theorem 2.1. *The stack $\overline{\mathcal{M}}_{g,n}$ is rigid, that is, has no infinitesimal deformations.*

Let $\partial\overline{\mathcal{M}}_{g,n} \subset \overline{\mathcal{M}}_{g,n}$ denote the boundary of the moduli stack, that is, the complement of the locus of smooth curves (with its reduced structure). The locus $\partial\overline{\mathcal{M}}_{g,n}$ is a normal crossing divisor in $\overline{\mathcal{M}}_{g,n}$.

Theorem 2.2. *The pair $(\overline{\mathcal{M}}_{g,n}, \partial\overline{\mathcal{M}}_{g,n})$ has no locally trivial deformations.*

Let $\overline{M}_{g,n}$ denote the coarse moduli space of the stack $\overline{\mathcal{M}}_{g,n}$. The space $\overline{M}_{g,n}$ is a projective variety with quotient singularities.

Theorem 2.3. *The variety $\overline{M}_{g,n}$ has no locally trivial deformations if $(g, n) \neq (1, 2), (2, 0), (2, 1), (3, 0)$.*

Remark 2.4. In the exceptional cases, the projection $\overline{\mathcal{M}}_{g,n} \rightarrow \overline{M}_{g,n}$ is ramified in codimension one over the interior of $\overline{M}_{g,n}$, and an additional calculation is needed to relate the deformations of the stack and the deformations of the coarse moduli space (cf. Prop. 5.2). Presumably the result still holds.

3 Proof of Theorem 2.2

Let $\Omega_{\overline{\mathcal{M}}_{g,n}}(\log)$ denote the sheaf of 1-forms on $\overline{\mathcal{M}}_{g,n}$ with logarithmic poles along the boundary, and $T_{\overline{\mathcal{M}}_{g,n}}(\log)$ the dual of $\Omega_{\overline{\mathcal{M}}_{g,n}}(\log)$. The sheaf $T_{\overline{\mathcal{M}}_{g,n}}(\log)$ is the subsheaf of the tangent sheaf $T_{\overline{\mathcal{M}}_{g,n}}$ consisting of vector fields on $\overline{\mathcal{M}}_{g,n}$ which are tangent to the boundary. In other words, it is the sheaf of first order infinitesimal automorphisms of the pair $(\overline{\mathcal{M}}_{g,n}, \partial\overline{\mathcal{M}}_{g,n})$. Hence the first order locally trivial deformations of the pair $(\overline{\mathcal{M}}_{g,n}, \partial\overline{\mathcal{M}}_{g,n})$ are identified with the space $H^1(T_{\overline{\mathcal{M}}_{g,n}}(\log))$. To prove Thm. 2.2, we show $H^1(T_{\overline{\mathcal{M}}_{g,n}}(\log)) = 0$.

Let $\pi : \mathcal{U}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$ denote the universal family over $\overline{\mathcal{M}}_{g,n}$. That is, $\mathcal{U}_{g,n}$ is the stack of n -pointed stable curves of genus g together with an extra section (with no smoothness condition). There is an isomorphism of stacks

$$c : \overline{\mathcal{M}}_{g,n+1} \xrightarrow{\sim} \mathcal{U}_{g,n}$$

which identifies the morphism $p_{n+1} : \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ given by forgetting the last point with the projection $\pi : \mathcal{U}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$ [Knudsen83a, Sec. 1–2]. We define the boundary $\partial\mathcal{U}_{g,n} \subset \mathcal{U}_{g,n}$ as the image of $\partial\overline{\mathcal{M}}_{g,n+1} \subset \overline{\mathcal{M}}_{g,n+1}$ under the isomorphism c . Equivalently, $\partial\mathcal{U}_{g,n}$ is the union of $\pi^*\partial\overline{\mathcal{M}}_{g,n}$ and the n tautological sections of π .

We assume the reader is familiar with log structures at the level of [KatoK89]. If \mathcal{X} is a Deligne–Mumford stack, we can consider the étale site of \mathcal{X} , and define log structures on \mathcal{X} as in the case of schemes. If $(X, \mathcal{M}) \rightarrow (Y, \mathcal{N})$ is a morphism of log schemes (or stacks), we write $\Omega_{X/Y}(\log)$ for the sheaf of logarithmic 1-forms, and $T_{X/Y}(\log)$ for the dual sheaf.

The normal crossing divisors $\partial\overline{\mathcal{M}}_{g,n} \subset \overline{\mathcal{M}}_{g,n}$ and $\partial\mathcal{U}_{g,n} \subset \mathcal{U}_{g,n}$ define smooth log structures $\mathcal{M}_{\overline{\mathcal{M}}_{g,n}}$ and $\mathcal{M}_{\mathcal{U}_{g,n}}$. The morphism $\pi : \mathcal{U}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$ extends naturally to a log smooth morphism of log stacks $\pi : (\mathcal{U}_{g,n}, \mathcal{M}_{\mathcal{U}_{g,n}}) \rightarrow (\overline{\mathcal{M}}_{g,n}, \mathcal{M}_{\overline{\mathcal{M}}_{g,n}})$. There is an exact sequence

$$0 \rightarrow \pi^*\Omega_{\overline{\mathcal{M}}_{g,n}}(\log) \rightarrow \Omega_{\mathcal{U}_{g,n}}(\log) \rightarrow \Omega_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}(\log) \rightarrow 0. \quad (1)$$

Here $\Omega_{\overline{\mathcal{M}}_{g,n}}(\log)$ and $\Omega_{\mathcal{U}_{g,n}}(\log)$ are the sheaves of 1-forms with logarithmic poles along the boundary. Let ω_π denote the relative dualising sheaf of the morphism $\pi : \mathcal{U}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$ and Σ the union of the n tautological sections of π . The sheaf $\Omega_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}(\log)$ is identified with the line bundle $\omega_\pi(\Sigma)$. Indeed, the exact sequence above induces an isomorphism

$$\begin{aligned} \Omega_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}(\log) &\cong \det \Omega_{\mathcal{U}_{g,n}}(\log) \otimes \pi^* \det \Omega_{\overline{\mathcal{M}}_{g,n}}(\log)^\vee \\ &= \omega_{\mathcal{U}_{g,n}}(\partial\mathcal{U}_{g,n}) \otimes \pi^* \omega_{\overline{\mathcal{M}}_{g,n}}(\partial\overline{\mathcal{M}}_{g,n})^\vee. \end{aligned}$$

Now $\omega_{\mathcal{U}_{g,n}} \otimes \pi^* \omega_{\overline{\mathcal{M}}_{g,n}}^\vee \cong \omega_\pi$ since π is a Gorenstein morphism [Hartshorne66, p. 142–4] and $\partial \mathcal{U}_{g,n} - \pi^* \partial \overline{\mathcal{M}}_{g,n} = \Sigma$, so $\Omega_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}(\log) \cong \omega_\pi(\Sigma)$ as claimed.

Dualising the exact sequence (1) we obtain an exact sequence

$$0 \rightarrow T_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}(\log) \rightarrow T_{\mathcal{U}_{g,n}}(\log) \rightarrow \pi^* T_{\overline{\mathcal{M}}_{g,n}}(\log) \rightarrow 0.$$

Applying π_* , we obtain a long exact sequence of sheaves on $\overline{\mathcal{M}}_{g,n}$ with connecting homomorphism

$$\delta : T_{\overline{\mathcal{M}}_{g,n}}(\log) \rightarrow R^1 \pi_* T_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}(\log).$$

We claim that the map δ is an isomorphism. The idea is the following: The map δ is the Kodaira–Spencer map for the log smooth family $(\mathcal{U}_{g,n}, \mathcal{M}_{\mathcal{U}_{g,n}}) \rightarrow (\overline{\mathcal{M}}_{g,n}, \mathcal{M}_{\overline{\mathcal{M}}_{g,n}})$ (we make this precise below). The family $(\mathcal{U}_{g,n}, \mathcal{M}_{\mathcal{U}_{g,n}}) \rightarrow (\overline{\mathcal{M}}_{g,n}, \mathcal{M}_{\overline{\mathcal{M}}_{g,n}})$ is the universal log stable curve of type (g, n) (see [KatoF00] for the precise statement). Hence δ is an isomorphism.

Let us now be more explicit. Let $U \rightarrow \overline{\mathcal{M}}_{g,n}$ be étale and $p \in U$ a point. Then the restriction of the universal family of pointed stable curves to the germ $(p \in U)$ is the universal deformation of the fibre (C, x_1, \dots, x_n) over p . Let \mathcal{M}_U denote the log structure on U . Write $k = k(p)$, and let \mathcal{M}_k denote the log structure on $\text{Spec } k$ pulled back from U . Let $A = k[t]/(t^2)$, and let \mathcal{M}_A denote the log structure on $\text{Spec } A$ pulled back from $\text{Spec } k$ via the retraction $\text{Spec } A \rightarrow \text{Spec } k$. Then $T_{\overline{\mathcal{M}}_{g,n}}(\log) \otimes k$ is identified with the set of liftings of $(\text{Spec } k, \mathcal{M}_k) \rightarrow (U, \mathcal{M}_U)$ to a morphism $(\text{Spec } A, \mathcal{M}_A) \rightarrow (U, \mathcal{M}_U)$, by [KatoK89, Prop. 3.9, p. 203]. Let \mathcal{M}_C denote the log structure on C pulled back from the log structure on $\mathcal{U}_{g,n}$. The fibre $H^1(T_{C/k}(\log))$ of $R^1 \pi_* T_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}(\log)$ at p is identified with the set of log smooth deformations of $(C, \mathcal{M}_C)/(\text{Spec } k, \mathcal{M}_k)$ over $(\text{Spec } A, \mathcal{M}_A)$, i.e., log smooth morphisms $(C', \mathcal{M}_{C'})/(\text{Spec } A, \mathcal{M}_A)$ extending $(C, \mathcal{M}_C)/(\text{Spec } k, \mathcal{M}_k)$, by [KatoK89, Prop. 3.14(3), p. 205]. The map $\delta \otimes k : T_{\overline{\mathcal{M}}_{g,n}}(\log) \otimes k \rightarrow H^1(T_{C/k}(\log))$ sends a morphism $(\text{Spec } A, \mathcal{M}_A) \rightarrow (U, \mathcal{M}_U)$ to the log smooth deformation $(C', \mathcal{M}_{C'})/(\text{Spec } A, \mathcal{M}_A)$ obtained by pullback from the family over U . This is verified exactly as in the case of a smooth family of schemes (cf., e.g., [Voisin02, 9.1.2, p. 223–5]).

It remains to check that $\delta \otimes k$ is an isomorphism. The sheaf $T_{C/k}(\log)$ is the dual of $\omega_C(x_1 + \dots + x_n)$, where ω_C is the dualising sheaf, hence $\dim H^1(T_{C/k}(\log)) = 3g - 3 + n = \dim T_{\overline{\mathcal{M}}_{g,n}}(\log) \otimes k$ by Riemann Roch. So it is enough to show that $\delta \otimes k$ is injective. Suppose that a lifting $(\text{Spec } A, \mathcal{M}_A) \rightarrow (U, \mathcal{M}_U)$ induces the trivial log smooth deformation of $(C, \mathcal{M}_C)/(\text{Spec } k, \mathcal{M}_k)$. Then, since $(p \in U)$ is the universal deformation

space of (C, x_1, \dots, x_n) , it follows that the underlying morphism of schemes $\text{Spec } A \rightarrow U$ is the “constant” morphism $f : \text{Spec } A \rightarrow \text{Spec } k \rightarrow U$. The morphism $(\text{Spec } A, \mathcal{M}_A) \rightarrow (\text{Spec } U, \mathcal{M}_U)$ is then given by an automorphism $\sigma : f^* \mathcal{M}_U = \mathcal{M}_A \rightarrow \mathcal{M}_A$. So, the induced log deformation $(C', \mathcal{M}_{C'}) \rightarrow (\text{Spec } A, \mathcal{M}_A)$ is isomorphic to the deformation given by

$$(C, \mathcal{M}_C) \times_{(\text{Spec } k, \mathcal{M}_k)} (\text{Spec } A, \mathcal{M}_A) \rightarrow (\text{Spec } A, \mathcal{M}_A) \xrightarrow{g} (\text{Spec } A, \mathcal{M}_A),$$

where g is the automorphism given by the identity map on $\text{Spec } A$ and the automorphism σ^{-1} of \mathcal{M}_A . By [KatoF00, Prop. 2.1, p. 228] (or explicit calculation, cf. [Olsson01, Lem. 5.3.2, p. 112]), the log smooth deformation $(C', \mathcal{M}_{C'})/(\text{Spec } A, \mathcal{M}_A)$ is isomorphic to the trivial deformation only if σ equals the identity. Thus $\delta \otimes k$ is injective, as required.

We have $H^0(T_{C/k}(\log)) = 0$ for each fibre C/k of $\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}$. Hence $\pi_* T_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}(\log) = 0$. Also $R^i \pi_* T_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}(\log) = 0$ for $i > 1$ by dimensions. So $H^{i+1}(T_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}) = H^i(R^1 \pi_* T_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}})$ by the Leray spectral sequence. Hence the isomorphism δ induces an isomorphism

$$H^i(T_{\overline{\mathcal{M}}_{g,n}}(\log)) \xrightarrow{\sim} H^{i+1}(T_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}(\log)) \quad (2)$$

for each i .

Recall that the sheaf $T_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}(\log)$ is the dual of the line bundle $\omega_\pi(\Sigma)$. Let $U_{g,n}$ denote the coarse moduli space of the stack $\mathcal{U}_{g,n}$ and $p : \mathcal{U}_{g,n} \rightarrow U_{g,n}$ the projection. The line bundle $\omega_\pi(\Sigma)$ on the stack $\mathcal{U}_{g,n}$ defines a \mathbb{Q} -line bundle $p_* \omega_\pi(\Sigma)$ on the coarse moduli space $U_{g,n}$ (see Sec. 5). We use the following important result, which can be viewed as a generalisation of a result of Arakelov [Arakelov71, Prop. 3.2, p. 1297], and depends on subsequent work by Fujita, Keel, Kollár, Viehweg and others.

Theorem 3.1. [Keel99, Thm 0.4, p. 255] *The \mathbb{Q} -line bundle $p_* \omega_\pi(\Sigma)$ is big and nef on $U_{g,n}$.*

It follows by Kodaira vanishing (see Thm 6.1) that $H^i(T_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}) = H^i(\omega_\pi(\Sigma)^\vee) = 0$ for $i < \dim \mathcal{U}_{g,n}$. Combining with (2), we deduce

Proposition 3.2. *$H^i(T_{\overline{\mathcal{M}}_{g,n}}(\log)) = 0$ for $i < \dim \overline{\mathcal{M}}_{g,n}$.*

In particular, $H^1(T_{\overline{\mathcal{M}}_{g,n}}(\log)) = 0$ if $\dim \overline{\mathcal{M}}_{g,n} > 1$. The remaining cases are easy to check. This completes the proof of Theorem 2.2.

4 Proof of Theorem 2.1

We now prove Theorem 2.1. Since $\overline{\mathcal{M}}_{g,n}$ is a smooth Deligne–Mumford stack, its first order infinitesimal deformations are identified with the space $H^1(T_{\overline{\mathcal{M}}_{g,n}})$, and we must show that $H^1(T_{\overline{\mathcal{M}}_{g,n}}) = 0$. Let \mathcal{B} denote the boundary $\partial\overline{\mathcal{M}}_{g,n}$, and let $\nu: \mathcal{B}^\nu \rightarrow \mathcal{B}$ denote its normalisation. We have an exact sequence

$$0 \rightarrow \Omega_{\overline{\mathcal{M}}_{g,n}} \rightarrow \Omega_{\overline{\mathcal{M}}_{g,n}}(\log) \rightarrow \nu_* \mathcal{O}_{\mathcal{B}^\nu} \rightarrow 0$$

where the last map is the Poincaré residue map. Dualising, we obtain an exact sequence

$$0 \rightarrow T_{\overline{\mathcal{M}}_{g,n}}(\log) \rightarrow T_{\overline{\mathcal{M}}_{g,n}} \rightarrow \nu_* \mathcal{N} \rightarrow 0$$

where \mathcal{N} is a line bundle on \mathcal{B}^ν . Roughly speaking, the line bundle \mathcal{N} is the normal bundle of the map $\mathcal{B}^\nu \rightarrow \overline{\mathcal{M}}_{g,n}$. More precisely, étale locally at a point p of \mathcal{B}^ν , the line bundle \mathcal{N} is identified with the pullback of the normal bundle of the corresponding branch of the boundary $\mathcal{B} \subset \overline{\mathcal{M}}_{g,n}$ at $\nu(p)$. Consider the associated long exact sequence of cohomology

$$\cdots \rightarrow H^i(T_{\overline{\mathcal{M}}_{g,n}}(\log)) \rightarrow H^i(T_{\overline{\mathcal{M}}_{g,n}}) \rightarrow H^i(\mathcal{N}) \rightarrow \cdots$$

We prove below that $H^i(\mathcal{N}) = 0$ for $i < \dim \mathcal{B}$. Now, since $H^i(T_{\overline{\mathcal{M}}_{g,n}}(\log)) = 0$ for $i < \dim \overline{\mathcal{M}}_{g,n}$ by Prop. 3.2, we deduce

Proposition 4.1. $H^i(T_{\overline{\mathcal{M}}_{g,n}}) = 0$ for $i < \dim \overline{\mathcal{M}}_{g,n} - 1$.

In particular, $H^1(T_{\overline{\mathcal{M}}_{g,n}}) = 0$ if $\dim \overline{\mathcal{M}}_{g,n} > 2$. In the remaining cases it is easy to check that $H^1(\mathcal{N}) = 0$, so again $H^1(T_{\overline{\mathcal{M}}_{g,n}}) = 0$.

The irreducible components of the normalisation \mathcal{B}^ν of the boundary $\mathcal{B} \subset \overline{\mathcal{M}}_{g,n}$ are finite images of the following stacks [Knudsen83a, Def. 3.8, Cor. 3.9]:

- (1) $\overline{\mathcal{M}}_{g_1, S_1 \cup \{n+1\}} \times \overline{\mathcal{M}}_{g_2, S_2 \cup \{n+2\}}$ where $g_1 + g_2 = g$ and S_1, S_2 is a partition of $\{1, \dots, n\}$.
- (2) $\overline{\mathcal{M}}_{g-1, n+2}$

Here $\overline{\mathcal{M}}_{h,S}$ denotes the moduli stack of stable curves of genus h with marked points labelled by a finite set S . In each case the map to \mathcal{B}^ν is given by identifying the points labelled by $n+1$ and $n+2$. The map is an isomorphism

onto the component of \mathcal{B}^ν except in case (1) for $g_1 = g_2$ and $n = 0$ and case (2), when it is étale of degree 2.

For $\overline{\mathcal{M}}_{h,S}$ a moduli stack of pointed stable curves as above, let $\pi: \mathcal{U}_{h,S} \rightarrow \overline{\mathcal{M}}_{h,S}$ denote the universal family, and $x_i: \overline{\mathcal{M}}_{h,S} \rightarrow \mathcal{U}_{h,S}$, $i \in S$, the tautological sections of π . Define $\psi_i = x_i^* \omega_\pi$, the pullback of the relative dualising sheaf of π along the section x_i .

Lemma 4.2. *The pullback of \mathcal{N}^\vee to $\overline{\mathcal{M}}_{g_1, S_1 \cup \{n+1\}} \times \overline{\mathcal{M}}_{g_2, S_2 \cup \{n+2\}}$ is identified with $\text{pr}_1^* \psi_{n+1} \otimes \text{pr}_2^* \psi_{n+2}$. Similarly, the pullback of \mathcal{N}^\vee to $\overline{\mathcal{M}}_{g-1, n+2}$ is identified with $\psi_{n+1} \otimes \psi_{n+2}$.*

Proof. Let $U \rightarrow \overline{\mathcal{M}}_{g,n}$ be étale and $p \in U$ a point. Let (C, x_1, \dots, x_n) be the fibre of the universal family of pointed stable curves over p . The tangent space at $p \in U$ is identified with the space of first order deformations of (C, x_1, \dots, x_n) , which equals $\text{Ext}^1(\Omega_C(\sum x_i), \mathcal{O}_C)$, cf. [DM69, p. 79–82]. The components of the boundary at $p \in U$ are labelled by the nodes of C — the component B_q labelled by a node q is the locus where the induced deformation of the germ $(q \in C)$ is trivial. The fibre of the normal bundle $\mathcal{N}_{B_q/U}$ at p is thus identified with the 1-dimensional k -vector space $\text{Ext}^1(\Omega_{\hat{\mathcal{O}}_{C,q}}, \hat{\mathcal{O}}_{C,q})$. Let $C^\nu \rightarrow C$ denote the normalisation of C and q_1, q_2 the pre-images of q . The space $\text{Ext}^1(\Omega_{\hat{\mathcal{O}}_{C,q}}, \hat{\mathcal{O}}_{C,q})$ is canonically isomorphic to $\mathcal{O}_{C^\nu}(q_1)|_{q_1} \otimes \mathcal{O}_{C^\nu}(q_2)|_{q_2}$ [Friedman83, Prop. 2.3, p. 85], and there are isomorphisms $\omega_{C^\nu} \otimes \mathcal{O}_{C^\nu}(q_i)|_{q_i} \cong \mathcal{O}_{q_i}$ given by the residue map. Combining, we obtain a canonical isomorphism $\mathcal{N}_{B_q/U}^\vee \otimes k(p) \cong \omega_{C^\nu}|_{q_1} \otimes \omega_{C^\nu}|_{q_2}$. This gives the desired identifications. \square

Lemma 4.3. [Knudsen83b, Thm. 4.1(d), p. 202] *The line bundle ψ_{n+1} on $\overline{\mathcal{M}}_{g,n+1}$ is identified with the pullback of the line bundle $\omega_\pi(\Sigma)$ under the isomorphism $\phi: \overline{\mathcal{M}}_{g,n+1} \rightarrow \mathcal{U}_{g,n}$.*

Corollary 4.4. *The \mathbb{Q} -line bundle on the coarse moduli space of \mathcal{B}^ν defined by \mathcal{N}^\vee is big and nef on each component*

Proof. This follows immediately from Lem. 4.2, Lem. 4.3, and Thm. 3.1. \square

We deduce that $H^i(\mathcal{N}) = 0$ for $i < \dim \mathcal{B}$ by Thm 6.1. This completes the proof of Theorem 2.1.

5 Proof of Theorem 2.3

We first prove a basic lemma which relates the deformations of a smooth Deligne–Mumford stack and its coarse moduli space.

Let \mathcal{X} be a smooth proper Deligne–Mumford stack, X the coarse moduli space of \mathcal{X} , and $p : \mathcal{X} \rightarrow X$ the projection. Let $T_{\mathcal{X}}$ denote the tangent sheaf of \mathcal{X} . Let $D \subset X$ be the union of the codimension one components of the branch locus of $p : \mathcal{X} \rightarrow X$ (with its reduced structure). Let $T_X(\log D)$ denote the subsheaf of the tangent sheaf T_X consisting of derivations which preserve the ideal sheaf of D . It is the sheaf of first order infinitesimal automorphisms of the pair (X, D) .

Lemma 5.1. $p_*T_{\mathcal{X}} = T_X(\log D)$

Proof. The sheaves $p_*T_{\mathcal{X}}$ and $T_X(\log D)$ satisfy Serre’s S_2 condition, and are identified over the locus where p is étale. So it suffices to work in codimension 1. We reduce to the case $\mathcal{X} = [\mathbb{A}_x^1/\mu_e]$, where $\mu_e \ni \zeta : x \mapsto \zeta x$. Then $X = \mathbb{A}_x^1/\mu_e = \mathbb{A}_y^1$, where $y = x^e$, and $D = (y = 0) \subset X$. Let $\pi : \mathbb{A}_x^1 \rightarrow \mathbb{A}_y^1/\mu_e$ be the quotient map. We compute

$$p_*T_{\mathcal{X}} = \left(\pi_* \mathcal{O}_{\mathbb{A}_x^1} \cdot \frac{\partial}{\partial x} \right)^{\mu_e} = \mathcal{O}_{\mathbb{A}_y^1} \cdot x \frac{\partial}{\partial x} = \mathcal{O}_{\mathbb{A}_y^1} \cdot y \frac{\partial}{\partial y} = T_X(\log D),$$

as required. \square

Proposition 5.2. *The first order deformations of the stack \mathcal{X} are identified with the first order locally trivial deformations of the pair (X, D) .*

Proof. By the Lemma, $H^1(T_{\mathcal{X}}) = H^1(p_*T_{\mathcal{X}}) = H^1(T_X(\log D))$. \square

We now apply this result to relate deformations of the stack $\overline{\mathcal{M}}_{g,n}$ and its coarse moduli space $\overline{M}_{g,n}$.

A stable n -pointed curve of genus 0 has no non-trivial automorphisms. Hence the stack $\overline{\mathcal{M}}_{0,n}$ is equal to its coarse moduli space $\overline{M}_{0,n}$, and $\overline{M}_{0,n}$ is rigid by Thm. 2.1. Also, recall that $\overline{M}_{1,1}$ is isomorphic to \mathbb{P}^1 and therefore rigid. So, in the following, we assume that $g \neq 0$ and $(g, n) \neq (1, 1)$.

Let $\mathcal{D} \subset \overline{\mathcal{M}}_{g,n}$ be the component of the boundary whose general point is a curve with two components of genus 1 and $g - 1$ meeting in a node, with each of the n marked points on the component of genus $g - 1$. Note that each point of \mathcal{D} has a non-trivial automorphism given by the involution of the component of genus 1 fixing the node. Let $p : \overline{\mathcal{M}}_{g,n} \rightarrow \overline{M}_{g,n}$ be the projection, and $D \subset \overline{M}_{g,n}$ the coarse moduli space of \mathcal{D} .

Proposition 5.3. *If $g + n \geq 4$ then the automorphism group of a general point of $\overline{\mathcal{M}}_{g,n}$ is trivial, and the divisor $D \subset \overline{M}_{g,n}$ is the unique codimension 1 component of the branch locus of p .*

Proof. This is a straightforward dimension count. One shows that the locus of *smooth* curves in $\overline{\mathcal{M}}_{g,n}$ with a non-trivial automorphism has codimension at least $(g-2)+n$. Together with the description of the components of the boundary this implies the result. \square

Assume $g+n \geq 4$. Let $\nu : \mathcal{D}^\nu \rightarrow \mathcal{D}$ denote the normalisation, then $\mathcal{D}^\nu \cong \overline{\mathcal{M}}_{1,1} \times \overline{\mathcal{M}}_{g-1,n+1}$. Let \mathcal{N} denote the normal bundle of the map $\mathcal{D}^\nu \rightarrow \overline{\mathcal{M}}_{g,n}$, cf. Sec. 4.

Lemma 5.4. *There is an exact sequence*

$$0 \rightarrow T_{\overline{\mathcal{M}}_{g,n}}(\log D) \rightarrow T_{\overline{\mathcal{M}}_{g,n}} \rightarrow p_* \nu_* \mathcal{N}^{\otimes 2} \rightarrow 0.$$

Proof. We identify the cokernel of the map $T_{\overline{\mathcal{M}}_{g,n}}(\log D) \rightarrow T_{\overline{\mathcal{M}}_{g,n}}$. Étale locally over the coarse moduli space, the projection p is of the form $[U/G] \rightarrow U/G$, where U is a smooth affine variety and G a finite group acting on U . We reduce to the case $U = \mathbb{A}^N$ with G acting linearly. Let $V = U/G$ and $\pi : U \rightarrow V$ the quotient map. Let $H \subset G$ be the (normal) subgroup generated by pseudo-reflections. Write $U' = U/H$ and let $\alpha : U \rightarrow U'$ and $\beta : U' \rightarrow V$ be the quotient maps. Then U' is smooth and β is étale in codimension 1. Let $B \subset U$ be the union of the reflecting hyperplanes, B' its image in U' , and D its image in V . In our case B is a normal crossing divisor — it is the pullback of the divisor $\mathcal{D} \subset \overline{\mathcal{M}}_{g,n}$ by Prop 5.3. Hence we may assume $H = \mu_{m_1} \times \cdots \times \mu_{m_r}$, and H acts on $U = \mathbb{A}_{x_1, \dots, x_N}^N$ via

$$H \ni h = (\zeta_1, \dots, \zeta_r) : x_i \mapsto \begin{cases} \zeta_i x_i & \text{if } i \leq r \\ x_i & \text{if } i > r \end{cases}$$

Then $B = B_1 \cup \cdots \cup B_r$ where $B_i = (x_i = 0)$. Also $U' = \mathbb{A}_{y_1, \dots, y_N}^N$, where $y_i = x_i^{m_i}$ if $i \leq r$ and $y_i = x_i$ if $i > r$, and $B' = B'_1 \cup \cdots \cup B'_r$ where $B'_i = (y_i = 0)$. There is an exact sequence

$$0 \rightarrow T_{U'}(\log B') \rightarrow T_{U'} \rightarrow \oplus \mathcal{N}_{B'_i/U'} \rightarrow 0.$$

Moreover, there is a canonical isomorphism $\mathcal{N}_{B'_i/U'} \cong (\alpha_* \mathcal{N}_{B_i/U}^{\otimes m_i})^H$. Applying $(\beta_*(\cdot))^{G/H}$ we obtain an exact sequence

$$0 \rightarrow T_V(\log D) \rightarrow T_V \rightarrow \pi_*(\oplus \mathcal{N}_{B_i/U}^{\otimes m_i})^G \rightarrow 0.$$

In our case $m_i = 2$ for each i , since a general point of $\mathcal{D} \subset \overline{\mathcal{M}}_{g,n}$ has automorphism group $\mathbb{Z}/2\mathbb{Z}$. Hence the cokernel of $T_{\overline{\mathcal{M}}_{g,n}}(\log D) \rightarrow T_{\overline{\mathcal{M}}_{g,n}}$ is identified with $p_* \nu_* \mathcal{N}^{\otimes 2}$ as claimed. \square

We have $H^1(T_{\overline{M}_{g,n}}(\log D)) = H^1(T_{\overline{M}_{g,n}}) = 0$ by Prop. 5.2 and Sec. 4. Also $H^1(\mathcal{N}^{\otimes 2}) = 0$ by Thm. 6.1 because the \mathbb{Q} -line bundle defined by \mathcal{N}^\vee on the coarse moduli space of \mathcal{D}^ν is big and nef by Cor. 4.4. So $H^1(T_{\overline{M}_{g,n}}) = 0$, that is, $\overline{M}_{g,n}$ has no locally trivial deformations. This concludes the proof of Thm. 2.3.

6 Appendix: Kodaira vanishing for stacks

Let \mathcal{X} be a smooth proper Deligne–Mumford stack, X the coarse moduli space of \mathcal{X} , and $p : \mathcal{X} \rightarrow X$ the projection. Étale locally on X , $p : \mathcal{X} \rightarrow X$ is of the form $p : [U/G] \rightarrow U/G$, where U is a smooth affine variety and G is a finite group acting on U [AV02, Lemma 2.2.3, p. 32]. A sheaf \mathcal{F} on $[U/G]$ corresponds to a G -equivariant sheaf \mathcal{F}_U on U , and $p_*\mathcal{F} = (\pi_*\mathcal{F}_U)^G$, where $\pi : U \rightarrow U/G$ is the quotient map.

Let \mathcal{L} be a line bundle on \mathcal{X} . Then, for $N \in \mathbb{N}$ sufficiently divisible, the pushforward $p_*(\mathcal{L}^{\otimes N})$ is a line bundle on X . We define $p_*^\mathbb{Q}\mathcal{L} = \frac{1}{N}p_*(\mathcal{L}^{\otimes N}) \in \text{Pic}(X) \otimes \mathbb{Q}$, and call $p_*^\mathbb{Q}\mathcal{L}$ the \mathbb{Q} -line bundle on X defined by \mathcal{L} .

Theorem 6.1. *Assume that the coarse moduli space X is an algebraic variety. If the \mathbb{Q} -line bundle $p_*^\mathbb{Q}\mathcal{L}$ on X is big and nef then $H^i(\mathcal{L}^\vee) = 0$ for $i < \dim \mathcal{X}$.*

Theorem 6.1 is proved by reducing to the following generalisation of the Kodaira vanishing theorem.

Theorem 6.2. [KM98, Thm. 2.70, p. 73] *Let X be a proper normal variety and Δ a \mathbb{Q} -divisor on X such that the pair (X, Δ) is Kawamata log terminal (klt). Let N be a \mathbb{Q} -Cartier Weil divisor on X such that $N \equiv M + \Delta$, where M is a big and nef \mathbb{Q} -Cartier \mathbb{Q} -divisor. Then $H^i(X, \mathcal{O}_X(-N)) = 0$ for $i < \dim X$.*

Proof of Thm. 6.1. Observe first that X is a normal variety with quotient singularities. Consider the sheaf $p_*(\mathcal{L}^\vee)$ on X . If the automorphism group of a general point of \mathcal{X} acts nontrivially on \mathcal{L} , then $p_*\mathcal{L}^\vee = 0$, and so $H^i(\mathcal{L}^\vee) = H^i(p_*\mathcal{L}^\vee) = 0$ for each i . Suppose now that the automorphism group of a general point acts trivially on \mathcal{L} . Then $p_*\mathcal{L}^\vee$ is a rank 1 reflexive sheaf on X . Write $p_*\mathcal{L}^\vee = \mathcal{O}_X(-N)$, where N is a Weil divisor on X . Let $n \in \mathbb{N}$ be such that $p_*(\mathcal{L}^{\otimes n})$ is a line bundle on X . Let M be a \mathbb{Q} -divisor corresponding to the \mathbb{Q} -line bundle $p_*^\mathbb{Q}\mathcal{L} = \frac{1}{n}p_*(\mathcal{L}^{\otimes n})$. There is a natural map $(p_*\mathcal{L}^\vee)^{\otimes n} \rightarrow p_*(\mathcal{L}^{\vee \otimes n})$, i.e., a map $\mathcal{O}_X(-nN) \rightarrow \mathcal{O}_X(-nM)$,

which is an isomorphism over the locus where p is étale. So $N \equiv M + \Delta$, where Δ is an effective \mathbb{Q} -divisor supported on the branch locus of p . Let D_1, \dots, D_r be the codimension 1 components of the branch locus. Let e_i be the ramification index at D_i , and a_i the age of the line bundle \mathcal{L}^\vee along D_i . That is, over a general point of D_i , and after removing the automorphism group of a general point of \mathcal{X} , the stack \mathcal{X} is of the form $[\mathbb{A}_x^1/\mu_{e_i}]$, where $\mu_{e_i} \ni \zeta : x \mapsto \zeta \cdot x$, and μ_{e_i} acts on the fibre of \mathcal{L}^\vee by the character $\zeta \mapsto \zeta^{-a_i}$, where $0 \leq a_i \leq e_i - 1$. We compute that $\Delta = \sum \frac{a_i}{e_i} D_i$.

We claim that (X, Δ) is klt. Let $\Delta' = \sum \frac{e_i - 1}{e_i} D_i$, then $K_{\mathcal{X}} = p^*(K_X + \Delta')$, and \mathcal{X} is smooth, so (X, Δ') is klt by [KM98, Prop. 5.20(4), p. 160]. Now $\Delta \leq \Delta'$ and X is \mathbb{Q} -factorial, so (X, Δ) is also klt. We deduce that $H^i(\mathcal{L}^\vee) = H^i(p_* \mathcal{L}^\vee) = H^i(\mathcal{O}_X(-N)) = 0$ for $i < \dim \mathcal{X}$ by Thm. 6.2. \square

References

- [AV02] D. Abramovich, A. Vistoli, Compactifying the space of stable maps, J. Amer. Math. Soc. 15 (2002), no. 1, 27–75.
- [Arakelov71] S. Arakelov, Families of algebraic curves with fixed degeneracies (Russian), Izv. Akad. Nauk SSSR Ser. Mat. 35 (1971), 1269–1293, English translation in Math. USSR-Izv. 5 (1971), 1277–1302.
- [DM69] P. Deligne, D. Mumford, The irreducibility of the space of curves of given genus, Inst. Hautes Études Sci. Publ. Math. 36 (1969), 75–109.
- [Friedman83] R. Friedman, Global smoothings of varieties with normal crossings, Ann. of Math. (2) 118 (1983), no. 1, 75–114.
- [HM98] J. Harris, I. Morrison, Moduli of curves, Grad. Texts in Math. 187, Springer (1998).
- [Hartshorne66] R. Hartshorne, Residues and duality (Lecture notes of a seminar on the work of A. Grothendieck), with an appendix by P. Deligne, Lecture Notes in Math. 20, Springer (1966).
- [Kapranov97] M. Kapranov, Deformations of moduli spaces, unpublished manuscript, 1997.
- [KatoF00] F. Kato, Log smooth deformation and moduli of log smooth curves, Internat. J. Math. 11 (2000), no. 2, 215–232.

- [KatoK89] K. Kato, Logarithmic structures of Fontaine-Illusie, in Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988), 191–224, Johns Hopkins Univ. Press (1989).
- [Keel99] S. Keel, Basepoint freeness for nef and big line bundles in positive characteristic, *Ann. of Math. (2)* 149 (1999), no. 1, 253–286.
- [Knudsen83a] F. Knudsen, The projectivity of the moduli space of stable curves II : The stacks $M_{g,n}$, *Math. Scand.* 52 (1983), no. 2, 161–199.
- [Knudsen83b] F. Knudsen, The projectivity of the moduli space of stable curves III : The line bundles on $M_{g,n}$, and a proof of the projectivity of $\overline{M}_{g,n}$ in characteristic 0, *Math. Scand.* 52 (1983), no. 2, 200–212.
- [KM98] J. Kollár, S. Mori, Birational geometry of algebraic varieties, Cambridge Tracts in Math. 134, C.U.P. (1998).
- [Olsson01] M. Olsson, Log algebraic stacks and moduli of log schemes, PhD thesis, UC Berkeley (2001).
- [Siu80] Y-T. Siu, The complex-analyticity of harmonic maps and the strong rigidity of compact Kähler manifolds, *Ann. of Math. (2)* 112 (1980), no. 1, 73–111.
- [Voisin02] C. Voisin, Hodge theory and complex algebraic geometry I, Cambridge Stud. Adv. Math. 76, C.U.P. (2002).

Paul Hacking, Department of Mathematics, Yale University, PO Box 208283,
New Haven, CT 06520; paul.hacking@yale.edu