# A FEW REMARKS ON MIXING PROPERTIES OF $C^*$ -DYNAMICAL SYSTEMS

## FARRUKH MUKHAMEDOV AND SEYIT TEMIR

ABSTRACT. We consider strictly ergodic and strictly weak mixing  $C^*$ -dynamical systems. We prove that the system is strictly weak mixing if and only if its tensor product is strictly ergodic, moreover strictly weak mixing too. We also investigate some other mixing properties of the system.

Mathematics Subject Classification: 46L35, 46L55, 46L51, 28D05 60J99. Key words: strictly ergodic, strictly weak mixing, C\*-dynamical system.

## 1. Introduction

It is known [15],[13] that a notion of mixing for dynamical systems plays an important role in quantum statistical mechanics. A lot of papers (see, [5], [6],[9],[10],[16]) were devoted to the investigations of mixing properties of dynamical systems. Very recently in [11] certain relations between ergodicity, weak mixing and uniformly weak mixing conditions of  $C^*$ -dynamical systems have been investigated. It is known [16],[8] that strict ergodicity of a dynamical system is stronger than ergodicity. Therefore, it is natural to ask, how this notion is related with mixing conditions. The object of this paper is to investigate this question. Namely, we are going to consider strictly ergodic and strictly weak mixing  $C^*$ -dynamical systems. The paper organized as follows. In section 2 we recall some preliminaries on  $C^*$ -algebras and dynamical systems. Section 3 is devoted to the characterization of strictly ergodic  $C^*$ -dynamical systems. In the last section 4 we prove that the system is strictly weak mixing if and only if its tensor product is so. We also introduce a notion of  $\phi$ -ergodicity and compare it with known mixing conditions.

# 2. Preliminaries

In this section we recall some preliminaries concerning  $C^*$ -dynamical systems.

Let  $\mathfrak{A}$  be a  $C^*$ -algebra with unit  $\mathbf{1}$ . An element  $x \in \mathfrak{A}$  is called self-adjoint (resp. positive) if  $x = x^*$  (resp. there is an element  $y \in \mathfrak{A}$  such that  $x = y^*y$ ). The set of all self-adjoint (resp. positive) element will be denoted by  $\mathfrak{A}_{sa}$  (resp.  $\mathfrak{A}_+$ ). By  $\mathfrak{A}^*$  we denote the conjugate space to  $\mathfrak{A}$ . A linear functional  $\varphi \in \mathfrak{A}^*$  is called Hermitian if  $\varphi(x^*) = \overline{\varphi(x)}$  for every  $x \in \mathfrak{A}$ . A Hermitian functional  $\varphi$  is called positive if  $\varphi(x^*x) \geq 0$  for every  $x \in \mathfrak{A}$ . A positive functional  $\varphi$  is said to be a state if  $\varphi(\mathbf{1}) = 1$ . By S (resp.  $\mathfrak{A}_h^*$ ) we denote the set of all states (resp. Hermitian functionals) on  $\mathfrak{A}$ . Let  $\mathfrak{A} \odot \mathfrak{A}$  be the algebraic tensor product of  $\mathfrak{A}$ . By  $\mathfrak{A} \otimes \mathfrak{A}$  we denote a completion of  $\mathfrak{A} \odot \mathfrak{A}$  with respect to the minimal  $C^*$ -tensor norm on  $\mathfrak{A} \odot \mathfrak{A}$ . The set of all states on  $\mathfrak{A} \otimes \mathfrak{A}$  we denote by  $S^2$ . A linear operator  $T: \mathfrak{A} \mapsto \mathfrak{A}$  is called positive if  $Tx \geq 0$  whenever  $x \geq 0$ . A positive linear operator T is called a Markov operator if  $T = \mathbb{A}$ . A pair  $(\mathfrak{A}, T)$  consisting of a  $C^*$ -algebra  $\mathfrak{A}$  and a Markov operator  $T: \mathfrak{A} \mapsto \mathfrak{A}$ , is called a  $C^*$ -dynamical system. In the sequel, we will call any triplet  $(\mathfrak{A}, \varphi, T)$  consisting of a  $C^*$ -algebra  $\mathfrak{A}$ , a state  $\varphi$  on  $\mathfrak{A}$  and a Markov operator  $T: \mathfrak{A} \mapsto \mathfrak{A}$  with  $\varphi \circ T = \varphi$ , that is a dynamical system with an invariant state, a state preserving  $C^*$ -dynamical system. A state preserving  $C^*$ -dynamical system is a non-commutative  $C^*$ -probability space  $(\mathfrak{A}, \varphi)$  (see [4]) together with a Markov operator T of  $\mathfrak{A}$ 

preserving the non-commutative probability  $\varphi$ . We say that the state preserving  $C^*$ -dynamical system  $(\mathfrak{A}, \varphi, T)$  is ergodic (respectively, weakly mixing, strictly weak mixing) with respect to  $\varphi$  if

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} (\varphi(yT^k(x)) - \varphi(y)\varphi(x)) = 0, \text{ for all } x, y \in \mathfrak{A}.$$
 (2.1)

(respectively,

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} |\varphi(yT^k(x)) - \varphi(y)\varphi(x)| = 0, \text{ for all } x, y \in \mathfrak{A},$$
(2.2)

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^k(x)) - \varphi(x)| = 0, \text{ for all } x \in \mathfrak{A}, \psi \in S.)$$
 (2.3)

The state preserving  $C^*$ -dynamical system  $(\mathfrak{A}, \varphi, T)$  is called *strictly ergodic* with respect to  $\varphi$  if  $\varphi$  is the unique invariant state under T.

Given a  $C^*$ -algebra  $\mathfrak{A}$ , by  $M_n(\mathfrak{A})$  we denote the set of all  $n \times n$ -matrices  $a = (a_{ij})$  with entries  $a_{ij}$  in  $\mathfrak{A}$ . Recall that a linear mapping  $T : \mathfrak{A} \mapsto \mathfrak{A}$  is called n- positive if the linear operator  $T_n : M_n(\mathfrak{A}) \mapsto M_n(\mathfrak{A})$  given by  $T_n(a_{ij}) = (T(a_{ij}))$  is positive. If T is n-positive for all n then T is said to be completely positive. It is known [14] that if T is a completely positive map, then the linear operator  $T \otimes T : \mathfrak{A} \otimes \mathfrak{A} \mapsto \mathfrak{A} \otimes \mathfrak{A}$  defined by  $(T \otimes T)(x \otimes y) = Tx \otimes Ty$  is also completely positive.

# 3. Strictly ergodic dynamical systems

In this section we are going to characterize strictly ergodic  $C^*$ -dynamical systems. To do it we need the following

**Lemma 3.1.** Let  $(\mathfrak{A}, \varphi, T)$  be strictly ergodic. If  $h \in \mathfrak{A}^*$  is invariant with respect to T, i.e. h(Tx) = h(x) for all  $x \in \mathfrak{A}$ , then there is a number  $\lambda \in \mathbb{C}$  such that  $h = \lambda \varphi$ .

*Proof.* Let us first assume that h is positive, then  $\psi = h/h(\mathbb{1})$  is a state. According to the strict ergodicity of  $(\mathfrak{A}, \varphi, T)$  we have  $\psi = \varphi$ , which implies that  $h = h(\mathbb{1})\varphi$ . Now let h be a Hermitian functional. Then there is a unique Jordan decomposition [14] of h such that

$$h = h_{+} - h_{-}, \quad ||h||_{1} = ||h_{+}||_{1} + ||h_{-}||_{1},$$
 (3.1)

where  $\|\cdot\|_1$  is the norm on  $\mathfrak{A}^*$ . The invariance of h implies that

$$h \circ T = h_{+} \circ T - h_{-} \circ T = h_{+} - h_{-}.$$

Using  $||h_+ \circ T||_1 = h_+(\mathbb{1}) = ||h_+||_1$ , similarly  $||h_+ \circ T||_1 = ||h_+||_1$ , from uniqueness of the decomposition we find  $h_+ \circ T = h_+$  and  $h_- \circ T = h_-$ . Therefore, by the previous argument one gets  $h = \lambda \varphi$ . If h is an arbitrary functional, then there are Hermitian functionals  $h_1, h_2$  such that  $h = h_1 + ih_2$ . Again invariance of h implies that  $h_i \circ T = h_i$ , i = 1, 2. Consequently, we obtain that  $h = \lambda \varphi$ .

Now we are ready to formulate a criterion for the strict ergodicity of a dynamical system. The proof of the criterion is similar to the proof of Theorem 2, Ch.1, sec. 8 [8]. For the sake of completeness we will prove it.

**Theorem 3.2.** Let  $(\mathfrak{A}, \varphi, T)$  be a state preserving  $C^*$ -dynamical system. The following conditions are equivalent

(i)  $(\mathfrak{A}, \varphi, T)$  is strictly ergodic;

(ii) For every  $x \in \mathfrak{A}$  the following equality holds

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} T^k(x) = \varphi(x) \mathbf{1},$$

where convergence in norm of  $\mathfrak{A}$ ;

(iii) For every  $x \in \mathfrak{A}$  and  $\psi \in S$  the following equality holds

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} \psi(T^k(x)) = \varphi(x).$$

*Proof.* Let us consider the implication (i) $\Rightarrow$ (ii). It is clear that for every element of the form y = T(x) - x,  $x \in \mathfrak{A}$  we have

$$\left\| \frac{1}{n} \sum_{k=0}^{n-1} T^k(x) \right\| = \left\| \frac{1}{n} (T^n(x) - x) \right\| \le \frac{2}{n} \|x\| \to 0 \quad \text{as } n \to \infty.$$

So, as  $\varphi(y) = 0$  one gets

$$\lim_{n\to\infty}\frac{1}{n}\sum_{k=0}^{n-1}T^k(y)=\varphi(y)\mathbf{1}\!\!1.$$

It is evident that the set of elements of the form y = T(x) - x,  $x \in \mathfrak{A}$  forms a linear subspace of  $\mathfrak{A}$ . By  $\mathfrak{B}$  we denote the closure of this linear subspace. Set

$$\mathfrak{B}_0 = \{ x \in \mathfrak{A} : \ \varphi(x) = 0 \}.$$

It is clear that  $\mathfrak{B} \subseteq \mathfrak{B}_0$ . To show  $\mathfrak{B} = \mathfrak{B}_0$  assume that  $\mathfrak{B} \neq \mathfrak{B}_0$ , this means that there is an element  $x_0 \in \mathfrak{B}_0$  such that  $x_0 \notin \mathfrak{B}$ . Then according to the Hahn-Banach theorem there is a functional  $h \in \mathfrak{A}^*$  such that  $h \upharpoonright \mathfrak{B} = 0$  and  $h(x_0) = 1$ . The condition  $h \upharpoonright \mathfrak{B} = 0$  implies that h is invariant with respect to T. Therefore Lemma 3.1 yields that  $h = \lambda \varphi$ , which contradicts to  $\varphi(x_0) = 0$ . Hence  $\mathfrak{B} = \mathfrak{B}_0$ .

Let  $y \in \mathfrak{B}_0$ . Then for an arbitrary  $\epsilon > 0$  we can find  $y_{\epsilon} = T(x_{\epsilon}) - x_{\epsilon}$  such that  $||y - y_{\epsilon}|| < \epsilon/2$ . According to the following equality

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} T^k(y_{\epsilon}) = 0$$

there exists  $n_0 \in \mathbb{N}$  such that  $\left\| \frac{1}{n} \sum_{k=0}^{n-1} T^k(y_{\epsilon}) \right\| < \epsilon/2$  for all  $n \geq n_0$ . Hence, we have

$$\left\| \frac{1}{n} \sum_{k=0}^{n-1} T^k(y) \right\| \leq \left\| \frac{1}{n} \sum_{k=0}^{n-1} T^k(y - y_{\epsilon}) \right\| + \left\| \frac{1}{n} \sum_{k=0}^{n-1} T^k(y_{\epsilon}) \right\|$$

$$\leq \|y - y_{\epsilon}\| + \epsilon/2 < \epsilon \quad \text{for all} \quad n \geq n_0.$$

So,

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} T^k(y) = \varphi(y) \mathbf{1}$$

is valid for every  $y \in \mathfrak{B}_0$ .

Now let  $x \in \mathfrak{A}$ . Put  $y = x - \varphi(x) \mathbb{I}$ . Obviously that  $y \in \mathfrak{B}_0$ , and for y the last equality holds, whence we get the required relation.

The implication (ii) $\Rightarrow$ (iii) is evident. Let us prove (iii) $\Rightarrow$ (i). Assume that  $\nu$  is an invariant state with respect to T. According to the condition (iii) we find

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} \nu(T^k(x)) = \varphi(x)$$

for every  $x \in \mathfrak{A}$ . On the other hand, we have

$$\frac{1}{n} \sum_{k=0}^{n-1} \nu(T^k(x)) = \nu(x).$$

Whence  $\varphi = \nu$ . Thus the theorem is proved.

From this Theorem we immediately infer that strict ergodicity implies ergodicity of  $C^*$ -dynamical system. In the next section we will demonstrate an example of a dynamical system which is ergodic but not strictly ergodic. We mention that from Theorem 3.2 one gets that strict weak mixing trivially implies strict ergodicity.

# 4. Strictly weak mixing dynamical systems

In this section we are going to give a criterion characterizing strictly weak mixing  $C^*$ -dynamical systems.

Set

$$\mathfrak{A}_1^* = \{ g \in \mathfrak{A}^* : \|g\|_1 \le 1 \}, \quad \mathfrak{A}_{1,h}^* = \mathfrak{A}_1^* \cap \mathfrak{A}_h^*.$$

Before formulating a result we recall a well known fact (see for example [15])

**Lemma 4.1.** Let  $\{a_n\}$  be a bounded sequence of real numbers. Then the following are equivalent:

(i)

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=1}^{n} |a_k| = 0;$$

(ii) There exists a set  $J \subset \mathbb{N}$  of density zero (i.e.

$$\lim_{n\to\infty}\frac{\operatorname{cardinality}(J\cap[1,n])}{n}=0)$$

such that  $\lim_{n\to\infty} a_n = 0$  provided  $n \notin J$ ;

(iii)

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=1}^{n} |a_k|^2 = 0.$$

Now we are ready to formulate the following

**Theorem 4.2.** Let  $(\mathfrak{A}, \varphi, T)$  be a state preserving  $C^*$ -dynamical system and T be a completely positive map. The following conditions are equivalent:

- (i)  $(\mathfrak{A}, \varphi, T)$  is strictly weak mixing;
- (ii) The state preserving  $C^*$ -dynamical system  $(\mathfrak{A} \otimes \mathfrak{A}, \varphi \otimes \varphi, T \otimes T)$  is strictly weak mixing;
- (iii) The state preserving  $C^*$ -dynamical system  $(\mathfrak{A} \otimes \mathfrak{A}, \varphi \otimes \varphi, T \otimes T)$  is strictly ergodic;
- (iv) For every  $x \in \mathfrak{A}$  the following equality holds

$$\lim_{n \to \infty} \sup_{\psi \in \mathfrak{A}_1^*} \frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^k(x)) - \psi(\mathbf{1})\varphi(x)| = 0.$$

(v) For every  $x \in \mathfrak{A}$  and  $\psi \in \mathfrak{A}^*$  the following equality holds

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^k(x)) - \psi(\mathbf{1})\varphi(x)| = 0.$$
 (4.1)

*Proof.* Consider the implication (i) $\Rightarrow$ (ii). Recall that complete positivity of T implies that  $T \otimes T$  is so. It is clear that the state  $\varphi \otimes \varphi$  is invariant with respect to  $T \otimes T$ .

Let  $\psi, \phi \in S$  be arbitrary states and  $x, y \in \ker \varphi$ . Then according to (i) we have

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^k(x))| = 0, \quad \lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} |\phi(T^k(y))| = 0.$$

So according to Lemma 4.1 there exist two subsets  $J_1, J_2 \subset \mathbb{N}$  of density zero such that

$$\lim_{n\to\infty, n\notin J_1} |\psi(T^k(x))| = 0, \qquad \lim_{n\to\infty, n\notin J_2} |\phi(T^k(y))| = 0.$$

Then for the set  $J = J_1 \cup J_2$  we have

$$\lim_{n \to \infty, n \notin J} |\psi(T^k(x))\phi(T^k(y))| = 0,$$

and hence again using Lemma 4.1 one gets that

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^k(x))\phi(T^k(y))| = 0.$$

Thus,

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} |\psi \otimes \phi(T^k \otimes T^k(x \otimes y))| = 0.$$

By G we denote the convex hull of the set  $\{\psi \otimes \phi : \psi, \phi \in S\}$ . It is clear that the  $\|\cdot\|_1$ -closure of G is  $S^2$ . Therefore given  $\epsilon > 0$  and  $\omega \in S^2$  there is  $\zeta \in G$  such that  $\|\omega - \zeta\|_1 < \epsilon$ . For  $\zeta$  there is  $n_0 \in \mathbb{N}$  such that

$$\frac{1}{n} \sum_{k=0}^{n-1} |\zeta(T^k \otimes T^k(x \otimes y))| < \epsilon \quad \text{for all} \quad n \ge n_0.$$

Consequently,

$$\frac{1}{n} \sum_{k=0}^{n-1} |\omega(T^k \otimes T^k(x \otimes y))| \leq \frac{1}{n} \sum_{k=0}^{n-1} |(\omega - \zeta)(T^k \otimes T^k(x \otimes y))| 
+ \frac{1}{n} \sum_{k=0}^{n-1} |\zeta(T^k \otimes T^k(x \otimes y))| 
\leq \|\omega - \zeta\|_1 \|x \otimes y\| + \epsilon 
< \epsilon(\|x \otimes y\| + 1)$$
(4.2)

for all  $n \geq n_0$ .

Let  $x, y \in \mathfrak{A}$ . Denote  $x^0 = x - \varphi(x) \mathbb{I}$ ,  $y^0 = y - \varphi(y) \mathbb{I}$ . It is clear that  $x^0, y^0 \in \ker \varphi$ . By means of (4.2), for every  $\omega \in S^2$  we have

$$\frac{1}{n} \sum_{k=0}^{n-1} |\omega(T^k \otimes T^k(x^0 \otimes y^0))| < \epsilon \quad \text{for all } n \ge n_1.$$
 (4.3)

Denote  $\omega_1(x) = \omega(x \otimes \mathbb{1})$ ,  $\omega_2(x) = \omega(\mathbb{1} \otimes x)$ ,  $x \in \mathfrak{A}$ . Then according to the condition (i) there exist  $N_1, N_2 \in \mathbb{N}$  such that

$$\frac{1}{n} \sum_{k=0}^{n-1} |\omega_1(T^k(x)) - \varphi(x)| < \epsilon \quad \text{for all} \quad n \ge N_1,$$

$$\frac{1}{n} \sum_{k=0}^{n-1} |\omega_2(T^k(y)) - \varphi(y)| < \epsilon \quad \text{for all} \quad n \ge N_2.$$
(4.4)

Now using (4.3) and (4.4) we find

$$\frac{1}{n} \sum_{k=0}^{n-1} |\omega(T^k \otimes T^k(x \otimes y)) - \varphi(x)\varphi(y)| \leq |\varphi(y)| \left(\frac{1}{n} \sum_{k=0}^{n-1} |\omega_1(T^k(x)) - \varphi(x)|\right) + |\varphi(x)| \left(\frac{1}{n} \sum_{k=0}^{n-1} |\omega_2(T^k(y)) - \varphi(y)|\right) + \frac{1}{n} \sum_{k=0}^{n-1} |\omega(T^k \otimes T^k(x^0 \otimes y^0))| < \epsilon(|\varphi(x)| + |\varphi(y)| + 1) \tag{4.5}$$

for all  $n \ge \max\{n_1, N_1, N_2\}$ .

Now let  $z \in \mathfrak{A} \otimes \mathfrak{A}$ . Then there exists an element  $z_{\epsilon} \in \mathfrak{A} \odot \mathfrak{A}$  such that

$$||z-z_{\epsilon}||<\epsilon.$$

It follows from (4.5) that

$$\frac{1}{n}\sum_{k=0}^{n-1}|\omega(T^k\otimes T^k(z_{\epsilon}))-\varphi\otimes\varphi(z_{\epsilon})|<\epsilon$$

for all  $n \geq n_{\epsilon}$ . Therefore, we obtain

$$\frac{1}{n} \sum_{k=0}^{n-1} |\omega(T^k \otimes T^k(z)) - \varphi \otimes \varphi(z)| \leq \frac{1}{n} \sum_{k=0}^{n-1} |\omega(T^k \otimes T^k(z - z_{\epsilon}))| 
+ \frac{1}{n} \sum_{k=0}^{n-1} |\omega(T^k \otimes T^k(z_{\epsilon})) - \varphi \otimes \varphi(z_{\epsilon})| 
+ |\varphi \otimes \varphi(z_{\epsilon} - z))| 
\leq \epsilon + 2 ||z - z_{\epsilon}|| < 3\epsilon$$

for all  $n \geq n_{\epsilon}$ . The last relation implies that  $(\mathfrak{A} \otimes \mathfrak{A}, \varphi \otimes \varphi, T \otimes T)$  is strictly weak mixing.

The implication (ii) $\Rightarrow$ (iii) is obvious. Let us prove the implication (iii) $\Rightarrow$ (iv). Let  $(\mathfrak{A} \otimes \mathfrak{A}, \varphi \otimes \varphi, T \otimes T)$  be strictly ergodic. Let  $x \in \ker \varphi, x = x^*$ . Given  $\epsilon > 0$ , strict ergodicity of the dynamical system (see Theorem 3.2) implies that there is  $n_{0,x} \in \mathbb{N}$  such that

$$\left\| \frac{1}{n} \sum_{k=0}^{n-1} T^k \otimes T^k(x \otimes x) \right\| < \epsilon \quad \text{ for all } n \ge n_{0,x}.$$

Hence,

$$\left| \frac{1}{n} \sum_{k=0}^{n-1} \psi \otimes \psi(T^k \otimes T^k(x \otimes x)) \right| < \epsilon \quad \text{ for all } n \ge n_{0,x}, \forall \psi \in \mathfrak{A}_{1,h}^*.$$

As x is self-adjoint we get

$$\frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^k(x))|^2 < \epsilon \quad \text{for all } n \ge n_{0,x}, \forall \psi \in \mathfrak{A}_{1,h}^*.$$

According to Lemma 4.1 we infer that there is  $n_{1,x} \in \mathbb{N}$  such that

$$\frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^k(x))| < \epsilon \quad \text{ for all } n \ge n_{1,x}, \forall \psi \in \mathfrak{A}_{1,h}^*.$$

Consequently,

$$\sup_{\psi \in \mathfrak{A}_{1,h}^*} \frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^k(x))| < \epsilon \quad \text{for all} \quad n \ge n_{1,x}.$$
 (4.6)

Let  $x \in \ker \varphi$  be an arbitrary element. Then it can be represented as  $x = x_1 + ix_2$ , where  $x_1, x_2 \in \ker \varphi$ ,  $x_j^* = x_j$ , j = 1, 2. It then follows from (4.6) that

$$\sup_{\psi \in \mathfrak{A}_{1,h}^*} \frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^k(x))| < 2\epsilon \quad \text{ for all } n \ge n_{1,x} := \max\{n_{1,x_1}, n_{1,x_2}\}. \tag{4.7}$$

Let  $\psi \in \mathfrak{A}_{1}^{*}$ . Then  $\psi = \psi_{1} + i\psi_{2}$ , where  $\psi_{j} \in \mathfrak{A}_{1,h}^{*}, j = 1, 2$ . By means of (4.7) one yields

$$\sup_{\psi \in \mathfrak{A}_{1}^{*}} \frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^{k}(x))| \leq \sup_{\psi_{1} \in \mathfrak{A}_{1,h}^{*}} \frac{1}{n} \sum_{k=0}^{n-1} |\psi_{1}(T^{k}(x))| + \sup_{\psi_{2} \in \mathfrak{A}_{1,h}^{*}} \frac{1}{n} \sum_{k=0}^{n-1} |\psi_{2}(T^{k}(x))| < 4\epsilon, \quad \forall n \geq n_{1,x}. \tag{4.8}$$

Finally let  $x \in \mathfrak{A}$ . Then we have the last relation (4.8) for the element  $x^0 = x - \varphi(x) \mathbb{I}$ , which implies that

$$\lim_{n\to\infty} \sup_{\psi\in\mathfrak{A}_1^*} \frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^k(x)) - \psi(\mathbf{1})\varphi(x)| = 0.$$

So the implication (iii) $\Rightarrow$ (iv) is proved. The implications (iv) $\Rightarrow$ (v) $\Rightarrow$ (i) are obvious.

**Remark.** The implication (i) $\Leftrightarrow$ (v) can be proved directly using only positivity of the operator T. Indeed, it is enough to prove the implication (i) $\Rightarrow$ (v). Let  $x \in \ker \varphi$ . Assume that  $\psi \in \mathfrak{A}_h^*$  be a positive functional. Then  $\tilde{\psi}(x) = \frac{1}{\psi(\mathbf{I})}\psi(x)$  is a state. Hence, using (2.3) we get

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} |\tilde{\psi}(T^k(x))| = 0$$

which means

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^k(x))| = 0.$$
 (4.9)

Now let  $\psi \in \mathfrak{A}^*$  be an arbitrary functional, then it can be represented as  $\psi = \sum_{m=0}^{3} i^m \psi_m$ , where  $\psi_m \in \mathfrak{A}_h^*$ , (m = 0, 1, 2, 3) are positive functionals. By means of (4.9) we have

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} |\psi(T^k(x))| \le \lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} \sum_{m=0}^{3} |\psi_m(T^k(x))| = 0.$$

Using the same argument as in the finial part of the proof (iii) $\Rightarrow$ (iv) we obtain the required assertion. Therefore, if we take  $\psi(x) = \varphi(yx), x \in \mathfrak{A}$  in (v) we easily get (2.2), this means that strictly weak mixing implies weak mixing.

Using the same argument as the previous theorem one can prove the following

**Theorem 4.3.** Let  $(\mathfrak{A}, \varphi, T)$  be a state preserving  $C^*$ -dynamical system and T be a completely positive map. The following conditions are equivalent:

- (i)  $(\mathfrak{A}, \varphi, T)$  is weak mixing;
- (ii) The state preserving  $C^*$ -dynamical system  $(\mathfrak{A} \otimes \mathfrak{A}, \varphi \otimes \varphi, T \otimes T)$  is weak mixing.
- (iii) The state preserving  $C^*$ -dynamical system  $(\mathfrak{A} \otimes \mathfrak{A}, \varphi \otimes \varphi, T \otimes T)$  is ergodic.

**Remark.** It should be noted that Theorem 4.3 extends Theorem 6.3 in [16] to a  $C^*$ -algebra setting.

From Lemma 3.1 we infer that  $\varphi$  is a unique eigenvector with  $\lambda = 1$  eigenvalue of multiplicity one, for strictly ergodic dynamical system. Now what can we say about strictly weak mixing dynamical systems? We have the following

**Proposition 4.4.** Let  $(\mathfrak{A}, \varphi, T)$  be strictly weak mixing. If there exist a number  $\alpha \in \mathbb{C}$  with  $|\alpha| = 1$  and  $\alpha \neq 1$ , and  $h \in \mathfrak{A}^*$  such that

$$h \circ T = \alpha h,\tag{4.10}$$

then h = 0.

*Proof.* Assume that  $h \neq 0$ . Then  $h \neq \mu \varphi$  for all  $\mu \in \mathbb{C}$ . Then, using  $|\alpha| = 1$ , one gets

$$\frac{1}{n} \sum_{k=0}^{n-1} |h(T^k(x)) - h(\mathbf{I})\varphi(x)| = \frac{1}{n} \sum_{k=0}^{n-1} |\alpha^k h(x) - h(\mathbf{I})\varphi(x)|$$

$$> \left| |h(x)| - |h(\mathbf{I})\varphi(x)| \right| > 0 \quad \forall n \in \mathbb{N}$$

which contradicts to the strictly weak mixing condition.

Now we are going to give a concrete example of strictly mixing  $C^*$ -dynamical system.

**Example 1.** Let  $\mathfrak{A} = \ell^{\infty} = \{(x_n) : x_n \in \mathbb{C}, \sup |x_n| < \infty\}$ . Define an operator  $T : \ell^{\infty} \mapsto \ell^{\infty}$  by means of matrix  $(t_{ij})_{i,j\in\mathbb{N}}$  such that  $t_{ij} = 1/2^j$ ,  $i,j \geq 1$ . It is not hard to check that  $\varphi = (1/2, 1/2^2, \dots, 1/2^n, \dots)$  is an invariant state with respect to T. It is known from the Theory of Markov Chains with countable state space (see [12]) that  $\psi(T^n x)$  converges to  $\varphi$  in norm of  $\mathfrak{A}^*$  for every state  $\psi \in \mathfrak{A}^*$ . Consequently, T is strictly weak mixing.

The following example shows that strict ergodicity does not imply strict weak mixing.

**Example 2.** Let  $S^1 = \{z \in \mathbb{C} : |z| = 1\}$  and  $\lambda$  be the Lebesgue measure on  $S^1$  such that  $\lambda(S^1) = 1$ . Fix an element  $a = \exp(i2\pi\alpha)$ , where  $\alpha \in [0,1)$  is an irrational number. Define a transformation  $\tau: S^1 \mapsto S^1$  by  $\tau(z) = az$ . The measure induces a positive linear functional  $\varphi_{\lambda}(f) = \int_{S^1} f(z) d\lambda(z)$ 

such that  $\varphi_{\lambda}(\mathbf{1}) = 1$ . Consider a  $C^*$ -algebra  $\mathfrak{A} = C(S^1)$ , where  $C(S^1)$  is the space of all continuous functions on  $S^1$ . Now by means of  $\tau$  define a positive linear operator  $T_{\tau}: C(S^1) \to C(S^1)$  by

 $(T_{\tau}(f)(z)) = f(\tau(z))$  for all  $f \in C(S^1)$ . It is clear that  $(C(S^1), \varphi_{\lambda}, T_{\tau})$  is a state preserving  $C^*$ dynamical systems. Since  $\alpha$  is irrational, then according Theorem 2, Ch.3 [8] we find that the defined dynamical system is strictly ergodic. On the other hand, it is not strictly weak mixing. Indeed, take a linear functional  $h \in C(S^1)^*$  defined by  $h(f) = \int_{S^1} z f(z) d\lambda(z), f \in C(S^1)$ . Then we have

 $h(T_{\tau}(f)) = a^{-2}h(f)$  for all  $f \in C(S^1)$ . But this contradicts to Proposition 4.4. It should be noted that  $T_{\tau}$  is not also weak-mixing (see [15], Theorem 1.27).

Next example shows that strict ergodicity is stronger that ergodicity of  $C^*$ -dynamical system.

**Example 3.** Consider  $C^*$ -algebra  $\mathfrak{A} = \bigotimes M_2(\mathbb{C})$ , where  $M_2(\mathbb{C})$  is the algebra of  $2 \times 2$  matrices over the field  $\mathbb{C}$  of complex numbers. By  $e_{ij}^{(n)}$ ,  $n \in \mathbb{Z}$ ,  $i, j \in \{1, 2\}$  we denote the basis matrices of the algebra  $M_2(\mathbb{C})$  sited on nth place in the tensor product  $\bigotimes_{\mathbb{Z}} M_2(\mathbb{C})$ . The shift automorphism  $\theta : \mathfrak{A} \mapsto \mathfrak{A}$ 

of the algebra  $\mathfrak{A}$  is defined by  $\theta(e_{ij}^{(n)}) = e_{ij}^{(n+1)}$  for every  $n \in \mathbb{Z}$  and  $i, j \in \{1, 2\}$ . Let tr be the normalized trace on  $M_2(\mathbb{C})$ , i.e.  $tr(\mathbb{1}) = 1$ . Let  $\varphi_0(\cdot) = tr(\rho(\cdot))$  be a state on  $M_2(\mathbb{C})$ , where  $\rho \in M_2(\mathbb{C})$  is a positive operator such that  $tr(\rho) = 1$ . Such kind of  $\rho$  is called a density operator for  $\varphi_0$ . Now let  $K: M_2(\mathbb{C}) \mapsto M_2(\mathbb{C})$  be a completely positive Markov operator such that  $\varphi_0(x) = \varphi_0(Kx)$  for every  $x \in M_2(\mathbb{C})$ . On the algebra  $\mathfrak{A}_{[k,n]} = \bigotimes_{[k,n]} M_2(\mathbb{C})$  define the following linear

functional

$$\varphi_{[k,n]}(a_k \otimes a_{k+1} \otimes ... \otimes a_n) = \varphi_0(a_k K(a_{k+1}(\cdots K(a_n) \cdots))).$$

The defined functional  $\varphi_{[k,n]}$  is a state (see [2],[3]). If a compatibility condition holds

$$\varphi_{[k,n]} \upharpoonright \mathfrak{A}_{[k-1,n-1]} = \varphi_{[k-1,n-1]}$$

for the states  $\{\varphi_{[k,n]}\}$ , then there is a state  $\varphi_K$  on  $\mathfrak A$  such that  $\varphi_K \upharpoonright \mathfrak A_{[k,n]} = \varphi_{[k,n]}$  (see [1]), and  $\varphi$  is called a *Markov state*. We note that a more general definition of Markov state was given in [1],[2].

It is easy to see that the Markov state is invariant with respect to  $\theta$ . Define two Markov operators  $K_i: M_2(\mathbb{C}) \mapsto M_2(\mathbb{C}), i = 1, 2$  by

$$K_1 \left( \begin{array}{cc} a & b \\ c & d \end{array} \right) = \left( \begin{array}{cc} p_{11}a + p_{12}d & 0 \\ 0 & p_{21}a + p_{22}d \end{array} \right), \quad K_2 \left( \begin{array}{cc} a & b \\ c & d \end{array} \right) = \left( \begin{array}{cc} q_1a + q_2d & 0 \\ 0 & q_1a + q_2d \end{array} \right).$$

Here  $\mathbb{P} = (p_{ij})$  is a stochastic matrix, such that  $p_{ij} > 0$  for all i, j, and  $q_1 + q_2 = 1$ ,  $q_1, q_2 > 0$ . Now consider two states  $\varphi_{0,1}$  and  $\varphi_{0,2}$  defined on  $M_2(\mathbb{C})$ , whose density operators are given by

$$\rho_1 = \left(\begin{array}{cc} p_1 & 0 \\ 0 & p_2 \end{array}\right), \quad \rho_2 = \left(\begin{array}{cc} q_1 & 0 \\ 0 & q_2 \end{array}\right)$$

where  $\pi = (p_1, p_2)$  is a vector such that  $p_1 + p_2 = 1$ ,  $p_1 \ge 0$ ,  $p_2 \ge 0$  and  $\pi \mathbb{P} = \pi$ .

Note that for these operators and states the compatibility condition is satisfied, therefore there are two associated Markov states  $\varphi_{K_1}$  and  $\varphi_{K_2}$ .

Then  $(\mathfrak{A}, \varphi_{K_1}, \theta)$  and  $(\mathfrak{A}, \varphi_{K_2}, \theta)$  are weak mixing, and hance ergodic, state preserving  $C^*$ -dynamical systems (see [7], Th-ms 4.1 and 4.5). On the other hand, they are not strictly ergodic, because there exist two invariant states with respect to  $\theta$ .

**Remark.** From Examples 2 and 3 we conclude that weak mixing and strict ergodicity are not comparable. Therefore we may formulate the following

**Problem 4.5.** Let  $(\mathfrak{A}, \varphi, T)$  be a state preserving  $C^*$ -dynamical system. Are the following conditions equivalent?

(i)  $(\mathfrak{A}, \varphi, T)$  is weak mixing and strictly ergodic;

(ii)  $(\mathfrak{A}, \varphi, T)$  is strictly weak mixing.

Recall a state preserving dynamical system  $(\mathfrak{A}, \varphi, T)$  is called *exact* (see [9]) if, for each  $\psi \in \mathfrak{A}^*$ ,

$$\lim_{n \to \infty} \|\psi \circ T^n - \psi(\mathbf{1})\varphi\|_1 = 0$$

is valid, where  $\|\cdot\|_1$  is the norm in  $\mathfrak{A}^*$ . It is not hard to see that the exactness implies strict weak mixing. In [9] Luczak proved that exact and weak mixing conditions, for dynamical semi-groups on von Neumann algebras, are equivalent if and only if the von Neumann algebra is strongly  $\mathbb{R}_+$ -finite. Regarding this result we can formulate the following

**Problem 4.6.** Let  $(\mathfrak{A}, \varphi, T)$  be a state preserving  $C^*$ -dynamical system. When are the following conditions equivalent?

- (i)  $(\mathfrak{A}, \varphi, T)$  is exact;
- (ii)  $(\mathfrak{A}, \varphi, T)$  is strictly weak mixing.

Now by  $S_0$  denote the set of all continuous functionals  $f: \mathfrak{A}_+ \mapsto \mathbb{R}_+$  such that

$$f(\lambda x) = \lambda f(x)$$
 for all  $\lambda \in \mathbb{R}_+, x \in \mathfrak{A}_+,$ 

$$f(11) = 1.$$

Now we introduce a notion of  $\phi$ -ergodicity. Namely, a state preserving  $C^*$ -dynamical system  $(\mathfrak{A}, \varphi, T)$  is called  $\phi$ -ergodic if the equality

$$f(T(x)) = f(x)$$
 for all  $x \in \mathfrak{A}_+,$  (4.11)

where  $f \in S_0$ , implies that  $f(x) = \varphi(x)$  for all  $x \in \mathfrak{A}_+$ .

**Theorem 4.7.** Let  $(\mathfrak{A}, \varphi, T)$  be a state preserving  $C^*$ -dynamical system. Then for the conditions:

(i) For every  $x \in \mathfrak{A}$  the following equality holds

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} ||T^k(x) - \varphi(x) \mathbf{1}|| = 0;$$

(ii) For every  $x \in \mathfrak{A}$  the following equality holds

$$\lim_{n \to \infty} ||T^n(x) - \varphi(x)\mathbf{1}|| = 0;$$

- (iii)  $(\mathfrak{A}, \varphi, T)$  is  $\phi$ -ergodic;
- (iv) For every  $\psi \in S$  and  $x \in \mathfrak{A}$  the following equality holds

$$\lim_{n \to \infty} \psi(T^n(x)) = \varphi(x)$$

the following implications hold:  $(i)\Leftrightarrow(ii)\Rightarrow(iii)\Rightarrow(iv)$ 

*Proof.* The (i) $\Leftarrow$ (ii) implication is obvious. Consider the implication (i) $\Rightarrow$ (ii). Assume that  $x \in \ker \varphi$ , then we have

$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} ||T^k(x)|| = 0.$$
 (4.12)

On the other hand, one gets

$$||T^{n+1}(x)|| \le ||T^n(x)||$$

this means that the sequence  $\{\|T^nx\|\}$  is non-increasing. Hence, we have  $\lim_{n\to\infty} \|T^nx\| = \alpha$ . It follows from (4.12) that  $\alpha = 0$ . Let  $x \in \mathfrak{A}$ , then setting  $x^0 = x - \varphi(x) \mathbb{1}$  we find

$$\lim_{n \to \infty} ||T^n(x^0)|| = 0$$

which implies (ii).

(ii) $\Rightarrow$ (iii). Assume that (4.11) is valid for some  $f \in S_0$ . According to the condition (ii) we have

$$T^n(x) \to \varphi(x) \mathbf{1}$$
 as  $n \to \infty$ 

for  $x \in \mathfrak{A}_+$ , here the convergence in norm of  $\mathfrak{A}$ . By means of continuity of f one gets

$$f(T^n(x)) \to f(\varphi(x) \mathbb{1}) = \varphi(x)$$
 as  $n \to \infty$ .

On the other hand (4.11) implies that  $f(x) = \varphi(x), \forall x \in \mathfrak{A}_+$ . So  $(\mathfrak{A}, \varphi, T)$  is  $\phi$ -ergodic.

(iii) $\Rightarrow$ (iv). Let  $\psi \in S$ . Define functionals  $\hat{f}: \mathfrak{A}_+ \mapsto \mathbb{R}_+, \check{f}: \mathfrak{A}_+ \mapsto \mathbb{R}_+$  by

$$\hat{f}(x) = \limsup_{n \to \infty} \psi(T^n(x)), \quad x \in \mathfrak{A}_+,$$

$$\check{f}(x) = \liminf_{n \to \infty} \psi(T^n(x)), \quad x \in \mathfrak{A}_+.$$

It is clear that  $\hat{f}, \check{f} \in S_0$ . We have

$$\hat{f}(Tx) = \limsup_{n \to \infty} \psi(T^{n+1}(x)) = \hat{f}(x).$$

Similarly  $\check{f}(Tx) = \check{f}(x)$ . Hence,  $\phi$ -ergodicity of  $(\mathfrak{A}, \varphi, T)$  implies that

$$\hat{f}(x) = \varphi(x), \quad \check{f}(x) = \varphi(x) \quad \forall x \in \mathfrak{A}_+.$$

Consequently, we infer the existence of the following limit

$$\lim_{n \to \infty} \psi(T^n(x)) = \varphi(x), \quad x \in \mathfrak{A}_+.$$

Every  $x \in \mathfrak{A}$  can be written as  $x = \sum_{m=0}^{3} i^m x_m$ ,  $x_m \in \mathfrak{A}_+$ , (m = 0, 1, 2, 3), therefore by means of the last equality we get

$$\lim_{n \to \infty} \psi(T^n(x)) = \varphi(x), \quad x \in \mathfrak{A}.$$

This completes the proof.

From this theorem we have

**Problem 4.8.** Is the implication (iv) $\Rightarrow$ (iii) true?

It is clear that the exactness of a dynamical system implies the condition (iv). Therefore, it is natural to formulate the following

**Problem 4.9.** Let  $(\mathfrak{A}, \varphi, T)$  be a state preserving  $C^*$ -dynamical system. How are the following conditions related with each other?

- (i)  $(\mathfrak{A}, \varphi, T)$  is  $\phi$ -ergodic;
- (ii)  $(\mathfrak{A}, \varphi, T)$  is exact.

**Remark.** If  $C^*$ -algebra  $\mathfrak A$  is finite dimensional then all conditions in Theorem 4.7 are equivalent.

### ACKNOWLEDGEMENTS

The author (F.M.) acknowledges the TUBITAK-NATO PC- B programme for providing financial support, and Harran University for all facilities they provided and kind hospitality. Authors also thank with gratitude to a referee for useful observations and suggestions.

#### References

- [1] Accardi, L., On noncommutative Markov property, Funct. Anal. Appl. 9 (1975), 1–7.
- [2] Accardi, L. and Frigerio, R., Markovian cocycles, Proc. R. Ir. Acad. 83A (1983), 251–263.
- [3] Accardi, L. and Liebscher, V., Markovian KMS-states for one-dimensional spin chains, Infin. Dimens. Anal. Quantum Probab. Relat. Top. 2 (1999), 645–661.
- [4] Cuculescu, I. and Oprea, A.G., Noncommutative probability, Kluwer AP, Dordrecht, 1994. .
- [5] Frigerio, A. and Verri, M., Long-time asymptotic properties of dynamical semi-groups on  $W^*$ -algebras. Math. Z. **180**(1982), 275-286.
- [6] Fagnola, F. and Rebolledo R., Transience and recurrence of quantum Markov semi-groups. Probab. Theory Relat. Fields **126**(2003), 289-306.
- [7] Golodets, V.Ya. and Neshveyev, S.V., Non-Bernoullian quantum K-systems. Commun. Math. Phys. 195(1998), 213-232.
- [8] Kornfeld, I.P., Sinai, Ya. G. and Fomin, S.V., *Ergodic Theory*, Springer, Berlin–Heidelberg–New York, 1982
- [9] Luczak, A., Quantum dynamical semi-groups in strongly finite von Neumann algebras, Acta Math. Hungar. **92**(2001), 11-17.
- [10] Łuczak, A., Mixing and asymptotic properties of Markov semi-groups on von Neumann algebras, Math. Z. **235**(2000), 615-626.
- [11] Nicolescu, C., Ströh, A. and Zsidó, L., Noncommutative extension of classical and multiple recurrence theorems, J.Operator Theory, **50**(2003), 3-52.
- [12] Revuz D. Markov chains, North-Holland, Amsterdam 1984.
- [13] Ruelle, D., Statistical mechanics, Benjamin, Amsterdam-New York, 1969.
- [14] Takesaki, M., Theory of Operator algebras, I, Springer, Berlin-Heidelberg-New York, 1979.
- [15] Walters, P., An introduction to ergodic theory, Springer, Berlin-Heidelberg-New York, 1982.
- [16] Watanabe, S., Asymptotic behaviour and eigenvalues of dynamical semigroups on operator algebras, Jour. Math. Anal. Appl. 86(1982), 411-424.

FARRUKH MUKHAMEDOV, DEPARTMENT OF MECHANICS AND MATHEMATICS, NATIONAL UNIVERSITY OF UZBEKISTAN, VUZGORODOK, 700174, TASHKENT, UZBEKISTAN

E-mail address: far75m@yandex.ru

SEYIT TEMIR, DEPARTMENT OF MATHEMATICS, ART AND SCIENCE FACULTY, HARRAN UNIVERSITY, 63200, SANLIURFA, TURKEY

E-mail address: seyittemir67@hotmail.com