

PRE-COMPACT FAMILIES OF FINITE SETS OF INTEGERS AND WEAKLY NULL SEQUENCES IN BANACH SPACES

J. LOPEZ-ABAD AND S. TODORCEVIC

1. INTRODUCTION

In this paper we provide a somewhat general framework for studying weakly null sequences in Banach spaces using Ramsey theory of families of finite subsets of \mathbb{N} . Recall that the Ramsey theory on families of finite subsets of \mathbb{N} was developed in a series of papers of Nash-Williams in the 60's, a theory that is today naturally embedded in the more familiar infinite-dimensional Ramsey theory. The affinities between the infinite-dimensional Ramsey theory and some problems of the Banach space theory and especially those dealing with Schauder basic sequences have been explored for quite some time, starting perhaps with Farahat's proof of Rosenthal's ℓ_1 -theorem (see [12] and [17]). The Nash-Williams' theory though implicit in all this was not fully exploited in this context. We shall therefore try to demonstrate the usefulness of this theory by applying it to the classical problem of finding unconditional basic-subsequence of a given normalized weakly null sequence in some Banach space E . Recall that Bessaga and Pelczynski [6] have shown that every normalized weakly null sequence in a Banach space contains a subsequence forming a Schauder basis for its closed linear span. However, as demonstrated by Maurey and Rosenthal [14] there exist weakly null sequences in Banach spaces without unconditional basic subsequences. So one is left with a task of finding additional conditions on a given weakly null sequence guaranteeing the existence of unconditional subsequences. One such condition, given by Rosenthal himself around the time of publication of [14] (see also [17]), when put in a proper context reveals the connection with the Nash-Williams theory. It says that if a weakly null sequence (x_n) in some space of the form $\ell_\infty(\Gamma)$ is such that each x_n takes only the values 0 or 1, then (x_n) has an unconditional subsequence. To see the connection, consider the family

$$\mathcal{F} = \{\{n \in \mathbb{N} : x_n(\gamma) = 1\} : \gamma \in \Gamma\}$$

and note that \mathcal{F} is a pre-compact family of finite subsets of \mathbb{N} . As pointed out in [17], Rosenthal result is equivalent saying that there is an infinite subset M of \mathbb{N} such that the trace $\mathcal{F}[M] = \{t \cap M : t \in \mathcal{F}\}$ is *hereditary*, i.e., it is downwards closed under inclusion. On the other hand, recall that the basic notion of the Nash-Williams' theory is the notion of a *barrier*, which is simply a family \mathcal{F} of finite subsets of \mathbb{N} no two members of which are related under the inclusion which has the property that an arbitrary infinite subset of \mathbb{N} contains an initial segment in \mathcal{F} . Thus, in particular, \mathcal{F} is a pre-compact family of finite subsets of \mathbb{N} . Though the trace of an arbitrary pre-compact family might be hard to visualize, a trace $\mathcal{B}[M]$ of a barrier \mathcal{B} is easily to compute as it is simply equal to the downwards closure of its restriction $\mathcal{B} \upharpoonright M = \{t \in \mathcal{B} : t \subseteq M\}$. A further

2000 *Mathematics Subject Classification*. Primary 05D10, 46B20, 46A35; Secondary 03E05.

examination of Rosenthal's result shows that for every pre-compact family \mathcal{F} of finite subsets of \mathbb{N} there is an infinite set M such that the trace $\mathcal{F}[M]$ is actually equal to the downwards closure of a uniform barrier \mathcal{B} on M , or in other words that the \subseteq -maximal elements of $\mathcal{F}[M]$ form a uniform barrier on M . As it turns out, this fact holds considerably more information than the conclusion that $\mathcal{F}[M]$ is merely a hereditary family which is especially noticeable if one needs to perform further refinements of M while keeping track of the original family \mathcal{F} . This observation was the starting point of the research of this paper. Further extensions of Rosenthal's result required however analysis of not only pre-compact families of finite subsets of \mathbb{N} but also maps from barriers into pre-compact families of finite subsets of \mathbb{N} , or into weakly pre-compact subsets of c_0 . In fact, our more general results deal with partial maps from $\text{FIN} \times c_0$ into the reals whose domains project onto weakly pre-compact subsets of c_0 . Recall, that the equivalence relations associated to arbitrary maps defined on barriers have been characterized by Pudlak and Rödl [19]. Here we show that for certain maps one can say considerably more. For example, we show that for every mapping h from a barrier \mathcal{B} into a weakly pre-compact subset of c_0 and every $\varepsilon > 0$ there is an infinite subset M of \mathbb{N} such that $\sum_{i \in M \setminus s} |h(s)(i)| < \varepsilon$ for every $s \in \mathcal{B} \upharpoonright M$. This sort of a combinatorial result has shown to be quite useful in studying weakly-null sequences in Banach spaces. In fact, using a variation on this result, in Section 3 we show that if (x_n) is a normalized weakly-null sequence of $\ell_\infty(\Gamma)$ with the property that

$$\inf\{|x_n(\gamma)| : n \in \mathbb{N}, \gamma \in \Gamma\} = \delta > 0, \quad (1)$$

then (x_n) has an unconditional subsequence. More precisely, (x_n) has a subsequence which is $\delta/4$ -equivalent to the basis (e_i) of a \mathcal{F} -Schreier space associated with the downwards closure \mathcal{F} of a barrier on \mathbb{N} . (The original construction of Schreier uses the family $\{s : |s| \leq \min(s) + 1\}$ (see [8])). An exposition of this result appeared first in Part II of [3], a result that was proved independently from recent articles of Arvantakis [5] and Gasparis, Odell and Wahl [11] who use different approaches to prove a similar result.

Deeper applications of the combinatorics of finite sets of integers that we develop in Section 3 lead us to new forms of *near-unconditionality* and *convex-unconditionality* which we present in Section 4. Our near-unconditionality result says that for every normalized weakly-null sequence (x_n) and for every $\varepsilon > 0$ there is an infinite subset M of \mathbb{N} such that for every $(a_i)_{i \in M}$ such that $\sup_{i \in M} |a_i| \leq 1$ and every finite subset $s \subseteq M$,

$$\left\| \sum_{i \in s} a_i x_i \right\| \leq \frac{2 + \varepsilon}{\min_{i \in s} |a_i|} \left\| \sum_{i \in M} a_i x_i \right\|. \quad (2)$$

This should be compared with the well-known near-unconditionality result of Elton [9] (see also [17]). In the same section we prove our convex-unconditionality result which says that given a normalized weakly-null sequence (x_n) in some Banach space X then for every $\varepsilon > 0$ there is an infinite subset M of \mathbb{N} such that for every sequence of scalars $(a_i)_{i \in M}$ such that $\sup_{i \in M} |a_i| \leq 1$ and every subset $N \subseteq M$ such that $\sum_{i \in N} |a_i| \leq 1$,

$$\left\| \sum_{i \in N} a_i x_i \right\| \leq (4 + \varepsilon) \sqrt{\left\| \sum_{i \in M} a_i x_i \right\|}. \quad (3)$$

This in turn should be compared with the corresponding well-known convex-unconditionality result of Argyros, Mercourakis and Tsarpalis [2] originally obtained by dualizing the argument of Elton [9].

The Section 5 contains two kind of results about function spaces $C(K)$ for K is a countable compactum. The first part contains a version of c_0 -saturation for such Banach spaces $C(K)$ and this is given as a natural application of the theory of barriers developed in Section 3. (Recall, that the c_0 -saturation of Banach spaces $C(K)$ over countable compacta K is a result originally due to Pełczyński and Semadeni[18]. See also [4] and [11] for recent accounts on this result.) For example, we show that if $(x_i) \subseteq C(K)$ is a normalized weakly-null sequence, then there is $C \geq 1$, some infinite set M , some uniform barrier \mathcal{B} on M of rank at most the Cantor-Bendixson rank of K and some point-finite Lipschitz assignment $\mu : \mathcal{B} \rightarrow c_{00}^+$ with property $\text{supp } \mu(s) = s$ for every $s \in \mathcal{B}$, and such that for every block sequence (s_n) of elements of \mathcal{B} , the corresponding sequence $(x(s_n))$ of linear combinations,

$$x(s_n) = \sum_{i \in s_n} (\mu(s_n))(i) x_i,$$

is a normalized block sequence C -equivalent to the standard basis of c_0 .

The second part of Section 5 concerns the following natural measurement of unconditionality present in a given weakly null sequence (x_n) in a general Banach space E . Given a family \mathcal{F} of finite sets, we say that (x_n) is \mathcal{F} -unconditional with constant at most $C \geq 1$ iff for every sequence of scalars (a_n) ,

$$\sup_{s \in \mathcal{F}} \left\| \sum_{n \in s} a_n x_n \right\| \leq C \left\| \sum_{n \in \mathbb{N}} a_n x_n \right\|.$$

Thus, if for some infinite subset M of \mathbb{N} the trace $\mathcal{F}[M]$ contains the family of all finite subsets of M , the corresponding subsequence $(x_n)_{n \in M}$ is unconditional. Typically, one will not be able to find such a trace, so one is naturally led to study this notion when the family \mathcal{F} is pre-compact, or equivalently, when \mathcal{F} is a barrier. Since for every pair \mathcal{F}_0 and \mathcal{F}_1 of barriers on \mathbb{N} there is an infinite set M such that $\mathcal{F}_0[M] \subseteq \mathcal{F}_1[M]$ or $\mathcal{F}_1[M] \subseteq \mathcal{F}_0[M]$ and since the two alternatives depend on the ranks of \mathcal{F}_0 and \mathcal{F}_1 , one is also naturally led to the following measurement of unconditionality that refers only to a countable ordinal γ rather than a particular barrier of rank γ . Thus, we say that a normalized basic sequence (x_n) of a Banach space X is γ -unconditionally saturated with constant at most $C \geq 1$ if there is an γ -uniform barrier \mathcal{B} on \mathbb{N} such that for every infinite $M \subseteq \mathbb{N}$ there is infinite $N \subseteq M$ such that the corresponding subsequence $(x_n)_{n \in N}$ of (x_n) is $\overline{\mathcal{B} \upharpoonright N}$ -unconditional with constant at most C . (Here, $\overline{\mathcal{B} \upharpoonright N}$ denotes the topological closure of the restriction $\mathcal{B} \upharpoonright N$ which in turn is equal to the trace $\mathcal{B}[N]$, a pleasant property of any barrier.) It turns out that only indecomposable countable ordinals γ matter for this notion. We shall see, extending a result from [14], that every normalized basic sequence has a subsequence which is ω -unconditionally saturated, and that this cannot be extended further. For example, we show that for every indecomposable countable ordinal $\gamma > \omega$ there is a compactum K of Cantor-Bendixson rank $\gamma + 1$ and a normalized 1-basic weakly-null sequence $(x_n) \subseteq C(K)$ such that (x_n) is β -unconditionally saturated for all $\beta < \gamma$ but not γ -unconditionally saturated. More precisely, the summing basis of c_0 is finitely block-representable in every subsequence of (x_n) , and so in particular, no subsequence of (x_n) is unconditional.

2. PRELIMINARIES

We shall follow standard terminology and notation when dealing with sequences in Banach spaces (see [12]).

Definition 2.1. Let (x_i) be a sequence in a Banach space E .

- (a) (x_i) is called *weakly-null* iff for every $x^* \in E^*$, the sequence of scalars $(x^*(x_i))_i$ tends to 0.
- (b) (x_i) is called a *Schauder basis* of E iff for every $x \in E$ there is a unique sequence of scalars (a_i) such that $x = \sum_i a_i x_i$. This is equivalent to say that $x_i \neq 0$ for every i , the closed linear span of (x_i) is X , and there is a constant $\theta \geq 1$ such that for every sequence of scalars (a_i) , and every interval $I \subseteq \mathbb{N}$,

$$\left\| \sum_{i \in I} a_i x_i \right\| \leq \theta \left\| \sum_{i \in \mathbb{N}} a_i x_i \right\|. \quad (4)$$

- (c) (x_i) is called a *basic sequence* iff it is a Schauder basis of its closed linear span, i.e., $x_i \neq 0$ for every i , and there is $\theta \geq 1$ such that for every sequence of scalars (a_i) , and every interval $I \subseteq \mathbb{N}$, $\left\| \sum_{i \in I} a_i x_i \right\| \leq \theta \left\| \sum_{i \in \mathbb{N}} a_i x_i \right\|$. The infimum of those constants θ is called the *basic constant* of (x_i) .

- (d) (x_i) is called *θ -unconditional* ($\theta \geq 1$) iff for every sequence of scalars (a_i) , and every subset $A \subseteq \mathbb{N}$,

$$\left\| \sum_{i \in A} a_i x_i \right\| \leq \theta \left\| \sum_{i \in \mathbb{N}} a_i x_i \right\|. \quad (5)$$

(x_i) is called *unconditional* if it is θ -unconditional for some $\theta \geq 1$.

Given two basic sequences $(x_i)_{i \in M}$ and $(y_i)_{i \in N}$ of some Banach spaces E and F , indexed by the infinite sets $M, N \subseteq \mathbb{N}$, we say that $(x_i)_{i \in M} \subseteq E$ and $(y_i)_{i \in N} \subseteq F$ are θ -equivalent, denoted by $(x_i)_{i \in M} \sim_\theta (y_i)_{i \in N}$, if the order preserving bijection Φ between the two index-sets M and N lifts naturally to an isomorphism between the corresponding closed linear spans of these sequences sending x_i to $y_{\Phi(i)}$.

The *sequence of evaluation functionals* of c_0 is the biorthogonal sequence (p_i) of the natural basis (e_i) of c_0 , i.e. if $x = \sum_i a_i e_i \in c_0$, then $p_i(x) = a_i$. Note that weakly compact subsets K of c_0 are characterized by the property that every sequence in K has a pointwise converging subsequence to an element of K . We say that a subset X of c_0 is *weakly pre-compact* if its closure relative to the weak topology of c_0 is weakly compact. It is clear that for every weakly-compact subset $K \subseteq c_0$ the restrictions of evaluation mappings (p_i) to K is weakly-null in $C(K)$. The sequence of restrictions will also be denoted by (p_i) . Observe that (p_i) as a sequence in the Banach space $C(K)$ is a monotone basic sequence iff K is closed under restriction to initial intervals.

Let B_{E^*} denote the unit ball of the dual space E^* of E . Given a normalized weakly-null sequence (x_i) of a Banach space E , the set $K((x_i)) = \{(x^*(x_i)) : x^* \in B_{E^*}\}$ is clearly a symmetric, 1-bounded and weakly-compact subset of c_0 . Moreover, (x_i) is 1-equivalent to the evaluation mapping sequence $(p_i) \subseteq C(K((x_i)))$.

A classical result states that if (x_i) is a weakly-null sequence such that $\liminf \|x_i\| > 0$, then for every $\varepsilon > 0$ there is a subsequence $(x_i)_{i \in A}$ of (x_i) which is a basic sequence with basic constant $\leq 1 + \varepsilon$ (see [12]). In other words, for every $\varepsilon > 0$ there is some subsequence

$(x_i)_{i \in M}$ which is $1 + \varepsilon$ -equivalent to a monotone basic weakly-null sequence. Observe also that if (x_i) is a seminormalized monotone basic sequence, then for every interval I , and every sequence of scalars (a_i) , $\|\sum_{i \in I} a_i x_i\| \leq 2\|\sum_i a_i x_i\|$. So, if in addition (x_i) is normalized, then $\|\sum_i a_i x_i\| \geq (1/2)\|(a_i)\|_\infty$.

3. TOPOLOGY AND COMBINATORICS OF FAMILIES OF FINITE SETS

Let \mathbb{N} denote the set of all non-negative integers and let FIN denote the family of all finite sets of \mathbb{N} . The topology on FIN is the one induced from the Cantor cube ${}^{\mathbb{N}}2$ via the identification of subsets of \mathbb{N} with their characteristics function. Observe that this topology coincides with the one induced by c_0 with the same identification of finite sets and corresponding characteristic functions. Thus, we say that a family $\mathcal{F} \subseteq \text{FIN}$ is *compact* if it is a compact space under the induced topology. We say that $\mathcal{F} \subseteq \text{FIN}$ is *pre-compact* if its topological closure $\overline{\mathcal{F}}^{\text{top}}$ taken in the Cantor cube ${}^{\mathbb{N}}2$ consists only of finite subsets of \mathbb{N} . Given $X, Y \subseteq \mathbb{N}$ we write

- (1) $X < Y$ iff $\max X < \min Y$.
- (2) $X \sqsubseteq Y$ iff $X \subseteq Y$ and $X < Y \setminus X$.

A sequence (s_i) of finite sets of integers is called a *block sequence* iff $s_i < s_j$ for every $i < j$, and it is called a Δ -*sequence* iff there is some finite set s such that $s \sqsubseteq s_i$ ($i \in \mathbb{N}$) and $(s_i \setminus s)$ is a block sequence. The set s is called the *root* of (s_i) . Note that $s_i \rightarrow_i s$ iff for every subsequence of (s_i) has a Δ -subsequence with root s . It follows that the topological closure $\overline{\mathcal{F}}$ of a pre-compact family \mathcal{F} of finite subsets of \mathbb{N} is included in its downwards closure

$$\overline{\mathcal{F}}^{\sqsubseteq} = \{s \subseteq t : t \in \mathcal{F}\}$$

with respect to the inclusion relation and also included in its downwards closure

$$\overline{\mathcal{F}}^{\sqsubseteq\sqsubseteq} = \{s \sqsubseteq t : t \in \mathcal{F}\}$$

with respect to the relation \sqsubseteq . We say that a family $\mathcal{F} \subseteq \text{FIN}$ is \sqsubseteq -*hereditary* if $\mathcal{F} = \overline{\mathcal{F}}^{\sqsubseteq}$ and $\sqsubseteq\sqsubseteq$ -*hereditary* if $\mathcal{F} = \overline{\mathcal{F}}^{\sqsubseteq\sqsubseteq}$. The \sqsubseteq -hereditary families will simply be called *hereditary families*. We shall consider the following two restrictions of a given family \mathcal{F} of subsets of \mathbb{N} to a finite or infinite subset X of \mathbb{N}

$$\begin{aligned}\mathcal{F} \upharpoonright X &= \{s \in \mathcal{F} : s \subseteq X\}, \\ \mathcal{F}[X] &= \{s \cap X : s \in \mathcal{F}\}.\end{aligned}$$

Note the following simple facts regarding these notions.

Proposition 3.1. *Let $\mathcal{F} \subseteq \text{FIN}$.*

- (a) \mathcal{F} is pre-compact iff $\overline{\mathcal{F}}^{\sqsubseteq}$ is pre-compact iff $\overline{\mathcal{F}}$ is pre-compact.
- (b) Suppose further that \mathcal{F} is either \sqsubseteq -hereditary or $\sqsubseteq\sqsubseteq$ -hereditary. Then \mathcal{F} is compact iff it is pre-compact.
- (c) If \mathcal{F} is \sqsubseteq -hereditary then for every subset M of \mathbb{N} we have $\mathcal{F}[M] = \mathcal{F} \upharpoonright M$.
- (d) $\overline{\mathcal{F}}^{\sqsubseteq}[M] = \overline{\mathcal{F}[M]}^{\sqsubseteq}$.

PROOF. (d): Let $s \subseteq t \in \mathcal{F}$. Then $s \cap M \subseteq t \cap M$, and so $s \cap M \in \overline{\mathcal{F}[M]}^{\sqsubseteq}$. If $s \subseteq t \cap M$, with $t \in \mathcal{F}$, then $s \in \overline{\mathcal{F}}^{\sqsubseteq}$, and $s \subseteq M$, hence $s \in \overline{\mathcal{F}}^{\sqsubseteq}[M]$. \square

There are various ways to associate an ordinal index to a pre-compact family \mathcal{F} of finite subsets of \mathbb{N} . All these ordinal indices are based on the fact that for $n \in \mathbb{N}$, the index of the family

$$\mathcal{F}_{\{n\}} = \{s \in \text{FIN} : n < s, \{n\} \cup s \in \mathcal{F}\}$$

is smaller or equal from that of \mathcal{F} . For example, one may consider the *Cantor-Bendixson index* $r(\mathcal{F})$, the minimal ordinal α for which the iterated Cantor-Bendixson derivative $\partial^\alpha(\mathcal{F})$ is equal to \emptyset , then clearly $r(\mathcal{F}_{\{n\}}) \leq r(\mathcal{F})$ for all $n \in \mathbb{N}$. Recall that $\partial\mathcal{F}$ is the set of all proper accumulation points of \mathcal{F} and that $\partial^\alpha(\mathcal{F}) = \bigcap_{\xi < \alpha} \partial(\partial^\xi(\mathcal{F}))$. The rank is well defined since $\overline{\mathcal{F}}$ is countable and therefore a scattered compactum so the sequence $\partial^\xi(\mathcal{F})$ of iterated derivatives must vanish. Observe that if \mathcal{F} is a nonempty compact, then necessarily $r(\mathcal{F})$ is a successor ordinal.

We are now ready to introduce the basic combinatorial concepts of this section. For this we need the following piece of notation, where X and Y are subsets of \mathbb{N}

$$*_X = X \setminus \{\min X\} \text{ and } X/Y = \{m \in X : \max Y < m\}$$

The set $*X$ is called the *shift* of X . Given integer $n \in \mathbb{N}$, we write X/n to denote $X/\{n\} = \{m \in X : m > n\}$. The following notions have been introduced by Nash-Williams.

Definition 3.2. ([13]) Let $\mathcal{F} \subseteq \text{FIN}$.

- (1) \mathcal{F} is called *thin* if $s \not\sqsubseteq t$ for every pair s, t of distinct members of \mathcal{F} .
- (2) \mathcal{F} is called *Sperner* if $s \not\subseteq t$ for every pair $s \neq t \in \mathcal{F}$.
- (3) \mathcal{F} is called *Ramsey* if for every finite partition

$$\mathcal{F} = \mathcal{F}_0 \cup \dots \cup \mathcal{F}_k \tag{6}$$

there is an infinite set $M \subseteq \mathbb{N}$ such that at most one of the restrictions $\mathcal{F}_i \upharpoonright M$ is non-empty.

- (4) \mathcal{F} is called a *front* on M if $\mathcal{F} \subseteq \mathcal{P}(M)$, it is thin, and for every infinite $N \subseteq M$ there is some $s \in \mathcal{F}$ such that $s \subseteq N$.

- (5) \mathcal{F} is called a *barrier* on M if $\mathcal{F} \subseteq \mathcal{P}(M)$, it is Sperner, and for every infinite $N \subseteq M$ there is some $s \in \mathcal{F}$ such that $s \subseteq N$.

Clearly, every barrier is a front but not vice-versa. For example, the family $\mathbb{N}^{[k]}$ of all k -element subsets of \mathbb{N} is a barrier. The basic result of Nash-Williams [13] says that every front (and therefore every barrier) is Ramsey. Since as we will see soon there are many more barriers than those of the form $\mathbb{N}^{[k]}$ this is a far reaching generalization of the classical result of Ramsey. To see a typical application, let \mathcal{F} be a front on some infinite set M and consider its partition $\mathcal{F} = \mathcal{F}_0 \cup \mathcal{F}_1$, where \mathcal{F}_0 is the family of all \subseteq -minimal elements of \mathcal{F} . Since \mathcal{F} is Ramsey there is an infinite $N \subseteq M$ such that one of the restrictions $\mathcal{F}_i \upharpoonright M$ is empty. Note that $\mathcal{F}_1 \upharpoonright N$ must be empty. Since $\mathcal{F}_0 \upharpoonright N$ is clearly a Sperner family, it is a barrier on N . Thus we have shown that every front has a restriction that is a barrier. Since barrier are more pleasant to work with one might wonder why introducing the notion of front at all. The reason is that inductive constructions lead more naturally to fronts rather than barriers. To get an idea about this, it is instructive to consider the following notion introduced by Pudlak and Röd1.

Definition 3.3. ([19]) For a given countable ordinal α , the family \mathcal{F} is called α -uniform on M provided that:

- (a) $\alpha = 0$ implies $\mathcal{F} = \{\emptyset\}$,
- (b) $\alpha = \beta + 1$ implies that $\mathcal{F}_{\{n\}}$ is β -uniform on M/n ,
- (c) $\alpha > 0$ limit implies that there is an increasing sequence $\{\alpha_n\}_{n \in M}$ of ordinals converging to α such that $\mathcal{F}_{\{n\}}$ is α_n -uniform on M/n for all $n \in M$.

\mathcal{F} is called uniform on M if it is α -uniform on M for some countable ordinal α .

REMARK 3.4. (a) If \mathcal{F} is α -uniform (front, barrier) on M and $\Theta : M \rightarrow N$ is the unique order-preserving onto mapping between M and N , then $\Theta''\mathcal{F} = \{\Theta''s : s \in \mathcal{F}\}$ is α -uniform (front, barrier) on N .

- (b) If \mathcal{F} is a front on M , then $\overline{\mathcal{F}} = \overline{\mathcal{F}}^\square$.
- (c) If \mathcal{F} is uniform on M , then it is a front (though not necessarily a barrier) on M .
- (d) If \mathcal{F} is α -uniform on M then $\mathcal{F} \upharpoonright N$ is α -uniform on N for every $N \subseteq M$.
- (e) If \mathcal{F} is α -uniform on M , then $\partial^\alpha(\overline{\mathcal{F}}) = \{\emptyset\}$, hence $r(\mathcal{F}) = \alpha + 1$. (Hint: use that $\partial^\beta(\mathcal{F}_{\{n\}}) = (\partial^\beta(\mathcal{F}))_{\{n\}}$ for every β and every compact family \mathcal{F}).
- (f) An important example of a ω -uniform barrier on \mathbb{N} is the family $\mathcal{S} = \{s : |s| = \min(s) + 1\}$. We call \mathcal{S} a *Schreier barrier* since its downwards closure is commonly called a *Schreier family*. Note that unlike to the case of finite ranks there are many different ω -uniform families on \mathbb{N} . For example $\{s : |s| = 2\min(s) + 1\}$ is another such family.

The following result based on Nash-Williams' extension of Ramsey's theorem explains the relationship between the concepts introduced above (see [3] for proofs and fuller discussion).

Proposition 3.5. *The following are equivalent for a family \mathcal{F} of finite subsets of \mathbb{N} :*

- (a) \mathcal{F} is Ramsey.
- (b) There is an infinite $M \subseteq \mathbb{N}$ such that $\mathcal{F} \upharpoonright M$ is Sperner.
- (c) There is an infinite $M \subseteq \mathbb{N}$ such that $\mathcal{F} \upharpoonright M$ is either empty or uniform on M .
- (d) There is an infinite $M \subseteq \mathbb{N}$ such that $\mathcal{F} \upharpoonright M$ is either empty or a front on M .
- (e) There is an infinite $M \subseteq \mathbb{N}$ such that $\mathcal{F} \upharpoonright M$ is either empty or a barrier on M .
- (f) There is an infinite $M \subseteq \mathbb{N}$ such that $\mathcal{F} \upharpoonright M$ is thin.
- (g) There is an infinite $M \subseteq \mathbb{N}$ such that for every infinite $N \subseteq M$ the restriction $\mathcal{F} \upharpoonright N$ cannot be split into two disjoint families that are uniform on N . \square

We have also the following simple facts connecting these combinatorial notions with the topological concepts considered at the beginning of this section.

Proposition 3.6. *Fix a family $\mathcal{F} \subseteq \text{FIN}$.*

- (a) *If \mathcal{F} is a barrier on M then $\overline{\mathcal{F}}^\square = \overline{\mathcal{F}}^\square = \overline{\mathcal{F}}$, and hence $\overline{\mathcal{F}}^\square$ is a compact family.*
- (b) *If \mathcal{F} is a barrier on M then for every $N \subseteq M$, $\overline{\mathcal{F} \upharpoonright N}^\square = \overline{\mathcal{F}}^\square \upharpoonright N$.*
- (c) *Suppose that \mathcal{F} is a barrier on M . Then for every $N \subseteq M$ such that $M \setminus N$ is infinite we have that $\mathcal{F}[N] = \overline{\mathcal{F} \upharpoonright N}^\square$, and in particular $\mathcal{F}[N]$ is downwards closed.*
- (d) *A family $\mathcal{F} \subseteq M^{<\omega}$ is the topological closure of a barrier on M iff $\mathcal{F}^{\square-\max} = \mathcal{F}^{\square-\max}$ is a barrier on M .*

PROOF. (a): It is clear that $\overline{\mathcal{F}}^\square \supseteq \overline{\mathcal{F}}^\square \supseteq \overline{\mathcal{F}}$. Let us show that $\overline{\mathcal{F}}^\square \subseteq \overline{\mathcal{F}}$: Let $s \subsetneq t \in \mathcal{F}$, and fix an infinite subset N of M/s . Since \mathcal{F} is a barrier on M there is $u \in \mathcal{F} \upharpoonright N$. Then either

$u \sqsubseteq s$ or $s \sqsubseteq u$. The first case is impossible since it implies that $u \subsetneq t$ and both are elements of \mathcal{F} which is Sperner. So, the second alternative $s \sqsubseteq u$ holds. It is clear that now we can find Δ -sequence (u_k) of elements of \mathcal{F} with root s .

(b): It is clear that $\overline{\mathcal{F} \restriction N}^\subseteq \subseteq \overline{\mathcal{F}}^\subseteq \restriction N$. Now suppose that $s \in \overline{\mathcal{F}}^\subseteq \restriction N$. Let $t \in \mathcal{F}$ be such that $s \subseteq t$. Find $u \in \mathcal{F} \restriction N$ such that $u \sqsubseteq s \cup (N/s)$. If $u \sqsubseteq s \subseteq t$, then $u = s = t$, and hence $s \in \mathcal{F} \restriction N$. Otherwise $s \sqsubseteq u$, and hence $s \in \overline{\mathcal{F} \restriction N}^\subseteq$.

(c): Fix an infinite subset N of M such that $M \setminus N$ is infinite as well. Since \mathcal{B} is a barrier on N we obtain that $\overline{\mathcal{B} \restriction N}^\subseteq = \overline{\mathcal{B}} \restriction N^\subseteq$. Now let $s \sqsubseteq t \in \mathcal{B} \restriction N$, and consider the infinite subset $s \cup (M \setminus N)/t$ of M . Since \mathcal{B} is a barrier on M there is some $u \in \mathcal{B}$ such that $u \sqsubseteq s \cup (M \setminus N)/t$. Then $s \sqsubseteq u$ (use that otherwise $u \subsetneq s \subseteq t$ and $u, t \in \mathcal{B}$), $u \setminus s \subseteq (M \setminus N)/t$ and hence

$$u \cap M = s, \quad (7)$$

as desired. Now suppose that $s \in \mathcal{B}$, and let $t = s \cap N$. Consider the infinite subset $t \cup N/s$ of N . By similar reasons as above, there must be $u \in \mathcal{B} \restriction N$ such that $t \sqsubseteq u$, as desired.

(d) is not difficult to prove. We leave the details to the reader. \square

The next is a well known result. We extract its proof from [3].

Lemma 3.7. *Suppose that \mathcal{B} and \mathcal{C} are two barriers on M . Then there is some infinite $N \subseteq M$ such that either $\overline{\mathcal{B} \restriction N}^\subseteq \subseteq \overline{\mathcal{C} \restriction N}^\subseteq$ or else $\overline{\mathcal{C} \restriction N}^\subseteq \subseteq \overline{\mathcal{B} \restriction N}^\subseteq$*

PROOF. Let $\mathcal{B}_0 = \{s \in \mathcal{B} : s \in \overline{\mathcal{C}}\}$, $\mathcal{B}_1 = \mathcal{B} \setminus \mathcal{B}_0$, and in the same way $\mathcal{C}_0 = \{s \in \mathcal{C} : s \in \overline{\mathcal{B}}\}$, $\mathcal{C}_1 = \mathcal{C} \setminus \mathcal{C}_0$. By the Ramsey property of the uniform families, there is $N \subseteq M$ and $i, j \in \{0, 1\}$ such that $\mathcal{B} \restriction N = \mathcal{B}_i \restriction N$ and $\mathcal{C} \restriction N = \mathcal{C}_j \restriction N$. Let $s \in \mathcal{B} \restriction N$. Consider the infinite subset $P = s \cup (N/s) \subseteq N$, and let $t \in \mathcal{C} \restriction N$ be such that $t \sqsubseteq P$. If $s \sqsubseteq t$, then $i = 1$ and hence $\overline{\mathcal{B} \restriction N}^\subseteq \subseteq \overline{\mathcal{C} \restriction N}^\subseteq$. Otherwise, $t \sqsubseteq s$ and so $j = 1$ and $\overline{\mathcal{C} \restriction N}^\subseteq \subseteq \overline{\mathcal{B} \restriction N}^\subseteq$. \square

Corollary 3.8. *Suppose that \mathcal{B} and \mathcal{C} are respectively α and β -uniform some M , and suppose that $\alpha < \beta$. Then there is $N \subseteq M$ such that $\mathcal{B} \restriction N \subseteq \overline{\mathcal{C}}$.*

PROOF. It follows from Lemma 3.7 and Remark 3.4 (a) and (d). \square

In this kind of Ramsey theory one frequently performs diagonalisation arguments that can be formalized using the following notion.

Definition 3.9. An infinite sequence (M_i) of infinite subsets of \mathbb{N} is called a *fusion sequence of subsets of $M \subseteq \mathbb{N}$* if for all i :

- (a) $M_{i+1} \subseteq M_i \subseteq M$,
- (b) $\min M_i < \min M_{i+1}$.

The infinite set $M_\infty = \{m_i = \min M_i\}$ is called the *fusion* of the sequence (M_i) .

3.1. Mappings on barriers.

Lemma 3.10. *Suppose that \mathcal{B} is a uniform barrier on M , and suppose that $h : \mathcal{B} \rightarrow \text{FIN}$ is such that for every $s \in \mathcal{B}$, $h(s) \sqsubseteq s$. Then there is an infinite subset $N \subseteq M$ such that for every $s, t \in \mathcal{B} \restriction N$, if $h(s) \sqsubseteq t$ then $h(t) = h(s)$.*

PROOF. By the Ramsey property of \mathcal{B} we can find a fusion sequence (M_k) of subsets of M such that, setting $m_k = \min M_k$, then for every $k \in \mathbb{N}$, every $s \in \{m_0, \dots, m_{k-1}\}$ and every $t \in \mathcal{B} \restriction M_k$, if $h(t) = s$ then for every $u \in \mathcal{B} \restriction M_k$ $h(u) = s$. It is clear that $\{m_i\}$ has the desired property. \square

Corollary 3.11. *Suppose that \mathcal{B} is a uniform barrier on M , and suppose that $h : \mathcal{B} \rightarrow \text{FIN}$ is such that for every $s \in \mathcal{B}$, $h(s) \sqsubseteq s$. Then there is an infinite subset $N \subseteq M$ such that*

- (a) *for every $P \subseteq N$, $(h''(\mathcal{B} \restriction N)) \restriction P = h''(\mathcal{B} \restriction P)$.*
- (b) *$h''(\mathcal{B} \restriction N)$ is a uniform barrier on N .*
- (c) *For every $s \in \mathcal{B} \restriction N$ and every $t \in h''(\mathcal{B} \restriction N)$, if $t \sqsubseteq s$ then $h(s) = t$.*

PROOF. Let $N_0 \subseteq M$ given by the previous Lemma 3.10. We claim that N_0 satisfies (a) and (c): This last condition is clearly true. We show (a): the inverse inclusion is trivial to prove. Suppose now that $t \in (h''(\mathcal{B} \restriction N_0)) \restriction P$, let $s \in \mathcal{B} \restriction N_0$ be such that $h(s) = t$. Then consider the infinite set $t \cup (P/s) \subseteq P$. Find some $u \in \mathcal{B} \restriction P$ such that $u \sqsubseteq t \cup (P/s) \subseteq P$. Since \mathcal{B} is thin we obtain that $h(s) = t \sqsubseteq u$, so $h(u) = t$, as desired.

(b): Observe that from the conclusion of Lemma 3.10 it readily follows that $h''(\mathcal{B} \restriction N)$ is a front on N . So, by Proposition 3.5 and (a), there is some $N \subseteq N_0$ such that $(h''(\mathcal{B} \restriction N_0)) \restriction N = h''(\mathcal{B} \restriction N)$ is a uniform barrier on N . \square

Definition 3.12. Given $\mathcal{F} \subseteq \text{FIN}$, let

$$\begin{aligned} \mathcal{F}^{\sqsubseteq - \max} &= \{s \in \mathcal{F} : (\forall t \in \mathcal{F})(s \sqsubseteq t \rightarrow s = t)\} \\ \mathcal{F}^{\subseteq - \max} &= \{s \in \mathcal{F} : (\forall t \in \mathcal{F})(s \subseteq t \rightarrow s = t)\}. \end{aligned}$$

Lemma 3.13. *Suppose that $\mathcal{F} \subseteq \text{FIN}$ is compact and hereditary. Then there is an infinite set M such that $\mathcal{F} \restriction M$ is the closure of a uniform barrier on M , and therefore $(\mathcal{F} \restriction M)^{\sqsubseteq - \max} = (\mathcal{F} \restriction M)^{\subseteq - \max}$ is a uniform barrier on M .*

PROOF. Let \mathcal{B} be an arbitrary α -uniform barrier on \mathbb{N} , where $\alpha + 1 = r(\mathcal{F})$. Consider the coloring $c : \mathcal{B} \rightarrow \{0, 1\}$ defined for $s \in \mathcal{B}$ by $c(s) = 0$ iff there is some $t \in \mathcal{F}$ such that $s \subseteq t$. By the Ramsey property of \mathcal{B} there is some infinite set M such that c is constant on $\mathcal{B} \restriction M$ with value $i_0 \in \{0, 1\}$. We claim that $i_0 = 1$. Otherwise, since \mathcal{F} is hereditary, we have that $\overline{\mathcal{B}} \restriction M \subseteq \mathcal{F}$. But then $r(\mathcal{F}) \geq r(\overline{\mathcal{B}} \restriction M) = \alpha + 1 > r(\mathcal{F})$, a contradiction. Now observe that $i_0 = 1$ implies that every $t \in \mathcal{F} \restriction M$ is an initial part of some $s \in \mathcal{B} \restriction M$: Given $t \in \mathcal{F} \restriction M$ consider the infinite subset $N = t \cup (M/t)$ of M ; since $\mathcal{B} \restriction M$ is a barrier on M there is some $s \in \mathcal{B} \restriction N$. Hence $t \sqsubseteq s$, since otherwise $s \sqsubseteq t$, contradicting the fact that $i_0 = 1$. Now define $h : \mathcal{B} \restriction M \rightarrow \mathcal{F}$ by $h(s) = t \in \mathcal{F}$ iff t is the maximal initial part of s in \mathcal{F} . By Corollary 3.11 there is some $N \subseteq M$ such that $h''(\mathcal{B} \restriction N)$ is a uniform barrier on N . Observe that indeed $h''(\mathcal{B} \restriction N) = \mathcal{F} \restriction N$, so we are done. \square

We extend the previous result to an arbitrary pre-compact family \mathcal{F} of finite subsets of \mathbb{N} .

Theorem 3.14. *Suppose that $\mathcal{F} \subseteq \text{FIN}$ is pre-compact. Then there is an infinite set M such that $\mathcal{F}[M]$ is the closure of a uniform barrier on M , and hence $(\mathcal{F}[M])^{\sqsubseteq - \max} = (\mathcal{F}[M])^{\subseteq - \max}$ is a uniform barrier on M .*

PROOF. By Lemma 3.13 applied to the compact hereditary family $\mathcal{G} = \overline{\mathcal{F}}^\subseteq$ there is some M such that

$$(\mathcal{G} \upharpoonright M)^{\subseteq-\max} = (\mathcal{G}[M])^{\subseteq-\max} = ((\overline{\mathcal{F}}^\subseteq)[M])^{\subseteq-\max} = (\mathcal{F}[M])^{\subseteq-\max} \quad (8)$$

is a uniform barrier \mathcal{B} on M . Let $N \subseteq M$ be such that $M \setminus N$ is infinite. So, by Proposition 3.6 (b) and (c),

$$\begin{aligned} \mathcal{B}[N] &= (\mathcal{F}[M]^{\subseteq-\max})[N] \subseteq \mathcal{F}[N] \subseteq \mathcal{G}[N] = \mathcal{G} \upharpoonright N \subseteq (\overline{\mathcal{G}[M]^{\subseteq-\max}})^\subseteq \upharpoonright N = \\ &= \overline{\mathcal{B}}^\subseteq \upharpoonright N = \overline{\mathcal{B} \upharpoonright N}^\subseteq = \mathcal{B}[N], \end{aligned} \quad (9)$$

as required. \square

REMARK 3.15. (a) It follows that for every compact family \mathcal{F} there is an infinite set M such that $\mathcal{F}[M]$ is hereditary. As pointed out by Odell[17], this result is just another formulation of Rosenthal's result that every weakly-null sequence of characteristic functions of subsets of some index-set Γ has an unconditional subsequence.

(b) In general it is not possible to find for a given compact family \mathcal{F} an infinite set M such that $\mathcal{F}[M]$ is the closure of an α -uniform barrier, where $r(\mathcal{F}) = \alpha + 1$.

The following corollary generalizes the corresponding well-known result for compact hereditary families presented in [10].

Corollary 3.16. *Suppose that \mathcal{F}_0 and \mathcal{F}_1 are two pre-compact families. Then there is an infinite set M such that either $\mathcal{F}_0[M] \subseteq \mathcal{F}_1[M]$ or $\mathcal{F}_1[M] \subseteq \mathcal{F}_0[M]$.*

PROOF. Find an infinite set M such that for each $i = 0, 1$, the trace $\mathcal{F}_i[M]$ is equal to the closure of a uniform barrier \mathcal{B}_i on M . Apply now Lemma 3.7. \square

3.2. More on mappings on barriers. We begin by an extension of a result from [19].

Lemma 3.17. *Let \mathcal{B} be a uniform barrier on M , and suppose that $f : \mathcal{B} \rightarrow \text{FIN}$ is such that its range is a pre-compact family. Then there is some infinite subset $N \subseteq M$ such that $f(s) \cap N \subseteq s$ for every $s \in \mathcal{B} \upharpoonright N$.*

PROOF. Let $h : \mathcal{B} \rightarrow \text{FIN}$ be defined by $h(s) = f(s) \setminus s$ ($s \in \mathcal{B}$). It is clear that $h''\mathcal{B}$ is a pre-compact family, and, by definition, $h(s) \cap s = \emptyset$. We are going to show that there is some $N \subseteq M$ such that $h(s) \cap N = \emptyset$ for every $s \in \mathcal{B} \upharpoonright N$, that gives the desired conclusion for f . The proof is by induction on the rank of \mathcal{B} . For every $m \in M$, let $h_m : \mathcal{B}_{\{m\}} \rightarrow \text{FIN}$ be naturally defined by $h_m(s) = h(\{m\} \cup s)$ ($s \in \mathcal{B}_{\{m\}}$). It is clear that $h_m : \mathcal{B}_{\{m\}} \rightarrow \text{FIN}$ fulfills (a) and (b) above, so, by inductive hypothesis, we can find a fusion sequence $(M_k)_{k \in \mathbb{N}}$, $M_k = M$, and such that, setting $m_k = \min M_k$ ($k \in \mathbb{N}$), we have that for every $k \in \mathbb{N}$ and every $s \in \mathcal{B}_{\{m_k\}} \upharpoonright M_{k+1}$, $h_{m_k}(s_k) \cap M_{k+1} = \emptyset$. Let $M_\infty = \{m_k\}$. It is easy to check that for $s \in \mathcal{B} \upharpoonright M_\infty$, $h(s) \cap M_\infty \subseteq \{m_0, \dots, m_{k-1}\}$, where k is such that $m_k = \min s$. For $m \in M_\infty$, define $g_m : \mathcal{B}_{\{m\}} \upharpoonright M_\infty \rightarrow \mathcal{P}(M \cap \{0, \dots, m-1\})$ by $g_m(s) = h_m(s) \cap M_\infty$. Since the image of

g_m has only finitely many possibilities, we can find another fusion sequence (N_k) , $N_0 = M_\infty$, such that, setting $n_k = \min N_k$, for every k the mapping g_{n_k} is constant on $\mathcal{B}_{\{n_k\}} \upharpoonright N_{k+1}$ with value $s_{n_k} < n_k$. Let $N_\infty = \{n_k\}$. Notice that, by the properties of this last fusion sequence, for every $s \in \mathcal{B} \upharpoonright N_\infty$ we obtain that $h(s) \cap N_\infty \subseteq h(s) \cap M_\infty = s_{\min s}$. Since the range of h is a pre-compact family, there is some infinite set $I \subseteq N_\infty$ such that $(s_i)_{i \in I}$ is a Δ -sequence with root r . Take a thinner $N \subseteq I$ such that $N \cap \bigcup_{n \in N} s_n = \emptyset$. Then for every $s \in \mathcal{B} \upharpoonright N$ we have that $h(s) \cap N \subseteq s_{\min s} \cap N = \emptyset$, as desired. \square

We generalize the previous result.

Lemma 3.18. *Suppose that $\{\mathcal{B}_k\}_{k \in \mathbb{N}}$ is a collection of uniform barriers on M , and suppose that for every $k \in \mathbb{N}$ we have $h_k : \mathcal{B}_k \rightarrow \text{FIN}$ with pre-compact range. Then there is some infinite subset $N = \{n_i\}$ of M such that $h_k(s) \cap N \subseteq \{n_0, \dots, n_{k-1}\} \cup s$ for every $k \in \mathbb{N}$ and every $s \in \mathcal{B}_k$.*

PROOF. Again, we may assume that h_k , replacing it by $s \mapsto h_k(s) \setminus s$ if needed, has the property that for every k and every $s \in \mathcal{B}_k$, $h_k(s) \cap s = \emptyset$. Using previous Lemma 3.17 we can find a fusion sequence (M_k) of M such that, setting $m_k = \min M_k$ ($k \in \mathbb{N}$), we have that for every k and every $s \in \mathcal{B}_k \upharpoonright (\{m_0, \dots, m_k\} \cup M_{k+1})$, $h_k(s) \cap M_{k+1} = \emptyset$. It is clear that the fusion set $N = \{m_k\}$ fulfills the desired requirements. Let us explain how to find this fusion sequence. Suppose we have found $M_k \subseteq M_{k-1} \subseteq \dots \subseteq M_0$. For every $s \subseteq \{m_0, \dots, m_k\}$, let $h_{k,s} : (\mathcal{B}_k)_s \rightarrow \text{FIN}$ be naturally defined for $t \in (\mathcal{B}_k)_s$ by $h_{k,s}(t) = h_k(s \cup t)$. Using repeatedly Lemma 3.17 to each $h_{k,s}$ we can find $M_{k+1} \subseteq M_k$ with the property that for every $s \in \mathcal{B}_k \upharpoonright (\{m_0, \dots, m_k\} \cup M_{k+1})$, the intersection $h_k(s) \cap M_{k+1} = \emptyset$, as desired. \square

The previous results have consequences on mappings with domain a barrier and with range c_0 . We introduce the following natural extension to c_0 of pre-compactity.

Definition 3.19. We say that $K \subseteq c_0$ is *weakly-pre-compact* if its weak-closure \overline{K}^w is a weakly compact subset of c_0 . This is equivalent to say that every sequence $(\xi_n) \subseteq K$ has a weak convergent subsequence in c_0 .

Given $\xi \in c_0$, and $\varepsilon > 0$ we write $\text{supp}_\varepsilon \xi = \{i \in \text{supp } \xi : |\xi(i)| \geq \varepsilon\}$.

Proposition 3.20. *If $K \subseteq c_0$ is weakly-compact, then $\text{supp}_\varepsilon K \subseteq \text{FIN}$ is pre-compact for every $\varepsilon > 0$.* \square

PROOF. Otherwise there is some sequence $(\xi_i) \subseteq K$ such that $\text{supp}_\varepsilon \xi_i \rightarrow M$, M infinite. Then fix an accumulation point $\xi \in K$ of the sequence (ξ_i) . It follows that $|\xi(i)| \geq \varepsilon$ for every $i \in M$, a contradiction with the fact that $\xi \in c_0$. \square

Corollary 3.21. *Let \mathcal{B} be a uniform barrier on M and $h : \mathcal{B} \rightarrow c_0$ be such that $h''\mathcal{B}$ is weakly-pre-compact. Then for every $\varepsilon > 0$ there is an infinite set $N \subseteq M$ such that $\sum_{i \in N \setminus s} |h(s)(i)| < \varepsilon$ for every $s \in \mathcal{B} \upharpoonright N$.*

PROOF. Fix a summing sequence $\sum_i \varepsilon_i < \varepsilon$. For $k \in \mathbb{N}$ define $h_k : \mathcal{B} \rightarrow \text{FIN}$ for $s \in \mathcal{B}$ by $h_k(s) = \text{supp}_{\varepsilon_k} h(s) \setminus s$. Since $K \subseteq c_0$ is weakly-compact, h_k fulfills the conditions for Lemma 3.18 to be applied. So, there is some $N = \{n_k\} \subseteq M$ such that for every k and every

$s \in \mathcal{B} \upharpoonright M$ $h_k(s) \cap N \subseteq \{n_0, \dots, n_{k-1}\}$. In particular, fix $s \in \mathcal{B} \upharpoonright M$ and let $I \subseteq \mathbb{N}$ be such that $N \setminus s = \{n_i\}_{i \in I}$. Then for every $i \in I$, $n_i \notin \text{supp}_{\varepsilon_i} h(s)$, so $\sum_{i \in I} |h(s)(i)| \leq \sum_{i \in I} \varepsilon_i < \varepsilon$. \square

Definition 3.22. For $\mathcal{F} \subseteq \text{FIN}$, let

$$\mathcal{F}^{\sqsubseteq-\min} = \{s \in \mathcal{F} : \forall t \sqsubseteq s (t \in \mathcal{F} \rightarrow t = s)\}$$

$$\mathcal{F}^{\sqsubseteq-\max} = \{s \in \mathcal{F} : \forall t \sqsupseteq s (t \in \mathcal{F} \rightarrow t = s)\}.$$

REMARK 3.23. (a) $\mathcal{F}^{\sqsubseteq-\max}$ and $\mathcal{F}^{\sqsubseteq-\min}$ are thin families.

(b) For every infinite M and every $s \in \mathcal{F} \upharpoonright M$ there is some $t \in \mathcal{F}^{\sqsubseteq-\min} \upharpoonright M$ such that $t \sqsubseteq s$. In particular, $\mathcal{F} \upharpoonright M = \emptyset$ iff $\mathcal{F}^{\sqsubseteq-\min} \upharpoonright M = \emptyset$.

(c) Suppose that \mathcal{F} is pre-compact. Then for every infinite M and every $s \in \mathcal{F} \upharpoonright M$ there is some $t \in \mathcal{F}^{\sqsubseteq-\max} \upharpoonright M$ such that $s \sqsubseteq t$. In particular, $\mathcal{F} \upharpoonright M = \emptyset$ iff $\mathcal{F}^{\sqsubseteq-\max} \upharpoonright M = \emptyset$.

Given $\mathfrak{X} \subseteq \text{FIN} \times c_0$ and $M \subseteq \mathbb{N}$, let us denote $\mathfrak{X} \upharpoonright M = \{(s, \xi) \in K : s \subseteq M\}$.

Theorem 3.24. Suppose that \mathfrak{X} is a subset of $\text{FIN} \times c_0$ whose second projection $(\mathfrak{X})_2$ is a weakly pre-compact and bounded subset of c_0 . Let $\Phi : \mathfrak{X} \rightarrow \mathbb{R}$ is an arbitrary mapping. Then for every $\varepsilon > 0$, there is some M such that for every $(s, \xi) \in \mathfrak{X} \upharpoonright M$ there is some $(t, \eta) \in \mathfrak{X} \upharpoonright M$ such that $t \sqsubseteq s$ and

$$(a) |\Phi(s, \xi) - \Phi(t, \eta)| < \varepsilon,$$

$$(b) \sum_{i \in M \setminus t} |\eta(i)| < \varepsilon \max\{1, |\Phi(s, \xi)|\}.$$

PROOF. Fix $0 < \varepsilon < 1$. Suppose first that Φ is bounded, i.e. there is some bounded interval $I \subseteq \mathbb{R}$ such that $\Phi''\mathfrak{X} \subseteq I$. Let $I_0 \cup \dots \cup I_k$ be a partition of I into disjoint sets of diameter at most ε . Let $\mathcal{F}_i = \{s \in \text{FIN} : \exists \xi \in (\mathfrak{X})_2 \text{ such that } \Phi(s, \xi) \in I_i\}$, and let $\mathcal{B}_i = \mathcal{F}_i^{\sqsubseteq-\min}$ ($0 \leq i \leq k$). By a successive application of Proposition 3.5 we obtain some infinite set M_0 such that either $\mathcal{B}_i \upharpoonright M_0 = \emptyset$ or else $\mathcal{B}_i \upharpoonright M_0$ is a uniform barrier on M_0 ($0 \leq i \leq k$). Let $\Gamma \subseteq \{0, \dots, k\}$ be the set of those i such that $\mathcal{B}_i \upharpoonright M_0$ is a uniform barrier on M_0 . For $s \in \bigcup_{i \in \Gamma} \mathcal{B}_i \upharpoonright M_0$ let $\xi_s \in (\mathfrak{X})_2$ witnessing that $s \in \mathcal{F}_i$. Then, by Corollary 3.21, there is some $M \subseteq M_0$ such that for every $s \in \bigcup_{i \in \Gamma} \mathcal{B}_i \upharpoonright M$ we have that $\sum_{i \in M \setminus s} |\xi_s(i)| < \varepsilon \max\{1, \sup I\}$. We claim that M has the desired properties. So, for suppose that $(s, \xi) \in \mathfrak{X} \upharpoonright M$. Let i be such that $s \in \mathcal{F}_i \upharpoonright M$. Note that in this case $\mathcal{B}_i \upharpoonright M_0 \neq \emptyset$, so $i \in \Gamma$. Consider the infinite set $P = s \cup (M/s)$. Since $\mathcal{B}_i \upharpoonright M$ is a barrier on M there is some $t \in \mathcal{B}_i \upharpoonright M$ such that $t \sqsubseteq P$. So, either $t \sqsubseteq s$ or else $s \sqsubseteq t$. By the definition of \mathcal{B}_i , there is some $u \in \mathcal{B}_i \upharpoonright M$ such that $u \sqsubseteq s$. Since \mathcal{B}_i is thin, we conclude that $t \sqsubseteq s$ (otherwise $u \sqsubset t$). Now it is clear that (t, ξ_t) is the desired pair.

Suppose now that $\Phi : \mathfrak{X} \rightarrow \mathbb{R}$ is an arbitrary mapping. Let $L = \max\{\|\xi\|_\infty : \xi \in K\}$. Clearly we may assume that $\varepsilon < L$. For each $k \in \mathbb{N}$, let

$$\mathfrak{X}_k = \{(s, \xi) \in \mathfrak{X} : 2k\varepsilon^{-1}L \leq |\Phi(s, \xi)| < 2(k+1)\varepsilon^{-1}L\}.$$

Define $\Phi_k = \Phi \upharpoonright \mathfrak{X}_k$. For a finite set u and $k \in \mathbb{N}$, let

$$\mathfrak{X}_{k,u} = \{(s, \xi) \in \mathfrak{X}_k : u < s \text{ and } (u \cup s, \xi) \in \mathfrak{X}_k\},$$

and define $\Phi_{k,u} : \mathfrak{X}_{k,u} \rightarrow \mathbb{R}$ for $(s, \xi) \in \mathfrak{X}_{k,u}$ by $\Phi_{k,u}(s, \xi) = \Phi_k(u \cup s, \xi)$. Using that $\Phi_{k,u}$ is bounded, we can find a fusion sequence (M_k) of M such that, setting $m_k = \min M_k$, for every

$k \in \mathbb{N}$ every $u \subseteq \{m_0, \dots, m_{k-1}\}$ and every $(s, \xi) \in \mathfrak{X}_{k,u} \restriction M_k$ there is some $(t, \eta) \in \mathfrak{X}_{k,u} \restriction M_k$ such that $t \sqsubseteq s$,

$(a_{k,u}) \ |\Phi_k(u \cup s, \xi) - \Phi_k(u \cup t, \eta)| < \varepsilon$, and

$(b_{k,u}) \ \sum_{i \in M_k \setminus t} |\eta(i)| < (\varepsilon/2) \max\{1, \Phi_k(u \cup s, \xi)\}$.

Let us check that $N = \{m_k\}$ has the desired properties. For suppose that $(s, \xi) \in \mathfrak{X} \restriction M$. Let k be such that $2k\varepsilon^{-1}L \leq |\Phi(s, \xi)| < 2(k+1)\varepsilon^{-1}L$, set $u = s \cap \{m_0, \dots, m_{k-1}\}$, and $\bar{s} = s \setminus u \subseteq M_k$. Then there is some $(t, \eta) \in \mathfrak{X}_{k,u} \restriction M_k$ such that $t \sqsubseteq \bar{s}$ and $(a_{k,u})$ and $(b_{k,u})$ holds. This implies that $u \cup t \sqsubseteq s$, $|\Phi(s, \xi) - \Phi(u \cup t, \eta)| < \varepsilon$, and $\sum_{i \in M_k \setminus (u \cup t)} |\eta(i)| \leq \sum_{i \in M_k \setminus t} |\eta(i)| < (\varepsilon/2) \max\{1, \Phi(s, \xi)\}$. If $k = 0$ then we are done. Suppose that $k \geq 1$. Observe that in this case $|\Phi(s, \xi)| \geq 2kL\varepsilon^{-1} \geq 1$, so

$$\sum_{i \in N \setminus (u \cup t)} |\eta(i)| < \varepsilon + \sum_{i \leq k-1} |\eta(m_i)| \leq \frac{\varepsilon}{2} |\Phi(s, \xi)| + kL \leq \varepsilon |\Phi(s, \xi)| = \varepsilon \max\{1, |\Phi(s, \xi)|\}. \quad (10)$$

□

The following is a dual version of the previous result.

Theorem 3.25. *Suppose that $\mathfrak{X} \subseteq \mathcal{F} \times c_0$ and $\Phi : \mathfrak{X} \rightarrow \mathbb{R}$ is a bounded function such that $|\Phi|^{-1}(\delta, \infty) = \{(s, \xi) \in \mathfrak{X} : |\Phi(s, \xi)| > \delta\}$ is pre-compact for every $\delta > 0$. Then for every $\varepsilon > 0$, there is some M such that for every $N \subseteq M$ and every $(s, \xi) \in \mathfrak{X} \restriction N$ there is some $(t, \eta) \in \mathfrak{X} \restriction N$ such that $s \sqsubseteq t$ and*

(a) $|\Phi(s, \xi) - \Phi(t, \eta)| < \varepsilon$,

(b) $\sum_{i \in M \setminus t} |\eta(i)| < \varepsilon |\Phi(s, \xi)|^{-1}$.

PROOF. Fix $\varepsilon > 0$. Suppose first that Φ is strictly positive, i.e. there is some $0 < \delta < C \in \mathbb{R}$ such that $|\Phi|^{-1}(\delta, \infty) \subseteq (\mathfrak{X})_1$. Let $\mathcal{F} = (\mathfrak{X})_1$. By Theorem 3.14 we obtain some infinite set M_0 such that $(\mathcal{F}[M_0])^{\sqsubseteq-\max}$ is a uniform barrier \mathcal{B} on M . Let $I_0 \cup \dots \cup I_k$ be a partition of $(-C, -\delta) \cup (\delta, C)$ into disjoint sets of diameter at most ε . Consider the following coloring $c : \mathcal{B} \rightarrow \{0, 1\}^{\{0, \dots, k\}}$, defined by $c(s)(i) = 0$ iff there is some $t \sqsubseteq s$ and some ξ such that $(t, \xi) \in \mathfrak{X}$ and $\Phi(t, \xi) \in I_i$. By the Ramsey property of \mathcal{B} there is some $M_1 \subseteq M_0$ such that c is constant in $\mathcal{B} \restriction M_1$ with value $(a_i)_{i=0}^k$. Let $\Gamma = \{0 \leq i \leq k : a_i = 0\}$. For every $i \in \Gamma$, let $h_i : \mathcal{B} \restriction M_1 \rightarrow \text{FIN}$ be defined for $s \in \mathcal{B} \restriction M_1$ by $h_i(s) = t \in \mathcal{F} \restriction M_1$ iff t is the maximal initial part of s with the property that for some ξ , $(t, \xi) \in \mathfrak{X}$ and $\Phi(t, \xi) \in I_i$. By a successive application of Corollary 3.11 there is some $M_2 \subseteq M_1$ such that for every $i \in \Gamma$, $h_i''(\mathcal{B} \restriction M_2)$ is a uniform barrier \mathcal{B}_i on M_2 and if $t \in \mathcal{B}_i$ and $s \in \mathcal{B} \restriction M_2$ are such that $t \sqsubseteq s$, then $h_i(s) = t$. For $0 \leq i \leq k$ set $\mathcal{F}_i = \{s \in \text{FIN} : \text{there is some } \xi \text{ such that } (s, \xi) \in \mathfrak{X} \text{ and } \Phi(s, \xi) \in I_i\}$.

Claim. *If $i \notin \Gamma$ then $\mathcal{F}_i \restriction M_2 = \emptyset$. For every $N \subseteq M_2$ and every $i \in \Gamma$, $\mathcal{F}_i \restriction N = \overline{\mathcal{B}_i \restriction N}$, in particular hereditary.*

Proof of Claim: Suppose that $i \in \Gamma$, $N \subseteq M_2$ and suppose that $s \in \mathcal{F}_i \restriction N$. Let $t \in \mathcal{B} \restriction N$ be such that $s \sqsubseteq t$. Then $s \sqsubseteq h_i(t)$ by maximality of $h_i(t)$, so $s \in \overline{\mathcal{B}_i \restriction N}$. □

Now for $t \in \bigcup_{i \in \Gamma} \mathcal{B}_i$ let $\xi_t \in c_0$ be such that $(t, \xi_t) \in \mathfrak{X}$ and $\Phi(t, \xi_t) \in I_i$. Then, by Corollary 3.21, there is some $M \subseteq M_2$ such that for every $t \in \bigcup_{i \in \Gamma} \mathcal{B}_i \restriction M$ we have that $\sum_{i \in M \setminus t} |\xi_t(i)| < \varepsilon C^{-1}$. We claim that M has the desired properties. So, fix $N \subseteq M$, and $(s, \xi) \in \mathfrak{X} \restriction N$. Let i

be such that $\Phi(s, \xi) \in I_i$, and, by the previous claim, let $t \in \mathcal{B}_i \upharpoonright N$ be such that $s \subseteq t$. Then it is clear that $(t, \xi_t) \in \mathfrak{X} \upharpoonright M$ is the desired pair.

To show the general case, proceed as in the corresponding part of the proof of Theorem 3.24 using a suitable fast decreasing sequence to 0. \square

4. WEAKLY-NULL SEQUENCES IN BANACH SPACES

4.1. Weakly null sequences bounded away from zero. We start with the following classical construction of a Banach space.

Definition 4.1. Let \mathcal{F} be a compact hereditary family of finite subsets of \mathbb{N} containing all singletons $\{n\}$ ($n \in \mathbb{N}$) and let (e_n) denote the unit vector basis of c_{00} . We define a norm $\|\cdot\|_{\mathcal{F}}$ on c_{00} by the rule

$$\left\| \sum_n a_n e_n \right\|_{\mathcal{F}} = \sup \left\{ \sum_{n \in s} |a_n| : s \in \mathcal{F} \right\}, \text{ for all } (a_n) \in c_{00}. \quad (11)$$

The completion of $(c_{00}, \|\cdot\|_{\mathcal{F}})$ is a Banach space which we denote by $S_{\mathcal{F}}$ and call the \mathcal{F} -Schreier space because Schreier was the first to consider $S_{\mathcal{F}}$ when \mathcal{F} is equal to the Schreier family described above in Remark 3.4. Note that (e_i) as a normalized, 1-unconditional, shrinking, monotone Schauder basis (see [12] and [1]).

REMARK 4.2. Suppose that \mathcal{B} is a front on \mathbb{N} , $\alpha > 0$. Then for every $x = \sum_n a_n e_n \in S_{\overline{\mathcal{B}}}$ we have that

$$\|x\|_{\overline{\mathcal{B}}} = \sup \left\{ \sum_{n \in s} |a_n| : s \in \mathcal{B} \right\} \quad (12)$$

For a given $t \in \overline{\mathcal{B}}$. Let M be an infinite set of integers such that $t < M$. Then let $s \in \mathcal{B}$ be such that $s \subseteq t \cup M$. It is clear that $t \subseteq s$, so,

$$\sum_{n \in s} |a_n| = \sum_{n \in t} |a_n| + \sum_{n \in s \setminus t} |a_n| \geq \sum_{n \in t} |a_n|. \quad (13)$$

In other words, given an α -uniform barrier \mathcal{B} on \mathbb{N} , $\alpha > 0$, the corresponding rule (11) for $\mathcal{F} = \mathcal{B}$ produces a Banach space with the same properties as stated above, even if \mathcal{B} is obviously neither compact, nor hereditary.

Theorem 4.3. Suppose that $K \subseteq c_0$ is weakly-compact and bounded such that $(p_n) \subseteq C(K)$ is normalized and with the additional property that

$$\inf \{ |\xi(i)| : i \in \mathbb{N}, \xi \in K \} = \delta > 0. \quad (14)$$

Then there is some M and a uniform barrier \mathcal{C} on \mathbb{N} such that $(p_i)_{i \in M}$ is $\delta/4$ -equivalent to the natural basis (e_i) of the Schreier space $S_{\mathcal{C}}$. In particular, $(p_i)_{i \in M}$ is unconditional.

PROOF. By (14) the family $\text{supp}'' K \subseteq \text{FIN}$ is pre-compact. Let M_0 be such that $(\text{supp}'' K)[M_0]$ is the closure of a uniform barrier \mathcal{B} on M_0 . Define the partial mapping $\Phi : \mathcal{B} \times K \rightarrow \{-1, 1\}$ by $\Phi(s, \xi) = \sigma \in \{-1, 1\}$ iff ξ has constant sign σ on s . Apply Theorem 3.25 to Φ and to $\delta/2$ to obtain $M \subseteq M_0$ such that for every $(s, \xi) \in \overline{\mathcal{B}} \upharpoonright M \times K$ such that ξ has constant sign σ on s there is some $\eta \in K$ such that

(a) η has constant sign σ on s ,

(b) $\eta \upharpoonright (M \setminus s) = 0$ (this is because $\sum_{i \in M \setminus s} |\eta(i)| < 2\varepsilon \leq \delta$).

Let $\Theta : M \rightarrow \mathbb{N}$ be the unique order-preserving onto mapping between M and \mathbb{N} . Let $\mathcal{C} = \Theta''\mathcal{B}$. We are going to show that for every sequence of scalars $(a_i)_{i \in M}$

$$\frac{\delta}{4} \left\| \sum_{i \in M} a_i e_{\Theta(i)} \right\|_{S_{\mathcal{C}}} \leq \left\| \sum_{i \in M} a_i p_i \right\| \leq \left\| \sum_{i \in M} a_i e_{\Theta(i)} \right\|_{S_{\mathcal{C}}}. \quad (15)$$

So fix scalars $(a_i)_{i \in M}$. Then for a given $\xi \in K$ we obtain that

$$\left| \left(\sum_{i \in M} a_i p_i \right)(\xi) \right| = \left| \sum_{i \in \text{supp } \xi \cap M} a_i \xi(i) \right| \leq \sum_{i \in \text{supp } \xi \cap M} |a_i| \leq \left\| \sum_{i \in M} a_i e_{\Theta(i)} \right\|_{S_{\mathcal{C}}}, \quad (16)$$

since $\text{supp } \xi \upharpoonright M \in (\text{supp}''K)[M] = \overline{\mathcal{B}}^{\subseteq}$. This shows the right hand inequality of (15). Now fix $s \in \mathcal{B}$, and let $\xi \in \Gamma$ be such that $s = \text{supp } \xi \upharpoonright M$.

For $(j, k) \in \{0, 1\}^2$, let

$$s_{j,k} = \{i \in s : (-1)^j a_i, (-1)^k \xi(i) \geq 0\}. \quad (17)$$

Note that $s = s_{0,0} \cup s_{0,1} \cup s_{1,0} \cup s_{1,1}$, and that ξ has constant sign $(-1)^k$ on $s_{j,k}$, so for every $(j, k) \in \{0, 1\}^2$ there is $\eta_{j,k} \in \overline{\mathcal{B}} \upharpoonright M \times K$ such that $\eta_{j,k}$ has constant sign $(-1)^j$ on $s_{j,k}$, and $\eta_{j,k} \upharpoonright (M \setminus s_{j,k}) = 0$. Hence for every $(j, k) \in \{0, 1\}^2$ we obtain that

$$\begin{aligned} \left| \left(\sum_{i \in M} a_i p_i \right)(\eta_{j,k}) \right| &= \left| \sum_{i \in s_{j,k}} (-1)^j |a_i| (-1)^k |\eta_{j,k}(i)| \right| = \\ &= \sum_{i \in s_{j,k}} |a_i| |\eta_{j,k}(i)| \geq \delta \sum_{i \in s_{j,k}} |a_i|, \end{aligned} \quad (18)$$

and so

$$\left\| \sum_{i \in M} a_i p_i \right\| \geq \frac{\delta}{4} \left\| \sum_{i \in M} a_i t_{\Theta(i)} \right\|_{S_{\mathcal{C}}}, \quad (19)$$

that shows the left hand inequality of (15). \square

Corollary 4.4. *Suppose that (x_n) is a normalized weakly-null sequence of $\ell_{\infty}(\Gamma)$ with the property that*

$$\inf\{|x_n(\gamma)| : n \in \mathbb{N}, \gamma \in \Gamma\} = \delta > 0. \quad (20)$$

Then there is an infinite set N and a α -uniform barrier \mathcal{C} on \mathbb{N} ($\alpha > 0$) such that $(x_n)_{n \in M}$ is $\delta/4$ -equivalent to the natural basis (e_i) of the Schreier space $\mathcal{S}_{\mathcal{C}}$. In particular, $(x_n)_{n \in M}$ is unconditional. \square

REMARK 4.5. The unconditional constant $4 + \varepsilon$ obtained before can be improved to $2 + \varepsilon$.

4.2. Near unconditionality. Recall the result of Elton [9] (see also [17]) which says that if (x_n) is a weakly-null normalized sequence in some Banach space then it has an infinite subsequence $(x_n)_{n \in M}$ such that for every $\delta > 0$ there is a constant $C(\delta) < \infty$ such that for every sequence $(a_n)_{n \in M} \subseteq [-1, 1]$ and every finite $s \subseteq \{n : |a_n| \geq \delta\}$,

$$\left\| \sum_{n \in s} a_n y_n \right\| \leq C(\delta) \left\| \sum_{n \in M} a_n y_n \right\|. \quad (21)$$

The purpose of this subsection is to strengthen this result as follows.

Theorem 4.6 (Near unconditionality). *Suppose that (x_n) is a normalized weakly-null sequence. Then for every $\varepsilon > 0$ there is some M such that for every sequence of scalars $(a_i)_{i \in M}$ and every finite subset $s \subseteq M$,*

$$\left\| \sum_{i \in s} a_i x_i \right\| \leq (2 + \varepsilon) \frac{\sup_{i \in M} |a_i|}{\min_{i \in s} |a_i|} \left\| \sum_{i \in M} a_i x_i \right\| \quad (22)$$

PROOF. We assume that the sequence is the evaluation mapping sequence $(p_n) \subseteq C(K)$ for some symmetric weakly-compact $K \subseteq c_0$ that in addition is closed under restrictions on initial intervals. Let

$$\mathfrak{X} = \{(s, \xi) \in \text{FIN} \times K : \xi \text{ has constant sign on } s\},$$

and let $\Phi : \mathfrak{X} \rightarrow \mathbb{R}$ for $(s, \xi) \in \mathfrak{X}$ by $\Phi(s, \xi) = |\sum_{i \in s} \xi(i)|$. Now let $\delta_0 = 1$ and for $k \geq 1$ let $\delta_k = 2^{-(k+2)}\varepsilon$. For $k \in \mathbb{N}$ let $\varepsilon_k = \delta_{k+1}^2/\gamma$, where γ is such that $\gamma/(\gamma - 4) \leq 1 + \varepsilon/4$. Apply repeatedly Theorem 3.24 to $\Phi : \mathfrak{X} \rightarrow \mathbb{R}$ and ε_k to find a fusion sequence (M_k) where each M_k satisfies the conclusions there for Φ and ε_k . We claim that the fusion set $M = \{m_k\}$, $m_k = \min M_k$ ($k \in \mathbb{N}$), has the desired properties. For suppose that $(a_i)_{i \in M}$ are scalars. Without loss of generality we may assume that $\sup_{i \in M} |a_i| \leq 1$, and suppose that $s \subseteq M$ is finite. If $\min_{i \in s} |a_i| = 0$, then the desired result is trivially true. So, suppose that $\min_{i \in s} |a_i| > 0$ and let $k \in \mathbb{N}$ be such that $\delta_{k+1} < \min_{i \in s} |a_i| \leq \delta_k$. Let $u = s \cap \{m_0, \dots, m_{k-1}\}$, and $v = s \setminus u \subseteq M_k$. First observe that $k \min_{i \in s} |a_i| \leq k \delta_k \leq \varepsilon/4$ ($k \in \mathbb{N}$), so

$$\left\| \sum_{i \in u} a_i p_i \right\| \leq k \max_{i \in s} |a_i| \leq \frac{\varepsilon \max_{i \in s} |a_i|}{4 \min_{i \in s} |a_i|} \leq \frac{\varepsilon}{4 \min_{i \in s} |a_i|} 2 \left\| \sum_{i \in M} a_i p_i \right\| = \frac{\varepsilon}{\min_{i \in s} |a_i|} \left\| \sum_{i \in M} a_i p_i \right\|, \quad (23)$$

where we are using that $\left\| \sum_{i \in M} a_i p_i \right\| \geq (1/2) \max_{i \in s} |a_i|$, since (p_i) is monotone. Let $\xi \in K$ be such that $\left\| \sum_{i \in v} a_i p_i \right\| = \sum_{i \in v} a_i \xi(i)$. For $j = 0, 1$, let

$$v_j = \{i \in v : (-1)^j a_i, (-1)^j \xi(i) \geq 0\}$$

Then

$$\left\| \sum_{i \in v} a_i p_i \right\| = \sum_{i \in v} a_i \xi(i) \leq \sum_{i \in v_0} a_i \xi(i) + \sum_{i \in v_1} a_i \xi(i). \quad (24)$$

Let $j = 0, 1$ be such that $\left\| \sum_{i \in v} a_i p_i \right\| \leq 2 \sum_{i \in v_j} a_i \xi(i)$. Since $(v_j, \xi) \in \mathfrak{X} \upharpoonright M_k$ there is some $(t, \eta) \in \mathfrak{X} \upharpoonright M_k$ with $t \sqsubseteq v_j$ and such that

- (a) $|\sum_{i \in v_j} \xi(i) - \sum_{i \in t} \eta(i)| < \varepsilon_k$, and
- (b) $\sum_{i \in M_k \setminus t} |\eta(i)| < \varepsilon_k \max\{1, \sum_{i \in v_j} \xi(i)\}$. It follows that

$$\begin{aligned} \left\| \sum_{i \in M} a_i p_i \right\| &\geq \left\| \sum_{i \in M/m_{k-1}} a_i p_i \right\| \geq \left| \left(\sum_{i \in M/m_{k-1}} a_i p_i \right)(\eta) \right| \geq \left| \sum_{i \in t} a_i \eta(i) \right| - \varepsilon_k \max\{1, \sum_{i \in v_j} \xi(i)\} \geq \\ &\geq \min_{i \in s} |a_i| \left| \sum_{i \in t} \eta(i) \right| - \varepsilon_k \max\{1, \left| \sum_{i \in v_j} \xi(i) \right|\} \geq \\ &= \min_{i \in s} |a_i| \left(\left| \sum_{i \in v_j} \xi(i) \right| - \varepsilon_k \right) - \varepsilon_k \max\{1, \left| \sum_{i \in v_j} \xi(i) \right|\} \end{aligned} \quad (25)$$

Since $\delta_{k+1} \leq \min_{i \in s} |a_i| \leq 2 \|\sum_{i \in s} a_i p_i\| \leq 2 \|\sum_{i \in v_j} \xi(i)\|$ we have that

$$\min_{i \in s} |a_i| \left(\left| \sum_{i \in v_j} \xi(i) \right| - \varepsilon_k \right) = \min_{i \in s} |a_i| \left(\left| \sum_{i \in v_j} \xi(i) \right| - \frac{\delta_{k+1}^2}{\gamma} \right) \geq \frac{\gamma - 2}{\gamma} \min_{i \in s} |a_i| \sum_{i \in v_j} \xi(i), \quad (26)$$

while

$$\varepsilon_k \max\{1, \left| \sum_{i \in v_j} \xi(i) \right|\} \leq \frac{2}{\gamma} \min_{i \in s} |a_i| \sum_{i \in v_j} \xi(i). \quad (27)$$

It follows that

$$\left\| \sum_{i \in M} a_i p_i \right\| \geq \frac{\gamma - 4}{\gamma} \min_{i \in s} |a_i| \sum_{i \in v_j} \xi(i) \geq \frac{\gamma - 4}{2\gamma} \min_{i \in s} |a_i| \left\| \sum_{i \in v} a_i p_i \right\| \geq \frac{2}{4 + \varepsilon} \min_{i \in s} |a_i| \left\| \sum_{i \in v} a_i p_i \right\| \quad (28)$$

and so

$$\left\| \sum_{i \in s} a_i p_i \right\| \leq \left\| \sum_{i \in u} a_i p_i \right\| + \left\| \sum_{i \in v} a_i p_i \right\| \leq \frac{\varepsilon}{2 \min_{i \in s} |a_i|} \left\| \sum_{i \in M} a_i p_i \right\| + \quad (29)$$

$$+ \frac{4 + \varepsilon}{2} \frac{1}{\min_{i \in s} |a_i|} \left\| \sum_{i \in M} a_i p_i \right\| = \frac{2 + \varepsilon}{\min_{i \in s} |a_i|} \left\| \sum_{i \in M} a_i p_i \right\|. \quad (30)$$

□

4.3. Convex unconditionality. Recall the dual form of Elton's theorem proved by Argyros, Mercourakis and Tsarpalias [2] which says that every normalized weakly-null sequence (x_n) of a given Banach space has an infinite subsequence $(x_n)_{n \in M}$ such that for every $\delta > 0$ there is a constant $C(\delta) > 0$ such that if $\sum_{n \in M} |a_n| \leq 1$ and $\|\sum_{n \in M} a_n x_n\| > \delta$ then $\|\sum_{n \in M} \varepsilon_n a_n y_n\| > C(\delta)$ for all choices of signs ε_n . The purpose of this section is to extend this result as follows.

Theorem 4.7 (Convex unconditionality). *Suppose that (x_n) is a normalized weakly-null sequence of a Banach space E . Then for every $\varepsilon > 0$ there is some M such that for every $(a_i)_{i \in M}$ such that $\sup_{i \in M} |a_i| \leq 1$ and every subset $N \subseteq M$ such that $\sum_{i \in N} |a_i| \leq 1$,*

$$\left\| \sum_{i \in N} a_i x_i \right\| \leq (4 + \varepsilon) \sqrt{\left\| \sum_{i \in M} a_i x_i \right\|}. \quad (31)$$

PROOF. We assume that the sequence considered is the evaluation mapping sequence (p_n) of $C(K)$ where $K \subseteq c_0$ is weakly-compact, symmetric, bounded by 1, and closed under restriction on initial intervals (i.e. (x_i) is a weakly-null monotone basic sequence). Define the partial mapping $\Phi : \text{FIN} \times K \rightarrow \mathbb{R}$ by $\Phi(s, \xi) = \min_{i \in s} |\xi(i)|$ iff ξ has constant sign on s . It is not difficult to see that Φ fulfills the hypothesis of Theorem 3.25. Now let $\delta_0 = 1$, and for $k \geq 1$, let $\delta_k = \varepsilon/2^{k+2}$. And for $k \in \mathbb{N}$, let $\varepsilon_k = \delta_{k+1}^4/\gamma$, where γ is such that $\gamma/(\gamma - 2) \leq (2 + 2\varepsilon)/(2 + \varepsilon)$. Apply repeatedly Theorem 3.24 to Φ and ε_k to find a fusion sequence (M_k) where each M_k satisfies the conclusions there for Φ and ε_k . Let $M = \{m_k\}$ be the fusion set of (M_k) .

Claim. *Let $(a_i)_{i \in M}$ be a sequence of scalars with $\sup_{i \in M} |a_i| \leq 1$, and let $s \subseteq M$ be such that both $\xi \in K$ and a_i 's have constant signs on s . Then*

$$\left| \sum_{i \in s} a_i \xi(i) \right| \leq \frac{1 + \varepsilon}{\min\{\left| \sum_{i \in s} a_i \right|, \min_{i \in s} |\xi(i)|\}} \left\| \sum_{i \in M} a_i p_i \right\| \quad (32)$$

Proof of Claim: Set $C = \min\{|\sum_{i \in s} a_i|, \min_{i \in s} |\xi(i)|\}$. Let $k \in \mathbb{N}$ be such that $\delta_{k+1} < C \leq \delta_k$, and set $u = s \cap \{m_0, \dots, m_{k-1}\}$, and $v = s \setminus u \subseteq M_k$. First observe that $kC \leq k\delta_k \leq \varepsilon/4$ ($k \in \mathbb{N}$), so

$$|\sum_{i \in u} a_i \xi(i)| \leq k \max_{i \in s} |a_i| \leq \frac{\varepsilon \max_{i \in s} |a_i|}{4C} \leq \frac{\varepsilon}{4C} (2 \|\sum_{i \in M} a_i p_i\|) = \frac{\varepsilon}{2C} \|\sum_{i \in M} a_i p_i\|, \quad (33)$$

Since $(v, \xi) \in \text{dom } \Phi \upharpoonright M_k$ there is some $(t, \bar{\eta}) \in \text{dom } \Phi \upharpoonright M_k$ with $v_j \sqsubseteq t$ such that

(a) $\min_{i \in v} |\xi(i)| - \min_{i \in t} |\bar{\eta}(i)| < \varepsilon_k$, and

(b) $\sum_{i \in M_k \setminus t} |\bar{\eta}(i)| < \varepsilon_k (\min_{i \in v} |\xi(i)|)^{-1}$.

But K is closed under restriction on initial intervals, so it readily follows that, setting $\eta = \bar{\eta} \upharpoonright [0, \max v] \in K$, then we obtain that (a') $\min_{i \in v} |\eta(i)| \geq \min_{i \in v} |\xi(i)| - \varepsilon_k$, and (b') $\sum_{i \in M_k \setminus v} |\eta(i)| < \varepsilon_k (\min_{i \in v} |\xi(i)|)^{-1}$. Hence,

$$\begin{aligned} \|\sum_{i \in M} a_i p_i\| &\geq \|\sum_{i \in M/m_{k-1}} a_i p_i\| \geq |(\sum_{i \in M/m_{k-1}} a_i p_i)(\eta)| \geq |\sum_{i \in v} a_i \eta(i)| - \frac{\varepsilon_k}{\min_{i \in v} |\xi(i)|} \geq \\ &\geq |\sum_{i \in v} a_i| \min_{i \in v} \eta(i) - \frac{\varepsilon_k}{\min_{i \in v} |\xi(i)|} \geq |\sum_{i \in v} a_i| (\min_{i \in v} |\xi(i)| - \varepsilon_k) - \frac{\varepsilon_k}{\min_{i \in v} |\xi(i)|} \end{aligned} \quad (34)$$

Now since $\delta_{k+1} < C = \min\{|\sum_{i \in s} a_i|, \min_{i \in s} |\xi(i)|\}$ we obtain that

$$\min_{i \in v} |\xi(i)| - \varepsilon_k = \min_{i \in v} |\xi(i)| - \frac{\delta_{k+1}^4}{\gamma} \geq \min_{i \in v} |\xi(i)| \frac{\gamma - 1}{\gamma} \quad \text{and}, \quad (35)$$

$$\frac{\varepsilon_k}{\min_{i \in v} |\xi(i)|} \leq \frac{\min\{|\sum_{i \in s} a_i|, \min_{i \in s} |\xi(i)|\}^4}{\gamma \min_{i \in v} |\xi(i)|} \leq \frac{\min\{|\sum_{i \in s} a_i|, \min_{i \in s} |\xi(i)|\}^3}{\gamma}. \quad (36)$$

Hence,

$$\|\sum_{i \in M} a_i p_i\| \geq \frac{\gamma - 1}{\gamma} \min_{i \in v} |\xi(i)| \|\sum_{i \in v} a_i\| - \frac{(\min_{i \in s} |\xi(i)|)^2 |\sum_{i \in s} a_i|}{\gamma} \geq \quad (37)$$

$$\geq \frac{\gamma - 1}{\gamma} \min_{i \in v} |\xi(i)| \|\sum_{i \in v} a_i \xi(i)\| - \frac{\min_{i \in s} |\xi(i)| |\sum_{i \in s} a_i \xi(i)|}{\gamma} = \quad (38)$$

$$= \frac{1}{\gamma} \min_{i \in s} |\xi(i)| \left((\gamma - 1) \|\sum_{i \in v} a_i \xi(i)\| - \|\sum_{i \in s} a_i \xi(i)\| \right) \geq \quad (39)$$

$$\geq \frac{1}{\gamma} \min_{i \in s} |\xi(i)| \left((\gamma - 2) \|\sum_{i \in s} a_i \xi(i)\| - (\gamma - 1) \|\sum_{i \in u} a_i \xi(i)\| \right) \quad (40)$$

So,

$$\frac{\gamma - 2}{\gamma} \min_{i \in s} |\xi(i)| \|\sum_{i \in s} a_i \xi(i)\| \leq \|\sum_{i \in M} a_i p_i\| + \min_{i \in s} |\xi(i)| \|\sum_{i \in u} a_i \xi(i)\| \leq \quad (41)$$

$$\leq \|\sum_{i \in M} a_i p_i\| + \min_{i \in s} |\xi(i)| \frac{\varepsilon}{2C} \|\sum_{i \in M} a_i p_i\|, \quad (42)$$

and therefore,

$$|\sum_{i \in s} a_i \xi(i)| \leq \frac{\gamma}{\gamma - 2} \left(\frac{1}{\min_{i \in s} |\xi(i)|} \left\| \sum_{i \in M} a_i p_i \right\| + \frac{\varepsilon}{2C} \left\| \sum_{i \in M} a_i p_i \right\| \right) \leq \quad (43)$$

$$\leq \frac{\gamma}{\gamma - 2} \left(\frac{1}{C} \left\| \sum_{i \in M} a_i p_i \right\| + \frac{\varepsilon}{2C} \left\| \sum_{i \in M} a_i p_i \right\| \right) = \quad (44)$$

$$= \frac{1}{C} \frac{\gamma}{\gamma - 2} \left(1 + \frac{\varepsilon}{2} \right) \left\| \sum_{i \in M} a_i p_i \right\| \leq \frac{1 + \varepsilon}{\min\{|\sum_{i \in s} a_i|, \min_{i \in s} |\xi(i)|\}} \left\| \sum_{i \in M} a_i p_i \right\|. \quad (45)$$

□

Now fix scalars $(a_i)_{i \in M}$ with $\sup_{i \in M} |a_i| \leq 1$ and let $s \subseteq M$ be such that $\sum_{i \in s} |a_i| \leq 1$. Since (p_i) is a basic sequence, it is clear that we may assume that s is a finite set. Choose $\xi \in K$ such that $\|\sum_{i \in s} a_i p_i\| = \sum_{i \in s} a_i \xi(i) = \delta$. Let $\bar{s} = \{i \in s : |\xi(i)| \geq \delta/2\}$. Since $\sum_{i \in s} |a_i| \leq 1$ we obtain that $\sum_{i \in \bar{s}} a_i \xi(i) \geq (1/2) \|\sum_{i \in s} a_i p_i\|$. Consider the two disjoint subsets u and v of \bar{s} where both ξ and a_i have constant sign. Choose one of them, say u , such that $\sum_{i \in u} a_i \xi(i) \geq (1/2) \sum_{i \in \bar{s}} a_i \xi(i) \geq (1/4) \|\sum_{i \in s} a_i p_i\|$. Now we can apply the previous claim to u and ξ to obtain

$$\frac{1}{4} \left\| \sum_{i \in s} a_i p_i \right\| \leq \sum_{i \in u} a_i \xi(i) \leq \frac{1 + \varepsilon}{\min\{|\sum_{i \in u} a_i|, \min_{i \in u} |\xi(i)|\}} \left\| \sum_{i \in M} a_i p_i \right\|. \quad (46)$$

Since $u \subseteq \bar{s}$ we obtain that $\min_{i \in u} |\xi(i)| \geq (1/2) \|\sum_{i \in s} a_i p_i\|$, while $|\sum_{i \in u} a_i| \geq |\sum_{i \in u} a_i \xi(i)| \geq (1/4) \|\sum_{i \in s} a_i p_i\|$, so $\min\{|\sum_{i \in u} a_i|, \min_{i \in u} |\xi(i)|\} \geq (1/4) \|\sum_{i \in s} a_i p_i\|$, and hence, from (46),

$$\left\| \sum_{i \in s} a_i p_i \right\|^2 \leq 16(1 + \varepsilon) \left\| \sum_{i \in M} a_i p_i \right\|. \quad (47)$$

□

5. WEAKLY-NULL SEQUENCES IN $C(K)$

In this section we apply the results of Section 3 to study of weakly null sequences in Banach spaces of the form $C(K)$ for K a countable compactum. Recall that these spaces are isomorphic to spaces of the form $C(\omega^{\omega^\alpha})$ for α a countable ordinal (see [7]), so it is not surprising that the theory of barriers is relevant here.

5.1. Point-finite weakly-compact sets and Lipschitz assignments. We will use the following result that allows us to change, without losing generality, from a weakly compact subset of c_0 to another one which is more convenient to work with.

Proposition 5.1. *Suppose that $K \subseteq c_0$ is arbitrary. Then for every ε there is a point-finite $L \subseteq c_{00}$ such that*

$$d_{\ell_1}(K, L) = \max\{\sup\{d_{\ell_1}(K, \xi) : \xi \in L\}, \sup\{d_{\ell_1}(\eta, L) : \eta \in K\}\} < \varepsilon. \quad (48)$$

In case that K is weakly-compact, L can be found being also weakly-compact. Also, if $K \subseteq c_{00}$, then L can be chosen to be included in c_{00} .

PROOF. Fix (ε_i) such that $\sum_i \varepsilon_i < \varepsilon$. For each i , split \mathbb{R} into intervals $(I_j^{(i)})$ each one of them of diameter at most ε_i , and choose $r_j^{(i)} \in I_j^{(i)}$ in a way that if $0 \in I_j^{(i)}$ then the choice of $r_j^{(i)} = 0$. Define $\Phi : K \rightarrow c_{00}$ for $\xi \in K$, $\Phi(\xi)(i) = r_j^{(i)}$ iff $\xi(i) \in I_j^{(i)}$. Then the image $L = \Phi''K$ fulfills all the requirements. If K is weakly-compact, then it can be shown that the image $\Phi''K$ is also weakly-pre-compact, so its weak-closure is the desired compactum L . \square

REMARK 5.2. Note that if $d_{\ell_1}(K, L) \leq \varepsilon$ then the corresponding evaluation mapping sequences $(p_i^K) \subseteq C(K)$ and $(p_i^L) \subseteq C(L)$ are $1 + \varepsilon$ -equivalent. So for all our purposes here, we can change from $C(K)$ to $C(L)$.

Definition 5.3. We say that $K \subseteq c_0$ is *point-finite* if its projection $\{\xi(i) : \xi \in K\}$ on every coordinate $i \in \mathbb{N}$ is finite. A mapping $\Phi : \mathcal{F} \rightarrow c_0$ defined on some $\mathcal{F} \subseteq \text{FIN}$ is called point-finite if its range $\Phi''\mathcal{F}$ is a point finite subset of c_0 . We say that Φ is *Lipschitz* if $\Phi(s) \upharpoonright (s \cap t) = \Phi(t) \upharpoonright (s \cap t)$ for every $s, t \in \mathcal{F}$ such that $s \cap t \subseteq s, t$.

REMARK 5.4. (a) Note that if $K \subseteq c_{00}$ is weakly-compact and point-finite, then the Cantor-Bendixon rank of K coincides with the the Cantor-Bendixson rank of its support set $\text{supp } K \subseteq \text{FIN}$.

(b) If \mathcal{F} is a front on M and $h : \mathcal{F} \rightarrow c_{00}$ is Lipschitz, then its extension to the topological closure of \mathcal{F} is well defined and continuous. Hence, in this case, the image $h''\overline{\mathcal{F}}$ is also weakly-compact.

Proposition 5.5. Suppose that \mathcal{B} is a uniform barrier on M , and suppose that $\Phi : \mathcal{B} \rightarrow c_{00}$ is point-finite. Then there is an infinite subset N of M such that Φ restricted to $\mathcal{B} \upharpoonright N$ is Lipschitz.

PROOF. By the Ramsey property of \mathcal{B} we can find a fusion sequence (M_k) of subsets of M , setting $m_k = \min M_k$, then for every k and every $u \subseteq \{m_0, \dots, m_{k-1}\}$, if $s, t \in \mathcal{B}_u \upharpoonright M_k$ then $\Phi(u \cup s) \upharpoonright u = \Phi(u \cup t) \upharpoonright u$. Then it is clear that the desired result will hold for the fusion set $\{m_k\}$. So, suppose defined $(M_i)_{i \leq k-1}$. For a fixed $u \subseteq \{m_0, \dots, m_{k-1}\}$ consider the finite coloring

$$h_u : \mathcal{B}_u \upharpoonright M_{k-1} \rightarrow c_{00} \quad (49)$$

defined for $s \in \mathcal{B}_u \upharpoonright M_{k-1}$ by $h_u(s) = \Phi(u \cup s) \upharpoonright u$. Then there is some $P_u \subseteq M_{k-1}$ such that h_u is monochromatic. Since there are only finitely many $u \subseteq \{m_0, \dots, m_{k-1}\}$ we can find $M_k \subseteq M_{k-1}$ such that every h_u is constant on $\mathcal{B}_u \upharpoonright M_k$ ($u \subseteq \{m_0, \dots, m_{k-1}\}$). as desired. \square

Corollary 5.6. Suppose that \mathcal{B} is an uniform barrier on M , and suppose that $\Phi : \mathcal{B} \rightarrow c_{00}$. Then for every $\varepsilon > 0$ there is $N \subseteq M$, and a point-finite Lipschitz $\bar{\Phi} : \mathcal{B} \upharpoonright N \rightarrow c_{00}$ such that for every $s \in \mathcal{B} \upharpoonright N$, $\|\Phi(s) - \bar{\Phi}(s)\|_{\ell_1} \leq \varepsilon$. \square

5.2. c_0 -saturation. Recall the result of Pelczynski and Semadeni [18] which says that every Banach space of the form $C(K)$ for K a countable compactum is c_0 -saturated in the sense that every of its closed infinite-dimensional subspaces contains an isomorphic copy of c_0 . The purpose of this section is to examine the c_0 -saturation using the theory of mappings on barriers developed above in Section 3. We start with a convenient reformulation of the problem.

Lemma 5.7. *Suppose that K is a countable compactum. Suppose that $(x_i) \subseteq C(K)$ is a normalized weakly null sequence. Then for every $\varepsilon > 0$ there is subsequence $(x_i)_{i \in M}$ and a weakly-compact subset $L \subseteq c_0$ such that*

- (a) *L is point-finite, it is closed under restrictions on intervals and $\text{supp } L \subseteq \text{FIN}$ is the closure of an α -uniform barrier on \mathbb{N} with α smaller than the Cantor-Bendixson rank of K ,*
- (b) *$(x_i)_{i \in M}$ and $(p_i)_{i \in \mathbb{N}}$ are $(1 + \varepsilon)$ -equivalent.*

PROOF. Fix $\varepsilon > 0$. Since K is countable we can find an infinite set M and a decreasing sequence $(\varepsilon_i)_{i \in M}$ of positive reals such that $\sum_{i \in M} \varepsilon_i \leq \varepsilon$ and such that (x_i/ε_i) is also a weakly null sequence. Set $N = {}^*M$. Since K is zero-dimensional compactum, we can find clopen sets $C_i \subseteq K$ ($i \in N$) such that $x_i^{-1}(K \setminus (-\varepsilon_i, \varepsilon_i)) \subseteq C_i \subseteq y_i^{-1}(K \setminus [-\varepsilon_i, \varepsilon_i])$ for every $i \in N$, and where i^- denotes the immediate predecessor of i in M . Set $y_i = \chi_{C_i} x_i$ ($i \in N$). Observe that (a) $\|x_i - y_i\|_{C(K)} < \varepsilon_{i^-}^{-1}$, so $(x_i)_{i \in N}$ and $(y_i)_{i \in N}$ are $1 + \varepsilon$ -equivalent, and (b) for every $\xi \in K$ and every $i \in N$, if $|y_i(\xi)| \leq \varepsilon_i$, then $y_i(\xi) = 0$. This last condition implies that $\text{supp } K((y_i)_{i \in N})$ is a pre-compact family of finite sets, so by Theorem 3.14 there is some $P \subseteq N$ such that $\text{supp } K((y_i)_{i \in N})[P] = \text{supp } K((y_i)_{i \in P})$ is the closure of a uniform barrier on P . So the subsequence $(p_i)_{i \in P}$ of the evaluation mapping sequence of $C(K((y_i)_{i \in P}))$ is $(1 + \varepsilon)$ -equivalent to $(x_i)_{i \in P}$. Now, using Proposition 5.1, we can perturb $K((y_i)_{i \in P})$ to make it point-finite. Now using the order-preserving mapping between P and \mathbb{N} we can easily obtain the desired result. \square

Definition 5.8. A *mean* is an element $\mu \in c_{00}^+$ with the property that $\sum_{i \in \mathbb{N}} \mu(i) = 1$. A mapping $\mu : \mathcal{B} \rightarrow c_{00}^+$ defined on some $\mathcal{B} \subseteq \text{FIN}$ is called a *mean-assignment* if for every $s \in \mathcal{B}$, $\mu(s)$ is a mean and $\text{supp } \mu(s) = s$.

Theorem 5.9. *Suppose that K is a countable compactum and that $(x_i) \subseteq C(K)$ is a normalized weakly-null sequence. Then there is $C \geq 1$, an infinite set M , a uniform barrier \mathcal{B} on M of rank at most the Cantor-Bendixson rank of K , and some point-finite and Lipschitz assignment $\mu : \mathcal{B} \rightarrow c_{00}^+$ such that*

- (a) *$\text{supp } \mu(s) = s$ for every $s \in \mathcal{B}$, and*
- (b) *the sequence of linear combinations $(x(s_n))_n = (\sum_{i \in s_n} (\mu(s_n))(i) x_i)_n$ is a normalized block sequence C -equivalent to the unit vector basis of c_0 for every block sequence $(s_n) \subseteq \mathcal{B}$.*

PROOF. The proof is by induction on the Cantor-Bendixson rank of K . By Lemma 5.7 we may assume that $K \subseteq c_{00}$ is weakly-compact is such that $\text{supp } K = \overline{\mathcal{B}}$ with \mathcal{B} a α -uniform barrier, and that the weakly null sequence (x_i) is the evaluation mapping sequence (p_i) which is a monotone basic sequence. Set $\mathcal{F}_n = \overline{\bigcup_{m \leq n} \mathcal{B}_{\{n\}}}$ ($n \in \mathbb{N}$). Since \mathcal{B} is a α -uniform family, we have that for every n , $\partial^\alpha \mathcal{F}_n = \emptyset$, so its Cantor-Bendixson rank is strictly smaller than $\alpha + 1$. Set $K_n = \{f \upharpoonright s : s \in \mathcal{F}_n\}$ ($n \in \mathbb{N}$). This is a compactum K_n with $\text{supp } K_n = \mathcal{F}_n$ and whose rank is strictly smaller than $\alpha + 1$. So, the evaluation mapping sequence (p_i) is a weakly-null sequence of $C(K_n)$ for every n . Observe that for every sequence of scalars (a_i) we have that

$$\left\| \sum_i a_i p_i \right\|_{C(K_n)} = \sup \left\{ \left\| \sum_{i \in s} a_i p_i \right\|_{C(K)} : s \in \mathcal{F}_n \right\}. \quad (50)$$

Since in addition we are assuming that $\|p_i\|_{C(K)} = 1$ for every i , this implies that for every S , using that \mathcal{F}_n is hereditary, we obtain that (p_i) is 1-unconditional. Since we assume that

the singletons belong to \mathcal{F}_n , we obtain that (p_i) is a 1-unconditional normalized weakly null sequence in $C(K_n)$.

Fix $\varepsilon > 0$, and let $(\varepsilon_n)_n$ be a summable sequence with $\sum_n \varepsilon_n < \varepsilon/2$. By the Ramsey property of the uniform barrier \mathcal{B} , we can find a fusion sequence (M_k) such that, setting $n_k = \min M_k$ ($k \in \mathbb{N}$), we have that for every k the following dichotomy holds:

(1) Either for every $s \in \mathcal{B} \upharpoonright M_k$ there is some $\mu_k(s) \in c_{00}$ with $\text{supp } \mu_k(s) = s$, $\mu_k(s)(i) > 0$ ($i \in s$) and such that for every such that $\|\sum_{i \in s} \mu_k(s)(i)p_i\| = 1$ while $\|\sum_{i \in s} \mu_k(s)(i)p_i\|_{n_k} < \varepsilon_k$, or else

(2) $\|\sum_{i \in s} a_i p_i\| \leq 2\varepsilon_k^{-1} \|\sum_{i \in s} a_i p_i\|_{n_k}$ for every $s \in \mathcal{B} \upharpoonright M_k$ and every $(a_i)_{i \in s}$.

Suppose first that the (1) holds for every k . Let $M_\infty = \{n_k\}$ be the fusion set. Then let $\mathcal{C} = \mathcal{B} \upharpoonright M_\infty$. For $s \in \mathcal{C}$, define $\mu(s) = \mu_k(s)$, where $n_k = \min s$. This is well defined since $s \in \mathcal{B} \upharpoonright M_k$. Fix a block sequence $(s_i) \subseteq \mathcal{C}$, and let us work to show that $(x(s_i))$ is $2 + \varepsilon$ -equivalent to the c_0 -basis, where we recall that for a given $s \in \mathcal{C}$, $x(s) = \sum_{i \in s} \mu(s)(i)p_i$. Fix a sequence of scalars $(b_i)_{i \in M_\infty}$, $|b_i| \leq 1$ ($i \in \mathbb{N}$). Since each $\mu(s_i)$ is normalized and since (p_i) is monotone, we obtain that $\|\sum b_i \mu(s_i)\|_{C(K)} \geq (1/2) \|\sum b_i e_i\|_\infty$. Suppose that $\xi \in K$, and let $i_0 = \min\{i : s_i \cap \text{supp } \xi \neq \emptyset\}$. Fix $i > i_0$, and let k_i be such that $n_{k_i} = \min s_i$. Since $\text{supp } \xi \cap s_i \in \mathcal{F}_{\max s_{i_0}}$ we have that

$$|\mu(s_i)(\xi)| \leq \left\| \sum_{j \in s_i \cap \text{supp } \xi} a_i^{(k_i)} p_j \right\|_{C(K_{\max s_{i_0}})} < \varepsilon_{k_i}. \quad (51)$$

It follows that

$$\left| \sum_i b_i \mu(s_i)(\xi) \right| \leq |b_{i_0}| + \sum_{i > i_0} |b_i| |\mu(s_i)(\xi)| \leq |b_{i_0}| + \frac{\varepsilon}{2}. \quad (52)$$

So, $\|\sum_i b_i p_i\|_{C(K)} \leq (1 + \varepsilon/2) \|\sum b_i e_i\|_\infty$. Finally use Corollary 5.6 to perturb μ and make it point-finite and Lipschitz.

Suppose now that k_0 is the first k such that (2) holds for k . Set $M = M_k$. It readily follows that for every $x \in \langle f_n \rangle_{n \in M}$ we have that $\|x\|_{C(K)} \leq \varepsilon_{k_0}^{-1} \|x\|_{C(K_{n_{k_0}})}$. By inductive hypothesis applied to $(p_i) \subseteq C(K_{n_{k_0}})$, there is some $C \geq 1$, some uniform barrier \mathcal{C} on some $N \subseteq M$ of rank not bigger than the one of $K_{n_{k_0}}$ and some μ fulfilling the conclusions of the Lemma. Fix $s \in \mathcal{C}$. Then $\|\mu(s)\|_{n_{k_0}} = 1$, so we can find some $t \subseteq s$ such that $1 = \|\mu(s)\|_{C(K_{n_{k_0}})} = \|\mu(s) \upharpoonright t\|_{C(K)}$. Observe that, by 1-unconditionality of $\|\cdot\|_{C(K_{n_{k_0}})}$, $\|\mu(s) \upharpoonright t\|_{C(K_{n_{k_0}})} = 1$. Define $\nu(s) = \mu(s) \upharpoonright t$. Let us check that indeed $(\nu(s_n)) \subseteq C(K)$ is $C\varepsilon_{n_k}^{-1}$ -equivalent to the c_0 basis. Fix scalars (a_i) , $|a_i| \leq 1$ ($i \in \mathbb{N}$). We obtain the inequality $\|\sum_i a_i \nu(s_i)\|_{C(K)} \geq (1/2) \|\sum_i a_i e_i\|_\infty$ by the monotonicity of the basic sequence (p_i) . Now,

$$\left\| \sum_i a_i \nu(s_i) \right\|_{C(K)} \leq \frac{1}{\varepsilon_{n_{k_0}}} \left\| \sum_i a_i \nu(s_i) \right\|_{C(K_{n_{k_0}})} \leq \frac{1}{\varepsilon_{n_{k_0}}} \left\| \sum_i a_i \mu(s_i) \right\|_{C(K_{n_{k_0}})} \leq \frac{C}{\varepsilon_{n_{k_0}}} \left\| \sum_i a_i e_i \right\|_\infty. \quad (53)$$

□

5.3. Conditionality. We start with the following natural variation on the notion of an unconditional sequence in a Banach space E .

Definition 5.10. Let \mathcal{F} be a family of finite sets of integers. A normalized basic sequence (x_n) of a Banach space E is called \mathcal{F} -unconditional with constant at most $C \geq 1$ iff for every sequence of scalars (a_n) ,

$$\sup_{s \in \mathcal{F}} \left\| \sum_{n \in s} a_n x_n \right\| \leq C \left\| \sum_{n \in \mathbb{N}} a_n x_n \right\|.$$

Clearly, (x_n) is unconditional with constant C if it is FIN-unconditional with constant C . Thus, if the family \mathcal{F} has a trace $\mathcal{F}[M]$ on an infinite set M consisting of all finite subsets of M , the corresponding subsequence $(x_n)_{n \in M}$ is unconditional with constant C . A hereditary family with no such a trace is compact, so one is naturally led to examining the standard compact families of finite subsets of \mathbb{N} . Recall, the notion of a Schreier family introduced above in Remark 3.4: $\overline{\mathcal{S}} = \{s \subseteq \mathbb{N} : |s| \leq \min(s) + 1\}$.

Proposition 5.11. Suppose that (x_n) is a normalized weakly-null sequence of a Banach space E . For every $\varepsilon > 0$ there is some M such that $(x_n)_{n \in M}$ is $(2 + \varepsilon)$ -Schreier unconditional.

PROOF. We assume that (x_n) is the evaluation mapping sequence $(p_n) \subseteq C(K)$ for some $K \subseteq c_0$ weakly-compact bounded by 1 and closed under restriction on initial intervals. Using a finite ε -net of $[-1, 1]^n$ with the ℓ_1 -norm, we can use repeatedly Corollary 3.21 and find a fusion sequence (M_k) such that, setting $m_k = \min M_k$, for every k every $s \in M_k^{[m_k]}$ and every $\xi \in K$ there is some $\eta \in K$ such that $\|\xi \upharpoonright s - \eta \upharpoonright s\|_{\ell_1} \leq \varepsilon/2$, and $\sum_{i \in M_k \setminus s} |\eta(i)| < \varepsilon/2$. Let M be the fusion set of (M_k) . Notice that M has the following property: For every $s \in \mathcal{S} \upharpoonright M$ and every $\xi \in K$ there is $\eta \in K$ such that $\|\xi \upharpoonright s - \eta \upharpoonright s\|_{\ell_1} \leq \varepsilon/2$, and $\sum_{i \in (M/m_{k-1}) \setminus s} |\eta(i)| < \varepsilon/2$, where $m_k = \min s$.

We claim that M is the desired set: Fix scalars $(a_i)_{i \in M}$ and $s \in \mathcal{S} \upharpoonright M$. We may assume that $\|\sum_{i \in \mathbb{N}} a_i p_i\|_{C(K)} = 1$. Let $\xi \in K$ be such that $|\sum_{i \in s} a_i \xi(i)| = \|\sum_{i \in s} a_i p_i\|_{C(K)}$. Find η as above for the pair (s, ξ) . Then

$$\left| \sum_{i \in s} a_i \xi(i) \right| \leq \left| \sum_{i \in s} a_i \eta(i) \right| + \frac{\varepsilon}{2} \leq \left| \sum_{i \in M/m_{k-1}} a_i \eta(i) \right| + \varepsilon \leq (2 + \varepsilon) \left\| \sum_{i \in M} a_i p_i \right\|_{C(K)}, \quad (54)$$

where $m_k = \min s$. □

Recall that if \mathcal{F} is a barrier on some set M then its trace $\mathcal{F}[N]$ on any co-infinite subset N of M is hereditary and that for every pair \mathcal{F}_0 and \mathcal{F}_1 of barriers on the same domain M there is an infinite set $N \subseteq M$ such that $\mathcal{F}_0[N] \subseteq \mathcal{F}_1[N]$ or $\mathcal{F}_1[N] \subseteq \mathcal{F}_0[N]$. Since the two alternatives are dependent on the ranks of \mathcal{F}_0 and \mathcal{F}_1 , one is naturally led to the following measurement of unconditionality.

Definition 5.12. Suppose that γ is a countable ordinal. A normalized basic sequence (x_n) of a Banach space E is called γ -unconditionally saturated with constant at most $C \geq 1$ if for every γ -uniform barrier \mathcal{B} on \mathbb{N} and for every infinite M there is infinite $N \subseteq M$ such that the corresponding subsequence $(x_n)_{n \in N}$ of (x_n) is $\overline{\mathcal{B}} \upharpoonright \overline{N}$ -unconditional with constant at most C .

REMARK 5.13. (a) We have seen above in Proposition 5.11 that every normalized basic sequence has a subsequence which is ω -unconditionally saturated with constant at most $2 + \varepsilon$. An analysis of the Maurey-Rosenthal[14] example of a weakly-null sequence (x_n) with no unconditional basic

subsequence (see Example 5.14 below) reveals an ω^2 -uniform barrier \mathcal{B}_{MR} such that no infinite subsequence $(x_n)_{n \in \mathbb{N}}$ is $\mathcal{B}_{\text{MR}} \upharpoonright M$ -unconditional with any finite constant C .

(b) Recall that an ordinal γ is called indecomposable if for every $\beta < \gamma$, $\beta\omega \leq \gamma$. Equivalently, $\gamma = \omega^\beta$ for some β . Suppose that γ is the maximal indecomposable ordinal smaller than some fixed ordinal α . Then a normalized basic sequence (x_n) is α -unconditionally saturated if and only if it is γ -unconditionally saturated.

(c) If the normalized basic sequence (x_n) is monotone, then, it is $\overline{\mathcal{B}}$ -unconditional iff it is \mathcal{B} -unconditional for every uniform barrier \mathcal{B} on \mathbb{N} .

EXAMPLE 5.14. First of all, for a fixed $0 < \varepsilon < 1$ choose a fast increasing sequence (m_i) such that

$$\sum_{i=0}^{\infty} \sum_{j \neq i} \min\left(\left(\frac{m_i}{m_j}\right)^{1/2}, \left(\frac{m_j}{m_i}\right)^{1/2}\right) \leq \frac{\varepsilon}{2}. \quad (55)$$

Let $\text{FIN}^{[<\infty]}$ be the collection of all finite block sequences $E_0 < E_1 < \dots < E_k$ of nonempty finite subsets of \mathbb{N} . Now choose a 1 – 1 function

$$\sigma : \text{FIN}^{[<\infty]} \rightarrow \{m_i\} \quad (56)$$

such that $\varphi((s_i)_{i=0}^n) > s_n$ for all $(s_i) \in \text{FIN}^{[<\infty]}$. Now let \mathcal{B}_{MR} be the family of unions $s_0 \cup s_1 \cup \dots \cup s_n$ of finite sets such that

- (a) (s_i) is block and $s_0 = \{n\}$.
- (b) $|s_i| = \sigma(s_0, \dots, s_{i-1})$ ($1 \leq i \leq n$).

It turns out that \mathcal{B}_{MR} is a ω^2 -uniform barrier on \mathbb{N} (see Proposition 5.19 below), hence $\overline{\mathcal{B}_{\text{MR}}} = \overline{\mathcal{B}_{\text{MR}}}^\square$ is a compact family with rank $\omega^2 + 1$. Observe that by definition, every $s \in \mathcal{B}_{\text{MR}}$ has a unique decomposition $s = \{n\} \cup s_1 \cup \dots \cup s_n$ satisfying (a) and (b) above. Now define the mapping $\Phi : \mathcal{B}_{\text{MR}} \rightarrow c_{00}$,

$$\Phi(s) = e_n + \sum_{i=1}^n \frac{1}{|s_i|^{1/2}} \sum_{k \in s_i} e_k. \quad (57)$$

Observe that we have the property that if $s \cap t \subseteq s, t$, then $\Phi(s) \upharpoonright (s \cap t) = \Phi(t) \upharpoonright (s \cap t)$, so there is a unique extension $\Phi : \overline{\mathcal{B}_{\text{MR}}} \rightarrow c_{00}$, naturally defined by $\Phi(s) = \Phi(t) \upharpoonright t$, where $t \in \mathcal{B}_{\text{MR}}$ is (any) such that $s \subseteq t$. Now define $K = \Phi''\mathcal{B}_{\text{MR}} \subseteq c_{00}$. This is a weakly-compact subset of c_{00} , with rank the same than $\overline{\mathcal{B}_{\text{MR}}}$, i.e., $\omega^2 + 1$. Then $(p_i) \subseteq C(K)$ is a normalized weakly-null sequence with no unconditional subsequence. Moreover this weakly-null sequence has the property that the summing basis (S_i) of c , the Banach space of *convergent sequences of reals*, is finitely-block representable in the linear span of every subsequence of (p_i) (and so the summing basis of c_0), more precisely, for every M , every $n \in \mathbb{N}$ and every $\varepsilon > 0$ there is a normalized block subsequence $(x_i)_{i=0}^{n-1}$ of $(p_i)_{i \in M}$ such that for every sequence of scalars $(a_i)_{i=0}^{n-1}$,

$$\max\left\{\left|\sum_{i=0}^m a_i\right| : m < n\right\} \leq \left\|\sum_{i=0}^{n-1} a_i x_i\right\|_{C(K)} \leq (1 + \varepsilon) \max\left\{\left|\sum_{i=0}^m a_i\right| : m < n\right\}.$$

On the other hand, by Proposition 5.11 the sequence (p_i) is *Schreier-unconditionally saturated* with constant ~ 2 , i.e., for every M there is some $N \subseteq M$ such that for every element $s \in \mathcal{S}[N]$,

and every sequence of scalars $(a_i)_{i \in N}$,

$$\left\| \sum_{i \in s} a_i p_i \right\|_{C(K)} \leq (2 + \varepsilon) \left\| \sum_{i \in N} a_i p_i \right\|_{C(K)}.$$

Building on the idea of this Example, we are now going to find, for every countable indecomposable ordinal γ , a normalized weakly-null bimonotone basic sequence with no unconditional subsequences but β -unconditionally saturated for every $\beta < \gamma$. Before embarking into the construction, we need to recall a localized version of Pták's Lemma. For this we need the following notation: Given a family \mathcal{F} , and $n \in \mathbb{N}$, let $\mathcal{F} \otimes n = \{s_0 \cup \dots \cup s_{n-1} : (s_i)_{i=0}^{n-1} \subseteq \mathcal{F} \text{ is block}\}$. It can be shown that $\mathcal{F} \otimes n$ is a αn -uniform family if \mathcal{F} is an α -uniform family.

Given $\xi \in c_{00}$ we will write $\xi^{1/2}$ to denote $(\xi(i)^{1/2})$. Given $\xi \in c_{00}$ and a finite set s , let $\langle \xi, s \rangle = \langle \xi, \chi_s \rangle = \sum_{i \in s} \xi(i)$.

Lemma 5.15. *Suppose that \mathcal{B} is an α -uniform barrier on M , $\alpha \geq 1$. Let $\gamma = \gamma(\alpha)$ be the maximal indecomposable ordinal not bigger than α , and let $n = n(\alpha) \in \mathbb{N}$, $n \geq 1$, be such that $\gamma n \leq \alpha < \gamma(n+1)$. Then for every $k \in \mathbb{N}$, $k > 1$, every $\varepsilon > 0$, and every β -uniform barrier \mathcal{C} on M with $\beta > \alpha k$ there $N \subseteq M$ and some point-finite Lipschitz mean-assignment $\mu : \mathcal{C} \upharpoonright N \rightarrow c_{00}^+$ such that*

$$\sup\{\langle \mu(s)^{\frac{1}{2}}, t \rangle : t \in \mathcal{B}\} \leq \frac{(1 + \varepsilon)(n+1)}{(nk)^{\frac{1}{2}}} \quad (58)$$

for every $s \in \mathcal{C} \upharpoonright N$.

PROOF. The proof now is by induction on α . Fix $\varepsilon > 0$ and $k > 1$. Let \mathcal{C} be an β -uniform family on M such that $\beta > \alpha k$.

Notice that if we prove that for every $N \subseteq M$ there is one mean μ with support in $\mathcal{C} \upharpoonright N$ such that (58) holds, then the Ramsey property of the uniform barrier \mathcal{C} gives the existence for some $N \subseteq M$ of a mean-assignment $\mu : \mathcal{C} \upharpoonright N \rightarrow c_{00}$ satisfying (58) for every $t \in \mathcal{B}$. Then Corollary 5.6 gives the desired point-finite Lipschitz assignment.

Let \mathcal{D} be a γ -uniform barrier on M (if $n = 1$ we take $\mathcal{D} = \mathcal{B}$), and fix $N \subseteq M$. Find first $P \subseteq N$ be such that $(\mathcal{D} \otimes nk) \upharpoonright P \subseteq \overline{\mathcal{C}}$ as well as $\mathcal{B} \upharpoonright P \subseteq \overline{\mathcal{D} \otimes (n+1)}$. Consider $(\gamma_i)_{i \in P}$ such that $\mathcal{D}_{\{i\}} \upharpoonright P$ is γ_i -uniform on P/i . Observe that for every $i \in P$ we have that $\gamma_i < \gamma$, so, since γ is indecomposable, $\gamma_i \omega \leq \gamma$. Let μ_0 be any mean such that $\text{supp } \mu_0 \in \mathcal{B} \upharpoonright P$. By inductive hypothesis applied to appropriate α_i 's, we can find a block sequence $(\mu_j)_{j=0}^{nk-1}$ of means with support in $\mathcal{B} \upharpoonright P$ such that for every $1 \leq j \leq nk-1$,

$$\sup\{\langle \mu_j^{\frac{1}{2}}, t \rangle : t \in \mathcal{D}, \text{ and } \min t \leq \max \text{supp } \mu_{j-1}\} < \frac{\varepsilon}{2^{j+1}}. \quad (59)$$

Let $\nu = (1/(nk)) \sum_{j=0}^{nk-1} \mu_j$. Observe that $\text{supp } \nu \in \overline{(\mathcal{D} \otimes (nk)) \upharpoonright P} \subseteq \overline{\mathcal{C}}$. Then, for every $t \in \mathcal{B}$, by (59),

$$\langle \nu^{\frac{1}{2}}, t \rangle = \frac{1}{(nk)^{\frac{1}{2}}} \sum_{j=0}^{k-1} \sum_{i \in t} \mu_j(i)^{\frac{1}{2}} \leq \frac{1 + \frac{\varepsilon}{2}}{(nk)^{\frac{1}{2}}}. \quad (60)$$

Let us point out that $\text{supp } \nu$ is, possibly, not a set in \mathcal{C} . However it is easy to slightly perturb ν to a newer mean with support in \mathcal{C} and satisfying (60) for every $t \in \mathcal{B}$: Let $s \in \mathcal{C}$ be such that

$\text{supp } \nu \sqsubseteq s$, and set $u = s \setminus \text{supp } \nu$. Let $\delta > 0$ be such that

$$(1 + \frac{\varepsilon}{2})(1 - \delta)^{1/2} + (nk\delta|u|)^{1/2} \leq 1 + \varepsilon. \quad (61)$$

Now set

$$\mu = (1 - \delta)\nu + \frac{\delta}{|u|}\chi_u. \quad (62)$$

μ is a mean whose support is $s \in \mathcal{C}$. It can be shown now that for every $t \in \mathcal{B}$,

$$\sum_{i \in t} \mu(i)^{\frac{1}{2}} \leq \frac{1 + \varepsilon}{(nk)^{\frac{1}{2}}}, \quad (63)$$

by the choice of δ . Finally, let $t \in \mathcal{B}$ and let us compute $\sum_{i \in t} (\mu(i))^{1/2}$: First of all we have that $\sum_{i \in t} (\mu(i))^{1/2} = \sum_{i \in u} (\mu(i))^{1/2}$, where $u = t \cap P$. Now, since $u \in \overline{\mathcal{B} \upharpoonright P} \subseteq \overline{\mathcal{D} \otimes (n+1)}$, we can find $t_0 < \dots < t_n$ in \mathcal{D} such that $u \sqsubseteq t_0 \cup \dots \cup t_n$, and hence

$$\langle \mu^{1/2}, t \rangle = \sum_{j=0}^n \langle \mu^{1/2}, t_j \rangle \leq \frac{(n+1)(1 + \varepsilon)}{(nk)^{\frac{1}{2}}}, \quad (64)$$

as promised. \square

Corollary 5.16. *Suppose that \mathcal{B} is an α -uniform barrier on M , $\alpha \geq 1$. Then for every $\varepsilon > 0$ there is some $k = k(\alpha, \varepsilon)$ such that for every β -uniform barrier on M with $\beta > \alpha k$ there $N \subseteq M$ and some point-finite Lipschitz mean-assignment $\mu : \mathcal{C} \upharpoonright N \rightarrow c_{00}^+$ such that,*

$$\sup\{\langle \mu(s)^{1/2}, t \rangle : t \in \mathcal{B}\} \leq \varepsilon \quad (65)$$

for every $s \in \mathcal{B} \upharpoonright N$. \square

Lemma 5.17. *Fix an indecomposable countable α and a sequence (ε_n) of positive reals. Then:*
 (a) *there is a collection (\mathcal{B}_n) of α_n -uniform barriers on \mathbb{N}/n and a corresponding sequence of point-finite Lipschitz mean-assignments $\mu_n : \mathcal{B}_n \rightarrow c_{00}^+$ with the following properties:*

- (a.1) $\alpha_n > 0$, $\sup_n \alpha_n = \alpha$,
- (a.2) for every $m < n$ and every $s \in \mathcal{B}_n$

$$\sup\{\langle \mu_n(s)^{\frac{1}{2}}, t \rangle : t \in \mathcal{B}_m\} < \varepsilon_n. \quad (66)$$

(b) *Suppose that in addition $\alpha = \omega^\gamma$ with γ limit. Let $\alpha_n \uparrow \alpha$ be any sequence such that $\alpha_n \omega \leq \alpha_{n+1}$ ($n \in \mathbb{N}$). Then there is a double sequence (\mathcal{B}_i^n) such that*

- (b.1) \mathcal{B}_i^n is an $\alpha_i^{(n)}$ -uniform barrier on $\mathbb{N}/(n+i)$, with $\alpha_i^{(n)} > 0$ and $\alpha_i^{(n)} \uparrow_i \alpha_n$ ($n \in \mathbb{N}$).
- (b.2) *There are point-finite Lipschitz mean-assignments $\mu_{n,i} : \mathcal{B}_i^n \rightarrow c_{00}$ such that for every $s \in \mathcal{B}_i^n$, and every $(m, j) <_{\text{lex}} (n, i)$*

$$\sup\{\langle \mu_{n,i}(s)^{\frac{1}{2}}, t \rangle : t \in \mathcal{B}_j^m\} < \varepsilon_{n+i}, \quad (67)$$

where we recall that $<_{\text{lex}}$ denotes the lexicographical order on \mathbb{N}^2 defined by $(m, i) <_{\text{lex}} (n, j)$ iff $m < n$, or $m = n$ and $i < j$.

PROOF. (a): Choose $\alpha_n \uparrow_n \alpha$ such that for every $n \in \mathbb{N}$, $\alpha_{n+1} > \alpha_n k(\alpha_n, \varepsilon_n)$, that is possible since α is indecomposable. Let \mathcal{C}_n be an α_n -uniform family on \mathbb{N} ($n \in \mathbb{N}$). By Corollary 5.16 we can find a fusion sequence (M_n) such that

(c) $\mathcal{C}_m \upharpoonright M_m \subseteq \overline{\mathcal{C}_n}$ if $m \leq n$, and

(d) for every $n \in \mathbb{N}$ there is a point-finite Lipschitz mean-assignment $\nu_n : \mathcal{C}_n \upharpoonright M_n \rightarrow c_{00}^+$ such that

$$\sup\{\langle \nu_n(s)^{\frac{1}{2}}, t \rangle : t \in \bigcup_{l < n} \mathcal{C}_l\} < \varepsilon_n \quad (68)$$

for every $s \in \mathcal{C}_n \upharpoonright M_n$. Let $M = \{m_n\}$ be the fusion set of (M_n) , and $\Theta : M \rightarrow \mathbb{N}$ be the corresponding order preserving onto mapping. It is not difficult to see that $\mathcal{C}_n = (\Theta'' \mathcal{B}_n) \upharpoonright (\mathbb{N}/n)$, and $\mu_n : \mathcal{C}_n \rightarrow c_{00}$ defined naturally out of ν_n and Θ fulfils all the requirements.

(b): Suppose that $\alpha = \omega^\gamma$ with γ limit. Let $\alpha_n \uparrow \alpha$ be any sequence such that $\alpha_n \omega \leq \alpha_{n+1}$ ($n \in \mathbb{N}$).

Claim. *There is a fusion sequence (M_n) , $M_n = \{m_i^{(n)}\}$, a double sequence $(\mathcal{B}_i^{(n)})$ of $\alpha_i^{(n)}$ -uniform barriers on $M_n/m_i^{(n)}$ and point-finite Lipschitz mean-assignments $\mu_{n,i} : \mathcal{B}_i^{(n)} \rightarrow c_{00}^+$ such that*

(e) $\alpha_i^{(n)} \uparrow_i \alpha_n$ ($n \in \mathbb{N}$), and

(f) for every $(m, j) <_{\text{lex}} (n, i)$, every $s \in \mathcal{B}_i^{(n)}$ and every $t \in \mathcal{B}_j^{(m)}$, $\langle (\mu_{n,i}(s))^{1/2}, t \rangle < \varepsilon_{n+i}$.

Proof of Claim: First, using Corollary 5.16 applied to α_0 to produce an infinite set $M_0 = \{m_i^{(0)}\}$ and a sequence $(\mathcal{B}_i^{(0)})$ of $\alpha_i^{(0)}$ -uniform barriers on $M_0/\{m_i^{(0)}\}$ with $\alpha_i^{(0)} \uparrow \alpha_0$ and point-finite Lipschitz mean-assignments $\mu_{0,i} : \mathcal{B}_i^{(0)} \rightarrow c_{00}$ such that for every i and every $s \in \mathcal{B}_i^{(0)}$, $\langle \mu_{0,i}(s)^{1/2}, t \rangle \leq \varepsilon_i$ for every $t \in \mathcal{B}_j^{(0)}$ with $j < i$. In general, suppose we have found for every $k \leq n$ $M_k = \{m_i^{(k)}\} \subseteq M_{k-1}$, $(\mathcal{B}_i^{(k)})$ $\alpha_i^{(k)}$ -uniform barriers on $M_k/m_i^{(k)}$ and point-finite Lipschitz mean-assignments $\mu_{k,i} : \mathcal{B}_i^{(k)} \rightarrow c_{00}$ such that for every $(k, j) <_{\text{lex}} (m, i)$ every $s \in \mathcal{B}_i^{(m)}$ and every $t \in \mathcal{B}_j^{(k)}$ $\langle \mu_{m,i}(s)^{1/2}, t \rangle \leq \varepsilon_{m+i}$. For each $k \leq n$ define the following families

$$\mathcal{B}_k = \{s \subseteq M_k : *s \in \mathcal{B}_{\min s}^k\}. \quad (69)$$

This is clearly an α_k -uniform family on M_k . Since $\alpha_n \omega \leq \alpha_{n+1}$, we can use again Corollary 5.16 and find an infinite subset $M_{n+1} = \{m_i^{(n+1)}\} \subseteq M_n$ and a sequence $(\mathcal{B}_i^{(n+1)})$ of $\alpha_i^{(n+1)}$ -uniform barriers on $M_{n+1}/m_i^{(n+1)}$ and point-finite Lipschitz mean-assignments $\mu_{n+1,i} : \mathcal{B}_i^{(n+1)} \rightarrow c_{00}$ such that for every $s \in \mathcal{B}_i^{(n+1)}$,

$$\sup\{\langle (\mu_{n+1,i}(s))^{\frac{1}{2}}, t \rangle : t \in \bigcup_{k \leq n} \mathcal{B}_m \cup \bigcup_{j < i} \mathcal{B}_j^{(n+1)}\} < \varepsilon_{n+i+1}, \quad (70)$$

so, in particular for every $k \leq n$ and every $t \in \mathcal{B}_j^{(k)}$, $\langle (\mu_{n+1,i}(s))^{\frac{1}{2}}, t \rangle < \varepsilon_{n+i+1}$. \square

Let M be the fusion set of (M_n) , i.e. $M = \{m_0^{(n)}\}$. Observe that $m_0^{(n+i)} \geq m_i^{(n)}$ for every n and i , so $M/m_0^{(n)} \subseteq M_n/m_i^{(n)}$. Set $\mathcal{C}_i^n = \mathcal{B}_i^{(n)} \upharpoonright (M/m_0^{(n+i)})$. This is an $\alpha_i^{(n)}$ -uniform barrier on $M/m_0^{(n+i)}$. Consider $\nu_{n,i} = \mu_{n,i} \upharpoonright \mathcal{C}_i^n : \mathcal{C}_i^n \rightarrow c_{00}$ has the property that for every $(m, j) <_{\text{lex}} (n, i)$, every $s \in \mathcal{C}_i^n$ and every $t \in \mathcal{C}_j^m$, $\langle (\nu_{n,i}(s))^{1/2}, t \rangle < \varepsilon_{n+i}$. Now use $\Theta : M \rightarrow \mathbb{N}$, $\Theta(m_0^{(n)}) = 0$, to define the desired mean-assignments and families. \square

REMARK 5.18. Observe that if \mathcal{B} is α -uniform on M with $\alpha > 0$, then $M^{[1]} \subseteq \overline{\mathcal{B}}$. It readily follows that the mean-assignments μ_n and $\mu_{n,i}$ obtained in Lemma 5.17 have the property that $\|\mu_n(s)^{1/2}\|_\infty \leq \varepsilon_n$ and $\|\mu_{n,i}(s)^{1/2}\|_\infty \leq \varepsilon_{n+i}$ for every s in the corresponding domains.

Proposition 5.19. (a) Suppose that \mathcal{C} and \mathcal{B}_i are β and α_i -uniform families on M ($i \in \mathbb{N}$) with $\alpha_i \uparrow \alpha$, $\alpha_i, \beta \geq 1$. Let $\sigma : \text{FIN}^{[<\infty]} \rightarrow \mathbb{N}$ be 1-1. Then for every $n \in \mathbb{N}$ the family

$$\mathcal{D} = \{s_0 \cup \dots \cup s_n : (s_i) \text{ is block, } s_0 \in \mathcal{C} \text{ and } s_i \in \mathcal{B}_{\sigma((s_0, \dots, s_{i-1}))} \text{ for every } 1 \leq i \leq n-1\}$$

is γ -uniform on M , where $\gamma = \alpha n + \beta^-$ if $1 \leq \beta < \omega$ and $n > 0$, and $\gamma = \alpha n + \beta$ if $\beta \geq \omega$ or $n = 0$.

(b) Suppose that \mathcal{B}_i is α_i -uniform on M ($i \in \mathbb{N}$) with $\alpha_i \uparrow \alpha$. Let $\sigma : \text{FIN}^{[<\infty]} \rightarrow \mathbb{N}$ be 1-1. Then the family

$$\begin{aligned} \mathcal{C} = \{ \{n\} \cup s_0 \cup \dots \cup s_{n-1} : (\{n\}, s_0, \dots, s_{n-1}) \text{ is block, and} \\ s_i \in \mathcal{B}_{\sigma((\{n\}, s_0, \dots, s_{i-1}))} \text{ for every } 0 \leq i \leq n-1 \} \end{aligned}$$

is $\alpha\omega$ -uniform on M .

PROOF. (a): The proof is by induction on n . If $n = 0$, the result is clear. So suppose that $n > 0$. Now the proof is by induction on β . Suppose first that $\beta = 1$. Then $\mathcal{C} = M^{[1]}$, and so, for every $m \in M$

$$\begin{aligned} \mathcal{D}_{\{m\}} = \{s_1 \cup \dots \cup s_n : (s_1, s_2, \dots, s_n) \text{ is block, } s_1 \in \mathcal{B}_{\sigma(\{m\})} \text{ and} \\ s_i \in \mathcal{B}_{\sigma(\{m\}, s_1, s_2, \dots, s_{i-1})} \text{ for every } 2 \leq i \leq n-1\}, \end{aligned}$$

so, by inductive hypothesis, $\mathcal{D}_{\{m\}}$ is $\alpha(n-1) + \gamma_m$ -uniform on M/m , depending whether α_m is finite or infinite, but in any case with $\gamma_m \uparrow \alpha$. Hence \mathcal{D} is αn -uniform on M . The general case for $1 \leq \beta < \omega$ is shown in the same way.

Suppose now that $\beta \geq \omega$. Then for every $m \in M$

$$\begin{aligned} \mathcal{D}_{\{m\}} = \{t \cup s_1 \cup \dots \cup s_n : (t, s_1, \dots, s_n) \text{ is block, } t \in \mathcal{C}_{\{m\}} \text{ and} \\ s_i \in \mathcal{B}_{\sigma(\{m\} \cup t, s_1, \dots, s_{i-1})} \text{ for every } 1 \leq i \leq n-1\}, \end{aligned}$$

By inductive hypothesis, $\mathcal{D}_{\{m\}}$ is $\alpha n + \gamma_m$ -uniform on M/m , with $\gamma_m \uparrow \beta$, so \mathcal{D} is $\alpha n + \beta$ -uniform on M , as desired.

(b) follows easily from (a). □

Theorem 5.20. For every indecomposable ordinal α there is a weakly-compact $K \subseteq c_{00}$ such that

- (a) $K \subseteq B_{c_0}$ is point-finite with Cantor-Bendixson rank $\alpha + 1$, and with support included in the downwards closure of an α -uniform family on \mathbb{N} .
- (b) $(p_n) \subseteq C(K)$ is a normalized weakly-null 1-basic sequence, and
- (c) The Summing basis of c is $2 + \varepsilon$ -finitely representable in every subsequence of (p_n) ; hence no subsequence of (p_n) is unconditional, but
- (d) (p_n) is β -unconditionally saturated for every $\beta < \alpha$.

PROOF. Fix $\varepsilon > 0$, and let $\varepsilon_n = \varepsilon/2^{n+3}$. Suppose that $\alpha = \omega^\gamma$. There are two cases to consider. Suppose first that $\gamma = \beta + 1$. We apply Lemma 5.17 (a) to the indecomposable ordinal ω^β and (ε_n) to produce the corresponding sequences of barriers (\mathcal{C}_n) and point-finite Lipschitz mean-assignments $\nu_n : \mathcal{C}_n \rightarrow c_{00}$ ($n \in \mathbb{N}$) satisfying the conclusions (a.1) and (a.2) of the Lemma. If γ is limit, then we use the part (b) of that lemma to produce a double sequence (\mathcal{B}_i^n) and mean-assignments $\nu_{n,i} : \mathcal{C}_i^n \rightarrow c_{00}$ satisfying (b.1) and (b.2). In order to unify the two cases we set for n, i ,

$$\mathcal{B}_i^n = \begin{cases} \mathcal{C}_i & \text{if } \gamma \text{ is successor ordinal} \\ \mathcal{C}_i^n & \text{if } \gamma \text{ is limit ordinal} \end{cases}$$

and

$$\mu_{n,i} = \begin{cases} \nu_i & \text{if } \gamma \text{ is successor ordinal} \\ \nu_{n,i} & \text{if } \gamma \text{ is limit ordinal.} \end{cases}$$

Let $\sigma : \text{FIN}^{[<\infty]} \rightarrow \mathbb{N}$ be 1-1 mapping such that $\sigma((s_0, \dots, s_n)) > \max s_n$ for every block sequence (s_0, \dots, s_n) of finite sets. For each n define

$$\mathcal{C}_n = \{s_0 \cup \dots \cup s_{n-1} : (s_i) \text{ is block and } s_i \in \mathcal{B}_{\sigma((\{n\}, s_0, \dots, s_{i-1}))}^n \text{ for every } 0 \leq i \leq n\},$$

So, by Proposition 5.19, if $\alpha = \omega^{\beta+1}$, then \mathcal{C}_n is a $\omega^\beta(n-1) + \zeta$ -uniform family on \mathbb{N} , where ζ is such that $\mathcal{B}_{\sigma((\{n\}))}^n$ is ζ -uniform; while if $\alpha = \omega^\gamma$ with γ limit, then it is $\alpha_n(n-1) + \zeta$ where ζ is such that $\mathcal{B}_{\sigma((\{n\}))}^n$ is ζ -uniform. Now let

$$\mathcal{C} = \{s \in \text{FIN} : *s \in \mathcal{C}_{\min s}\}. \quad (71)$$

It turns out that \mathcal{C} is an α -uniform family on \mathbb{N} . Observe that every $s \in \mathcal{C}$ has a unique decomposition $s = \{n\} \cup s(0) \cup \dots \cup s(n-1)$ with $n = \min s$ and $s(i) \in \mathcal{B}_{\sigma(s[i])}$, and where $s[i] = (\{n\}, s_0, \dots, s_{i-1})$ ($0 \leq i \leq n-1$). Define $\Phi : \mathcal{C} \rightarrow c_{00}$ for $s = \{n\} \cup s(0) \cup \dots \cup s(n-1) \in \mathcal{C}$ by

$$\Phi(s) = e_n + \sum_{i=0}^{n-1} (\mu_{n, \sigma(s[i])}(s(i)))^{\frac{1}{2}}, \quad (72)$$

It is not difficult to see that Φ is point-finite and Lipschitz. So, since \mathcal{C} is a front, $\overline{\mathcal{C}} = \overline{\mathcal{C}}^\square$, hence Φ extends naturally to a continuous $\Phi : \overline{\mathcal{C}} \rightarrow c_{00}$. Let $K \subseteq c_{00}$ be the set of those ξ of the form

$$K = \{\Phi(s) \upharpoonright (I \cap t) : s \in \mathcal{B}, I \text{ is an interval and } t \subseteq s(i) \text{ for some } i \leq \min s\}$$

It is not difficult to see, using that $\Phi''\overline{\mathcal{C}}$ is weakly-compact, that $K \subseteq c_{00}$ is point-finite, weakly compact, closed under restrictions on intervals. Since Φ is point-finite we obtain that the Cantor-Bendixson rank of K is $\alpha + 1$.

For every $s \in \mathcal{B}$ and every $i \leq s$, set

$$\xi(s, i) = (\mu_{\min s, \sigma(s[i])}(s(i)))^{1/2}.$$

Notice that $\xi(s, i) \upharpoonright t \in K$ for every t .

Claim. For every $s, t \in \mathcal{C}$ and every $i \leq \min s$ and $j \leq \min t$, we have that

$$\langle \xi(t, j), \xi(s, i) \rangle \leq \begin{cases} \varepsilon_{\max\{\min s, \min t\}} & \text{if } t[j] \neq s[i] \\ 1 & \text{if } t[j] = s[i]. \end{cases}$$

Proof of Claim: Set $n = \min s$, $m = \min t$, and assume that $t[j] \neq s[i]$. Suppose first that $\alpha = \omega^{\beta+1}$. Then, by definition of the mean assignments, $\langle \xi(t, j), \xi(s, i) \rangle \leq \varepsilon_{\max\{\sigma(t[j]), \sigma(s[i])\}}$, but $\sigma(u_0, \dots, u_k) \geq \max u_k$ for every block sequence (u_i) , which derives into the desired inequality. Assume now that $\alpha = \omega^\gamma$, γ limit ordinal. If $\min s = \min t$, then $\langle \xi(t, j), \xi(s, i) \rangle \leq \varepsilon_{\min s + \max\{\sigma(t[j]), \sigma(s[i])\}} \leq \varepsilon_{\min s}$. While if $\min t \neq \min s$, say $\min t < \min s$, then $\langle \xi(t, j), \xi(s, i) \rangle \leq \varepsilon_{\min s + \sigma(s[i])} \leq \varepsilon_{\min s}$.

If $\sigma(s[i]) = \sigma(t[j]) = l$, then $\min s = \min t = n$, and

$$\langle \xi(s, i), \xi(t, j) \rangle \leq \|(\mu_{n,l}(s(i)))^{1/2}\|_{\ell_2} \|(\mu_{n,l}(t(j)))^{1/2}\|_{\ell_2} \leq 1, \quad (73)$$

since both are means. \square

It is clear that (p_n) is a normalized bimonotone basic sequence.

Claim. *The summing basis (S_n) of c is finitely block represented in any subsequence of (p_n) .*

Proof of Claim: Fix an infinite set M of integers, and $l \in \mathbb{N}$. Let $s \in \mathcal{B} \upharpoonright M/l$, $s = \{n\} \cup s(0) \cup \dots \cup s(n-1)$ its canonical decomposition, and set

$$x_i = \sum_{j \in s(i)} (\mu_{n, \sigma(s[i])}(s(i))(j))^{\frac{1}{2}} p_j \quad (74)$$

Observe that $x_i(\xi(s, i)) = \langle \xi(s, i), \xi(s, i) \rangle = 1$, so from the previous claim we obtain that $\|x_i\| = 1$. Now consider scalars $(a_i)_{i \leq n-1}$ with $\|\sum_{i \leq n-1} a_i S_i\|_\infty = 1$. Observe that this implies that $\max_{i \leq n-1} |a_i| \leq 2$. We want to show that

$$\left\| \sum_{0 \leq i \leq n-1} a_i S_i \right\|_\infty \leq \left\| \sum_{0 \leq i \leq n-1} a_i x_i \right\|_{C(K)} \leq (1 + \varepsilon) \left\| \sum_{0 \leq i \leq n-1} a_i S_i \right\|_\infty. \quad (75)$$

To get the left hand inequality, suppose that $\|\sum_{0 \leq i \leq n-1} a_i S_i\|_\infty = |\sum_{i \in I} a_i|$, where I is some initial interval of integers less than n . Let $\xi = \sum_{i \in I} \xi(s, i) \in K$, and observe that

$$\left(\sum_{i \leq n-1} a_i x_i \right)(\xi) = \left| \sum_{i \in I} a_i \right|. \quad (76)$$

Next, consider a typical element $\xi = \Phi(t) \upharpoonright (I \cap u) \in K$, where $t \in \mathcal{C}$, $j \leq \min s$, $u \subseteq t(j)$ and I is an interval of integers. Suppose first that $\min s = \min t$. Let $i_0 = \max\{i \leq n-1 : s(i) = t(i)\}$. Then by the previous claim we obtain

$$\begin{aligned} \left| \left(\sum_{i \leq n-1} a_i x_i \right)(\xi) \right| &\leq \left| \sum_{i \leq i_0, s(i) \cap I \neq \emptyset} a_i \right| + \sum_{i_0 < i, j \leq n-1} 2|\langle \xi(s, i), \xi(t, j) \rangle| \leq 2 \left\| \sum_{i \leq n-1} a_i S_i \right\|_\infty + 2n^2 \varepsilon_n \leq \\ &\leq (2 + \varepsilon) \left\| \sum_{i \leq n-1} a_i S_i \right\|_\infty. \end{aligned} \quad (77)$$

Suppose now that $n = \min s \neq \min t$, say $\min t < \min s$. Let $i_0 < n$, if possible, be such that $\min t \in s(i_0)$. Then,

$$\left| \left(\sum_{i \leq n-1} a_i x_i \right)(\xi) \right| \leq |a_{i_0}| \|\xi(s, i_0)\|_\infty + 2 \sum_{i_0 \leq i < n} \sum_{0 \leq j < \min t} \langle \xi(t, j), \xi(s, i) \rangle \leq 2\varepsilon_n + 2n^2 \varepsilon_n \leq \varepsilon. \quad (78)$$

\square

Finally, we are going to show that (p_n) is β -unconditionally saturated for every $\beta < \alpha$. We consider two cases:

CASE 1. $\alpha = \omega^{\beta+1}$. Let $\mathcal{D} = \{s \subseteq \mathbb{N} : *s \in \mathcal{B}_{\min s}^0\}$. This is an α -uniform family on \mathbb{N} . The next result implies that (p_n) is β -unconditionally saturated for every $\beta < \alpha$.

Claim. *The sequence $(p_n) \subseteq C(K)$ is \mathcal{D} -unconditional with constant at most $1 + \varepsilon$.*

Proof of Claim: Fix $t \in \mathcal{D}$, and let $(a_i)_{i \in \mathbb{N}}$ be scalars such that $\|\sum_{i \in \mathbb{N}} a_i p_i\|_{C(K)} = 1$. Fix a typical element $\xi = \Phi(s) \upharpoonright (I \cap u)$. Suppose first that $\min s \in t$. Then since $\sigma(s[i]) > \min s \geq \min t$ and $*t \in \mathcal{B}_{\min t}^0$ we obtain that

$$|(\sum_{j \in t} a_j \xi(j))| \leq |a_{\min s}| + \varepsilon \leq (1 + \varepsilon) \|\sum_i a_i p_i\|_{C(K)}. \quad (79)$$

Now suppose that $\min s \notin t$, but $s \cap t \neq \emptyset$ (otherwise $|\sum_{j \in t} a_j \xi(j)| = 0$). Let $i_0 = \min\{i \leq \min s : s(i) \cap t \neq \emptyset\}$. Then for every $i_0 < i < \min s$ we have that $\sigma(s[i]) > \max s_{i_0} \geq \min t$, so

$$|\sum_{j \in t} a_j \xi(s, i)(j)| < \varepsilon_{\sigma(s[i])}, \quad (80)$$

hence

$$|(\sum_{j \in t} a_j \xi(j))| \leq |\sum_{j \in t \cap s(i_0)} a_j \xi(s, i_0)(j)| + \sum_{i_0 < i < \min s} |\sum_{j \in t} a_j \xi(s, i)(j)| \leq (1 + \varepsilon) \|\sum_{i \in \mathbb{N}} a_i p_i\|_{C(K)}, \quad (81)$$

the last inequality because $\xi(s, i_0) \upharpoonright t \in K$. \square

CASE 2. $\alpha = \omega^\gamma$, γ a countable limit ordinal. The desired result follows from the following fact.

Claim. *For every $n \in \mathbb{N}$, the sequence (p_i) is \mathcal{B}_0^n -unconditional with constant at most $\max\{n, 1 + \varepsilon\}$.*

Proof of Claim: Fix $n \in \mathbb{N}$ and $t \in \mathcal{B}_0^n$. Let $(a_i)_{i \in \mathbb{N}}$ be scalars such that $\|\sum_{i \in \mathbb{N}} a_i p_i\|_{C(K)} = 1$. Fix a typical element $\xi = \Phi(s) \upharpoonright (I \cap t)$. Suppose first that $n \leq \min s$. Then in a similar manner that in CASE 1 one can show that

$$|\sum_{j \in t} a_j \xi(j)| \leq |a_{\min s}| + \varepsilon \leq (1 + \varepsilon) \|\sum_{j \in \mathbb{N}} a_j p_j\|_{C(K)}. \quad (82)$$

Suppose that $m = \min s < n$, then

$$|\sum_{j \in t} a_j \xi(j)| \leq |a_{\min s}| + \sum_{i=1}^m |\sum_{j \in s(i) \cap t \cap I} a_j \xi(s, i)(j)| \leq n \|\sum_{j \in \mathbb{N}} a_j p_j\|_{C(K)}, \quad (83)$$

the last inequality because $\xi(s, i) \upharpoonright (t \cap I) \in K$. \square

\square

REFERENCES

- [1] D. Alspach and S.A. Argyros, *Complexity of weakly null sequences*, Dissertationes Mathematicae, **321**, (1992), 1–44.
- [2] S.A. Argyros, S. Mercourakis and A. Tsarpalias, *Convex unconditionality and summability of weakly null sequences*, Israel J. Math. **107** (1998), 157–193
- [3] S. A. Argyros and S. Todorćević, *Ramsey methods in analysis*. Birkhäuser Verlag, Basel 2005.
- [4] S. A. Argyros and V. Kanellopoulos, *Determining c_0 in $C(K)$ spaces*, Preprint 2005.
- [5] A. Arvanitakis, *Weakly null sequences with an unconditional subsequence*. Preprint 2004.
- [6] C. Bessaga and A. Pełczyński, *On bases and unconditional convergence of series in Banach spaces*. Studia Math. **17** 1958 151–164.
- [7] C. Bessaga and A. Pełczyński, *Spaces of continuous functions. IV. On isomorphical classification of spaces of continuous functions*. Studia Math. **19** 1960 53–62.
- [8] P.G. Casazza and T.J. Shura, *Tsirelson's space*. Lecture Notes in Math., Vol.1363, Springer-Verlag, Berlin 1989.
- [9] J. Elton, *Thesis*, Yale University (1978).
- [10] I. Gasparis, *A dichotomy theorem for subsets of the power set of the natural numbers*. Proc. Amer. Math. Soc. **129** (2001), no. 3, 759–764.
- [11] I. Gasparis, E. Odell and B. Wahl, *Weakly null sequences in the Banach space $C(K)$* . Preprint 2004.
- [12] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces. I. Sequence spaces*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Vol. 92.
- [13] C. St. J. A. Nash-Williams, *On well-quasi-ordering transfinite sequences*, Proc. Cambridge Philos. Soc. **61** (1965), 33–39.
- [14] B. Maurey and H. P. Rosenthal, *Normalized weakly null sequence with no unconditional subsequence*. Studia Math. **61** (1977), no. 1, 77–98.
- [15] B. Maurey, *Une suite faiblement convergente vers zero sans sous-suite inconditionnelle*, Séminaire Maurey-Schwartz 1975-76, exposé IX (1976).
- [16] B. Maurey, *Quelques resultats concernant l'inconditionnalité*, Séminaire Maurey-Schwartz 1975-76, exposé XVI (1976).
- [17] E. Odell, *Applications of Ramsey theorems to Banach space theory*, Notes in Banach spaces, pp. 379–404, Univ. Texas Press, Austin, Tex., 1980.
- [18] A. Pełczyński and Z. Semadeni, *Spaces of continuous functions III. Spaces $C(\Omega)$ for Ω without perfect subsets*, Studia Math. **18** (1959), 211–222.
- [19] P. Pudlak and V. Rödl, *Partition theorems for systems of finite subsets of integers*, Discrete Math., **39**, (1982), 67–73.

EQUIPE DE LOGIQUE MATHÉMATIQUE, UNIVERSITÉ PARIS 7, 2 PLACE JUSSIEU, 75251 PARIS CEDEX 05, FRANCE.

E-mail address: abad@logique.jussieu.fr

C.N.R.S., U.M.R. 7056 ET UNIVERSITÉ PARIS 7, U.F.R. DE MATHÉMATIQUES, CASE 7012, 2 PLACE JUSSIEU, 75251 PARIS, CEDEX 5, FRANCE.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, CANADA, M5S 3G3

E-mail address: stevo@math.toronto.edu