

A FOURTH ORDER CURVATURE FLOW ON A CR 3-MANIFOLD

*SHU-CHENG CHANG¹, *JIH-HSIN CHENG², AND HUNG-LIN CHIU³

ABSTRACT. Let $(\mathbf{M}^3, J, \theta_0)$ be a closed pseudohermitian 3-manifold. Suppose the associated torsion vanishes and the associated Q -curvature has no kernel part with respect to the associated Paneitz operator. On such a background pseudohermitian 3-manifold, we study the change of the contact form according to a certain version of normalized Q -curvature flow. This is a fourth order evolution equation. We prove that the solution exists for all time and converges smoothly to a contact form of zero Q -curvature. We also consider other background conditions and obtain a priori bounds up to high orders for the solution.

1. Introduction

Let (M, J, θ) be a closed (i.e., compact with no boundary) pseudohermitian 3-manifold (see Appendix A for basic notions in pseudohermitian geometry). In the papers [FH], [GG], [H], and [GL], the (CR) Paneitz operator P (acting on a smooth real function λ) with respect to (J, θ) is defined by

$$P\lambda = \Delta_b^2 \lambda + T^2 \lambda + 4 \operatorname{Im}(A_{\overline{11}} \lambda_{11} + A_{\overline{11},1} \lambda_1),$$

and the so-called Q -curvature is defined by

$$(1.1) \quad Q = -\frac{2}{3}(\Delta_b W + 2 \operatorname{Im} A_{11, \overline{11}})$$

where Δ_b , T , W , and $A_{\overline{11}}$ denote the sub-Laplacian, the characteristic vector field, the Tanaka-Webster (scalar) curvature, and the torsion with respect to (J, θ) , respectively. Moreover, for a contact form change $\theta = e^{2\lambda} \theta_0$, we have the following transformation laws:

$$(1.2) \quad P = e^{-4\lambda} P_0$$

1991 *Mathematics Subject Classification.* Primary 32V20; Secondary 53C44.

Key words and phrases. CR Manifold, Tanaka-Webster Curvature, Torsion, Moser inequality, Q -curvature flow, Paneitz operator, Sub-Laplacian, Kohn Laplacian.

*Research supported in part by the NSC of Taiwan.

and

$$(1.3) \quad Q = e^{-4\lambda}(Q_0 + 2P_0\lambda)$$

where P_0 and Q_0 denote the (CR) Paneitz operator and the Q -curvature with respect to (J, θ_0) , respectively. Note that

$$(1.4) \quad \int_M Q d\mu = 0.$$

Here the volume form $d\mu = \theta \wedge d\theta$.

Then we ask if we can always choose θ so that Q vanishes pointwise. The related problems for the Q -curvature on a Riemannian manifold are also addressed and studied in [B], [C1], and [CW] by using the method of the Q -curvature flow. In this paper we study a similar evolution equation and apply it to the problem addressed above on a CR 3-manifold.

We consider the functional \mathcal{E} on a closed CR 3-manifold (M, J) with a given contact class $[\theta_0]$ (which consists of all contact forms annihilating the underlying contact bundle):

$$(1.5) \quad \mathcal{E}(\theta) = \int_M P_0\lambda \cdot \lambda d\mu_0 + \int_M Q_0\lambda d\mu_0,$$

for $\theta = e^{2\lambda}\theta_0$, where $d\mu_0 = \theta_0 \wedge d\theta_0$. Then, for minimizing $\mathcal{E}(\theta)$ in $[\theta_0]$, it is natural to consider the following fourth order (Q -curvature) flow on a closed CR 3-manifold (M, J) :

$$(1.6) \quad \begin{cases} \frac{\partial \lambda}{\partial t} = -(Q_0 + 2P_0\lambda) + r = -e^{4\lambda}Q + r, \\ \theta = e^{2\lambda}\theta_0; \lambda(p, 0) = \lambda_0(p), \\ \int_M e^{4\lambda_0} d\mu_0 = \int_M d\mu_0, \end{cases}$$

where

$$r = \frac{\int_M (Q_0 + 2P_0\lambda) d\mu}{\int_M d\mu} = \frac{\int_M e^{4\lambda} Q d\mu}{\int_M d\mu}$$

and λ_0 is an initial real C^∞ smooth function. Note that $d\mu = e^{4\lambda}d\mu_0$ and the volume $V = \int_M d\mu$ is kept invariant under the flow (1.6). Indeed (1.6) is the (volume normalized) negative gradient flow of $\mathcal{E}(\theta)$. That is, by using (1.4) and (1.3), one can

check that

$$\begin{aligned}
(1.7) \quad & \frac{d}{dt} \mathcal{E}(\theta) \\
&= 2 \int_M P_0 \lambda \cdot \frac{\partial \lambda}{\partial t} d\mu_0 + \int_M Q_0 \frac{\partial \lambda}{\partial t} d\mu_0 \\
&= -2 \int_M (Q_0 + 2P_0 \lambda) P_0 \lambda d\mu_0 - \int_M Q_0 (Q_0 + 2P_0 \lambda) d\mu_0 \\
&= - \int_M (Q_0 + 2P_0 \lambda)^2 d\mu_0 \\
&= - \int_M e^{4\lambda} Q^2 d\mu.
\end{aligned}$$

under the flow (1.6). We will often denote a CR 3-manifold by $(M, J, [\theta_0])$ with its contact class (or equivalently its underlying contact bundle) indicated. Let $\overset{0}{A}_{11}$ denote the torsion with respect to (J, θ_0) . For the long time solution to the flow (1.6), we have the following result.

Theorem 1.1. *Let $(M, J, [\theta_0])$ be a closed CR 3-manifold with $\overset{0}{A}_{11} = 0$. Then the solution of (1.6) exists on $M \times [0, \infty)$.*

There are many torsion free examples. On the other hand, this condition implies strong topological obstruction (see the Appendix in [CH]). We hope this condition can be weakened in the future study.

To motivate the definitions of some analytic conditions, let us first examine the standard CR 3-sphere $\mathbf{S}^3 = \{(z_0, z_1) \mid \sum_{j=0}^1 z_j \bar{z}_j = 1\} \subset \mathbf{C}^2$ with the induced CR structure J from \mathbf{C}^2 and the contact form $\theta_0 = \frac{i(\bar{\partial}u - \partial u)}{2}|_{\mathbf{S}^3}$, where $u = \left(\sum_{j=0}^1 z_j \bar{z}_j\right) - 1$ is a defining function. With respect to (J, θ_0) , \mathbf{S}^3 is torsion free and $Q_0 = 0$. On the other hand, if $f_{p,q}$ is a bigraded spherical harmonic of type (p, q) on \mathbf{C}^2 (i.e. a harmonic polynomial which is a linear combination of terms of the form $z^\rho \bar{z}^\gamma$), then we have ([Chi]) (Δ_0 denotes the sub-Laplacian with respect to (J, θ_0))

$$(1.8) \quad \Delta_0 f_{p,q} = -(2pq + p + q) \cdot f_{p,q}$$

and

$$(1.9) \quad P_0 f_{p,q} = 4pq(p+1)(q+1) \cdot f_{p,q}.$$

Definition 1.1. *On a closed pseudohermitian manifold (M, J, θ) , we call the Paneitz operator P with respect to (J, θ) essentially positive if there exists a constant $\Upsilon > 0$*

such that

$$(1.10) \quad \int_M P\varphi \cdot \varphi d\mu \geq \Upsilon \int_M \varphi^2 d\mu.$$

for all real C^∞ smooth functions $\varphi \perp \text{Ker}(P)$ (i.e. perpendicular to the kernel of P in the L^2 norm with respect to the volume form $d\mu = \theta \wedge d\theta$).

Remark 1.1. We note that P being essentially positive is a CR invariant property by (1.2), i.e., it is independent of the choice of contact form. For M being the boundary of a bounded strictly pseudoconvex domain in C^2 , P appears in the transformation law of the first invariant in the logarithmic term of Fefferman's asymptotic expansion of the Szegő kernel ([H]). Also P appears as the compatibility operator for the degenerate Laplacian in the paper [GL]. On the other hand, the kernel of P is infinite dimensional, containing all CR-pluriharmonic functions (see Section 5). So even for the short time solution to (1.6), we need to treat the kernel part separately. The condition that P is essentially positive was also used to study the problem of the first eigenvalue of the sub-Laplacian (see [Chi]).

Since the restrictions of bigraded spherical harmonics to \mathbf{S}^3 span a dense subspace of $L^2(\mathbf{S}^3)$ (see Proposition 12.3.3 in [CS]), we conclude that the Paneitz operator P_0 of $(\mathbf{S}^3, J, \theta_0)$ is essentially positive by (1.9). Combining (1.8) and (1.9), we get

$$(2P_0 - \Delta_0^2)f_{p,q} = \lambda(p, q)f_{p,q},$$

where $\lambda(p, q) = p^2(4q - 1) + q^2(4p - 1) + 6pq + 4p^2q^2$. It is clear that if $P_0 f_{p,q} \neq 0$, i.e., $pq \neq 0$, then $\lambda(p, q) > 0$. This means that the operator $2P_0 - \Delta_0^2$ is positive on the orthogonal complement of the Kernel of P_0 . Therefore if φ is a real C^∞ smooth function such that $\varphi \perp \text{Ker}(P_0)$, then

$$(1.11) \quad 2 \int_{\mathbf{S}^3} P_0 \varphi \cdot \varphi d\mu_0 \geq \int_{\mathbf{S}^3} \Delta_0^2 \varphi \cdot \varphi d\mu_0 = \int_{\mathbf{S}^3} (\Delta_0 \varphi)^2 d\mu_0.$$

Let $\overset{0}{\nabla}$ and $\overset{0}{\nabla}^2$ denote the sub-gradient and the sub-Hessian with respect to (J, θ_0) , respectively. Now based on the Bochner formula, we have (Lemma 3.2 in Section 3;

see [Chi] also)

$$(1.12) \quad \begin{aligned} \int_M |\overset{0}{\nabla}^2 \varphi|^2 d\mu_0 &= 3 \int_M (\Delta_0 \varphi)^2 d\mu_0 - 2 \int_M P_0 \varphi \cdot \varphi d\mu_0 \\ &\quad - \int_M \overset{0}{W} |\overset{0}{\nabla} \varphi|^2 d\mu_0 - 6 \operatorname{Im} \int_M \overset{0}{A}_{\overline{11}} \varphi_1 \varphi_1 d\mu_0 \end{aligned}$$

where $\overset{0}{W}$ and $\overset{0}{A}_{\overline{11}}$ denote the Tanaka-Webster curvature and the torsion with respect to (J, θ_0) , respectively.

Observe that $\overset{0}{W}$ is a positive constant and $\overset{0}{A}_{\overline{11}} = 0$ for $(\mathbf{S}^3, J, \theta_0)$. It follows from (1.11) and (1.12) that

$$(1.13) \quad \int_{\mathbf{S}^3} |\overset{0}{\nabla}^2 \varphi|^2 d\mu_0 \leq 2 \int_{\mathbf{S}^3} (\Delta_0 \varphi)^2 d\mu_0$$

for $\varphi \perp \operatorname{Ker}(P_0)$.

Inspired by the inequality (1.13), we make the following definition.

Definition 1.2. *We say that the condition $(*)$ is satisfied on a closed CR 3-manifold $(M, J, [\theta_0])$ if there exist constants $0 \leq \varepsilon_0 < 1$ and $C(\varepsilon_0) \geq 0$ such that*

$$(*) \quad \int_M |\overset{0}{\nabla}^2 \varphi|^2 d\mu_0 \leq (2 + \varepsilon_0) \int_M (\Delta_0 \varphi)^2 d\mu_0 + C(\varepsilon_0) \int_M \varphi^2 d\mu_0$$

for all real C^∞ smooth functions $\varphi \perp \operatorname{Ker}(P_0)$.

Theorem 1.2. *Let $(M, J, [\theta_0])$ be a closed CR 3-manifold. Suppose that P_0 is essentially positive and the condition $(*)$ holds. Then the solution λ of (1.6) satisfies an a priori $S^{2,2}$ estimate:*

$$||\lambda||_{S^{2,2}} \leq C(T)$$

for $t \in [0, T)$, where $C(T)$ is a constant depending on T . Moreover, if in addition $(Q_0)_{\ker} = 0$, then the bound $C(T)$ can be replaced by a constant C independent of the time.

In the paper [B], the (essential) positivity of the Riemannian Paneitz operator is also needed for a similar result on the Q -curvature flow on a closed conformal 4-manifold $(M, [g_0])$ with a given conformal class $[g_0]$. However, in the Riemannian case, the condition analogous to the condition $(*)$ holds always. We wonder to what extent the condition $(*)$ is valid for a closed pseudohermitian manifold.

In Section 5, we show that if the torsion $\overset{0}{A}_{11}$ of (M, J, θ_0) is zero, then the *CR* Paneitz operator P_0 is essentially positive. In that case, the *CR* Paneitz operator $P_0 = \square_b \overline{\square}_b$ and the Kohn Laplacian \square_b and $\overline{\square}_b$ commute (see Section 5). We can write an L^2 function $\varphi = \varphi_{\ker} + \varphi^\perp$ where $\varphi_{\ker} \in \text{Ker}(P_0)$ and $\varphi^\perp \in \text{Ker}(P_0)^\perp$. We have the following a priori estimates of higher orders.

Theorem 1.3. *Let $(M, J, [\theta_0])$ be a closed CR 3-manifold. Suppose that P_0 is essentially positive and the condition $(*)$ holds. In addition, suppose also that $\Delta_0(\text{Ker}(P_0)) \subset \text{Ker}(P_0)$. Then for any nonnegative integer k , the solution λ of (1.6) satisfies an a priori $S^{2k,2}$ estimate:*

$$||\lambda||_{S^{2k,2}} \leq C(k, T)$$

for $t \in [0, T]$, where $C(k, T)$ is a constant depending on k and T . Moreover, if in addition $(Q_0)_{\ker} = 0$, then the bound $C(k, T)$ can be replaced by a constant $C(k)$ independent of the time.

We remark that in the torsion free case, the condition $\Delta_0(\text{Ker}(P_0)) \subset \text{Ker}(P_0)$ holds true (also P_0 is essentially positive as mentioned above). We have the following asymptotic convergence of solutions of (1.6).

Theorem 1.4. *Let $(M, J, [\theta_0])$ be a closed CR 3-manifold with $(Q_0)_{\ker} = 0$. Suppose that $\overset{0}{A}_{11} = 0$. Then the solution of (1.6) exists on $M \times [0, \infty)$ and converges smoothly to $\lambda_\infty \equiv \lambda(\cdot, \infty)$ as $t \rightarrow \infty$. Moreover, the contact form $e^{2\lambda_\infty} \theta_0$ has zero Q -curvature.*

Remark 1.2. 1. Let W_0 denote the Tanaka-Webster (scalar) curvature with respect to (J, θ_0) . If $\overset{0}{A}_{11} = 0$, then P_0 commutes with Δ_0 and hence there holds

$$(Q_0)_{\ker} = -\frac{2}{3}(\Delta_0 W_0)_{\ker} = -\frac{2}{3}\Delta_0(W_0)_{\ker}$$

by (1.1). It follows that

$$(Q_0)_{\ker} = 0 \iff (W_0)_{\ker} \text{ is a constant.}$$

2. On the standard CR 3-sphere $(\mathbf{S}^3, J, [\theta_0])$, we have

(i) $\overset{0}{A}_{11} = 0$ and $Q_0 = 0$,

(ii) the condition $(*)$ holds with $\varepsilon_0 = 0$ and $C(\varepsilon_0) = 0$.

As a consequence of Theorem 1.4, we have

Corollary 1.5. *Let $(\mathbf{S}^3, J, [\theta_0])$ be the standard CR 3-sphere. Then the solution of (1.6) exists on $\mathbf{S}^3 \times [0, \infty)$ and converges smoothly to λ_∞ such that $e^{2\lambda_\infty} \theta_0$ is a contact form of zero Q -curvature.*

We recall that θ_0 is called an invariant contact form on a CR 3-manifold M if it is locally volume-normalized with respect to a closed $(2, 0)$ -form on M ([FH], [L1], [Fa]). In the paper [FH], the authors proved that the Q -curvature of an invariant contact form vanishes. Indeed if M is a real hypersurface in \mathbf{C}^2 , then M admits an invariant contact form θ_0 so that $Q_0 = 0$ on M . In general, there is a topological obstruction for the global existence of an invariant contact form θ_0 ([L1]). However, on the standard CR 3-sphere, $Q_0 = 0$ if and only if θ_0 is an invariant contact form.

About other curvature flows, we notice that it is still an open problem whether we have the long-time existence and convergence for solutions of the CR Yamabe flow on a closed CR 3-manifold ([CC1], [JL]). This is a second order subparabolic equation. On the other hand, the flow (1.6) which we are dealing with is a fourth-order subparabolic (at least under a certain condition) equation. Because there seems to be no suitable maximum principle available for fourth-order subelliptic operators, we need to invoke a priori L^2 estimates for solutions to (1.6) in place of the pointwise estimates used for second-order (sub)elliptic operators ([B], [C], [C1], [C2], [CW]). In case M is a surface, the Q -curvature flow corresponds to the 2-dimensional Calabi flow which is solved completely by P. T. Chruściel ([C]) and the first author ([C3], [C4]; see also [CW]).

In order to get the $S^{2,2}$ -estimate (see Appendix A for the definition of Folland-Stein norms $S^{k,p}$), we need an additional analytic condition $(*)$ (which holds for the standard pseudohermitian 3-sphere) plus a trick from S. Brendle's work ([B]) for the Q -curvature flow on Riemannian 4-manifolds. This is because the pseudohermitian version of a Bochner-type estimate is entirely different from the Riemannian version (see Lemma 3.1).

We briefly describe the methods used in our proofs. In Section 2, in order to have the $S^{k,2}$ -estimates for λ , we need to derive the analogue of the Moser inequality on pseudohermitian 3-manifolds. In Section 3, based on a pseudohermitian version

of the Bochner formula, we first derive a key estimate for the equation (1.6) as in Lemma 3.2, which involves the CR Paneitz operator P_0 . We show the subellipticity of P_0 on $(\ker P_0)^\perp$ under the torsion free condition to get the short time solution. For the long-time solution of (1.6) under the same condition (i.e., Theorem 1.1), we obtain the higher order bounds for the solution.

In Section 4, we derive the $S^{2,2}$ -estimate and higher-order $S^{2k,2}$ -estimates for λ under the flow (1.6), and hence prove Theorem 1.2 and Theorem 1.3. Then we prove the smooth convergence of solutions of (1.6) (i.e. Theorem 1.4) by a method analogous to that in [S]. In Section 5, we show the essential positivity of the CR Paneitz operator P on (M, J, θ) with the zero torsion.

2. Moser's Inequality on Pseudohermitian 3-manifolds

In this section, based on [A], [CL], [FS1], and [SC], we derive the analogue of Moser's inequality on pseudohermitian 3-manifolds.

Let $H^1 = \mathbf{C} \times \mathbf{R}$ be the (3-dimensional) Heisenberg group with coordinates (z, t) . For each real number $r \in \mathbf{R}$, there is a dilation naturally associated with H^1 , which is usually denoted as

$$\delta_r u = \delta_r(z, t) = (rz, r^2 t).$$

The anisotropic dilation structure on H^1 introduces a homogeneous norm

$$|u| = |(z, t)| = (|z|^4 + t^2)^{1/4}.$$

With this norm, we can define the Heisenberg ball centered at $u = (z, t)$ with radius R by $B(u, R) = \{v \in H^1 : |u^{-1} \cdot v| < R\}$.

In their paper ([CL]), William S. Cohn and Guozhen Lu show that for all $\varphi \in C_0^\infty(H^1)$,

$$(2.1) \quad |\varphi(v)| \leq L^{-1} \int_{H^1} \frac{|\nabla_b \varphi(u)|}{|v^{-1} \cdot u|^3} dV(u),$$

where $L = \frac{2\pi \Gamma(1/2)\Gamma(3/4)}{\Gamma(1)\Gamma(5/4)}$ and $dV(u) = dx \wedge dy \wedge dt$, $z = x + iy$. Note that we only take the case $\beta = 1$ in Theorem 1.2 of ([CL]).

Let B be the unit Heisenberg ball of H^1 , centered at $(0, 0)$. Let $\|\cdot\|_p$ denote the L^p norm with respect to the volume form $dV(u)$.

Lemma 2.1. *For $\varphi \in C_0^\infty(B)$ and $p \geq 4$, we have*

$$(2.2) \quad \|\varphi\|_p \leq L^{-1} \|\nabla_b \varphi\|_4 \sup_{v \in B} \left[\int_B |v^{-1} \cdot u|^{-3k} dV(u) \right]^{\frac{1}{k}}$$

where $\frac{1}{k} = \frac{1}{p} + \frac{3}{4}$.

Proof. We write

$$\begin{aligned} & |v^{-1} \cdot u|^{-3} |\nabla_b \varphi(u)| \\ &= \left(|v^{-1} \cdot u|^{-3k} |\nabla_b \varphi(u)|^4 \right)^{\frac{1}{p}} \left(|v^{-1} \cdot u|^{-3k} \right)^{\frac{3}{4}} \left(|\nabla_b \varphi(u)|^4 \right)^{\frac{1}{4} - \frac{1}{p}}. \end{aligned}$$

Since $\frac{1}{p} + \frac{1}{4} + \left(\frac{1}{4} - \frac{1}{p}\right) = 1$, applying Hölder's inequality to (2.1), we have

$$\begin{aligned} |\varphi(v)| &\leq L^{-1} \left(\int_B |v^{-1} \cdot u|^{-3k} |\nabla_b \varphi(u)|^4 dV(u) \right)^{\frac{1}{p}} \left(\int_B |v^{-1} \cdot u|^{-3k} dV(u) \right)^{\frac{3}{4}} \\ &\quad \left(\int_B |\nabla_b \varphi(u)|^4 dV(u) \right)^{\frac{1}{4} - \frac{1}{p}}. \end{aligned}$$

This implies that

$$\begin{aligned} \|\varphi\|_p &= \left(\int_B |\varphi(v)|^p dV(v) \right)^{\frac{1}{p}} \\ &\leq \left(L^{-1} \|\nabla_b \varphi\|_4^{1 - \frac{4}{p}} \cdot \Lambda^{\frac{3k}{4}} \right) \left[\int_B \int_B |v^{-1} \cdot u|^{-3k} |\nabla_b \varphi(u)|^4 dV(u) dV(v) \right]^{\frac{1}{p}} \\ &\leq L^{-1} \|\nabla_b \varphi\|_4 \cdot \Lambda, \end{aligned}$$

where $\Lambda = \sup_{v \in B} \left[\int_B |v^{-1} \cdot u|^{-3k} dV(u) \right]^{\frac{1}{k}}$.

■

Corollary 2.2. *For all $\varphi \in C_0^\infty(B)$,*

$$(2.3) \quad \int_B e^\varphi dV(u) \leq C \exp(\varkappa \|\nabla_b \varphi\|_4^4),$$

where C and \varkappa are two positive constants.

Proof. Let $\Sigma = \partial B$ be the unit Heisenberg sphere and dA be the unique Radon measure on Σ ([CL], [FS1]). We denote $A_1 = \int_\Sigma dA$. Note that there exists a

number δ such that $|v^{-1} \cdot u| < \delta$ for all $u, v \in B$. This means that $B \subset B(v, \delta)$ for all $v \in B$.

By (2.2), we have

$$\begin{aligned}
 \|\varphi\|_p &\leq L^{-1} \|\nabla_b \varphi\|_4 \sup_{v \in B} \left[\int_B |v^{-1} \cdot u|^{-3k} dV(u) \right]^{\frac{1}{k}} \\
 &\leq L^{-1} \|\nabla_b \varphi\|_4 \sup_{v \in B} \left[\int_{B(v, \delta)} |v^{-1} \cdot u|^{-3k} dV(v^{-1} \cdot u) \right]^{\frac{1}{k}} \\
 (2.4) \quad &= L^{-1} \|\nabla_b \varphi\|_4 \left[\int_{B(0, \delta)} |u|^{-3k} dV(u) \right]^{\frac{1}{k}} \\
 &= L^{-1} A_1^{\frac{1}{k}} \|\nabla_b \varphi\|_4 \left(\int_0^\delta r^{3-3k} dr \right)^{\frac{1}{k}} \\
 &= L^{-1} A_1^{\frac{1}{k}} \delta^{\frac{4-3k}{k}} \|\nabla_b \varphi\|_4 \left(\frac{1}{4-3k} \right)^{\frac{1}{k}}.
 \end{aligned}$$

Since $\frac{1}{k} = \frac{1}{p} + \frac{3}{4}$ and $p \geq 4$, we immediately get that $\left(\frac{1}{4-3k}\right)^{\frac{1}{k}} \leq Cp^{\frac{3}{4}}$ and $-3 \leq \frac{4-3k}{k} \leq 1$ for some constant C . Thus, by (2.4), there exists a constant K such that for all $p \geq 1$,

$$(2.5) \quad \|\varphi\|_p \leq K \|\nabla_b \varphi\|_4 \cdot p^{\frac{3}{4}}.$$

It follows that

$$\begin{aligned}
 \int_B e^\varphi dV(u) &= \int_B \left(\sum_{p=0}^{\infty} \frac{\varphi^p}{p!} \right) dV(u) \\
 &\leq \sum_{p=0}^{\infty} \frac{\|\varphi\|_p^p}{p!} \\
 (2.6) \quad &\leq \sum_{p=0}^{\infty} K^p \|\nabla_b \varphi\|_4^p (p!)^{-1} p^{\frac{3p}{4}} \\
 &= \sum_{p=0}^{\infty} \frac{(K^4)^{\frac{p}{4}} (\|\nabla_b \varphi\|_4^4)^{\frac{p}{4}} (\frac{p}{4})!}{(\frac{p}{4})!} \frac{1}{p!} p^{\frac{3p}{4}}.
 \end{aligned}$$

Here $x! = \Gamma(x+1)$ for all real number $x \geq 0$.

According to Stirling's formula, when $p \rightarrow \infty$, we can estimate

$$\begin{aligned} \frac{\left(\frac{p}{4}\right)!}{p!} p^{\frac{3p}{4}} &\approx \frac{\left(2\pi\frac{p}{4}\right)^{\frac{1}{2}} \left(\frac{p}{4e}\right)^{\frac{p}{4}}}{(2\pi p)^{\frac{1}{2}} \left(\frac{p}{e}\right)^p} p^{\frac{3p}{4}} \\ &\approx \frac{1}{2} \left(\frac{1}{4}\right)^{\frac{p}{4}} e^{\frac{3p}{4}}. \end{aligned}$$

Therefore, from (2.6), we get

$$\begin{aligned} \int_B e^\varphi dV(u) &\leq C \sum_{p=0}^{\infty} \frac{\left(\frac{e^3}{4} K^4\right)^{\frac{p}{4}} (\|\nabla_b \varphi\|_4^4)^{\frac{p}{4}}}{\left(\frac{p}{4}\right)!} \\ &\leq C \cdot \exp\left(\varkappa \|\nabla_b \varphi\|_4^4\right), \end{aligned}$$

for some constants C and \varkappa .

■

Now we are ready to prove an analogue of the Moser inequality in pseudohermitian geometry. Let $C^\infty(M)$ denote the space of all real valued C^∞ smooth functions on M .

Theorem 2.3. (*Pseudohermitian Moser inequality*) *Let (M, J, θ) be a closed pseudohermitian 3-manifold. Then there exist constants C , \varkappa , and ν such that for all $\varphi \in C^\infty(M)$, there holds*

$$(2.7) \quad \int_M e^\varphi d\mu \leq C \exp\left(\varkappa \|\nabla_b \varphi\|_4^4 + \nu \|\varphi\|_4^4\right)$$

where $d\mu = \theta \wedge d\theta$ and the L^4 norm $\|\cdot\|_4$ respects the volume form $d\mu$.

Proof. For each point $x \in M$, there exists a neighborhood U_x of x such that U_x is diffeomorphic to a Heisenberg ball $B(r_x)$ centered at the origin with radius $r_x < 1$. Choose a cut-off function η_x such that $\eta_x(v) = 1$ for $|v| < \frac{r_x}{2}$ and $\eta_x(v) = 0$ for $|v| \geq r_x$.

Since M is closed, there exist finite balls $(B(\frac{r_j}{2}), \eta_j)$, $j = 1, \dots, m$ such that $\{B(\frac{r_j}{2})\}$ is an open covering of M . Let $\varphi_j = \eta_j \varphi$. Making use of (2.3), we compute

$$\begin{aligned} \int_M e^\varphi d\mu &\leq C \cdot \sum_{j=1}^m \int_{B(\frac{r_j}{2})} e^\varphi dV(u) \\ &\leq C \cdot \sum_{j=1}^m \int_B e^{\varphi_j} dV(u) \\ &\leq C \cdot \sum_{j=1}^m \exp\left(\varkappa \|\nabla_b \varphi_j\|_{4;B}^4\right) \\ &\leq C \cdot \sum_{j=1}^m \exp\left(\varkappa \|\nabla_b \varphi\|_4^4 + \nu \|\varphi\|_4^4\right), \end{aligned}$$

for some constants C , \varkappa , and ν .

■

3. The Long-time Existence

Let T be the maximal time for a solution of the flow (1.6) on $M \times [0, T)$. We will derive the $S^{k,2}$ -norm estimate for λ under the flow (1.6) for all $0 \leq t < T$. It then follows that we have the long-time existence for solutions of (1.6) on $M \times [0, \infty)$.

First we have an integral version of the Bochner formula on a pseudohermitian 3-manifold.

Lemma 3.1. *Let (M, J, θ_0) be a closed pseudohermitian 3-manifold. Then for any $\lambda \in C^\infty(M)$, there holds*

$$\begin{aligned} (3.1) \quad 0 &= \int_M (\Delta_0 \lambda)^2 d\mu_0 - \int_M |\overset{0}{\nabla}^2 \lambda|^2 d\mu_0 + 2 \int_M (\lambda_0)^2 d\mu_0 \\ &\quad - \int_M \overset{0}{W} |\overset{0}{\nabla} \lambda|^2 d\mu_0 + 2 \operatorname{Im} \int_M \overset{0}{A}_{\overline{11}} \lambda_1 \lambda_1 d\mu_0 \end{aligned}$$

where $\overset{0}{\nabla}$ and T_0 denote the sub-gradient and the characteristic vector field with respect to (J, θ_0) , respectively, and $\lambda_0 = T_0 \lambda$.

Proof. We first show that (see Appendix A for definitions of the notations)

$$\begin{aligned} (3.2) \quad \frac{1}{2} \Delta_0 |\overset{0}{\nabla} \lambda|^2 &= |\overset{0}{\nabla}^2 \lambda|^2 + \langle \overset{0}{\nabla} \lambda, \overset{0}{\nabla} (\Delta_0 \lambda) \rangle_{J, \theta_0} + \overset{0}{W} |\overset{0}{\nabla} \lambda|^2 \\ &\quad + \operatorname{Tor}(\overset{0}{\nabla} \lambda, \overset{0}{\nabla} \lambda) - 2i \lambda_1 \lambda_{0\overline{1}} + 2i \lambda_{\overline{1}} \lambda_{01}, \end{aligned}$$

where $Tor(\overset{0}{\nabla}\lambda, \overset{0}{\nabla}\lambda) \equiv i\overset{0}{A}_{\overline{11}}\lambda_1\lambda_1 + \text{conjugate of } i\overset{0}{A}_{\overline{11}}\lambda_1\lambda_1$.

We compute

$$|\overset{0}{\nabla}\lambda|^2 = 2|\lambda_1|^2 = 2\lambda_1\lambda_{\overline{1}}, \quad |\overset{0}{\nabla}^2\lambda|^2 = 2\lambda_{\overline{11}}\lambda_{11} + 2\lambda_{\overline{1}\overline{1}}\lambda_{1\overline{1}}$$

and

$$\begin{aligned} \frac{1}{2}\overset{0}{\Delta}_0|\overset{0}{\nabla}\lambda|^2 &= (\lambda_1\lambda_{\overline{1}})_{1\overline{1}} + (\lambda_1\lambda_{\overline{1}})_{\overline{1}1} \\ (3.3) \quad &= \lambda_{\overline{1}1\overline{1}}\lambda_1 + 2\lambda_{\overline{1}\overline{1}}\lambda_{1\overline{1}} + 2\lambda_{\overline{11}}\lambda_{11} + \lambda_{\overline{1}}\lambda_{11\overline{1}} + \lambda_1\lambda_{\overline{1}\overline{1}1} + \lambda_{\overline{1}}\lambda_{1\overline{1}1} \\ &= |\overset{0}{\nabla}^2\lambda|^2 + \lambda_1(\lambda_{\overline{1}\overline{1}1} + \lambda_{\overline{1}1\overline{1}}) + \text{conjugate of } \lambda_1(\lambda_{\overline{1}\overline{1}1} + \lambda_{\overline{1}1\overline{1}}). \end{aligned}$$

Observe that

$$\begin{aligned} (3.4) \quad \langle \overset{0}{\nabla}\lambda, \overset{0}{\nabla}(\overset{0}{\Delta}_0\lambda) \rangle_{J, \theta_0} &= \lambda_1(\lambda_{1\overline{1}} + \lambda_{\overline{1}1})_{\overline{1}} + \text{conjugate of } \lambda_1(\lambda_{1\overline{1}} + \lambda_{\overline{1}1})_{\overline{1}} \\ &= \lambda_1(\lambda_{1\overline{1}\overline{1}} + \lambda_{\overline{1}\overline{1}1}) + \text{conjugate of } \lambda_1(\lambda_{1\overline{1}\overline{1}} + \lambda_{\overline{1}\overline{1}1}). \end{aligned}$$

Taking the covariant differentiation of

$$(3.5) \quad \lambda_{\overline{1}1} = \lambda_{1\overline{1}} - i\lambda_0$$

([L1]) in the $Z_{\overline{1}}$ direction gives

$$(3.6) \quad \lambda_{\overline{1}1\overline{1}} = \lambda_{1\overline{1}\overline{1}} - i\lambda_{0\overline{1}}.$$

A commutation relation ([L1]) for covariant derivatives of a 1-form gives

$$(3.7) \quad \lambda_{\overline{1}\overline{1}1} = \lambda_{\overline{1}1\overline{1}} - i\lambda_{\overline{1}0} + \overset{0}{W}\lambda_{\overline{1}}.$$

Now from (3.3), (3.4), (3.6), and (3.7), we obtain

$$\begin{aligned} (3.8) \quad \frac{1}{2}\overset{0}{\Delta}_0|\overset{0}{\nabla}\lambda|^2 &= |\overset{0}{\nabla}^2\lambda|^2 + \langle \overset{0}{\nabla}\lambda, \overset{0}{\nabla}(\overset{0}{\Delta}_0\lambda) \rangle + (\overset{0}{W}\lambda_1\lambda_{\overline{1}} + \text{conjugate}) \\ &\quad + [-i\lambda_1(\lambda_{\overline{1}0} + \lambda_{0\overline{1}}) + \text{conjugate}]. \end{aligned}$$

Then (3.2) follows from (3.8) and the following commutation relation ([L1])

$$\lambda_{0\overline{1}} = \lambda_{\overline{1}0} + \overset{0}{A}_{\overline{11}}\lambda_1.$$

Finally integrating both sides of (3.2) and applying (3.5), we obtain

$$\begin{aligned} 0 &= \int_M (\overset{0}{\Delta}_0\lambda)^2 d\mu_0 - \int_M |\overset{0}{\nabla}^2\lambda|^2 d\mu_0 + 2 \int_M (i\lambda_1\lambda_{0\overline{1}} - i\lambda_{\overline{1}}\lambda_{01}) d\mu_0 \\ &\quad - \int_M \overset{0}{W}|\overset{0}{\nabla}\lambda|^2 d\mu_0 + 2 \operatorname{Im} \int_M \overset{0}{A}_{\overline{11}}\lambda_1\lambda_1 d\mu_0. \end{aligned}$$

By integrating by parts and (3.5), we compute

$$\begin{aligned} \int_M (i\lambda_1\lambda_{0\bar{1}} - i\lambda_{\bar{1}}\lambda_{01})d\mu_0 &= \int_M i\lambda_0(-\lambda_{1\bar{1}} + \lambda_{\bar{1}1})d\mu_0 \\ &= \int_M (\lambda_0)^2 d\mu_0. \end{aligned}$$

Combining the above two formulas gives (3.1).

■

Lemma 3.2. *Let (M, J, θ_0) be a closed pseudohermitian 3-manifold. Then for any $\lambda \in C^\infty(M)$, there holds*

$$\begin{aligned} 2 \int_M P_0 \lambda \cdot \lambda d\mu_0 &= 3 \int_M (\Delta_0 \lambda)^2 d\mu_0 - \int_M |\overset{0}{\nabla}^2 \lambda|^2 d\mu_0 \\ &\quad - \int_M \overset{0}{W} |\overset{0}{\nabla} \lambda|^2 d\mu_0 - 6 \operatorname{Im} \int_M \overset{0}{A}_{\bar{1}1} \lambda_1 \lambda_1 d\mu_0. \end{aligned}$$

Proof. Multiplying both sides of the formula $P_0 \lambda = \Delta_0^2 \lambda + T_0^2 \lambda + 4 \operatorname{Im}(\overset{0}{A}_{\bar{1}1} \lambda_{11} + \overset{0}{A}_{\bar{1}1,1} \lambda_1)$ by λ and integrating, we compute

$$\begin{aligned} 2 \int_M P_0 \lambda \cdot \lambda d\mu_0 &= 2 \int_M (\Delta_0 \lambda)^2 d\mu_0 - 2 \int_M (\lambda_0)^2 d\mu_0 - 8 \operatorname{Im} \int_M \overset{0}{A}_{\bar{1}1} \lambda_1 \lambda_1 d\mu_0 \\ &= 3 \int_M (\Delta_0 \lambda)^2 d\mu_0 - \int_M |\overset{0}{\nabla}^2 \lambda|^2 d\mu_0 - \int_M \overset{0}{W} |\overset{0}{\nabla} \lambda|^2 d\mu_0 \\ &\quad - 6 \operatorname{Im} \int_M \overset{0}{A}_{\bar{1}1} \lambda_1 \lambda_1 d\mu_0 \end{aligned}$$

by integrating by parts and Lemma 3.1.

■

As a consequence of Lemma 3.2 and (1.11), we have

Corollary 3.3. *Let $(\mathbf{S}^3, J, \theta_0)$ be the standard pseudohermitian 3-sphere. Then*

$$\int_{\mathbf{S}^3} |\overset{0}{\nabla}^2 \lambda|^2 d\mu_0 \leq 2 \int_{\mathbf{S}^3} (\Delta_0 \lambda)^2 d\mu_0$$

for $\lambda \perp \ker(P_0)$.

It follows from (1.7) that

Lemma 3.4. *Let $(M, J, [\theta_0])$ be a closed CR 3-manifold. Let λ be a solution of the flow (1.6) on $M \times [0, T)$. Then there exists a positive constant $\beta = \beta(Q_0, \theta_0)$ such*

that

$$(3.9) \quad \mathcal{E}(\theta) = \int_M P_0 \lambda \cdot \lambda d\mu_0 + \int_M Q_0 \lambda d\mu_0 \leq \beta^2,$$

for all $t \in [0, T)$.

Lemma 3.5. *Let $f : [0, T) \rightarrow \mathbb{R}$ be a C^1 smooth function satisfying $f' \leq -C_1 f + C_2$ for some positive constants $C_1, C_2 > 0$. Then $f(t) \leq f(0)e^{-C_1 t} + \frac{C_2}{C_1}$ for $t \in [0, T)$.*

We will often use the above lemma (whose proof is left to the reader) to obtain the higher order estimates. Now we write $\lambda = \lambda_{\ker} + \lambda^\perp$. $Q_0 = (Q_0)_{\ker} + Q_0^\perp$ with respect to P_0 . Comparing both sides of the following formula:

$$\frac{\partial \lambda_{\ker}}{\partial t} + \frac{\partial \lambda^\perp}{\partial t} = \frac{\partial \lambda}{\partial t} = -(Q_0 + 2P_0 \lambda) + r(t).$$

we obtain

$$(3.10) \quad \frac{\partial \lambda^\perp}{\partial t} = -(Q_0^\perp + 2P_0 \lambda^\perp)$$

and

$$(3.11) \quad \frac{\partial \lambda_{\ker}}{\partial t} = -(Q_0)_{\ker} + r(t).$$

From now on, C or C_j ($C(k, T)$, $C(T)$, etc., respectively) denotes a generic constant (with emphasis on depending on (k, T) , T , etc., respectively) which may vary from line to line.

Proposition 3.6. *Let $(M, J, [\theta_0])$ be a closed CR 3-manifold with $A_{11}^0 = 0$. Then under the flow (1.6) (or equivalently (3.10) and (3.11)), there exists a positive constant $C(k, \Upsilon, \theta_0, \beta, T) > 0$ such that*

$$\|\lambda\|_{S^{2k,2}} \leq C(k, \Upsilon, \theta_0, \beta, T)$$

for all $0 \leq t < T$.

Proof. From (3.11), we have

$$(3.12) \quad \lambda_{\ker}(t, p) - \lambda_{\ker}(0, p) = -(Q_0)_{\ker}(p)t + \int_0^t r dt.$$

Since $(P_0$ being self-adjoint)

$$\int_M P_0 \lambda^\perp \cdot \lambda_{\ker} d\mu_0 = 0$$

and

$$\begin{aligned} & \int_M Q_0 \lambda_{\ker} d\mu_0 \\ &= \int_M Q_0 [\lambda_{\ker}(0, p) - (Q_0)_{\ker}(p)t + \int_0^t r dt] d\mu_0 \\ &= \int_M Q_0 \lambda_{\ker}(0, p) d\mu_0 - t \int_M [(Q_0)_{\ker}(p)]^2 d\mu_0 \\ &\geq -C - Ct, \end{aligned}$$

we compute

$$\begin{aligned} & \int_M P_0 \lambda \cdot \lambda d\mu_0 + \int_M Q_0 \lambda d\mu_0 \\ &= \int_M P_0 \lambda^\perp \cdot (\lambda_{\ker} + \lambda^\perp) d\mu_0 + \int_M Q_0 (\lambda_{\ker} + \lambda^\perp) d\mu_0 \\ &\geq \int_M P_0 \lambda^\perp \cdot \lambda^\perp d\mu_0 + \int_M Q_0 \lambda^\perp d\mu_0 - C - Ct. \end{aligned}$$

It then follows from Lemma 3.4 that

$$\int_M P_0 \lambda^\perp \cdot \lambda^\perp d\mu_0 + \int_M Q_0 \lambda^\perp d\mu_0 \leq (\beta^2 + C) + Ct$$

for all $t \in [0, T)$. Since the torsion of (M, J, θ_0) vanishes, the CR Paneitz operator P_0 is essentially positive in the sense that

$$\int_M P_0 \lambda^\perp \cdot \lambda^\perp d\mu_0 \geq \Upsilon \int_M (\lambda^\perp)^2 d\mu_0$$

by Theorem 5.3. All these imply that there exists a positive constant $C(\Upsilon, Q_0, \theta_0) > 0$ such that

$$\begin{aligned} (\beta^2 + C) + Ct &\geq \int_M P_0 \lambda^\perp \cdot \lambda^\perp d\mu_0 + \int_M Q_0 \lambda^\perp d\mu_0 \\ &\geq \frac{\Upsilon}{2} \int_M (\lambda^\perp)^2 d\mu_0 - C(\Upsilon, Q_0, \theta_0) \end{aligned}$$

for all $t \in [0, T)$. So there exists a positive constant $C(\Upsilon, Q_0, \theta_0, \beta, T)$ such that

$$(3.13) \quad \int_M (\lambda^\perp)^2 d\mu_0 \leq C(\Upsilon, Q_0, \theta_0, \beta, T),$$

for all $t \in [0, T)$.

On the other hand, we observe from the condition $\overset{0}{A}_{11} = 0$ that

$$(3.14) \quad \Delta_0^k T_0^2 = T_0^2 \Delta_0^k$$

and hence

$$(3.15) \quad \Delta_0^k P_0 = P_0 \Delta_0^k.$$

It follows that

$$(3.16) \quad \Delta_0^k \lambda^\perp \perp \ker P_0, \quad \Delta_0^k \lambda_{\ker} \in \ker P_0$$

Next we compute, for all positive integers k ,

$$(3.17) \quad \begin{aligned} & \frac{d}{dt} \int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 \\ &= 2 \int_M (\Delta_0^k \lambda^\perp) (\Delta_0^k \frac{\partial \lambda^\perp}{\partial t}) d\mu_0 \\ &= -2 \int_M (\Delta_0^k \lambda^\perp) [\Delta_0^k (Q_0^\perp + 2P_0 \lambda^\perp)] d\mu_0 \\ &= -2 \int_M \lambda^\perp [\Delta_0^{2k} (Q_0^\perp)] d\mu_0 - 4 \int_M (\Delta_0^k \lambda^\perp) (P_0 \Delta_0^k \lambda^\perp) d\mu_0. \end{aligned}$$

Here we have used (3.15). By (3.16) and essential positivity of P_0 , we obtain

$$(3.18) \quad \int_M (\Delta_0^k \lambda^\perp) (P_0 \Delta_0^k \lambda^\perp) d\mu_0 \geq \Upsilon \int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0.$$

Therefore from (3.17), (3.18), and (3.13), we conclude that

$$(3.19) \quad \frac{d}{dt} \int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 \leq -4\Upsilon \int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 + C(k, T).$$

By applying Lemma 3.5 to the O.D.E. $f'(t) \leq -4\Upsilon f(t) + C(k, T)$ from (3.19), we obtain

$$\int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 \leq C(k, T).$$

and hence

$$\|\lambda^\perp - \overline{\lambda^\perp}\|_{S^{2k,2}} \leq C(k, T)$$

for all $0 \leq t < T$.

From the definition of the Q -curvature, $\overset{0}{A}_{11} = 0$, and (3.16), we compute

$$\int_M (Q_0)_{\ker} d\mu_0 = -\frac{2}{3} \int_M \Delta_0 (R_0)_{\ker} d\mu_0 = 0.$$

This and (3.12) imply

$$\lambda - \overline{\lambda} = (\lambda^\perp - \overline{\lambda^\perp}) + [\lambda_{\ker}(0, p) - (Q_0)_{\ker}(p)t] - \overline{\lambda_{\ker}(0, p)},$$

and hence

$$(3.20) \quad \|\lambda - \bar{\lambda}\|_{S^{2k,2}} \leq C((Q_0)_{\ker}, k, T) + \|\lambda^\perp - \bar{\lambda}^\perp\|_{S^{2k,2}} \leq C(k, T)$$

for all $0 \leq t < T$. Recall that the average \bar{f} of a function f is defined by $\bar{f} = \frac{\int_M f d\mu_0}{\int_M d\mu_0}$.

In particular, there holds

$$\|\lambda - \bar{\lambda}\|_{S^{2,2}} \leq C(T).$$

Therefore by the Sobolev embedding theorem, we have $S^{2,2} \subset S^{1,4}$ and

$$(3.21) \quad \|\lambda - \bar{\lambda}\|_{S^{1,4}} \leq C(T).$$

Now using Theorem 2.3 (pseudohermitian Moser inequality), we get

$$\int_M e^{4(\lambda - \bar{\lambda})} d\mu_0 \leq C \exp(C\|\lambda - \bar{\lambda}\|_{S^{1,4}}) \leq C(T).$$

Together with $\int_M e^{4\lambda} d\mu_0$ being invariant under the flow, we conclude that

$$(3.22) \quad C \geq \bar{\lambda} \geq -C(T)$$

(the upper bound is obtained by observing that $\int \lambda d\mu_0 \leq \int e^{4\lambda} d\mu_0$). By (3.20) and (3.22), we finally obtain

$$\|\lambda\|_{S^{2k,2}} \leq C(k, T)$$

for all $0 \leq t < T$.

■

Proof of Theorem 1.1:

When the torsion is zero, we can write $P_0 = \square_0 \overline{\square_0} = \overline{\square_0} \square_0$ (\square_0 and $\overline{\square_0}$ commute) where $\square_0 \equiv \Delta_0 + iT_0$ is the Kohn Laplacian (acting on functions) with respect to (J, θ_0) , and T_0 is an infinitesimal CR diffeomorphism. This implies that (M, J) is embeddable (see Theorem 2.1 in [Le]). Therefore \square_0 and $\overline{\square_0}$ have subelliptic estimates on the orthogonal complements of $\text{Ker}(\square_0)$ and $\text{Ker}(\overline{\square_0})$ in L^2 , denoted as $(\text{Ker}(\square_0))^\perp$ and $(\text{Ker}(\overline{\square_0}))^\perp$, respectively ([K1], [K2]). Since P_0 commutes with \square_0 and $\text{Ker}(\overline{\square_0}) \subset \text{Ker} P_0$, we have

$$(3.23) \quad \square_0 \varphi \in (\text{Ker} P_0)^\perp \subset (\text{Ker}(\overline{\square_0}))^\perp$$

for $\varphi \in (\text{Ker} P_0)^\perp$. We can then estimate

$$\begin{aligned}
(3.24) \quad \|\varphi\|_{S^{k+4,2}} &\leq C\|\square_0\varphi\|_{S^{k+2,2}} \\
&\leq C\|\overline{\square}_0\square_0\varphi\|_{S^{k,2}} = C\|P_0\varphi\|_{S^{k,2}}
\end{aligned}$$

for $\varphi \in (Ker P_0)^\perp (\subset (Ker(\square_0))^\perp)$ by (3.23). By (3.24), we mean that P_0 is subelliptic on $(Ker P_0)^\perp$. It follows that there exists a unique C^∞ smooth solution λ^\perp of (3.10) for a short time (noting that P_0 is essentially positive by Theorem 5.3, which explains the negative sign in (3.10)). On the other hand, we can apply the contraction mapping principle to show the short time existence of a unique C^∞ smooth solution λ_{ker} to (3.11) (observing that λ_{ker} satisfies (3.12)). The long time solution then follows from Proposition 3.6, the Sobolev embedding theorem for $S^{k,2}$, and the standard argument for extending the solution at the maximal time T .

4. A Priori Estimates and Asymptotic Convergence

We can obtain an a priori $S^{2,2}$ estimate under some analytic assumption replacing the torsion free condition. Also a condition on the background Q -curvature will assure that the bound is uniform, i.e., independent of the time. Under the same condition together with the vanishing torsion, we will then have the higher-order estimates with uniform bounds. Therefore we are able to prove the asymptotic convergence of solutions of (1.6).

Lemma 4.1. *Suppose $(Q_0)_{ker} = 0$. Then the solution λ^\perp of (3.10) satisfies the following a priori estimate*

$$(4.1) \quad \mathcal{E}(\lambda^\perp) \equiv \int_M P_0 \lambda^\perp \cdot \lambda^\perp d\mu_0 + \int_M Q_0 \lambda^\perp d\mu_0 \leq \beta^2$$

for a constant β independent of the time.

Proof. The condition $(Q_0)_{ker} = 0$ implies that

$$\int_M Q_0 \lambda_{ker} d\mu_0 = 0.$$

It follows that

$$\begin{aligned}
& \int_M P_0 \lambda \cdot \lambda d\mu_0 + \int_M Q_0 \lambda d\mu_0 \\
&= \int_M P_0 \lambda^\perp \cdot (\lambda_{\ker} + \lambda^\perp) d\mu_0 + \int_M Q_0 (\lambda_{\ker} + \lambda^\perp) d\mu_0 \\
&= \int_M P_0 \lambda^\perp \cdot \lambda^\perp d\mu_0 + \int_M Q_0 \lambda^\perp d\mu_0.
\end{aligned}$$

Hence from Lemma 3.4, we obtain (4.1).

■

Proof of Theorem 1.2:

The essential positivity of P_0 assures that for some constant Υ , there holds

$$(4.2) \quad \int_M P_0 \lambda^\perp \cdot \lambda^\perp d\mu_0 \geq \Upsilon \int_M (\lambda^\perp)^2 d\mu_0.$$

From the proof of Proposition 3.6, we have (3.13), i.e., a bound for the L^2 norm of λ^\perp , depending on the maximal time T . If in addition we assume $(Q_0)_{\ker} = 0$, then by (4.1), (4.2), and the Young inequality, we obtain

$$(4.3) \quad \int_M (\lambda^\perp)^2 d\mu_0 \leq C(\Upsilon, Q_0, \theta_0, \beta)$$

and

$$(4.4) \quad \int_M P_0 \lambda \cdot \lambda d\mu_0 = \int_M P_0 \lambda^\perp \cdot \lambda^\perp d\mu_0 \leq C(\Upsilon, Q_0, \theta_0, \beta)$$

for all $t \geq 0$.

Now if the condition $(*)$ holds on $(M, J, [\theta_0])$, that is,

$$\int_M |\overset{0}{\nabla}^2 \lambda^\perp|^2 d\mu_0 \leq (2 + \varepsilon_0) \int_M (\Delta_0 \lambda^\perp)^2 d\mu_0 + C(\varepsilon_0) \int_M |\overset{0}{\nabla} \lambda^\perp|^2 d\mu_0.$$

From this and Lemma 3.2, we have

$$2 \int_M P_0 \lambda^\perp \cdot \lambda^\perp d\mu_0 \geq (1 - \varepsilon_0) \int_M (\Delta_0 \lambda^\perp)^2 d\mu_0 - C \int_M |\overset{0}{\nabla} \lambda^\perp|^2 d\mu_0.$$

By applying (4.3), integrating by parts and the Young inequality, we compute

$$\begin{aligned}
(4.5) \quad 2 \int_M P_0 \lambda^\perp \cdot \lambda^\perp d\mu_0 &= (1 - \varepsilon_0) \int_M (\Delta_0 \lambda^\perp)^2 d\mu_0 + C \int_M \lambda^\perp \Delta_0 \lambda^\perp d\mu_0 \\
&\geq (1 - \varepsilon_0 - \varepsilon) \int_M (\Delta_0 \lambda^\perp)^2 d\mu_0 - C(\varepsilon_0, \tfrac{1}{\varepsilon}),
\end{aligned}$$

for small $\varepsilon > 0$. Now if we choose ε small enough such that $(1 - \varepsilon_0 - \varepsilon) = \delta > 0$, then

$$(4.6) \quad 2 \int_M P_0 \lambda^\perp \cdot \lambda^\perp d\mu_0 \geq \delta \int_M (\Delta_0 \lambda^\perp)^2 d\mu_0 - C(\varepsilon_0, \frac{1}{\varepsilon}).$$

Finally from (4.4) and (4.6), there exists a positive constant $C = C(\delta, \Upsilon, Q_0, \theta_0, \beta)$ such that

$$\int_M (\Delta_0 \lambda^\perp)^2 d\mu_0 \leq C,$$

and hence

$$(4.7) \quad \|\lambda^\perp\|_{S^{2,2}} \leq C$$

for all $t \geq 0$ by (4.3) (the constant C depends on the maximal time T by (3.13) if we don't assume $(Q_0)_{\ker} = 0$).

To deal with λ_{\ker} , we consider $\lambda - \bar{\lambda}$. Without the condition $(Q_0)_{\ker} = 0$, we have obtained the bound for $\|\lambda\|_{S^{2,2}}$, depending on T , from the later part of the proof of Proposition 3.6. Let us assume $(Q_0)_{\ker} = 0$ below and see how to get an uniform bound.

Under the condition $(Q_0)_{\ker} = 0$, the equation (3.11) reads

$$\frac{\partial \lambda_{\ker}}{\partial t} = r,$$

and hence we have

$$(4.8) \quad \lambda_{\ker}(t, p) = \lambda_{\ker}(0, p) + \int_0^t r dt.$$

It follows that

$$\lambda - \bar{\lambda} = (\lambda^\perp - \bar{\lambda}^\perp) + [\lambda_{\ker}(0, p)]$$

and from (4.7) and (4.3) that

$$\|\lambda - \bar{\lambda}\|_{S^{2,2}} \leq C_1 + \|\lambda^\perp - \bar{\lambda}^\perp\|_{S^{2,2}} \leq C_2$$

for all $t \geq 0$. Therefore by the Sobolev embedding theorem for $S^{k,p}$, we have

$$(4.9) \quad \|\lambda - \bar{\lambda}\|_{S^{1,4}} \leq C_3$$

for all $t \geq 0$.

Now again by using Theorem 2.3, we have

$$\int_M e^{4(\lambda - \bar{\lambda})} d\mu_0 \leq C \exp(C \|\lambda - \bar{\lambda}\|_{S^{1,4}}) \leq C_4$$

for all $t \geq 0$. Together with $\int_M e^{4\lambda} d\mu_0$ being invariant under the flow, we conclude that

$$(4.10) \quad C_5 \geq \bar{\lambda} \geq -C_5.$$

Thus

$$\|\lambda\|_{S^{2,2}} \leq C$$

(and also (from Theorem 2.3))

$$(4.11) \quad \int_M e^{\alpha\lambda} d\mu_0 \leq C$$

for all real number α and all $t \geq 0$).

Since Δ_0 is self adjoint, we can deduce the following result by the Young inequality.

Lemma 4.2. *(interpolation inequality for Δ_0) On a closed pseudohermitian manifold (M, J, θ_0) , given $\varepsilon > 0$, there exists a constant $C(\varepsilon)$ such that*

$$(4.12) \quad \int_M (\Delta_0 \varphi)^2 d\mu_0 \leq \varepsilon \int_M (\Delta_0^2 \varphi)^2 d\mu_0 + C(\varepsilon) \int_M \varphi^2 d\mu_0$$

for all real valued functions $\varphi \in C^\infty(M)$.

Proof of Theorem 1.3:

We compute

$$(4.13) \quad \begin{aligned} & \frac{d}{dt} \int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 \\ &= 2 \int_M (\Delta_0^k \lambda^\perp) (\Delta_0^k \frac{\partial \lambda^\perp}{\partial t}) d\mu_0 \\ &= -2 \int_M (\Delta_0^k \lambda^\perp) [\Delta_0^k (Q_0 + 2P_0 \lambda^\perp)] d\mu_0 \quad (\text{by (3.10)}) \\ &= -2 \int_M (\Delta_0^k \lambda^\perp) [\Delta_0^k (Q_0)] d\mu_0 - 4 \int_M (\Delta_0^k \lambda^\perp) [\Delta_0^k (P_0 \lambda^\perp)] d\mu_0. \end{aligned}$$

Next we will estimate the second term in the last line of (4.13). From the Bochner formula (3.1), (3.16), and (3.14), we compute

$$\begin{aligned}
(4.14) \quad & -4 \int_M (\Delta_0^k \lambda^\perp) [\Delta_0^k (P_0 \lambda^\perp)] d\mu_0 \\
& = -4 \int_M (\Delta_0^k \lambda^\perp) [\Delta_0^k (\Delta_0^2 \lambda^\perp + T_0^2 \lambda^\perp)] d\mu_0 + E_1 \\
& = -4 \int_M (\Delta_0^{k+1} \lambda^\perp)^2 d\mu_0 - 4 \int_M (\Delta_0^k \lambda^\perp) [\Delta_0^k T_0^2 \lambda^\perp] d\mu_0 + E_1 \\
& = -4 \int_M (\Delta_0^{k+1} \lambda^\perp)^2 d\mu_0 - 4 \int_M (\Delta_0^k \lambda^\perp) [T_0^2 \Delta_0^k \lambda^\perp] d\mu_0 + E_1 + E_2 \\
& = -4 \int_M (\Delta_0^{k+1} \lambda^\perp)^2 d\mu_0 + 4 \int_M (T_0 \Delta_0^k \lambda^\perp)^2 d\mu_0 + E_1 + E_2 \\
& = 2 \int_M |\nabla^2 (\Delta_0^k \lambda^\perp)|^2 d\mu_0 - 3 \int_M (\Delta_0^{k+1} \lambda^\perp)^2 d\mu_0 + E_1 + E_2 + E_3
\end{aligned}$$

where

$$\begin{aligned}
E_1 &= \int_M (\Delta_0^k \lambda^\perp) \Delta_0^k (O_2(\lambda^\perp)) d\mu_0, \\
E_2 &= \int_M (\Delta_0^k \lambda^\perp) O_{2k}(\lambda^\perp) d\mu_0, \\
E_3 &= \int_M (\Delta_0^k \lambda^\perp) O_{2k+2}(\lambda^\perp) d\mu_0,
\end{aligned}$$

with O_j being some differential operator of weight j (see Appendix A for the definition). Using the subelliptic estimate and the Young inequality, we can estimate the "error terms" E'_j s by

$$\begin{aligned}
(4.15) \quad & |E| \\
& = |E_1 + E_2 + E_3| \\
& \leq C(k, \varepsilon_1) \int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 + \varepsilon_1 (\int_M (\Delta_0^{k+1} \lambda^\perp)^2 d\mu_0 + \int_M (\lambda^\perp)^2 d\mu_0).
\end{aligned}$$

Note that Δ_0 (and hence Δ_0^k) maps $\text{Ker}(P_0)$, and hence $\text{Ker}(P_0)^\perp$, into itself by assumption. So we can apply the condition (*) to the first term of the last line in (4.14). Together with substituting (4.15) into (4.14), we obtain

$$\begin{aligned}
(4.16) \quad & -4 \int_M (\Delta_0^k \lambda^\perp) [\Delta_0^k (P_0 \lambda^\perp)] d\mu_0 \\
& \leq -2(1 - \varepsilon_0 - \frac{\varepsilon_1}{2}) \int_M (\Delta_0^{k+1} \lambda^\perp)^2 d\mu_0 + C(k, \varepsilon_1) \int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 + \varepsilon_1 \int_M (\lambda^\perp)^2 d\mu_0.
\end{aligned}$$

Taking ε_1 small so that $1 - \varepsilon_0 - \frac{\varepsilon_1}{2} > 0$. From the interpolation inequality (4.12) with $\varphi = \Delta_0^{k-1} \lambda^\perp$, we get

$$(4.17) \quad \int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 \leq \varepsilon \int_M (\Delta_0^{k+1} \lambda^\perp)^2 d\mu_0 + C(\varepsilon) \int_M (\Delta_0^{k-1} \lambda^\perp)^2 d\mu_0.$$

By choosing ε (depending on k, ε_1) small enough, we can absorb the middle term of the second line into the first term of the second line in (4.16). Substituting (4.17) into (4.16) and then substituting the result into (4.13), we obtain

$$(4.18) \quad \begin{aligned} & \frac{d}{dt} \int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 \\ & \leq -C \int_M (\Delta_0^{k+1} \lambda^\perp)^2 d\mu_0 + C(k) \int_M (\Delta_0^{k-1} \lambda^\perp)^2 d\mu_0 + C \int_M (\lambda^\perp)^2 d\mu_0. \end{aligned}$$

In view of the subellipticity of Δ_0^k , we have

$$\int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 \leq C(k) \left(\int_M (\Delta_0^{k+1} \lambda^\perp)^2 d\mu_0 + \int_M (\lambda^\perp)^2 d\mu_0 \right).$$

Therefore we can reduce (4.18) to

$$(4.19) \quad \begin{aligned} & \frac{d}{dt} \int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 \\ & \leq -C(k) \int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 + C(k) \int_M (\Delta_0^{k-1} \lambda^\perp)^2 d\mu_0 + C(k) \int_M (\lambda^\perp)^2 d\mu_0. \end{aligned}$$

Now we apply induction on $k \geq 2$ to (4.18). When $k = 2$, the last two terms of (4.19) are bounded due to Theorem 1.2. So by Lemma 3.5, we get

$$\int_M (\Delta_0^2 \lambda^\perp)^2 d\mu_0 \leq C(T)$$

and $C(T)$ is independent of T if the condition $(Q_0)_{\ker} = 0$ is imposed. By induction hypothesis, $\int_M (\Delta_0^{k-1} \lambda^\perp)^2 d\mu_0$ is bounded. Also recall in the proof of Proposition 3.6 that the essential positivity of P_0 and the energy estimate (3.9) imply the L^2 bound of λ^\perp (see (3.13) and (4.3) if, in addition, $(Q_0)_{\ker} = 0$). So the last two terms of (4.19) are bounded. Applying Lemma 3.5 again, we conclude that

$$(4.20) \quad \int_M (\Delta_0^k \lambda^\perp)^2 d\mu_0 \leq C(k, T)$$

where $C(k, T)$ is independent of T if the condition $(Q_0)_{\ker} = 0$ is imposed. From (3.13) ((4.3) if, in addition, $(Q_0)_{\ker} = 0$), (4.20), and the subelliptic estimate, we can then have

$$\|\lambda^\perp\|_{S^{2k,2}} \leq C(k, T)$$

($C(k, T)$ is independent of T if the condition $(Q_0)_{\ker} = 0$ is imposed). Observe that the argument from (3.20) to (3.22) in the proof of Proposition 3.6 still works without torsion free condition. We therefore have the following estimate

$$||\lambda||_{S^{2k,2}} \leq C(k, T)$$

with $C(k, T)$ being independent of T if the condition $(Q_0)_{\ker} = 0$ is imposed.

Proposition 4.3. *Let $(M, J, [\theta_0])$ be a CR 3-manifold with $\overset{0}{A}_{11} = 0$ and $(Q_0)_{\ker} = 0$. There exists a constant $C = C(\Upsilon, Q_0, \theta_0, \beta)$ such that*

$$||\lambda||_{S^{4,2}} \leq C$$

for all $t \geq 0$. In particular, there holds

$$|\lambda| \leq C$$

for all $t \geq 0$. Moreover, we have

$$(4.21) \quad ||\lambda||_{S^{2k,2}} \leq C(k)$$

for all $t \geq 0$.

Proof. Observe that λ has an uniform L^2 bound if $(Q_0)_{\ker} = 0$ in the proof of Proposition (3.6), and hence the constants $C(\dots, T)$, $C(T)$, and $C(k, T)$ can be replaced by C or $C(k)$ (independent of T) from (3.13) to the end of the proof.

■

Proof of Theorem 1.4:

By Theorem 1.1, we have a long time solution. Starting from (1.7), we compute

$$\begin{aligned} (4.22) \quad \frac{d^2}{dt^2} \mathcal{E}(\theta) &= -4 \int (Q_0 + 2P_0\lambda) P_0 \frac{\partial \lambda}{\partial t} d\mu_0 \\ &= 4 \int (Q_0^\perp + 2P_0\lambda) P_0 (Q_0^\perp + 2P_0\lambda) d\mu_0 \text{ by (1.6)} \\ &\geq 4\Upsilon \int (Q_0^\perp + 2P_0\lambda)^2 \end{aligned}$$

by essential positivity of P_0 (which is implied by the torsion free condition; see Theorem 5.3). Therefore $\frac{d}{dt} \mathcal{E}(\theta)$ is nondecreasing, and hence

$$(4.23) \quad \int_M e^{4\lambda} Q^2 d\mu \ (\geq 0) \text{ is nonincreasing.}$$

By (4.21), we can find a sequence of times t_j such that $\lambda_j \equiv \lambda(\cdot, t_j)$ converges to λ_∞ in C^∞ topology as $t_j \rightarrow \infty$. On the other hand, integrating (1.7) gives

$$(4.24) \quad \mathcal{E}(\lambda_\infty) - \mathcal{E}(\lambda_0) = - \int_0^\infty \int_M e^{4\lambda} Q^2 d\mu dt.$$

In view of (4.23), (4.24), we obtain

$$(4.25) \quad 0 = \lim_{t \rightarrow \infty} \int_M e^{4\lambda} Q^2 d\mu = \int_M e^{4\lambda_\infty} Q_\infty^2 d\mu_\infty$$

where Q_∞ denotes the Q -curvature with respect to (J, θ_∞) , $\theta_\infty = e^{2\lambda_\infty} \theta_0$, and $d\mu_\infty = e^{4\lambda_\infty} d\mu_0$. It follows that $Q_\infty = 0$.

In the following, we are going to prove the smooth convergence for all time. First we want to prove that λ converges to λ_∞ in L^2 . Write $\lambda_\infty = \lambda_\infty^\perp + (\lambda_\infty)_{\ker}$. Observe that $(\|\cdot\|_2$ denotes the L^2 norm with respect to the volume form $d\mu_0$)

$$(4.26) \quad \begin{aligned} \|\lambda_\infty^\perp - \lambda^\perp\|_2 &\leq \|\lambda_\infty^\perp - \lambda^\perp\|_{S^{4,2}} \\ &\leq C \|2P_0(\lambda_\infty^\perp - \lambda^\perp)\|_2 \\ &= C \|2P_0\lambda^\perp + Q_0^\perp\|_2 \end{aligned}$$

by the subellipticity of P_0 on $(\text{Ker } P_0)^\perp$ and $0 = Q_\infty = e^{-4\lambda_\infty} (2P_0\lambda_\infty^\perp + Q_0^\perp)$ ($(Q_0)_{\ker} = 0$ by assumption). We compute

$$(4.27) \quad \begin{aligned} &| \mathcal{E}(\lambda_\infty^\perp) - \mathcal{E}(\lambda^\perp) | \\ &= | \int_0^1 \frac{d}{ds} \mathcal{E}(\lambda^\perp + s(\lambda_\infty^\perp - \lambda^\perp)) ds | \\ &= | \int_0^1 \int_M [2P_0(\lambda^\perp + s(\lambda_\infty^\perp - \lambda^\perp)) + Q_0^\perp] \cdot (\lambda_\infty^\perp - \lambda^\perp) d\mu_0 ds | \\ &\leq \int_0^1 \|2P_0(\lambda^\perp + s(\lambda_\infty^\perp - \lambda^\perp)) + Q_0^\perp\|_2 \|\lambda_\infty^\perp - \lambda^\perp\|_2 ds \\ &\leq C_1 \|2P_0\lambda^\perp + Q_0^\perp\|_2^2 \end{aligned}$$

by the Cauchy inequality and (4.26). Let ϑ be a number between 0 and $\frac{1}{2}$. It follows from (4.27) that

$$(4.28) \quad | \mathcal{E}(\lambda_\infty^\perp) - \mathcal{E}(\lambda^\perp) |^{1-\vartheta} \leq C_2 \|2P_0\lambda^\perp + Q_0^\perp\|_2^{2(1-\vartheta)} \leq C_2 \|2P_0\lambda^\perp + Q_0^\perp\|_2$$

for t large by noting that $2(1 - \vartheta) > 1$ and $\|2P_0\lambda^\perp + Q_0^\perp\|_2$ tends to 0 as $t \rightarrow \infty$ by (4.25). Next we compute

$$\begin{aligned}
 & -\frac{d}{dt}(\mathcal{E}(\lambda^\perp) - \mathcal{E}(\lambda_\infty^\perp))^\vartheta \\
 (4.29) \quad & = -\vartheta(\mathcal{E}(\lambda^\perp) - \mathcal{E}(\lambda_\infty^\perp))^{\vartheta-1} \frac{d}{dt}(\mathcal{E}(\lambda^\perp) - \mathcal{E}(\lambda_\infty^\perp)) \\
 & = +\vartheta(\mathcal{E}(\lambda^\perp) - \mathcal{E}(\lambda_\infty^\perp))^{\vartheta-1} \|2P_0\lambda^\perp + Q_0^\perp\|_2 \|\dot{\lambda}^\perp\|_2 \\
 & \geq \vartheta C_2^{-1} \|\dot{\lambda}^\perp\|_2
 \end{aligned}$$

by (1.7), (4.28), and noting that $\dot{\lambda}^\perp = -(2P_0\lambda^\perp + Q_0^\perp)$ (see (3.10)) and hence the left side is nonnegative. We learned the above trick of raising the power to ϑ from [S]. Integrating (4.29) with respect to t gives

$$(4.30) \quad \int_0^\infty \|\dot{\lambda}^\perp\|_2 dt < +\infty.$$

Observing that $\frac{d}{dt}(\lambda^\perp - \lambda_\infty^\perp) = \dot{\lambda}^\perp$ and $-\frac{d}{dt}\|\lambda^\perp - \lambda_\infty^\perp\|_2^2 \leq C\|\dot{\lambda}^\perp\|_2$, we can then deduce

$$(4.31) \quad \lim_{t \rightarrow \infty} \|\lambda^\perp - \lambda_\infty^\perp\|_2^2 = 0$$

by (4.30). On the other hand, we can estimate $|r(t)| \leq C\|2P_0\lambda^\perp + Q_0^\perp\|_2 = C\|\dot{\lambda}^\perp\|_2$ by (4.11). It follows from (4.30) that

$$(4.32) \quad \int_0^\infty |r(t)| dt < +\infty.$$

So in view of (4.8) and (4.32), λ_{\ker} converges to $(\lambda_\infty)_{\ker} = (\lambda_{\ker})_\infty$ as $t \rightarrow \infty$ (not just a sequence of times). Since $\lambda_{\ker} - (\lambda_\infty)_{\ker} = -\int_t^\infty r(t) dt$ is a function of time only, we also have

$$(4.33) \quad \lim_{t \rightarrow \infty} \|\lambda_{\ker} - (\lambda_\infty)_{\ker}\|_{S^{k,2}} = 0$$

for any nonnegative integer k . With λ^\perp replaced by $\lambda^\perp - \lambda_\infty^\perp$ in the argument to deduce (3.19), we obtain

$$(4.34) \quad \frac{d}{dt} \int_M (\Delta_0^k(\lambda^\perp - \lambda_\infty^\perp))^2 d\mu_0 \leq -4\Upsilon \int_M (\Delta_0^k(\lambda^\perp - \lambda_\infty^\perp))^2 d\mu_0 + C(k) \|\lambda^\perp - \lambda_\infty^\perp\|_2.$$

By Lemma 3.5 and (4.31), we get

$$(4.35) \quad \lim_{t \rightarrow \infty} \int_M (\Delta_0^k(\lambda^\perp - \lambda_\infty^\perp))^2 d\mu_0 = 0.$$

Hence by the subellipticity of Δ_0^k , we conclude from (4.35) and (4.31) that

$$\lim_{t \rightarrow \infty} \|\lambda^\perp - \lambda_\infty^\perp\|_{S^{2k,2}} = 0.$$

Together with 4.33, we have proved that λ converges to λ_∞ smoothly as $t \rightarrow \infty$.

5. Essential Positivity of the CR Paneitz Operator

Let (M, J, θ) be a closed pseudohermitian 3-manifold with zero torsion. We will prove the essential positivity of the CR Paneitz operator P .

Since the torsion of (M, J, θ) is zero, the Lie derivative of the CR structure J with respect to T (characteristic vector field) is zero. Hence M admits a smooth CR action of R , which is transverse to the contact bundle. This implies that (M, J) is embeddable by Theorem 2.1. in [Le]. Therefore the L^2 -closure of the Kohn Laplacian \square_b has the closed range ([K1]) and hence subelliptic estimates on the orthogonal complement of the kernel of \square_b in L^2 ([K1], [K2]). We use the same symbol \square_b to denote the L^2 -closure.

On the other hand, since \square_b is a closed operator, $\text{Ker}(\square_b)$ is closed, and we have the following direct sum decomposition

$$(5.1) \quad L^2 = \text{Ker}(\square_b) \oplus R(\square_b),$$

where $R(\square_b)$ is the range of \square_b .

Let $C^\infty(M, C)$ be the space of all C^∞ smooth complex-valued functions on M and $C_K^\infty = C^\infty(M, C) \cap \text{Ker}(\square_b)$. Then we have $C^\infty(M, C) = C_K^\infty \oplus (C_K^\infty)^\perp$. By (5.1) and subelliptic estimates on the orthogonal complement of the kernel of \square_b in L^2 , we see that $(C_K^\infty)^\perp = \{\square_b \varphi \mid \varphi \in C^\infty(M, C)\}$.

Let $H : C^\infty(M, C) \rightarrow C_K^\infty$ be the projection. Define the Green operator $G : C^\infty(M, C) \rightarrow (C_K^\infty)^\perp$ by

$$G(\alpha) = \omega \in (C_K^\infty)^\perp,$$

where ω is the unique solution of $\square_b \omega = \alpha - H(\alpha)$. It is easy to check that G is symmetric and positive on $(C_K^\infty)^\perp$. Moreover, since \square_b has subelliptic estimates, G is compact on the orthogonal complement of $\text{Ker}(\square_b)$.

Lemma 5.1. *Let $\eta = \sup \{ \|G\varphi\| : \varphi \in C^\infty(M, C), \|\varphi\| = 1, \varphi \in (C_K^\infty)^\perp \}$. Then $\frac{1}{\eta}$ is an eigenvalue of \square_b .*

Proof. Let $\{\varphi_j\} \subset (C_K^\infty)^\perp$ be a maximizing sequence for η , that is, $\|\varphi_j\| = 1$ and $\|G\varphi_j\| \rightarrow \eta$. Since G is symmetric, we have

$$\begin{aligned} \|G^2\varphi_j - \eta^2\varphi_j\|^2 &= \|G^2\varphi_j\|^2 - 2\eta^2 \langle G^2\varphi_j, \varphi_j \rangle + \eta^4 \\ &\leq \eta^2 \|G\varphi_j\|^2 - 2\eta^2 \|G\varphi_j\|^2 + \eta^4. \end{aligned}$$

This inequality means that $\|G^2\varphi_j - \eta^2\varphi_j\| \rightarrow 0$ as $j \rightarrow \infty$.

Let $\psi_j = G\varphi_j - \eta\varphi_j$. Since G is positive on $(C_K^\infty)^\perp$, we have

$$\begin{aligned} \langle \psi_j, G^2\varphi_j - \eta^2\varphi_j \rangle &= \langle \psi_j, G\psi_j + \eta\psi_j \rangle \\ &= \langle \psi_j, G\psi_j \rangle + \eta \|\psi_j\|^2 \\ &\geq \eta \|\psi_j\|^2. \end{aligned}$$

This inequality, together with the above, implies that $\|G\varphi_j - \eta\varphi_j\| \rightarrow 0$.

On the other hand, since G is compact on the orthogonal complement of the kernel of \square_b , there exists a subsequence of $\{\varphi_j\}$, also denoted by $\{\varphi_j\}$, such that $\{G\varphi_j\}$ converges to a function $f \in L^2$. We have, for all $\varphi \in C^\infty(M, C)$,

$$\begin{aligned} \left\langle \left(\square_b - \frac{1}{\eta}\right)f, \varphi \right\rangle &= \left\langle f, \left(\square_b - \frac{1}{\eta}\right)\varphi \right\rangle \\ &= \lim_{j \rightarrow \infty} \left\langle G\varphi_j, \left(\square_b - \frac{1}{\eta}\right)\varphi \right\rangle \\ &= \lim_{j \rightarrow \infty} \left\langle \varphi_j - \frac{1}{\eta}G\varphi_j, \varphi \right\rangle \\ &= \lim_{j \rightarrow \infty} \frac{1}{\eta} \langle \eta\varphi_j - G\varphi_j, \varphi \rangle \\ &= 0. \end{aligned}$$

That is, $f \in L^2$ is a nontrivial weak solution of $(\square_b - \frac{1}{\eta})f = 0$. Actually $f \perp \text{Ker}\square_b$, so f is smooth and $\frac{1}{\eta}$ is an eigenvalue of \square_b .

■

Lemma 5.2. *Let (M, J, θ) be a closed pseudohermitian 3-manifold. Suppose that (M, J) is embeddable. Then \square_b has discrete eigenvalues: $0 = \mu_0 < \mu_1 \leq \mu_2 \leq \dots$*

$\cdots \leq \mu_n \rightarrow \infty$ and the corresponding eigenfunctions $\{\varphi_i \mid i \geq 1\}$ satisfying $\square_b \varphi_i = \mu_i \varphi_i$, $\varphi_i \in C^\infty(M, C)$ can be chosen so that $\{\varphi_i\}$ forms an orthonormal basis of $(\text{Ker} \square_b)^\perp$ in L^2 .

Proof. Let $E(\mu)$ be the eigenspace with respect to the eigenvalue μ . By the subelliptic estimates, $\dim E(\mu) < \infty$ if $\mu > 0$.

For $j \geq 1$, let $\eta_j \equiv \sup \left\{ \|G\varphi\| : \varphi \in C^\infty(M, C), \|\varphi\| = 1, \varphi \in \left(\bigoplus_{i=0}^{j-1} E(\frac{1}{\eta_i}) \right)^\perp \right\}$. Here we denote $E(\frac{1}{\eta_0}) = C_K^\infty$. A similar argument as in Lemma 5.1 shows that $\frac{1}{\eta_j}$ is an eigenvalue of \square_b . Therefore we have the discrete eigenvalues $0 = \mu_0 < \mu_1 \leq \mu_2 \leq \cdots \leq \mu_n$, $n \rightarrow \infty$. By the subelliptic estimates, $\{\mu_n\}$ can not have a finite limit. Thus $\mu_n \rightarrow \infty$ as $n \rightarrow \infty$.

Let $0 = \mu_0 < \mu_1 \leq \mu_2 \leq \cdots$ be the eigenvalues of \square_b , where each eigenvalue is included as many times as the dimension of its eigenspace, with a corresponding orthonormal sequence of eigenfunctions $\{\varphi_i\}$. Let $\alpha \in C^\infty(M, C)$, there exists $\beta \in (C_K^\infty)^\perp$ such that $G\beta = \alpha - H(\alpha)$. It is easy to check that $\langle \alpha, \varphi_i \rangle \varphi_i = G(\langle \beta, \varphi_i \rangle \varphi_i)$, so we have

$$\begin{aligned} \left\| \alpha - \sum_{i=1}^n \langle \alpha, \varphi_i \rangle \varphi_i - H(\alpha) \right\| &= \left\| G\beta - \sum_{i=1}^n \langle \alpha, \varphi_i \rangle \varphi_i \right\| \\ &= \left\| G\beta - G \left(\sum_{i=1}^n \langle \beta, \varphi_i \rangle \varphi_i \right) \right\| \\ &= \left\| G \left(\beta - \sum_{i=1}^n \langle \beta, \varphi_i \rangle \varphi_i \right) \right\| \\ &\leq \frac{1}{\mu_{n+1}} \left\| \beta - \sum_{i=1}^n \langle \beta, \varphi_i \rangle \varphi_i \right\| \\ &\leq \frac{1}{\mu_{n+1}} \|\beta\|. \end{aligned}$$

Then $\mu_n \rightarrow \infty$ implies that

$$\lim_{n \rightarrow \infty} \left\| \alpha - \sum_{i=1}^n \langle \alpha, \varphi_i \rangle \varphi_i - H(\alpha) \right\| = 0.$$

■

Theorem 5.3. *If the torsion of (M, J, θ) is zero, then the CR Paneitz operator P is essentially positive.*

Proof. First, zero torsion implies that (i) the CR Paneitz operator $P = \square_b \overline{\square_b}$, and (ii) the Kohn Laplacian \square_b and $\overline{\square_b}$ commute, so they are diagonalized simultaneously on the finite dimensional eigenspace of \square_b with respect to any nonzero eigenvalue.

Therefore, we can choose an orthonormal basis $\{\varphi_i\}$ such that each eigenfunction $\in \{\varphi_i\}$ is also an eigenfunction of $\overline{\square}_b$, and hence, of P . We know that the eigenvalues of \square_b (and hence of $\overline{\square}_b$) are all nonnegative. Therefore by Lemma 5.2, P is essentially positive.

■

APPENDIX A.

We will give a brief introduction to pseudohermitian geometry (see [L1], [L2] for more details). Let M be a closed 3-manifold with an oriented contact structure ξ . There always exists a global contact form θ , obtained by patching together local ones with a partition of unity. The characteristic vector field of θ is the unique vector field T such that $\theta(T) = 1$ and $\mathcal{L}_T\theta = 0$ or $d\theta(T, \cdot) = 0$. A CR structure compatible with ξ is a smooth endomorphism $J : \xi \rightarrow \xi$ such that $J^2 = -\text{identity}$. A pseudohermitian structure compatible with ξ is a CR -structure J compatible with ξ together with a global contact form θ .

Given a pseudohermitian structure (J, θ) , we can choose a complex vector field Z_1 , an eigenvector of J with eigenvalue i , and a complex 1-form θ^1 such that $\{\theta, \theta^1, \theta^{\bar{1}}\}$ is dual to $\{T, Z_1, Z_{\bar{1}}\}$. It follows that $d\theta = ih_{1\bar{1}}\theta^1 \wedge \theta^{\bar{1}}$ for some nonzero real function $h_{1\bar{1}}$. If $h_{1\bar{1}}$ is positive, we call such a pseudohermitian structure (J, θ) positive, and we can choose a Z_1 (hence θ^1) such that $h_{1\bar{1}} = 1$. That is to say

$$d\theta = i\theta^1 \wedge \theta^{\bar{1}}.$$

We will always assume our pseudohermitian structure (J, θ) is positive and $h_{1\bar{1}} = 1$ throughout the paper. The pseudohermitian connection of (J, θ) is the connection $\nabla^{\psi.h.}$ on $TM \otimes C$ (and extended to tensors) given by

$$\nabla^{\psi.h.} Z_1 = \omega_1^1 \otimes Z_1, \nabla^{\psi.h.} Z_{\bar{1}} = \omega_{\bar{1}}^{\bar{1}} \otimes Z_{\bar{1}}, \nabla^{\psi.h.} T = 0$$

in which the 1-form ω_1^1 is uniquely determined by the following equation with a normalization condition:

$$\begin{aligned} (A.1) \quad d\theta^1 &= \theta^1 \wedge \omega_1^1 + A_{\bar{1}}^1 \theta \wedge \theta^{\bar{1}} \\ 0 &= \omega_1^1 + \omega_{\bar{1}}^{\bar{1}}. \end{aligned}$$

The coefficient $A^1_{\bar{1}}$ in (A.1) is called the (pseudohermitian) torsion. Since $h_{1\bar{1}} = 1$, $A_{\bar{1}\bar{1}} = h_{1\bar{1}}A^1_{\bar{1}} = A^1_{\bar{1}}$. And A_{11} is just the complex conjugate of $A_{\bar{1}\bar{1}}$. Differentiating ω_1^1 gives

$$d\omega_1^1 = W\theta^1 \wedge \theta^{\bar{1}} + 2i\text{Im}(A_{11,\bar{1}}\theta^1 \wedge \theta)$$

where W is the Tanaka-Webster curvature.

We can define the covariant differentiations with respect to the pseudohermitian connection. For instance, $f_1 = Z_1 f$, $f_{1\bar{1}} = Z_{\bar{1}}Z_1 f - \omega_1^1(Z_{\bar{1}})Z_1 f$ for a (smooth) function f . We define the sub-gradient operator ∇_b and the sub-Laplacian operator Δ_b by

$$\begin{aligned}\nabla_b f &= f_{\bar{1}}Z_1 + f_1Z_{\bar{1}}, \\ \Delta_b f &= f_{1\bar{1}} + f_{\bar{1}1}.\end{aligned}$$

respectively. We also define the Levi form $\langle, \rangle_{J,\theta}$ by

$$\langle V, U \rangle_{J,\theta} = 2d\theta(V, JU) = v_1 u_{\bar{1}} + v_{\bar{1}} u_1,$$

for $V = v_1 Z_{\bar{1}} + v_{\bar{1}} Z_1, U = u_1 Z_{\bar{1}} + u_{\bar{1}} Z_1$ in ξ and

$$(V, U)_{J,\theta} = \int_M \langle V, U \rangle_{J,\theta} \theta \wedge d\theta.$$

For a vector $X \in \xi$, we define $|X|^2 \equiv \langle X, X \rangle_{J,\theta}$. It follows that $|\nabla_b f|^2 = 2f_1 f_{\bar{1}}$ for a real valued smooth function f . Also the square modulus of the sub-Hessian $\nabla_b^2 f$ of f reads $|\nabla_b^2 f|^2 = 2f_{11} f_{\bar{1}\bar{1}} + 2f_{1\bar{1}} f_{\bar{1}1}$. We recall below what the Folland-Stein space $S^{k,p}$ is. Let D denote a differential operator acting on functions. We say D has weight m , denoted $w(D) = m$, if m is the smallest integer such that D can be locally expressed as a polynomial of degree m in vector fields tangent to the contact bundle ξ . We define the Folland-Stein space $S^{k,p}$ of functions on M by

$$S^{k,p} = \{f \in L^p : Df \in L^p \text{ whenever } w(D) \leq k\}.$$

We define the L^p norm of $\nabla_b f, \nabla_b^2 f, \dots$ to be $(\int |\nabla_b f|^p \theta \wedge d\theta)^{1/p}, (\int |\nabla_b^2 f|^p \theta \wedge d\theta)^{1/p}, \dots$, respectively, as usual. So it is natural to define the $S^{k,p}$ norm of $f \in S^{k,p}$ as follows:

$$\|f\|_{S^{k,p}} \equiv \left(\sum_{0 \leq j \leq k} \|\nabla_b^j f\|_{L^p}^p \right)^{1/p}.$$

The function space $S^{k,p}$ with the above norm is a Banach space for $k \geq 0$, $1 < p < \infty$. There are also embedding theorems of Sobolev type. For instance, $S^{2,2} \subset S^{1,4}$ (for $\dim M = 3$). We refer the reader to, for instance, [FS2] and [Fo] for more discussions on these spaces.

REFERENCES

- [A] T. Aubin, *Some Nonlinear Problems in Riemannian Geometry*, Springer-Verlag, Berlin, 1998.
- [B] S. Brendle, Global Existence and Convergence for a Higher-Order Flow in Conformal Geometry, *Ann. Math.*, 158 (2003), 323-343.
- [C] P. T. Chruściel, Semi-Global Existence and Convergence of solutions of the Robinson-Trautman (2-Dimensional Calabi) Equation, *Commun. Math. Phys.* 137 (1991), 289-313.
- [C1] S.- C. Chang, The Q -Curvature Flow on a Closed 3-Manifold of Positive Q -Curvature, preprint.
- [C2] S.- C. Chang, Recent Developments on the Calabi Flow, *Contemporary Mathematics*, 367, 17-42, A.M.S., 2005.
- [C3] S.- C. Chang, Global Existence and Convergence of Solutions of the Calabi Flow on Riemann Surfaces of Genus $g \geq 2$, *J. Math. Kyoto Univ.* Vol. 40, No. 2 (2000), 363-377.
- [C4] S.- C. Chang, The 2-dimensional Calabi Flow, *Nagoya Math. J.*, to appear.
- [CC1] S.- C. Chang and J.- H. Cheng, The Harnack Estimate for the Yamabe Flow on CR Manifolds of Dimension 3, *Annals of Global Analysis and Geometry*, 21 (2002), 111-121.
- [CH] S.- S. Chern and R. Hamilton, On Riemannian Metrics Adapted to Three-dimensional Contact Manifolds, *Lecture Notes in Math.*, 1111, 279-305, Springer-Verlag, 1984.
- [Chi] H.-L. Chiu, The Sharp Lower Bound for the First Positive Eigenvalue of a Sub-Laplacian on a Three-Dimensional Pseudo-hermitian Manifold, submitted.
- [CL] W. S. Cohn and G. Lu, Best Constants for Moser-Trudinger Inequalities on the Heisenberg Group, *Indiana Univ. Math. J.* 50 (2001), 1567-1591.
- [CLe] J.- H. Cheng and J. M. Lee, The Burns-Epstein Invariant and Deformation of CR Structures, *Duke Math. J.*, 60 (1990), 221-254.
- [CS] S.- C. Chen and M.- C. Shaw, *Partial Differential Equations in several complex variables*, *Studies in Advan. Math.*, 19, AMS/IP, 2001.
- [CW] S.- C. Chang and C.- T. Wu, The Fourth-Order Q -Curvature Flow on Closed 3-Manifolds, *Nagoya Math. J.*, to appear.
- [Fa] F. Farris, An Intrinsic Construction of Fefferman's CR Metric, *Pacific J. Math.*, 123 (1986), 33-45.

- [FH] C. Fefferman and K. Hirachi, Ambient Metric Construction of Q -Curvature in Conformal and CR Geometries, to appear in Math. Res. Lett..
- [Fo] G. B. Folland, Subelliptic Estimates and Function Spaces on Nilpotent Lie Groups, Arkiv for Mat. 13 (1975), 161-207.
- [FS1] G. B. Folland and E. M. Stein, *Hardy spaces on homogeneous groups*, Princeton U. Press, 1980.
- [FS2] G. B. Folland and E. M. Stein, Estimates for the $\bar{\partial}_b$ Complex and Analysis on the Heisenberg Group, Comm. Pure Appl. Math., 27 (1974), 429-522.
- [GG] A. R. Gover and C. R. Graham, CR Invariant Powers of the Sub-Laplacian, preprint.
- [GL] C. R. Graham and J. M. Lee, Smooth Solutions of Degenerate Laplacians on Strictly Pseudoconvex Domains, Duke Math. J., 57 (1988), 697-720.
- [H] K. Hirachi, Scalar Pseudo-hermitian Invariants and the Szegő Kernel on 3-dimensional CR Manifolds, Lecture Notes in Pure and Appl. Math. 143, pp. 67-76, Dekker, 1992.
- [JL] D. Jerison and J. M. Lee, The Yamabe Problem on CR Manifolds, J. Diff. Geom. 25 (1987), 167-197.
- [K1] J. J. Kohn, Estimates for $\bar{\partial}_b$ on Compact Pseudoconvex CR Manifolds, Proc. of Symposia in Pure Math., 43 (1985), 207-217.
- [K2] J. J. Kohn, The Range of the Tangential Cauchy-Riemann Operator, Duke. Math. J., 53 (1986) 525-545.
- [L1] J. M. Lee, Pseudo-Einstein Structure on CR Manifolds, Amer. J. Math. 110 (1988), 157-178.
- [L2] J. M. Lee, The Fefferman Metric and Pseudohermitian Invariants, Trans. A.M.S. 296 (1986), 411-429.
- [Le] L. Lempert, On Three-Dimensional Cauchy-Riemann Manifolds. J. of Amer. Math. Soc., 5 (1992), 923-969.
- [P] S. Paneitz, A Quartic Conformally Covariant Differential Operator for Arbitrary Pseudo-Riemannian Manifolds, preprint, 1983.
- [S] L. Simon, Asymptotics for a Class of Nonlinear Evolution Equations, with Applications to Geometric Problems, Ann. of Math., 118 (1983), 525-571.
- [SC] Sánchez-Calle A., Fundamental Solutions and Geometry of the Sum of Squares of Vector Fields, Invent. Math., 78 (1984), 143-160.

¹DEPARTMENT OF MATHEMATICS, NATIONAL TSING HUA UNIVERSITY, HSINCHU 30013, TAIWAN, R.O.C.

E-mail address: `scchang@math.nthu.edu.tw`

²INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, NANKANG, TAIPEI 11529, TAIWAN, R.O.C.

E-mail address: `cheng@math.sinica.edu.tw`

³INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, NANKANG, TAIPEI 11529, TAIWAN, R.O.C.

E-mail address: `hlchiu@math.sinica.edu.tw`