Regularity of the distance function to the boundary

YanYan Li^{*} Department of Mathematics Rutgers University 110 Frelinghuysen Road Piscataway, NJ 08854

> Louis Nirenberg Courant Institute 251 Mercer Street New York, NY 10012

Dedicated to the memory of Luigi Amerio

Abstract

Let Ω be a domain in a smooth complete Finsler manifold, and let G be the largest open subset of Ω such that for every x in G there is a unique closest point from $\partial\Omega$ to x (measured in the Finsler metric). We prove that the distance function from $\partial\Omega$ is in $C_{loc}^{k,\alpha}(G\cup\partial\Omega), k \geq 2$ and $0 < \alpha \leq 1$, if $\partial\Omega$ is in $C^{k,\alpha}_{loc}$.

1 Introduction

In [1] we studied the singular set of viscosity solutions of some Hamilton-Jacobi equations. This was reduced to the study of the singular set of the distance function to the boundary of a domain Ω — for a Finsler metric. The singular set was defined as the complement of the following open set

 $G := \text{ the largest open subset of } \Omega \text{ such that for every } x \text{ in } G \text{ there is a}$ unique closest point from $\partial \Omega$ to x (measured in the Finsler metric). (1)

^{*}Partially supported by NSF grant DMS-0401118.

In [1] we stated that if $\partial\Omega$ is in $C^{k,\alpha}$, $k \geq 2$ and $0 < \alpha \leq 1$, then the distance function from the boundary belongs to $C^{k-1,\alpha}(G \cup \partial\Omega)$. Recently Joel Spruck asked to see the proof for a Riemannian metric and pointed out that the result would imply that the distance function would be in $C^{k,\alpha}(G \cup \partial\Omega)$. In this paper we provide a proof of that in the Finsler case. This paper can be regarded as an addendum to [1].

We present two proofs of the $C^{k,\alpha}$ result. We use the notation, as in [1],

$$\int_0^T \varphi(\xi(t); \dot{\xi}(t)) dt$$

for the length of a curve $\xi(t)$. $\varphi(\xi; v)$ is homogeneous of degree one in v. For fixed ξ , the level surface $\varphi(\xi; v) = 1$ is smooth, closed, strictly convex, with positive principal curvatures.

The first proof uses very little of [1] and is essentially self-contained. The second proof uses some structure from [1], and may be of some interest to some readers.

We actually prove a more general result here, involving conjugate points from the boundary.

Definition. Conjugate Point. Consider a point y on $\partial\Omega$, and consider the geodesic $\xi(y, s)$ from y going inside Ω "normal" to $\partial\Omega$ (explained below) with s as arclength. The conjugate point to y is the first point \bar{x} on the normal geodesic such that any point x'' on the geodesic beyond \bar{x} has, in any neighborhood of the geodesic, a shorter join from $\partial\Omega$ to x'' than our normal geodesic to x''.

Normal. A geodesic Γ from a point $y \in \partial \Omega$ is "normal" to $\partial \Omega$ if for x on Γ close to y, the geodesic is the shortest join from $\partial \Omega$ to x.

To obtain the regularity in G we prove a slightly more general result which is local on $\partial\Omega$. Namely, suppose C is a neighborhood on $\partial\Omega$ of a point y and that the normal geodesic Γ from y to a point X in Ω is the unique shortest join from C to X. If the conjugate point to y is beyond X then there is a neighborhood A of X such that the distance from C to any point in A belongs to $C^{k,\alpha}$. See Theorem 1 below.

We shall make use of special coordinates introduced in section 3 of [1] about a given normal geodesic Γ , going from a point $y \in \partial \Omega$ into Ω . In these coordinates y is the origin and the x_n -axis is normal to $\partial \Omega$ there and is the geodesic Γ . Furthermore, in these coordinates, φ has the following properties, see (4.1)-(4.6) in [1]. Here Greek letters α, β range from 1 to n - 1, and Latin letters i, j range from 1 to n.

$$\varphi(te_n; e_n) \equiv 1, \tag{2}$$

$$\varphi_{\xi^j}(te_n; e_n) \equiv 0, \tag{3}$$

$$\varphi_{v^{\alpha}}(te_n; e_n) \equiv 0, \qquad \varphi_{v^n}(te_n; e_n) \equiv 1,$$
(4)

$$\varphi_{\xi^{j}v^{k}}(te_{n};e_{n}) \equiv 0, \tag{5}$$

$$\varphi_{v^j v^n}(te_n; e_n) \equiv 0, \tag{6}$$

$$\varphi_{\xi^j\xi^n}(te_n;e_n) \equiv 0. \tag{7}$$

In these coordinates for $y \in \partial \Omega$ near the origin the geodesic $\xi(y, s)$ from y "normal" to $\partial \Omega$ there satisfies

$$\dot{\xi}(y,0) = V(y)$$

where V(y) is the unique vector-valued function on $\partial\Omega$ satisfying (here $\nu(y)$ is the Euclidean interior unit normal to $\partial\Omega$ at y)

$$\begin{cases} V(y) \cdot \nu(y) > 0\\ \varphi(y; V(y)) = 1\\ \nabla_v \varphi(y; V(y)) \text{ is parallel to } \nu(y). \end{cases}$$
(8)

Using these special coordinates, near the origin, $\partial \Omega$ has the form

$$y = (x', f(x')), \quad f(0') = 0, \quad \nabla f(0') = 0'.$$
 (9)

We assume that $f \in C^{k,\alpha}, k \ge 2, 0 < \alpha \le 1$.

The result we prove is

Theorem 1 Assume that the conjugate point of the origin on the geodesic $\Gamma = \{te_n\}$ is beyond e_n , and that there exists a neighborhood C of 0' on $\partial\Omega$ such that $\{te_n \mid 0 \leq t \leq 1\}$ is the unique shortest geodesic from C to e_n . Then there exist neighborhoods A of e_n and A of 0' on $\partial\Omega$ such that for any X in A there is a unique $y \in A$ and geodesic from y to X which is the shortest join from A to A. Furthermore, if d(X) is its length, then the Jacobian of the map $X \to (d, y)$ is nonsingular at e_n , and d lies in $C_{loc}^{k,\alpha}$ in A, and y lies in $C_{loc}^{k-1,\alpha}$ in A.

2 Second Variation

Consider one parameter family of curves $\tau(\epsilon, t)$ from \mathcal{A} to $\bar{t}e_n$, $\bar{t} > 0$, with $\tau(0, t) = te_n$. We look at the second variation of its length $I[\tau(\epsilon, \cdot)]$. For \bar{t} small, it is clearly positive definite. The first \hat{t} for which it fails to be strictly positive definite is the conjugate point. For if $\tilde{t} = \hat{t} + \delta$, $\delta > 0$, then the second variation of curves to \tilde{t} cannot be semipositive definite, and there would then be a shorter connection from \mathcal{A} to $\tilde{t}e_n$ near Γ . The standard computation of second variation yields

$$\frac{d^2}{d\epsilon^2} I[\tau(\epsilon,\cdot)]\Big|_{\epsilon=0} = J(\tau_\epsilon|_{\epsilon=0}) - f_{x_\alpha x_\beta}(0')\tau_\epsilon^\alpha(0,0)\tau_\epsilon^\beta(0,0).$$

Here J is the usual expression of the second variation if the bottom point were kept at the origin. Namely,

$$J(\tau_{\epsilon}|_{\epsilon=0}) = \int_0^{\bar{t}} \left\{ \varphi_{\xi^{\alpha}\xi^{\beta}}(te_n; e_n) \tau_{\epsilon}^{\alpha}(0, t) \tau_{\epsilon}^{\beta}(0, t) + \varphi_{v^{\alpha}v^{\beta}}(te_n; e_n) \dot{\tau}_{\epsilon}^{\alpha}(0, t) \dot{\tau}_{\epsilon}^{\beta}(0, t) \right\} dt.$$
(10)

Note that τ_{ϵ}^{n} and $\dot{\tau}_{\epsilon}^{n}$ do not occur in J.

3 First proof of Theorem 1

3.1

Recalling (9) we shall denote the normal geodesic from y = (x', f(x')) by $X = \xi(x', s)$; this is a slight change of notation. The geodesic ξ and ξ_s depend smoothly on s and their initial data, while the initial data depend $C^{k-1,\alpha}$ on x'. To prove the theorem, it suffices to show that the Jacobian of the mapping (x', s) to X at (0', 1) is nonsingular. It follows that dand y belong to $C^{k-1,\alpha}$. Since $\nabla_X d = X_s$, it follows that $\nabla_X d$ is in $C^{k-1,\alpha}$ and hence d is in $C^{k,\alpha}$ — as Spruck pointed out to us.

We now prove the Jacobian is nonsingular.

Write $X = (X', X^n)$. Since $X_s(0', 1) = (0', 1)$, the Jacobian of the mapping (x', s) to X at (0', 1) is simply

$$M := \frac{\partial X'}{\partial x'}(0', 1).$$

Assume M is singular, without loss of generality we may suppose that

$$X'_{x_1}(0',1) = 0'. (11)$$

We construct a perturbation $\tau(\epsilon, t)$ of $\Gamma = \{te_n \mid 0 \le t \le 1\}$ such that $\zeta(t) := \tau_{\epsilon}|_{\epsilon=0}$ satisfies

$$J[\zeta] = f_{x_{\alpha}x_{\beta}}(0')\zeta^{\alpha}(0)\zeta^{\beta}(0).$$
(12)

3.2

Consider the geodesic $\xi(\delta e_1, t)$ of length 1 starting at $(\delta e_1, f(\delta e_1))$, $0 < \delta$ small and "normal" to $\partial \Omega$ there. Set

$$\zeta(t) = \frac{\partial}{\partial \delta} \xi(\delta e_1, t) \Big|_{\delta=0}.$$
(13)

By (11),

$$\zeta(1) = 0. \tag{14}$$

We obtain an equation for $\zeta(t)$ by differentiating the geodesic equation

$$\varphi_{\xi^i} = \frac{d}{dt} \varphi_{v^i}(\xi; \dot{\xi})$$

with respect to δ , and setting $\delta = 0$. We find

$$\varphi_{\xi^i\xi^j}(te_n;e_n)\zeta^j = \frac{d}{dt} \left(\varphi_{v^iv^j}(te_n;e_n)\dot{\zeta}^j\right).$$

Here we have used property (5) of our special coordinates. By (7) and (6),

$$\varphi_{\xi^{\alpha}\xi^{\beta}}\zeta^{\beta} = \frac{d}{dt} \left(\varphi_{v^{\alpha}v^{\beta}}\dot{\zeta}^{\beta}\right). \tag{15}$$

We have

$$\zeta^{\alpha}(0) = \delta_1^{\alpha}. \tag{16}$$

In addition,

$$\dot{\zeta}(0) = \frac{\partial}{\partial\delta} \dot{\xi}(\delta e_1, 0) \Big|_{\delta=0} = \frac{\partial}{\partial\delta} V(\delta e_1) \Big|_{\delta=0} = V_{x_1}(0').$$
(17)

By the last formula in (8) we have

$$\nabla_v \varphi \left((\delta e_1, f(\delta e_1)); V(\delta e_1) \right) \cdot (e_1 + f_{x_1}(\delta e_1) e_n) = 0.$$

Differentiating in δ and setting $\delta = 0$, we find, using properties of our special coordinates,

$$\varphi_{v^1 v^\beta}(0'; e_n) V_{x_1}^\beta(0') + f_{x_1 x_1}(0') = 0.$$
(18)

Now we introduce the perturbation $\tau(\epsilon, t)$ as follows

$$\begin{aligned} \tau^{\alpha}(\epsilon,t) &= \epsilon \zeta^{\alpha}(t), \\ \tau^{n}(\epsilon,t) &= te_{n} + (1-t)f(\epsilon e_{1}). \end{aligned}$$

The definition of τ^n is just to ensure that $\tau(\epsilon, 0)$ lies on $\partial\Omega$.

According to (10),

$$J[\tau_{\epsilon}|_{\epsilon=0}] = \int_0^1 \left\{ \varphi_{\xi^{\alpha}\xi^{\beta}}(te_n; e_n) \zeta^{\alpha} \zeta^{\beta} + \varphi_{v^{\alpha}v^{\beta}}(te_n; e_n) \dot{\zeta}^{\alpha} \dot{\zeta}^{\beta} \right\} dt.$$

Integrating the last expression by parts we find, using (15), (16), (17) and (18),

$$J[\tau_{\epsilon}|_{\epsilon=0}] = \int_{0}^{1} \left\{ \varphi_{\xi^{\alpha}\xi^{\beta}} \zeta^{\alpha} \zeta^{\beta} - \zeta^{\alpha} \frac{d}{dt} \left(\varphi_{v^{\alpha}v^{\beta}} \dot{\zeta}^{\beta} \right) \right\} - \zeta^{\alpha}(0) \varphi_{v^{\alpha}v^{\beta}}(0'; e_{n}) \dot{\zeta}^{\beta}(0)$$
$$= -\varphi_{v^{1}v^{\beta}}(0'; e_{n}) V_{x_{1}}^{\beta}(0') = f_{x_{1}x_{1}}(0') = f_{x_{\alpha}x_{\beta}}(0') \tau_{\epsilon}^{\alpha}(0, 0) \tau_{\epsilon}^{\beta}(0, 0).$$

It follows from Section 2 that the second variation is zero.

4 Second proof of Theorem 1

Second proof of Theorem 1. For X near e_n and for small $\sigma' = (\sigma_1, \dots, \sigma_{n-1}) \in \mathbb{R}^{n-1}$, let $\tau = \tau(\sigma', X)$ be defined by $\varphi(X; (\sigma', \tau)) = 1$ and $\tau(0', e_n) = 1$. Since $\varphi_{v^n}(e_n; e_n) = 1$, by the implicit function theorem, τ exists as a smooth function of (σ', X) near $(0', e_n)$.

Let, as on page 111 of [1], $\eta = \eta(\sigma', X, t)$ be the unique smooth function of, with $\psi = \varphi^2$,

$$\psi_{\xi^i}(\eta;\dot{\eta}) = \frac{d}{dt}\psi_{v^i}(\eta;\dot{\eta}), \qquad t \le 1,$$

satisfying

$$\eta(\sigma',X,1)=X,\qquad \dot{\eta}(\sigma',X,1)=(\sigma',\tau(\sigma',X)).$$

As explained in the last two lines of page 108 in [1], $\eta(\sigma', X, t)$ is a geodesic with t the arclength.

In the special coordinates described in Section 1, $\partial\Omega$ has the form (9) near the origin with $f \in C^{k,\alpha}$, $k \ge 2$, $0 < \alpha \le 1$. Since $\{te_n \mid 0 \le t \le 1\}$ is the unique shortest geodesic from C to e_n , we know that for X close to e_n , there exists x' close to 0' such that the "normal geodesic" starting from (x', f(x')) will reach X as a shortest join from C to X. It follows that for some σ' close to 0' and t close to 0, we have

$$\begin{cases} \eta(\sigma', X, t) - (x', f(x')) = 0, \\ \dot{\eta}^{\mu}(\sigma', X, t) - V^{\mu}(x') = 0, \end{cases}$$
(19)

where V(x') := V(x', f(x')) satisfies (8). Note that 1 - t is the distance from C to X.

To prove Theorem 1, we only need to show that the left hand side of (19), denoted as LHS, has nonsingular Jacobian $\frac{\partial(LHS)}{\partial(t,\sigma',x')}$ at $(t,\sigma',x',X) = (0,0',0',e_n)$. Indeed, this would allow the use of the implicit function theorem to show that for X close to e_n and in a neighborhood of (0,0',0'), there exists a unique $C^{k-1,\alpha}$ solution $(t,\sigma',x') =$ $(t(X),\sigma'(X),x'(X))$ of (19). Thus, Theorem 1 follows as explained at the beginning of Section 3.

Clearly,

$$\frac{\partial(LHS)}{\partial t}(0,0',0',e_n) = \begin{pmatrix} \dot{\eta}(0',e_n,0)\\ (\ddot{\eta}^{\mu}(0',e_n,0)) \end{pmatrix} = \begin{pmatrix} 0'\\ 1\\ 0' \end{pmatrix}$$

a $(2n-1) \times 1$ column vector,

$$\frac{\partial(LHS)}{\partial\sigma'}(0,0',0',e_n) = \begin{pmatrix} (\eta_{\sigma_{\alpha}}(0',e_n,0))\\ (\dot{\eta}^{\mu}_{\sigma_{\alpha}}(0',e_n,0)) \end{pmatrix} = \begin{pmatrix} (\eta^{\mu}_{\sigma_{\alpha}}(0',e_n,0))\\ (\eta^{n}_{\sigma_{\alpha}}(0',e_n,0))\\ (\dot{\eta}^{\mu}_{\sigma_{\alpha}}(0',e_n,0)) \end{pmatrix},$$
(20)

a $(2n-1) \times (n-1)$ matrix,

$$\frac{\partial(LHS)}{\partial x'}(0,0',0',e_n) = \begin{pmatrix} -I\\ 0\\ -\nabla V'(0') \end{pmatrix},$$
(21)

a $(2n-1) \times (n-1)$ matrix, where I is the $(n-1) \times (n-1)$ identity matrix and $\nabla V' := (V^{\mu}_{x_{\beta}})$. Thus

$$det\left(\frac{\partial(LHS)}{\partial(t,\sigma',x')}(0,0',0',e_n)\right) = (-1)^{n-1}det\left(\begin{array}{cc}\frac{\partial\eta'}{\partial\sigma'}(0',e_n,0) & -I\\\frac{\partial\eta'}{\partial\sigma'}(0',e_n,0) & -\nabla V'(0')\end{array}\right),\tag{22}$$

where $\frac{\partial \eta'}{\partial \sigma'} := (\eta^{\mu}_{\sigma_{\alpha}})$ and $\frac{\partial \dot{\eta}'}{\partial \sigma'} := (\dot{\eta}^{\mu}_{\sigma_{\alpha}})$. By the last line in (8),

$$\nabla_v \varphi((x', f(x')); V(x'))[e_\beta + f_{x_\beta}(x')e_n] = 0,$$

i.e.

$$\varphi_{v^{\beta}}((x', f(x')); V(x')) + \varphi_{v^{n}}((x', f(x')); V(x')) f_{x_{\beta}}(x') = 0.$$

Differentiating in x_{α} and setting x' = 0' we find, using properties of our special coordinates (4), (5) and (6),

$$D_{v'}^2 \varphi(0'; e_n) \cdot \nabla V'(0') + D^2 f(0') = 0, \qquad (23)$$

where $D_{v'}^2 \varphi := (\varphi_{v^\beta v^\mu})$. We now evaluate $\frac{\partial \eta'}{\partial \sigma'}(0', e_n, 0)$ and $\frac{\partial \eta'}{\partial \sigma'}(0', e_n, 0)$. It is proved in section 4.4 of [1] that there exists a $C^{2,1}$ function \tilde{f} near 0' satisfying

$$\tilde{f}(0') = 0, \qquad \nabla \tilde{f}(0') = 0',$$

$$\left(D^2 \tilde{f}(0') - D^2 f(0') \right) > 0,$$
(24)

$$\eta(\sigma', e_n, 0) = (y', \tilde{f}(y')), \tag{25}$$

$$\dot{\eta}^{\mu}(\sigma', e_n, 0) = \tilde{V}^{\mu}(y'), \qquad (26)$$

where $\tilde{V}(y') := \tilde{V}((y', \tilde{f}(y')))$ is determined by (8) with f replaced by \tilde{f} , and $y' = y'(\sigma')$ satisfies

$$det\left(\frac{\partial y'}{\partial \sigma'}(0')\right) \neq 0.$$
(27)

Note that (27) is given by (4.9) in [1], while (24) follows from corollary 4.15 in [1] together with the fact that e_n is not a conjugate point.

Differentiating (26) in σ_{α} and setting $\sigma' = 0'$ we find

$$\dot{\eta}^{\mu}_{\sigma_{\alpha}}(0', e_n, 0) = \tilde{V}^{\mu}_{y_{\beta}}(0') \frac{\partial y_{\beta}}{\partial \sigma_{\alpha}}(0'),$$

i.e.

$$\frac{\partial \dot{\eta}'}{\partial \sigma'}(0', e_n, 0) = \nabla \tilde{V}'(0') \frac{\partial y'}{\partial \sigma'}(0').$$
(28)

Differentiating (25) in σ_{α} and setting $\sigma' = 0'$ we find

$$\frac{\partial \eta'}{\partial \sigma'}(0', e_n, 0) = \frac{\partial y'}{\partial \sigma'}(0').$$
(29)

Since

$$\begin{pmatrix} \frac{\partial y'}{\partial \sigma'}(0') & -I \\ \nabla \tilde{V}'(0') \frac{\partial y'}{\partial \sigma'}(0') & -\nabla V'(0') \end{pmatrix} = \begin{pmatrix} I & -I \\ \nabla \tilde{V}'(0') & -\nabla V'(0') \end{pmatrix} \begin{pmatrix} \frac{\partial y'}{\partial \sigma'}(0') & I \end{pmatrix},$$

we have, by putting (28) and (29) into (22),

$$\det\left(\frac{\partial(LHS)}{\partial(t,\sigma',x')}(0,0',0',e_n)\right) = (-1)^{n-1}\det\left(\frac{\partial y'}{\partial\sigma'}(0')\right)\det\left(\begin{array}{cc}I & -I\\\nabla\tilde{V}'(0') & -\nabla V'(0')\end{array}\right).$$
(30)

The proof of (23), applied to \tilde{f} instead of f, yields

$$D_{v'}^2 \varphi(0'; e_n) \nabla \tilde{V}'(0') + D^2 \tilde{f}(0') = 0.$$
(31)

Thus, by (23) and (31),

$$\begin{pmatrix} I & -I \\ -D^2 \tilde{f}(0') & D^2 f(0') \end{pmatrix} = \begin{pmatrix} I & -I \\ D_{v'}^2 \varphi(0'; e_n) \nabla \tilde{V}'(0') & -D_{v'}^2 \varphi(0'; e_n) \nabla V'(0') \end{pmatrix}$$
$$= \begin{pmatrix} I \\ D_{v'}^2 \varphi(0'; e_n) \end{pmatrix} \begin{pmatrix} I & -I \\ \nabla \tilde{V}'(0') & -\nabla V'(0') \end{pmatrix},$$

and therefore

$$\det \left(D^2 f(0') - D^2 \tilde{f}(0') \right) = \det D^2_{v'} \varphi(0'; e_n) \det \left(\begin{array}{c} I & -I \\ \nabla \tilde{V}'(0') & -\nabla V'(0') \end{array} \right).$$
(32)

Since $D_{v'}^2 \varphi(0'; e_n)$ is positive definite, we deduce from (30), (27) and (32) that

$$\det\left(\frac{\partial(LHS)}{\partial(t,\sigma',x')}(0,0',0',e_n)\right) \neq 0.$$

	Г		
	L		1

References

 Y.Y. Li and L. Nirenberg, The distance function to the boundary, Finsler geometry and the singular set of viscosity solutions of some Hamilton-Jacobi equations, Comm. Pure Appl. Math. 58 (2005), 85-146.