

The Chow rings of generalized Grassmannians

Haibao Duan and Xuezhi Zhao

Institute of Mathematics, Chinese Academy of Sciences,
dhd@math.ac.cn

Department of Mathematics, Capital Normal University
zhaoxve@mail.cnu.edu.cn

February 8, 2020

Abstract

08

2000 Mathematical Subject Classification: 14M15; 57T15.

1 Introduction

Let G be a compact connected Lie group and $H \subset G$ a closed subgroup. The space G/H of left cosets of H in G is called a *homogeneous space*. In the special case where H is the centralizer of a 1-parameter subgroup, the G/H is a smooth, projective complex algebraic variety, known as a *flag variety*.

A classical problem in algebraic geometry (resp. topology) is to characterize the Chow ring $A^*(G/H)$ of a flag variety (resp. the integral cohomology $H^*(G/H)$ of a homogeneous space) by a minimal system of generators and relations. The traditional method dealing with this problem is due to A. Borel [B₁, B, HMS, T, Wo]. It utilizes Leray spectral sequence in which the topology of Lie groups is requested at the beginning by the E_2 -terms. This approach is effective when $H^*(G)$ is torsion free. However, efforts to apply it to the remaining cases have encountered considerable computational difficulties, in particular, when G is one of the five exceptional Lie groups [I, IT, T, TW, W₁, W₂, N] (cf. §7).

We introduce a new method for calculating the Chow ring of flag varieties (resp. integral cohomology of homogeneous spaces). This is based on two fundamental results from Schubert's enumerative calculus [Sch; BGG]. The first is the Basis Theorem due to Bruhat–Chevalley [C] stating that the classical *Schubert classes* on a flag variety G/H constitute an additive basis for the Chow ring $A^*(G/H)$; while the second is the formula obtained in [D₂] for multiplying Schubert classes. Since these two results have all been programed from the Cartan matrix of G in [DZ₁], our approach boils down

the problem directly to such primary and well known invariants of Lie groups as *Cartan numbers* and, therefore, is self-contained in the sense that no knowledge on the topology of Lie groups is assumed.

As an initial step of this project we restrict ourself to a family of flag varieties (resp. homogeneous spaces) that are of classical interest. Let G be a compact connected semi-simple Lie group with Lie algebra $L(G)$, exponential map $\exp : L(G) \rightarrow G$ and a fixed maximal torus T in G . Let $\Omega = \{\omega_1, \dots, \omega_n\} \subset L(T)$ be the set of *fundamental dominant weights* of G relative to a system Φ of simple roots of G (cf. **2.1**). For a $\omega \in \Omega$ the centralizer of the 1-dimensional subgroup $\{\exp(t\omega) \mid t \in \mathbb{R}\}$ in G , denoted by H , is called the *parabolic subgroup of G* corresponding to ω . Let H_s be the semi-simple part of H .

Definition 1. The flag variety G/H is called *the Grassmannian of G associated to the weight ω* and G/H_s , a rank 1 homogeneous space of G .

Grassmannians (resp. rank 1 homogeneous spaces) are many. To see this we recall that, up to local isomorphisms, all compact connected semi-simple Lie groups fall into four infinite sequences of classical groups

- $A_n = SU(n)$: the special unitary group of order n ;
- $D_n = Spin(2n)$: the spinor group of order $2n$;
- $B_n = Spin(2n + 1)$: the spinor group of order $2n + 1$;
- $C_n = Sp(n)$: the symplectic group of order n

as well as the five exceptional ones: G_2, F_4, E_6, E_7, E_8 . Assume that, if G is one of these groups, a set $\Omega = \{\omega_1, \dots, \omega_n\}$ of fundamental dominant weights of G is given and ordered by the root-vertices in the Dynkin diagram of G in [Hu, p.58]. With this convention we tabulate, for given G and ω , some parabolic H indicated by its semi-simple part H_s .

G	$SU(n)$	$Spin(2n)$	F_4	F_4	E_6	E_6	E_7	E_7
ω	ω_k	ω_2	ω_1	ω_4	ω_2	ω_6	ω_1	ω_7 .
H_s	$A_k \times A_{n-k}$	A_n	C_3	B_3	A_6	D_5	D_6	E_6

In the first two cases, the G/H correspond respectively to *the Grassmannian $G_{n,k}(\mathbb{C})$* of k -planes through the origin in the complex n -space \mathbb{C}^n , and *the Grassmannian $\mathbb{C}S_n$* of complex structures on the $2n$ -Euclidean space \mathbb{R}^{2n} [D₁,DP]. These originate the notion *Grassmannian* in Definition 1.

As application of the method, calculation is performed for 6 cases among 27 exceptional Grassmannians (results for the remaining cases can be derived analogously). The general strategy is to select in the set of all Schubert classes on G/H a minimal subset, whose elements may be termed as the *special Schubert classes* on G/H , so that the ring $A^*(G/H)$ admits a presentation as a quotient of the polynomial ring in the special Schubert

classes. More precisely, granted with the *Weyl coordinates* for Schubert classes introduced in **2.2**, the following results are established.

Theorem 1. $A^*(F_4/C_3 \cdot S^1) = \mathbb{Z}[y_1, y_3, y_4, y_6]/\langle r_3, r_6, r_8, r_{12} \rangle$, where y_1, y_3, y_4, y_6 are the Schubert classes specified by their Weyl coordinates

$$\sigma[1], \sigma[3, 2, 1], \sigma[4, 3, 2, 1], \sigma[3, 2, 4, 3, 2, 1]$$

respectively, and where

$$\begin{aligned} r_3 &= 2y_3 - y_1^3; \\ r_6 &= 2y_6 + y_3^2 - 3y_1^2y_4; \\ r_8 &= 3y_4^2 - y_1^2y_6; \\ r_{12} &= y_6^2 - y_4^3. \end{aligned}$$

Theorem 2. $A^*(F_4/B_3 \cdot S^1) = \mathbb{Z}[y_1, y_4]/\langle r_8, r_{12} \rangle$, where y_1, y_4 are the Schubert classes specified by their Weyl coordinates

$$\sigma[4], \sigma[3, 2, 3, 4]$$

respectively, and where

$$r_8 = 3y_4^2 - y_1^8; \quad r_{12} = 26y_4^3 - 5y_1^{12}.$$

Theorem 3. $A^*(E_6/A_6 \cdot S^1) = \mathbb{Z}[y_1, y_3, y_4, y_6]/\langle r_6, r_8, r_9, r_{12} \rangle$, where y_1, y_3, y_4, y_6 are the Schubert classes specified by their Weyl coordinates

$$\sigma[2], \sigma[3, 4, 2], \sigma[1, 3, 4, 2], \sigma[1, 3, 6, 5, 4, 2]$$

respectively, and where

$$\begin{aligned} r_6 &= 2y_6 + y_3^2 - 3y_1^2y_4 + 2y_1^3y_3 - y_1^6; \\ r_8 &= 3y_4^2 - 6y_1y_3y_4 + y_1^2y_6 + 5y_1^2y_3^2 - 2y_1^5y_3; \\ r_9 &= 2y_3y_6 - y_1^3y_6; \\ r_{12} &= y_4^3 - y_6^2. \end{aligned}$$

Theorem 4. $A^*(E_6/D_5 \cdot S^1) = \mathbb{Z}[y_1, y_4]/\langle r_9, r_{12} \rangle$, where y_1, y_4 are the Schubert classes specified by their Weyl coordinates

$$\sigma[6], \sigma[2, 4, 5, 6],$$

respectively, and where

$$\begin{aligned} r_9 &= 2y_1^9 + 3y_1y_4^2 - 6y_1^5y_4; \\ r_{12} &= y_4^3 - 6y_1^4y_4^2 + y_1^{12}. \end{aligned}$$

Theorem 5. $A^*(E_7/E_6 \cdot S^1) = \mathbb{Z}[y_1, y_5, y_9]/\langle r_{10}, r_{14}, r_{18} \rangle$, where y_1, y_5, y_9 are the Schubert classes specified by their Weyl coordinates

$$\sigma[7], \sigma[2, 4, 5, 6, 7], \sigma[1, 5, 4, 2, 3, 4, 5, 6, 7]$$

respectively, and where

$$\begin{aligned} r_{10} &= y_5^2 - 2y_1y_9; \\ r_{14} &= 2y_5y_9 - 9y_1^4y_5^2 + 6y_1^9y_5 - y_1^{14}; \\ r_{18} &= y_9^2 + 10y_1^3y_5^3 - 9y_1^8y_5^2 + 2y_1^{13}y_5. \end{aligned}$$

Theorem 6. $A^*(E_7/D_6 \cdot S^1) = \mathbb{Z}[y_1, y_4, y_6, y_9]/\langle r_9, r_{12}, r_{14}, r_{18} \rangle$, where y_1, y_4, y_6, y_9 are the Schubert classes specified by their Weyl coordinates

$$\sigma[1], \sigma[2, 4, 3, 1], \sigma[2, 6, 5, 4, 3, 1], \sigma[3, 4, 2, 7, 6, 5, 4, 3, 1],$$

respectively, and where

$$\begin{aligned} r_9 &= 2y_9 + 3y_1y_4^2 + 4y_1^3y_6 + 2y_1^5y_4 - 2y_1^9; \\ r_{12} &= 3y_6^2 - y_4^3 - 3y_1^4y_4^2 - 2y_1^6y_6 + 2y_1^8y_4; \\ r_{14} &= 3y_4^2y_6 + 3y_1^2y_6^2 + 6y_1^2y_4^3 + 6y_1^4y_4y_6 + 2y_1^5y_9 - y_1^{14}; \\ r_{18} &= 5y_9^2 + 29y_6^3 - 24y_1^6y_6^2 + 45y_1^2y_4y_6^2 + 2y_1^9y_9. \end{aligned}$$

Theorem 1-6 can be interpreted as integral cohomology of the corresponding Grassmannian for, by a classical result of Chow, the $H^*(G/H)$ is canonically isomorphic to $A^*(G/H)$. Moreover, by presenting the ring in terms of Schubert classes, Theorem 1-6 can be interpreted as the *Schubert presentations* of the ring $A^*(G/H)$, hence are directly applicable to the intersection theory on G/H (cf. [IM], [DP, Section 9.6]).

Traditionally, Schubert calculus deals with intersection theory on flag varieties. Statements and proofs of Theorem 7-12 in §5 illustrate how this calculation is extended to yield the integral cohomology of homogeneous spaces such as the G/H_s .

The paper is arranged as follows. §2 contains a brief introduction to what we need from Schubert calculus. §3 develops two results concerning computing with ideals in a polynomial ring. By resorting to the Gysin sequence of the fibration $G/H_s \rightarrow G/H$, relationship between cohomologies of a Grassmannian G/H and its allied space G/H_s is discussed in §4.

Certain theoretical notation and results in §2–§4 are algorithmic in nature. Their effective computability is emphasized by referring to appropriate sections of [DZ₂], which serves also the purpose to tabulated the intermediate data used in establishing Theorem 1–12 in §5 and §6 (The data plays no further role once Theorem 1-12 are established. So we choose to exclude them from the main body of the paper for the sake of simplicity).

Certain cases of the homogeneous spaces concerned in this paper have previously been investigated by many authors. Comparisons between our results and those archived by classical means are made in §7, where a mistake occurring in earlier computation is corrected.

2 Elements of Schubert calculus

Assume throughout that the Lie group G under consideration is compact and 1-connected. Fixed a maximal torus T in G and equip the Lie algebra $L(G)$ with an inner product $(,)$, so that the adjoint representation acts as isometries of $L(G)$. Let $\Phi = \{\beta_1, \dots, \beta_n\} \subset L(T)$ be a *set of simple roots* of G [Hu, p.47] (which is so ordered as the root-vertices in the Dynkin diagram given in [Hu, p.58] when G is one of the semi-simple Lie groups). The *Cartan matrix* of G is the $n \times n$ integral matrix $C = (c_{ij})_{n \times n}$, where the c_{ij} is the *Cartan integer* defined by

$$\beta_i \circ \beta_j =: 2(\beta_i, \beta_j) / (\beta_j, \beta_j), \quad 1 \leq i, j \leq n \text{ [Hu, p.55].}$$

We recall two algorithms “*Decomposition*” and “*L-R coefficients*” developed from the Cartan matrix in [DZ₁]. The first presents the Weyl group of G by minimized decompositions of its elements from which the Schubert varieties on the flag variety G/H can be constructed. The second expands a polynomial in the Schubert classes as a linear combination of Schubert classes. Both algorithms play a fundamental role throughout the paper.

2.1. Preliminaries in Weyl group. Since $\Phi = \{\beta_1, \dots, \beta_n\}$ is a basis for the vector space $L(T)$, we may introduce another basis $\Omega = \{\omega_1, \dots, \omega_n\}$ of $L(T)$ by the rule

$$(\omega_i, \beta_j) / (\beta_j, \beta_j) = \delta_{i,j}, \quad 1 \leq i, j \leq n.$$

The ω_i is known as the i^{th} *fundamental dominant weight* relative to Φ [Hu, p.67]. With respect to the basis Ω the entries of the Cartan matrix C gives rise to n isometries of the Euclidean space $L(T)$ by

$$\sigma_i(\omega_k) = \begin{cases} \omega_i & \text{if } k \neq i; \\ \omega_i - \sum_{1 \leq j \leq n} c_{ij} \omega_j & \text{if } k = i, \end{cases} \quad 1 \leq i \leq n.$$

Geometrically, σ_i is the reflection in the hyperplane L_i perpendicular to β_i and through the origin.

Definition 2. The subgroup $W(G) \subset \text{Aut}(L(T))$ generated by $\sigma_i, 1 \leq i \leq n$, is called the *Weyl group* of G .

By Definition 2, any $w \in W(G)$ admits a factorization of the form

$$(2.1) \quad w = \sigma_{i_1} \circ \dots \circ \sigma_{i_r}, \quad 1 \leq i_1, \dots, i_r \leq n.$$

The *length* $l(w)$ of a $w \in W(G)$ is the least number of factors in all decompositions of w in the form (2.1). The decomposition (2.1) is said *reduced*, written by $w =: \sigma[i_1, \dots, i_r]$, if $r = l(w)$.

The reduced decompositions of a $w \in W(G)$ may not be unique. However, this ambiguity can be eliminated by employing the following notion. For a $w \in W(G)$ with $l(w) = r$, consider the set of all reduced decompositions of w

$$D(w) = \{I = (i_1, \dots, i_r) \mid w = \sigma[I]\}.$$

It can be ordered by $I = (i_1, \dots, i_r) < J = (j_1, \dots, j_r)$ if there exists $s \leq r$ such that $i_t = j_t$ for all $t < s$ but $i_s < j_s$.

Definition 3. If $I \in D(w)$ is minimum with respect to the order \leq , the decomposition $w = \sigma[I]$ is called the *minimized decomposition* of w .

Clearly one has

Corollary 1. *Every $w \in W(G)$ admits a unique minimized decomposition.*

For a subset $K \subset [1, \dots, n]$ let $H_K \subset G$ be the centralizer of the one-parameter subgroup $\{\exp(tb) \in G \mid t \in \mathbb{R}\}$, $b = \sum_{i \in K} \omega_i$. Its Weyl group $W(H_K)$ is then the subgroup of $W(G)$ generated by $\{\sigma_j \mid j \notin K\}$. Resorting to the length function l on $W(G)$ one may embed the set $W(H_K; G)$ of left cosets of $W(H_K)$ in $W(G)$ as the subset of $W(G)$ (cf. [BGG, 5.1])

$$(2.2) \quad W(H_K; G) = \{w \in W(G) \mid l(w_1) \geq l(w), w_1 \in wW(H_K)\}.$$

We put $W^r(H_K; G) = \{w \in W(H_K; G) \mid l(w) = r\}$.

According to Corollary 1, every $w \in W^r(H_K; G)$ admits a unique minimized decomposition as $w = \sigma[I]$. As a result, the $W^r(H_K; G)$ becomes an ordered set with the order specified by $\sigma[I] < \sigma[J]$ if $I < J$ and therefore, can be presented as

$$(2.3) \quad W^r(H_K; G) = \{w_{r,i} \mid 1 \leq i \leq \beta(r)\}, \quad \beta(r) =: |W^r(H_K; G)|$$

where $w_{r,i}$ is the i^{th} element with respect to the order on $W^r(H_K; G)$.

In [DZ₁] a program entitled “*Decomposition*” has been composed, whose function is summarized below:

Algorithm: *Decomposition.*

Input: *The Cartan matrix $C = (c_{ij})_{n \times n}$ of G , and a subset $K \subset [1, \dots, n]$.*

Output: *The set $W(H_K; G)$ being presented by the minimized decompositions for all its elements, together with the index system (2.3) imposed by the decompositions.*

Example 1. For those $H \subset G$ concerned by Theorem 1-6, the corresponding results coming from the *Decomposition* are tabulated in [DZ₂, 1.1–6.1]. These will be used in the proofs of Theorem 1–12.

2.2. Schubert varieties and Basis Theorem. While studying the geometry of a flag variety G/H we may assume that the subgroup H is of the form H_K for some $K \subset [1, \dots, n]$, since the centralizer of any one-parameter subgroup is conjugate in G to one of the H_K (cf. [BH, 13.5-13.6]).

For a simple root $\beta_i \in \Phi$ let $L_i \subset L(T)$ be the hyperplane perpendicular to β_i and through the origin, and let $K_i \subset G$ be the centralizer of $\exp(L_i)$. For a $w \in W(H; G)$ with the minimized decomposition $w = \sigma[i_1, \dots, i_r]$, write X_w for the image of the map

$$(2.4) \quad K_{i_1} \times \dots \times K_{i_r} \rightarrow G \xrightarrow{p} G/H \quad \text{by } (k_1, \dots, k_r) \rightarrow p(k_1 \cdots k_r),$$

where p is the obvious projection, and where the product \cdot takes place in G . The next result is essentially due to Chevalley [C], except that our description for the X_w follows from Hansen [H], Bott and Samelson [BS]:

Lemma 2. 1) *The subspace $X_w \subset G/H$ is a subvariety with $\dim X_w = 2l(w)$ (known as the Schubert variety in G/H associated to $w \in W(H; G)$).*
 2) *The union $\cup_{w \in W(H; G)} X_w$ dominates G/H by a cell complex.*

Since only even dimensional cells are involved in the decomposition $G/H = \cup_{w \in W(H; G)} X_w$, we may introduce *Schubert class* $s_w \in A^{2l(w)}(G/H)$ as the cocycle class Kronecker dual to the fundamental classes $[X_u]$ as

$$\langle s_w, [X_u] \rangle = \delta_{w,u}, \quad w, u \in W(H; G).$$

Lemma 2 implies that (cf. [BGG, §5])

Corollary 2 (Basis Theorem). *The set of Schubert classes $\{s_w \mid w \in W(H; G)\}$ constitutes an additive basis for the Chow ring $A^*(G/H)$.*

Referring to the index system (2.3) on $W^r(H; G)$, the notion $s_{r,i}$ will be used to simplify $s_{w_r,i}$. We create also a definition emphasizing the role that the minimized decomposition of w has played in the construction (2.4) of the Schubert variety X_w :

Definition 4. The minimized decomposition $\sigma[I]$ of a $w \in W(H; G)$ will be referred to as the *Weyl coordinate* of s_w .

2.3. Multiplying Schubert classes Let f be a polynomial of homogeneous degree $2r$ in Schubert classes $\{s_w \mid w \in W(H; G)\}$. By considering f as an element in $A^{2r}(G/H)$ one has the expression

$$(2.5) \quad f = \sum_{w \in W^r(H; G)} a_w(f) s_w, \quad a_w(f) \in \mathbb{Z}$$

in view of the Basis Theorem. Effective computation in the ring $A^*(G/H)$ amounts to find a method to evaluate the integer $a_w(f)$ for any f and w . In the special case $f = s_u s_v$ (i.e. product of two Schubert classes), the $a_w(f)$ are well known as *the structure constants* for multiplying Schubert classes [Br; Bu; L; P].

A unified formula evaluating $a_w(f)$ can be given in term of the minimized decomposition of $w \in W(H; G)$. To explain this we need a few notations.

Let $\mathbb{Z}[x_1, \dots, x_k] = \bigoplus_{r \geq 0} \mathbb{Z}[x_1, \dots, x_k]^{(r)}$ be the ring of polynomials in x_1, \dots, x_k with integer coefficients, graded by $|x_i| = 1$.

Given a $k \times k$ strictly upper triangular integer matrix $A = (a_{i,j})$ define a homomorphism $T_A : \mathbb{Z}[x_1, \dots, x_k]^{(k)} \rightarrow \mathbb{Z}$ recursively by the following

Elimination laws:

- 1) if $h \in \mathbb{Z}[x_1, \dots, x_{k-1}]^{(k)}$, then $T_A(h) = 0$;
- 2) if $k = 1$ (consequently $A = (0)$), then $T_A(x_1) = 1$;
- 3) if $h \in \mathbb{Z}[x_1, \dots, x_{k-1}]^{(k-r)}$ with $r \geq 1$, then

$$T_A(hx_k^r) = T_{A'}(h(a_{1,k}x_1 + \dots + a_{k-1,k}x_{k-1})^{r-1}),$$

where A' is the $((k-1) \times (k-1))$ strictly upper triangular matrix obtained from A by deleting the k^{th} column and k^{th} row.

By additivity, T_A is defined for every $f \in \mathbb{Z}[x_1, \dots, x_k]^{(k)}$ using the unique expansion $f = \sum h_r x_k^r$ with $h_r \in \mathbb{Z}[x_1, \dots, x_{k-1}]^{(k-r)}$.

For a $w \in W(H; G)$ with minimized decomposition $w = \sigma[i_1, \dots, i_k]$, let $A_w = (a_{s,t})$ be the $k \times k$ (strictly upper triangular) with

$$a_{s,t} = \begin{cases} 0 & \text{if } s \geq t; \\ -\beta_{i_t} \circ \beta_{i_s} & \text{if } s < t. \end{cases}$$

Definition 5. The additive map $T_w =: T_{A_w} : \mathbb{Z}[x_1, \dots, x_k]^{(k)} \rightarrow \mathbb{Z}$ is called the *triangular operator* associated to w .

The next result is seen as a natural generalization of the theorem in [D₂].

Lemma 3. For any $w \in W^r(H; G)$ we have

$$a_w(f) = T_w(g(x_1, \dots, x_k)),$$

where $g(x_1, \dots, x_k)$ is the polynomial obtained from f by substituting the Schubert class s_u by $\sum x_I$, where the sum is over all $I \subset [i_1, \dots, i_k]$ with $u = \sigma[I]$, and where $x_I = x_{j_1} \cdots x_{j_r}$ if $I = [j_1, \dots, j_r]$.

Based on Lemma 3, a program entitled “*Littlewood-Richardson Coefficients*” (abbreviated as *L-R Coefficients* in sequel) implementing $a_w(f)$ has been compiled (see also [DZ₁]), whose function is briefed below:

Algorithm. *L-R coefficients.*

Input: A polynomial f in Schubert classes on G/H ; a $w \in W(H; G)$ given by its minimized decomposition.

Output: $a_w(f) \in \mathbb{Z}$.

Example 2. The data in [DZ₂, 1.2–6.2; 1.3–6.3; 1.4–6.4] are generated by the *L-R coefficients*.

3 The quotient of a polynomial ring

3.1. The problems. Let A be a finitely generated commutative ring, graded by $A = \bigoplus_{r \geq 0} A^r$. An element $y \in A$ is called *homogeneous* of degree r if $y \in A^r$. All elements y in a graded ring (e.g. cohomology ring; the quotient of a polynomial ring) concerned in this paper are homogeneous, and their degree is denoted by $|y|$.

An ordered subset $S = \{y_1, \dots, y_n\}$ of A is called a *set of generators* if the ordering on S satisfies $|y_1| \leq \dots \leq |y_n|$; and A is generated multiplicatively by elements in S .

Given two sets $S = \{y_1, \dots, y_n\}, T = \{z_1, \dots, z_{n'}\}$ of generators of A , the notion $S \leq T$ is adopted to indicate the statement that “one has either $n < n'$ or, $n = n'$ but $|y_1| = |z_1|, \dots, |y_{k-1}| = |z_{k-1}|, |y_k| < |z_k|$ for some $k \leq n$ ”.

Definition 6. A set S of generators of A is said to be *minimal* if $S \leq T$ for any other set T of generators of A .

Problem 1. Given a flag variety G/H , find a minimal set $S = \{y_1, \dots, y_n\}$ of generators of $A^*(G/H)$ that consists of Schubert classes on G/H .

Suppose that a solution to Problem 1 is afforded by $S = \{y_1, \dots, y_n\}$, and let $\mathbb{Z}[y_1, \dots, y_n]$ be the ring of integral polynomials in y_1, \dots, y_n . The inclusion $\{y_1, \dots, y_n\} \subset A^*(G/H)$ then induces a surjective ring map

$$\pi : \mathbb{Z}[y_1, \dots, y_n] \rightarrow A^*(G/H),$$

whose kernel $\ker \pi \subset \mathbb{Z}[y_1, \dots, y_n]$ is an ideal.

Problem 2. Find a set $\{r_1, \dots, r_m\} \subset \mathbb{Z}[y_1, \dots, y_n]$ of polynomials so that the ideal $\langle r_1, \dots, r_m \rangle$ generated by r_1, \dots, r_m agrees with $\ker \pi$.

If solutions to both problems are archived, one may arrive at the desired *Schubert presentation* of the ring $A^*(G/H)$ [IM]:

$$A^*(G/H) = \mathbb{Z}[y_1, \dots, y_n] / \langle r_1, \dots, r_m \rangle.$$

In comparison, Problem 1 is relatively easy to deal with by geometric means. On the other hand, difficulties in working with Problem 2 may arise from great variety of choices of the subset $\{r_1, \dots, r_m\}$ with $\langle r_1, \dots, r_m \rangle = \ker \pi$, any particular choice giving rise to artificial looking expressions. So, while looking for a solution to Problem 2, two additional requests should be concerned:

- 1) *the m should be as less as possible; and at the same time,*
- 2) *each r_i should have the simplest expression.*

This section is devoted to two machineries (Lemma 4 and 5) that take care of these two requirements respectively.

3.2. Eliminating relations. Let $\mathbb{Z}[y_1, \dots, y_n]$ be the graded ring of polynomials in n variables y_1, \dots, y_n with preassigned degrees $|y_i| > 0$, and let \mathbb{N}^n be the set of all n -tuples $\alpha = (b_1, \dots, b_n)$ of non-negative integers. Assign to each $\alpha = (b_1, \dots, b_n) \in \mathbb{N}^n$ the monomial $y^\alpha = y_1^{b_1} \dots y_n^{b_n} \in \mathbb{Z}[y_1, \dots, y_n]$ furnished with the degree $|y^\alpha| =: \sum b_k |y_k|$. In this way the $\mathbb{Z}[y_1, \dots, y_n]$ becomes a graded ring $\bigoplus_{m \geq 0} \mathbb{Z}[y_1, \dots, y_n]^{(m)}$ with

$$\mathbb{Z}[y_1, \dots, y_n]^{(m)} = \text{Span}_{\mathbb{Z}}\{y^\alpha \mid |y^\alpha| = m\}.$$

This suggests us to introduce the *monomial basis* of $\mathbb{Z}[y_1, \dots, y_n]^{(m)}$ as

$$(3.1) \quad B(m) = \{y^\alpha \mid |y^\alpha| = m\},$$

regarded as an ordered set with respect to the lexicographical order on α 's. The rank of $\mathbb{Z}[y_1, \dots, y_n]^{(m)}$ (the cardinality of $B(m)$) is denoted by $b(m)$.

Let $\{r_1, \dots, r_k\} \subset \mathbb{Z}[y_1, \dots, y_n]$ be a set of polynomials. The kernel of the quotient map

$$\psi : \mathbb{Z}[y_1, \dots, y_n] \rightarrow \mathbb{Z}[y_1, \dots, y_n] / \langle r_1, \dots, r_k \rangle$$

in degree m , denoted by $\Phi_m(r_1, \dots, r_k)$, is spanned additively by the set of polynomials

$$\Sigma_m(r_1, \dots, r_k) = \{y^\alpha r_i \mid |y^\alpha| + |r_i| = m\},$$

whose cardinality is easily seen to be $c_m(r_1, \dots, r_k) = b(m - |r_1|) + \dots + b(m - |r_k|)$. In terms of the ordered basis $B(m)$, every $y^\alpha r_i \in \Sigma_m(r_1, \dots, r_k)$ admits a unique expansion as

$$y^\alpha r_i = \sum_{y^\beta \in B(m)} a_{(\alpha, i), \beta} y^\beta, \quad a_{(\alpha, i), \beta} \in \mathbb{Z}.$$

Write $M_m(r_1, \dots, r_k)$ for the matrix $(a_{(\alpha, i), \beta})_{c(r_1, \dots, r_k; m) \times b(m)}$ (with respect to some order on $\Sigma_m(r_1, \dots, r_k)$) so obtained.

Definition 7. The *deficiency* of the set $\{r_1, \dots, r_k\}$ in degree m , denoted by $\delta_m(r_1, \dots, r_k)$, is the invariant of the matrix $M_m(r_1, \dots, r_k)$ computed as follows (cf. [S, p.163-166])

- 1) diagonalize $M_m(r_1, \dots, r_k)$ using integral row and column operations;
- 2) set $\delta_m(r_1, \dots, r_k)$ to be the numbers of 1's appearing in the resulting diagonal matrix.

Example 3. Based on the algorithm on integral row and column reductions given in [S, p.163], a program computing the $\delta_m(r_1, \dots, r_k)$ has been composed. However, when $b(m)$ is relatively small, the $\delta_m(r_1, \dots, r_k)$ can of course be computed directly. As an example, consider the ring $\mathbb{Z}[y_1, y_5, y_9]$ with $|y_i| = 2i$, and let $r_{10}, r_{14}, r_{18} \in \mathbb{Z}[y_1, y_5, y_9]$ be given respectively by

$$\begin{aligned}
r_{10} &= y_5^2 - 2y_1y_9; \\
r_{14} &= 2y_5y_9 - 18y_1^5y_9 + 6y_1^9y_5 - y_1^{14}; \\
r_{18} &= y_9^2 + 20y_1^4y_5y_9 + 2y_1^{13}y_5 - 18y_1^9y_9,
\end{aligned}$$

(cf. Theorem 5). If $m = 36$ we find that

$$\begin{aligned}
B(36) &= \{y_9^2, y_1^3y_5^3, y_1^4y_5y_9, y_1^8y_5^2, y_1^9y_9, y_1^{13}y_5, y_1^{18}\}; \\
\Sigma_{36}(r_{10}, r_{14}, r_{18}) &= \{r_{18}, y_1^4r_{14}, y_1^3y_5r_{10}, y_1^8r_{10}\};
\end{aligned}$$

and that

$$M_{36}(r_{10}, r_{14}, r_{18}) = \begin{pmatrix} 1 & 0 & 20 & 0 & -18 & 2 & 0 \\ 0 & 0 & 2 & 0 & -18 & 6 & -1 \\ 0 & 1 & -2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -2 & 0 & 0 \end{pmatrix}.$$

These yield $b(36) = 7$, $\delta_{36}(r_{10}, r_{14}, r_{18}) = 4$ (for, as is clear, the $M_{36}(r_{10}, r_{14}, r_{18})$ has a 4×4 minor that equals to 1).

For another subset $\{g_1, \dots, g_s\} \subset \mathbb{Z}[y_1, \dots, y_n]$ consider the quotient map

$$\varphi : \mathbb{Z}[y_1, \dots, y_n] / \langle r_1, \dots, r_k \rangle \rightarrow A^* = \bigoplus_{m \geq 0} A^m,$$

where $A^* = \mathbb{Z}[y_1, \dots, y_n] / \langle r_1, \dots, r_k, g_1, \dots, g_s \rangle$. The next result tells the manner by which the integers $b(m)$, $\delta_m(r_1, \dots, r_k)$ are used in eliminating relations in a quotient of $\mathbb{Z}[y_1, \dots, y_n]$.

Lemma 4. *Assume that, for all $m = |g_i|$, $1 \leq i \leq s$,*

$$\text{rank}(A^m) = b(m) - \delta_m(r_1, \dots, r_k).$$

Then, $\{g_1, \dots, g_s\} \subset \langle r_1, \dots, r_k \rangle$. That is, φ is a ring isomorphism.

Proof. For a g_i we set $m = |g_i|$, $\delta_m(r_1, \dots, r_k) = t$. Then, there is a subset $\{f_1, \dots, f_t\} \subset \Phi_m(r_1, \dots, r_k)$ of cardinality t that can be extended to a basis of $\mathbb{Z}[y_1, \dots, y_n]^{(m)}$ (cf. [S, Theorem 13,1]). That is, there exist $h_1, \dots, h_{b(m)-t} \in \mathbb{Z}[y_1, \dots, y_n]^{(m)}$ so that the union

$$\Lambda = \{f_1, \dots, f_t\} \cup \{h_1, \dots, h_{b(m)-t}\}$$

is a basis of $\mathbb{Z}[y_1, \dots, y_n]^{(m)}$.

Expanding g_i in terms of Λ gives rise to

$$g_i = a_1h_1 + \dots + a_{b(m)-t}h_{b(m)-t} + c_1f_1 + \dots + c_tf_t, \quad a_i, b_j \in \mathbb{Z}.$$

Assume on the contrary that $g_i \notin \langle r_1, \dots, r_k \rangle$. Then the coefficients a_k 's are not all zero. One gets from $\varphi(g_i) = 0$ in A^m and $\{f_1, \dots, f_t\} \subset \langle r_1, \dots, r_k \rangle$ that $\text{rank}(A^m) \leq b(m) - t - 1$, a contradiction to the assumption.

3.3. The Nullspace. Let $S = \{y_1, \dots, y_n\}$ be any subset of Schubert classes on a flag variety G/H . Assign to y_i the degree $|y_i|$ = the dimension of y_i as a Schubert class. The inclusion $S \subset A^*(G/H)$ induces a ring map $\pi : \mathbb{Z}[y_1, \dots, y_n] \rightarrow A^*(G/H)$ whose restriction on degree $2m$ is denoted by

$$\pi_m : \mathbb{Z}[y_1, \dots, y_n]^{(2m)} \rightarrow A^{2m}(G/H).$$

Combining the L - R coefficients (cf. **2.3**) with the function “Nullspace” in *Mathematica*, a basis for $\ker \pi_m$ can be explicitly exhibited.

Since $A^{2m}(G/H)$ has the canonical basis given by the set of Schubert classes $\{s_{m,i} \mid 1 \leq i \leq \beta(m)\}$ (cf. (2.3) and **2.2**), for each $y^\alpha \in B(2m)$ one has the expression in $A^{2m}(G/H)$

$$\pi_m(y^\alpha) = c_{\alpha,1}s_{m,1} + \dots + c_{\alpha,t_m}s_{m,\beta(m)}, \quad c_{\alpha,i} \in \mathbb{Z}.$$

where the coefficients $c_{\alpha,i}$ can be evaluated by the L - R coefficients as $c_{\alpha,i} = a_{w_{m,i}}(y^\alpha)$ since every y^α is a monomial in the Schubert classes (cf. Lemma 3). The matrix $M(\pi_m) = (c_{\alpha,i})_{b(2m) \times \beta(m)}$ so obtained will be referred to as the structure matrix of π_m .

The built-in function *Nullspace* in *Mathematica* transforms the $M(\pi_m)$ to another matrix $N(\pi_m)$ in the fashion

$$\text{In:} = \text{Nullspace}[M(\pi_m)]$$

$$\text{Out:} = \text{a matrix } N(\pi_m) = (b_{j,\alpha})_{(b(2m)-\beta(m)) \times b(2m)}.$$

The significance of $N(\pi_m)$ is shown in the next result.

Lemma 5. *The set of polynomials*

$$k_i = \sum_{y^\alpha \in B(2m)} b_{i,\alpha} y^\alpha, \quad 1 \leq i \leq (b(2m) - \beta(m)),$$

is a basis for $\ker \pi_m$.

Example 4. See in [DZ₂, 1.4–6.4; 1.5–6.5] for examples of structure matrices and their *Nullspaces*.

4 Computing with Gysin sequence

Assume from now on that G/H is a Grassmannian of G associated to the k^{th} weight $\omega_k \in \Omega$. Based on Gysin sequence of oriented circle bundles, we derive partial solutions to problem 1 and 2 from information on $H^*(G/H_s)$ in Lemma 7 and 8; and develop a procedure to compute the ring $H^*(G/H_s)$.

For a topological space X we put

$$H^{\text{even}}(X) = \bigoplus_{r \geq 0} H^{2r}(X), \quad H^{\text{odd}}(X) = \bigoplus_{r \geq 0} H^{2r+1}(X).$$

Note that $H^{even}(X) \subset H^*(X)$ is always a subring.

4.1. The generators of $H^*(G/H)$. Since the set $W^1(H, G)$ consists of the single element $\{w_{1,1} = \sigma[k]\}$, the Basis Theorem implies that:

Lemma 6. $H^2(G/H) = \mathbb{Z}$ is generated by $\omega =: s_{1,1}$.

The natural projection $p : G/H_s \rightarrow G/H$ is an oriented circle bundle over G/H with Euler class $\omega \in H^2(G/H)$. Since $H^{odd}(G/H) = 0$ by the basis Theorem, the Gysin sequence [MS, p.143] of p yields the short exact sequence

$$(4.1) \quad 0 \rightarrow \omega H^{2r-2}(G/H) \rightarrow H^{2r}(G/H) \xrightarrow{p^*} H^{2r}(G/H_s) \rightarrow 0$$

as well as the isomorphism ($\omega \cdot$ means taking cup-product with ω)

$$(4.2) \quad \beta : H^{2r-1}(G/H_s) \xrightarrow{\cong} \text{Ker}\{H^{2r-2}(G/H) \xrightarrow{\omega} H^{2r}(G/H)\}.$$

We observe from (4.1) that a minimal set of generators of $H^*(G/H)$ can be selected from the simpler ring $H^{even}(G/H_s)$:

Lemma 7. If $S = \{y_1, \dots, y_m\} \subset H^*(G/H)$ is a subset so that the $p^*S = \{p^*(y_1), \dots, p^*(y_m)\}$ is a minimal set of generators of $H^{even}(G/H_s)$, then $S' = \{\omega, y_1, \dots, y_m\}$ is a minimal set of generators of $H^*(G/H)$.

Proof. Firstly, with the assumption that p^*S is a minimal set of generators, we show by induction on r that

$$(4.3) \quad \text{each } y \in H^{2r}(G/H) \text{ can be expressed as a polynomial in } \{\omega, y_1, \dots, y_m\}.$$

The case $r = 1$ has been done by Lemma 6. So suppose that (4.3) holds for all $r < n$. Consider next the case $r = n$.

Since p^*S is a set of generators of $H^{even}(G/H_s)$, there exists a polynomial f in the $p^*(y_1), \dots, p^*(y_m)$ so that $p^*(y) = f(p^*(y_1), \dots, p^*(y_m))$. Clearly, $y - f(y_1, \dots, y_m) \in \text{ker } p^*$. It follows from (4.1) as well as the inductive hypothesis that $y - f(y_1, \dots, y_m) = \omega g(\omega, y_1, \dots, y_m)$ for some $g(\omega, y_1, \dots, y_m) \in H^{2(n-1)}(G/H)$. (4.3) is verified by the expression

$$y = f(y_1, \dots, y_m) + \omega g(\omega, y_1, \dots, y_m).$$

Next, let $T = \{z_0, z_1, \dots, z_{m'}\}$ be any set of generators of $H^*(G/H)$. We may assume $z_0 = \omega$ by Lemma 6. Since $p^* : H^{even}(G/H) \rightarrow H^{even}(G/H_s)$ is surjective and annihilates ω , $p^*T = \{p^*(z_1), \dots, p^*(z_{m'})\}$ is a set of generators of $H^{even}(G/H_s)$. From $p^*S \leq p^*T$ (by the minimal assumption on p^*S) one gets $S \leq T$. This finishes the proof.

4.2. Locating the degrees of relations. Let $S = \{y_1, \dots, y_m\} \subset H^*(G/H)$ be a subset so that $p^*S = \{p^*(y_1), \dots, p^*(y_m)\}$ is a minimal set of generators of $H^{even}(G/H_s)$. The inclusions $\{\omega\} \cup S \subset H^*(G/H)$, $p^*S \subset H^*(G/H_s)$ extend to surjective ring maps π and $\bar{\pi}$ that fit in the commutative diagram

$$(4.4) \quad \begin{array}{ccccccc} \mathbb{Z}[\omega, y_1, \dots, y_m]^{(2r)} & \xrightarrow{\varphi} & \mathbb{Z}[y_1, \dots, y_m]^{(2r)} & & & & \\ \pi \downarrow & & \bar{\pi} \downarrow & & & & \\ \rightarrow H^{2r-2}(G/H) \rightarrow & & H^{2r}(G/H) & \xrightarrow{p^*} & H^{2r}(G/H_s) & \rightarrow & 0 \end{array}$$

where $\mathbb{Z}[\omega, y_1, \dots, y_m]$ is graded by $|\omega|, |y_1|, \dots, |y_m|$, and where

$$\varphi(\omega) = 0, \varphi(y_i) = y_i; \bar{\pi}(y_i) = p^*(y_i).$$

The graded group $H^{odd}(G/H_s)$ is always free by (4.2), and will be considered a module over $H^{even}(G/H_s)$ via cup-product

$$H^{even}(G/H_s) \times H^{odd}(G/H_s) \rightarrow H^{odd}(G/H_s), (x, y) \rightarrow x \cup y.$$

Lemma 8. *If $\{h_1, \dots, h_n\} \subset \mathbb{Z}[y_1, \dots, y_m]$ is a subset such that*

$$(4.5) \quad H^{even}(G/H_s) = \mathbb{Z}[p^*(y_1), \dots, p^*(y_m)] / \langle p^*(h_1), \dots, p^*(h_n) \rangle,$$

and if $\{d_1, \dots, d_t\}$ is a basis for $H^{odd}(G/H_s)$ as an $H^{even}(G/H_s)$ -module; then, for any two subsets $\{r_1, \dots, r_n\}, \{g_1, \dots, g_t\} \subset \mathbb{Z}[\omega, y_1, \dots, y_m]$ that satisfy

- 1) $r_i \in \ker \pi$ with $r_i|_{\omega=0} = h_i$, $1 \leq i \leq n$; and
- 2) $\pi(g_i) = \beta(d_i)$, $1 \leq i \leq t$,

respectively, one has

$$(4.6) \quad H^*(G/H) = \mathbb{Z}[\omega, y_1, \dots, y_m] / \langle r_1, \dots, r_n; \omega g_1, \dots, \omega g_t \rangle.$$

Proof. Observe that

(a) the condition $r_i|_{\omega=0} = h_i$ is equivalent to $r_i = h_i + \omega f_i$ for some $f_i \in \mathbb{Z}[\omega, y_1, \dots, y_m]$;

(b) the (4.5) implies that in (4.4), $\ker \bar{\pi} = \langle h_1, \dots, h_n \rangle$.

It suffices for us to show that

$$(4.7) \quad \text{for any } \alpha \in \ker \pi, \alpha \in \langle r_1, \dots, r_n; \omega g_1, \dots, \omega g_t \rangle$$

for, as is clear, $\ker \pi \supseteq \langle r_1, \dots, r_n; \omega g_1, \dots, \omega g_t \rangle$. This is done by induction on $2r = |\alpha|$. The case $r = 1$ trivial by Lemma 6. So suppose that (4.7) holds for all α with $|\alpha| \leq 2r - 2$, and consider the case $|\alpha| = 2r$.

The α can be uniquely expressed as $\alpha = \alpha_1 + \omega \alpha_2$ with $\alpha_1 \in \mathbb{Z}[y_1, \dots, y_m]$. From $p^*\pi(\alpha) = \bar{\pi}\alpha_1 = 0$ and (b) we get $\alpha_1 = a_1 h_1 + \dots + a_n h_n$ for some $a_i \in \mathbb{Z}[y_1, \dots, y_m]$. We can rewrite, in view of (a), that

$$(4.8) \quad \alpha = a_1 r_1 + \cdots + a_n r_n + \omega \alpha_3, \text{ where } \alpha_3 = \alpha_2 - (a_1 f_1 + \cdots + a_n f_n).$$

From $\pi(\alpha) = 0$, $\pi(r_k) = 0$ we find $\pi(\alpha_3) \in \text{Ker}\{H^{2r-2}(G/H) \xrightarrow{\cup\omega} H^{2r}(G/H)\}$. Since $\beta : H^{2r-1}(G/H_s) \rightarrow \text{Ker}\{H^{2r-2}(G/H) \xrightarrow{\cup\omega} H^{2r}(G/H)\}$ is an isomorphism by (4.2), and since $H^{odd}(G/H_s)$ (as an $H^{even}(G/H_s)$ -module) has the basis $\{d_1, \cdots, d_t\}$ by the assumption, one has

$$(4.9) \quad \pi(\alpha_3) = \sum b_i \beta(d_i) \text{ for some } b_i \in H^*(G/H).$$

Since π is surjective, $b_i = \pi(q_i)$ for some $q_i \in \mathbb{Z}[\omega, y_1, \cdots, y_m]$. Set $\gamma = \alpha_3 - \sum q_i g_i$, where g_1, \cdots, g_t are given as that in the lemma. The (4.8) becomes

$$(4.10) \quad \alpha = a_1 r_1 + \cdots + a_n r_n + \omega \gamma + \sum b_i (\omega g_i).$$

Since $|\gamma| = |\alpha| - 2$ with $\pi(\gamma) = 0$ by (4.9), the inductive hypothesis concludes $\gamma \in \langle r_1, \cdots, r_n; \omega g_1, \cdots, \omega g_t \rangle$. (4.7) is verified by (4.10).

4.3. Algorithm for computing $H^*(G/H_s)$. We conclude this section with a procedure computing the integral cohomology of G/H_s . This will be applied, in the coming section, to determine $H^*(G/H_s)$ for the (G, H) concerned by Theorem 1-6.

The method begins with finding an additive basis of $H^*(G/H_s)$; followed by deriving multiplication formulae for the subring $H^{even}(G/H_s)$; and completed by describing $H^{odd}(G/H_s)$ as an module over $H^{even}(G/H_s)$.

Step 1. Finding a basis for $H^*(G/H_s)$. According to (4.1) and (4.2), the additive groups $H^{2k-1}(G/H_s)$ and $H^{2k}(G/H_s)$ are completely determined by the homomorphism $H^{2k-2}(G/H) \xrightarrow{\cup\omega} H^{2k}(G/H)$.

Set $\beta(r) = |W^r(H; G)|$ (as in (2.3)). With respect to the basis $\{s_{r,1}, \cdots, s_{r,\beta(r)}\}$ of $H^{2r}(G/H)$ for $r = k-1, k$ one has the expressions

$$\omega s_{k-1,i} = \sum a_{i,j} s_{k,j}, \quad a_{i,j} \in \mathbb{Z}.$$

Equivalently,

$$(4.11) \quad \begin{pmatrix} \omega s_{k-1,1} \\ \omega s_{k-1,2} \\ \vdots \\ \omega s_{k-1,\beta(k-1)} \end{pmatrix} = A_k \begin{pmatrix} s_{k,1} \\ s_{k,2} \\ \vdots \\ s_{k,\beta(k)} \end{pmatrix}, \quad A_k = (a_{i,j})_{\beta(k-1) \times \beta(k)}.$$

Since each $\omega s_{k-1,i}$ is a monomial in Schubert classes, the entries of A_k can be evaluated by using the *L-R coefficients* (cf. **2.3**). Diagonalizing A_k by using the standard integral row and column reductions (cf. [S, p.162-166])

enables one to specify bases for $H^{2k}(G/H_s)$ and $H^{2k-1}(G/H_s)$ (together with orders of the basis elements) in terms of Schubert classes on G/H .

Example 5. For those (G, H) concerned by Theorem 1-6, the matrices A_k have all been computed and tabulated in [DZ₂, 1.2–6.2]. See also the tables in the proofs of Theorem 7-12 in §5 for the basis of $H^*(G/H_s)$ so derived.

Step 2. The ring structure on $H^{even}(G/H_s)$. It has been shown in step 1 that a basis for $H^*(G/H_s)$ can be selected in terms of the matrix A_k in (4.11). In practice, in view of the surjective ring map $p^*: H^{even}(G/H) \rightarrow H^{even}(G/H_s)$, it is possible to find a subset Λ of the Schubert classes $s_{k,i}$ on G/H , so that

$$(4.12) \quad p^*\Lambda = \{\bar{s}_{r,i} =: p^*(s_{k,i}) \mid s_{k,i} \in \Lambda\} \text{ constitutes a basis for } H^{even}(G/H_s).$$

Given two basis elements $\bar{s}_{r,i}, \bar{s}_{k,j} \in p^*\Lambda$ consider their corresponding product in $H^*(G/H)$:

$$s_{r,i}s_{k,j} = \sum b_{(r,i),(k,j)}^t s_{r+k,t},$$

where, again, the constants $b_{(r,i),(k,j)}^t$ can be computed by the *L-R coefficients* (i.e. Lemma 3). Applying p^* yields the equation in $H^{even}(G/H_s)$

$$\bar{s}_{r,i}\bar{s}_{k,j} = \sum b_{(r,i),(k,j)}^t p^* s_{r+k,t}.$$

Expressing the $p^* s_{r+k,t}$ in the right hand side in terms of the elements in $p^*\Lambda$ gives rise to the *multiplicative rule* of the basis elements in $p^*\Lambda$

$$(4.13) \quad \bar{s}_{r,i}\bar{s}_{k,j} = \sum_{s_{r+k,t} \in \Lambda} c_{(r,i),(k,j)}^t \bar{s}_{r+k,t}$$

Clearly, (4.13) suffices to characterize $H^{even}(G/H_s)$ as a ring.

Example 6. For those (G, H) concerned by Theorem 1-6, the formulae (4.13) have been decided and are listed in [DZ₂, 1.3–6.3].

Step 3. The $H^{odd}(G/H_s)$ as an $H^{even}(G/H_s)$ -module. Since the graded group $H^{odd}(G/H_s)$ is torsion free by (4.2), one has $y \cdot H^{odd}(G/H_s) = 0$ for all $y \in \text{Tor}(H^{even}(G/H_s))$. For this reason the pairing $H^{even}(G/H_s) \times H^{odd}(G/H_s) \rightarrow H^{odd}(G/H_s)$ in 4.2 is reduced to

$$(4.14) \quad [H^{even}(G/H_s)/\text{Tor}(H^{even}(G/H_s))] \times H^{odd}(G/H_s) \rightarrow H^{odd}(G/H_s).$$

The G/H_s is an orientable manifold with odd dimension. The Poincaré duality tells that

Lemma 9. *If $\dim_{\mathbb{R}} G/H_s = 2b+1$, the product (4.14) in the complementary dimensions $[H^{2r}/\text{Tor}(H^{2r})] \times H^{2(b-r)+1} \rightarrow H^{2b+1} = \mathbb{Z}$ are all non-singular.*

We shall see in the proof of Theorem 7-12 that, practically, Lemma 9 suffices to characterize $H^{odd}(G/H_s)$ as an $H^{even}(G/H_s)$ module

5 Integral cohomology of G/H_s

Following the instruction in 4.3, we calculate the rings $H^*(G/H_s)$ for the (G, H) concerned by Theorem 1-6. The results are stated in Theorem 7-12, where relevance of the generators with Schubert classes on G/H is emphasized.

Given a set $\{d_1, \dots, d_t\}$ of elements with preassigned degrees $|d_i| > 0$, let $\Gamma(1, d_1, \dots, d_t)$ be the free abelian group generated by $1, d_1, \dots, d_t$, considered as a graded ring with the trivial products $1 \cdot d_i = d_i$; $d_i \cdot d_j = 0$.

Let A be a graded commutative ring. Denote by $A \widehat{\otimes} \Gamma(1, d_1, \dots, d_t)$ the quotient ring of the tensor product $A \otimes \Gamma(1, d_1, \dots, d_t)$ subject to the relations $Tor(A) \cdot d_i = 0$, $1 \leq i \leq t$.

If $y \in H^*(G/H)$ we set $\bar{y} = p^*(y) \in H^*(G/H_s)$.

Theorem 7. *Let $y_3, y_4, y_6 \in H^*(F_4/C_3 \cdot S^1)$ be the Schubert class with Weyl coordinates $\sigma[3, 2, 1]$, $\sigma[4, 3, 2, 1]$, $\sigma[3, 2, 4, 3, 2, 1]$ respectively, and let $d_{23} \in H^{23}(F_4/C_3)$ be with $\beta(d_{23}) = 2s_{11,1} - s_{11,2}$. Then*

$$H^*(F_4/C_3) = \frac{\mathbb{Z}[\bar{y}_3, \bar{y}_4, \bar{y}_6]}{\langle h_3; h_6; h_8; h_{12} \rangle} \widehat{\otimes} \Gamma(1, d_{23}),$$

where $h_6 : 2\bar{y}_3 = 0$; $h_6 : 2\bar{y}_6 - \bar{y}_3^2 = 0$; $h_8 : 3\bar{y}_4^2 = 0$; $h_{12} = \bar{y}_6^2 - \bar{y}_4^3$.

Proof. Step 1. With the matrices A_k in (4.11) being computed by the L - R coefficients and presented in [DZ₂, 1.2], row and column reduction yield results in the first two columns of the following table, which characterizes $H^*(F_4/C_3)$ as a graded group:

nontrivial $H^k(F_4/C_3)$	basis elements	relations
$H^6 = \mathbb{Z}_2$	$\bar{s}_{3,1}$	
$H^8 = \mathbb{Z}$	$\bar{s}_{4,2}$	
$H^{12} = \mathbb{Z}_4$	$\bar{s}_{6,2}$	$-2\bar{s}_{6,2} = \bar{s}_{3,1}^2$
$H^{14} = \mathbb{Z}_2$	$\bar{s}_{7,1}$	$= \bar{s}_{3,1}\bar{s}_{4,2}$
$H^{16} = \mathbb{Z}_3$	$\bar{s}_{8,1}$	$= -\bar{s}_{4,2}^2$
$H^{18} = \mathbb{Z}_2$	$\bar{s}_{9,2}$	$= \bar{s}_{3,1}\bar{s}_{6,2}$
$H^{20} = \mathbb{Z}_4$	$\bar{s}_{10,2}$	$= \bar{s}_{4,2}\bar{s}_{6,2}$
$H^{26} = \mathbb{Z}_2$	$\bar{s}_{13,1}$	$= \bar{s}_{3,1}\bar{s}_{4,2}\bar{s}_{6,2}$
$H^{23} = \mathbb{Z}$	$d_{23} = \beta^{-1}(2s_{11,1} - s_{11,2})$	
$H^{31} = \mathbb{Z}$	$d_{31} = \beta^{-1}(s_{15,1})$	$= \pm \bar{s}_{4,2}d_{23}$

Step 2. Items in the second column tell that H^{even} has additive basis of the form $p^*\Lambda$ with $\Lambda = \{s_{3,1}, s_{4,2}, s_{6,2}, s_{7,1}, s_{8,1}, s_{9,2}, s_{10,2}, s_{13,1}\}$. By algorithm given in 4.3, the multiplicative rule (4.13) for the basis elements in $p^*\Lambda$ have been determined in [DZ₂, 1.3], and recorded in the last column of the table corresponding to H^{even} . These imply that, if we put $y_3 = s_{3,1}$, $y_4 = s_{4,2}$, $y_6 = s_{6,2}$, then

a) y_3, y_4, y_6 are the Schubert classes whose Weyl coordinates are given as that as in the theorem by [DZ₂, 1.1];

b) $H^{even}(F_4/C_3)$ is generated by $\bar{y}_3, \bar{y}_4, \bar{y}_6$ subject to the relations $h_3; h_6; h_8$ (cf. the theorem).

Combining these with the obvious relations $\bar{y}_6^2 = \bar{y}_4^3 = 0$ (because of $H^{24} = 0$ by the first column), together with the fact that, as ideals in $\mathbb{Z}[\bar{y}_3, \bar{y}_4, \bar{y}_6]$,

$$\langle h_3; h_6; h_8; \bar{y}_6^2; \bar{y}_4^3 \rangle = \langle h_3; h_6; h_8; h_{12} \rangle,$$

one obtains

$$(5.1) \quad H^{even}(F_4/C_3) = \frac{\mathbb{Z}[\bar{y}_3, \bar{y}_4, \bar{y}_6]}{\langle h_3; h_6; h_8; h_{12} \rangle}.$$

Step 3. The proof is completed by $d_{31} = \pm \bar{s}_{4,2} d_{23}$ (Lemma 9) and $d_{23}^2 \in H^{46} = 0$ (in view of the first column of the table).

Theorem 8. Let $y_4 \in H^8(F_4/B_3 \cdot S^1)$ be the Schubert class with Weyl coordinate $\sigma[3, 2, 3, 4]$; and let $d_{23} \in H^{23}(F_4/B_3)$ be with $\beta(d_{23}) = -s_{11,1} + s_{11,2}$. Then

$$H^*(F_4/B_3) = \frac{\mathbb{Z}[\bar{y}_4]}{\langle h_8; h_{12} \rangle} \widehat{\otimes} \Gamma(1, d_{23}),$$

where $h_8 = 3\bar{y}_4^2; h_{12} = \bar{y}_4^3$.

Proof. Step 1. With the matrices A_k in (4.11) being computed by the L - R coefficients and presented in [DZ₂, 2.2], row and column reduction yield results in the first two columns of the following table, which characterizes $H^*(F_4/B_3)$ as a graded group:

nontrivial H^k	basis elements	relations
$H^8 = \mathbb{Z}$	$\bar{s}_{4,2}$	
$H^{16} = \mathbb{Z}_3$	$\bar{s}_{8,1}$	$= \bar{s}_{4,2}^2$
$H^{23} = \mathbb{Z}$	$d_{23} = \beta^{-1}(-s_{11,1} + s_{11,2})$	
$H^{31} = \mathbb{Z}$	$d_{31} = \beta^{-1}s_{15,1}$	$= \pm \bar{s}_{4,2} d_{23}$

Step 2. The second column implies that $H^{even}(F_4/B_3)$ has additive basis of the form $p^* \Lambda$, with $\Lambda = \{s_{4,2}, s_{8,1}\}$ a subset of Schubert classes. The corresponding (4.13) consists of the single equation $\bar{s}_{8,1} = \bar{s}_{4,2}^2$ (cf. [DZ₂, 2.3]). These implies that, if we put $y_4 = s_{4,2}$, then

a) y_4 is the Schubert class whose Weyl coordinate is given as that as in the theorem by [DZ₂, 2.1];

b) the ring $H^{even}(F_4/B_3)$ is generated by \bar{y}_4 subject to the relation h_8 .

Combining a) and b) with the obvious relation $h_{12} : \bar{y}_4^3 = 0$ (in view of $H^{24} = 0$ by the first column), implies that

$$(5.2) \quad H^{even}(F_4/B_3) = \frac{\mathbb{Z}[\bar{y}_4]}{\langle h_8; h_{12} \rangle}.$$

Step 3. The proof is completed by $d_{31} = \pm \bar{s}_{4,2} d_{23}$ (Lemma 9) and $d_{23}^2 \in H^{46} = 0$ (in view of the first column of the table).

Remark 1. In the ring $\mathbb{Z}[\bar{y}_4]$ one has $\langle h_8; h_{12} \rangle = \langle h_8; 26\bar{y}_4^3 \rangle$

Theorem 9. Let $y_3, y_4, y_6 \in H^*(E_6/A_6 \cdot S^1)$ be the Schubert class with Weyl coordinates $\sigma[3, 4, 2]$, $\sigma[1, 3, 4, 2]$, $\sigma[1, 3, 6, 5, 4, 2]$ respectively, and let $d_{23}, d_{29} \in H^{odd}(E_6/A_6)$ be with

$$\beta(d_{23}) = 2s_{11,1} - s_{11,2}; \quad \beta(d_{29}) = s_{14,1} + s_{14,2} + s_{14,4} - s_{14,5}.$$

Then

$$H^*(E_6/A_6) = \left\{ \frac{\mathbb{Z}[\bar{y}_3, \bar{y}_4, \bar{y}_6]}{\langle h_6; h_8; h_9; h_{12} \rangle} \widehat{\otimes} \Gamma(1, d_{23}, d_{29}) \right\} / \langle 2d_{29} = \pm \bar{y}_3 d_{23} \rangle^1,$$

where $h_6 : 2\bar{y}_6 + \bar{y}_3^2 = 0$; $h_8 : 3\bar{y}_4^2 = 0$; $h_9 : 2\bar{y}_3\bar{y}_6 = 0$; $h_{12} : \bar{y}_6^2 - \bar{y}_4^3 = 0$.

Proof. Step 1. From the matrices A_k presented in [DZ₂, 3.2], one obtains the results in the first two columns of the table below.

nontrivial H^k	basis elements	relations
$H^6 = \mathbb{Z}$	$\bar{s}_{3,1}$	
$H^8 = \mathbb{Z}$	$\bar{s}_{4,1}$	
$H^{12} = \mathbb{Z}$	$\bar{s}_{6,1}$	$-2\bar{s}_{6,1} = \bar{s}_{3,1}^2$
$H^{14} = \mathbb{Z}$	$\bar{s}_{7,1}$	$\bar{s}_{3,1}\bar{s}_{4,1}$
$H^{16} = \mathbb{Z}_3$	$\bar{s}_{8,1}$	$\bar{s}_{4,1}^2$
$H^{18} = \mathbb{Z}_2$	$\bar{s}_{9,1}$	$\bar{s}_{3,1}\bar{s}_{6,1}$
$H^{20} = \mathbb{Z}$	$\bar{s}_{10,1}$	$-\bar{s}_{4,1}\bar{s}_{6,1}$
$H^{22} = \mathbb{Z}_3$	$\bar{s}_{11,1}$	$-\bar{s}_{3,1}^2\bar{s}_{4,1}$
$H^{26} = \mathbb{Z}_2$	$\bar{s}_{13,2}$	$-\bar{s}_{3,1}\bar{s}_{4,1}\bar{s}_{6,1}$
$H^{28} = \mathbb{Z}_3$	$\bar{s}_{14,1}$	$\bar{s}_{4,1}^2\bar{s}_{6,1}$
$H^{23} = \mathbb{Z}$	$d_{23} = \beta^{-1}(s_{11,1} - s_{11,2} - s_{11,3} + s_{11,4} - s_{11,5} + s_{11,6})$	
$H^{29} = \mathbb{Z}$	$d_{29} = \beta^{-1}(s_{14,1} + s_{14,2} + s_{14,4} - s_{14,5})$	$2d_{29} = \pm \bar{s}_{3,1} d_{23}$
$H^{31} = \mathbb{Z}$	$d_{31} = \beta^{-1}(s_{15,1} - 2s_{15,2} + s_{15,3} - s_{15,4})$	$\pm \bar{s}_{4,1} d_{23}$
$H^{35} = \mathbb{Z}$	$d_{35} = \beta^{-1}(s_{17,1} - s_{17,2} - s_{17,3})$	$\pm \bar{s}_{6,1} d_{23}$
$H^{37} = \mathbb{Z}$	$d_{37} = \beta^{-1}(s_{18,1} - s_{18,2})$	$\pm \bar{s}_{4,1} d_{29}$
$H^{43} = \mathbb{Z}$	$d_{43} = \beta^{-1}(s_{22,1})$	$\pm \bar{s}_{4,1}\bar{s}_{6,1} d_{23}$

Step 2. From the second column of the table one finds that an additive basis of $H^{even}(E_6/A_6)$ is given as $p^*\Lambda$, where

$$\Lambda = \{s_{3,1}, s_{4,1}, s_{6,1}, s_{7,1}, s_{8,1}, s_{9,1}, s_{10,1}, s_{11,1}, s_{13,1}, s_{14,1}\}$$

consists of Schubert classes. With the multiplicative rule (4.13) for the basis elements being determined in [DZ₂, 3.3], the items in the last column

¹The choice in the sign \pm appearing in the relation $2d_{29} = \pm \bar{y}_3 d_{23}$ does not effect the ring structure.

corresponding to H^{even} are verified. These imply that, if we put $y_3 = s_{3,1}, y_4 = s_{4,1}, y_6 = s_{6,1}$, then

a) y_3, y_4, y_6 are the Schubert classes whose Weyl coordinates are as that as given in the theorem by [DZ₂, 3.1];

b) $H^{even}(E_6/A_6)$ is generated by $\bar{y}_3, \bar{y}_4, \bar{y}_6$ subject to the relations $h_6; h_8; h_9$ (cf. the theorem).

Combining these with the obvious relations $\bar{y}_6^2 = \bar{y}_4^3 = 0$ (because of $H^{24} = 0$ by the first column), together with the fact that in $\mathbb{Z}[\bar{y}_3, \bar{y}_4, \bar{y}_6]$

$$\langle h_6; h_8; h_9; \bar{y}_6^2, \bar{y}_4^3 \rangle = \langle h_6; h_8; h_9; h_{12} \rangle,$$

one obtains

$$(5.3) \quad H^{even}(E_6/A_6) = \frac{\mathbb{Z}[\bar{y}_3, \bar{y}_4, \bar{y}_6]}{\langle h_6; h_8; h_9, h_{12} \rangle}.$$

Step 3. In view of the second column of the table, the $H^{2k+1} = \mathbb{Z}$ is generated by the d_{2k+1} for $k = 23, 29, 31, 35, 37, 43$. According to the first column of the table, we have also

$$d_{2k+1}d_{2k'+1} \in H^{2(k+k'+1)} = 0, \text{ for all } k, k' = 23, 29, 31, 35, 37, 43.$$

Further, we may assume, for the degree reasons, that

$$\bar{s}_{3,1}d_{23} = a_1d_{29}; \quad \bar{s}_{4,1}d_{23} = a_2d_{31}; \quad \bar{s}_{6,1}d_{23} = a_3d_{35}; \quad \bar{s}_{4,1}d_{29} = a_4d_{37};$$

Lemma 9 suffices to determine the $a_i \in \mathbb{Z}$ up to sign. For instance, applying it to the pairings $H^{20} \times H^{23} \rightarrow H^{43}$, $H^{14} \times H^{29} \rightarrow H^{43}$ yield respectively that

$$d_{43} = \pm \bar{s}_{4,1}\bar{s}_{6,1}d_{23}, \quad d_{43} = \pm \bar{s}_{3,1}\bar{s}_{4,1}d_{29}.$$

These implies that

$$\begin{aligned} \bar{s}_{4,1}\bar{s}_{6,1}d_{23} &= \pm \bar{s}_{3,1}\bar{s}_{4,1}d_{29} \\ &= \pm a_1^{-1}\bar{s}_{3,1}^2\bar{s}_{4,1}d_{23} \text{ (by the assumption } \bar{s}_{3,1}d_{23} = a_1d_{29}) \\ &= \pm 2a_1^{-1}\bar{s}_{4,1}\bar{s}_{6,1}d_{23} \text{ (by } h_6). \end{aligned}$$

Coefficients comparison gives $a_1 = \pm 2$.

Similarly, applying Lemma 9 to the pairings $H^{20} \times H^{23} \rightarrow H^{43}$, $H^{12} \times H^{31} \rightarrow H^{43}$ yield respectively that $d_{43} = \pm \bar{s}_{4,1}\bar{s}_{6,1}d_{23}$, $d_{43} = \pm \bar{s}_{6,1}d_{31}$. These implies that

$$\bar{s}_{6,1}d_{31} = \pm \bar{s}_{4,1}\bar{s}_{6,1}d_{23} = \pm a_2\bar{s}_{6,1}d_{31}$$

(by the assumption $\bar{s}_{4,1}d_{23} = a_2d_{31}$). Coefficients comparison gives $a_2 = \pm 1$.

The same method is applicable to show $a_i = \pm 1$, $i = 3, 4$. These verify the items in the third column of the table corresponding to H^{odd} , and therefore, completes the proof of Theorem 9.

Theorem 10. Let $y_4 \in H^*(E_6/D_5 \cdot S^1)$ be the Schubert class with Weyl coordinate $\sigma[2, 4, 5, 6]$, and let $d_{17} \in H^{odd}(E_6/D_5)$ be with

$$\beta(d_{17}) = s_{8,1} - s_{8,2} - s_{8,3}.$$

Then

$$H^*(E_6/D_5) = \frac{\mathbb{Z}[\bar{y}_4]}{\langle h_{12} \rangle} \widehat{\otimes} \Gamma(1, d_{17}),$$

where $h_{12} = \bar{y}_4^3$.

Proof. Step 1. With the matrices A_k being presented in [DZ₂, 4.2], one obtains the results in the first two columns of the table:

nontrivial H^k	basis elements	relations
$H^8 = \mathbb{Z}$	$\bar{s}_{4,1}$	
$H^{16} = \mathbb{Z}$	$\bar{s}_{8,1}$	$\bar{s}_{4,1}^2$
$H^{17} = \mathbb{Z}$	$d_{17} = \beta^{-1}(s_{8,1} - s_{8,2} - s_{8,3})$	
$H^{25} = \mathbb{Z}$	$d_{25} = \beta^{-1}(s_{12,1} - s_{12,2})$	$\pm \bar{s}_{4,1} d_{17}$
$H^{33} = \mathbb{Z}$	$d_{33} = \beta^{-1}(s_{16,1})$	$\pm \bar{s}_{4,1}^2 d_{17}$

Step 2. From the second column one finds that an additive basis of $H^{even}(E_6/D_5)$ is given as $p^*\Lambda$, where $\Lambda = \{s_{4,1}, s_{8,1}\}$ is a subset of Schubert classes. The multiplicative rule (4.13) of this basis elements consists the single equation $\bar{s}_{8,1} = \bar{s}_{4,1}^2$ by [DZ₂, 4.3]. These imply that, if we let $y_4 = s_{4,1}$, then

a) y_4 is the Schubert classes whose Weyl coordinates is as that as given in the theorem by [DZ₂, 4.1];

b) $H^{even}(E_6/D_5)$ is generated by \bar{y}_4 subject to the relation $h_{12} : \bar{y}_4^3 = 0$ (because of $H^{24} = 0$ by the first column).

As a result,

$$(5.4) \quad H^{even}(E_6/D_5) = \frac{\mathbb{Z}[\bar{y}_4]}{\langle h_{12} \rangle}.$$

Step 3. Since $H^{25} = \mathbb{Z}$ is generated d_{25} , $\bar{s}_{4,1}d_{17} = ad_{25}$ for some $a \in \mathbb{Z}$ for the degree reason. Applying Lemma 9 to the pairings $H^8 \times H^{25} \rightarrow H^{33}$, $H^{16} \times H^{17} \rightarrow H^{33}$ yield respectively that

$$d_{33} = \pm \bar{s}_{4,1}d_{25}, \quad d_{33} = \pm \bar{s}_{4,1}^2d_{17}.$$

These imply that $a = \pm 1$. The proof is completed by $d_{25}^2 \in H^{50} = 0$ according to the first column of the table.

Theorem 11. Let $y_5, y_9 \in H^*(E_7/E_6 \cdot S^1)$ be the Schubert class with Weyl coordinates $\sigma[2, 4, 5, 6, 7]$, $\sigma[1, 5, 4, 2, 3, 4, 5, 6, 7]$ respectively, and let $d_{37}, d_{45} \in H^{odd}(E_7/E_6)$ be with

$$\beta(d_{37}) = s_{18,1} - s_{18,2} + s_{18,3}, \quad \beta(d_{45}) = s_{22,1} - s_{22,2}.$$

Then

$$H^*(E_7/E_6) = \left\{ \frac{\mathbb{Z}[\bar{y}_5, \bar{y}_9]}{\langle h_{10}; h_{14}; h_{18} \rangle} \widehat{\otimes} \Gamma(1, d_{37}, d_{45}) \right\} / \langle \bar{y}_9 d_{37} = \pm \bar{y}_5 d_{45} \rangle^2,$$

where $h_{10} : \bar{y}_5^2 = 0$; $h_{14} : 2\bar{y}_5\bar{y}_9 = 0$; $h_{18} : \bar{y}_9^2 = 0$.

Proof. Step 1. With the matrices A_k being presented in [DZ₂, 5.2], one obtains the results in the first two columns of the table:

²See the footnote in Theorem 10.

nontrivial H^k	basis elements	relations
$H^{10} = \mathbb{Z}$	$\bar{s}_{5,1}$	$\bar{s}_{5,1}$
$H^{18} = \mathbb{Z}$	$\bar{s}_{9,1}$	$\bar{s}_{9,1}$
$H^{28} = \mathbb{Z}_2$	$\bar{s}_{14,1}$	$\bar{s}_{5,1}\bar{s}_{9,1}$
$H^{37} = \mathbb{Z}$	$d_{37} = \beta^{-1}(s_{18,1} - s_{18,2} + s_{18,3})$	
$H^{45} = \mathbb{Z}$	$d_{45} = \beta^{-1}(s_{22,1} - s_{22,2})$	
$H^{55} = \mathbb{Z}$	$d_{55} = \beta^{-1}(s_{27,1})$	$\bar{s}_{9,1}d_{37} = \pm\bar{s}_{5,1}d_{45}$

Step 2. By the second column of the table, a basis of $H^{even}(E_7/E_6)$ is given as $p^*\Lambda$, where $\Lambda = \{s_{5,1}, s_{9,1}, s_{14,1}\}$ is a subset of Schubert classes. The multiplicative rule (4.13) of the basis elements consists of the single equation $\bar{s}_{14,1} = \bar{s}_{5,1}\bar{s}_{9,1}$ by [DZ₂, 5.3]. These imply that, if we put $y_5 = s_{5,1}$, $y_9 = s_{9,1}$, then

a) y_4, y_9 are the Schubert classes whose Weyl coordinates are as that as given in the theorem by [DZ₂, 5.1];

b) $H^{even}(E_7/E_6)$ is generated by \bar{y}_4, \bar{y}_9 subject to the relation h_{14} .

Combining these with the obvious relations h_{10}, h_{18} (because of $H^{20} = H^{36} = 0$ by the first column) one obtains

$$(5.5) \quad H^{even}(E_7/E_6) = \frac{\mathbb{Z}[\bar{y}_5, \bar{y}_9]}{\langle h_{10}; h_{14}; h_{18} \rangle}.$$

Step 3. Applying Lemma 9 to the pairing $H^{10} \times H^{45} \rightarrow H^{55}$, $H^{18} \times H^{37} \rightarrow H^{55}$ yields $\bar{s}_{9,1}d_{37} = \pm\bar{s}_{5,1}d_{45}$. One has also $d_{37}^2 = d_{45}^2 = 0$ because of $H^{74} = H^{90} = 0$ (cf. in the first column of the table).

Theorem 12. Let $y_4, y_6, y_9 \in H^*(E_7/D_6 \cdot S^1)$ be the Schubert class with Weyl coordinates $\sigma[2, 4, 3, 1]$, $\sigma[2, 6, 5, 4, 3, 1]$, $\sigma[3, 4, 2, 7, 6, 5, 4, 3, 1]$ respectively, and let $d_{35}, d_{51} \in H^{odd}(E_7/D_6)$ be with

$$\beta(d_{35}) = s_{17,1} - s_{17,2} - s_{17,3} + s_{17,4} - s_{17,5} + s_{17,6} - s_{17,7};$$

$$\beta(d_{51}) = s_{25,1} - s_{25,2} - s_{25,4}.$$

Then

$$H^*(E_7/D_6) = \left\{ \frac{\mathbb{Z}[\bar{y}_4, \bar{y}_6, \bar{y}_9]}{\langle h_9; h_{12}; h_{14}; h_{18} \rangle} \widehat{\otimes} \Gamma(1, d_{35}, d_{51}) \right\} / \langle 3d_{51} = \pm\bar{y}_4^2 d_{35} \rangle^3,$$

where $h_9 : 2\bar{y}_9 = 0$; $h_{12} : 3\bar{y}_6^2 - \bar{y}_4^3 = 0$; $h_{14} : 3\bar{y}_4^2\bar{y}_6 = 0$; $h_{18} : \bar{y}_9^2 - \bar{y}_6^3 = 0$.

Proof. Step 1. The matrices A_k presented in [DZ₂, 6.2] yield the results in the first two columns of the table below:

³See the footnote in Theorem 10.

nontrivial H^k	basis elements	relations
$H^8 = \mathbb{Z}$	$\bar{s}_{4,1}$	
$H^{12} = \mathbb{Z}$	$\bar{s}_{6,1}$	
$H^{16} = \mathbb{Z}$	$\bar{s}_{8,1}$	$\bar{s}_{4,1}^2$
$H^{18} = \mathbb{Z}_2$	$\bar{s}_{9,2}$	
$H^{20} = \mathbb{Z}$	$\bar{s}_{10,1}$	$\bar{s}_{4,1}\bar{s}_{6,1}$
$H^{24} = \mathbb{Z}$	$\bar{s}_{12,2}$	$\bar{s}_{12,2} = \bar{s}_{6,1}^2;$ $3\bar{s}_{12,2} = \bar{s}_{4,1}^3$
$H^{26} = \mathbb{Z}_2$	$\bar{s}_{13,1}$	$\bar{s}_{4,1}\bar{s}_{9,2}$
$H^{28} = \mathbb{Z}_3$	$\bar{s}_{14,1}$	$-\bar{s}_{4,1}^2\bar{s}_{6,1}$
$H^{30} = \mathbb{Z}_2$	$\bar{s}_{15,1}$	$\bar{s}_{6,1}\bar{s}_{9,2}$
$H^{32} = \mathbb{Z}$	$\bar{s}_{16,1}$	$\bar{s}_{4,1}\bar{s}_{6,1}^2$
$H^{34} = \mathbb{Z}_2$	$\bar{s}_{17,2}$	$\bar{s}_{4,1}^2\bar{s}_{9,2}$
$H^{38} = \mathbb{Z}_2$	$\bar{s}_{19,2}$	$\bar{s}_{4,1}\bar{s}_{6,1}\bar{s}_{9,2}$
$H^{40} = \mathbb{Z}_3$	$\bar{s}_{20,1}$	$\bar{s}_{4,1}^2\bar{s}_{6,1}^2$
$H^{42} = \mathbb{Z}_2$	$\bar{s}_{21,3}$	$\bar{s}_{4,1}^3\bar{s}_{9,2}$
$H^{50} = \mathbb{Z}_2$	$\bar{s}_{25,1}$	$\bar{s}_{4,1}^4\bar{s}_{9,2}$
$H^{35} = \mathbb{Z}$	$d_{35} = \beta^{-1}(s_{17,1} - s_{17,2} - s_{17,3}$ $+s_{17,4} - s_{17,5} + s_{17,6} - s_{17,7})$	
$H^{43} = \mathbb{Z}$	$\beta^{-1}(s_{21,1} - 2s_{21,2} + s_{21,3}$ $-3s_{21,4} + 2s_{21,5} - s_{21,6})$	$\pm\bar{s}_{4,1}d_{35}$
$H^{47} = \mathbb{Z}$	$\beta^{-1}(2s_{23,1} - s_{23,2} + s_{23,3}$ $-s_{23,4} + s_{23,5})$	$\pm\bar{s}_{6,1}d_{35}$
$H^{51} = \mathbb{Z}$	$d_{51} = \beta^{-1}(s_{25,1} - s_{25,2} - s_{25,4})$	$3d_{51} = \pm\bar{s}_{4,1}^2d_{35}$
$H^{55} = \mathbb{Z}$	$\beta^{-1}(s_{27,1} + s_{27,2} - s_{27,3})$	$\pm\bar{s}_{4,1}\bar{s}_{6,1}d_{35}$
$H^{59} = \mathbb{Z}$	$\beta^{-1}(s_{29,1} - s_{29,2})$	$\pm\bar{s}_{6,1}^2d_{35}, \pm\bar{s}_{4,1}d_{51}$
$H^{67} = \mathbb{Z}$	$\beta^{-1}(s_{33,1})$	$\bar{s}_{4,1}\bar{s}_{6,1}^2d_{35} = \pm\bar{s}_{4,1}^2d_{51}$

Step 2. By results in the second column of the table, an additive basis of $H^{even}(E_7/D_6)$ is given as $p^*\Lambda$, where

$$\Lambda = \{s_{4,1}, s_{6,1}, s_{8,1}, s_{9,2}, s_{10,1}, s_{12,2}, s_{13,1}, s_{14,1}, s_{15,1}, s_{16,1}, \\ s_{17,2}, s_{19,2}, s_{20,1}, s_{21,3}, s_{25,1}\}$$

is a subset of Schubert classes. With the multiplicative rule (4.13) for the basis elements being determined in [DZ₂, 6.3], the items in the last column corresponds to H^{even} are verified. These imply that, if we put $y_4 = s_{4,1}$, $y_6 = s_{6,1}$, $y_9 = s_{9,2}$, then

a) y_4, y_6, y_9 are Schubert classes whose Weyl coordinates are as that as given in the theorem by [DZ₂, 6.1];

b) $H^{even}(E_7/D_6)$ is generated by $\bar{y}_4, \bar{y}_6, \bar{y}_9$ subject to the relations $h_9; h_{12}; h_{14}$ (cf. the theorem).

Combining these with the obvious relations $\bar{y}_9^2 = \bar{y}_6^3 = 0$ (because of $H^{36} = 0$ by the first column), together with the fact that, as ideals in $\mathbb{Z}[\bar{y}_4, \bar{y}_6, \bar{y}_9]$,

$$\langle h_9; h_{12}; h_{14}; \bar{y}_9^2, \bar{y}_6^3 \rangle = \langle h_9; h_{12}; h_{14}; h_{12} \rangle,$$

one obtains

$$(5.6) \quad H^{even}(E_7/D_6) = \frac{\mathbb{Z}[\bar{y}_4, \bar{y}_6, \bar{y}_9]}{\langle h_9; h_{12}; h_{14}; h_{18} \rangle}.$$

Step 3. The same method as that in Step 3 in the proof of Theorem 9 verifies the items in the last column of the table corresponding to H^{odd} . We omit the details.

Remark 2. Let $h_9; h_{12}; h_{14}; h_{18}$ be the polynomials in Theorem 12. It can be shown that, as ideals in $\mathbb{Z}[\bar{y}_4, \bar{y}_6, \bar{y}_9]$,

$$\langle h_9; h_{12}; h_{14}; h_{18} \rangle = \langle h_9; h_{12}; h_{14}; 5\bar{y}_9^2 + 29\bar{y}_6^3 \rangle.$$

6 Proofs of Theorem 1-6

Since a Grassmannian G/H is naturally a flag variety, its integral cohomology $H^*(G/H)$ can be identified with the Chow ring $A^*(G/H)$. For this reason Lemma 7 and 8 are directly applicable in the proofs of Theorem 1–6.

Proof of Theorem 1. Combining Theorem 7 with Lemma 7 and 8, we get the partial description for $A^*(F_4/C_3 \cdot S^1)$ as

$$(6.1) \quad A^*(F_4/C_3 \cdot S^1) = \mathbb{Z}[y_1, y_3, y_4, y_6] / \langle r_3, r_6, r_8, r_{12}, y_1 g_{11} \rangle,$$

where y_1, y_3, y_4, y_6 are the Schubert classes as asserted by the theorem, and where if we let

$$\pi_m : \mathbb{Z}[y_1, y_3, y_4, y_6]^{(2m)} \rightarrow A^{2m}(F_4/C_3 \cdot S^1)$$

be the map induced by the inclusion $\{y_1, y_3, y_4, y_6\} \subset A^*(F_4/C_3 \cdot S^1)$ (cf. **3.3**), then

- 1) for $m = 3, 6, 8, 12$, $r_m \in \ker \pi_m$ with

$$\begin{aligned} r_3 \mid_{y_1=0} &= 2y_3; & r_6 \mid_{y_1=0} &= 2y_6 + y_3^2; \\ r_8 \mid_{y_1=0} &= 3y_4^2; & r_{12} \mid_{y_1=0} &= y_6^2 - y_4^3; \end{aligned}$$
- 2) $\pi(g_{11}) = 2s_{11,1} - s_{11,2}$.

With respect to the ordered basis $B(2m)$ of $\mathbb{Z}[y_1, y_3, y_4, y_6]^{(2m)}$, $m = 3, 6, 8, 12$, the structure matrix $M(\pi_m)$ has been computed by the *L–R coefficients* and presented in [DZ₂, 1.4]. Applying the *Nullspace* in *Mathematica* yield respectively that (cf. [DZ₂, 1.5])

$$N(\pi_3) = (-2, 1), \quad N(\pi_6) = \begin{pmatrix} 0 & -4 & 0 & 0 & 1 \\ 0 & -2 & 0 & 1 & 0 \\ -2 & -1 & 3 & 0 & 0 \end{pmatrix},$$

see in [DZ₂, 1.5] for $N(\pi_8)$, $N(\pi_{12})$. If we take, in view of Lemma 5, that

$$\begin{aligned} r_3 &= 2y_3 - y_1^3 \quad (= -\kappa_1 \text{ in } N(\pi_3)); \\ r_6 &= 2y_6 + y_3^2 - 3y_1^2 y_4 \quad (= -\kappa_3 \text{ in } N(\pi_6)) \\ r_8 &= 3y_4^2 - y_1^2 y_6 \quad (= -\kappa_5 \text{ in } N(\pi_8)) \\ r_{12} &= y_6^2 - y_4^3 \quad (= \kappa_{15} \text{ in } N(\pi_{12})) \end{aligned}$$

then condition 1) is met by the set $\{r_3, r_6, r_8, r_{12}\}$ of polynomials.

The proof will be completed once we show

$$(6.2) \quad y_1 g_{11} \in \langle r_3, r_6, r_8, r_{12} \rangle.$$

For this purpose we examine, in view of (6.1), the quotient map (cf. **3.2**)

$$\varphi : \mathbb{Z}[y_1, y_3, y_4, y_6] / \langle r_3, r_6, r_8, r_{12} \rangle \rightarrow A^*(F_4/C_3 \cdot S^1) = \bigoplus_{m \geq 0} A^m.$$

With the r_3, r_6, r_8, r_{12} being obtained explicitly, it is straightforward to find that (cf. Example 3)

$$b(24) = 16; \quad \delta_{24}(r_3, r_6, r_8, r_{12}) = 15.$$

On the other hand, granted with the Basis Theorem, we read from [DZ₂, 1.1] that $\text{rank}(A^{24}) = 1$. (6.2) is verified by Lemma 4.

Proof of Theorem 2. Combining Theorem 8 with Lemma 7 and 8, we get the partial description of $A^*(F_4/B_3 \cdot S^1)$ as

$$(6.3) \quad A^*(F_4/B_3 \cdot S^1) = \mathbb{Z}[y_1, y_4] / \langle r_8, r_{12}, y_1 g_{11} \rangle,$$

where the generators y_1, y_4 are the Schubert classes as asserted in Theorem 1, and where if we let

$$\pi_m : \mathbb{Z}[y_1, y_4]^{(2m)} \rightarrow A^{2m}(F_4/B_3 \cdot S^1)$$

be the map induced by the inclusion $\{y_1, y_4\} \subset A^*(F_4/B_3 \cdot S^1)$ (cf. **3.3**), then

1) for $m = 8, 12$, $r_m \in \ker \pi_m$ with $r_8|_{y_1=0} = 3y_4^2$; $r_{12}|_{y_1=0} = 26y_4^3$ (cf. Remark 1 after the proof of Theorem 8);

2) $\pi(g_{11}) = -s_{11,1} + s_{11,2}$.

With respect to the ordered basis $B(2m)$ for $m = 8, 12$ of $\mathbb{Z}[y_1, y_4]^{(2m)}$, the structure matrix $M(\pi_m)$ has been computed by the *L-R coefficients* and are presented in [DZ₂, 2.4]. Applying the *Nullspace* in *Mathematica* yield respectively that (cf. [DZ₂, 2.5])

$$N(\pi_8) = \begin{pmatrix} -3 & 0 & 1 \end{pmatrix}, \quad N(\pi_{12}) = \begin{pmatrix} -26 & 0 & 0 & 5 \\ -3 & 0 & 1 & 0 \\ -26 & 15 & 0 & 0 \end{pmatrix}.$$

Therefore, if we take, in view of Lemma 5, that

$$\begin{aligned} r_8 &= 3y_4^2 - y_1^8 \quad (= -\kappa_1 \text{ in } N(\pi_8)), \\ r_{12} &= 26y_4^3 - 5y_1^{12} \quad (= -\kappa_1 \text{ in } N(\pi_{12})) \end{aligned}$$

then condition 1) is met by the set $\{r_8, r_{12}\}$ of polynomials.

The proof will be completed once we show

$$(6.4) \quad y_1 g_{11} \in \langle r_8, r_{12} \rangle.$$

For this purpose we examine, in view of (6.1), the quotient map (cf. **3.2**)

$$\varphi : \mathbb{Z}[y_1, y_4] / \langle r_8, r_{12} \rangle \rightarrow A^*(F_4/C_3 \cdot S^1) = \bigoplus_{m \geq 0} A^m.$$

With the r_8, r_{12} being obtained explicitly it is straightforward to find that (cf. Example 3)

$$b(24) = 4; \quad \delta_{24}(r_8, r_{12}) = 3.$$

On the other hand, granted with the Basis Theorem, we read from [DZ₂, 2.1] that $\text{rank}(A^{24}) = 1$. (6.4) is verified by Lemma 4.

Proof of Theorem 3. Combining Theorem 9 with Lemma 7 and 8, we get the partial description for $A^*(E_6/A_6 \cdot S^1)$ as

$$(6.5) \quad A^*(E_6/A_6 \cdot S^1) = \mathbb{Z}[y_1, y_3, y_4, y_6] / \langle r_6, r_8, r_9, r_{12}, y_1 g_{11}, y_1 g_{14} \rangle,$$

where the y_1, y_3, y_4, y_6 are the Schubert classes as asserted by the theorem, and where if we let

$$\pi_m : \mathbb{Z}[y_1, y_3, y_4, y_6]^{(2m)} \rightarrow A^{2m}(F_4/B_3 \cdot S^1)$$

be the map induced by $\{y_1, y_3, y_4, y_6\} \subset A^{2m}(F_4/B_3 \cdot S^1)$ (cf. **3.3**), then

- 1) for $m = 6, 8, 9, 12$, $r_m \in \ker \pi_m$ with

$$\begin{aligned} r_6 \big|_{y_1=0} &= 2y_6 + y_3^2; \quad r_8 \big|_{y_1=0} = 3y_4^2; \\ r_9 \big|_{y_1=0} &= 2y_3 y_6; \quad r_{12} \big|_{y_1=0} = y_4^3 - y_6^2; \end{aligned}$$
- 2) $\pi(g_{11}) = s_{11,1} - s_{11,2} - s_{11,3} + s_{11,4} - s_{11,5} + s_{11,6}$,
 $\pi(g_{14}) = s_{14,1} + s_{14,2} + s_{14,4} - s_{14,5}$.

With respect to the ordered basis $B(2m)$ of $\mathbb{Z}[y_1, y_3, y_4, y_6]$, $m = 6, 8, 9, 12$, the structure matrix $M(\pi_m)$ has been computed by the *L-R coefficients* and presented in [DZ₂, 3.4]. Applying the *Nullspace* in *Mathematica* yield respectively that (cf. [DZ₂, 3.5])

$$N(\pi_6) = \begin{pmatrix} -2 & -1 & 3 & -2 & 1 \end{pmatrix}, \quad N(\pi_8) = \begin{pmatrix} -3 & 6 & -3 & -6 & 3 & 0 & 1 \\ -3 & 6 & -1 & -5 & 0 & 2 & 0 \end{pmatrix},$$

see in [DZ₂, 3.5] for $N(\pi_9)$, $N(\pi_{12})$. If we take, in view of Lemma 5, that

$$\begin{aligned} r_6 &= 2y_6 + y_3^2 - 3y_1^2y_4 + 2y_1^3y_3 - y_1^6 \quad (= -\kappa_1 \text{ in } N(\pi_6)), \\ r_8 &= 3y_4^2 - 6y_1y_3y_4 + y_1^2y_6 + 5y_1^2y_3^2 - 2y_1^5y_3 \quad (= -\kappa_2 \text{ in } N(\pi_8)), \\ r_9 &= 2y_3y_6 - y_1^3y_6 \quad (= -\kappa_4 \text{ in } N(\pi_9)), \\ r_{12} &= y_4^3 - y_6^2 \quad (= \kappa_{11} \text{ in } N(\pi_{12})), \end{aligned}$$

then condition 1) is satisfied by the set $\{r_6, r_8, r_9, r_{12}\}$ of polynomials.

The proof will be completed once we show

$$(6.6) \quad y_1g_{11}, y_1g_{14} \in \langle r_6, r_8, r_9, r_{12} \rangle.$$

For this purpose we examine, in view of (6.5), the quotient map (cf. **3.2**)

$$\varphi : \mathbb{Z}[y_1, y_3, y_4, y_6] / \langle r_6, r_8, r_9, r_{12} \rangle \rightarrow A^*(E_6/A_6 \cdot S^1) = \bigoplus_{m \geq 0} A^m$$

With the r_6, r_8, r_9, r_{12} being obtained explicitly, it is straightforward to find that (cf. Example 3)

$$\begin{aligned} b(24) &= 16; \quad \delta_{24}(r_6, r_8, r_9, r_{12}) = 11; \\ b(30) &= 24; \quad \delta_{30}(r_6, r_8, r_9, r_{12}) = 20. \end{aligned}$$

On the other hand, granted with the Basis Theorem, we read from [DZ₂, 3.1] that $\text{rank}(A^{24}) = 5$, $\text{rank}(A^{30}) = 4$. (6.6) is verified by Lemma 4.

Proof of Theorem 4. Combining Theorem 10 with Lemma 7 and 8, we get the partial description for $A^*(E_6/D_5 \cdot S^1)$ as

$$(6.7) \quad A^*(E_6/D_5 \cdot S^1) = \mathbb{Z}[y_1, y_4] / \langle y_1g_8, r_{12}, \rangle$$

where y_1, y_4 are the Schubert classes as asserted by the Theorem, and where

- 1) $\pi(r_{12}) = 0$ with $r_{12}|_{y_1=0} = y_4^3$;
- 2) $\pi(g_8) = s_{8,1} - s_{8,2} - s_{8,3}$.

Let us find the $g_8 \in \mathbb{Z}[y_1, y_4]$ required to specify the first relation y_1g_8 . Assume, with respect to the basis $B(16)$ of $\mathbb{Z}[y_1, y_4]^{(16)}$, that

$$(6.8) \quad g_8 = a_1y_1^8 + a_2y_1^4y_4 + a_3y_4^2.$$

According to [DZ₂, 4.1] there are three Schubert classes in dimension 16, whose Weyl coordinates are respectively

$$\begin{aligned} w_{8,1} &= \sigma[1, 5, 4, 2, 3, 4, 5, 6]; \quad w_{8,2} = \sigma[3, 1, 4, 2, 3, 4, 5, 6]; \\ w_{8,3} &= \sigma[6, 5, 4, 2, 3, 4, 5, 6]. \end{aligned}$$

The constraint 2) implies that the g_8 must satisfy the system

$$T_{w_{8,1}}(g_8) = 1; \quad T_{w_{8,2}}(g_8) = -1; \quad T_{w_{8,3}}(g_8) = 1.$$

Thus, applying the *L-R Coefficients* (cf. **2.3**) to (6.8) yields

$$\begin{cases} 1 = 7a_1 + 3a_2 + a_3 \\ -1 = 5a_1 + 2a_2 + a_3 \\ -1 = 2a_1 + a_2 + a_3. \end{cases}$$

From this we find that $(a_1, a_2, a_3) = (-2, 6, -3)$, and consequently

$$y_1 g_8 = 2y_1^9 + 3y_1 y_4^2 - 6y_1^5 y_4 \text{ (cf. Theorem 4).}$$

To find r_{12} we consider the map

$$\pi_{12} : \mathbb{Z}[y_1, y_4]^{(24)} \rightarrow A^{24}(E_6/D_5 \cdot S^1)$$

induced by $\{y_1, y_4\} \subset A^*(E_6/D_5 \cdot S^1)$ (cf. **3.3**). With respect to the ordered basis $B(24)$ of $\mathbb{Z}[y_1, y_4]^{(24)}$, the structure matrix $M(\pi_{12})$ is presented in [DZ₂, 4.4]. Applying the *Nullspace* in *Mathematica* yields that (cf. [DZ₂, 4.5])

$$N(\pi_{12}) = \begin{pmatrix} 1 & -6 & 0 & 1 \\ 2 & -15 & 6 & 0 \end{pmatrix},$$

If we take, in view of Lemma 5, that

$$r_{12} = y_4^3 - 6y_1^4 y_4^2 + y_1^{12} \text{ (= } \kappa_1 \text{ in } N(\pi_{12})),$$

then condition 1) is satisfied by r_{12} . This finishes the proof.

Proof of Theorem 5. Combining Theorem 11 with Lemma 7 and 8, we get the partial description of $A^*(E_7/E_6 \cdot S^1)$ as

$$(6.9) \quad A^*(E_7/E_6 \cdot S^1) = \mathbb{Z}[y_1, y_5, y_9] / \langle r_{10}, r_{14}, r_{18}, y_1 g_{18}, y_1 g_{22} \rangle,$$

where y_1, y_5, y_9 are the Schubert classes as asserted by the theorem, and where if we let

$$\pi_m : \mathbb{Z}[y_1, y_5, y_9]^{(2m)} \rightarrow A^{2m}(F_4/B_3 \cdot S^1)$$

be induced by $\{y_1, y_5, y_9\} \subset A^*(E_7/E_6 \cdot S^1)$ (cf. **3.3**), then

1) For $m = 10, 14, 18$, $r_m \in \ker \pi_m$ and

$$r_{10} \mid_{y_1=0} = y_5^2; \quad r_{14} \mid_{y_1=0} = 2y_5 y_9; \quad r_{18} \mid_{y_1=0} = y_9^2;$$

2) $\pi(g_{18}) = s_{18,1} - s_{18,2} + s_{18,3}$, $\pi(g_{22}) = s_{22,1} - s_{22,2}$.

With respect to the ordered basis $B(2m)$ of $\mathbb{Z}[y_1, y_5, y_9]^{(2m)}$, $m = 10, 14, 18$, the structure matrix $M(\pi_m)$ has been computed by the *L-R coefficients* and presented in [DZ₂, 5.4]. Applying the *Nullspace* in *Mathematica* yield respectively that (cf. [DZ₂, 5.5])

$$N(\pi_{10}) = \begin{pmatrix} -1 & 2 & 0 & 0 \end{pmatrix}, \quad N(\pi_{14}) = \begin{pmatrix} -2 & 9 & 0 & -6 & 1 \\ 0 & -1 & 2 & 0 & 0 \end{pmatrix},$$

see in [DZ₂, 5.5] for $N(\pi_{18})$. If we take, in view of Lemma 5, that

$$r_{10} = y_5^2 - 2y_1 y_9 \text{ (= } -\kappa_1 \text{ in } N(\pi_{10}));$$

$$r_{14} = 2y_5 y_9 - 9y_1^4 y_5^2 + 6y_1^9 y_5 - y_1^{14} \text{ (= } -\kappa_1 \text{ in } N(\pi_{14}));$$

$$r_{18} = y_9^2 + 10y_1^3 y_5^3 - 9y_1^8 y_5^2 + 2y_1^{13} y_5 \text{ (= } \kappa_2 \text{ in } N(\pi_{18})).$$

then condition 1) is satisfied by the set $\{r_{10}, r_{14}, r_{18}\}$ of polynomials.

The proof will be completed once we show

$$(6.10) \quad y_1 g_{18}, y_1 g_{22} \in \langle r_{10}, r_{14}, r_{18} \rangle.$$

For this purpose we examine, in view of (6.9), the quotient map (cf. **3.2**)

$$\varphi : \mathbb{Z}[y_1, y_5, y_9] / \langle r_{10}, r_{14}, r_{18} \rangle \rightarrow A^*(E_7/E_6 \cdot S^1) = \bigoplus_{m \geq 0} A^m$$

With the r_{10}, r_{14}, r_{18} being obtained explicitly, it is straightforward to find that (cf. Example 3)

$$\begin{aligned} b(38) &= 8; \quad \delta_{38}(r_{10}, r_{14}, r_{18}) = 6; \\ b(46) &= 10; \quad \delta_{46}(r_{10}, r_{14}, r_{18}) = 9. \end{aligned}$$

On the other hand, granted with the Basis Theorem, we read from [DZ₂, 5.1] that $\text{rank}(A^{38}) = 2$, $\text{rank}(A^{46}) = 1$. (6.10) is verified by Lemma 4.

Proof of Theorem 6. Combining Theorem 12 with Lemma 7 and 8, we get the partial description of $A^*(E_7/E_6 \cdot S^1)$ as

$$(6.11) \quad A^*(E_7/D_6 \cdot S^1) = \mathbb{Z}[y_1, y_4, y_6, y_9] / \langle r_9, r_{12}, r_{14}, r_{18}, y_1 g_{17}, y_1 g_{25} \rangle,$$

where y_1, y_4, y_6, y_9 are the Schubert classes as asserted in Theorem 6, and where if we let

$$\pi_m : \mathbb{Z}[y_1, y_4, y_6, y_9]^{(2m)} \rightarrow A^{2m}(E_7/D_6 \cdot S^1)$$

induced by $\{y_1, y_4, y_6, y_9\} \subset A^*(E_7/D_6 \cdot S^1)$ (cf. **3.3**), then

- 1) for $m = 9, 12, 14, 18$, $r_m \in \ker \pi_m$ with

$$\begin{aligned} r_9 \big|_{y_1=0} &= 2y_9; & r_{12} \big|_{y_1=0} &= 3y_6^2 - y_4^3; & r_{14} \big|_{y_1=0} &= 3y_4^2 y_6; \\ r_{18} \big|_{y_1=0} &= 5y_9^2 + 29y_6^3 \quad (\text{cf. Remark 2 after the proof of Theorem 12}) \end{aligned}$$
- 2) $\pi(g_{17}) = s_{17,1} - s_{17,2} - s_{17,3} + s_{17,4} - s_{17,5} + s_{17,6} - s_{17,7}$,
 $\pi(g_{25}) = s_{25,1} - s_{25,2} - s_{25,4}$.

With respect to the ordered basis $B(2m)$ of $\mathbb{Z}[y_1, y_4, y_6, y_9]^{(2m)}$, $m = 9, 12, 14, 18$, the structure matrix $M(\pi_m)$ has been computed by the L - R coefficients and presented in [DZ₂, 6.4]. Applying the Nullspace in *Mathematica* yield respectively that (cf. [DZ₂, 6.5])

$$\begin{aligned} N(\pi_9) &= \begin{pmatrix} -2 & -3 & -4 & -2 & 2 \end{pmatrix}; \\ N(\pi_{12}) &= \begin{pmatrix} 3 & -1 & 0 & -2 & -6 & -6 & 0 & 2 \\ 3 & -1 & 0 & 0 & -3 & -2 & 2 & 0 \end{pmatrix}; \end{aligned}$$

see in [DZ₂, 6.5] for $N(\pi_{14})$, $N(\pi_{18})$. If we take, in view of Lemma 5, that

$$\begin{aligned} r_9 &= 2y_9 + 3y_1 y_4^2 + 4y_1^3 y_6 + 2y_1^5 y_4 - 2y_1^9 \quad (= -\kappa_1 \text{ in } N(\pi_9)); \\ r_{12} &= 3y_6^2 - y_4^3 - 3y_1^4 y_4^2 - 2y_1^6 y_6 + 2y_1^8 y_4 \quad (= \kappa_1 \text{ in } N(\pi_{12})); \\ r_{14} &= 3y_4^2 y_6 + 3y_1^2 y_6^2 + 6y_1^2 y_4^3 + 6y_1^4 y_4 y_6 + 2y_1^5 y_9 - y_1^{14} \quad (= -\kappa_1 \text{ in } N(\pi_{14})); \\ r_{18} &= 5y_9^2 + 29y_6^3 - 24y_1^6 y_6^2 + 45y_1^2 y_4 y_6^2 + 2y_1^9 y_9 \quad (= \kappa_5 - 2\kappa_8 \text{ in } N(\pi_{18})). \end{aligned}$$

then condition 1) is satisfied by the set $\{r_9, r_{12}, r_{14}, r_{18}\}$ of polynomials.

The proof will be completed once we show

$$(6.12) \quad y_{19}y_{17}, y_{19}y_{25} \in \langle r_9, r_{12}, r_{14}, r_{18} \rangle.$$

For this purpose we examine, in view of (6.11), the quotient map (cf. **3.2**)

$$\varphi : \mathbb{Z}[y_1, y_4, y_6, y_9] / \langle r_9, r_{12}, r_{14}, r_{18} \rangle \rightarrow A^*(E_7/D_6 \cdot S^1) = \bigoplus_{m \geq 0} A^m$$

With the r_{10}, r_{14}, r_{18} being obtained explicitly, it is straightforward to find that (cf. Example 3)

$$\begin{aligned} b(36) &= 17; \quad \delta_{36}(r_9, r_{12}, r_{14}, r_{18}) = 11; \\ b(52) &= 32; \quad \delta_{52}(r_9, r_{12}, r_{14}, r_{18}) = 29. \end{aligned}$$

On the other hand, granted with the Basis Theorem, we read from [DZ₂, 6.1] that $\text{rank}(A^{36}) = 6$, $\text{rank}(A^{52}) = 3$. (6.12) is verified by Lemma 4.

7 Historical remarks

In [Co, 1964] Conlon computed the ring $H^*(E_6/D_5)$ as well as the additive homology of $E_6/D_5 \cdot S^1$. His method amounts to apply Morse theory to the space $\Omega(E_6/D_5 \cdot S^1, x, W)$ of paths to yield a cell decomposition of $E_6/D_5 \cdot S^1$ relative to W in dimensions less than 32, here W is the Cayley projective plane canonically embedded in $E_6/D_5 \cdot S^1$. Indeed, the Basis Theorem (cf. Corollary 2) implies already the additive homology of any flag variety G/H .

In [IM, 2005] Iliev and Manivel described the ring $A^*(E_6/D_5 \cdot S^1)$ in terms of three Schubert classes subject to three relations (cf. [IM, Proposition 5.1–5.2]) by using divided difference operators due to Demazure and Bernstein-Gelfand-Gelfand [D; BGG], where the space $E_6/D_5 \cdot S^1$ is called *the complex Cayley plane* and is denoted by $\mathbb{O}\mathbb{P}^2$. Theorem 4 indicates that two Schubert classes and two relations suffice to present the ring.

Historically, the mod p cohomologies $H^*(G; \mathbb{Z}_p)$ for exceptional G were archived by using case by case computations [K]. In 1974, H. Toda [T] initiated a project for computing the integral cohomology of homogeneous spaces G/H with G an exceptional Lie group. This amounts to combine Borel's method [B₁] with the previous results on $H^*(G; \mathbb{Z}_p)$ (as a module over Steenrod algebra) for all prime p . After Toda, the cohomologies of the G/H considered in Theorem 1, 3–6 have been studied by Toda, Watanabe, Ishitoya and Nakagawa (cf. [I, IT, TW, W₁, W₂, N]). In their presentation of the rings the geometry of the generators appears un-transparent. Our method does not resorting to $H^*(G; \mathbb{Z}_p)$, whilst the geometry of the generators are made transparent in the context of Schubert calculus (cf. (2.4)).

Theorem 3 corrects a mistake occurring in earlier computation. Toda and Ishitoya asserted in [IT, 1977] that the ring $H^*(E_6/A_6 \cdot S^1)$ is the quotient of a polynomial ring in eight variables modulo an ideal generated by eight polynomials (with those eight polynomials not being computed explicitly).

Watanabe claimed in [W,1998] that it was generated by three elements in degrees 2, 6 and 8 respectively. However, according to the proof of Theorem 3, *four* is the minimal number of generators of $H^*(E_6/A_6 \cdot S^1)$. This issue witnesses the subtleness in the traditional approach.

Turning to discussion in **2.3**, the classical *Littlewood–Richardson rule* is a combinatorial description of the structure constants for multiplying Schubert classes in the Grassmannian $G_{n,k}(\mathbb{C})$ [M, p.148]. In recent years, a major theme in Schubert calculus is to find an analogue of the rule, for flag varieties of other types, by describing structure constants as the cardinalities of some sets [P; Br; Bu; L].

On the other hand, effective calculation in the cohomology theories of such classical manifolds as the G/H is decidedly required by many problems from geometry and topology [BH]. With our Theorem 1–6 (resp. 7–12) being derived in a unified pattern, we hope very much that we have been able to demonstrate another prospect of the calculus originated in the classical works of H. Schubert [Sch]: the integral cohomology of G/H can be effectively computed without resorting to any information on the topology of Lie groups.

This paper is by no means a final exposition on the topic. Our results and method are ready to extend to flag varieties G/H of more general types. In particular, Theorem 1, 3, 5 will be applied in [DZ₃] to determine the Chow rings of the complete flag manifolds G/T , as well as the integral Chow rings $A(G, \mathbb{Z})$, for all exceptional G [K, Mar].

References

- [B] P. Baum, On the cohomology of homogeneous spaces, *Topology* 7(1968), 15-38.
- [B₁] A. Borel, Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts, *Ann. Math.* 57(1953), 115-207.
- [BH] A. Borel and F. Hirzebruch, Characteristic classes and homogeneous spaces (I), *Amer. J. Math.* 80, 1958, 458–538.
- [BGG] I. N. Bernstein, I. M. Gelfand, S. I. Gelfand, Schubert cells and cohomology of the spaces G/P , *Russian Math. Surveys* 28 (1973), 1-26.
- [Br] M. Brion, Lectures on the Geometry of flag Varieties, *Topics in Cohomological studies of algebraic varieties*, P. Pragacz ed., 33-85. Trends in Math., Birkhauser Verlag Basel, 2005.
- [BS] R. Bott and H. Samelson, Application of the theory of Morse to symmetric spaces, *Amer. J. Math.*, Vol. LXXX, no. 4 (1958), 964-1029.

- [Bu] A.S. Buch, A Littlewood-Richardson rule for the K -theory of Grassmannians, *Acta Math.* 189 (2002), no. 1, 37–78.
- [C] C. Chevalley, Sur les Décompositions Cellulaires des Espaces G/B , in Algebraic groups and their generalizations: Classical methods, W. Haboush ed. Proc. Symp. in Pure Math. 56 (part 1) (1994), 1-26.
- [Co] L. Conlon, On the topology of $EIII$ and EIV , Proc. Amer. Math. Soc., 16(1965), 575-581.
- [D] M. Demazure, Désingularization des variétés de Schubert généralisées, *Ann. Sci. École. Norm. Sup.* (4) 7(1974), 53-88.
- [D₁] H. Duan, Self-maps of the Grassmannian of complex structures, *Compositio Math.*, 132 (2002), 159-175.
- [D₂] H. Duan, Multiplicative rule of Schubert classes, *Invent. Math.* 159 (2005), 407-436.
- [DP] H. Duan and P. Pragacz, Divided differences of Type D and the Grassmannian of complex structures, *Algebraic combinatorics and quantum groups*, 31–60, World Sci. Publishing, River Edge, NJ, 2003.
- [DZ₁] H. Duan and Xuezhi Zhao, Algorithm for multiplying Schubert classes, to appear in *International Journal of Algebra and Computation*; available on arXiv: math.AG/0309158.
- [DZ₂] H. Duan and Xuezhi Zhao, Appendix to “The Chow rings of generalized Grassmannians”, available on arXiv: math.AG/0510085.
- [DZ₃] H. Duan and Xuezhi Zhao, The Chow rings of complete flag varieties, *in preparation*.
- [FP] W. Fulton and P. Pragacz, Schubert Varieties and Degeneracy Loci, *Lecture Notes in Math.*, 1689, Springer, 1998.
- [H] H.C. Hansen, On cycles in flag manifolds, *Math. Scand.* 33 (1973), 269-274.
- [HMS] D. Husemoller, J. Moore, J. Stasheff, Differential homological algebra and homogeneous spaces, *J. Pure Appl. Algebra* 5 (1974), 113–185.
- [Hu] J. E. Humphreys, Introduction to Lie algebras and representation theory, *Graduated Texts in Math.* 9, Springer-Verlag New York, 1972.
- [IM] A. Iliev and L. Manivel, The Chow ring of the Cayley plane, *Compositio Math.*, 141(2005), 146-160.
- [I] K. Ishitoya, Integral cohomology ring of the symmetric space EII . *J. Math. Kyoto Univ.* 17(1977), no.2, 375–397.

- [IT] K. Ishitoya, H. Toda, On the cohomology of irreducible symmetric spaces of exceptional type. *J. Math. Kyoto Univ.* 17(1977), 225–243.
- [K] V.G. Kac, Torsion in cohomology of compact Lie groups and Chow rings of reductive algebraic groups, *Invent. Math.* 80(1985), 69-79.
- [L] C. Lenart, A K -theory version of Monk’s formula and some related multiplication formulas. *J. Pure Appl. Algebra* 179 (2003), no. 1-2, 137–158.
- [M] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford Mathematical Monographs, Oxford University Press, 1995.
- [Mar] R. Marlin, Une conjecture sur les anneaux de Chow $A(G, \mathbb{Z})$ renforcée par un calcul formel, *Effective methods in algebraic geometry, Proc. Symp*, *Prog. Math.* 94(1991) 299-311.
- [MS] J. Milnor and J. Stasheff, *Characteristic classes*, *Ann. of Math. Studies* 76, Princeton Univ. Press, 1975.
- [N] M. Nakagawa, The integral cohomology ring of E_7/T . *J. Math. Kyoto Univ.* 41(2001), 303–321.
- [P] P. Pragacz, Multiplying Schubert classes, *Topics in Cohomological studies of algebraic varieties*, 163–174. *Trends in Math.*, Birkhauser Verlag Basel, 2005.
- [S] J. Scherk, *Algebra. A computational introduction*, *Studies in Advanced Mathematics*, Chapman & Hall/CRC, Boca Raton, FL, 2000.
- [Sch] H. Schubert, *Kalkül de abzählenden Geometrie*, Reprint from the 1879 original, Springer–Verlag, Berlin–New York, 1979.
- [T] H. Toda, On the cohomology ring of some homogeneous spaces. *J. Math. Kyoto Univ.* 15(1975), 185–199.
- [TW] H. Toda, T. Watanabe, The integral cohomology ring of F_4/T and E_6/T , *J. Math. Kyoto Univ.* 14(1974), 257–286.
- [W₁] T. Watanabe, The integral cohomology ring of the symmetric space $EVII$. *J. Math. Kyoto Univ.* 15(1975), 363–385
- [W₂] T. Watanabe, Cohomology of a homogeneous space, *Group representations: cohomology, group actions and topology*, 511–518, *Proc. Sympos. Pure Math.*, 63, Amer. Math. Soc., Providence, RI, 1998.
- [Wo] J. Wolf, The cohomology of homogeneous spaces. *Amer. J. Math.* 99 (1977), no. 2, 312–340.