THE NEIGHBORHOOD COMPLEX OF A RANDOM GRAPH

MATTHEW KAHLE

ABSTRACT. For a graph G, the neighborhood complex $\mathcal{N}[G]$ is the simplicial complex having all subsets of vertices with a common neighbor as its faces. It is a well known result of Lovász that if $\|\mathcal{N}[G]\|$ is k-connected, then the chromatic number of G is at least k + 3.

We prove that the connectivity of the neighborhood complex of a random graph is tightly concentrated, almost always between 1/2 and 2/3 of the expected clique number. We also show that the number of dimensions of nontrivial homology is almost always small, $O(\log d)$, compared to the expected dimension d of the complex itself.

1. INTRODUCTION

In 1978, László Lovász proved Kneser's conjecture [8], that if the *n*-subsets of a (2n+k)set are partitioned into k + 1 families, at least one family contains a disjoint pair. He restated the problem graph theoretically and then proved a more general theorem about graph coloring.

All our graphs will be simple undirected graphs, with no loops or multiple edges. For a graph G, a k-coloring is a function $f: V(G) \to \{1, 2, ..., k\}$ such that $f(x) \neq f(y)$ whenever $\{x, y\} \in E(G)$. The chromatic number $\chi(G)$ is the minimum k such that Gadmits a k-coloring. Define the Kneser graph KG(n, k) to have all n-subsets of a (2n+k)set as its vertices, with edges between disjoint pairs. The Kneser conjecture is equivalent to the claim that $\chi(KG(n, k)) \geq k + 2$.

There are several simplicial complexes naturally associated with a graph G. The clique complex X(G) is the simplicial complex X(G) on vertex set V(G) whose simplices are all complete subgraphs of G. In another article [7], we study the clique complex of a random graph. The *neighborhood complex* $\mathcal{N}[G]$ is the simplicial complex on V(G) which has all subsets of V(G) with a common neighbor for its faces. For example, the neighborhood complex of the complete graph K_n has all proper subsets of the vertices for its faces, so it is the boundary of an (n-1)-dimensional simplex. Its geometric realization $\|\mathcal{N}[K_n]\|$ is homeomorphic to an (n-2)-dimensional sphere \mathbb{S}^{n-2} . (We denote the geometric realization of a simplicial complex Δ by $\|\Delta\|$.)

A topological space X is said to be k-connected if every map from a sphere $\mathbb{S}^n \to X$ extends to a map from the ball $\mathbb{B}^{n+1} \to X$ for $n = 0, 1, \ldots, k$. On the way to proving the Kneser conjecture, Lovász proved the following [8].

Date: January 31,2006.

Supported by NSF grant DMS-9983797.

MATTHEW KAHLE

Theorem 1.1 (Lovász). If $\|\mathcal{N}[G]\|$ is k-connected, then $\chi(G) \ge k+3$.

In the case of the Kneser graphs this lower bound is tight, matching an easy upper bound to give an exact answer. It seems natural to ask how good is the bound for "typical" graphs. For our purposes, a typical graph is the random graph G(n, p) [4].

The random graph G(n, p) is the probability space of all graphs on a vertex set of size n with each edge inserted independently with probability p. Frequently, one considers p to be a function of n and asks whether the graph is likely to have some property as $n \to \infty$. We say that G(n, p) almost always (a.a.) has property \mathcal{P} if $\operatorname{Prob}[G(n, p) \in \mathcal{P}] \to 1$ as $n \to \infty$. The main goal of this article is to understand some of the most basic topological features of $\|\mathcal{N}[G(n, p)]\|$.

2. Statement of results

Let p = p(n) be a monotone function of n, and let i, j, k, and l be integer valued monotone functions of n. In the asymptotic notation that follows, $n \to \infty$ is the free variable. Homology is understood to be reduced with coefficients in \mathbb{Z} throughout.

Theorem 2.1. If $\binom{n}{i}(1-p^i)^{n-i} = o(1)$ then $\mathcal{N}[G(n,p)]$ is a.a. (i-2)-connected.

Theorem 2.2. If $\binom{n}{j}\binom{n}{k}p^{jk} = o(1)$ then $\mathcal{N}[G(n,p)]$ a.a. strong deformation retracts to a simplicial complex of dimension at most j + k - 3.

In proving Theorem 2.2, we make use of the following lemma, which might be of independent interest.

Lemma 2.3. If H is any graph not containing a complete bipartite subgraph $K_{a,b}$ then $||\mathcal{N}[H]||$ strong deformation retracts to a complex of dimension at most a + b - 3.

A *d*-connected complex has trivial homology through dimension *d* by the Hurewicz theorem. Also, $\tilde{H}_k(\Delta) = 0$ whenever *k* is greater than the dimension of Δ , and strong deformation retracts are homotopy equivalences and preserve homology. Hence Theorems 2.1 and 2.2 bound the possible dimensions of nontrivial homology from below and above. We give special cases of the theorems as corollaries. First fix p = 1/2, where G(n, p) is the uniform distribution on all graphs on vertex set [n].

Corollary 2.4. If p = 1/2 and $\epsilon > 0$ then a.a. $\widetilde{H}_l(||\mathcal{N}[G(n,p)]||) = 0$ for $l \leq (1-\epsilon)\log_2 n$ and $l \geq (4+\epsilon)\log_2 n$.

Note for comparison that the dimension of the neighborhood complex is one less than the maximum vertex degree, so when p = 1/2 we expect it to be slightly more than n/2. Next, fix l and check that \tilde{H}_l is trivial outside a certain range of p.

Corollary 2.5. Let $p = n^{\alpha}$ with $\alpha \in [-2, 0]$. If $\alpha > \frac{-1}{l+2}$ then a.a. $\widetilde{H}_l(\|\mathcal{N}[G(n, p)]\|) = 0$. For l even, if $\alpha < \frac{-4}{l+2}$ then a.a. $\widetilde{H}_l(\|\mathcal{N}[G(n, p)]\|) = 0$. For l odd, if $\alpha < \frac{-4(l+2)}{(l+1)(l+3)}$ then a.a. $\widetilde{H}_l(\|\mathcal{N}[G(n, p)]\|) = 0$. For a partial converse to Corollaries 2.4 and 2.5, we exhibit explicit nontrivial homology classes by retracting onto random spheres. Recall that a *clique* of order n is a complete subgraph on n vertices.

Definition 2.6. Let the graph X_n have vertex set $\{u_1, u_2, \ldots, u_n\} \coprod \{v_1, v_2, \ldots, v_n\}$, such that $\{u_1, u_2, \ldots, u_n\}$ spans a clique, and u_i is adjacent to v_j whenever $i \neq j$.

Theorem 2.7. If H is any graph containing a maximal clique of order n that can't be extended to an X_n subgraph, then $\|\mathcal{N}[H]\|$ retracts onto a sphere \mathbb{S}^{n-2} .

Corollary 2.8. If p = 1/2, $\epsilon > 0$, and $(4/3 + \epsilon) \log_2 n < k < (2 - \epsilon) \log_2 n$, then a.a. $\widetilde{H}_k(||\mathcal{N}[G(n,p)]||) \neq 0$.

Corollary 2.9. Let $p = n^{\alpha}$ with $\frac{-2}{k+1} < \alpha < \frac{-4}{3(k+1)}$, then a.a. $\widetilde{H}_k(\|\mathcal{N}[G(n,p)]\|) \neq 0$.

3. Proofs

We first prove that if $\binom{n}{i}(1-p^i)^{n-i} = o(1)$ then $\mathcal{N}[G(n,p)]$ is a.a. (i-2)-connected.

Proof of Theorem 2.1. A simplex complex is *i*-neighborly if every *i* vertices span a face. By simplicial approximation, if a complex is *i*-neighborly then it is (i-2)-connected. The probability that a given set of *i* vertices in G(n,p) has no common neighbor is $(1-p^i)^{n-i}$. Then the total probability that any set of *i* vertices doesn't have a common neighbor is bounded above by $\binom{n}{i}(1-p^i)^{n-i} = o(1)$. So a.a. every such set has some common neighbor, hence spans a face in the neighborhood complex. So $\mathcal{N}[G(n,p)]$ is a.a. *i*-neighborly and (i-2)-connected.

Next we prove that if H is any graph not containing a complete bipartite subgraph $K_{a,b}$ then $||\mathcal{N}[H]||$ strong deformation retracts to a complex of dimension at most a + b - 3.

Proof of Lemma 2.3. For a poset Q, the order complex $\Delta(Q)$ is the simplicial complex of all chains in Q. For a simplicial complex S, let P(S) denote its face poset. To avoid proliferation of notation, we denote the geometric realization of the order complex of a poset Q by ||Q|| rather than $||\Delta(Q)||$.

For a vertex x of H, let $\Gamma(x)$ denote the set of common neighbors of x. Similarly, for any face in the neighborhood complex $X \in \mathcal{N}[H]$, let $\Gamma(X) = \bigcap_{x \in X} \Gamma(x)$. (This map is used in Lovász's paper [8].) Note that Γ is an order reversing self-map of $P(\mathcal{N}[H])$, abbreviated for the rest of this proof by P. So we can define an order preserving poset map $v: P \to P$ by $v(X) = \Gamma(\Gamma(X))$. It's also easy to check that $\Gamma^3 = \Gamma$, so $\Gamma^4 = \Gamma^2$, and $v^2 = v$. Since $v(X) \supseteq X$ for every X, a standard theorem in combinatorial homotopy theory [5] gives that ||v(P)|| is a strong deformation retract of ||P||.

An m+1-dimensional face in v(P) is a chain of faces in $\mathcal{N}[H], X_1 \subsetneq X_2 \subsetneq \cdots \subsetneq X_{m+2}$. Set $Y_i = \Gamma(X_i)$ and we have $Y_1 \supsetneq Y_2 \supsetneq \cdots \supsetneq Y_{m+2}$. (The inclusions $Y_i \subsetneq Y_{i+1}$ are strict since $\Gamma^3 = \Gamma$.)

Suppose $a + b - 3 \leq m$. Since the inclusions $X_i \subsetneq X_{i+1}$ are strict and X_1 is nonempty, X_a contains at least a vertices. Similarly, Y_a contains at least $m + 3 - a \geq b$ vertices. But $Y_p = \Gamma(X_p)$, so X_a and Y_b span a complete bipartite subgraph $K_{a,b}$. Then if the dimension of v(P) is at least m + 1, H contains $K_{a,b}$ subgraphs for every a and b such that $a + b - 3 \leq m$, which is the claim.

Now we apply Lemma 2.3 to check that if $\binom{n}{j}\binom{n}{k}p^{jk} = o(1)$ then $\mathcal{N}[G(n,p)]$ a.a. strong deformation retracts to a simplicial complex of dimension at most j + k - 3.

Proof of Theorem 2.2. Let U and V be vertex subsets of G(n, p) of order j and k respectively. The probability that they span a complete bipartite graph with parts U and V is p^{jk} . So the total probability that there are any $K_{j,k}$ subgraphs is bounded above by $\binom{n}{j}\binom{n}{k}p^{jk} = o(1)$. There are a.a. no such subgraphs, so the claim follows by Lemma 2.3.

Checking Corollaries 2.4 and 2.5 is now a straightforward computation.

Proof of Corollary 2.4. Let p = 1/2 and $\epsilon > 0$. If $l \leq (1-\epsilon) \log_2 n$, then $\binom{n}{l} (1-p^l)^{n-l}$

$$= \binom{n}{l} (1 - (1/2)^l)^{n-l}$$
$$\leq n^l e^{-(1/2)^l (n-l)}$$
$$\leq n^{\log_2 n} e^{-n^{-1+\epsilon} (n-\log_2 n)}$$
$$= \exp\left(\log n \log_2 n - n^{\epsilon} + n^{-1+\epsilon} \log_2 n\right)$$

$$= o(1).$$

Then Theorem 2.1 gives that $\mathcal{N}[G(n,p)]$ is a.a. (l-2)-connected. Since ϵ doesn't appear anywhere in the conclusion of the theorem, we can replace it be a slightly smaller ϵ and for large enough n, $l+2 \leq (1-\epsilon)\log_2 n$ and this gives that $\mathcal{N}[G(n,p)]$ is a.a. *l*-connected.

On the other hand, suppose $l \ge (4 + \epsilon) \log_2 n$ and let $j = \lfloor l/2 \rfloor$.

$$\binom{n}{j}^{2} \left(\frac{1}{2}\right)^{j^{2}} \leq n^{2(2+\epsilon/2)\log_{2}n} \left(\frac{1}{2}\right)^{(2+\epsilon/2)^{2}(\log_{2}n)^{2}} = n^{2(2+\epsilon/2)\log_{2}n-(2+\epsilon/2)^{2}\log_{2}n} = o(n^{-\epsilon\log_{2}n})$$

= o(1).

Then Theorem 2.2 gives that $\mathcal{N}[G(n,p)]$ a.a. strong deformation retracts to a complex of dimension at most $2j - 3 \leq l - 1$.

Proof of Corollary 2.5. Let $p = n^{\alpha}$ and suppose first that $\alpha > \frac{-1}{l+2}$.

$$\binom{n}{l+2} (1-p^{l+2})^{n-l} \le n^{l+2} e^{-n^{\alpha(l+2)}(n-l)}$$

$$\leq \exp\left[(l+2)\log n - n^{1+\alpha(l+2)} + n^{\alpha(l+2)}l\right]$$

= o(1),

since l is constant and $1 + \alpha(l+2) > 0$. Then Theorem 2.1 gives that $\mathcal{N}[G(n,p)]$ is a.a. *l*-connected. So a.a. $\widetilde{H}_l(||\mathcal{N}[G(n,p)]||) = 0.$ Now suppose *l* is even and $\alpha < \frac{-4}{l+2}$. Set $j = \frac{l+2}{2}$.

$$\binom{n}{j}^{2} (n^{\alpha})^{j^{2}}$$

$$\leq n^{l+2} n^{\alpha(l+2)^{2}/4}$$

$$= n^{(l+2)(4+\alpha(l+2))/4}$$

since $4 + \alpha(l+2) < 0$. So Theorem 2.2 gives that $\mathcal{N}[G(n,p)]$ a.a. strong deformation retracts to a complex of dimension at most 2j - 3 = l - 1. Similarly, suppose l is odd and $\alpha < \frac{-4(l+2)}{(l+1)(l+3)}$. Set $j = \frac{l+1}{2}$.

= o(1),

$$\binom{n}{j}\binom{n}{j+1}(n^{\alpha})^{j(j+1)}$$
$$\leq n^{2j+1}n^{\alpha j(j+1)}$$
$$\leq n^{l+2}n^{\alpha \frac{l+1}{2}\frac{l+3}{2}}$$

= o(1).

Then Theorem 2.2 gives that $\mathcal{N}[G(n,p)]$ a.a. strong deformation retracts to a complex of dimension at most $2j-2 \leq l-1$. In both the even and odd cases $\widetilde{H}_l(||\mathcal{N}[G(n,p)]||) = 0$.

MATTHEW KAHLE

Recall that the graph X_n has vertex set $\{u_1, u_2, \ldots, u_n\} \coprod \{v_1, v_2, \ldots, v_n\}$, such that $\{u_1, u_2, \ldots, u_n\}$ spans a clique, $\{v_1, v_2, \ldots, v_n\}$ is an independent set, and u_i is adjacent to v_j whenever $i \neq j$. We show now that if H is any graph containing a maximal clique $\{u_1, u_2, \ldots, u_n\}$ that isn't contained in an X_n subgraph, then $\|\mathcal{N}[H]\|$ retracts onto a sphere \mathbb{S}^{n-2} .

Proof of Theorem 2.7. Suppose H contains a clique $X = \{u_1, u_2, \ldots, u_n\}$ that isn't contained in any larger clique or X_n subgraph. The induced subcomplex of $\mathcal{N}[G(n,p)]$ on X is a topological sphere \mathbb{S}^{n-2} , since X itself is not a face by assumption of maximality of the clique. Define a map on vertices $r_1 : V(H) \to V(H)$ by $r_1(x) = x$ for $x \in X$ and $r_1(x) = u_1$ otherwise.

The only possible obstruction to r_1 extending to a simplicial map $\tilde{r_1} : \mathcal{N}[H] \to \mathcal{N}[H]$, is an (n-1)-dimensional face getting mapped onto X. This happens only if for some vertex $u_1^*, X \cup \{u_1^*\} - u_1$ has a common neighbor v_1 . Note that u_1 isn't adjacent to v_1 since then $X \cup \{v_1\}$ would be an extension of X to a larger clique. Similarly, replacing u_1 with u_i for $i = 2, \ldots, n$. If none of the candidate maps r_1, \ldots, r_n extends to a simplicial map, then the v_i are clearly distinct, since the u_i are distinct and u_i is adjacent to v_j if and only if $i \neq j$. But this yields an X_n subgraph containing X. Otherwise $\|\mathcal{N}[H]\|$ retracts onto $\|X\| = \mathbb{S}^{n-2}$ as claimed, via one of these maps.

Proof of Corollary 2.8. Let p = 1/2 and $\epsilon > 0$. It is well known that G(n, p) a.a. contains maximal cliques of every order k with $(1+\epsilon)\log_2 n < k < (2-\epsilon)\log_2 n$ [4]. We need only check that there are a.a. no X_k subgraphs when $k > (4/3 + \epsilon)\log_2 n$. Note that X_k has 2k vertices and 3k(k-1)/2 edges. Then the probability that G(n, p) contains a copy of X_k is bounded above by

$$(2k)! \binom{n}{2k} \left(\frac{1}{2}\right)^{3k(k-1)/2} \\ \leq n^{2k} \left(\frac{1}{2}\right)^{3k(k-1)/2} \\ \leq n^{(8/3+2\epsilon)\log_2 n} n^{(-3/2)(4/3+\epsilon)((4/3+\epsilon)\log_2 n-1)} \\ = n^{(8/3+2\epsilon)\log_2 n} n^{-(8/3+4\epsilon+3\epsilon^2/2)\log_2 n+2+3\epsilon/2} \\ = n^{-(2+3\epsilon/2)(\epsilon\log_2 n-1)}$$

= o(1).

Proof of Corollary 2.9. Define the density of a graph with v vertices and e edges to be $\lambda = e/v$. We say a graph is strictly balanced if the density of the graph itself is strictly greater than the density of any of its subgraphs.

Let *H* be any strictly balanced graph of density λ . It is classical that $p = n^{-1/\lambda}$ is a sharp threshold for G(n, p) containing *H* as a subgraph [4]. In particular, if $p = n^{\alpha}$ and $\alpha > -1/\lambda$ then G(n, p) a.a. contains *H* as a subgraph, and if $\alpha < -1/\lambda$ then G(n, p) a.a. doesn't contain *H*.

Since K_k and X_k are both strictly balanced we may apply this result twice. The density of K_k is (k-1)/2, and the density of X_k is 3(k-1)/4. So if $p = n^{\alpha}$ with $\frac{-2}{k+1} < \alpha < \frac{-4}{3(k+1)}$, then G(n,p) a.a. contains K_{k+2} but not X_{k+2} subgraphs. This implies that $\widetilde{H}_k(||\mathcal{N}[G(n,p)]||) \neq 0$ by Theorem 2.7 once we check the detail that at least one of these K_{k+2} subgraphs can't be extended to a K_{k+3} . In fact a randomly chosen clique will do the job. The conditional probability that a given K_{k+2} extends to a K_{k+3} is easily seen to be bounded above by np^{k+2} , and $np^{k+2} = o(1)$ since $p = n^{\alpha}$ with $\alpha < \frac{-4}{3(k+1)}$.

4. Connectivity, cliques, and chromatic number

The chromatic number $\chi(G(n, 1/2))$ is tightly concentrated around $n/\log_2 n$. For comparison, the clique number is almost always close to $2\log_2 n$. As a corollary of what we've shown here, the connectivity of the neighborhood complex, somewhere between $\log_2 n$ and $(4/3)\log_2 n$, is almost always less than the clique number.

Similar remarks hold for all monotone functions p = p(n). The asymptotic picture that emerges is the following. The neighborhood complex strong deformation retracts to a complex of dimension d, which is d/4-connected, with nonvanishing homology between dimensions d/3 and d/2, where the clique number is d/2. We see that the connectedness of the neighborhood complex won't do better than the clique number as a lower bound on chromatic number for random graphs; the maximal cliques themselves actually represent nontrivial homology classes.

Recent work of Eric Babson and Dmitry Kozlov [1; 2; 3] provides new examples of topological lower bounds on chromatic number and a more general setting in which to work. However, Carsten Schultz put bounds on the strength of these bounds [10]. In particular, the \mathbb{Z}_2 -index of the neighborhood complex provides the strongest known topological bound on chromatic number. This may in general be higher than the connectivity of the neighborhood complex. But by what we've shown here, even the \mathbb{Z}_2 -index won't do much better for random graphs as a lower bound on chromatic number than connectivity, since the dimension of the retract is an upper bound on the index.

5. RANDOM SIMPLICIAL COMPLEXES AND UNIMODALITY

One justification for random graph theory is that it provides models for "typical" graphs. This can be made precise in a few ways. For example, G(n, 1/2) is the uniform distribution on all graphs on vertex set [n]. Any property that G(n, 1/2) a.a. has is a

MATTHEW KAHLE

property of almost all graphs. Or for another example, the Szemerédi Regularity Lemma states that every graph is well approximated by random graphs.

Every neighborhood complex is homotopy equivalent to a free \mathbb{Z}_2 -complex, via Lovasz's retract. Up to homotopy, the converse also holds [6].

Theorem 5.1 (Csorba). Given a finite simplicial complex Δ with a free \mathbb{Z}_2 -action, there exists a graph G such that $\|\mathcal{N}[G]\|$ is homotopy equivalent to $\|\Delta\|$.

So neighborhood complexes of random graphs asymptotically give a probability distribution on all finite triangulable \mathbb{Z}_2 -spaces as $n \to \infty$, at least up to homotopy type.

Little seems to be written so far about random simplicial complexes. However, Nathan Linial and Roy Meshulam recently studied $H_1(||Y(n,p)||,\mathbb{Z}_2)$ for random 2-dimensional simplicial complexes Y(n,p) [9]. Their definition of Y(n,p) is a natural extension of the Erdős-Rényi random graph G(n,p); Y(n,p) has vertex set [n] and edge set $\binom{[n]}{2}$, with each 2-face appearing independently with probability p. One advantage of the Linial-Meshulam model is that vanishing of homology is a monotone property. That is, once enough 2-faces have been added that $H_1(||Y(n,p)||,\mathbb{Z}_2)$ vanishes, adding more 2-faces can't ever make it nonvanishing. (This particular fact doesn't depend on the coefficients of homology, but only on the definition of random 2-complex. Simple connectivity is also a monotone property but it's still not known where the threshold function lies.)

Most properties of G(n, p) that have been studied to date are monotone graph properties, in contrast to what we've studied in this article, where vanishing of homology is clearly not monotone. But in this setting unimodality seems like a natural substitute for montonicity.

In another article [7], we study the *clique complex* of a random graph, which is the simplicial complex with all complete subgraphs for its faces. The results are analogous to what we find here, although we also study the expectation of the Betti numbers. Denote the kth Betti number by β_k . We conjecture that for both random neighborhood and clique complexes, for any fixed k and large enough n depending on k, the expectation $E[\beta_k]$ is a unimodal function of p.

6. Acknowledgements

The author wishes to thank his advisor Eric Babson for inspiration, guidance, and patience; fellow graduate students, particularly Anton Dochtermann, Alex Papazoglou, and David Rosoff, for many helpful conversations; and Sara Billey for generous support. Any mistakes are his own.

References

 E. Babson and D. Kozlov, Topological obstructions to graph coloring, Electron. Res. Announc. Amer. Math. Soc. 9, 61-68, 2003.

[2] E. Babson and D. Kozlov, Complexes of graph homomorphisms, arXiv:math.CO/0310056, 2003.

- [3] E. Babson and D. Kozlov, Proof of the Lovász conjecture, arXiv:math.CO/0402395, 2003.
- [4] B. Bollobás, Random Graphs, Second Edition, Cambridge University Press, 2001.
- [5] A. Björner, Topological methods, Handbook of combinatorics (vol. 2), 1819-1872, 1996.
- [6] P. Csorba, Homotopy types of box complexes, arXiv:math.CO/0406118, to appear in Combinatorica.
- [7] M. Kahle, Topology of random clique complexes, in preparation.
- [8] L. Lovász, Kneser's conjecture, chromatic number, and homotopy, J. Comb. Th. A, **25**, 319-324, 1978.
- [9] N. Linial and R. Meshulam, Homological connectivity of random 2-complexes, to appear in Combinatorica.
- [10] C. Schultz, The relative strength of graph coloring obstructions, preprint, 2006.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON *E-mail address*: kahle@math.washington.edu