FILTERED ENDS OF INFINITE COVERS AND GROUPS

TOM KLEIN

ABSTRACT. Let $f:A\to B$ be a covering map. We say A has e filtered ends with respect to f (or B) if for some filtration $\{K_n\}$ of B by compact subsets, $A-f^{-1}(K_n)$ "eventually" has e components. The main theorem states that if Y is a (suitable) free H-space, if K < H has infinite index, and if Y has a positive finite number of filtered ends with respect to $H \setminus Y$, then Y has one filtered end with respect to $K \setminus Y$. This implies that if G is a finitely generated group and K < H < G are subgroups each having infinite index in the next, then $0 < \tilde{e}(G, H) < \infty$ implies $\tilde{e}(G, K) = 1$, where $\tilde{e}(\cdot, \cdot)$ is the number of filtered ends of a pair of groups in the sense of Kropholler and Roller.

1. Introduction

The number of relative ends of a group G and subgroup H, denoted $\tilde{e}(G,H)$, was originally introduced algebraically by Kropholler and Roller [2]. Geoghegan [1] gives a topological description of $\tilde{e}(G,H)$ (when G is finitely generated) as the number of filtered ends of a certain filtration (derived from H) of the universal cover of a presentation complex for G. To be more precise (see also section 2), let $f:A\to B$ be a covering map of CW complexes of locally finite type. We say A has $e=\tilde{e}(A,f)$ filtered ends with respect to f (or B) if for some filtration $\{C_n\}$ of B by finite subcomplexes, the cardinality of $\varprojlim_n \left\{\pi_0\left(A-f^{-1}(C_n)\right)\right\}$ is e (this is independent of the filtration). $\tilde{e}(G,H)$, also called

the number of filtered ends of the pair, is then $\tilde{e}(\widetilde{X}^1,q)$, where X is a path connected CW complex with $\pi_1(X) = G$, \widetilde{X} is the universal cover of X, and $q: \widetilde{X}^1 \to \bar{X}(H)^1$ is the restriction to 1-skeleta of the natural map from \widetilde{X} to the covering space of X corresponding to H. The following theorem on filtered ends of covering spaces (proved in section 3) and its corollary on filtered ends of pairs of groups (proved in section 5) are the main results of the article.

Theorem 1. Let H be a group and let Y be a path connected free H-CW complex of locally finite type. Let K < H with $[H : K] = \infty$, and let $p : Y \to H \backslash Y$, $q : Y \to K \backslash Y$ be the quotient covering maps. Then $0 < \tilde{e}(Y, p) < \infty$ implies $\tilde{e}(Y, q) = 1$.

Applied to 1-skeleta with Y equal to the universal cover of a presentation complex (with finite 1-skeleton) of a finitely generated group G, we get the following corollary for filtered ends of pairs of groups:

Corollary 2. Let G be a finitely generated group with $K \leq H < G$ and assume $\tilde{e}(G, H) = n$ is finite and non-zero. Then

$$\tilde{e}(G,K) = \left\{ \begin{array}{ll} n & \textit{if} \ [H:K] < \infty \\ 1 & \textit{if} \ [H:K] = \infty. \end{array} \right.$$

Note also the contrapositive of Corollary 2: if $\tilde{e}(G,K)>1$ and $[G:H]=[H:K]=\infty$, then $\tilde{e}(G,H)=\infty$. This contrapositive combined with other results can be used to show that Thompson's groups T and V are not Kähler (cf. [3], Lemma 0.8 and Theorem 0.3): T and V are known to have subgroups with at least two filtered ends and then the contrapositive provides subgroups with at least three filtered ends so that the theorems of [3] can be applied. It seems likely that similar arguments could be applied to other groups with subgroups having two filtered ends.

ACKNOWLEDGEMENTS

Theorem 1 is in answer to a question of Mohan Ramachandran. I would like to thank Ross Geoghegan for suggestions which improved the exposition.

To appear in the Journal of Pure and Applied Algebra.

2 TOM KLEIN

2. Definitions

We now briefly review the necessary definitions (sections 4.7 and 4.9 of [1]). We are heading for "the number of ends of a CW complex with respect to a filtration" (Definition 3). We work with closed cells and use the term graph for a 1-dimensional CW complex. Let X be an arbitrary CW complex.

X is locally finite if a given cell intersects non-trivially with only finitely many other cells, and X has locally finite type if the n-skeleton X^n of X is locally finite for all n.

The carrier of a cell e of X, denoted C(e), is the intersection of all subcomplexes of X which contain e.

The CW neighborhood of a subcomplex A of X, denoted N(A), is the union of all cell carriers that meet A non-trivially.

The CW complement of a subcomplex A of X, denoted X - A, is the largest subcomplex of X with 0-skeleton $X^0 - A^0$.

All filtrations of a CW complex X will be by subcomplexes of X indexed by the natural numbers. A finite filtration is a filtration by finite subcomplexes¹. If X has locally finite type and $\mathcal{K} = \{K_n\}$ is a filtration of X, we say (X, \mathcal{K}) is well filtered if for each n and i there exists j such that $N_{X^n}(K_i^n) \subseteq K_j$.

Now let $(Y, \{L_i\})$ be a well filtered path connected CW complex of locally finite type. The number

of filtered ends of $(Y, \{L_i\})$ is the cardinality of the set $\varprojlim_i \left\{ \pi_0 \left(Y - L_i \right) \right\}$. (We are skipping the

intermediate notion of filtered end, which is what the number of filtered ends counts – briefly, a filtered end of Y is an equivalence class of filtered rays $\omega : [0, \infty) \to Y$, where two filtered rays are equivalent if their restrictions to \mathbb{N} are filtered homotopic. Then each $[\omega]$ picks out an element of the inverse limit, and this correspondence is a bijection. See [1] section 4.7 for more details.)

Definition 3. Let Y and X be path connected CW complexes of locally finite type and let $p: Y \to X$ be a covering map. We say Y has e filtered ends over p $(0 \le e \le \infty)$, and write $\tilde{e}(Y,p) = e$, if the pair $(Y, \{p^{-1}(C_n)\})$ has e filtered ends for some finite filtration $\{C_n\}$ of X. [By Lemma 5.i) below there exists a finite filtration and $(Y, \{p^{-1}(C_n)\})$ is well filtered. Also it is known that in this case Y has e filtered ends with respect to every finite filtration of X.]

Remark 4. As noted in section 4.7 of [1], if (Y, L) is a well filtered path connected CW complex of locally finite type then the number of filtered ends of $(Y^1, \{L_i \cap Y^1\})$ equals the number of filtered ends of $(Y, \{L_i\})$ (by the "CW-filtered cellular approximation theorem", [1] 3.12.3).

We use this in the proof of Theorem 1 to reduce to the case where Y is a graph.

Let A be a subgraph of a path connected locally finite graph X. In this case N(A) is simply $A \cup \bigcup \{ \text{edges of } X \text{ with at least one endpoint in } A \}$. Then $X = N(A) \cup (X - A) = A \cup B \cup (X - A)$, where $B = \bigcup \{ \text{edges not contained in } A \text{ or } X - A \}$. We refer to the edges of B as $bridge \ edges$ (of A).

If (Y, \mathbb{L}) is a filtered CW complex we say $S \subseteq Y$ is L-bounded if $S \subseteq L_j$ for some j, and (Y, \mathbb{L}) is regular if for each i the union of all L-bounded path components of $Y - L_i$ is L-bounded ([1], section 4.7).

The next proposition shows that these notions all work well with respect to covering spaces of CW complexes of locally finite type.

Lemma 5. i) If X is a path connected CW complex of locally finite type then there exists a finite filtration $\{K_n\}$, and $(X, \{K_n\})$ is well filtered (for any finite filtration).

ii) If $p: Y \to X$ is a covering map of CW complexes then Y has locally finite type if and only if X does.

iii) Let $p: Y \to X$ be a covering map of CW complexes of locally finite type. If $(X, \{K_n\})$ is well filtered then so is $(Y, \{p^{-1}(K_n)\})$.

iv) Let X be a locally finite graph with $p: Y \to X$ a covering map, and let $\{C_n\}$ be a finite filtration of X. Then $(Y, \{p^{-1}(C_n)\})$ is well filtered and regular.

¹This differs from [1], section 3.6, which requires a finite filtration to be by full finite subcomplexes.

Proof. i) By [1] 3.6.3 X is countable. Well order the cells as e_1, e_2, \ldots and let $E_n = e_1 \cup \cdots \cup e_n$. Since E_n is compact it is contained in a (minimal) finite subcomplex, say K_n . Then $\{K_n\}$ is a finite filtration of X. By [1], 3.1.12 and 3.6.9, $N_{X^n}(K_i^n)$ is finite, so $(X, \{K_n\})$ is well filtered.

- ii) $p|_{Y^n}:Y^n\to X^n$ is a covering map, so Y^n is locally finite if and only if X^n is locally finite.
- iii) Let $q = p|_{Y^n}: Y^n \to X^n$ and let $L_i = p^{-1}(K_i)$. $q(N_{Y^n}(L_i^n)) = N_{X^n}(K_i^n)$ is contained in some K_i (j > i).
- iv) Well filtered follows from i) and iii). For regular, note that $p(Y-p^{-1}(C_n))=X-C_n$ and that if A is an L-bounded path component of $Y-p^{-1}(C_n)$ ($\mathbbm{L}=\{p^{-1}(C_n)\}$) then p(A) is contained in a finite component of $X-C_n$ since otherwise we could construct an unbounded ray in $X-C_n$ starting in p(A) which would lift to an L-unbounded ray in $Y-p^{-1}(C_n)$ starting in A, a contradiction. But since $N(C_n)$ is finite and X is locally finite, there are only finitely many finite components of $X-C_n$. \square

We now turn to filtered ends of pairs of groups ([1], section 4.9). Let G be a finitely generated group and $H \leq G$. Let X be a path connected CW complex with fundamental group isomorphic to G and having finite 1-skeleton. Let $\bar{X}(H)$ be the covering space of X corresponding to H and let \tilde{X} be the universal cover of X. We work on the 1-skeleta of \tilde{X} and $\bar{X}(H)$, both locally finite graphs. Let $p:\tilde{X}^1\to \bar{X}(H)^1$ be the covering projection. The number of filtered ends of (G,H) is $\tilde{e}(G,H):=\tilde{e}(\tilde{X}^1,p)$ (see Definition 3). In other words, start with any finite filtration $\{C_n\}$ of $\bar{X}(H)^1$, then count the number of $\{p^{-1}(C_n)\}$ -unbounded components of $\tilde{X}^1-p^{-1}(C_k)$ and take the sup of that count as k goes to infinity.

3. Proof of Theorem 1

Remark 6. The case $\tilde{e}(Y,p)=1$ implies $\tilde{e}(Y,q)=1$ of the theorem follows from the following "monotonicity" property of $\tilde{e}(Y,\cdot)$: if the covering map p (of CW complexes of locally finite type) factors through the covering map q as $p=r\circ q$ then $\tilde{e}(Y,q)\leq \tilde{e}(Y,p)$. However, we will not need this fact.

PROOF of Theorem 1. By Remark 4 we may assume Y is a graph, which we do. Define $X := H \setminus Y$ and $\bar{X} := K \setminus Y$, and let $r : \bar{X} \to X$ be the induced covering map. The first step of the proof is to produce compatible filtrations of \bar{X} and X along with a rough "fundamental domain" in Y (called \tilde{C}_n below) for the H and K actions. The specifics of the filtrations are stated in the following lemma, whose proof will be postponed to section 4.

Lemma 7. There exists a finite filtration $\{\bar{C}_n\}$ of \bar{X} and a finite filtration $\{C_n\}$ of X with $r(\bar{C}_n) = C_n$ and $U_n := Y - p^{-1}(C_n)$ the disjoint union of $e\{p^{-1}(C_n)\}$ -unbounded path components. There are also finite subgraphs $\tilde{C}_n \subseteq p^{-1}(C_n)$ and $\hat{C}_n \subseteq N(p^{-1}(C_n))$ in Y with $\hat{C}_n \cap p^{-1}(C_n) = \tilde{C}_n$, $q(\tilde{C}_n) = \bar{C}_n$, $p(\tilde{C}_n) = C_n$, and $\hat{C}_n \cup U_n$ path connected.

Let \mathcal{K} be the filtration $\{p^{-1}(C_n)\}$ of Y and let \mathcal{L} be the filtration $\{q^{-1}(\bar{C}_n)\}$. Define U^1, U^2, \ldots, U^e to be the e \mathcal{K} -unbounded path components of $U_n = Y - p^{-1}(C_n)$, so that $U_n = U^1 \sqcup \cdots \sqcup U^e$.

H preserves $p^{-1}(C_n)$, so also bridge edges of $p^{-1}(C_n)$ and $U_n = Y - p^{-1}(C_n)$. Then $\hat{C}_n \cup U_n$ path connected implies

- (†) $h(\hat{C}_n \cup U_n) = h\hat{C}_n \cup U_n$ is path connected for all $h \in H$ and all n.
- $p^{-1}(C_n) = H\widetilde{C}_n$ and $q^{-1}(\overline{C}_n) = K\widetilde{C}_n$ (since $p(\widetilde{C}_n) = C_n$ and $q(\widetilde{C}_n) = \overline{C}_n$). Let $A = \bigcup \left\{ h\widetilde{C}_n : h \in H \text{ and } h\widetilde{C}_n \cap K\widetilde{C}_n = \emptyset \right\}$, $B = \bigcup \left\{ h\widetilde{C}_n : h \in H \text{ and } h\widetilde{C}_n \cap K\widetilde{C}_n \neq \emptyset \right\}$. The proof of the next claim contains the main argument of the proof of the theorem (compare with Figure 1).
- Claim 8. i) $A \neq \emptyset$. $\hat{U}_n := A \cup U_n \cup \{\text{all bridge edges of } p^{-1}(C_n) \text{ that intersect } A \text{ but not } q^{-1}(\bar{C}_n) \}$ is path connected in $Y = q^{-1}(\bar{C}_n)$ and L-unbounded.
- ii) $B \cup \{all \ bridge \ edges \ of \ p^{-1}(C_n) \ that \ intersect \ B\}$ is L-bounded.

4 TOM KLEIN




FIGURE 1. The decomposition of $N(p^{-1}(C_n))$ from Claim 8.

Proof. If $h\widetilde{C}_n \cap k\widetilde{C}_n \neq \emptyset$ for some $h \in H, k \in K$, then $k^{-1}h\widetilde{C}_n \cap \widetilde{C}_n \neq \emptyset$, so letting $S = \{g \in H : g\widetilde{C}_n \cap \widetilde{C}_n \neq \emptyset\}$ we get $k^{-1}h =: s_0 \in S$. Now let T be a set of right coset representatives for K in H. For each $s \in S$ write $s = k_s t_s$, where $t_s \in T$ and $k_s \in K$. Then $k^{-1}h = k_{s_0}t_{s_0}$ and $h = (kk_{s_0})t_{s_0}$, so if $h\widetilde{C}_n \cap K\widetilde{C}_n \neq \emptyset$ then h lies in one of finitely many right cosets of K in H.

i) By assumption $[H:K]=\infty$, so there are in fact infinitely many $h\in H$ with $h\widetilde{C}_n\cap K\widetilde{C}_n=\emptyset$. So A is non-empty. Recall that $\widetilde{C}_n=\hat{C}_n\cap p^{-1}(C_n)$, so bridge edges of $p^{-1}(C_n)$ contained in \hat{C}_n intersect $p^{-1}(C_n)$ in \widetilde{C}_n . So if $h\widetilde{C}_n\subseteq A$ then the bridge edges of $p^{-1}(C_n)$ contained in $h\widehat{C}_n$ only intersect $p^{-1}(C_n)$ in $h\widetilde{C}_n\subseteq A$ and not in $q^{-1}(\bar{C}_n)$. Then for $h\widetilde{C}_n\subseteq A$ we get $h\widehat{C}_n\cup U_n=h\widetilde{C}_n\cup \{\text{bridge edges of }p^{-1}(C_n)\text{ in }h\widehat{C}_n\}\cup U_n\subseteq \hat{U}_n$, and $h\widehat{C}_n\cup U_n$ is path connected by (\dagger) , so \widehat{U}_n is path connected. $(A\cup U_n)\cap q^{-1}(\bar{C}_n)=\emptyset$ by definition of A and U_n , so \widehat{U}_n is path connected in $Y-q^{-1}(\bar{C}_n)$. \widehat{U}_n is L-unbounded since it contains U_n which is \mathcal{K} -unbounded (\mathcal{K} -unbounded implies L-unbounded).

ii) First note that q(B) is finite since $q(K\widetilde{C}_n) = \overline{C}_n$ is finite and the other edges of B are within a bounded edge distance of $K\widetilde{C}_n$ (and \overline{X} is locally finite). But then the union of all edges intersecting B has finite image.

Now write $Y = q^{-1}(\bar{C}_n) = \bar{U}_n \sqcup \bar{D}_n$, where \bar{U}_n is the path component containing \hat{U}_n and \bar{D}_n is the union of the other components (possibly empty). \bar{U}_n is L-unbounded since \hat{U}_n is. We can write Y as $Y = p^{-1}(C_n) \cup \{\text{bridge edges of } p^{-1}(C_n)\} \cup U_n$, and $A \cup B = p^{-1}(C_n)$, so $Y = (A \cup U_n \cup \{\text{bridge edges intersecting } A \text{ but not } q^{-1}(\bar{C}_n)\}) \cup (B \cup \{\text{bridge edges intersecting } B\})$, and \bar{U}_n contains the first parenthetical term, so \bar{D}_n must be contained in the second parenthetical term, which by Claim 8.ii) is L-bounded. So for each n, $Y = q^{-1}(\bar{C}_n) = \bar{U}_n \sqcup \bar{D}_n$ where \bar{U}_n is L-unbounded and \bar{D}_n is L-bounded.

Finally as in the proof of Lemma 9 we can adjust the $\{\bar{C}_n\}$ filtration so that $q^{-1}(\bar{C}_n)$ is a single L-unbounded component for each n, from which it follows that $\varprojlim_n \left\{ \pi_0 \left(Y - q^{-1}(\bar{C}_n) \right) \right\}$ has cardinality 1. i.e. $\tilde{e}(Y,q) = 1$.

4. Construction of compatible filtrations

We now undertake the proof of Lemma 7. Recall that we are working under the hypotheses of Theorem 1 with Y a locally finite graph, $p:Y\to X,\ q:Y\to \bar X,\ r:\bar X\to X,\ {\rm and}\ 0<\tilde e(Y,p)=e<\infty.$ We restate Lemma 7:

Lemma 7. There exists a finite filtration $\{\bar{C}_n\}$ of \bar{X} and a finite filtration $\{C_n\}$ of X with $r(\bar{C}_n) = C_n$ and $Y - p^{-1}(C_n)$ the disjoint union of $e\{p^{-1}(C_n)\}$ -unbounded path components. There are also finite subgraphs $\tilde{C}_n \subseteq p^{-1}(C_n)$ and $\hat{C}_n \subseteq N(p^{-1}(C_n))$ in Y with $\hat{C}_n \cap p^{-1}(C_n) = \tilde{C}_n$, $q(\tilde{C}_n) = \bar{C}_n$, $p(\tilde{C}_n) = C_n$, and $\hat{C}_n \cup U_n$ path connected.

The proof will start from the following general lemma.

Lemma 9. Let A be a locally finite graph and $f: A \to B$ a covering map. If $\tilde{e}(A, f) = e$, $0 < e < \infty$, then there exists a finite filtration $\{D_n\}$ of B such that $A = p^{-1}(D_n)$ is the union of $e \{p^{-1}(D_n)\}$ -unbounded path components for all n.

Proof. Let $\{K_n\}$ be any finite filtration of B and let $\mathcal{L} = \{p^{-1}(K_n)\}$. By Corollary 4.7.5 of [1] we may assume $A - p^{-1}(K_n)$ has exactly e \mathcal{L} -unbounded components for all n. Now for each n let $D_n = K_n \bigcup \bigcup \{\text{finite components of } B - K_n\}$. The union is finite (since B is locally finite), so D_n is finite. The image of a $\{p^{-1}(D_n)\}$ -bounded component of $A - p^{-1}(D_n)$ would have to be contained in a finite component of $B - D_n$ (see the proof of Lemma 5.iv)), so $A - p^{-1}(D_n)$ consists exactly of e $\{p^{-1}(D_n)\}$ -unbounded components.

Remark 10. If $e = \infty$ in the hypotheses of Lemma 9, then there exists a finite filtration $\{C_n\}$ of X such that $Y - p^{-1}(C_n)$ equals the union of $\{p^{-1}(C_n)\}$ -unbounded path components, with the number of path components an increasing function of n.

PROOF of Lemma 7. Let $\{\bar{C}_n\}$ be a finite filtration of \bar{X} . Define $C_n = r(\bar{C}_n)$. $\{C_n\}$ is a finite filtration of X. By the (proof of) Lemma 9 (after renumbering) for each n there exists a finite subgraph F_n of X with the property that $\{C_n \cup F_n\}$ is a finite filtration of X and $Y - p^{-1}(C_n \cup F_n)$ is the union of e $\{p^{-1}(C_n \cup F_n)\}$ -unbounded path components. Now for each n choose a finite subgraph \bar{F}_n in \bar{X} with $r(\bar{F}_n) = F_n$, choose a subsequence filtration of $\{\bar{C}_n \cup \bar{F}_n\}$, and then rename the \bar{X} subsequence filtration to be $\{\bar{C}_n\}$ and the X filtration to be $\{C_n = r(\bar{C}_n)\}$, a subsequence of the old $\{C_n \cup F_n\}$ filtration. To summarize: we have a finite filtration $\{\bar{C}_n\}$ of \bar{X} and a finite filtration $\{C_n\}$ of X with the property that $x(\bar{C}_n) = C_n$ and $x - p^{-1}(C_n)$ is the union of $x \in \{p^{-1}(C_n)\}$ -unbounded path components for all $x \in \mathbb{R}$.

Let
$$K = \{p^{-1}(C_n)\}.$$

Choose a finite subgraph \widetilde{C}_n in Y with $q(\widetilde{C}_n) = \overline{C}_n$. The given lifts \widetilde{C}_n will not in general satisfy the properties we will require, but we can use them to construct all new filtrations that will, which we do now. Fix n. Let U^1, \ldots, U^e be the \mathcal{K} -unbounded components of $Y - p^{-1}(C_n)$ and define $U_n = U^1 \cup \cdots \cup U^e$. Our next aim is to define new \widetilde{C}_n so that $U_n \cup \widetilde{C}_n$ is path connected (up to bridge edges of $p^{-1}(C_n)$). Claim 11 tells us how to adjust our original choices (compare with Figure 2).

Claim 11. There exists a finite subgraph \hat{C}_n of $N(p^{-1}(C_n))$ containing \widetilde{C}_n such that $\hat{C}_n \cup U_n$ is path connected.




FIGURE 2. The construction of \hat{C}_n in Claim 11.

Proof. Let α be a finite edge path (i.e. union of edges) in Y starting in U_n with the property that α intersects each U^i non-trivially. Let $\tilde{\alpha}$ be the edges of α contained in $N(p^{-1}(C_n))$. Then $\tilde{\alpha} \cup U_n$ is path connected. Now for each cell e of \tilde{C}_n choose a finite edge path β_e in $N(p^{-1}(C_n))$ to U_n . Take


$$\hat{C}_n = \widetilde{C}_n \cup \widetilde{\alpha} \cup \left(\bigcup_{e \subset \widetilde{C}_n} \beta_e\right).$$

6 TOM KLEIN

Let $\widetilde{C}'_n = \widehat{C}_n \cap p^{-1}(C_n) \supseteq \widetilde{C}_n$, and let $\overline{C}'_n = q(\widetilde{C}'_n)$ $(p(\widetilde{C}'_n) = C_n)$, so we do not rename for X). $\{\overline{C}'_n\}$ may no longer be a chain of subsets, but since we only added finitely many cells to each term of the original filtration we can find a subsequence which is a finite filtration. Now renumber and then rename all filtrations to get finite filtrations $\{\overline{C}_n\}$ for X, $\{C_n\}$ for X, and a finite lift \widetilde{C}_n for each \overline{C}_n with the property that there is a finite $\widehat{C}_n \subseteq N(p^{-1}(C_n))$ containing \widetilde{C}_n with $U_n \cup \widehat{C}_n$ path connected – note that we only took a subsequence of our previous X filtration $\{C_n\}$, so we still have $Y - p^{-1}(C_n) = U_n$, where $U_n = U^1 \cup \cdots \cup U^e$ is the same union of K-unbounded path components as before (of course numbering and the meaning of K have changed, but the components and their K-unboundedness have not).

5. FILTERED ENDS OF PAIRS OF GROUPS


We now apply this to groups. The general situation will be as follows:



G is finitely generated infinite, $K \leq H \leq G$, and X(G) has finite 1-skeleton and fundamental group G (we write " $\bar{X}(L)$ " for the covering space of X corresponding to $L \leq G$). We apply Theorem 1 to the dotted triple of covering spaces to obtain Corollary 2 on filtered ends of pairs of groups.

PROOF of Corollary 2. The case $[H:K] < \infty$ is standard ([2], 2.4.v)); the case $[H:K] = \infty$ is the statement of Theorem 1.

Corollary 2 and its contrapositive give the following picture of filtered end behavior above and below a subgroup H with $\tilde{e}(G, H) = n$:



References

- [1] R. Geoghegan, Topological Methods in Group Theory, forthcoming book.
- [2] P. Kropholler, M. Roller, Relative ends and duality groups, J. Pure Appl. Algebra 61 (1989), no. 2, 197-210.
- [3] T. Napier, M. Ramachandran, Filtered ends, proper holomorphic mappings of Kähler manifolds to Riemann surfaces, and Kähler groups, arXiv:math.DG/0506254.