

L_∞ BIALGEBRAS AND L_∞ QUASIBIALGEBRAS

OLGA KRAVCHENKO

This paper is dedicated to Jean-Louis Loday on the occasion of his 60th birthday with admiration and gratitude

Abstract

Structures of Lie algebras, Lie coalgebras, Lie bialgebras and Lie quasibialgebras are presented as solutions of Maurer-Cartan equations on corresponding governing differential graded Lie algebras. There are several directions in this paper. Firstly, cohomology theories of all these structures are described in a concise way using the big bracket construction of Kosmann-Schwarzbach. Secondly, this approach provides a definition of an L_∞ -(quasi)bialgebra (also known as strong homotopy Lie (quasi)bialgebra). We recover an L_∞ -algebra structure as a particular case of our construction. Thirdly, formal geometry interpretation leads us to a definition of an L_∞ (quasi)bialgebra structure on V as a differential operator Q on V , autocommuting with respect to the Poisson bracket. Finally, we establish an L_∞ -version of Manin triple (and Manin pair) and get a correspondence theorem with L_∞ -bialgebras (L_∞ -(quasi)bialgebra).

1. INTRODUCTION.

Algebraic structures are often defined as certain maps which must satisfy quadratic relations. One of the examples is a Lie algebra structure: a Lie bracket satisfies the Jacobi identity (indeed the Jacobi identity is a quadratic relation since the bracket appears twice in each summand). Other examples include an associative multiplication (associativity condition is quadratic), L_∞ and A_∞ algebras (also called strong homotopy Lie and strong homotopy associative algebras) and many others.

The subject of this article is a description of Lie (quasi) bialgebras and their L_∞ -versions. The main philosophy is that the axioms of Lie (quasi)bialgebras (and their L_∞ -versions) could be written in the form of a quadratic relation on a certain governing differential graded Lie algebra. We find the governing dgLie algebras for Lie bialgebra and Lie quasi-bialgebra structures using Kosmann-Schwarzbach's big-bracket construction [6].

In general, the governing Lie algebra of a structure could be defined as follows.

Definition 1.1. *Consider a graded Lie algebra*

$$\mathfrak{g} = \bigoplus_k \mathfrak{g}^k, \quad [\cdot, \cdot] : \mathfrak{g}^k \wedge \mathfrak{g}^l \rightarrow \mathfrak{g}^{k+l}.$$

a: *Let there exist an element $\theta \in \mathfrak{g}$, satisfying $[\theta, \theta] = 0$. Call $Q = \text{ad}_\theta$ an algebraic structure.*

Then $\mathfrak{g}$ is called the governing graded Lie algebra of the structure Q .

b: *Let, moreover, $\mathfrak{g}$ be a differential graded Lie (dgLie) algebra, that is there exist an operator*

$$d : \mathfrak{g}^k \rightarrow \mathfrak{g}^{k+1}, \quad d^2 = 0, \quad \forall k,$$

which is a derivation for the Lie bracket. Let $\theta \in \mathfrak{g}$ satisfy $[\theta, \theta] = 0$, and $d\theta = 0$. Then $Q = d + \text{ad}_\theta$ is also called an algebraic structure and $\mathfrak{g}$ is its governing dgLie algebra.

In fact, one could get from [a] to [b] by setting $d = \text{ad}_\theta$ and passing from a graded Lie algebra to a differential graded Lie algebra. The quadratic relation on the structure could be expressed in the form of a Maurer-Cartan equation (MC equation): $d\gamma + \frac{[\gamma, \gamma]}{2} = 0$, in other words, $ad_{m+\gamma}$ must be a differential on $\tilde{\mathfrak{g}}$.

Well studied examples of such a setup include an associative multiplication on the space A . The governing graded Lie algebra of an associative structure $\mathfrak{g} = \sum_{k \geq 0} \mathfrak{g}^k$, $\mathfrak{g}^k = \text{Hom}(A^{\otimes k+1}, A)$, so that a multiplication $m \in \text{Hom}(A^{\otimes 2}, A) = \mathfrak{g}^1$. The Lie structure on $\mathfrak{g}$ is given by the Gerstenhaber bracket $[\cdot, \cdot]_G$. The role of θ takes an element m from $\text{Hom}(A^{\otimes 2}, A)$, such that the associativity condition reads as $(m \otimes \text{id} + \text{id} \otimes m)m = 0$, which in terms of the Gerstenhaber bracket is written as an equation of second order: $[m, m]_G = 0$. In other words, $d = \text{ad}_m$ is a differential. In fact, any other associative structure on A is described by an auto-commuting element of degree 1 in the governing differential graded Lie algebra (in this case this dgLie algebra is a Lie algebra of coderivations of the tensor algebra of A , see for example [12]).

Usually, solutions of the Maurer-Cartan equation are considered only from $\mathfrak{g}_1$, however if we allow terms in the whole $\mathfrak{g}$ the corresponding structure is a strong homotpy version of the one obtained from $\mathfrak{g}_1$.

Let us give here the definitions of Lie bialgebras and Lie quasibialgebras to start with. Let x, y, z be elements of V . Denote $\text{Alt}(x \otimes y \otimes z) = x \otimes y \otimes z + y \otimes z \otimes x + z \otimes x \otimes y$.

Definition 1.2. A Lie bialgebra structure on a vector space V is the following data:

- a: a Lie bracket, $\{\cdot, \cdot\} : V \wedge V \rightarrow V$;
- b: a Lie cobracket, that is an element $\delta : V \rightarrow V \wedge V$, satisfying coJacobi identity:

$$\text{Alt}(\delta \otimes 1)\delta(x) = 0$$

- c: a compatibility condition between $\{\cdot, \cdot\}$ and δ , meaning that δ is a 1-cocycle: $\delta(\{x, y\}) = \{\delta(x), 1 \otimes y + y \otimes x\} + \{1 \otimes x + x \otimes 1, \delta(y)\}$.

Definition 1.3. A Lie quasibialgebra structure on a vector space V is the following data:

- a: a Lie bracket $\{\cdot, \cdot\}$;
- b: an element $\delta \in \text{Hom}(V, V \wedge V)$, and an element $\phi \in V \wedge V \wedge V$ satisfying a modified coJacobi identity:

$$\frac{1}{2} \text{Alt}(\delta \otimes 1)\delta(x) = [x \otimes 1 \otimes 1 + 1 \otimes x \otimes 1 + 1 \otimes 1 \otimes x, \phi]$$

$$\text{and } (\delta \otimes 1 \otimes 1 + 1 \otimes \delta \otimes 1 + 1 \otimes 1 \otimes \delta)(\phi) = 0.$$

- c: δ is a 1-cocycle.

Good reference on Lie (quasi)bialgebras is a book by Etingof and Shiffmann ([4], pages 32–34 and 150–152). The notion of (finite dimensional) Lie bialgebra is equivalent to the notion of a Manin triple.

Definition 1.4. A Manin triple $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ is a triple of finite dimensional Lie algebras, where $\mathfrak{g}_+ \oplus \mathfrak{g}_- = \mathfrak{g}$ as a vector space and $\mathfrak{g}$ is equipped with a nondegenerate symmetric invariant bilinear form $\langle \cdot, \cdot \rangle$ such that $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are Lagrangian subalgebras (that is maximal isotropic subspaces which are Lie subalgebras).

Lie quasi-bialgebras turn out to be described by the notion of a Manin pair.

Definition 1.5. A Manin pair is a pair $(\mathfrak{g}, \mathfrak{g}_+)$ where $\mathfrak{g}$ is a finite dimensional Lie algebra equipped with a non-degenerate symmetric invariant bilinear form $\langle \cdot, \cdot \rangle$ and $\mathfrak{g}_+$ is a Lagrangian subalgebra.

The main theorem here is by Drinfeld which states that Manin pairs are in one-to-one correspondence with equivalence classes (under twisting) of Lie quasi-bialgebras.

It should be mentioned that properade (or PROP) approach is not used in this paper and it will be really interesting to make explicite connections with the work of Bruno Vallette [15].

We use the Koszul sign convention: in a graded algebra whenever there is a change of places of its elements there should be a corresponding sign. Most of the time we do not write it explicitly, instead we write “ $(-1)^{sgn}$ ”.

Throughout this paper, the summation convention is understood: indices $\alpha, \beta, \dots$ once as superscript and once as subscript in a formula are to be summed over.

Acknowledgements. My interest in homotopy structures flourished when I was a postdoc in Strasbourg with Jean-Louis Loday. One of his inventions, his Leibniz algebras, I encounter over and over again, and in particular in this article. I consider myself extremely lucky to have profited from his amazing intellectual generosity and his joy of mathematics.

I thank Ezra Getzler for useful discussions. I am grateful to Vladimir Hinich who helped me to see cohomological implications and encouraged me to write it up. I am indebted to Boris Tsygan for a discussion crucial to the completion of this article (in particular for Remark 4.3 and for $\hbar$) and also for making my work look more exciting by pointing out possible connections to [5].

2. KOSMANN-SCHWARZBACH'S BIG BRACKET

In 1991 Yvette Kosmann-Schwarzbach published an article [6] where the notion of a so called big bracket was introduced in order to describe proto-bialgebras (a notion generalizing Lie bialgebras). It turns out that this big bracket defines, in particular, the Lie structure of the governing dgLie algebra of Lie bialgebras (and, in fact, Lie algebras, Lie coalgebras and Lie quasibialgebras).

Here is the construction from [6] in a graded context.

On any $\mathbb{Z}$ -graded space $X = \sum X^i$ there is an operation called desuspension, mapping X to the space $X[1]$ so that $X[1]_i = X_{i+1}$. Let $\tilde{V}$ be a finite dimensional k -vector space ($k = \mathbb{R}$ or $\mathbb{C}$), and $\tilde{V}^*$ its dual. A non-graded space $\tilde{V}^* \oplus \tilde{V}$ could be considered as a graded one by assigning a 0-degree to each of its elements. Consider $V^* \oplus V = (\tilde{V}^* \oplus \tilde{V})[1]$, this means that “points” of $V^* \oplus V$ are in degree 1. The purpose of this shift of the degree is to see symmetric powers of the space $\tilde{V}^* \oplus \tilde{V}$ as exterior powers of the shifted space $V^* \oplus V$.

Algebraic (symmetric polynomial) functions on $\tilde{V} \oplus \tilde{V}^*$ are

(2.1)

$$B = \Lambda(V^* \oplus V) = \bigoplus_{j \geq -2} B^j, \quad \text{where for } B^j = \bigoplus_{p+q=j} (\Lambda^{p+1} V^* \otimes \Lambda^{q+1} V), \quad j \geq -1, \quad \text{and } B^{-2} = k.$$

From this point of view B is an algebra of exterior powers of the odd space $V^* \oplus V$.

Grading in B is given by the sum $p + q$. That is, B^j consists of terms of the diagonal j , with $j = p + q$, for $p, q \geq -1$, so that first few terms are as follows:

$$\begin{aligned} B^{-2} &= k, \\ B^{-1} &= V \oplus V^*, \\ B^0 &= V \wedge V \oplus V^* \otimes V \oplus V^* \wedge V^*, \\ B^1 &= V \wedge V \wedge V \oplus V^* \otimes V \wedge V \oplus V^* \wedge V^* \otimes V \oplus V^* \wedge V^* \wedge V^*. \end{aligned}$$

Let us write the elements of B in the table, using the bigrading on the number of V and V^* involved:

		...				
2		$V^* \wedge V^* \wedge V^*$	...			
1		$V^* \wedge V^*$	$V^* \wedge V^* \otimes V$	...		
0		V^*	$V^* \otimes V$	$V^* \otimes V \wedge V$	...	
-1		k	V	$V \wedge V$	$V \wedge V \wedge V$	...
$p \quad / \quad q$		-1	0	1	2	

Let $\langle \cdot, \cdot \rangle$ be the natural pairing of $\tilde{V}$ and $\tilde{V}^*$. We extend it to a symmetric form on $\tilde{V} \oplus \tilde{V}^*$ as follows: for $x, y \in \tilde{V}$ and $v, w \in \tilde{V}^*$:

$$\langle x + v, y + w \rangle = \langle x, w \rangle + \langle v, y \rangle.$$

This symmetric form on $\tilde{V} \oplus \tilde{V}^*$ could be considered as an antisymmetric form on the desuspended space $V \oplus V^*$ and more generally as a Lie algebra structure on B .

Definition 2.1. *The big bracket is the graded Lie algebra structure on algebraic functions on $(V \oplus V^*)$ defined as follows.*

- For $u, v \in B^{-2} \oplus B^{-1} = k \oplus V \oplus V^*$:

$$[u, v] = \langle u, v \rangle.$$

In other words,

$$[u, v] = \begin{cases} \langle u, v \rangle & \text{if } u \in V \oplus V^* \text{ and } v \in V \oplus V^* \\ 0 & \text{if } u \in k \text{ or } v \in k \end{cases}$$

- The bracket on other terms is defined by linearity and the graded Leibniz rule: for $u \in B^k$, $v \in B^l$, $w \in B^m$

$$[u, v \wedge w] = [u, v] \wedge w + (-1)^{kl} v \wedge [u, w]$$

Remark 2.1. *Notice that*

- 1) $[u, v] = -(-1)^{kl}[v, u]$; $u \in B^k$, $v \in B^l$, that is the big bracket is skew-symmetric in the graded sense;
- 2) $[\cdot, \cdot] : B^i \wedge B^j \rightarrow B^{i+j}$, for $i, j \geq -1$, in other words: the bracket is of degree zero;
- 3) moreover,

$$[\cdot, \cdot] : (\Lambda^k V^* \otimes \Lambda^l V) \wedge (\Lambda^m V^* \otimes \Lambda^n V) \longrightarrow \Lambda^{k+m-1} V^* \otimes \Lambda^{l+n-1} V;$$

- 4) B^0 is a Lie subalgebra of B ;
- 5) B^{-2} is the center of $(B, [\cdot, \cdot])$.

3. GOVERNING GRADED LIE ALGEBRAS AND MAURER-CARTAN EQUATIONS

Graded Lie algebra B has several graded Lie subalgebras. Verification that they are indeed Lie subalgebras is easy using Remark 2.1 (3). Some of these Lie subalgebras describe well known structures. We arrange them in the following proposition with a table of governing Lie algebras with the first three graded components:

$\mathfrak{g}$ - Lie subalgebra of B	$\mathfrak{g}^{-1}$	$\mathfrak{g}^0$	$\mathfrak{g}^1$	Solutions of Maurer-Cartan equation
$\mathcal{C}$: row $p = 0$	$u^* \in V^*$	$f \in V^* \otimes V$	$c \in V^* \otimes V \wedge V$	$[c, c] = 0$ Lie coalgebra structure on V
$\mathcal{L}$: column $q = 0$	$u \in V$	$f \in V^* \otimes V$	$l \in V^* \wedge V^* \otimes V$	$[l, l] = 0$ Lie algebra structure on V
$\mathcal{B}$: $p, q \neq -1$	—	$f \in V^* \otimes V$	$c \in V^* \otimes V \wedge V \oplus$ $l \in V^* \wedge V^* \otimes V$	$[c + l, c + l] = 0$ Lie bialgebra structure on V
$\mathcal{QB}$: $q \neq -1$	$u \in V$	$f \in V^* \otimes V \oplus$ $g \in V \wedge V$	$c \in V^* \otimes V \wedge V \oplus$ $l \in V^* \wedge V^* \otimes V \oplus$ $\phi \in V \wedge V \wedge V$	$[c + l + \phi, c + l + \phi] = 0$ Lie quasi-bialgebra structure on V

Let ΛV denote the sum $k \oplus V \oplus \Lambda^2 V \oplus \Lambda^3 V \oplus \dots$. Then $\mathcal{C} = V^* \otimes \Lambda V \simeq \text{Hom}(V, \Lambda V)$, and $\mathcal{L} = \Lambda V^* \otimes V \simeq \text{Hom}(\Lambda V, V)$.

In other words,

Proposition 3.1. *On a vector space V*

Coalgebras Lie: $\mathcal{C} = V^* \otimes (\oplus_{l \geq 0} \Lambda^l V)$ is the governing graded Lie algebra of Lie coalgebra structures on V . In particular, an element of degree 1, c such that $[c, c] = 0$ defines a Lie coalgebra structure $\delta \in \text{Hom}(V, V \wedge V)$ as follows: $\delta(x) = [c, x]$.

Algebras Lie: $\mathcal{L} = \oplus_{k \geq 0} \Lambda^k V^* \otimes V$ is the governing graded Lie algebra of Lie algebra structures on V . In particular, an element of degree 1, l such that $[l, l] = 0$ defines a Lie algebra structure $\{\cdot, \cdot\} \in \text{Hom}(V \wedge V, V)$ as follows: $\{x, y\} = [[l, x], y]$.

Lie bialgebras: $\mathcal{B} = \oplus_{k \geq 0, l \geq 1} \Lambda^k V^* \otimes \Lambda^l V$ is the governing graded Lie algebra of Lie bialgebra structures on V . In particular, elements of degree 1, $\theta \in V^* \otimes V \wedge V$ and $l \in V^* \wedge V^* \otimes V$ such that $[c + l, c + l] = 0$ define a Lie bialgebra structure with the cobracket $\delta(x) = [c, x]$ and the bracket $\{x, y\} = [[l, x], y]$.

Lie quasi-bialgebras: $\mathcal{QB} = \oplus_{k \geq 0} \Lambda^k V^* \otimes \oplus_{l \geq 1} \Lambda^l V$ is the governing graded Lie algebra of Lie quasibialgebra structures on V . In particular, elements of degree 1, $c \in V^* \otimes V \wedge V$, $l \in V^* \wedge V^* \otimes V$ and $\phi \in V \wedge V \wedge V$ such that

$$(3.1) \quad [c + l + \phi, c + l + \phi] = 0$$

define a Lie quasibialgebra structure with the 1-cocycle $\delta(x) = [c, x]$, the Lie bracket $\{x, y\} = [[l, x], y]$ and the 3-form ϕ .

Proof. Let us write out in coordinates the condition for $\delta \in \text{Hom}(V, V \wedge V)$ to be a Lie cobracket. Let $\{e_i\}$ constitute a basis in V , and $\{e^i\}$ the dual basis in V^* . Then a Lie cobracket with structure coefficients c_k^{ij} :

$$\delta(e_k) = c_k^{ij} e_i e_j = [c_l^{ij} e_i e_j e^l, e_k]$$

(we omit the signs of $\wedge$ and $\otimes$). The condition $[c, c] = 0$ thus becomes a condition on structure coefficients c_k^{ij} of a Lie coalgebra:

$$\langle c_k^{ij} e_i e_j e^k, c_k^{ij} e_i e_j e^k \rangle = (c_m^{ij} c_l^{mk} + c_m^{jk} c_l^{mi} + c_m^{ki} c_l^{mj}) e_i e_j e^l = 0,$$

which is exactly the coJacobi identity on $\delta = ad_c$, where $c = c_l^{ij} e_i e_j e^l$.

Similarly, in the case of Lie algebra structure, the condition $[l, l] = 0$ is exactly the Jacobi identity.

The commutator $[l + c, l + c]$ lies in $B^2 : \Lambda^3 V^* \otimes V, V^* \otimes \Lambda^3 V$, and $\Lambda^2 V^* \otimes \Lambda^2 V$. Then $l + c$ defines a Lie bialgebra structure if the commutator is 0, that is all three components of it in B^2 must be equal to 0. This leads to three axioms of a Lie bialgebra:

- a:** Jacobi identity follows from $[l, l] = 0$, for the derived bracket given by $\{x, y\} = [[l, x]y]$.
- b:** Co-Jacobi identity follows from $[c, c] = 0$.
- c:** The cocycle condition translates as: $[l, c] = 0$.

In fact, a presentation of a Lie bialgebra as a square zero element in $\Lambda(V^* \oplus V)$ appeared already in [10].

In the same manner the equation 3.1 gives independent equations: $[l, l] = 0$, $[c, c] + 2[l, \phi] = 0$, $[c, \phi] = 0$ and $[c, l] = 0$, which give the axioms of a Lie quasibialgebra.

One could look for complete proves in [6]. $\square$

In all the cases of Proposition 3.1 we get a graded Lie algebra with a differential d given by the adjoint action of an auto-commuting element in the first degree. Hence $\mathcal{C}, \mathcal{L}, \mathcal{B}$ and $\mathcal{QB}$ become dgLie algebras. On a dgLie algebra one can define a derived bracket. Here is a general fact:

Proposition 3.2. [7] *Differential Graded Lie algebra $(\mathfrak{g}, [\cdot, \cdot], D)$ gives rise to a new bracket of degree +1, called a derived bracket: for $a, b \in \mathfrak{g}$, $\{a, b\} = [Da, b]$. This is a Loday's Leibniz algebra [11]. If W is an Abelian subalgebra of $\mathfrak{g}$, such that $[DW, W]$ is in W , then $\{\cdot, \cdot\}$ is a Lie bracket on W . Moreover, $D : (W, \{\cdot, \cdot\}) \rightarrow (\mathfrak{g}, [\cdot, \cdot])$ is a Lie algebra morphism.*

In our case, the algebra $\mathfrak{g}$ is one of $\mathcal{C}, \mathcal{L}, \mathcal{B}$ and $\mathcal{QB}$, while the subalgebra Lie W in all cases is the same $B^{-1} = V^* \oplus V$. The derived bracket with corresponding differentials on $\mathcal{C}, \mathcal{L}, \mathcal{B}$ will define the Lie bracket on $V^* \oplus V$, leading to Manin triples (we could see a Lie structure on V as a particular case of a Manin triple with a zero cobracket, as well as a Lie coalgebra structure - a Manin triple with an Abelian Lie structure). In the same way, we get a Manin pair from $\mathcal{QB}$.

However, we need a certain refinement of Proposition 3.2, since the Lie subalgebra B^{-1} is not Abelian.

Proposition 3.3. *Differential Graded Lie algebra $(B, [\cdot, \cdot], d)$, gives rise to a derived bracket of degree 1 : $\{a, b\} = [da, b]$.*

This derived bracket restricted on B^{-1} is a Lie bracket and $d : (B^{-1}, \{\cdot, \cdot\}) \rightarrow (B^0, [\cdot, \cdot])$ is a Lie algebra morphism.

Proof. Notice that $\{B^{-1}, B^{-1}\} \subset B^{-1}$. The subspace $B^{-1} = V^* \oplus V$ is not an Abelian Lie subalgebra of $(B, [\cdot, \cdot])$, and we cannot use Proposition 3.2. However, since the bracket $[\cdot, \cdot]$ on B^{-1} takes values in the center of B (namely, $B^{-2} = k$), the Jacobi identity on B gives us the skew-symmetry of the derived bracket $\{\cdot, \cdot\}$. Hence the derived bracket defines a Lie algebra structure (not just Loday's) on $V^* \oplus V$.

The Lie algebra morphism part is immediate: $da, b = d[da, b] = [da, db]$. $\square$

Depending on the terms involved in d we get several known cohomology theories:

Proposition 3.4. *Let us take the Lie algebra $B = \oplus_{p \geq -1, q \geq -1} \Lambda^{p+1} V^* \otimes \Lambda^{q+1} V$ with a differential*

$$d = ad_{\lambda + \theta + \phi + \psi}, \quad d \circ d = 0, \quad l \in \Lambda^2 V^* \otimes V, \quad c \in V^* \otimes \Lambda^2 V, \quad \phi \in \Lambda^3 V, \quad \psi \in \Lambda^3 V^*.$$

Cohomology of the Differential Graded Lie algebra $(B, [\cdot, \cdot], d)$ gives rise to

- the Lie algebra cohomology defined with the differential $d = ad_l$, on its Lie subalgebra

$$\mathcal{L} = \bigoplus_{p \geq 0} \Lambda^{p+1} V^* \otimes V$$

$$(c = 0, \phi = 0, \psi = 0),$$

- the Lie coalgebra cohomology for $d = ad_c$, on the Lie subalgebra

$$\mathcal{C} = \bigoplus_{q \geq 0} V^* \otimes \Lambda^{q+1} V$$

$$(l = 0, \phi = 0, \psi = 0),$$

- the Lie bialgebra cohomology for $d = ad_{l+c}$ on the Lie subalgebra

$$\mathcal{B} = \bigoplus_{p+q \geq 0} \Lambda^{p+1} V^* \otimes \Lambda^{q+1} V$$

$$(\phi = 0, \psi = 0),$$

- the quasi Lie bialgebra cohomology for $d = ad_{l+c+\phi}$ on the Lie subalgebra

$$\mathcal{QB} = \bigoplus_{p \geq -1, q \geq 0} \Lambda^{p+1} V^* \otimes \Lambda^{q+1} V$$

$$(\psi = 0).$$

Finally, to make explicit the connection to Manin pairs and triples we have the following

Proposition 3.5. *Let $l \in \Lambda^2 V^* \otimes V$, $c \in V^* \otimes \Lambda^2 V$, $\phi \in \Lambda^3 V$. Then $(V \oplus V^*, \{a, b\} = [da, b])$, for*

- $d = ad_{l+c}$, with a condition $[l+c, l+c] = 0$ defines a Manin triple $(V \oplus V^*, V, V^*)$.
- $d = ad_{l+c+\phi}$, satisfying $[l+c+\phi, l+c+\phi] = 0$ defines a Manin pair $(V \oplus V^*, V)$.

4. L_∞ STRUCTURES

In the previous section we have seen that a Lie algebra structure on a space V is obtained as a derived bracket defined on the graded Lie algebra $\mathcal{L}$ with a differential given by an adjoint action of an element from $V^* \wedge V^* \otimes V \subset B^1$ whose bracket with itself is 0. This way we are in the setup of Definition 1.1.

If we take an element from $\mathcal{L} = \sum_{k=0}^\infty \Lambda^k V^* \otimes V$ (not just from $V^* \wedge V^* \otimes V$) whose bracket with itself is 0 we could define an L_∞ structure on V using higher derived brackets of Th. Voronov and Akman-Ionescu ([17, 1]).

The same idea works for other Lie subalgebras of B : an autocommuting element from $\mathcal{B}$ (or $\mathcal{QB}$) defines an L_∞ bialgebra (or L_∞ quasi-bialgebra) structure. Here we develop this theory.

A certain subtlety here is in defining what an element of degree 1 would mean in this context. For that we need to introduce a new notion of degree.

4.1. Degree. A starting point of any homotopy construction is a graded space with a differential of degree 1 (a complex). Let the space V be graded. Then the space B , as well as the space of maps from $\Lambda V \rightarrow \Lambda V$, inherit the grading from V in a consistent way. We will define a grading and a differential on these spaces as follows.

A differential graded $(V = \bigoplus_a V_a, \delta)$, space over an ungraded field $\mathbb{K}$ is in other words a complex

$$(4.1) \quad \cdots \xrightarrow{\delta} V_a \xrightarrow{\delta} V_{a+1} \xrightarrow{\delta} \cdots$$

Lemma 4.1. *Let $(V = \bigoplus_a V_a, \delta)$ be a graded differential space over an ungraded ground field $\mathbb{K}$. Assume that each V_a is finite dimensional. Then on the dual space V^* there is a corresponding grading with the opposite sign and a differential (also denoted δ) of degree 1, $\delta : V^*_{-(a+1)} \rightarrow V^*_{-a}$:*

$$(4.2) \quad \cdots \xleftarrow{\delta} V^*_{-a} \xleftarrow{\delta} V^*_{-(a+1)} \xleftarrow{\delta} \cdots$$

Proof. If v is in V_a and $u = \delta v$ then their dual elements $v^*, u^* \in V^*$ are related as follows: $v^* = \delta u^*$. If u has a degree a then degree of $v = \delta u$ is $a + 1$, while degree of v^* should be $-(a + 1)$ and of u^* then is of $-a$. This relation is obtained from the condition:

$$(4.3) \quad [v^*, v] = 1 = [v^*, \delta u] = [\delta v^*, u] = [u^*, u].$$

If $\delta : V \rightarrow V$ given by an adjoint action of a corresponding element $d \in V^* \otimes V$, $\delta = ad_d : V \rightarrow V$ or $ad_d : V^* \rightarrow V^*$ then the condition $[v^*, \delta u] = [\delta v^*, u]$ from 4.3 is just a consequence of the Jacobi identity and a fact that constants are in the center. $\square$

This way $V^* \oplus V$ becomes a complex. We could extend the action of δ on $\Lambda^p V^* \otimes \Lambda^q V$ for any positive p, q by the Leibniz rule, so that the whole B becomes a complex.

Let us define a new grading on the complex B , taking into account the grading on V .

Definition 4.1. Given a graded vector space $V = \oplus_a V_a$, consider elements from $\Lambda^p V^* \otimes \Lambda^q V$. The degree of elements from V^* is defined according Lemma 4.1, and $\Lambda^0 V = \Lambda^0 V^* = \mathbb{K}$.

Then the inner degree of $x_1^* x_2^* \dots x_p^* y_1 y_2 \dots y_q \in \Lambda^p V^* \otimes \Lambda^q V$ is $\sum \widetilde{x_i} + \sum \widetilde{y_i} - (p + q - 2)$ where the degree of each element is denoted $\sim$.

A map $\tau_{pq} : \Lambda^p V \rightarrow \Lambda^q V$ sending $x_1 x_2 \dots x_p$ to $y_1 y_2 \dots y_q$ has the inner degree n if $(\sum \widetilde{y_i}) - (\sum \widetilde{x_j}) = n - (p + q - 2)$, which corresponds to the inner degree of $x_1^* x_2^* \dots x_p^* y_1 y_2 \dots y_q \in \Lambda^p V^* \otimes \Lambda^q V$, where $[x_i^*, x_i] = 1$.

We will now define solutions of Maurer-Cartan equation of inner degree 1 from B . It is this correction of $(p + q - 2)$ which takes into account that now elements participating in the Maurer-Cartan equation are not necessarily from B^1 (that is when $p + q = 3$) but from any B^{p+q-2} .

4.2. L_∞ algebra structure from multiple adjoint action. For a finite dimensional space V there is a following isomorphism $\text{Hom}(\Lambda^k V, \Lambda^l V) \simeq \Lambda^k V^* \otimes \Lambda^l V$. Any operator $\tau_{kl} \in \text{Hom}(\Lambda^k V, \Lambda^l V)$ is represented by $t_{kl} \in \Lambda^k V^* \otimes \Lambda^l V$ in the following way:

Definition 4.2. Consider an element $t_{kl} \in \Lambda^k V^* \otimes \Lambda^l V$. Multiple adjoint action of t_{kl} on $\Lambda^n V$ is defined as follows. For $n = k$ it defines a higher derived bracket as in [1, 16]:

$$\tau_{kl}(v_1 \wedge \dots \wedge v_k) = \mathbf{mad}(t_{kl})(v_1 \wedge \dots \wedge v_k) = [[\dots [t_{kl}, v_1], v_2] \dots], v_k],$$

thus defining a map $\Lambda^k V^* \otimes \Lambda^l V \rightarrow \text{Hom}(\Lambda^k, \Lambda^l V) : t_{kl} \mapsto \tau_{kl}$. More generally, we define an action of t_{kl} on any exterior power of V , by making it a map $\Lambda V \rightarrow \Lambda V^* \otimes \Lambda V$ as follows

$$(4.4) \quad \begin{aligned} \mathbf{mad}(t_{kl}) & (v_1 \wedge \dots \wedge v_n) \\ &= \begin{cases} [[\dots [t_{kl}, v_1], v_2] \dots], v_n], & \text{if } k \geq n \\ \sum_{\sigma \in Sh_n^k} (-1)^{sgn\sigma} \underbrace{[\dots [t_{kl}, v_{\sigma(1)}], v_{\sigma(2)}] \dots]_{k \text{ times}}}_{k \text{ times}}, v_{\sigma(k)}] \wedge v_{\sigma(k+1)} \dots \wedge v_{\sigma(n)} & \text{otherwise.} \end{cases} \end{aligned}$$

The set of permutations Sh_n^k is as in Definition 4.3. The case $n = k$ fits with keeping in mind that $\Lambda^0 V^* = \mathbb{K}$.

It is shown already in [17] that L_∞ -algebra could be obtained as a particular case of this multiple adjoint action. First we recall an L_∞ -algebra definition here.

Definition 4.3. Consider a map $\lambda_k \in \text{Hom}(\Lambda^k V, V)$. This map λ_k acts on $\Lambda^n V$ for any $n \in \mathbb{N}$ by coderivation of the unshuffle coproduct on the algebra of exterior powers of V :

$$(4.5) \quad \lambda_k(v_1 \wedge \dots \wedge v_n) = \begin{cases} 0, & \text{if } k > n \\ \sum_{\sigma \in Sh_n^k} (-1)^{sgn\sigma} \lambda_k(v_{\sigma(1)} \wedge \dots \wedge v_{\sigma(k)}) \wedge v_{\sigma(k+1)} \dots \wedge v_{\sigma(n)} & \text{otherwise.} \end{cases}$$

The set Sh_n^k is a set of all k -unshuffles in the permutation group of n elements, that is such permutations that have $\sigma(1) < \sigma(2) < \dots < \sigma(k)$ and $\sigma(k+1) < \dots < \sigma(n)$.

Definition 4.4. An L_∞ -structure on a graded space V is a set of maps of inner degree 1,

$$\lambda_p : \Lambda^p V \rightarrow V[2-p], \quad p \in \mathbb{N}$$

such that the following generalized form of Jacobi identity is satisfied for any $n \geq 2$:

$$(4.6) \quad \sum_{p=1}^{n-1} [\lambda_p, \lambda_{n-p}] = 0.$$

The bracket $[\cdot, \cdot]$ here is a commutator of λ_p and λ_{n-p} considered as operators on $\Lambda^n V$. These equations are higher Jacoby identities and could be summarized in one equation:

$$(4.7) \quad \lambda^2 = 0, \quad \text{where } \lambda = \sum_{p \geq 1} \lambda_p$$

We now state two theorems which are L_∞ analogues of results of Proposition 3.2.

Theorem 4.1. (see [17]) Consider an element $L \in \mathcal{L}$, $L = \sum_{k=1}^{\infty} l_k$, $l_k \in \Lambda^k V^* \otimes V$ of inner degree 1, such that $[L, L] = 0$. The set of maps $\lambda_k = \mathbf{mad} \, l_k \in \text{Hom}(\Lambda^k V, V)$ form an L_∞ -algebra.

In other words, $[L, L] = 0$ implies that $\lambda^2 = 0$ where $\lambda = \mathbf{mad} \, L$.

The proof is a direct albeit tedious computation reducing the condition on the maps λ_k to $[L, L] = 0$ using the fact that ΛV is an Abelian Lie subalgebra of $\mathcal{L}$ and also that $[L, v] \in \Lambda V^*$ for any $v \in \Lambda V^*$.

We could also prove a statement analogous to the one about the Lie algebra morphism in Proposition 3.2.

Definition 4.5. An L_∞ -morphism from an L_∞ -algebra $(V, \sum \lambda_k)$ to a Lie algebra $(W, [\cdot, \cdot])$ is a series of maps $\phi_l : \Lambda^l V \rightarrow W$ of inner degree 0 such that for all $n \geq 2$ we have the following equality for all $n \geq 2$:

$$\sum_{k+l=n} \phi_l \lambda_k = \sum_{k+l=n+1} [\phi_k, \phi_l]$$

Proposition 4.1. Consider a graded space V with an L_∞ -structure given by the multiple adjoint action of $L = \sum l_k$, $l_k \in \Lambda^k V^* \otimes V$, $[L, L] = 0$. Consider a graded Lie algebra structure on $V^* \otimes V$ with a bracket $[\cdot, \cdot]$ given by the natural coupling.

Then maps

$$\mathbf{mad} \, (l_{k+1}) : \Lambda^k V \rightarrow V^* \otimes V$$

define an L_∞ morphism from $(V, \sum \mathbf{mad} \, (l_k)) \rightarrow (V^* \otimes V, [\cdot, \cdot])$.

Checking this proposition amounts just to writing the higher Jacobi identities (4.6) on V .

4.3. L_∞ -coalgebras.

Definition 4.6. L_∞ coalgebra structure on V is a sequence of maps

$$\delta_{1l} : V \rightarrow \Lambda^l V \quad [2-l], \quad l \geq 1$$

of inner degree 1 such that

$$\sum_{p \geq 1, q \geq 1} \delta_{1p} \delta_{1q} = 0.$$

In other words,

$$\delta^2 = 0, \quad \text{where } \delta = \sum_{p \geq 1} \delta_{1p}$$

Consider the Lie algebra governing the Lie coalgebra structure on $V : \mathcal{C} = \oplus_{q \geq 0} V^* \otimes \Lambda^{q+1}V$. The adjoint action of $c_{1l} \in V^* \otimes \Lambda^l V$ on $\Lambda^n V$ is a map from $\Lambda^n V$ to $\Lambda^{n+l-1}V$:

$$ad_{c_{1l}}(v_1 \wedge \cdots \wedge v_n) = \sum_{i=1}^n (-1)^{sgn} [c_{1l}, v_i] \wedge v_1 \wedge \cdots \check{v}_i \cdots \wedge v_n$$

Theorem 4.2. *Let*

$$C = \sum_{l=1}^{\infty} c_{1l}, \quad c_{1l} \in V^* \otimes \Lambda^l V$$

such that $[C, C] = 0$. Then the adjoint action of C defines an L_{∞} -coalgebra structure on V .

Moreover, the multiple adjoint action of C on ΛV^ defines an L_{∞} -algebra structure on V^* , the dual space to V .*

Proof. First statement follows from the definition of an L_{∞} coalgebra, while an L_{∞} structure on V^* is a consequence of Theorem 4.1. $\square$

4.4. L_{∞} -bialgebras. Following the general philosophy of structures given by solutions of Maurer-Cartan equations on differential graded Lie algebras, we consider solutions of an equation $[Q, Q] = 0$ where Q belongs to Lie subalgebras of $B : \mathcal{C}, \mathcal{L}, \mathcal{B}$ and $\mathcal{QB}$ (3.4). They respectively define an L_{∞} -coalgebra, L_{∞} -algebra, L_{∞} -bialgebra and L_{∞} -quasibialgebra structures on V . In all cases we consider the multiple adjoint action 4.2 of corresponding Q on ΛV .

Definition 4.7. *An L_{∞} bialgebra structure on V is a set of maps*

$$\tau_{kl} : \Lambda^k V \rightarrow \Lambda^l V[3 - (k + l)], \quad k, l \geq 1$$

of inner degree 1 such that each $\tau_{kl} = \mathbf{mad} \, t_{kl}$, $t_{kl} \in \Lambda^k V^ \otimes \Lambda^l V$ and*

$$(4.8) \quad \sum_{k+k'=p, \quad l+l'=q} [t_{kl}, t_{k'l'}] = 0$$

for all $p \geq 2, q \geq 2$.

In other words,

Definition 4.7.1. *Consider an element of $\mathcal{B}$ of inner degree 1 :*

$$(4.9) \quad T = \sum_{k \geq 1, \quad l \geq 1} t_{kl}, \quad t_{kl} \in \Lambda^k V^* \otimes \Lambda^l V$$

such that

$$(4.10) \quad [T, T] = 0.$$

Then the set of operators $\tau_{kl} = \mathbf{mad} \, (t_{kl})$ defines an L_{∞} -bialgebra structure on V .

We could see that T lives in the following double graded space:

		...				
4	$\Lambda^4 V^* \otimes V$	...				
3	$\Lambda^3 V^* \otimes V$	$\Lambda^3 V^* \otimes \Lambda^2 V$	...			
2	$\Lambda^2 V^* \otimes V$	$\Lambda^2 V^* \otimes \Lambda^2 V$	$\Lambda^2 V^* \otimes \Lambda^3 V$	...		
1	$V^* \otimes V$	$V^* \otimes \Lambda^2 V$	$V^* \otimes \Lambda^3 V$	$V^* \otimes \Lambda^4 V$	...	
$k \quad \diagdown \quad l$	1	2	3	4		

This way the condition $[T, T] = 0$ gives a set of equations indexed by two numbers. For each pair $p \geq 1, q \geq 1$ we get an equation:

$$(4.11) \quad \sum_{k+k'=p+1} \sum_{l+l'=q+1} [t_{kl}, t_{k'l'}] = 0,$$

The first one is

$$(4.12) \quad [t_{11}, t_{11}] = 0$$

providing the equation which defines a differential $d = \mathbf{ad}_{t_{11}}$ on V . The next couple of equations for $p = 1, q = 2$ and $p = 2, q = 1$ gives respectively:

$$(4.13) \quad [t_{11}, t_{12}] = 0, \quad [t_{11}, t_{21}] = 0,$$

which is a condition that d is a derivation of the cobracket and of the bracket. Next one $p = 2, q = 2$:

$$(4.14) \quad [t_{11}, t_{22}] + [t_{12}, t_{21}] = 0$$

is the condition that the bracket is the cocycle with respect to the cobracket satisfied up-to homotopy. The homotopy is given by an element $t_{22} \in \Lambda^2 V^* \otimes \Lambda^2 V$.

The Jacobi identity on the bracket $\mathbf{mad}(t_{21})$ also holds only up-to homotopy, it is given by the equation for $p = 3, q = 1$:

$$(4.15) \quad [t_{11}, t_{31}] + \frac{1}{2}[t_{21}, t_{21}] = 0,$$

as well as the coJacobi identity - equation with $p = 1, q = 3$:

$$(4.16) \quad [t_{11}, t_{13}] + \frac{1}{2}[t_{12}, t_{12}] = 0.$$

Remark 4.1. 1) The homology of V , $H^*(V, d = \mathbf{ad} t_{11})$ is a Lie bialgebra.

2) If V is not graded, that is all $V_n = \emptyset$ for all $n \neq 0$ then $V = V_0$ is an ordinary Lie bialgebra. Indeed, since t_{kl} is of inner degree 1, on an ungraded space only terms with k, l satisfying $1 - (k + l - 2) = 0$ survive. The result is that all t_{kl} except for t_{12} and t_{21} have to vanish. The Maurer-Cartan equation then gives the axioms of a Lie bialgebra from Definition 1.2: $\tau_{21} = \mathbf{mad} t_{21}$ defines a bracket and $\tau_{12} = \mathbf{mad} t_{12}$ a cobracket.

3) If all $t_{kl} = 0$ for all $k + l \geq 3$ then V is a differential graded Lie bialgebra.

4.5. L_∞ -quasibialgebra. To have an L_∞ -quasibialgebra structure we need to allow terms in $\Lambda^q V$. Hence Maurer-Cartan equation of the previous subsection (4.10) becomes an equation on $S = \sum_{k \geq 0, l \geq 1} t_{kl}$, $t_{kl} \in \Lambda^k V^* \otimes \Lambda^l V$ which differs from T contains elements $t_{0l} \in \Lambda^l V$. This S must satisfy a set of equations indexed by $p \geq 1, q \geq 2$:

$$[S, S] = \sum_{k+k'=p} \sum_{l+l'=q} [t_{kl}, t_{k'l'}] = 0,$$

		...				
4		$\Lambda^4 V^* \otimes V$	...			
3		$\Lambda^3 V^* \otimes V$	$\Lambda^3 V^* \otimes \Lambda^2 V$	...		
2		$\Lambda^2 V^* \otimes V$	$\Lambda^2 V^* \otimes \Lambda^2 V$	$\Lambda^2 V^* \otimes \Lambda^3 V$	...	
1		$V^* \otimes V$	$V^* \otimes \Lambda^2 V$	$V^* \otimes \Lambda^3 V$	$V^* \otimes \Lambda^4 V$	...
0		$\mathbf{ad}^2 V$	$\mathbf{ad}^3 V$	$\Lambda^3 V$	$\Lambda^4 V$	...
$(p+1) \quad / \quad (q+1)$		1	2	3	4	

Terms t_{0l} do not act on ΛV , however they change the coJacobi condition and other equations as well in comparison with equations on T .

Moreover, terms $t_{01} \in V$ and $t_{02} \in V \wedge V$ change the nature of certain equations: for example, $[t_{11}, t_{11}] = 0$ is no longer true in the presence of t_{01} . Allowing non-zero terms t_{01} and t_{02} would lead us to completely different setup and to avoid that we should impose that these terms are 0.

Definition 4.8. Consider an element of $\mathcal{B}$ of inner degree 1 :

$$(4.17) \quad S = \sum_{k \geq 1, l \geq 1} t_{kl} + \sum_{l \geq 3} t_{0l}, \quad t_{kl} \in \Lambda^k V^* \otimes \Lambda^l V,$$

such that

$$(4.18) \quad [S, S] = 0.$$

An L_∞ quasi-bialgebra structure on V is a set of maps

$$\tau_{kl} : \Lambda^k V \rightarrow \Lambda^l V[3 - (k + l)], \quad k \geq 0, \quad l \geq 1$$

of inner degree 1 such that $\tau_{kl} = \mathbf{mad} \, t_{kl}$.

Let us state properties of an L_∞ -quasibialgebra similar to Remark 4.1:

Remark 4.2. 1) The homology of $H^*V = H^*(V, \mathbf{ad}t_{11})$ is a Lie quasi-bialgebra.

2) If V is not graded, that is all $V_n = \emptyset$ for all $n \neq 0$ then $V = V_0$ is an ordinary Lie quasibialgebra. Indeed, since t_{kl} is of degree $2 - (k + l - 1)$ all t_{kl} except for t_{12}, t_{21} and t_{03} have to vanish. Then $\mathbf{mad} \, t_{12}$ defines a cobracket $\mathbf{mad} \, t_{21}$ a cobracket and $t_{03} \in \Lambda^3 V$ satisfy the axioms of Lie quasibialgebra from Definition 1.3.

3) If all $t_{kl} = 0$ for all $k + l \geq 3$ then V is a differential graded Lie quasi-bialgebra.

Remark 4.3. For L_∞ -(quasi)bialgebras there is no analogue of Theorems 4.1, 4.2. The condition $[Q, Q] = 0$ in general does not imply $(\mathbf{mad} \, Q)^2 = 0$ for some $Q \in B$. Although it is true for Q either in $\mathcal{L}$ or in $\mathcal{C}$ in a more general case the equation $(\mathbf{mad} \, Q)^2 = 0$ force higher order conditions on Q which are not necessarily satisfied even for an ordinary Lie bialgebra or a Lie quasibialgebra (not L_∞).

5. FORMAL GEOMETRY OF L_∞ -STRUCTURES

We now move to a geometric definition of an L_∞ algebra structure on a finite dimensional space (see for example [2] or [13]). Consider a finite dimensional graded vector space $V = \oplus_k V_k$ with a

chosen basis $\{e_\alpha\}$. Then a Lie algebra structure is defined by structure constants $c_{\alpha,\beta}^\gamma$:

$$[e_\alpha, e_\beta] = \sum c_{\alpha,\beta}^\gamma e_\gamma,$$

satisfying the structure equation which boils down to the Jacobi identity. We could also see the bracket acting from $V \wedge V$ to V as a differential operator $Q_{21} = c_{\alpha,\beta}^\gamma e_\gamma \otimes \frac{\partial}{\partial e_\alpha} \wedge \frac{\partial}{\partial e_\beta}$. We use the index 21 for this operator to underline that it is quadratic in $\frac{\partial}{\partial e}$'s and linear in e 's.

Shifting the degree of V by 1 makes Q_{21} look like a second order differential operator acting on polynomial functions. There is a following natural identification of exterior powers of V with symmetric powers of $V[1]$ namely,

$$(5.1) \quad \text{Sym}^n(V[1]) \simeq \Lambda^n(V)[n].$$

In particular under this identification a Lie bracket $\Lambda^2 V \rightarrow V$ is a map of degree 1 from $\text{Sym}^2(V[1]) \rightarrow V[1][1]$. On the other hand, the dual of the free cocommutative coalgebra $\text{Sym}(V[1])$ could be identified with the algebra of formal polynomials on the $\mathbb{Z}$ graded space $V[1]$. This way a Lie bracket becomes a degree 1 quadratic vector field on it. The dual space to $V[1]$ is $V^*[-1]$. Elements of $\text{Sym}(V^*[-1])$ are coordinate functions on $V[1]$. Let us chose corresponding coordinates $\{x_i\}$. So in the space $V^*[-1]$ x_i has a degree $\tilde{x}_i - 1$.

Very natural generalization of this construction is a definition of an L_∞ algebra structure:

Definition 5.1. *An L_∞ -algebra structure on a graded space V is given by vector field Q of degree 1 on a $\mathbb{Z}$ -graded formal manifold corresponding to $V[1]$ such that $[Q, Q] = 0$.*

Definition 5.2. *(following [Va]) An autocommuting operator (and in particular a vector field) is called homological.*

In coordinates, an L_∞ -algebra structure is given by a homological vector field:

$$Q = b_\alpha^\beta x^\alpha \frac{\partial}{\partial x^\beta} + c_{\alpha\beta}^\gamma x^\alpha x^\beta \frac{\partial}{\partial x^\gamma} + d_{\alpha\beta\gamma}^\delta x^\alpha x^\beta x^\gamma \frac{\partial}{\partial x^\delta} + \dots,$$

$$Q = \sum_{k=1}^{\infty} Q_k, \quad Q_{k1} = f_{\alpha_1 \dots \alpha_k}^\beta x^{\alpha_1} \dots x^{\alpha_k} \frac{\partial}{\partial x^\beta}.$$

In fact, each term in Q corresponds to a map of degree 1 on the dual space:

$$Q = \sum Q_{k1}, \quad Q_{k1} : V^*[-1] \rightarrow \text{Sym}^k(V^*[-1])[1].$$

Hence we are naturally brought to a geometric definition of an L_∞ -bialgebra structure on V . It is given by a differential operator on $V[1]$, which means that it acts on functions on it belonging to $\text{Sym}(V^*[-1])$.

To keep track of degrees, we need to introduce a formal variable $\hbar$ of degree -2 .

We consider the ring of differential operators on $V[1]$, weighted with $\hbar$. That is we consider

$$\text{Sym}(V^*[-1])[\xi^1, \dots, \xi^n, \hbar]$$

where $\xi_i = \hbar \frac{\partial}{\partial x_\alpha}$. We take $x_\alpha \in V^*[-1]$, then the dual space is $V[1]$ and $\xi^\alpha \in V[1][-2] = V[-1]$. Let $Op_\hbar$ be the ring of operators being sums of monomials a map from $\text{Sym} V^*[-1] \otimes \text{Sym} V[1] \rightarrow \text{Sym}(V^*[-1])[\xi^1, \dots, \xi^n, \hbar]$ defined as follows:

$$(5.2) \quad Op(v_1 \dots v_p w_1^* \dots w_q^*) = \frac{1}{\hbar} (v_1 \dots v_p \frac{\partial}{\partial w_1} \frac{\partial}{\partial w_p} \text{ where } v_i \in V^*[-1], w^* \in V[1])$$

Let us call the space of such operators $Op_\hbar$.

On $Op_\hbar$ there is a Poisson bracket defined in coordinates as follows:

$$\{E, F\} = \hbar \left(\frac{\partial E}{\partial x_\alpha} \frac{\partial F}{\partial \xi^\alpha} - (-1)^{s_{gn}} \frac{\partial F}{\partial x_\alpha} \frac{\partial E}{\partial \xi^\alpha} \right)$$

Proposition 5.1. *Then the L_∞ -bialgebra structure on V corresponds to an operator from $Op_{\hbar}$ of degree 1 such that its Poisson bracket with itself is 0.*

Proof. A differential operator of degree 1 on $V[1]$ is sum of maps:

$$Q = \sum_{k,l \geq 1} Q_{pq}, \quad Q_{pq} : \text{Sym}^p(V^*[-1]) \rightarrow \text{Sym}^q(V^*[-1])[1].$$

The grading of an operator Q_{pq} in $Op(v_1 \cdots v_p w_1^* \cdots w_q^*) = \frac{1}{\hbar}(v_1 \cdots v_p \frac{\partial}{\partial w_1} \cdots \frac{\partial}{\partial w_p})$ is $2 + \sum(\tilde{v}_i - 1) + \sum(\tilde{w}_j - 1) = \sum(\tilde{v}_i + \tilde{w}_j) + 2 - (p + q)$, which is exactly what we need to get the right degree.

Thus is after the identification (5.1) we get that $Q_{pq} : \Lambda^p V^* \rightarrow \Lambda^q V^*[1][2 - (p + q)]$ which on the dual space gives $\Lambda^q V[p + q - 3] \rightarrow \Lambda^p V$ leading to the right degrees.

The Poisson bracket being 0 corresponds to the autocummuting condition (4.9). $\square$

6. MANIN L_∞ -TRIPLE AND L_∞ -PAIR

We give a definition of an Manin L_∞ -triple being a natural generalization of a Manin triple.

For this we need a notion of an L_∞ -subalgebra which is immediate in the formal geometry terms:

Definition 6.1. *Let (W, Q) be an L_∞ -algebra then a subspace $V \subset W$ is an L_∞ -subalgebra if V is Q -invariant.*

That is V is an L_∞ -subalgebra of W if the operator Q acting on $V^*[-1]$ lands in $\text{Sym} V^*[-1][1]$. In other words it means that $Q = \sum_{k,l \geq 1} Q_{kl}$ restricted on V consists of maps $V^* \rightarrow \Lambda^k V^*[2 - k]$ or, on the dual it gives an L_∞ structure on V , $Q_{kl} : \Lambda^k V \rightarrow V[2 - k]$.

Definition 6.2. *A finite dimensional Manin L_∞ -triple is a triple of finite dimensional L_∞ -algebras $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$ equipped with a nondegenerate bilinear form $\langle, \rangle$ such that*

- $\mathfrak{g}_+, \mathfrak{g}_-$ are L_∞ -subalgebras of $\mathfrak{g}$ such that $\mathfrak{g} = \mathfrak{g}_+ + \mathfrak{g}_-$ as a vector space;
- $\mathfrak{g}_+$ and $\mathfrak{g}_-$ are isotropic with respect to $\langle, \rangle$;
- the n -brackets constituting the L_∞ -algebras structure $\lambda_n : \Lambda^n(V^* \oplus V) \rightarrow V^* \oplus V$ are invariant with respect to the form $\langle, \rangle$, that is

$$(6.1) \quad \langle \lambda_n(v_1 \wedge \cdots \wedge v_n), v_0 \rangle = (-1)^{\tilde{v}_n \tilde{v}_0} \langle \lambda_n(v_1 \wedge \cdots \wedge v_{n-1} \wedge v_0), v_n \rangle$$

Notice that a Manin triple is an example of a Manin L_∞ -triple.

The invariance is in fact cyclic, since l_n is antisymmetric in all variables being a map on $\Lambda^n V$.

Theorem 6.1. *The notions of a (finite dimensional) Manin L_∞ -triple and an L_∞ -bialgebra are equivalent.*

Proof. Consider a Manin L_∞ -triple $(\mathfrak{g}, \mathfrak{g}_+, \mathfrak{g}_-)$. Let us show that there is an L_∞ -bialgebra structure on $\mathfrak{g}_+$. Let the L_∞ -algebra structure on $\mathfrak{g}$ be given by maps $\lambda_k : \Lambda^k \mathfrak{g} \rightarrow \mathfrak{g}$, that is $\lambda_k : \Lambda^k(\mathfrak{g}_+ \oplus \mathfrak{g}_-) \rightarrow (\mathfrak{g}_+ \oplus \mathfrak{g}_-)$. These maps could be split as follows:

$$\lambda_{mn+} : \Lambda^m \mathfrak{g}_+ \otimes \Lambda^n \mathfrak{g}_- \rightarrow \mathfrak{g}_+ \text{ and } \lambda_{mn-} : \Lambda^m \mathfrak{g}_+ \otimes \Lambda^n \mathfrak{g}_- \rightarrow \mathfrak{g}_- \text{ with } m + n = k.$$

Using nondegeneracy and invariance (6.1) of the inner product $\langle, \rangle$ and also a natural isomorphism $\text{Hom}(U, W) = U^* \otimes W$, we get the equivalences

$$\begin{aligned} \text{Hom}_{\text{inv}}(\Lambda^m \mathfrak{g}_+ \otimes \Lambda^n \mathfrak{g}_-, \mathfrak{g}_+) &\simeq \text{Hom}(\Lambda^m \mathfrak{g}_+, \Lambda^{n+1} \mathfrak{g}_+) \simeq \Lambda^m \mathfrak{g}_+^* \otimes \Lambda^{n+1} \mathfrak{g}_+^* \quad \text{and} \\ \text{Hom}_{\text{inv}}(\Lambda^m \mathfrak{g}_+ \otimes \Lambda^n \mathfrak{g}_-, \mathfrak{g}_-) &\simeq \text{Hom}(\Lambda^{m+1} \mathfrak{g}_+, \Lambda^n \mathfrak{g}_+) \simeq \Lambda^m \mathfrak{g}_+^* \otimes \Lambda^{n+1} \mathfrak{g}_+^*. \end{aligned}$$

Let $l_{mn+} \in \Lambda^m \mathfrak{g}_+^* \otimes \Lambda^{n+1} \mathfrak{g}_+^*$ correspond to λ_{mn+} . Using nondegeneracy of the inner product, we get an isomorphism of $\mathfrak{g}_+^*$ with $\mathfrak{g}_+$ and hence $l_{mn+} \in \Lambda^m \mathfrak{g}_+^* \otimes \Lambda^{n+1} \mathfrak{g}_+$. This way, l_{mn+} would act by a multiple adjoint action:

$$\text{mad}(l_{mn+}) : \Lambda^m \mathfrak{g}_+ \rightarrow \Lambda^{n+1} \mathfrak{g}_+.$$

On the other hand $\lambda_{mn-} : \Lambda^m \mathfrak{g}_+ \otimes \Lambda^n \mathfrak{g}_- \rightarrow \mathfrak{g}_-$ again is given by a multiple adjoint action of an element $l_{mn-} \in \Lambda^{m+1} \mathfrak{g}_+^* \otimes \Lambda^n \mathfrak{g}_+$ (after an identification of $\mathfrak{g}_-^*$ with $\mathfrak{g}_+$) and

$$mad(l_{mn+}) : \Lambda^{m+1} \mathfrak{g}_+ \rightarrow \Lambda^n \mathfrak{g}_+.$$

Now, the $L = \sum_{m,n \geq 1} (l_{mn+} + l_{mn-})$ satisfies $[L, L] = 0$ from the condition on λ .

In the other direction it is actually easier. Let L_∞ -bialgebra structure on V be defined by the multiple adjoint action of $T = \sum_{k \geq 1, l \geq 1} t_{kl}$, where $t_{kl} \in \Lambda^k V^* \otimes \Lambda^l V$. Then the corresponding Manin L_∞ -triple is $(V \oplus V^*, V, V^*)$ with nondegenerate form given by natural coupling of V and V^* .

The L_∞ structure on $V^* \oplus V$ is given by $\tau = mad(T)$. In

$$ad(t_{kl}) : (V^* \oplus V) \rightarrow \Lambda^{k-1} V^* \otimes \Lambda^l V \oplus \Lambda^k V^* \otimes \Lambda^{l-1} V$$

we recognize an L_∞ -coalgebra structure dual to an L_∞ -algebra structure we are looking for.

To get the L_{infy} -subalgebra structure on V we consider the quadratic equations (4.11) for $q = 2$. We get:

$$\sum_{k+k'=p} t_{k1} t_{k'1} = 0$$

in which is set of equations defining an L_∞ -structure (4.1) on V by taking $l_k = t_{k1}$.

The operator $\tau_V = \sum_{k \geq 1} ad(t_{k1}) : V^* \rightarrow \Lambda V^*$ is a restriction of τ on V^* making V an L_∞ -subalgebra of $V^* \oplus V$, since T_V is autocommuting.

In the same way, elements $t_{1l} \in V^* \otimes \Lambda^l V$ satisfy the following quadratic equation (the equations (4.11) for $p = 2$):

$$\sum_{l+l'=q} t_{1l} t_{1l'} = 0$$

which induces maps $mad(t_{1l}) : V \rightarrow \Lambda^l V$, on V thus defining an L_∞ -algebra structure on V^* . $\square$

The notion which corresponds to an L_∞ -quasibialgebra is a Manin L_∞ -pair:

Definition 6.3. A finite dimensional Manin L_∞ -pair is a couple of finite dimensional L_∞ -algebras $(\mathfrak{g} \supset \mathfrak{g}_+)$ equipped with a nondegenerate bilinear form $\langle, \rangle$, invariant as in (6.1), such that $\mathfrak{g}_+$ is an L_∞ -subalgebra of $\mathfrak{g}$ isotropic with respect to $\langle, \rangle$.

There is also a correspondence like in Theorem 6.1:

Theorem 6.2. The notions of a (finite dimensional) Manin L_∞ -pair and an L_∞ -quasibialgebra are equivalent.

The proof is similar to the one of Theorem 6.1.

REFERENCES

- [1] F.Akman, and L.Ionescu, *Higher Derived Brackets and Deformation Theory I* Available at <http://xxx.lanl.gov/abs/math.QA/0504541>
- [2] M. Alexandrov; M. Kontsevich; A. Schwarz; O. Zaboronsky, Internat. J. Modern Phys. A **12** (1997), no. 7, 1405–1429
- [3] N. Ciccoli and L. Guerra, J. Lie Theory **13** (2003), no. 2, 579–590
- [4] P. Etingof and O. Schiffmann, *Lectures on quantum groups*, Internat. Press, Boston, MA, 1998
- [5] Eliashberg, Y.; Givental, A.; Hofer, H. *Introduction to symplectic field theory* GAFA 2000 (Tel Aviv, 1999) Geom. Funct. Anal. 2000, Special Volume, Part II, 560–673.
- [6] Y. Kosmann-Schwarzbach, *em* Jacobian quasi-bialgebras and quasi-Poisson Lie groups in Mathematical aspects of classical field theory (Seattle, WA, 1992), 459–489, Contemp. Math., 132, Amer. Math. Soc., Providence, RI
- [7] Y. Kosmann-Schwarzbach, Ann. Inst. Fourier (Grenoble) **46** (1996), no. 5, 1243–1274
- [8] Y. Kosmann-Schwarzbach, *Quasi, twisted, and all that ... in Poisson geometry and Lie algebroid theory*. The breadth of symplectic and Poisson geometry, 363–389, Progr. Math., 232, Birkhuser Boston, Boston, MA, 2005.
- [9] T. Lada and J. Stasheff, Internat. J. Theoret. Phys. **32** (1993), no. 7, 1087–1103
- [10] P.B.A.Lecomte and C.Roger, C. R. Acad. Sci. Paris Sér. I Math. **310** (1990), 405–410 and **311** (1990), 893–894

- [11] J.-L.Loday, in *Cyclic cohomology and noncommutative geometry (Waterloo, ON, 1995)*, 91–102, Amer. Math. Soc., Providence, RI, 1997
- [12] M.Markl, S.Shnider and J.Stasheff, *Operads in algebra, topology and physics*, Amer. Math. Soc., Providence, RI, 2002
- [13] S.Merkulov, Intern. Math. Research Notices 3 (2000) 147-164
- [14] A.Nijenhuis and R.W.Richardson, Jr., Bull. Amer. Math. Soc. **70** (1964), 406–411
- [15] B. Vallette, *A Koszul duality for props*, (2004) to appear in Transactions of A.M.S. Available at <http://math.unice.fr/~brunov/publications/Koszuldualityforprops.pdf>
- [16] A. Yu. Vaintrob, Lie algebroids and homological vector fields. *Russian Math. Surveys* 52 (1997), no. 2, 428–429
- [17] Th. Voronov, *Higher derived brackets and homotopy algebras*. <http://xxx.lanl.gov/abs/math.QA/0304038>

DEPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ LYON-I, 69622 VILLEURBANNE CEDEX, FRANCE

E-mail address: `ok@alum.mit.edu`