

KÄHLER DECOMPOSITION OF 4-MANIFOLDS

R. İNANÇ BAYKUR

ABSTRACT. In this article we show that every closed oriented smooth 4-manifold can be decomposed into two codimension zero submanifolds (one with reversed orientation) so that both pieces are exact Kähler manifolds with strictly pseudoconvex boundaries and that induced contact structures on the common boundary are isotopic. Meanwhile, matching pairs of Lefschetz fibrations with bounded fibers are offered as the geometric counterpart of these structures. We also provide a simple topological proof of the existence of folded symplectic forms on 4-manifolds.

0. INTRODUCTION

One possible strategy for understanding oriented smooth 4-manifolds is to break them up into more tractable classes of manifolds in a controlled manner. Situated in the intersection of complex, symplectic and Riemannian geometries, Kähler manifolds are the best known candidates to be pieces of such a decomposition. The main goal of this article is to show that this can be achieved for any closed oriented smooth 4-manifold X . We decompose X into two exact Kähler manifolds with strictly pseudoconvex boundaries, up to orientation, such that contact structures on the common boundary induced by the maximal complex distributions are isotopic.

The decomposition gives rise to a globally defined 2-form on X , which we will call a (*nicely*) *folded Kähler structure*, and it belongs to a larger family of 2-forms: *folded symplectic structures*. Cannas da Silva showed in [C] that any closed smooth oriented 4-manifold can be equipped with a folded symplectic form, by using a version of the h-principle defined for folding maps by Eliashberg. In Section 2, we introduce a way to construct some simple examples of folded symplectic 4-manifolds. Afterwards we reprove the existence fact by constructing a folded symplectic form ω for a given handlebody decomposition of X , essentially by means of simple Kirby calculus and contact topology (Theorem 3.1). The main ingredient there will be achiral Lefschetz fibrations, and recent work of Etnyre and Fuller [EF] will play a key role in our construction.

Next, we switch gears, and using several results on compact Stein surfaces and Lefschetz fibrations with bounded fibers (mainly [Ha], [El1], [Go2], [LP], [AO]), we prove the aforementioned decomposition theorem (Theorem 4.2). In fact we obtain a stronger result, as the pieces of this decomposition are actually Stein manifolds with strictly pseudoconvex boundaries. It was first shown by Akbulut and Matveyev in [AM] that a closed oriented smooth 4-manifold X could always be decomposed into Stein pieces, but there was no particular information one could use to argue for matching the induced contact structures

The author was partially supported by NSF Grant DMS0305818.

on the separating hypersurface. Our proof follows an alternative way via open book decompositions, and we conclude that the Stein structures can be chosen to agree on the common contact boundary.

In the last section, we introduce folded Kähler structures, and discuss some properties they enjoy, after showing that all closed oriented smooth 4-manifolds admit them (Theorem 5.2). This improves the folded symplectic existence result, and indeed both structures we construct are shown to be equivalent on the symplectic level. The collection of these discussions yield us to define *folded Lefschetz fibrations*, roughly speaking, pairs of positive and negative Lefschetz fibrations over disks with bounded fibers which agree on the common boundary through induced open book decompositions. We prove that any nicely folded Kähler 4-manifold, possibly after an orientation preserving diffeomorphism, admits compatible folded Lefschetz fibrations (Proposition 5.6).

Acknowledgments. The author would like to thank Ron Fintushel for his support and for helpful conversations. The author would also like to thank Burak Ozbagci for commenting on the draft of this paper.

1. PRELIMINARIES

1.1. Lefschetz fibrations. All manifolds and the maps in this article are assumed to be smooth. A *Lefschetz fibration* on an oriented 4-manifold X , possibly with boundary, is a map $f: X \rightarrow \Sigma$, where Σ is a compact oriented surface, such that each critical point of f lies in the interior of X and has a local model $f(z_1, z_2) = z_1 z_2$, given by orientation preserving charts both on X and Σ . These singularities are obtained by attaching 2-handles to regular fibers with framing -1 with respect to the framing induced by the fiber. An *achiral Lefschetz fibration* is defined the same way, except that the given charts around critical points are allowed to reverse orientation. In other words, the 2-handles can be glued with framing $+1$ with respect to fiber framing, too. Also recall that a Lefschetz pencil is a map $f: X \setminus \{b_1, \dots, b_k\} \rightarrow S^2$, such that around any *base point* b_i it has a local model $f(z_1, z_2) = z_1/z_2$, preserving the orientations, and that f is a Lefschetz fibration elsewhere. An *achiral Lefschetz pencil* is then defined by allowing orientation reversing charts around the base points as well. Critical points or base points with orientation reversing charts are called *negative critical points* or *negative base points*, whereas the other critical points or base points are *positive*. For a detailed treatment of this topic and proofs of some facts quoted below, the reader is advised to turn to [GS].

A Lefschetz fibration is said to be *allowable* if all its vanishing cycles are homologically nontrivial in the fiber. Particularly, we will be interested in allowable Lefschetz fibrations over D^2 with bounded fibers. In the literature, this type of Lefschetz fibration having only positive critical points is called a *PALF*. Similarly, when the critical points are instead all negative, we will call the fibration a *NALF*.

Given a compact oriented genus g surface F with m boundary components and r marked points on it, the *mapping class group* $\Gamma_{g,m}^r$ is defined as the group of orientation preserving self-diffeomorphisms of F fixing marked points and ∂F pointwise, modulo isotopies of F fixing marked points and ∂F pointwise. It can be shown that $\Gamma_{g,m}^r$ is generated by positive (right handed) and negative (left handed) Dehn twists. Importantly, isotopy type of a surface bundle over S^1 with fiber closed oriented surface F is determined by the return

map of a flow transverse to the fibers, which can be identified with an element $\mu \in \Gamma_g$, called *monodromy* of this fibration.

Let $f : X \rightarrow D^2$ be an achiral Lefschetz fibration, where the regular fiber F is an oriented genus g surface with m boundary components, and suppose all critical points of the fibration lie on various fibers. Select a regular value p in the interior of D^2 , an identification of $f^{-1}(p) \cong F$, and a collection of arcs s_1, \dots, s_k in the interior of D^2 with each s_i connecting p to a distinct critical value, and all disjoint except at p . We index the critical values as well, so that each arc s_i is connected to a critical value q_i and that they appear in a counterclockwise order around the point p . Now if we take a regular neighborhood of each arc away from remaining critical points and consider the union of these, we obtain a disk V and an F -bundle over $\partial V = S^1$. The monodromy of this fibration is an element $\mu \in \Gamma_{g,m}$, which is called the *global monodromy* of the achiral Lefschetz fibration f . It is well-known that this data gives a handlebody description of X , and vice versa. We call the ordered set of arcs $\{s_1, \dots, s_k\}$ a *representation* of the achiral Lefschetz fibration f .

Next is a standard fact which was first observed by Harer:

Theorem 1.1 (Harer [Ha]). *Let X be a 4-manifold with boundary. Then X admits an achiral Lefschetz fibration over D^2 with bounded fibers if and only if it admits a handlebody decomposition with no handle of index greater than two.*

1.2. Open book decompositions. An *open book decomposition* of a 3-manifold M is a pair (B, f) where B is an oriented link in M , called the *binding*, and $f : M \setminus B \rightarrow S^1$ is a fibration such that $f^{-1}(t)$ is the interior of a compact oriented surface $F_t \subset M$ and $\partial F_t = B$ for all $t \in S^1$. The surface $F = F_t$, for any t , is called the *page* of the open book. The *monodromy* of an open book is given by the return map of a flow transverse to the pages and meridional near the binding, which is an element $\mu \in \Gamma_{g,m}$, where g is the genus of the page F , and m is the number of components of $B = \partial F$.

Suppose we have an achiral Lefschetz fibration $f : X \rightarrow D^2$ with bounded regular fiber F , and let p be a regular value in the interior of the base D^2 . Composing f with the radial projection $D^2 \setminus \{p\} \rightarrow \partial D^2$ we obtain an open book decomposition on ∂X with binding $\partial f^{-1}(p)$. Identifying $f^{-1}(p) \cong F$, we can write $\partial X = (\partial F \times D^2) \cup f^{-1}(\partial D^2)$. Thus we view $\partial F \times D^2$ as the tubular neighborhood of the binding $B = \partial f^{-1}(p)$, and the fibers over ∂D^2 as its *truncated pages*. The monodromy of this open book is prescribed by that of the achiral fibration [Ha]. In this case, we say the open book $(B, f|_{\partial X \setminus B})$ *bounds* or *is induced by* the achiral Lefschetz fibration $f : X \rightarrow D^2$. Recalling that any closed oriented 3-manifold can be bounded by a 4-manifold with only 0-, 1- and 2- handles, it is fairly easy to see that any open book decomposition bounds such an achiral Lefschetz fibration over a disk.

We would like to describe an elementary modification of these structures: Let $f : X \rightarrow D^2$ be an achiral Lefschetz fibration with bounded regular fiber F . Attach a 1-handle to ∂F to obtain F' , and then attach a positive (resp. negative) Lefschetz 2-handle along an embedded loop in F' that goes over the new 1-handle exactly once. This is called a *positive stabilization* (resp. *negative stabilization*) of f . A positive (resp. negative) Lefschetz handle is attached with framing -1 (resp. $+1$) with respect to the fiber, and thus it introduces a positive (resp. negative) Dehn twist on F' . If the focus is on the 3-manifold, one

can totally forget the bounding 4-manifold and view all the handle attachments in the 3-manifold. Either way, stabilizations correspond to adding canceling handle pairs, so diffeomorphism types of the underlying manifolds do not change, whereas the achiral Lefschetz fibration and the open book decomposition change in the obvious way. It turns out that stabilizations preserve more than the underlying topology, as we will discuss shortly.

1.3. Contact structures and compatibility. A 1-form $\alpha \in \Omega^1(M)$ on a $(2n - 1)$ -dimensional oriented manifold M is called a *contact form* if it satisfies $\alpha \wedge (d\alpha)^n \neq 0$. An *oriented contact structure* on M is then a hyperplane field ξ which can be globally written as kernel of a contact 1-form α . In dimension three, this is equivalent to asking that $d\alpha$ be nondegenerate on the plane field ξ .

A contact structure ξ on a 3-manifold M is said to be *supported by an open book* (B, f) if ξ is isotopic to a contact structure given by a 1-form α satisfying $\alpha > 0$ on positively oriented tangents to B and $d\alpha$ is a positive volume form on every page. When this holds, we say that the open book (B, f) is *compatible with the contact structure* ξ on M .

Improving results of Thurston and Winkelnkemper [TW], Giroux proved the following groundbreaking theorem regarding compatibility of open books and contact structures:

Theorem 1.2 (Giroux [Gi2]). *Let M be a closed oriented 3-manifold. Then there is a one-to-one correspondence between oriented contact structures on M up to isotopy and open book decompositions of M up to positive stabilizations and isotopy.*

Considering contact 3-manifolds as boundaries of certain 4-manifolds together with some compatibility conditions is a current focus of research in low dimensional topology. From the contact topology point of view, it is the study of different types of *fillings* of a fixed contact manifold. In dimension four, there are essentially two considerations, yet we formulate them for all dimensions: Let (X^{2n}, ω) be a symplectic manifold with cooriented nonempty boundary $M = \partial X$. If there exists a *Liouville vector field* (aka *symplectic dilation*) ν defined on a neighborhood of ∂X pointing out along ∂X , then we obtain a positive contact structure ξ on ∂X , which can be written as the kernel of contact 1-form $\alpha = \iota_\nu \omega|_{\partial X}$. When this holds, we say (M, ξ) is the ω -*convex boundary* or *strongly convex boundary* of (X, ω) . For the sake of entirety, note when ν points inside, we obtain a negative contact structure instead, and in this case we say (M, ξ) is the ω -*concave boundary* of (X, ω) .

Now if (X^{2n}, J) is almost-complex, then the complex tangencies on $M = \partial X$ give a uniquely defined oriented hyperplane field. It follows that there is a 1-form α on M such that $\xi = \text{Ker} \alpha$. We define the *Levi form* on M as $d\alpha|_\xi(\cdot, J\cdot)$. If L is positive definite then (M, ξ) is said to be *strictly J -convex boundary* of (X, J) , and if it is J -convex for an unspecified J (for instance when J is tamed by a given symplectic form), we say (M, ξ) is *strictly pseudoconvex boundary*. If (X, ω, J) is an almost-Kähler manifold, i.e. a manifold equipped with a symplectic form ω and a compatible almost-complex structure J , then it can be shown that strict pseudoconvexity of the boundary is equivalent to the condition that $\omega|_\xi > 0$ in dimension $2n = 4$. We would like to remark that all these definitions can be formulated in more generality for hypersurfaces in X^{2n} , not necessarily for ∂X only.

For detailed and comparative discussions of these concepts, as well as proofs of some facts mentioned in the next subsection, the reader can turn to [ElGr] and [E]. Also for further basic notions from contact topology of 3-manifolds such as Legendrian knots, Thurston-Bennequin framing, or convex surfaces, which we will occasionally use in this paper, see for example [OS].

1.4. Stein manifolds. A smooth function $\psi : X \rightarrow \mathbb{R}$ on a complex manifold X of real dimension $2n$ is called *strictly plurisubharmonic* if ψ is strictly subharmonic on every holomorphic curve in X . We call a complex manifold X *Stein*, if it admits a proper strictly plurisubharmonic function $\psi : X \rightarrow [0, \infty)$ (after Grauert [Gra]). Thus a compact manifold X with boundary which is equipped with a complex structure in its interior is called *compact Stein* if it admits a proper strictly plurisubharmonic function which is constant on the boundary.

Given a function $\psi : X \rightarrow \mathbb{R}$ on a Stein manifold, we can define a 2-form $\omega_\psi = -dJ^*d\psi$. It turns out that ψ is a strictly plurisubharmonic function if and only if the symmetric form $g_\psi(\cdot, \cdot) = \omega_\psi(\cdot, J\cdot)$ is positive definite. So every Stein manifold X admits a Kähler structure ω_ψ , for any strictly plurisubharmonic function $\psi : X \rightarrow [0, \infty)$. It is easy to see that the restriction of ω_ψ to each level set $\psi^{-1}(t)$ gives a Levi form on $\psi^{-1}(t)$, implying that all nonsingular level sets of ψ are strictly pseudoconvex hypersurfaces. Thus in this article, we equivalently call a Stein manifold a *strictly pseudoconvex manifold*. Moreover, it was observed in [ElGr] that the gradient vector field of ψ defines a (global) Liouville vector field $\nu = \nabla_\psi$, making all nonsingular level sets ω_ψ -convex. Hence, Stein manifolds exhibit strongest filling properties for a contact manifold which can be realized as their boundary.

In this article, we are mainly interested in compact Stein surfaces. Another characterization of these manifolds, which might be called “the topologist’s fundamental theorem of compact Stein surfaces”, is due to Eliashberg, and was made explicit by Gompf in dimension four:

Theorem 1.3 (Eliashberg [El1], Gompf [Go2]). *A smooth oriented compact 4-manifold with boundary is a Stein surface, up to orientation preserving diffeomorphisms, if and only if it has a handle decomposition $X_0 \cup h_1 \cup \dots \cup h_m$, where X_0 consists of 0- and 1-handles and each h_i , $1 \leq i \leq m$, is a 2-handle attached to $X_i = X_0 \cup h_1 \cup \dots \cup h_i$ along a Legendrian circle L_i with framing $tb(L_i) - 1$.*

All structures we have introduced so far meet in the following theorem:

Theorem 1.4 (Loi-Piergallini [LP], also see [AO]). *An oriented compact 4-manifold with boundary is a Stein surface, up to orientation preserving diffeomorphisms, if and only if it admits a PALF.*

Throughout the article, we give ourselves the freedom of using the prefix ‘anti’ as a shorthand, whenever an oriented manifold X admits a structure when the orientation on X is reversed; like anti-symplectic, anti-Kähler, or anti-Stein. For Lefschetz fibrations and open books though, we use ‘positive’ and ‘negative’ adjectives to distinguish two possible cases.

2. SIMPLE FOLDED SYMPLECTIC STRUCTURES

The definition of symplectic (or anti-symplectic) structures can be enlarged as follows in order to cover a larger family of manifolds, which was shown in [C] to contain entire family of closed oriented smooth 4-manifolds:

Definition 2.1. *A folded symplectic form on a smooth $2n$ -dimensional manifold X is a closed 2-form ω such that ω^n is transverse to the 0-section of $\Lambda^{2n}T^*X$, and whenever this intersection is nonempty, ω^{n-1} does not vanish on the hypersurface $H = (\omega^n)^{-1}(0)$, called the fold.*

For an oriented X , the kernel of ω on H integrates to a foliation called *null-foliation*. Martinet's singular form $x_1 dx_1 \wedge dy_1 + dx_2 \wedge dy_2 + \dots + dx_n \wedge dy_n$ on \mathbb{R}^{2n} defines the standard folded symplectic structure, as every folded symplectic form can be expressed in this way in an appropriate Darboux coordinate system around any point on the fold. There is also a simple folded structure that every even dimensional sphere carries: We think of S^{2n} sitting in \mathbb{R}^{2n+1} , then pull back the standard symplectic form $dx_1 \wedge dy_1 + \dots + dx_n \wedge dy_n$ on the unit disk bounded by the equator in \mathbb{R}^{2n} to S^{2n} by the projection maps along the last coordinate, and finally glue them along the fold S^{2n-1} to obtain ω_0 . This is equivalent to doubling the unit disk equipped with its standard symplectic form (by reversing the orientation on one of the disks). We call this form the *standard folded symplectic form on S^{2n}* .

For more on folded symplectic structures, the reader is referred to [CGW], [C]. Here we only consider these forms on Riemann surfaces and compact 4-manifolds, possibly with boundaries. For the former class, folded symplectic forms form an open and dense set in the space of 2-forms, whereas in dimension four openness remains but the nonvanishing condition implies that they are nongeneric. We say an embedded surface $\Sigma \subset X^4$ is a *folded symplectic submanifold* of (X, ω) if $\omega|_\Sigma$ is a folded symplectic form on Σ . Observe that S^2 equipped with the standard form obtained by pulling back $dx_1 \wedge dy_1$ embeds as a folded symplectic submanifold of S^4 with the standard folded symplectic form defined as the pullback of $dx_1 \wedge dy_1 + dx_2 \wedge dy_2$ as above.

The following proposition provides several examples of folded symplectic 4-manifolds:

Proposition 2.2. *Let X be a closed oriented smooth 4-manifold and Σ be a closed oriented surface. If $f : X \rightarrow \Sigma^2$ is an achiral Lefschetz fibration such that the regular fiber is a closed oriented surface F which is nonzero in $H_2(X; \mathbb{R})$, then X admits a folded symplectic structure ω such that fibers are symplectic and the fold H is an F -bundle over S^1 . The fold H splits X into pieces X_+ and X_- , and f induces symplectic Lefschetz fibrations on $(X_+, \omega|_{X_+})$ and on $(-X_-, \omega|_{X_-})$, respectively. Furthermore, any finite set of sections can be made folded symplectic for an appropriate choice of ω . This form is canonical up to deformation equivalence of folded symplectic forms.*

We will call this type of folded symplectic structures *simple* (after Thurston [T]). Base spaces of the fibrations defined on X_+ and $-X_-$ are determined by an arbitrary splitting $\Sigma = \Sigma_+ \cup \Sigma_-$. Here we take $\Sigma_- = D^2$ for simplicity. Observe that the fibration induces an exact sequence $\pi_1(F) \rightarrow \pi_1(X) \rightarrow \pi_1(\Sigma) \rightarrow \pi_0(F) \rightarrow 0$. It follows that fibers are connected if the base is simply-connected. Otherwise we can define a new achiral Lefschetz fibration

from X to the finite cover of Σ corresponding to the finite-index subgroup $f_{\#}(\pi_1(X))$ in $\pi_1(\Sigma)$, which has connected fibers. Finally, one can perturb f to get a fibration which has at most one critical point on each fiber. Hence, without loss of generality, we will assume that the fibers of f are connected and critical values are distinct.

Proof of Proposition 2.2. Start by connecting all negative critical points in the base by an embedded arc in the complement of positive critical points, and cover it by the images of orientation reversing charts so that we get a closed neighborhood $\Sigma_- \cong D^2$ of this arc away from the positive critical points. This can be done because around the regular points we have freedom to take charts of either orientation. After we reverse the orientation on $f^{-1}(\Sigma_-)$, the map $f : f^{-1}(\Sigma_-) \rightarrow \Sigma_-$ defines a negative Lefschetz fibration. Set $\Sigma_+ = \Sigma \setminus \Sigma_-$, $C = \Sigma_+ \cap \Sigma_-$, $X_+ = f^{-1}(\Sigma_+)$, $X_- = f^{-1}(\Sigma_-)$, and $H = f^{-1}(C)$. If there are no negative critical points, we can choose Σ_- as a small disk around a regular value which does not contain any critical values. Now let β be a folded symplectic form on Σ which folds over C , such that it is a positive area form on Σ_+ and a negative area form on Σ_- . These forms always exist: For example take S^2 with its standard folded form ω_0 , and suppose Σ_{\pm} has genus g_{\pm} . Symplectic connect sum the upper-hemisphere of S^2 with a closed genus g_+ surface equipped with a positive symplectic form, and the lower-hemisphere with a closed genus g_- surface equipped with a negative symplectic form. This yields a folded symplectic form on Σ , folded along C .

We will construct a folded symplectic form on X by mimicking Gompf's proof which generalizes Thurston's result for symplectic fibrations to symplectic Lefschetz fibrations ([T], [GS]). Let ζ be a closed 2-form on X which evaluates positively on any closed surface contained in a fiber with the induced orientation. (We have not made any assumptions on the type of vanishing cycles, so one might have more than one closed surface on a fiber if there are separating vanishing cycles.) First we wish to define a closed 2-form η on all over X which is symplectic on each $F_y = f^{-1}(y)$, for all $y \in \Sigma$.

Let A be a tubular neighborhood of C in Σ which does not contain any critical values. Choose disjoint open balls $U_{+,k}$ around each positive and $V_{-,l}$ around each negative critical point so that these sets do not intersect $f^{-1}(A)$ in X and that in appropriate charts the fibration map can be written as $f(z_1, z_2) = z_1 z_2$ and $f(z_1, z_2) = \bar{z}_1 z_2$, respectively. Take the standard forms $\omega_{+,k} = dx_1 \wedge dy_1 + dx_2 \wedge dy_2 = -\frac{i}{2} dz_1 \wedge d\bar{z}_1 - \frac{i}{2} dz_2 \wedge d\bar{z}_2$ on $U_{+,k}$ and $\omega_{-,l} = -dx_1 \wedge dy_1 + dx_2 \wedge dy_2 = \frac{i}{2} dz_1 \wedge d\bar{z}_1 - \frac{i}{2} dz_2 \wedge d\bar{z}_2$ on $V_{-,l}$ for all k, l . For any $y \in f(U_{+,k})$, $F_y \cap U_{+,k}$ is a $J_{+,k}$ -holomorphic curve, where $J_{+,k}$ is an almost-complex structure compatible with $\omega_{+,k}$. Similarly for any $y \in f(V_{-,l})$, $F_y \cap V_{-,l}$ is $J_{-,l}$ -holomorphic curve, where $J_{-,l}$ is an almost-complex structure compatible with $\omega_{-,l}$. Having expressed $\omega_{+,k}$ and $\omega_{-,l}$ in terms of Kähler forms, we can take these almost-complex structures as (i, i) and $(-i, i)$, respectively. It follows that $\omega_{+,k}|_{F_y \cap U_{+,k}}$ is symplectic, so we can extend it to a symplectic form ω_y on the entire fiber and get ω_y defined for all points in each $f(U_{+,k})$ this way. Do the same for all points in $f(V_{-,l})$, for every j . Finally, for all remaining $y \in \Sigma$ take any symplectic form ω_y on the fiber, and rescale every ω_y we have defined away from all $U_{+,k}$ and $V_{-,l}$ so that they are in the same cohomology class as the restriction of ζ to each F_y . Next, cover Σ with finitely many balls B_s containing at most one critical value, and whenever they do contain a critical value, assume they are centered at that point. Reindex $U_{+,k}$ and $V_{-,l}$, and shrink them if necessary to make sure they lie in $f^{-1}(B_s)$ for some k . Define η_s on each $f^{-1}(B_s)$ as the pullback of $\omega_{+,s}$, $\omega_{-,s}$, or ω_y by r_s , where r_s is

the retraction of $f^{-1}(B_s)$ to the fiber F_y over the center of B_s , or the union of F_y either with closure of $U_{+,s}$ or with closure of $V_{-,s}$, whenever B_s contains a positive or negative critical value, respectively. Now we can glue these forms to construct the 2-form η we wanted, by using a partition of unity and that each η_s is cohomologous to $\zeta|_{f^{-1}(B_s)}$ as in [GS].

We claim that $\omega_\kappa = \kappa\eta + f^*(\beta)$ is a folded symplectic form on X , where κ is a small enough positive real number. ω_κ is clearly closed and symplectic in the fiber direction. It follows that for any noncritical point $x \in F_y$, $T_x X = T_x F_y \oplus (T_x F_y)^\perp$. Here $f^*(\beta)$ is nondegenerate over $(T_x F_y)^\perp$ for all $x \notin H$, implying that for sufficiently small κ , ω_κ is nondegenerate on $X \setminus (H \cup_s (U_{+,s} \cup V_{-,s}))$. On the other hand $\omega_\kappa|_{U_{+,s}} = \kappa\omega_{+,s} + f^*(\beta)$ and $\omega_\kappa|_{V_{-,s}} = \kappa\omega_{-,s} + f^*(\beta)$. Therefore for any nonzero $v \in TU_{+,s}$, we have $\omega(v, J_{+,s}v) = \kappa g(v, v)_{+,s} + \beta(f_*(v), if_*(v)) > 0$, where $g(-, -)_{+,s}$ is the metric induced from $\omega_{+,s}$ and $J_{+,s}$. Likewise, for any nonzero $v \in TV_{-,s}$, we will have $\omega(v, J_{-,s}v) = \kappa g(v, v)_{-,s} + \beta(f_*(v), -if_*(v)) > 0$, $g(-, -)_{-,s}$ being the metric induced from $\omega_{-,s}$ and $J_{-,s}$. (Recall that β is negative on Σ_- .) Hence ω_κ is symplectic everywhere on X except H , where it vanishes transversely. Moreover, $f^*(\beta)$ is a folded symplectic form on any section, so taking κ even smaller, we can as well assume that any finite collection of sections of f are folded symplectic. It is easy to check that the folded symplectic form we get satisfies all the other declared properties. (Also see Remark 2.3). \square

The homological assumption in the theorem is a very mild one. If S is the set of critical points of the achiral fibration $f : X \rightarrow \Sigma$, then the tangencies of the fibers define a complex line bundle $L = \text{Ker}(df)$ on $X \setminus S$, which extends uniquely over X . It follows that unless we have a torus fibration, the regular fiber F is essential, since $\langle c_1(L), F \rangle = \chi(F)$. Also if the fibration is obtained from a pencil by blowing up the base points, the exceptional spheres will become sections of the fibration, guaranteeing that the fibers are essential in the homology.

Remark 2.3. Alternatively, the folded symplectic form in Theorem 2.2 can be constructed by using the *folding* operation described in [CGW]. Restrictions of β on Σ_+ and on Σ_- give well-defined area forms β_+ and β_- , respectively. Gompf's method can be used to define a symplectic form $\kappa_+\eta + f^*(\beta_+)$ on X_+ , where η is a 2-form on X that restricts to the fibers as a (positive) symplectic form and κ_+ is a small enough positive real number. The orientation on the base together with the orientation on the regular fiber determines the orientation of the total space, and thus by taking the opposite orientation on Σ_- but keeping the orientation on F , one orients $-X_-$. Let $\bar{f} : -X_- \rightarrow \Sigma_-$ be the fibration defined by taking orientation-preserving charts for $f : X_- \rightarrow \Sigma_-$, then we can define a symplectic form $\kappa_-\eta + \bar{f}^*(-\beta_-)$ on $-X_-$ (as $-\beta_-$ is the area form on Σ_-) by following the same construction method. Observe that $\bar{f}^*(-\beta_-) = f^*(\beta_-)$. Hence, setting $\kappa = \min\{\kappa_+, \kappa_-\}$, we obtain two symplectic manifolds (X_+, ω_+) and $(-X_-, \omega_-)$, where $\omega_\pm = \kappa\eta + f^*(\beta_\pm)$. Let ι_\pm be the inclusions of boundaries into $\pm X_\pm$, then $\iota_+^*(\omega_+) = \kappa\eta = \iota_-^*(\omega_-)$ and the orientations of both null-foliations agree. Thus we can glue these pieces to obtain a folded symplectic structure on $X_+ \cup X_- = X$, which agrees with ω_+ and ω_- in the complement of a tubular neighborhood of the fold $\partial X_+ = H = -\partial X_-$. (See [CGW] for details.) This form is deformation equivalent to the form $\kappa\eta + f^*(\beta)$ in Proposition 2.2.

3. EVERY CLOSED ORIENTED 4-MANIFOLD IS FOLDED SYMPLECTIC

Here we show that any closed oriented smooth 4-manifold X can be equipped with a folded symplectic form. For the sake of completeness, we start by outlining Etnyre and Fuller's proof that every 4-manifold admits an achiral Lefschetz fibration after a surgery along a framed circle [EF] : Take a handlebody decomposition of X with one 0– and one 4-handle, let X_1 denote the union of the 0–handle, 1–handles and 2–handles, and X_2 denote the union of the 3–handles and the 4–handle. By Theorem 1.1 there exist achiral Lefschetz fibrations $f_i : X_i \rightarrow D^2$, which necessarily have bounded fibers, and stabilizing both fibrations we may as well assume the fibers have connected boundaries. After a possible slight modification of the handlebody decomposition, Etnyre and Fuller manipulate the contact structures on the boundaries so that they are both overtwisted and homotopic as plane fields. Then it follows from results of Eliashberg and Giroux that we have isotopic contact structures, and thus the induced open books are the same, possibly after some stabilizations and isotopies. Denoting the final manifolds and fibrations with X_i and f_i again, we may therefore assume that the open book decompositions induced by these fibrations on the common boundary $H = \partial X_1 = -\partial X_2$ are the same, so we can glue both pieces of X back along the truncated pages, and obtain an achiral Lefschetz fibration

$$f_1 \cup f_2 : W = X_1 \bigcup_{f_1^{-1}(\partial D^2)=f_2^{-1}(\partial D^2)} X_2 \longrightarrow S^2.$$

To recover X we need to glue $S^1 \times D_1^2$ to $S^1 \times D_2^2$, where $S^1 \times D_i^2 = \partial X_i \setminus f_i^{-1}(\partial D^2)$. Filling the boundary of W with an $S^1 \times D^3$ gives the same result, so we can view W as $X \setminus N$ where N is a neighborhood of an embedded curve $\gamma \subset X$. Now, if we instead add on a $D^2 \times S^2$ so that each $\partial D^2 \times \{pt\}$ is identified with $S^1 \times \{pt\}$, we can extend the fibration on W by the projection on the S^2 component of $D^2 \times S^2$. Hence, we obtain an achiral Lefschetz fibration over S^2 on the resulting manifold Y , where the section S of this fibration discussed in [EF] can be taken as $0 \times S^2$ coming from the glued in $D^2 \times S^2$, implying S has trivial normal bundle in Y .

We will refer the following as the *standard model*: Consider S^4 with the standard folded symplectic structure ω_0 described before, and take $S^4 \cap \{x_4 = 0\} \cong S^3$ vertical to the fold $H_0 = S^4 \cap \{x_5 = 0\} \cong S^3$. Take $S_0 = S^4 \cap \{x_4 = 0 = x_3\} \cong S^2$ which intersects the fold along the circle $C_0 = \{x_1^2 + x_2^2 = 1 \mid x_3 = x_4 = x_5 = 0\}$. It is easy to see that ω_0 restricts to this S_0 as the standard folded symplectic form on S^2 , folded along C_0 , and symplectic on the normal disks to S_0 . Fix a disk neighborhood M_0 of S_0 so that ω_0 evaluates as 1 on each normal disk. That is, each normal disk projects onto unit disk $\{x_3^2 + x_4^2 \leq 1 \mid x_1 = x_2 = x_5 = 0\}$ symplectomorphically. By restricting ω_0 , we get two folded symplectic manifolds $M_0 \cong S^2 \times D^2$ and $N_0 = S^4 \setminus M_0 \cong D^3 \times S^1$, with folds $S^1 \times D^2$ and $D^2 \times S^1$, respectively.

The existence of the section $s : S^2 \rightarrow S \subset X$ guarantees that the fiber of the achiral Lefschetz fibration $f : Y \rightarrow S^2$ is homologically essential and therefore there exists a folded symplectic form ω as described in Theorem 2.2. This restricts to $Y \setminus M$, where $M \cong S^2 \times D^2$ is a neighborhood of S . We may assume ω is constructed such that M is identified with M_0 in the standard model above as follows: Let $\phi : M \rightarrow M_0$ be an orientation preserving diffeomorphism such that ϕ is orientation preserving on the spheres

(and on the normal directions as well), and that it maps the upper-hemisphere of S_0 (where ω_0 is positive) to the positive part of S . Then one can start the construction in the proof of Theorem 2.2 with the folded symplectic form $s^*\phi^*(\omega_0)$ on the base sphere, which naturally restricts to an area form on each hemisphere. We can also modify the symplectic form $\kappa\eta$ on the fibers so that it is symplectomorphic to $\phi^*(\omega_0)$ on the normal disks to S , each of which lies on a fiber.

Hence we obtain a folded symplectic form ω on X such that $(M, \omega|_M)$ is folded symplectomorphic to $(M_0, \omega_0|_{M_0})$. This allows us to trade M for $N \cong S^1 \times D^3$ and extend the folded symplectic structure to $(Y \setminus M) \cup N \cong X$. The effect of this surgery on the fold of Y is to turn the surface fibration over S^1 into an open book decomposition on the resulting fold. The core curve of N sits in the 3-manifold as the binding of this open book and therefore it carries a canonical framing. We have proved:

Theorem 3.1. *Every closed oriented smooth 4-manifold X admits a folded symplectic structure. Furthermore, there exist folded symplectic forms on X with connected folds, such that a surgery along a framed curve which lies in the fold results in a simple folded symplectic manifold.*

Remark 3.2. Away from the framed curve γ in X , the folded symplectic model we have constructed is the restriction of the simple model discussed in the previous section, and as we will see shortly, the pieces are Stein and anti-Stein. So for any sort of pseudo-holomorphic curve counting with respect to this folded symplectic structure, the focus would be understanding the limit behaviors around γ of the curves in the moduli space, where we do have a standard model, namely $(N_0, \omega_0|_{N_0})$ above. (For a digression on this topic, see [vB].) We would like to point out that both the knot type of γ in the fold and its framing depend on the achiral Lefschetz fibrations we use in the construction, so does the simple model we get. The following example illustrates this phenomenon.

Example 3.3. If we construct S^4 following the recipe given in the proof of Theorem 3.1, we get $W = D^2 \times D^2 \bigcup_{C \times D^2} D^2 \times D^2 = S^2 \times D^2$, which can be identified with M_0 , and the simple folded symplectic form on $Y = S^2 \times S^2 = M_0 \bigcup_{S^2 \times \partial D^2} M_0$ can be constructed so that its restriction to each copy of M_0 is indeed the standard form ω_0 . Note that here both open books already agree, so we do not need to alter the contact structures and change the initial fibrations. Now if we undo the surgery, that is if we trade $N = N_0$ and M in the proof, what we get is the standard folded symplectic form ω_0 on S^4 .




FIGURE 1. On the left: 0-surgery along the binding yields a trivial S^2 fibration over D^2 on each piece, which make up $S^2 \times S^2$. On the right: 0-surgery along the new binding yields a cusp neighborhood on both sides.

It is a standard fact that surgery along a framed curve in a simply-connected 4-manifold will result in connect summing with either $S^2 \times S^2$ or $S^2 \tilde{\times} S^2$, depending on the framing, which can be thought as an element of $\pi_1(\mathrm{SO}(3)) = \mathbb{Z}_2$. In [EF] (also see [Ha]) it is described how one can homotope the framed knot in the 3-manifold to another framed knot, which is isotopic in the ambient 4-manifold to the original one, so that their framings differ by one and that surgering the new curve yields an achiral Lefschetz fibration on the resulting manifold as well. Applying this trick to our example, we can instead pass to an achiral Lefschetz fibration on $S^2 \tilde{\times} S^2 \cong \mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$, which is a torus fibration with two cusp fibers of opposite signs (Figures 1 and 2). The monodromy of this achiral fibration is $t_a t_b t_b^{-1} t_a^{-1}$, and the corresponding Kirby diagram is depicted in Figure 2 (cf. [Go1]). To verify that this manifold is $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$, we first slide one of the vertical 2-handles over the other one, and then separate this pair from the rest of the diagram by sliding over the 0 framed 2-handle. Now the rest of the diagram can be shown to be S^4 after obvious handle cancelations. It is possible to see that our construction method will give a folded symplectic structure on S^4 equivalent to the standard one again. Note that the first simple model above is obtained by surgering the unknot, whereas the second comes from surgering the right trefoil in S^3 . Surgery framings on them differ by one in S^4 .




FIGURE 2. The achiral Lefschetz fibration on the second associated model. The total space is shown to be $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$.

4. DECOMPOSITION THEOREM

While we shift our attention to Stein structures, we would like to have only non-separating vanishing cycles in our constructions, as it is suggested by the correspondence between PALFs and compact Stein surfaces established in [LP] and [AO]. We start with the following lemma:

Lemma 4.1. *Let X be a closed oriented 4-manifold. Then it can be decomposed into two handlebodies, each of which admits an allowable achiral Lefschetz fibration over D^2 , such that the fibers have connected boundaries and that the induced open books are the same.*

Proof. We follow the construction of Etnyre and Fuller with more care given to having fibrations allowable. Take a handlebody decomposition of X with one 0– and one 4–handle, let X_1 be the union of the 0–handle, 1–handles and 2–handles, and X_2 be the union of 3–handles and the 4–handle. As it was implicitly present in [Ha], and was also observed in [AO], one can always build an achiral Lefschetz fibration on a given 2–handlebody so that all vanishing cycles are non-separating. Therefore, we can start with allowable fibrations and then proceed with stabilizations as described in [EF] to match the induced open books. A stabilization is given by gluing a positive or a negative Lefschetz 2–handle along a new 1–handle added to a regular fiber, and in order to keep the binding connected, we always introduce another adjacent stabilization. Therefore, all vanishing cycles introduced during stabilizations are also nonseparating. Induction concludes the proof. \square

Theorem 4.2. *Let X be a closed oriented smooth 4-manifold. Then X admits a decomposition into two codimension zero submanifolds X_+ and X_- , such that X_+ and $-X_-$ are both compact Stein manifolds with strictly pseudoconvex boundaries. These Stein structures can be chosen so that the induced contact structures ξ_+ on ∂X_+ and ξ_- on $-\partial X_-$ are isotopic. Furthermore, there are PALFs on each piece such that the open book decompositions they induce on ∂X_+ and $-\partial X_-$ are compatible with ξ_+ and ξ_- , respectively, and they coincide. In short, all data match on the hypersurface $H = \partial X_+ = -\partial X_-$.*

Proof. The lemma above gives us a decomposition of X into two pieces X_1 and X_2 with allowable achiral Lefschetz fibrations f_1 and f_2 on them, such that induced open books on the boundaries match. As in the proof of Theorem 3.1, we glue these two pieces along the truncated pages to get: $W = X_1 \bigcup_{f_1^{-1}(\partial D^2)=f_2^{-1}(\partial D^2)} X_2$. Next we glue in $S^2 \times D^2$ to cap off the fibers and establish an achiral Lefschetz fibration $\hat{f} : Y \rightarrow S^2$ with closed fibers.

We wish to split the base of this fibration into two disks D_+ and D_- so that all the positive critical values lie in the interior of D_+ and all the negative ones lie in the interior of D_- . As discussed earlier, restrictions of f give a positive Lefschetz fibration on $X_+ = f^{-1}(D_+)$ and a negative Lefschetz fibration on $X_- = f^{-1}(D_-)$, respectively. It also prescribes a surface bundle over $S^1 = \partial D_+ = -\partial D_-$ on the hypersurface separating Y_+ and Y_- . This time we would like to describe the splitting more carefully by taking into consideration how the global monodromies of the new fibrations are related to the original ones.

Let μ_1 be the monodromy of the achiral Lefschetz fibration f_1 on X_1 and μ_2 be the monodromy of the fibration f_2 on X_2 . Fix a representation of μ_1 by using arcs $s_1^1, \dots, s_{k_1}^1$ and a representation of μ_2 by using arcs $s_1^2, \dots, s_{k_2}^2$. Corresponding critical values are




FIGURE 3. New monodromies from old ones. On the left: μ_1 is given by solid arcs, and μ_2 by dotted ones. On the right: Solid arcs are the *positive arcs* representing μ_+ , whereas conjugated dotted arcs are *negative*, providing a representation of μ_- after closing the base back to S^2 .

denoted by y_i^1 and y_j^2 . Monodromies of the open book decompositions bounding each fibration are given by μ_1 and μ_2 as well, and they coincide under an orientation reversing diffeomorphism, so $\mu_1 = (\mu_2)^{-1}$. Let V be a small neighborhood of a regular value in the base S^2 . We obtain an achiral Lefschetz fibration $f : W \setminus f^{-1}(V) \rightarrow D = S^2 \setminus V$, which closes up to a fibration over S^2 . If g is the genus of the page, then this fibration is determined by the relation $\mu_1\mu_2 = 1$ in $\Gamma_{g,1}$ and is mapped under the maps $\Gamma_{g,1} \rightarrow \Gamma_g^1 \rightarrow \Gamma_g$ to the relation that describes the achiral Lefschetz fibration $\hat{f} : Y \setminus \hat{f}^{-1}(V) \rightarrow D$. Since this map factors through Γ_g^1 , the achiral Lefschetz fibration \hat{f} comes naturally with a section S of self-intersection zero. We denote images of the elements in $\Gamma_{g,1}$ under this map with the same elements, so $\mu_1\mu_2 = 1$ is the global monodromy of the achiral Lefschetz fibration on $Y \setminus \hat{f}^{-1}(V)$. Note that we can use the same arcs s_i^1 and s_j^2 to represent the global monodromy of this fibration. Now, if we move counterclockwise and choose only the arcs that run through positive Dehn twists, we establish a monodromy μ_+ . Call these arcs *positive*. Next, we choose a nearby base point, and move counterclockwise by running through the negative Dehn twists only, while avoiding intersecting any positive arc. This way, we obtain a monodromy μ_- . The new set of arcs involved in this monodromy will be referred as *negative arcs*. Observe that each negative arc is obtained by traveling around some old arcs s_i^1 and s_j^2 in order to avoid intersecting positive arcs, then going around the aimed negative critical point once, and finally going all the way back on the same detour (Figure 3). That is, each negative arc corresponds to a conjugate of a negative Dehn twist in Γ_g , which defines a negative Dehn twist, too. By taking regular neighborhoods of these arcs such that positive and negative arcs stay apart, we get a disk enclosing only positive critical points, and an annulus containing only negative critical points. Closing the fibration to a fibration over S^2 , the latter becomes a disk as well. Now we can enlarge anyone of these disks so both disks share the same boundary, and call the one containing positive values D_+ , and the other one D_- . So we have a new factorization of the global

monodromy of \hat{f} , given by the relation $\mu_+\mu_- = 1$. The section S prescribes how to lift the new elements μ_\pm of Γ_g to Γ_g^1 uniquely.

We proceed with taking out the tubular neighborhood $S^2 \times D^2$ of the section from $Y = Y_+ \cup Y_-$, and we get an inherited splitting $X_+ \cup X_-$. The discussion above shows that μ_+ defines a positive Lefschetz fibration on X_+ and μ_- defines a negative fibration on X_- . To recover the original 4-manifold X we need to put back in $S^1 \times D^3$, which has the same effect as gluing each other the tubular neighborhoods $S^1 \times D_+^2$ and $S^1 \times D_-^2$ of the bindings of open books on ∂X_+ and ∂X_- , respectively. Therefore we can think of X as decomposing into X_+ and X_- . We claim that this decomposition possesses the desired properties.

When we take out a tubular neighborhood of S from Y , we turn the positive and negative Lefschetz fibrations on Y_+ and Y_- into a PALF on X_+ and a NALF on X_- , respectively. In the meantime the surgery converts the surface fibration that separates Y_+ and Y_- to an open book decomposition on the common boundary $H = \partial X_+ = -\partial X_-$. The binding of this open book is the identified bindings of ∂X_+ and $-\partial X_-$, the page F is the bounded closed surface obtained by cutting off a disk from the regular fiber of \hat{f} , and the monodromy is induced from the fibration on either side. Noting that the NALF on X_- becomes a PALF on $-X_-$, we see that both PALFs induce the same open book decomposition on their boundaries.

By Theorem 1.4, both X_+ and $-X_-$ admit Stein structures. We will construct these Stein structures using Eliashberg's characterization so that they match on the common boundary. The technique we are going to use is the same as the one which was presented in [AO]: The PALF on X_+ is obtained by attaching positive Lefschetz handles $h_1 \cdots h_m$ to $X_0 = F \times D^2$, which has the obvious PALF defined by projection onto D^2 component. The same is true for the PALF on $-X_-$. $F \times D^2$ has a natural Stein structure by Theorem 1.3. We can assume all vanishing cycles (coming from either side) sit in various pages of the open book on H . Read backwards, we can think of the fibrations as being constructed by attaching positive and negative Lefschetz handles to H on either side in a sequence following the monodromy of the open book. Thus we can induct on the number of handles. Assume that the PALF on $X_{i-1} = X_0 \cup h_1 \cup \dots \cup h_{i-1}$ ($i \leq m$) induces an open book decomposition on its boundary, and it carries a Stein structure such that the contact structure induced on the boundary is compatible with this open book. Let C be the vanishing cycle of the positive Lefschetz handle h_i lying on a page F of ∂X_{i-1} . We open up the open book decomposition and choose a page against F , and glue them together along the binding B to form a smooth closed convex surface Σ in the 3-manifold ∂X_{i-1} . As C is non-separating, $\Sigma \setminus C$ is connected and it contains the dividing set, namely B . So we can use the *Legendrian realization principle* ([Gi1], [Ho]) to isotope Σ through convex surfaces to make C Legendrian. Note that this is done by a small C^∞ isotopy of the contact structure supported in a neighborhood of Σ , which fixes the binding pointwise. Hence the framing of C relative to the fiber F is the same as its contact framing, implying that the Lefschetz handle h_i is attached along a Legendrian curve with framing $tb - 1$. By Theorem 1.3 the Stein structure extends over this handle, and as shown by Gay in [Ga], the new open book on ∂X_i will be compatible with the new induced contact structure on ∂X_i . This completes the induction. Repeating the same argument dually for $-X_-$, we see that the compatible open books on ∂X_+ and $\partial(-X_-)$ are isotopic, and therefore the

induced contact structures ξ_+ on ∂X_+ and ξ_- on $\partial(-X_-) = -\partial X_-$ are isotopic as well. So we fulfill all the matching conditions listed in the statement of the theorem. \square

Remark 4.3. In [AM] it was asked if one could decompose a given closed oriented smooth 4-manifold into Stein pieces so that the induced contact structures on the separating 3-manifold coincide. Our theorem gives an affirmative answer to this question. In the same article authors remark that it is possible to alter their Stein decomposition to make the induced contact plane distributions homotopic, but the tightness of the contact structure precludes the use of Eliashberg's celebrated theorem on overtwisted contact structures to conclude more. Considering the underlying PALFs and isotopies of open books gives a way around this difficulty, thanks to Giroux's Theorem.

Remark 4.4. In [Q], Quinn studied so-called *dual decompositions* of 4-manifolds: descriptions of 4-manifolds as a union of two 2-handlebodies. The author formulates the same question as in [AM] in terms of necessary sequence of Kirby moves to relate a possibly nonmatching Stein decomposition. Theorem 4.2 provides an *implicit* answer to this question, and we would like to take this as an opportunity to summarize the handle calculus behind our construction: An arbitrary Stein decomposition $X = X_1 \cup X_2$ comes with some PALF pair. Using the stabilization moves of Etnyre and Fuller, we first change this PALF pair with a matching pair. This corresponds to adding cancelling 1- and 2- handles to each X_i , or in other words, we add cancelling handle pairs of index 1-, 2- and 3- in the original handlebody decomposition of X . In the next step, we pass to a cobordant 4-manifold Y so that we can split the positive and negative Lefschetz handles. Then we 'undo' the surgery and get the decomposition $X = X_+ \cup X_-$ with Stein structures on each piece that coincide on the common boundary. Having the simply-connected case in mind, this intermediate step can be seen as a stabilization. Let $W \cong X \setminus S^1 \times D^3$ be the complement of a regular neighborhood of the framed knot γ in X , then the first surgery defines a cobordism

$$[0, 2] \times W \bigcup_{[0,2] \times S^1 \times S^2} ([0, 1] \times S^1 \times D^3) \cup_{1 \times S^1 \times S^2} ([1, 2] \times D^2 \times S^2),$$

which is identity on the first component. We trade 2-handles of X_1 and X_2 in Y by making use of the two extra handles of index 2. Finally, the composition of two cobordisms that gives back X can be seen as the double of the cobordism above, and thus it deformation retracts to

$$W \bigcup_{S^1 \times S^2} (S^4 \cup_{D^2 \times S^2} S^4).$$

This cobordism is built by attaching cells to $\partial W = S^1 \times S^2$, where $D^2 \times S^2$ is attached uniquely and the framing of γ indicates in which one of the two ways we shall glue the other two $S^1 \times D^3$ pieces. Although here we started with a (nonmatching) Stein decomposition, it is clear that the same discussion can be carried out in our main construction as well. Therefore we have a well-defined process, during which we first inflate the number of handles in a given decomposition of X , and then trade some of the 2-handles through a cobordism to achieve the desired decomposition at the end.

5. FOLDED KÄHLER STRUCTURES

Unlike symplectic structures, random folded symplectic structures do not need to bear any information about the geometry or topology of the manifold they are defined on. In order to specify more meaningful members of this family, one first of all needs to impose some boundary conditions on the folding hypersurface. We would like to acknowledge a result of Kronheimer and Mrowka: In [KM], the authors prove that a compact symplectic 4-manifold (Y, ω) with strictly pseudoconvex boundary has $SW_Y(K) = 1$, where K is the canonical class of ω . This motivates us to see such manifolds as building blocks of 4-manifolds, and yields a good boundary constraint for folded symplectic structures, at least in this dimension. Henceforth, we assume that the fold $H = (\omega^n)^{-1}(0)$ of a given folded symplectic manifold (X^{2n}, ω) is always connected and nonempty. We will generalize the notion of a Kähler structure on a smooth $2n$ -manifold by considering a distinguished subset of the family of folded symplectic structures, and we then present some properties of these structures:

Definition 5.1. *A folded symplectic form ω on an oriented $2n$ -dimensional manifold X is called a folded Kähler structure, if there is a tubular neighborhood N of H such that:*

- (1) *The closure of each component of $X \setminus N$ is a compact Kähler manifold $(\pm X_{\pm}, \omega|_{\pm X_{\pm}})$ with strictly pseudoconvex boundary,*
- (2) *$(N, \omega|_N)$ is folded symplectomorphic to $([-1, +1] \times H, d((t^2 + 1)\pi^*(\alpha)))$, where α is a contact 1-form on H , and π is the projection $\pi : [-1, +1] \times H \rightarrow H$.*

In addition, if each $(\pm X_{\pm}, \omega|_{\pm X_{\pm}})$ is strictly pseudoconvex, we say ω is a nicely folded Kähler structure on X .

In the definition above, *nice folding* can be reformulated as folding Stein manifolds along matching strictly pseudoconvex boundaries. Recall that if ψ is a proper strictly plurisubharmonic function on a complex manifold S , then the associated 2-form $\omega_{\psi} = -dJ^*d\psi$ is Kähler, and importantly, the symplectic class of (X, ω_{ψ}) is independent of the choice of ψ [ElGr]. Therefore, to complete our alternative formulation, we ask that each piece $\pm X_{\pm}$ should admit some proper strictly plurisubharmonic function ψ_{\pm} , so that $\omega|_{\pm X_{\pm}} = \omega_{\psi_{\pm}}$. In short, it is built in the definition that a nicely folded Kähler manifold is folded Kähler. Finally note that, due to a theorem of Bogomolov [B], any compact folded Kähler manifold X can be made nicely folded after deforming the complex structure and blowing down any exceptional curves. Even though these definitions narrow the family of folded symplectic structures quite a lot, it is important to note, at least in dimension four, that we still have an adequately large family in the light of the following result:

Theorem 5.2. *Any closed oriented 4-manifold X admits a nicely folded Kähler structure.*

Proof. By Theorem 4.2, X can be decomposed into two compact Stein manifolds X_+ and $-X_-$ with strictly pseudoconvex boundaries such that both induce the same contact structure on the common boundary $H = \partial X_+ = -\partial X_-$. We begin with adding collars $\pm U_{\pm}$ to $(\pm X_{\pm}, \omega_{\pm})$, and extending the symplectic structures to ω'_{\pm} on $\pm X'_{\pm} = \pm(X_{\pm} \cup U_{\pm})$ so that new boundaries $\partial(\pm X'_{\pm})$ are still convex and contactomorphic. Let ξ_{\pm} be the induced contact structures on $\partial(\pm X'_{\pm})$ and ψ be a contactomorphism between them. Using the symplectic cut-and-paste argument of Etnyre [E], we can add a symplectic collar to $(\partial X'_+, \omega'_+)$ so that the new boundary is not only contactomorphic to $-\partial X'_-$ but also the

induced *contact forms* agree up to a multiple $k \in \mathbb{R}^+$. For the sake of brevity, let us assume that U_+ above contains this collar part as well. So after rescaling ω'_- (and ω_-) by k if necessary, we see that restrictions of symplectic forms $\omega'_+|_{\partial X'_+}$ and $k\omega'_-|_{-\partial X'_-}$ agree via ψ , and orientations of null-foliations (which correspond to Reeb directions) are the same. Therefore, once again we can apply the folding technique of [CGW] to obtain a folded symplectic structure ω on $X'_+ \cup X'_-$ such that ω agrees with ω'_\pm on the complement of a small tubular neighborhood of the fold H . We enlarge this neighborhood to include U_+ and U_- and call it N . It follows that $X = X_+ \cup N \cup X_- \cong X_+ \cup X_-$, and $\omega|_{X_+} = \omega_+$, whereas $\omega|_{X_-} = k\omega_-$. Also note that, the folding operation provides us with the desired local model on N , that is, $(N, \omega|_N)$ is folded symplectomorphic to $([-1, +1] \times H, d((t^2 + 1)\pi^*(\alpha)))$ by construction [CGW].

Lastly, suppose $\psi_\pm : \pm X_\pm \rightarrow [0, \infty)$ are proper strictly plurisubharmonic functions such that $\pm \partial X_\pm$ correspond to the maximum points of ψ_\pm , and $\omega_\pm = -dJ^* d\psi_\pm$, respectively. If $k \neq 1$, we can replace ψ_- with $k\psi_-$ and obtain $k\omega_-$ above as a Kähler form of a strictly pseudoconvex manifold. Equipped with these properties, ω is a nicely folded Kähler form on X . \square

Remark 5.3. It is clear that Theorem 5.2 is a refinement of Theorem 3.1. Since the folded forms we have constructed in both proofs are obtained through similar steps, one expects that these structures are actually equivalent. Next, we verify this fact, and this way we get an insight of how folded Kähler forms are ‘supported’ (precise definition is given below) by Lefschetz fibrations as was illustrated in Proposition 2.2:

Take the PALF on X_+ in the proof of Theorem 3.1, and attach a symplectic 2-handle along the binding of the induced open book on ∂X_+ as described by Eliashberg in [El2]. This yields a symplectic Lefschetz fibration over D_+^2 . Dually the same argument for the NALF on X_- gives an anti-symplectic Lefschetz fibration over D_-^2 , and these handle attachments can be done so that two fibrations agree on the common boundary. Moreover, we can assume that these fibrations have genus at least two, so the fibrations can be matched as symplectic surface fibrations over a circle, as it was pointed out in [El2]. At the end we get a *simple folded symplectic* manifold Y obtained from X after a surgery along a framed curve γ . However, any two simple folded symplectic forms compatible with a fixed achiral Lefschetz fibration are deformation equivalent by Proposition 2.2. Moreover, we can normalize both forms on the disks which are parallel copies of cocores of new 2-handles that were used to cap off the fibers. Hence, these two folded forms are deformation equivalent on $Y \setminus S^2 \times D^2$. As the folded symplectic structure on $D^3 \times S^1$ which is glued back in to recover X is standard, the folded symplectic form constructed in the proof of Theorem 3.1 and the folded Kähler form obtained in Theorem 5.2 are indeed *equivalent as folded symplectic structures*. \square

Motivated by symplectic and near-symplectic cases ([D], [ADK]), we can conclude our discussion above by defining the Lefschetz fibration analogue for our structures:

Definition 5.4. Let X be a closed oriented 4-manifold, and decompose S^2 as the union of the upper-hemisphere D_+ and the lower-hemisphere D_- which are glued along the equator $C = \partial D_+ = -\partial D_-$. Then a smooth map $f : X \rightarrow S^2$ is said to be a folded Lefschetz fibration on X , if it restricts to a PALF over D_+ , to a NALF over D_- , and to an open book over C bounding both fibrations.

Definition 5.5. *Let X be a closed oriented 4-manifold equipped with a nicely folded Kähler form ω . Then a folded Lefschetz fibration $f : X \rightarrow S^2$ is said to be compatible with ω if each Stein piece X_{\pm} corresponds to $f^{-1}(D_{\pm})$, and if the contact structure they induce on $H = f^{-1}(C)$ is compatible with the open book decomposition coming from the fibration f . In this case, we also say that nicely folded Kähler manifold (X, ω) is supported by the folded Lefschetz fibration f .*

The compatibility in the above definition is completely *on the symplectic level*. This becomes more visible if once again we recall that surgering the binding γ of the open book $f|_{H \setminus \gamma} : H \setminus \gamma \rightarrow S^1$, we pass to a *simple model* where the folded Lefschetz fibration can be extended to a folded *symplectic* achiral Lefschetz fibration \hat{f} with closed fibers. Also note that, since Stein manifolds harbor less topological obstructions in complex dimensions > 2 , it is very likely that they admit higher dimensional analogues of PALFs with similar topological correspondences. If that is established, last two definitions, as well as several results in this paper, can be generalized to all $2n$ -dimensions.

The complete statement of Theorem 4.2 combined with Theorem 5.2 shows that, given a closed oriented 4-manifold X , one can always find a nicely folded Kähler structure ω on X together with a compatible folded Lefschetz fibration. Next, we prove that this property in fact holds for any nicely folded Kähler structure ω :

Proposition 5.6. *Any nicely folded Kähler structure ω on X , up to orientation preserving diffeomorphism, admits a compatible folded Lefschetz fibration.*

Proof. Each Stein piece X_+ and $-X_-$ admits a PALF by Theorem 1.4. If we construct these fibrations following the algorithm of [AO] and keep track of the associated open books, the work of Plamenevskaya [P] shows that we can establish PALFs $f_{\pm} : \pm X_{\pm} \rightarrow D^2$ with the property that the open book decomposition on the boundaries are compatible with the contact structures induced from the Stein structures on $\pm X_{\pm}$, respectively. As the contact structures are assumed to be the same, Theorem 1.2 tells us that we can match these open books after positive stabilizations. Consequently, we get a compatible folded Lefschetz fibration. \square

Remark 5.7. A folded Lefschetz fibration that supports a given folded Kähler structure fails to be unique. In fact, one can find infinitely many pairwise inequivalent such fibrations. This can be seen for example from the construction of [AO], by using different (p, q) torus knots in their algorithm which we adopt for building our achiral Lefschetz fibrations.

Example 5.8. The easiest examples are doubles. If Y^4 is a compact Kähler manifold with strictly pseudoconvex boundary, then $X = Y \cup -Y$ is equipped with a folded Kähler structure. When Y is indeed Stein, we get a nicely folded structure. The first folded structure constructed in Example 3.3 is the double of standard $D^4 \subset \mathbb{C}^2$, whereas the latter is a ‘monodromy double’ of a cusp neighborhood minus a section. Here by ‘monodromy double’ we mean that the pieces are first glued along the pages of the open books, and if the monodromy of the folded Lefschetz fibration on one piece is μ , then the monodromy on the other one is μ^{-1} .

Example 5.9. There is a construction which also allows us to see the nicely folded Kähler structure together with a compatible folded Lefschetz fibration. Take a contact 3-manifold (H, ξ) , and fix a positive open book decomposition (B, f) compatible with ξ . Different

PALFs bounding this open book describe (possibly) different Stein fillings of (H, ξ) . Indeed there are examples of infinitely many pairwise non-diffeomorphic contact 3-manifolds each of which admit infinitely many pairwise non-diffeomorphic Stein fillings constructed this way [OS1]. Thus for every pair of PALFs X_1 and X_2 that fill the same open book, we can construct a nicely folded Kähler form on $X = X_1 \cup -X_2$, as designated in the proof of Theorem 5.2.

Example 5.10. The main steps of our construction are depicted in the following simple, albeit instructive example: Start with classical handlebody decomposition of $X = \#_8 S^2 \times S^2$ with one 0-handle, sixteen 2-handles, and a 4-handle. Let X_1 be the union of 0-, 2-handles, and X_2 be the 4-handle. Each piece admits a D^2 fibration over D^2 . However we wish to construct allowable fibrations, so we introduce two 1- and 2- cancelling handle pairs and two 2- and 3- cancelling handle pairs in the original handlebody decomposition of X . We start building the fibrations from the scratch: Add the 1-handles to the 0-handle and 3-handles to the 4-handle. Attach the two canceling 2-handles with framing -1 to the union of the 0- handle and 1- handles. Attach the other two the same way to the handlebody X_2 , which is the union of 3- handles and the 4- handle. To simplify our description, we will label the 1-handles of the first handlebody as a and b , which generate π_1 of the torus fiber with one boundary component, and we do the same for the 1-handles of X_2 under the obvious identification. So we obtain two achiral Lefschetz fibrations over disks with bounded torus fibers; one with monodromy $t_a^{-1} t_b^{-1}$, and one with $t_b t_a$. One can verify by Kirby calculus that each time we insert a pair of Lefschetz handles prescribed by $t_a t_a^{-1}$ or $t_b t_b^{-1}$, we introduce an $S^2 \times S^2$. (See Figure 2, and observe that here we slide off the 2-handle pair over a $+1$ framed 2-handle instead.) Doing this consecutively, we attach all the remaining 2-handles to the first handlebody, and obtain an achiral Lefschetz fibration on X_1 with monodromy

$$\mu_1 = t_a^{-1} t_b^{-1} t_b t_a t_b t_a t_b t_a t_a^{-1} t_b^{-1} t_a^{-1} t_b^{-1} t_a^{-1} t_b^{-1} t_a^{-1} t_b^{-1},$$

whereas X_2 still has the monodromy

$$\mu_2 = t_b t_a = (t_a^{-1} t_b^{-1})^{-1}.$$

Both open books that bound these fibrations contain negative Dehn twists (recall that on $-\partial X_2$, the monodromy is μ_2^{-1}), and therefore the contact structures they support are overtwisted. As we have already manipulated the monodromy that way, contact structures and open books are isotopic, so we can glue X_1 and X_2 along the truncated pages. Putting in $S^2 \times D^2$ we pass to a torus fibration $\hat{f} : Y \rightarrow S^2$ with global monodromy $\mu_1 \cdot \mu_2$. (Applying the handle slides given in Example 3.3 repeatedly, and proceeding with the same handle cancelations, one can indeed check that $Y = \#_8 S^2 \times S^2 \# \mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$.) Now the monodromy splits easily as explained in the proof of Theorem 4.2, and we get $\mu_+ = (t_b t_a)^5$ and $\mu_- = (t_a^{-1} t_b^{-1})^5$. It is not hard to see that when we take out the section now, we get pieces X_+ and X_- , which are diffeomorphic to $-E_8$ and E_8 , respectively. So X decomposes into a Stein piece $-E_8$ and an anti-Stein piece E_8 . This defines a nicely folded Kähler structure ω on X , folded along the Poincaré homology sphere $\Sigma(2, 3, 5)$, and it is supported by a folded Lefschetz fibration which is the monodromy double of a torus fibration over D^2 minus a section.

REFERENCES

- [ADK] D. Auroux, S.K. Donaldson, L. Katzarkov, *Singular Lefschetz pencils*, Geom. Topol. **9** (2005), 1043–114.
- [AM] S. Akbulut, R. Matveyev, *A convex decomposition theorem for 4-manifolds*, International Math Research Notices **7** (1998), 371–381.
- [AO] S. Akbulut, B. Ozbagci, *Lefschetz fibrations on compact Stein surfaces*, Geom. Topol. **5** (2001), 319–334 (electronic).
- [B] F. Bogomolov, *Fillability of contact pseudoconvex manifolds*, Göttingen Univ. preprint, Heft **13** (1993), 1–13.
- [C] A. Cannas da Silva, *Fold-forms on four-folds*, preprint, 2002.
- [CGW] A. Cannas da Silva, V. Guillemin, C. Woodward, *On the unfolding of folded symplectic structures*, Mathematical Research Letters **7** (2000), 35–53.
- [D] S.K. Donaldson, *Lefschetz pencils on symplectic manifolds*, J. Differential Geom., **53** (2) (1999), 205–236.
- [E] J.B. Etnyre, *Symplectic convexity in low-dimensional topology*, Topology and its Appl. **88** (1998), 3–25.
- [EF] J.B. Etnyre, T. Fuller, *Realizing 4-Manifolds as Achiral Lefschetz Fibrations*, preprint, math.GT/0510008.
- [El1] Y. Eliashberg, *Topological characterization of Stein manifolds in dimension > 2* , Int. Journ. of Math. **1** (1990), 29–46.
- [El2] Y. Eliashberg, *A few remarks about symplectic filling*, Geom. Topol. **8** (2004), 277–293 (electronic).
- [ElGr] Y. Eliashberg, M. Gromov *Convex symplectic manifolds*, Proc. of Symp. in Pure Math. **57** (1991), 135–162.
- [Ga] D. Gay, *Explicit concave fillings of contact 3-manifolds*, Math. Proc. Cambridge. Philos. Soc. **133** (2002), 431–441.
- [Gi1] E. Giroux, *Structures de contact en dimension trois et bifurcations des feuilletages de surfaces*, Invent. Math. **141** no. 3 (2000), 615–689.
- [Gi2] E. Giroux, *Géométrie de contact: de la dimension trois vers les dimensions supérieures*, Proceedings of the ICM, Beijing 2002, vol. 2, 405–414.
- [Gra] H. Grauert, *On Levi’s problem*, Ann. of Math. **68** (1958), 460–472.
- [Go1] R. E. Gompf, *Sums of elliptic surfaces*, J. Differential Geom. **34** no.1 (1991), 93–114.
- [Go2] R. E. Gompf, *Handlebody construction of Stein surfaces*, Ann. of Math. **148** (1998), 619–693.
- [GS] R. E. Gompf, A.I. Stipsicz, *4-manifolds and Kirby calculus*, Graduate Studies in Math. **20**, Amer. Math. Soc., Providence, RI, 1999.
- [Ha] J. L. Harer, *Pencils of curves of 4-manifolds*. PhD thesis, Univeristy of California, Berkeley, 1979.
- [Ho] K. Honda, *On the classification of tight contact structures -I*, Geom. Topol. **4** (2000), 309–368 (electronic).
- [KM] P. B. Kronheimer, T. S. Mrowka, *Monopoles and contact structures*, Invent. Math. **130** no.2 (1997), 209–255.
- [LP] A. Loi, R. Piergallini, *Compact Stein surfaces with boundary as branched covers of B^4* , Invent. Math. **143** (2001), 325–348.
- [OS] B. Ozbagci, A. I. Stipsicz, *Surgery on contact 3-manifolds and Stein surfaces*, Bolyai Society Mathematical Studies, **13** (2004), Springer-Verlag, Berlin.
- [OS1] B. Ozbagci, A.I. Stipsicz, *Contact 3-manifolds with infinitely many Stein fillings*, Proc. Amer. Math. Soc. **132** no. 5 (2004), 1549–1558 (electronic).
- [P] O. Plamenevskaya, *Contact structures with distinct Heegaard Floer invariants*, Math. Res. Lett. **11** no. 4 (2004), 547–561.
- [Q] F. Quinn, *Dual decompositions of 4-manifolds*, Trans. Amer. Math. Soc. 354 **4** (2002), 1373–1392 (electronic).
- [T] W. P. Thurston, *Some simple examples of symplectic manifolds*, Proc. Amer. Math. Soc. **55** (1976), 467–468.
- [TW] W. P. Thurston, H. E. Winkelnkemper, *On the existence of contact forms*, Proc. Amer. Math. Soc. **52** (1975), 345–347.

- [vB] J. von Bergmann, *Pseudo-holomorphic maps into folded symplectic four-manifolds*, preprint, math.SG/0511597.

DEPARTMENT OF MATHEMATICS, MSU, EAST LANSING MI 48824, USA
E-mail address: baykur@msu.edu