

Noncommutative instantons from twisted conformal symmetries

Giovanni Landi¹, Walter van Suijlekom²

¹ Dipartimento di Matematica e Informatica, Università di Trieste
Via A. Valerio 12/1, I-34127 Trieste, Italy
landi@univ.trieste.it

² Max Planck Institute for Mathematics
Vivatsgasse 7, D-53111 Bonn, Germany
waltervs@mpim-bonn.mpg.de

23 January 2006

Abstract

We construct a five-parameter family of gauge-nonequivalent $SU(2)$ instantons on a noncommutative four sphere S_θ^4 and of topological charge equal to -1 . These instantons are critical points of a gauge functional and satisfy self-duality equations with respect to a Hodge star operator on forms on S_θ^4 . They are obtained by acting with a twisted conformal symmetry on a basic instanton canonically associated with a noncommutative instanton bundle on the sphere. A completeness argument for this family is obtained by means of index theorems. The dimension of the “tangent space” to the moduli space is computed as the index of a twisted Dirac operator and turns out to be equal to five, a number that survives deformation.

Contents

1	Introduction	3
2	Connections and gauge transformations	4
2.1	Connections on modules	4
2.2	Gauge transformations	7
3	Toric noncommutative manifolds M_θ	8
3.1	Deforming a torus action	9
3.2	The manifold M_θ as a fixed point algebra	11
3.3	Vector bundles on M_θ	12
3.4	Differential calculus on M_θ	14
4	Gauge theory on the sphere S_θ^4	17
4.1	The principal fibration $S_{\theta'}^7 \rightarrow S_\theta^4$	17
4.2	Associated bundles	19
4.3	Yang-Mills theory on S_θ^4	22
5	Construction of $SU(2)$-instantons on S_θ^4	24
5.1	The basic instanton	24
5.2	Twisted infinitesimal symmetries	26
5.3	Twisted conformal transformations	29
5.4	Local expressions	34
5.5	Moduli space of instantons	35
5.6	Dirac operator associated to the complex	37
6	Towards Yang-Mills theory on M_θ	38
7	Final remarks	40
A	Local index formula	41

1 Introduction

The importance of Yang-Mills instantons in physics and mathematics needs not be stressed. They have played a central role since their first appearance [9] and are most elegantly described via the so-called ADHM construction [5, 4]. The generalization in [36] of this method for instantons on a noncommutative space \mathbb{R}^4 has found several important applications notably in brane and superconformal theories.

Toric noncommutative manifolds M_Θ were constructed and studied in [19]. One starts with any (Riemannian spin) manifold M carrying a torus action and then deforms the torus to a noncommutative one governed by a real antisymmetric matrix Θ of deformation parameters. The starting example of [19] – the archetype of all these deformations – was a four dimensional sphere S_θ^4 , which came with a natural noncommutative instanton bundle endowed with a natural connection. At the classical value of the deformation parameter, $\theta = 0$, the bundle and the connection reduces to the one of [9]. The present sphere S_θ^4 can be thought of [18] as a one point compactification of a noncommutative \mathbb{R}_θ which is structurally different from the one considered in [36].

In [31] this basic noncommutative instanton was put in the context of an $SU(2)$ noncommutative principal fibration $S_\theta^7 \rightarrow S_\theta^4$ over S_θ^4 . In the present paper, we continue the analysis by first putting it in the setting of a noncommutative Yang-Mills theory. We will then construct a five-parameter family of (infinitesimal) gauge-nonequivalent instantons, by acting with twisted conformal symmetries on the basic instanton. All these instantons will be gauge configurations satisfying self-duality equations – with a suitable defined Hodge $*_\theta$ -operator on forms $\Omega(S_\theta^4)$ – and will have a ‘topological charge’ of values -1 . A completeness argument on the family of instantons is provided by index theoretical arguments, similar to the one in [6] for undeformed instantons on S^4 . The dimension of the “tangent” of the moduli space can be computed as the index of a twisted Dirac operator which turns out to be equal to its classical value that is five. The twisting of the conformal symmetry is implemented with a Drinfel’d twist [23] and gives rise to a deformed Hopf algebra $U_\theta(so(5, 1))$. That these are conformal infinitesimal transformations is stressed by the fact that the Hopf algebra $U_\theta(so(5, 1))$ leaves the Hodge $*_\theta$ -structure of $\Omega(S_\theta^4)$ invariant.

The paper is organized as follows. In Section 2 we recall the setting of gauge theories (connections) and gauge transformations on finite projective modules (the substitute for vector bundles) over algebras (the substitute for spaces). We also show how to implement a Bianchi identity that will be crucial later on for the self-duality equations.

Section 3 deals with toric noncommutative manifolds. These were indeed named isospectral deformations in that they can be endowed with the structure of a noncommutative Riemannian spin manifold via a spectral triple $(C^\infty(M_\theta), D, \mathcal{H})$ with the properties of [16]. For this class of examples, the Dirac operator D is the classical one and $\mathcal{H} = L^2(M, \mathcal{S})$ is the usual Hilbert space of spinors on which the algebra $C^\infty(M_\theta)$ acts in a twisted manner. Thus one twists the algebra and its representation while keeping the geometry unchanged. The resulting noncommutative geometry is isospectral and all spectral properties are preserved including the dimension. Both the algebra and its action on spinors can be given via a “star-type” or “Moyal-type” product.

In Section 4 we specialize to gauge theories on the sphere S_θ^4 and introduce a Yang-Mills functional from which we derive field equations (equations for critical points) as

well as a topological action.

The heart of the paper is Section 5 where we explicitly construct instantons. As usual, these are gauge configurations which are solutions of (anti)self-duality equations and realize absolute minima of the Yang-Mills functional. We start from a basic instanton which is shown to be invariant under twisted orthogonal transformations in $U_\theta(\mathfrak{so}(5))$. We then perturb it by the action of conformal operators in $U_\theta(\mathfrak{so}(5, 1)) - U_\theta(\mathfrak{so}(5))$ producing a five parameter family of new, not gauge equivalent instantons. A completeness argument is obtained by using an index theorem to compute the dimension of the tangent space of the moduli space of instantons on S_θ^4 , which is shown to be just five. The relevant material from noncommutative index theory is recalled in the appendix.

Section 6 sketches a general scheme for gauge theories on four dimensional toric non-commutative manifolds.

Finally in Section 7 we mention on the recent work of twisting symmetries on the usual Moyal planes and their use in the context of noncommutative field theories on these deformed spaces.

It is worth stressing that in the present paper we change orientation with respect to our previous papers [19, 31, 28]. This results in changing sign to the topological charges and into self-dual solutions being converted into anti self-dual solutions and viceversa.

2 Connections and gauge transformations

We first review the notion of a (gauge) connection on a (finite projective) module \mathcal{E} over an algebra \mathcal{A} with respect to a given calculus; we take a right module structure. Also, we recall gauge transformations in this setting.

2.1 Connections on modules

Let us suppose we have an algebra \mathcal{A} with a differential calculus $(\Omega\mathcal{A} = \bigoplus_p \Omega^p \mathcal{A}, d)$. A *connection* on the right \mathcal{A} -module \mathcal{E} is a \mathbb{C} -linear map

$$\nabla : \mathcal{E} \otimes_{\mathcal{A}} \Omega^p \mathcal{A} \longrightarrow \mathcal{E} \otimes_{\mathcal{A}} \Omega^{p+1} \mathcal{A},$$

defined for any $p \geq 0$, and satisfying the Leibniz rule

$$\nabla(\omega\rho) = (\nabla\omega)\rho + (-1)^p \omega d\rho, \quad \forall \omega \in \mathcal{E} \otimes_{\mathcal{A}} \Omega^p \mathcal{A}, \quad \rho \in \Omega\mathcal{A}.$$

A connection is completely determined by its restriction

$$\nabla : \mathcal{E} \rightarrow \mathcal{E} \otimes_{\mathcal{A}} \Omega^1 \mathcal{A}, \tag{2.1}$$

which satisfies

$$\nabla(\eta a) = (\nabla\eta)a + \eta \otimes_{\mathcal{A}} da, \quad \forall \eta \in \mathcal{E}, \quad a \in \mathcal{A}, \tag{2.2}$$

and which is extended by the Leibniz rule. It is again the latter property that implies the $\Omega\mathcal{A}$ -linearity of the composition,

$$\nabla^2 = \nabla \circ \nabla : \mathcal{E} \otimes_{\mathcal{A}} \Omega^p \mathcal{A} \longrightarrow \mathcal{E} \otimes_{\mathcal{A}} \Omega^{p+2} \mathcal{A}.$$

Indeed, for any $\omega \in \mathcal{E} \otimes_{\mathcal{A}} \Omega^p \mathcal{A}$, $\rho \in \Omega \mathcal{A}$ one has $\nabla^2(\omega\rho) = \nabla((\nabla\omega)\rho + (-1)^p \omega d\rho) = (\nabla^2\omega)\rho + (-1)^{p+1}(\nabla\omega)d\rho + (-1)^p(\nabla\omega)d\rho + \omega d^2\rho = (\nabla^2\omega)\rho$. The restriction of ∇^2 to \mathcal{E} is the *curvature*

$$F : \mathcal{E} \rightarrow \mathcal{E} \otimes_{\mathcal{A}} \Omega^2 \mathcal{A}, \quad (2.3)$$

of the connection. It is \mathcal{A} -linear, $F(\eta a) = F(\eta)a$ for any $\eta \in \mathcal{E}$, $a \in \mathcal{A}$, and satisfies

$$\nabla^2(\eta \otimes_{\mathcal{A}} \rho) = F(\eta)\rho, \quad \forall \eta \in \mathcal{E}, \rho \in \Omega \mathcal{A}. \quad (2.4)$$

Thus, $F \in \text{End}_{\mathcal{A}}(\mathcal{E}, \mathcal{E} \otimes_{\mathcal{A}} \Omega^2 \mathcal{A})$, the latter being the collection of (right) \mathcal{A} -linear endomorphisms of \mathcal{E} , with values in the two-forms $\Omega^2 \mathcal{A}$.

In order to have the notion of a Bianchi identity we need some generalization. Let $\text{End}_{\Omega \mathcal{A}}(\mathcal{E} \otimes_{\mathcal{A}} \Omega \mathcal{A})$ be the collection of all $\Omega \mathcal{A}$ -linear endomorphisms of $\mathcal{E} \otimes_{\mathcal{A}} \Omega \mathcal{A}$. It is an algebra under composition. The curvature F can be thought of as an element of $\text{End}_{\Omega \mathcal{A}}(\mathcal{E} \otimes_{\mathcal{A}} \Omega \mathcal{A})$. There is then a well-defined map

$$\begin{aligned} [\nabla, \cdot] &: \text{End}_{\Omega \mathcal{A}}(\mathcal{E} \otimes_{\mathcal{A}} \Omega \mathcal{A}) \longrightarrow \text{End}_{\Omega \mathcal{A}}(\mathcal{E} \otimes_{\mathcal{A}} \Omega \mathcal{A}) \\ [\nabla, T] &:= \nabla \circ T - (-1)^{|T|} T \circ \nabla. \end{aligned} \quad (2.5)$$

where $|T|$ denotes the degree of T with respect to the \mathbb{Z}^2 -grading of $\Omega \mathcal{A}$. Indeed, for any $\omega \in \mathcal{E} \otimes_{\mathcal{A}} \Omega^p \mathcal{A}$, $\rho \in \Omega \mathcal{A}$, it follows that

$$\begin{aligned} [\nabla, T](\omega\rho) &= \nabla(T(\omega\rho)) - (-1)^{|T|} T(\nabla(\omega\rho)) \\ &= \nabla(T(\omega)\rho) - (-1)^{|T|} T((\nabla\omega)\rho + (-1)^p \omega d\rho) \\ &= (\nabla(T(\omega)))\rho + (-1)^{p+|T|} T(\omega)d\rho - (-1)^{|T|} T(\nabla\omega)\rho - (-1)^{p+|T|} T(\omega)d\rho \\ &= (\nabla(T(\omega)) - (-1)^{|T|} T(\nabla\omega))\rho = ([\nabla, T](\omega))\rho. \end{aligned}$$

It is easily checked that $[\nabla, \cdot]$ is a graded derivation for the algebra $\text{End}_{\Omega \mathcal{A}}(\mathcal{E} \otimes_{\mathcal{A}} \Omega \mathcal{A})$,

$$[\nabla, S \circ T] = [\nabla, S] \circ T + (-1)^{|S|} S \circ [\nabla, T]. \quad (2.6)$$

Proposition 1. *The curvature F satisfies the Bianchi identity,*

$$[\nabla, F] = 0. \quad (2.7)$$

Proof. Since $F \in \text{End}_{\Omega \mathcal{A}}^0(\mathcal{E} \otimes_{\mathcal{A}} \Omega \mathcal{A})$, the map $[\nabla, F]$ makes sense. Furthermore,

$$[\nabla, F] = \nabla \circ \nabla^2 - \nabla^2 \circ \nabla = \nabla^3 - \nabla^3 = 0.$$

□

In Section II.2 of [14], such a Bianchi identity was implicitly used in the construction of a so-called canonical cycle from a connection on a finite projective \mathcal{A} -module \mathcal{E} .

Connections always exist on a projective module. On the free module $\mathcal{E} = \mathbb{C}^N \otimes_{\mathbb{C}} \mathcal{A} \simeq \mathcal{A}^N$, a connection is given by the operator

$$\nabla_0 = \mathbb{I} \otimes d : \mathbb{C}^N \otimes_{\mathbb{C}} \Omega^p \mathcal{A} \longrightarrow \mathbb{C}^N \otimes_{\mathbb{C}} \Omega^{p+1} \mathcal{A}.$$

With the canonical identification $\mathbb{C}^N \otimes_{\mathbb{C}} \Omega \mathcal{A} = (\mathbb{C}^N \otimes_{\mathbb{C}} \mathcal{A}) \otimes_{\mathcal{A}} \Omega \mathcal{A} \simeq (\Omega \mathcal{A})^N$, one thinks of ∇_0 as acting on $(\Omega \mathcal{A})^N$ as the operator $\nabla_0 = (d, d, \dots, d)$ (N -times). For

a generic projective module \mathcal{E} one has a canonical inclusion map, $\lambda : \mathcal{E} \rightarrow \mathcal{A}^N$, which identifies \mathcal{E} as a direct summand of the free module \mathcal{A}^N and a canonical idempotent $p : \mathcal{A}^N \rightarrow \mathcal{E}$ which allows one to identify $\mathcal{E} = p\mathcal{A}^N$. Using these maps and their natural extensions to \mathcal{E} -valued forms, a connection ∇_0 on \mathcal{E} (called *Levi-Civita* or *Grassmann*) is the composition,

$$\mathcal{E} \otimes_{\mathcal{A}} \Omega^p \mathcal{A} \xrightarrow{\lambda} \mathbb{C}^N \otimes_{\mathbb{C}} \Omega^p \mathcal{A} \xrightarrow{\mathbb{I} \otimes d} \mathbb{C}^N \otimes_{\mathbb{C}} \Omega^{p+1} \mathcal{A} \xrightarrow{p} \mathcal{E} \otimes_{\mathcal{A}} \Omega^{p+1} \mathcal{A},$$

that is

$$\nabla_0 = p \circ (\mathbb{I} \otimes d) \circ \lambda. \quad (2.8)$$

One indicates it simply by $\nabla_0 = pd$. The existence of a connection on the module \mathcal{E} is completely equivalent to it being projective [21]. Furthermore, the space $C(\mathcal{E})$ of all connections on \mathcal{E} is an affine space modeled on $\text{End}_{\mathcal{A}}(\mathcal{E}, \mathcal{E} \otimes_{\mathcal{A}} \Omega^1 \mathcal{A})$. Indeed, if ∇_1, ∇_2 are two connections on \mathcal{E} , their difference is \mathcal{A} -linear,

$$(\nabla_1 - \nabla_2)(\eta a) = ((\nabla_1 - \nabla_2)(\eta))a, \quad \forall \eta \in \mathcal{E}, a \in \mathcal{A},$$

so that $\nabla_1 - \nabla_2 \in \text{End}_{\mathcal{A}}(\mathcal{E}, \mathcal{E} \otimes_{\mathcal{A}} \Omega^1 \mathcal{A})$. Thus, any connection can be written as

$$\nabla = pd + \alpha, \quad (2.9)$$

where α is any element in $\text{End}_{\mathcal{A}}(\mathcal{E}, \mathcal{E} \otimes_{\mathcal{A}} \Omega^1 \mathcal{A})$. The ‘‘matrix of 1-forms’’ α as in (2.9) is called the *gauge potential* of the connection ∇ . The corresponding curvature F of ∇ is

$$F = pdpd + pd\alpha + \alpha^2. \quad (2.10)$$

Next, let the algebra \mathcal{A} have an involution $*$ extended to the whole of $\Omega\mathcal{A}$ by the requirement $(da)^* = da^*$ for any $a \in \mathcal{A}$. A *Hermitian structure* on the module \mathcal{E} is a map $\langle \cdot, \cdot \rangle : \mathcal{E} \otimes \mathcal{E} \rightarrow \mathcal{A}$ with the properties

$$\begin{aligned} \langle \eta a, \xi \rangle &= a^* \langle \xi, \eta \rangle, & \langle \eta, \xi \rangle^* &= \langle \xi, \eta \rangle, \\ \langle \eta, \eta \rangle &\geq 0, \langle \eta, \eta \rangle = 0 &\iff \eta = 0, \end{aligned} \quad (2.11)$$

for any $\eta, \xi \in \mathcal{E}$ and $a \in \mathcal{A}$ (an element $a \in \mathcal{A}$ is positive if it is of the form $a = b^*b$ for some $b \in \mathcal{A}$). The Hermitian structure is naturally extended to a linear map from $\mathcal{E} \otimes_{\mathcal{A}} \Omega\mathcal{A} \times \mathcal{E} \otimes_{\mathcal{A}} \Omega\mathcal{A}$ to $\Omega\mathcal{A}$ by

$$\langle \eta \otimes_{\mathcal{A}} \omega, \xi \otimes_{\mathcal{A}} \rho \rangle = (-1)^{|\eta||\omega|} \omega^* \langle \eta, \xi \rangle \rho, \quad \forall \eta, \xi \in \mathcal{E} \otimes_{\mathcal{A}} \Omega\mathcal{A}, \omega, \rho \in \Omega\mathcal{A}. \quad (2.12)$$

A connection ∇ on \mathcal{E} and a Hermitian structure $\langle \cdot, \cdot \rangle$ on \mathcal{E} are said to be compatible if the following condition is satisfied [15],

$$\langle \nabla \eta, \xi \rangle + \langle \eta, \nabla \xi \rangle = d \langle \eta, \xi \rangle, \quad \forall \eta, \xi \in \mathcal{E}. \quad (2.13)$$

It follows directly from the Leibniz rule and (2.12) that this extends to

$$\langle \nabla \eta, \xi \rangle + (-1)^{|\eta|} \langle \eta, \nabla \xi \rangle = d \langle \eta, \xi \rangle, \quad \forall \eta, \xi \in \mathcal{E} \otimes_{\mathcal{A}} \Omega\mathcal{A}. \quad (2.14)$$

Compatible connections always exist. Indeed, any Hermitian structure on $\mathcal{E} = p\mathcal{A}^N$ can be written as $\langle \eta, \xi \rangle = \sum_{j=1}^N \eta_j^* \xi_j$ with $\eta = p\eta = (\eta_1, \dots, \eta_N)$ and the same for ξ . Then the Grassmann connection (2.8) is easily seen to be compatible,

$$d \langle \eta, \xi \rangle = \langle \nabla_0 \eta, \xi \rangle + \langle \eta, \nabla_0 \xi \rangle. \quad (2.15)$$

For a general connection (2.9), the compatibility with the Hermitian structure reduces to

$$\langle \alpha \eta, \xi \rangle + \langle \eta, \alpha \xi \rangle = 0, \quad \forall \eta, \xi \in \mathcal{E}, \quad (2.16)$$

which just says that the gauge potential is skew-hermitian,

$$\alpha^* = -\alpha. \quad (2.17)$$

We still use the symbol $C(\mathcal{E})$ to denote the space of compatible connections on \mathcal{E} .

Let $\text{End}_{\Omega\mathcal{A}}^s(\mathcal{E} \otimes_{\mathcal{A}} \Omega\mathcal{A})$ denote the space of elements T in $\text{End}_{\Omega\mathcal{A}}(\mathcal{E} \otimes_{\mathcal{A}} \Omega\mathcal{A})$ which are skew-hermitian with respect to the Hermitian structure (2.12), i.e. satisfying

$$\langle T\eta, \xi \rangle + \langle \eta, T\xi \rangle = 0, \quad \forall \eta, \xi \in \mathcal{E}. \quad (2.18)$$

Proposition 2. *The map $[\nabla, \cdot]$ in (2.5) restricts to $\text{End}_{\Omega\mathcal{A}}^s(\mathcal{E} \otimes_{\mathcal{A}} \Omega\mathcal{A})$ as a derivation*

$$[\nabla, \cdot] : \text{End}_{\Omega\mathcal{A}}^s(\mathcal{E} \otimes_{\mathcal{A}} \Omega\mathcal{A}) \longrightarrow \text{End}_{\Omega\mathcal{A}}^s(\mathcal{E} \otimes_{\mathcal{A}} \Omega\mathcal{A}), \quad (2.19)$$

Proof. Let $T \in \text{End}_{\Omega\mathcal{A}}^s(\mathcal{E} \otimes_{\mathcal{A}} \Omega\mathcal{A})$ be of order $|T|$; it then satisfies

$$\langle T\eta, \xi \rangle + (-1)^{|\eta||T|} \langle \eta, T\xi \rangle = 0, \quad (2.20)$$

for $\eta, \xi \in \mathcal{E} \otimes_{\mathcal{A}} \Omega\mathcal{A}$. Since $[\nabla, T]$ is $\Omega\mathcal{A}$ -linear, it is enough to show that

$$\langle [\nabla, T]\eta, \xi \rangle + \langle \eta, [\nabla, T]\xi \rangle = 0, \quad \forall \eta, \xi \in \mathcal{E}.$$

This follows from equations (2.20) and (2.14),

$$\begin{aligned} \langle [\nabla, T]\eta, \xi \rangle + \langle \eta, [\nabla, T]\xi \rangle &= \langle \nabla T\eta, \xi \rangle - (-1)^{|T|} \langle T\nabla\eta, \xi \rangle + \langle \eta, \nabla T\xi \rangle - (-1)^{|T|} \langle \eta, T\nabla\xi \rangle \\ &= \langle \nabla T\eta, \xi \rangle - \langle \nabla\eta, T\xi \rangle + \langle \eta, \nabla T\xi \rangle - (-1)^{|T|} \langle T\eta, \nabla\xi \rangle \\ &= d(\langle T\eta, \xi \rangle + \langle \eta, T\xi \rangle) = 0. \end{aligned}$$

□

2.2 Gauge transformations

An \mathcal{A} -linear map $T : \mathcal{E} \rightarrow \mathcal{E}$ is said to be adjointable if it admits an adjoint, i.e. there exists an \mathcal{A} -linear map $T^* : \mathcal{E} \rightarrow \mathcal{E}$ such that

$$\langle T^*\eta, \xi \rangle = \langle \eta, T\xi \rangle, \quad \forall \eta, \xi \in \mathcal{E}.$$

The collection $\text{End}_{\mathcal{A}}(\mathcal{E})$ of all \mathcal{A} -linear adjointable maps is an algebra with involution; its elements are also called endomorphisms of \mathcal{E} . The group $\mathcal{U}(\mathcal{E})$ of unitary endomorphisms of \mathcal{E} is given by

$$\mathcal{U}(\mathcal{E}) := \{u \in \text{End}_{\mathcal{A}}(\mathcal{E}) \mid uu^* = u^*u = \text{id}_{\mathcal{E}}\}. \quad (2.21)$$

This group plays the role of the *infinite dimensional group of gauge transformations*. It naturally acts on compatible connections by

$$(u, \nabla) \mapsto \nabla^u := u^* \nabla u, \quad \forall u \in \mathcal{U}(\mathcal{E}), \nabla \in C(\mathcal{E}), \quad (2.22)$$

where u^* is really $u^* \otimes \text{id}_{\Omega_{\mathcal{A}}}$; this will always be understood in the following. Then the curvature transforms in a covariant way

$$(u, F) \mapsto F^u = u^* F u, \quad (2.23)$$

since, evidently, $F^u = (\nabla^u)^2 = u^* \nabla u u^* \nabla u^* = u^* \nabla^2 u = u^* F u$.

As for the gauge potential, one has the usual affine transformation

$$(u, \alpha) \mapsto \alpha^u := u^* p du + u^* \alpha u. \quad (2.24)$$

Indeed, $\nabla^u(\eta) = u^*(pd + \alpha)u\eta = u^*pd(u\eta) + u^*\alpha u\eta = u^*pud\eta + u^*p(du)\eta + u^*\alpha u\eta = pd\eta + (u^*pdu + u^*\alpha u)\eta$ for any $\eta \in \mathcal{E}$, which yields (2.24) for the transformed potential.

The ‘tangent vectors’ to the gauge group $\mathcal{U}(\mathcal{E})$ constitute the vector space of infinitesimal gauge transformations. For $X \in \text{End}(\mathcal{E})$ we define a family $\{u_t\}_{t \in \mathbb{R}}$ of elements in $\mathcal{U}(\mathcal{E})$ by $u_t = 1 + tX + \mathcal{O}(t^2)$, so that $X = (\partial u_t / \partial t)_{t=0}$. Unitarity of u_t becomes $(1 + t(X + X^*) + \mathcal{O}(t^2)) = 1$. If we take derivatives with respect to t , putting $t = 0$ afterwards, we find $X = -X^*$. In other words, for u_t to be a gauge transformation, X should be a skew-hermitian endomorphisms of \mathcal{E} . In this way, we understand $\text{End}_{\mathcal{A}}^s(\mathcal{E})$ as the collection of *infinitesimal gauge transformations*. It is a real vector space whose complexification $\text{End}_{\mathcal{A}}^s(\mathcal{E}) \otimes_{\mathbb{R}} \mathbb{C}$ can be identified with $\text{End}_{\mathcal{A}}(\mathcal{E})$.

Infinitesimal gauge transformations act on a connection in a natural way. Let the above gauge transformation u_t act on ∇ as in (2.22). Since $(\partial(u_t \nabla u_t^*) / \partial t)_{t=0} = [\nabla, X]$, we conclude that an element $X \in \text{End}_{\mathcal{A}}^s(\mathcal{E})$ acts infinitesimally on a connection ∇ by the addition of $[\nabla, X]$,

$$(X, \nabla) \mapsto \nabla^X = \nabla + t[\nabla, X] + \mathcal{O}(t^2), \quad \forall X \in \text{End}_{\mathcal{A}}^s(\mathcal{E}), \nabla \in C(\mathcal{E}). \quad (2.25)$$

As a consequence, for the transformed curvature one has

$$(X, F) \mapsto F^X = F + t[F, X] + \mathcal{O}(t^2), \quad (2.26)$$

since $F^X = (\nabla + t[\nabla, X]) \circ (\nabla + t[\nabla, X]) = \nabla^2 + t[\nabla^2, X] + \mathcal{O}(t^2)$.

3 Toric noncommutative manifolds M_θ

We start by recalling the general construction of toric noncommutative manifolds given in [19] where they were called isospectral deformations. These are deformations of a classical Riemannian manifold and satisfy all the properties of noncommutative spin geometry [16]. They are the content of the following result taken from [19],

Theorem 3. *Let M be a compact spin Riemannian manifold whose isometry group has rank $r \geq 2$. Then M admits a natural one parameter isospectral deformation to noncommutative geometries M_θ .*

The idea of the construction is to deform the standard spectral triple describing the Riemannian geometry of M along a torus embedded in the isometry group, thus obtaining a family of spectral triples describing noncommutative geometries.

3.1 Deforming a torus action

Let M be an m dimensional compact Riemannian manifold equipped with an isometric smooth action σ of an n -torus \mathbb{T}^n , $n \geq 2$. We denote by σ also the corresponding action of \mathbb{T}^n by automorphisms on the algebra $C^\infty(M)$ of smooth functions on M , obtained by pull-back. The algebra $C^\infty(M)$ may be decomposed into spectral subspaces which are indexed by the dual group $\mathbb{Z}^n = \widehat{\mathbb{T}^n}$. Now, with $s = (s_1, \dots, s_n) \in \mathbb{T}^n$, each $r \in \mathbb{Z}^n$ labels a character $e^{2\pi i s} \mapsto e^{2\pi i r \cdot s}$ of \mathbb{T}^n , with the scalar product $r \cdot s := r_1 s_1 + \dots + r_n s_n$. The r -th spectral subspace for the action σ of \mathbb{T}^n on $C^\infty(M)$ consists of those smooth functions f_r for which

$$\sigma_s(f_r) = e^{2\pi i r \cdot s} f_r, \quad (3.1)$$

and each $f \in C^\infty(M)$ is the sum of a unique (rapidly convergent) series $f = \sum_{r \in \mathbb{Z}^n} f_r$. Let now $\theta = (\theta_{jk} = -\theta_{kj})$ be a real antisymmetric $n \times n$ matrix. The θ -deformation of $C^\infty(M)$ may be defined by replacing the ordinary product by a deformed product, given on spectral subspaces by

$$f_r \times_\theta g_{r'} := f_r \sigma_{r \cdot \theta}(g_{r'}) = e^{2\pi i \theta \cdot r r'} f_r g_{r'}, \quad (3.2)$$

where $r \cdot \theta = (r_j \theta_{j1}, \dots, r_j \theta_{jn}) \in \mathbb{T}^n$. This product is then extended linearly to all functions in $C^\infty(M)$. We denote $C^\infty(M_\theta) := (C^\infty(M), \times_\theta)$ and note that the action σ of \mathbb{T}^n on $C^\infty(M)$ extends to an action on $C^\infty(M_\theta)$ given again by (3.1) on the homogeneous elements.

Next, let us take M to be a spin manifold with $\mathcal{H} := L^2(M, \mathcal{S})$ the Hilbert space of spinors and D the usual Dirac operator of the metric of M . Smooth functions act on spinors by pointwise multiplication thus giving a representation $\pi : C^\infty(M) \rightarrow \mathcal{B}(\mathcal{H})$, the latter being the algebra of bounded operators on \mathcal{H} .

There is a double cover $c : \widetilde{\mathbb{T}^n} \rightarrow \mathbb{T}^n$ and a representation of $\widetilde{\mathbb{T}^n}$ on \mathcal{H} by unitary operators $U(s)$, $s \in \widetilde{\mathbb{T}^n}$, so that

$$U(s)DU(s)^{-1} = D, \quad (3.3)$$

since the torus action is assumed to be isometric, and such that for all $f \in C^\infty(M)$,

$$U(s)\pi(f)U(s)^{-1} = \pi(\sigma_{c(s)}(f)). \quad (3.4)$$

Recall that an element $T \in \mathcal{B}(\mathcal{H})$ is called smooth for the action of $\widetilde{\mathbb{T}^n}$ if the map

$$\widetilde{\mathbb{T}^n} \ni s \mapsto \alpha_s(T) := U(s)TU(s)^{-1},$$

is smooth for the norm topology. From its very definition, α_s coincides on $\pi(C^\infty(M)) \subset \mathcal{B}(\mathcal{H})$ with the automorphism $\sigma_{c(s)}$. Much as it was done before for the smooth functions, we shall use the torus action to give a spectral decomposition of smooth elements of $\mathcal{B}(\mathcal{H})$. Any such a smooth element T is written as a (rapidly convergent) series $T = \sum T_r$ with $r \in \mathbb{Z}$ and each T_r is homogeneous of degree r under the action of $\widetilde{\mathbb{T}^n}$, i.e.

$$\alpha_s(T_r) = e^{2\pi i r \cdot s} T_r, \quad \forall s \in \widetilde{\mathbb{T}^n}. \quad (3.5)$$

Let (p_1, p_2, \dots, p_n) be the infinitesimal generators of the action of $\widetilde{\mathbb{T}}^n$ so that we can write $U(s) = \exp 2\pi i s \cdot p$. Now, with θ a real $n \times n$ anti-symmetric matrix as above, one defines a twisted representation of the smooth elements $\mathcal{B}(\mathcal{H})$ on \mathcal{H} by

$$L_\theta(T) := \sum_r \text{Tr}_r U(r_j \theta_{j_1}, \dots, r_j \theta_{j_n}) = \sum_r \text{Tr}_r \exp \{2\pi i r_j \theta_{jk} p_k\}, \quad (3.6)$$

Taking smooth functions on M as elements of $\mathcal{B}(\mathcal{H})$, via the representation π , the previous definition gives an algebra $L_\theta(C^\infty(M))$ which we may think of as a representation (as bounded operators on \mathcal{H}) of the algebra $C^\infty(M_\theta)$. Indeed, by the very definition of the product \times_θ in (3.2) one establishes that

$$L_\theta(f \times_\theta g) = L_\theta(f)L_\theta(g), \quad (3.7)$$

proving that the algebra $C^\infty(M)$ equipped with the product \times_θ is isomorphic to the algebra $L_\theta(C^\infty(M))$. It is shown in [39] that there is a natural completion of the algebra $C^\infty(M_\theta)$ to a C^* -algebra $C(M_\theta)$ whose smooth subalgebra – under the extended action of \mathbb{T}^n – is precisely $C^\infty(M_\theta)$. Thus, we can understand L_θ as a *quantization map* from $C^\infty(M)$ to $C^\infty(M_\theta)$ which provides a strict deformation quantization in the sense of Rieffel. More generally, in [39] one considers a (not necessarily commutative) C^* -algebra A carrying an action of \mathbb{R}^n . For an anti-symmetric $n \times n$ matrix θ , one defines a star product \times_θ between elements in A much as we did before. The algebra A equipped with the product \times_θ gives rise to a C^* -algebra denoted by A_θ . Then the collection $\{A_{\hbar\theta}\}_{\hbar \in [0,1]}$ is a continuous family of C^* -algebras providing a strict deformation quantization in the direction of the Poisson structure on A defined by the matrix θ .

Our case of interest corresponds to the choice $A = C(M)$ with an action of \mathbb{R}^n that is periodic or, in other words, an action of \mathbb{T}^n . The smooth elements in the deformed algebra make up the algebra $C^\infty(M_\theta)$.

The quantization map will play a key role in what follows, allowing us to extend differential geometric techniques from M to the noncommutative space M_θ .

It was shown in [19] that $(L_\theta(C^\infty(M)), \mathcal{H}, D)$ satisfies all axioms of a noncommutative spin geometry [17, 26]; there is also a grading γ (for the even case) and a real structure J . In particular, boundedness of the commutators $[D, L_\theta(f)]$ for $f \in C^\infty(M)$ follows from $[D, L_\theta(f)] = L_\theta([D, f])$, D being of degree 0 (since \mathbb{T}^n acts by isometries, each p_μ commutes with D). This noncommutative geometry is an isospectral deformation of the classical Riemannian geometry of M , in that the spectrum of the operator D coincides with that of the Dirac operator D on M . Thus all spectral properties are unchanged. In particular, the triples are m^+ -summable and there is a noncommutative integral as a Dixmier trace [22],

$$\int L_\theta(f) := \text{Tr}_\omega(f|D|^{-m}), \quad (3.8)$$

with $f \in C^\infty(M_\theta)$ understood in its representation on \mathcal{H} . A drastic simplification of this noncommutative integral is given by the Lemma [24].

Lemma 4. *If $f \in C^\infty(M)$ then*

$$\int L_\theta(f) = \int_M f d\nu.$$

Proof. Any element $f \in C^\infty(M)$ is given as an infinite sum of functions that are homogeneous under the action of \mathbb{T}^n . Let us therefore assume that f is homogeneous of degree k so that $\sigma_s(L_\theta(f)) = L_\theta(\sigma_s(f)) = e^{2\pi i k \cdot s} L_\theta(f)$. From the tracial property of the noncommutative integral and the invariance of D under the action of \mathbb{T}^n , we see that

$$\mathrm{Tr}_\omega(\sigma_s(L_\theta(f))|D|^{-m}) = \mathrm{Tr}_\omega(U(s)L_\theta(f)U(s)^{-1}|D|^{-m}) = \mathrm{Tr}_\omega(L_\theta(f)|D|^{-m}).$$

In other words, $e^{2\pi i k \cdot s} \mathrm{Tr}_\omega(L_\theta(f)|D|^{-m}) = \mathrm{Tr}_\omega(L_\theta(f)|D|^{-m})$ from which we infer that this trace vanishes if $k \neq 0$. If $k = 0$, then $L_\theta(f) = f$, leading to the desired result. \square

3.2 The manifold M_θ as a fixed point algebra

A different but equivalent approach to these noncommutative manifolds M_θ was introduced in [18]. In there the algebra $C^\infty(M_\theta)$ is identified as a fixed point subalgebra of $C^\infty(M) \otimes C^\infty(\mathbb{T}_\theta^n)$ where $C^\infty(\mathbb{T}_\theta^n)$ is the algebra of smooth functions on the noncommutative torus. This identification allows one to extend techniques from commutative differential geometry on M to the noncommutative space M_θ .

We recall the definition of the noncommutative n -torus \mathbb{T}_θ^n [38]. Let $\theta = (\theta_{\mu\nu})$ be a real $n \times n$ anti-symmetric matrix as before, and $\lambda^{\mu\nu} = e^{2\pi i \theta_{\mu\nu}}$. The unital $*$ -algebra $\mathcal{A}(\mathbb{T}_\theta^n)$ of polynomial functions on \mathbb{T}_θ^n is generated by n unitary elements U^μ , $\mu = 1, \dots, n$, with relations

$$U^\mu U^\nu = \lambda^{\mu\nu} U^\nu U^\mu, \quad \mu, \nu = 1, \dots, n. \quad (3.9)$$

The polynomial algebra is extended to the universal C^* -algebra with the same generators. There is a natural action of \mathbb{T}^n on $\mathcal{A}(\mathbb{T}_\theta^n)$ by $*$ -automorphisms given by $\tau_s(U^\mu) = e^{2\pi i s_\mu} U^\mu$ with $s = (s_\mu) \in \mathbb{T}^n$. The corresponding infinitesimal generators X_μ of the action are algebra derivations given explicitly on the generators by $X_\mu(U^\nu) = 2\pi i \delta_\mu^\nu$. They are used [13] to construct the pre- C^* -algebra $C^\infty(\mathbb{T}_\theta^n)$ of smooth functions on \mathbb{T}_θ^n , which is the completion of $\mathcal{A}(\mathbb{T}_\theta^n)$ with respect to the locally convex topology generated by the seminorms,

$$\|u\|_r := \sup_{r_1 + \dots + r_n \leq r} \|X_1^{r_1} \cdots X_n^{r_n}(u)\|, \quad (3.10)$$

and $\|\cdot\|$ is the C^* -norm. The algebra $C^\infty(\mathbb{T}_\theta^n)$ turns out to be a nuclear Fréchet space and one can unambiguously take the completed tensor product $C^\infty(M) \overline{\otimes} C^\infty(\mathbb{T}_\theta^n)$. Then, one defines $(C^\infty(M) \overline{\otimes} C^\infty(\mathbb{T}_\theta^n))^{\sigma \otimes \tau^{-1}}$ as the fixed point subalgebra of $C^\infty(M) \overline{\otimes} C^\infty(\mathbb{T}_\theta^n)$ consisting of elements a in the tensor product that are invariant under the diagonal action of \mathbb{T}^n , i.e. such that $\sigma_s \otimes \tau_{-s}(a) = a$ for all $s \in \mathbb{T}^n$. The noncommutative manifold M_θ is defined by ‘duality’ by setting

$$C^\infty(M_\theta) := (C^\infty(M) \overline{\otimes} C^\infty(\mathbb{T}_\theta^n))^{\sigma \otimes \tau^{-1}}.$$

As the notation suggests, the algebra $C^\infty(M_\theta)$ turns out to be isomorphic to the algebra $L_\theta(C^\infty(M))$ defined in the previous section.

Next, let \mathcal{S} be a spin bundle over M and D the Dirac operator on $\Gamma(M, \mathcal{S})$, the $C^\infty(M)$ -module of smooth sections of \mathcal{S} . The action of the group \mathbb{T}^n on M does not lift directly to the spinor bundle. Rather, there is a double cover $c : \tilde{\mathbb{T}}^n \rightarrow \mathbb{T}^n$ and a group homomorphism $\tilde{s} \rightarrow V_{\tilde{s}}$ of $\tilde{\mathbb{T}}^n$ into $\mathrm{Aut}(\mathcal{S})$ covering the action of \mathbb{T}^n on M ,

$$V_{\tilde{s}}(f\psi) = \sigma_{c(s)}(f)V_{\tilde{s}}(\psi), \quad (3.11)$$

for $f \in C^\infty(M)$ and $\psi \in \Gamma(M, \mathcal{S})$. According to [18], the proper notion of smooth sections $\Gamma(M_\theta, \mathcal{S})$ of a spinor bundle on M_θ is given by the subalgebra of $\Gamma(M, \mathcal{S}) \widehat{\otimes} C^\infty(\mathbb{T}_{\theta/2}^n)$ made of elements which are invariant under the diagonal action $V \times \tilde{\tau}^{-1}$ of $\tilde{\mathbb{T}}^n$. Here $\tilde{s} \mapsto \tilde{\tau}_{\tilde{s}}$ is the canonical action of $\tilde{\mathbb{T}}^n$ on $\mathcal{A}(\mathbb{T}_{\theta/2}^n)$. Since the Dirac operator D commutes with $V_{\tilde{s}}$ (remember that the torus action is by isometries) one can restrict $D \otimes \text{id}$ to the fixed point subalgebra $\Gamma(M_\theta, \mathcal{S})$.

Then, let $L^2(M, \mathcal{S})$ be the space of square integrable spinors on M and let $L^2(\mathbb{T}_{\theta/2}^n)$ be the completion of $C^\infty(\mathbb{T}_{\theta/2}^n)$ in the norm $a \mapsto \|a\| = \text{tr}(a^*a)^{1/2}$, with tr the usual trace on $C^\infty(\mathbb{T}_{\theta/2}^n)$. The diagonal action $V \times \tilde{\tau}^{-1}$ of $\tilde{\mathbb{T}}^n$ extends to $L^2(M, \mathcal{S}) \otimes L^2(\mathbb{T}_{\theta/2}^n)$ (where it becomes $U \times \tau$) and one defines $L^2(M_\theta, \mathcal{S})$ to be the fixed point Hilbert subspace. If D also denotes the closure of the Dirac operator on $L^2(M, \mathcal{S})$, one still denotes the operator $D \otimes \text{id}$ on $L^2(M, \mathcal{S}) \otimes L^2(\mathbb{T}_{\theta/2}^n)$ when restricted to $L^2(M_\theta, \mathcal{S})$ by D . The triple $(C^\infty(M_\theta), L^2(M_\theta, \mathcal{S}), D)$ is an m^+ -summable noncommutative spin geometry.

3.3 Vector bundles on M_θ

Noncommutative vector bundles on M_θ , i.e. finite projective modules over $C^\infty(M_\theta)$, were obtained in [18] as fixed point submodules of $\Gamma(M, E) \otimes C^\infty(\mathbb{T}_\theta^n)$ under some diagonal action of the torus \mathbb{T}^n . We will presently give an equivalent description of them in terms of a kind of $*$ -product.

Let E be a σ -equivariant vector bundle M , that is a bundle which carries an action V of \mathbb{T}^n by automorphisms, covering the action σ of \mathbb{T}^n on M ,

$$V_s(f\psi) = \sigma_s(f)V_s(\psi), \quad \forall f \in C^\infty(M), \psi \in \Gamma(M, E). \quad (3.12)$$

The $C^\infty(M_\theta)$ -bimodule $\Gamma(M_\theta, E)$ is defined as the vector space $\Gamma(M, E)$ but with the bimodule structure given by

$$f \triangleright_\theta \psi = \sum_k f_k V_{k,\theta}(\psi), \quad (3.13)$$

$$\psi \triangleleft_\theta f = \sum_k V_{-k,\theta}(\psi) f_k, \quad (3.14)$$

where $f = \sum_k f_k$ with $f_k \in C^\infty(M)$ homogeneous of degree k under the action of \mathbb{T}^n – as in (3.5) – and ψ is a smooth section of E . By using the explicit expression (3.2) for the star product and eq. (3.12), one checks that these are indeed actions of $C^\infty(M_\theta)$.

The $C^\infty(M_\theta)$ -bimodule $\Gamma(M_\theta, E)$ is finite projective [18] and still carries an action of \mathbb{T}^n by V with equivariance as in (3.12) for both the left and right action of $C^\infty(M_\theta)$. Indeed, the group \mathbb{T}^n being abelian, one establishes that

$$V_s(f \triangleright_\theta \psi) = \sigma_s(f) \triangleright_\theta V_s(\psi), \quad \forall f \in C^\infty(M_\theta), \psi \in \Gamma(M_\theta, E), \quad (3.15)$$

and a similar property for the right structure \triangleleft_θ . In fact, due to the fact that the category of σ -equivariant finite projective module over $C^\infty(M)$ is equivalent to the category of σ -equivariant finite projective modules over $C^\infty(M_\theta)$ [27], all finite projective modules on $C^\infty(M_\theta)$ are of the above type. This also reflects the result in [40] that the K-groups of a C^* -algebra deformed by an action of \mathbb{R}^n are isomorphic to the K-groups of the original

C^* -algebra: as mentioned above, the noncommutative manifolds M_θ are a special case – in which the starting algebra is commutative and the action periodic – of the general formulation in [39] of deformations of C^* -algebras under an action of \mathbb{R}^n .

Although we defined the above left and right actions on the sections with respect to an action of \mathbb{T}^n on E , the same construction can be done for vector bundles carrying instead an action of the double cover $\widetilde{\mathbb{T}}^n$. We have already seen an example of this in the case of the spinor bundle, where we defined a left action of $C^\infty(M_\theta)$ in terms of (3.6).

From the very definition of $\Gamma(M_\theta, E)$ the following lemma is true.

Lemma 5. *If $E \simeq F$ as σ -equivariant vector bundles, then $\Gamma(M_\theta, E) \simeq \Gamma(M_\theta, F)$ as $C^\infty(M_\theta)$ -bimodules.*

To proceed, we require a technical condition on the undeformed vector bundles.

Definition 6. *Let E be a σ -equivariant vector bundle on M with σ an action of \mathbb{T}^n . The $C^\infty(M)$ -module $\Gamma(M, E)$ is said to have the homogeneous decomposition property if any section $\phi \in \Gamma(M, E)$ can be decomposed as $\phi = \sum_r \phi_r$ where $\phi_r \in \Gamma(M, E)$ is homogeneous of degree r under the action of \mathbb{T}^n , i.e. $V_s(\phi_r) := e^{2\pi s \cdot r} \phi_r$.*

With this hypothesis we have analogues of ‘classical properties’ for tensor products of $C^\infty(M_\theta)$ -modules.

Lemma 7. *Suppose E and F are σ -equivariant vector bundles on M such that $\Gamma(M, E)$ and $\Gamma(M, F)$ satisfy the homogeneous decomposition property for the actions V^E, V^F of \mathbb{T}^n , respectively. Then $E \otimes F$ is a σ -equivariant vector bundle for the diagonal action of \mathbb{T}^n and we have,*

$$\Gamma(M_\theta, E \otimes F) \simeq \Gamma(M_\theta, E) \otimes_{C^\infty(M_\theta)} \Gamma(M_\theta, F),$$

as both left and right $C^\infty(M_\theta)$ -modules.

Proof. Note first that $\Gamma(M_\theta, E \otimes F) \simeq \Gamma(M, E) \otimes_{C^\infty(M)} \Gamma(M, F)$ as σ -equivariant modules if the action of \mathbb{T}^n on the tensor product $\Gamma(M, E) \otimes_{C^\infty(M)} \Gamma(M, F)$ is defined as the diagonal action $V_t^E \otimes V_t^F$, $t \in \mathbb{T}^n$. Consequently, on the tensor product the left and right action of $C^\infty(M_\theta)$ on $\Gamma(M_\theta, E \otimes F)$ takes the form,

$$\begin{aligned} f \triangleright_\theta (\phi \otimes_{C^\infty(M)} \psi) &= \sum_k f_k V_{k\theta}^E(\phi) \otimes_{C^\infty(M)} V_{k\theta}^F(\psi), \\ (\phi \otimes_{C^\infty(M)} \psi) \triangleleft_\theta f &= \sum_k V_{-k\theta}^E(\phi) \otimes_{C^\infty(M)} V_{-k\theta}^F(\psi) f_k. \end{aligned}$$

We construct an explicit isomorphism of the left and right $C^\infty(M_\theta)$ -modules $\Gamma(M_\theta, E \otimes F)$ and $\Gamma(M_\theta, E) \otimes_{C^\infty(M_\theta)} \Gamma(M_\theta, F)$. First the left module structure: since as vector spaces $\Gamma(M_\theta, E) = \Gamma(M, E)$, it makes sense to define a linear map

$$T : \Gamma(M_\theta, E) \otimes \Gamma(M_\theta, F) \rightarrow \Gamma(M, E) \otimes \Gamma(M, F),$$

first on homogeneous sections ϕ_r and ψ_s of degree r and s respectively – i.e. such that $V_t^E(\phi_r) = e^{2\pi i t \cdot r} \phi_r$ and $V_t^F(\psi_s) = e^{2\pi i t \cdot s} \psi_s$ for $t \in \mathbb{T}^n$ – by

$$T : \phi_r \otimes \psi_s \mapsto \phi_r \otimes V_{r\theta}^F(\psi_s) \tag{3.16}$$

This is then extended linearly on generic elements $\phi = \sum_r \phi_r$ and $\psi = \sum_s \psi_s$. Clearly, the above map T is an isomorphism of vector spaces. It also maps isomorphically the ideal $I_\theta = \{\phi \triangleleft_\theta f \otimes \psi - \phi \otimes f \triangleright_\theta \psi\}$ to the ideal $I_0 = \{f \cdot \phi \otimes \psi - \phi \otimes f \cdot \psi\}$, where \cdot denotes pointwise multiplication. Indeed, for $f \in C^\infty(M_\theta)$ homogeneous of degree k , we find that

$$T(\phi_r \triangleleft_\theta f \otimes \psi_s - \phi \otimes f \triangleright_\theta \psi) = e^{2\pi(k+r)\theta(k+s)}(f \cdot \phi \otimes \psi - \phi \otimes f \cdot \psi). \quad (3.17)$$

Hence T becomes a *vector space* isomorphism between $\Gamma(M_\theta, E) \otimes_{C^\infty(M_\theta)} \Gamma(M_\theta, F)$ and $\Gamma(M, E) \otimes_{C^\infty(M)} \Gamma(M, F)$. From the very definition of T , one checks that it is in fact a left $C^\infty(M_\theta)$ -module map, i.e.

$$T((f \triangleright_\theta \phi) \otimes_{C^\infty(M_\theta)} \psi) = f \triangleright_\theta T(\phi \otimes_{C^\infty(M_\theta)} \psi). \quad (3.18)$$

Similarly, there is an isomorphism of right $C^\infty(M)$ -modules

$$T' : \Gamma(M_\theta, E) \otimes_{C^\infty(M_\theta)} \Gamma(M_\theta, F) \rightarrow \Gamma(M, E) \otimes_{C^\infty(M)} \Gamma(M, F), \quad (3.19)$$

defined on homogeneous elements by

$$T' : \phi_r \otimes_{C^\infty(M_\theta)} \psi_s \mapsto V_{-s\theta}^E \phi_r \otimes_{C^\infty(M)} \psi_s. \quad (3.20)$$

□

Corollary 8. *Let E and F be vector bundles on M such that $\Gamma(M, E)$ and $\Gamma(M, F)$ satisfy the homogeneous decomposition property. Then there is an isomorphism,*

$$\Gamma(M_\theta, E) \otimes_{C^\infty(M_\theta)} \Gamma(M_\theta, F) \simeq \Gamma(M_\theta, F) \otimes_{C^\infty(M_\theta)} \Gamma(M_\theta, E),$$

of both left and right $C^\infty(M_\theta)$ modules.

Proof. Classically, such an isomorphism is given by the tensor flip. Since the action of \mathbb{T}^n on $E \otimes F$ is diagonal, the tensor flip commutes with the action of \mathbb{T}^n and extends to the deformed bundles. □

3.4 Differential calculus on M_θ

It is straightforward to construct a differential calculus on M_θ . This can be done in two equivalent manners, either by extending to forms the quantization maps, or by using the general construction in [15] by means of the Dirac operator.

Firstly, let $(\Omega(M), d)$ be the usual differential calculus on M , with d the exterior derivative. The quantization map $L_\theta : C^\infty(M) \rightarrow C^\infty(M_\theta)$ is extended to $\Omega(M)$ by imposing that it commutes with d . The image $L_\theta(\Omega(M))$ will be denoted $\Omega(M_\theta)$. Equivalently, $\Omega(M_\theta)$ could be defined to be $\Omega(M)$ as a vector space but equipped with an “exterior star product” which is the extension of the product (3.2) to $\Omega(M)$ by the requirement that it commutes with d . Indeed, since the action of \mathbb{T}^n commutes with d , an element in $\Omega(M)$ can be decomposed into a sum of homogeneous elements for the action of \mathbb{T}^n – as was done for $C^\infty(M)$. Then one defines a star product \times_θ on homogeneous elements in $\Omega(M)$ as in (3.2) and denotes $\Omega(M_\theta) = (\Omega(M), \times_\theta)$. This construction is in

concordance with the previous section, when $\Omega(M)$ is considered as the $C^\infty(M)$ -bimodule of sections of the cotangent bundle. The extended action of \mathbb{T}^n from $C^\infty(M)$ to $\Omega(M)$ is used to endow the space $\Omega(M_\theta)$ with the structure of a $C^\infty(M_\theta)$ -bimodule with the left and right action given by (3.13) and (3.14).

A differential calculus $\Omega_D(C^\infty(M_\theta))$ on $C^\infty(M_\theta)$ can also be obtained from a general procedure [15] by means of the isospectral Dirac operator D defined above. The $C^\infty(M_\theta)$ -bimodule $\Omega_D^p(C^\infty(M_\theta))$ of p -forms is made of classes of operators of the form

$$\omega = \sum_j a_0^j [D, a_1^j] \cdots [D, a_p^j], \quad a_i^j \in C^\infty(M_\theta), \quad (3.21)$$

modulo the sub-bimodule of operators

$$\left\{ \sum_j [D, b_0^j] [D, b_1^j] \cdots [D, b_{p-1}^j] : b_i^j \in C^\infty(M_\theta), b_0^j [D, b_1^j] \cdots [D, b_{p-1}^j] = 0. \right\} \quad (3.22)$$

The exterior differential d_D is given by

$$d_D \left[\sum_j a_0^j [D, a_1^j] \cdots [D, a_p^j] \right] = \left[\sum_j [D, a_0^j] [D, a_1^j] \cdots [D, a_p^j] \right], \quad (3.23)$$

and satisfies $d_D^2 = 0$. One also introduces an inner product on forms by declaring that forms of different degree are orthogonal, while for two p -forms ω_1, ω_2 , the inner product is given by

$$(\omega_1, \omega_2)_D = \int \omega_1^* \omega_2. \quad (3.24)$$

Here the noncommutative integral is the natural extension of the one in (3.8),

$$\int T := \text{Tr}_\omega(T|D|^{-m}), \quad (3.25)$$

with T an element in a suitable class of operators. Not surprisingly, these two constructions of forms agree [18], that is, the differential calculi $\Omega(M_\theta)$ and $\Omega_D(C^\infty(M_\theta))$ are isomorphic. This allows us in particular to integrate forms of top dimension, by defining

$$\int_{M_\theta} \omega := \int \omega_D, \quad \omega \in \Omega(M_\theta), \quad (3.26)$$

where ω_D denotes the element in $\Omega_D(C^\infty(M_\theta))$ corresponding to ω (replacing every d in ω by d_D). We have the following noncommutative Stokes theorem.

Lemma 9. *If $\omega \in \Omega^{\dim M - 1}(M_\theta)$ then*

$$\int_{M_\theta} d\omega = 0$$

Proof. By definition of the noncommutative integral,

$$\int_{M_\theta} d\omega = \int d_D \omega_D = \int d_D L_\theta(\omega_D^{(0)})$$

with $\omega_D^{(0)}$ the classical analogue of ω , i.e. $\omega = L_\theta(\omega_D^{(0)})$. At this point one remembers that D commutes with L_θ (see Section 3.1), and realizes that there is an analogue of Lemma 4 for forms, i.e. $\int L_\theta(T) = \int_M T$. One concludes that the above integral vanishes since it vanishes in the classical case. \square

The next ingredient is a Hodge star operator on $\Omega(M_\theta)$. Classically, the Hodge star operator is a map $*$: $\Omega^p(M) \rightarrow \Omega^{m-p}(M)$ depending only on the conformal class of the metric on M . On the one end, since \mathbb{T}^n acts by isometries, it leaves the conformal structure invariant and therefore, it commutes with $*$. On the other hand, with the isospectral deformation one does not change the metric. Thus it makes sense to define a map $*_\theta : \Omega^p(M_\theta) \rightarrow \Omega^{m-p}(M_\theta)$ by

$$*_\theta L_\theta(\omega) = L_\theta(*\omega), \quad \text{for } \omega \in \Omega(M_\theta). \quad (3.27)$$

With this Hodge operator, there is an alternative definition of an inner product on $\Omega(M_\theta)$. Given that $*_\theta$ maps $\Omega^p(M_\theta)$ to $\Omega^{m-p}(M_\theta)$, we can define for $\alpha, \beta \in \Omega^p(M_\theta)$

$$(\alpha, \beta)_2 = \int *_\theta(\alpha^* *_\theta \beta), \quad (3.28)$$

since $*_\theta(\alpha^* *_\theta \beta)$ is an element in $C^\infty(M_\theta)$.

Lemma 10. *Under the isomorphism $\Omega_D(C^\infty(M_\theta)) \simeq \Omega(M_\theta)$, the inner product $(\cdot, \cdot)_2$ coincides with $(\cdot, \cdot)_D$.*

Proof. Let ω_1, ω_2 be two forms in $\Omega_D(C^\infty(M))$, so that $L_\theta(\omega_i)$ are two generic forms in $\Omega_D(C^\infty(M_\theta)) \simeq L_\theta(\Omega_D(C^\infty(M)))$. Then, using Lemma 4 it follows that

$$\int L_\theta(\omega_1)^* L_\theta(\omega_2) = \int L_\theta(\omega_1^* \times_\theta \omega_2) = \int \omega_1^* \times_\theta \omega_2, \quad (3.29)$$

Now, the inner product $(\cdot, \cdot)_D$ coincides with $(\cdot, \cdot)_2$ as defined by (3.28) in the classical case – under the above isomorphism $\Omega_D(C^\infty(M)) \simeq \Omega(M)$; see for example [15, VI.1]. It follows that the above expression equals

$$\int *(\omega_1^* \times_\theta (*\omega_2)) = \int *_\theta(L_\theta(\omega_1)^* (*_\theta L_\theta(\omega_2))), \quad (3.30)$$

using Lemma 4 for forms once more, together with the defining property of $*_\theta$. \square

Lemma 11. *The formal adjoint d^* of d with respect to the inner product $(\cdot, \cdot)_2$ – i.e. so that $(d^*\alpha, \beta)_2 = (\alpha, d\beta)_2$, – is given on $\Omega^p(M_\theta)$ by*

$$d^* = (-1)^{m(p+1)+1} *_\theta d *_\theta$$

Proof. Just as in the classical case, this follows from Stokes lemma 9, together with the observation that

$$\int_{M_\theta} \omega = \int *_\theta \omega, \quad \omega \in \Omega^m(M_\theta),$$

which follows again from the classical case using the mentioned analogue of Lemma 4 for forms. \square

Remark 12. *The algebra $\Omega(M_\theta)$ can also be defined as a fixed point algebra [18]. The action σ of \mathbb{T}^n on $\Omega(M)$ allows one to define $\Omega(M_\theta)$ by $(\Omega(M) \overline{\otimes} C^\infty(\mathbb{T}_\theta^n))^{\sigma \otimes \tau^{-1}}$. Furthermore, since the exterior derivative d on $\Omega(M)$ commutes with the action of \mathbb{T}^n , the differential d_θ , for the fixed point algebra can be defined as $d_\theta = d \otimes \text{id}$. Similarly, the Hodge star operator takes the form $*_\theta = * \otimes \text{id}$ with $*$ the classical Hodge operator.*

4 Gauge theory on the sphere S_θ^4

We apply the general theory of noncommutative gauge field theories – as developed in Section 2 – to the case of the $SU(2)$ noncommutative principal bundle $S_{\theta'}^7 \rightarrow S_\theta^4$ constructed in [31]. This will make more explicit all the constructions above. It is worth stressing that the following holds for any θ -deformed G -principal bundle. We will come back to this point later.

4.1 The principal fibration $S_{\theta'}^7 \rightarrow S_\theta^4$

The $SU(2)$ noncommutative principal fibration $S_{\theta'}^7 \rightarrow S_\theta^4$ is given by an algebra inclusion $\mathcal{A}(S_\theta^4) \hookrightarrow \mathcal{A}(S_{\theta'}^7)$. The algebra $\mathcal{A}(S_\theta^4)$ of polynomial functions on the sphere S_θ^4 is generated by elements $z_0 = z_0^*$, z_j, z_j^* , $j = 1, 2$, subject to relations

$$z_\mu z_\nu = \lambda_{\mu\nu} z_\nu z_\mu, \quad z_\mu z_\nu^* = \lambda_{\nu\mu} z_\nu z_\mu^*, \quad z_\mu^* z_\nu^* = \lambda_{\mu\nu} z_\nu^* z_\mu^*, \quad \mu, \nu = 0, 1, 2, \quad (4.1)$$

together with the spherical relation $\sum_\mu z_\mu^* z_\mu = 1$. Here θ is a real parameter and the deformation parameters are given by

$$\lambda_{12} = \bar{\lambda}_{21} =: \lambda = e^{2\pi i \theta}, \quad \lambda_{j0} = \lambda_{0j} = 1, \quad j = 1, 2. \quad (4.2)$$

For $\theta = 0$ one recovers the $*$ -algebra of complex polynomial functions on the usual S^4 .

The differential calculus $\Omega(S_\theta^4)$ is generated as a graded differential $*$ -algebra by the elements z_μ, z_μ^* in degree 0 and elements dz_μ, dz_μ^* in degree 1 satisfying the relations,

$$\begin{aligned} dz^\mu dz^\nu + \lambda^{\mu\nu} dz^\nu dz^\mu &= 0, & d\bar{z}^\mu dz^\nu + \lambda^{\nu\mu} dz^\nu d\bar{z}^\mu &= 0; \\ z^\mu dz^\nu &= \lambda^{\mu\nu} dz^\nu z^\mu, & \bar{z}^\mu dz^\nu &= \lambda^{\nu\mu} dz^\nu \bar{z}^\mu. \end{aligned} \quad (4.3)$$

with $\lambda^{\mu\nu}$ as before. There is a unique differential d on $\Omega(S_\theta^4)$ such that $d : z_\mu \mapsto dz_\mu$ and the involution on $\Omega(S_\theta^4)$ is the graded extension of $z_\mu \mapsto z_\mu^*$: $(d\omega)^* = d\omega^*$ and $(\omega_1 \omega_2)^* = (-1)^{d_1 d_2} \omega_2^* \omega_1^*$ for ω_j a form of degree d_j .

With $\lambda'_{ab} = e^{2\pi i \theta'_{ab}}$ and (θ'_{ab}) a real antisymmetric matrix, the algebra $\mathcal{A}(S_{\theta'}^7)$ of polynomial functions on the sphere $S_{\theta'}^7$ is generated by elements ψ_a, ψ_a^* , $a = 1, \dots, 4$, subject to relations

$$\psi_a \psi_b = \lambda'_{ab} \psi_b \psi_a, \quad \psi_a \psi_b^* = \lambda'_{ba} \psi_b^* \psi_a, \quad \psi_a^* \psi_a^* = \lambda'_{ab} \psi_b^* \psi_a^*, \quad (4.4)$$

and with the spherical relation $\sum_a \psi_a^* \psi_a = 1$. At $\theta = 0$ it is the $*$ -algebra of complex polynomial functions on the sphere S^7 . As before, a differential calculus $\Omega(S_{\theta'}^7)$ can be defined to be generated by the elements ψ_a, ψ_a^* in degree 0 and elements $d\psi_a, d\psi_a^*$ in degree 1 satisfying relations similar to the one in (4.3).

In order to construct the noncommutative fibration over the given 4-sphere S_θ^4 we need to select a particular noncommutative 7 dimensional sphere $S_{\theta'}^7$. We take the one corresponding to the following deformation parameters

$$\lambda'_{ab} = \begin{pmatrix} 1 & 1 & \bar{\mu} & \mu \\ 1 & 1 & \mu & \bar{\mu} \\ \mu & \bar{\mu} & 1 & 1 \\ \bar{\mu} & \mu & 1 & 1 \end{pmatrix}, \quad \mu = \sqrt{\lambda}, \quad \text{or} \quad \theta'_{ab} = \frac{\theta}{2} \begin{pmatrix} 0 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 \\ 1 & -1 & 0 & 0 \\ -1 & 1 & 0 & 0 \end{pmatrix}. \quad (4.5)$$

The previous choice is essentially the only one that allows the algebra $\mathcal{A}(S_{\theta'}^7)$ to carry an action of the group $SU(2)$ by automorphisms and such that the invariant subalgebra coincides with $\mathcal{A}(S_{\theta}^4)$. There are indeed two possibilities corresponding to $\mu = \pm\sqrt{\lambda}$. The best way to see this is by means of the matrix-valued function on $\mathcal{A}(S_{\theta'}^7)$,

$$\Psi = \begin{pmatrix} \psi_1 & -\psi_2^* \\ \psi_2 & \psi_1^* \\ \psi_3 & -\psi_4^* \\ \psi_4 & \psi_3^* \end{pmatrix}. \quad (4.6)$$

Then $\Psi^\dagger\Psi = \mathbb{I}_2$ and $p = \Psi\Psi^\dagger$ is a projection, $p^2 = p = p^\dagger$, with entries in $\mathcal{A}(S_{\theta}^4)$. Indeed, the right action of $SU(2)$ on $\mathcal{A}(S_{\theta'}^7)$ is simply given by

$$\alpha_w(\Psi) = \Psi w, \quad w = \begin{pmatrix} w_1 & -\bar{w}_2 \\ w_2 & \bar{w}_1 \end{pmatrix} \in SU(2), \quad (4.7)$$

from which the invariance of the entries of p follows at once. Explicitly,

$$p = \frac{1}{2} \begin{pmatrix} 1+z_0 & 0 & z_1 & -\bar{\mu}z_2^* \\ 0 & 1+z_0 & z_2 & \mu z_1^* \\ z_1^* & z_2^* & 1-z_0 & 0 \\ -\mu z_2 & \bar{\mu}z_1 & 0 & 1-z_0 \end{pmatrix}, \quad (4.8)$$

with the generators of $\mathcal{A}(S_{\theta}^4)$ given by

$$\begin{aligned} z_0 &= \psi_1^*\psi_1 + \psi_2^*\psi_2 - \psi_3^*\psi_3 - \psi_4^*\psi_4 \\ &= 2(\psi_1^*\psi_1 + \psi_2^*\psi_2) - 1 = 1 - 2(\psi_3^*\psi_3 + \psi_4^*\psi_4), \\ z_1 &= 2(\mu\psi_3^*\psi_1 + \psi_2^*\psi_4) = 2(\psi_1\psi_3^* + \psi_2^*\psi_4), \\ z_2 &= 2(-\psi_1^*\psi_4 + \bar{\mu}\psi_3^*\psi_2) = 2(-\psi_1^*\psi_4 + \psi_2\psi_3^*). \end{aligned} \quad (4.9)$$

One straightforwardly computes that $z_1^*z_1 + z_2^*z_2 + z_0^2 = 1$ and the commutation rules $z_1z_2 = \lambda z_2z_1$, $z_1z_2^* = \bar{\lambda}z_2^*z_1$, and that z_0 is central.

The relations (4.9) can also be expressed in the form,

$$z_\mu = \sum_{ab} \psi_a^*(\gamma_\mu)_{ab}\psi_b, \quad z_\mu^* = \sum_{ab} \psi_a^*(\gamma_\mu^*)_{ab}\psi_b, \quad (4.10)$$

with γ_μ twisted 4×4 Dirac matrices given by

$$\gamma_0 = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & -1 & \\ & & & -1 \end{pmatrix}, \quad \gamma_1 = 2 \begin{pmatrix} 0 & 0 & 0 & \\ \mu & 0 & 0 & \\ 0 & 0 & 0 & \\ 0 & 0 & 0 & \end{pmatrix}, \quad \gamma_2 = 2 \begin{pmatrix} 0 & 0 & -1 & \\ 0 & \bar{\mu} & 0 & \\ 0 & 0 & 0 & \\ 0 & 0 & 0 & \end{pmatrix}. \quad (4.11)$$

Note that as usual γ_0 is the grading

$$\gamma_0 = -\frac{1}{4}[\gamma_1, \gamma_1^*][\gamma_2, \gamma_2^*].$$

These matrices satisfy twisted Clifford algebra relations [18],

$$\gamma_\mu\gamma_\nu + \lambda_{\mu\nu}\gamma_\nu\gamma_\mu = 0, \quad \gamma_\mu\gamma_\nu^* + \lambda_{\nu\mu}\gamma_\nu^*\gamma_\mu = 4\delta_{\mu\nu}; \quad (\mu, \nu = 1, 2). \quad (4.12)$$

There are compatible toric actions on S_θ^4 and $S_{\theta'}^7$. The torus \mathbb{T}^2 acts on $\mathcal{A}(S_\theta^4)$ as

$$\sigma_s(z_0, z_1, z_2) = (z_0, e^{2\pi i s_1} z_1, e^{2\pi i s_2} z_2), \quad s \in \mathbb{T}^2. \quad (4.13)$$

This action is lifted to a double cover action on $\mathcal{A}(S_{\theta'}^7)$. The double cover map $p : \widetilde{\mathbb{T}}^2 \rightarrow \mathbb{T}^2$ is given explicitly by $p : (s_1, s_2) \mapsto (s_1 + s_2, -s_1 + s_2)$. Then $\widetilde{\mathbb{T}}^2$ acts on the ψ_a 's as:

$$\tilde{\sigma} : (\psi_1, \psi_2, \psi_3, \psi_4) \mapsto (e^{2\pi i s_1} \psi_1, e^{-2\pi i s_1} \psi_2, e^{-2\pi i s_2} \psi_3, e^{2\pi i s_2} \psi_4) \quad (4.14)$$

Equation (4.9) shows that $\tilde{\sigma}$ is indeed a lifting to $S_{\theta'}^7$ of the action of \mathbb{T}^2 on S_θ^4 . Clearly, this compatibility is built in the construction of the Hopf fibration $S_{\theta'}^7 \rightarrow S_\theta^4$ as a deformation of the classical Hopf fibration $S^7 \rightarrow S^4$ with respect to an action of \mathbb{T}^2 , a fact that also dictated the form of the deformation parameter λ' in (4.5). As we shall see, the previous double cover of tori comes from a spin cover $\text{Spin}_\theta(5)$ of $\text{SO}_\theta(5)$ deforming the usual action of $\text{Spin}(5)$ on S^7 as a double cover of the action of $\text{SO}(5)$ on S^4 .

In which sense the algebra inclusion $\mathcal{A}(S_\theta^4) \hookrightarrow \mathcal{A}(S_{\theta'}^7)$ is a noncommutative principal bundle was explained in [31] to which we refer for more details. Presently we shall recall the construction of associated bundles.

4.2 Associated bundles

We shall work in the context of smooth functions on $S_{\theta'}^7$ and S_θ^4 as defined in general in Section 3. Let ρ be any representation of $\text{SU}(2)$ on an n -dimensional vector space V . The corresponding equivariant maps are defined as

$$\mathcal{E} := C^\infty(S_{\theta'}^7) \boxtimes_\rho V := \{ \eta \in C^\infty(S_{\theta'}^7) \otimes V : (\alpha_w \otimes \text{id})(\eta) = (\text{id} \otimes \rho(w)^{-1})(\eta) \}, \quad (4.15)$$

where α_w is the $\text{SU}(2)$ action given in (4.7). The previous is clearly a $C^\infty(S_\theta^4)$ bimodule. We have proved in [31] that \mathcal{E} is also a finite projective module. It is worth stressing that the choice of a projection for a finite projective module requires the choice of one of the two (left or right) module structures. Similarly, the definition of a Hermitian structure requires the choice of the left or right module structure. In the following, we will always work with the right structure for the associated modules. There is a natural (right) Hermitian structure on \mathcal{E} , defined in terms of the inner product of V as:

$$\langle \eta, \eta' \rangle := \sum_i \bar{\eta}_i \eta'_i. \quad (4.16)$$

where we denoted $\eta = \sum_i \eta_i \otimes e_i, \eta' = \sum_i \eta'_i \otimes e_i$ for a basis $\{e_i\}_{i=1}^n$ in V . One quickly checks that $\langle \eta, \eta' \rangle$ is an element in $C^\infty(S_\theta^4)$, and that $\langle \cdot, \cdot \rangle$ satisfies all conditions of a right Hermitian structure.

The bimodules $C^\infty(S_{\theta'}^7) \boxtimes_\rho V$ are of the type described in Section 3.3. Indeed, the associated vector bundle $E = S^7 \times_\rho V$ on S^4 carries an action of $\widetilde{\mathbb{T}}^2$ induced from its action on S^7 , which is obviously σ -equivariant. By the very definition of $C^\infty(S_{\theta'}^7)$ and of $\Gamma(S_\theta^4, E)$ Section 3.3, it follows that $C^\infty(S_{\theta'}^7) \boxtimes_\rho V \simeq \Gamma(S_\theta^4, E)$. Indeed, from the undeformed isomorphism, $\Gamma(S^4, E) \simeq C^\infty(S^7) \boxtimes_\rho V$, the quantization map $L_{\theta'}$ of $C^\infty(S^7)$, acting only on the first leg of the tensor product, establishes this isomorphism,

$$L_{\theta'} : C^\infty(S^7) \boxtimes_\rho V \rightarrow C^\infty(S_{\theta'}^7) \boxtimes_\rho V. \quad (4.17)$$

The above is well defined since the action of $\widetilde{\mathbb{T}}^2$ commutes with the action of $SU(2)$. Also, it is such that $L_{\theta'}(f\eta) = L_{\theta'}(f) \triangleright_{\theta'} L_{\theta'}(\eta) = L_{\theta}(f) \triangleright_{\theta'} L_{\theta'}(\eta)$ for $f \in C^\infty(S^4)$ and $\eta \in C^\infty(S^7) \boxtimes_{\rho} V$ – due to the identity $L_{\theta'} = L_{\theta}$ on $C^\infty(S^4) \subset C^\infty(S^7)$. A similar result holds for the action $\triangleleft_{\theta'}$.

Remark 13. *The $C^\infty(S^4)$ -modules $\Gamma(S^4, E)$ have the homogeneous decomposition property. For instance, if $V = \mathbb{C}^2$ a basis $\{e_i\}_{i=1}^4$ of the module $\Gamma(S^4, E)$ is given in terms of the (classical counterpart of the) matrix Ψ of the matrix (4.6) by*

$$e_1 := \begin{pmatrix} \psi_1^* \\ -\psi_2 \end{pmatrix}, \quad e_2 := \begin{pmatrix} \psi_2^* \\ \psi_1 \end{pmatrix}, \quad e_3 := \begin{pmatrix} \psi_3^* \\ -\psi_4 \end{pmatrix}, \quad e_4 := \begin{pmatrix} \psi_4^* \\ \psi_3 \end{pmatrix}. \quad (4.18)$$

These are homogeneous elements under the action (4.14) of $\widetilde{\mathbb{T}}^2$ on S^7 . A generic element $\eta = \sum_i e_i f_i$ in $\Gamma(S^4, E)$ with $f_i \in C^\infty(S^4)$ can then clearly be decomposed into homogeneous elements – as every component f_i can. More generally, for $V^{(n)} = \mathbb{C}^{n+1}$, the homogeneous decomposition property follows from the existence of similar (but larger) matrices $\Psi_{(n)}$, constructed in [31]. Any element in the module $\Gamma(S^4, E)$ can be written in a module basis $\{e_i\}_{i=1}^{4^n}$ – homogeneous under the action of $\widetilde{\mathbb{T}}^2$ – given by the columns of the $(n+1) \times 4^n$ -matrix $\Psi_{(n)}^*$.

Given the right $C^\infty(S_\theta^4)$ -module \mathcal{E} , its dual module of \mathcal{E} is defined by

$$\mathcal{E}' := \{\phi : \mathcal{E} \rightarrow C^\infty(S_\theta^4) : \phi(\eta f) = \phi(\eta) f, f \in C^\infty(S_\theta^4)\}, \quad (4.19)$$

and is naturally a left $C^\infty(S_\theta^4)$ -module. If $\mathcal{E} := C^\infty(S_{\theta'}^7) \boxtimes_{\rho} V$ comes from the $SU(2)$ -representation (V, ρ) , by using the induces dual representation ρ' on the dual vector space V' given by

$$(\rho'(w)v')(v) := v'(\rho(w)^{-1}v); \quad \forall v' \in V', v \in V, \quad (4.20)$$

we have that

$$\mathcal{E}' \simeq C^\infty(S_{\theta'}^7) \boxtimes_{\rho'} V' := \{\phi \in C^\infty(S_{\theta'}^7) \otimes V' : (\alpha_w \otimes \text{id})(\phi) = (\text{id} \otimes \rho'(w)^{-1})(\phi)\}. \quad (4.21)$$

Next, let $L(V)$ denote the space of linear maps on V , so that $L(V) = V \otimes V'$. The adjoint action of $SU(2)$ on $L(V)$ is the tensor product representation $\text{ad} := \rho \otimes \rho'$ on $V \otimes V'$. We define

$$C^\infty(S_{\theta'}^7) \boxtimes_{\text{ad}} L(V) := \{T \in C^\infty(S_{\theta'}^7) \otimes L(V) : (\alpha_w \otimes \text{id})(T) = (\text{id} \otimes \text{ad}(w)^{-1})(T)\}, \quad (4.22)$$

and write $T = T_{ij} \otimes e_{ij}$ with respect to the basis $\{e_{ij}\}$ of $L(V)$ induced from the basis $\{e_i\}_{i=1}^n$ of V and the dual one $\{e'_j\}_{j=1}^n$ of V' .

On the other hand, there is the endomorphism algebra

$$\text{End}_{C^\infty(S_\theta^4)}(\mathcal{E}) := \{T : \mathcal{E} \rightarrow \mathcal{E} : T(\eta f) = T(\eta) f, f \in C^\infty(S_\theta^4)\}. \quad (4.23)$$

We will suppress the subscript $C^\infty(S_\theta^4)$ from End in the following.

Proposition 14. *There is an isomorphism of algebras*

$$\text{End}(\mathcal{E}) \simeq C^\infty(S_{\theta'}^7) \boxtimes_{\text{ad}} L(V).$$

Proof. Recall that $\mathcal{E} \otimes_{C^\infty(S_\theta^4)} \mathcal{E}' \subset \text{End}(\mathcal{E})$ densely (in the operator norm, cf. for instance [26]). We define a map from $\text{End}(\mathcal{E})$ to $C^\infty(S_{\theta'}^7) \boxtimes_{\text{ad}} L(V)$ on this dense subalgebra by

$$\eta \otimes \phi \mapsto \eta_i \phi_j \otimes e_{ij},$$

with $\eta = \eta_i \otimes e_i$ and $\phi = \phi_i \otimes e'_i$. On the other hand, $T \in C^\infty(S_{\theta'}^7) \boxtimes_{\text{ad}} L(V)$ acts on $\eta \in \mathcal{E}$ by

$$(T, \eta) \mapsto (T_{ij} \eta_j) \otimes e_i,$$

which is clearly a right $C^\infty(S_\theta^4)$ -linear map with image in \mathcal{E} . Hence, the opposite inclusion $C^\infty(S_{\theta'}^7) \boxtimes_{\text{ad}} L(V) \subset \text{End}(\mathcal{E})$. \square

We see that the algebra of endomorphisms of \mathcal{E} can be understood as the space of sections of the noncommutative vector bundle associated to the adjoint representation on $L(V)$ – exactly as it happens in the classical case. This also allows an identification of skew-hermitian endomorphisms $\text{End}^s(\mathcal{E})$ – which were defined in general in (2.18) – for the toric deformations at hand.

Corollary 15. *There is an identification*

$$\text{End}^s(\mathcal{E}) \simeq C_{\mathbb{R}}^\infty(S_{\theta'}^7) \boxtimes_{\text{ad}} u(n),$$

with $C_{\mathbb{R}}^\infty(S_{\theta'}^7)$ denoting the algebra of self-adjoint elements in $C^\infty(S_{\theta'}^7)$ whereas $u(n)$ consists of skew-adjoint matrices in $M_n(\mathbb{C}) \simeq L(V)$.

Proof. Note that the involution $T \mapsto T^*$ in $\text{End}(\mathcal{E})$ reads in components $T_{ij} \mapsto \overline{T_{ji}}$ so that, with the identification of Proposition 14, the algebra $\text{End}^s(\mathcal{E})$ is made of elements $X \in C^\infty(S_{\theta'}^7) \boxtimes_{\text{ad}} L(V)$ satisfying $\overline{X_{ji}} = -X_{ij}$. Since any element in $C^\infty(S_{\theta'}^7)$ can be written as the sum of two self-adjoint elements, $X_{ij} = X_{ij}^{\Re} + iX_{ij}^{\Im}$, we can write

$$X = \sum_i X_{ii}^{\Im} \otimes ie_{ii} + \sum_{i \neq j} X_{ij}^{\Re} \otimes (e_{ij} - e_{ji}) + X_{ij}^{\Im} \otimes (ie_{ij} + ie_{ji}) = \sum_a X_a \otimes \sigma^a,$$

where X_a are generic elements in $C_{\mathbb{R}}^\infty(S_{\theta'}^7)$ and $\{\sigma^a, a = 1, \dots, n^2\}$ are the generators of $u(n)$. \square

Example 16. *Of central interest in the following is the special case of the noncommutative instanton bundle first constructed in [19]. Now $V = \mathbb{C}^2$ and ρ is the defining representation of $\text{SU}(2)$. The projection p giving the $C^\infty(S_\theta^4)$ -module $C^\infty(S_{\theta'}^7) \boxtimes_{\rho} \mathbb{C}^2$ as a direct summand of $(C^\infty(S_\theta^4))^N$ for some N , is precisely given by the one in (4.8) and $N = 4$. Indeed, a generic element in $C^\infty(S_{\theta'}^7) \boxtimes_{\rho} \mathbb{C}^2$ is of the form $\Psi^\dagger f$ for some $f \in C^\infty(S_\theta^4) \otimes \mathbb{C}^4$ with Ψ defined in (4.6), and the correspondence is given by*

$$C^\infty(S_{\theta'}^7) \boxtimes_{\rho} \mathbb{C}^2 \simeq p(C^\infty(S_\theta^4))^4, \quad \Psi^\dagger f \leftrightarrow pf. \quad (4.24)$$

Furthermore, $\text{End}(\mathcal{E}) \simeq C^\infty(S_{\theta'}^7) \boxtimes_{\text{ad}} M_2(\mathbb{C})$. It is a known fact that $M_2(\mathbb{C})$ decomposes into the adjoint representation $\mathfrak{su}(2)$ and the trivial representation \mathbb{C} while it is easy to see that $C^\infty(S_{\theta'}^7) \boxtimes_{\text{id}} \mathbb{C} \simeq C^\infty(S_\theta^4)$. Thus, we conclude that

$$\text{End}(\mathcal{E}) \simeq \Gamma(\text{ad}(S_{\theta'}^7)) \oplus C^\infty(S_\theta^4), \quad (4.25)$$

where we have set $\Gamma(\text{ad}(S_{\theta'}^7)) := C^\infty(S_{\theta'}^7) \boxtimes_{\text{ad}} \mathfrak{su}(2)$. The latter $C^\infty(S_\theta^4)$ -bimodule will be understood as the space of (complex) sections of the adjoint bundle. It is the complexification of the traceless skew-hermitian endomorphisms $C_{\mathbb{R}}^\infty(S_{\theta'}^7) \boxtimes_{\text{ad}} \mathfrak{su}(2)$.

4.3 Yang-Mills theory on S_θ^4

Let us now move on to the main goal of this paper and discuss the Yang-Mills action functional on S_θ^4 together with its equation of motion. We will see that instantons naturally arise as the local minima of this action.

Before we proceed we recall the noncommutative spin structure $(C^\infty(S_\theta^4), \mathcal{H}, D, \gamma_5)$ of S_θ^4 with $\mathcal{H} = L^2(S^4, \mathcal{S})$ the Hilbert space of spinors, D the undeformed Dirac operator, and γ_5 – the even structure – the fifth Dirac matrix.

Let $\mathcal{E} = \Gamma(S_\theta^4, E)$ for some σ -equivariant vector bundle E on S^4 , so that there exists a projection $p \in M_N(C^\infty(S_\theta^4))$ such that $\mathcal{E} \simeq p(C^\infty(S_\theta^4))^N$. Recall from Section 2 that a connection ∇ on $\mathcal{E} = \Gamma(S_\theta^4, E)$ for some vector bundle E on S^4 , is a map from \mathcal{E} to $\mathcal{E} \otimes \Omega(S_\theta^4)$. The Yang-Mills action functional is defined in terms of the curvature of a connection on \mathcal{E} , which is an element in $\text{End}_{C^\infty(S_\theta^4)}(\mathcal{E}, \mathcal{E} \otimes \Omega^2(S_\theta^4))$. Equivalently, it is an element in $\text{End}_{\Omega(S_\theta^4)}(\mathcal{E} \otimes \Omega(S_\theta^4))$ of degree 2. We define an inner product on the latter algebra as follows [15, III.3]. An element $T \in \text{End}_{\Omega(S_\theta^4)}(\mathcal{E} \otimes \Omega(S_\theta^4))$ of degree k can be understood as an element in $pM_N(\Omega^k(S_\theta^4))p$, since $\mathcal{E} \otimes \Omega(S_\theta^4)$ is a finite projective module over $\Omega(S_\theta^4)$. A trace over internal indices together with the inner product defined in (3.28), defines the inner product $(\cdot, \cdot)_2$ on $\text{End}_{\Omega(S_\theta^4)}(\mathcal{E} \otimes \Omega(S_\theta^4))$. In particular, we can make the following definition.

Definition 17. *The Yang-Mills action functional for a connection ∇ on \mathcal{E} with curvature F is defined by*

$$\text{YM}(\nabla) = (F, F)_2 = \int *_\theta \text{tr}(F *_\theta F).$$

Recall from Section 2 that gauge transformations are given by unitary endomorphisms $\mathcal{U}(\mathcal{E})$ of \mathcal{E} .

Lemma 18. *The Yang-Mills action functional is gauge invariant, positive and quartic.*

Proof. From equation (2.23) the curvature F transforms under a gauge transformation $u \in \mathcal{U}(\mathcal{E})$ as $F \mapsto u^* F u$. Since $\mathcal{U}(\mathcal{E})$ can be identified with the unitary elements in $pM_N(\mathcal{A})p$, it follows that

$$\text{YM}(\nabla^u) = \int \sum_{i,j,k,l} *_\theta (\overline{u_{ji}} F_{jk} *_\theta F_{kl} u_{li}) = \text{YM}(\nabla)$$

using the tracial property of the Dixmier trace and the fact that $u_{li} \overline{u_{ji}} = \delta_{lj}$. Positiveness of the Yang-Mills action functional follows from Lemma 10 giving

$$(F, F)_2 = (F_D, F_D)_D = \int F_D^* F_D,$$

which is clearly positive. □

In physics, the Yang-Mills equations are obtained from the Yang-Mills action functional by a variational principle. Let us describe how this principle works in our case. We consider a linear perturbation $\nabla_t = \nabla + t\alpha$ of a connection ∇ on \mathcal{E} by an element $\alpha \in \text{End}(\mathcal{E}, \mathcal{E} \otimes_{C^\infty(S_\theta^4)} \Omega^1(S_\theta^4))$. The curvature F_t of ∇_t is readily computed as $F_t =$

$F + t[\nabla, \alpha] + \mathcal{O}(t^2)$. If we suppose that ∇ is an extremum of the Yang-Mills action functional, this linear perturbation should not affect the action. In other words, we should have

$$\left. \frac{\partial}{\partial t} \right|_{t=0} \text{YM}(\nabla_t) = 0. \quad (4.26)$$

If we substitute the explicit formula for F_t , we obtain

$$([\nabla, \alpha], F)_2 + \overline{([\nabla, \alpha], F)_2} = 0. \quad (4.27)$$

using the fact that $(\cdot, \cdot)_2$ defines a complex scalar product on $\text{End}(\mathcal{E}, \mathcal{E} \otimes \Omega(S_\theta^4))$. Positive definiteness of this scalar product implies that $(F_t, F_t) = \overline{(F_t, F_t)}$, which when differentiated with respect to t , setting $t = 0$ afterwards, yields $([\nabla, \alpha], F)_2 = \overline{([\nabla, \alpha], F)_2}$; hence, $([\nabla, \alpha], F)_2 = 0$. Using the fact that α was arbitrary, we derive the following equations of motion

$$[\nabla^*, F] = 0. \quad (4.28)$$

where the adjoint of $[\nabla, \cdot]$ is defined with respect to the inner product $(\cdot, \cdot)_2$, i.e.

$$([\nabla^*, \alpha], \beta)_2 = (\alpha, [\nabla, \beta])_2 \quad (4.29)$$

for $\alpha \in \text{End}(\mathcal{E}, \mathcal{E} \otimes \Omega^3(S_\theta^4))$ and $\beta \in \text{End}(\mathcal{E}, \mathcal{E} \otimes \Omega^1(S_\theta^4))$. From Lemma 11, it follows that $[\nabla^*, F] = *_\theta[\nabla, *_\theta F]$, so that the equations of motion can also be written as the more familiar *Yang-Mills equations*:

$$[\nabla, *_\theta F] = 0. \quad (4.30)$$

Note that connections with a self-dual or antiself-dual curvature $*_\theta F = \pm F$ are special solutions of the Yang-Mills equation. Indeed, in this case the latter equation follows directly from the Bianchi identity $[\nabla, F] = 0$. We call such connections instantons on S_θ^4 .

We will now establish a connection between the Yang-Mills action functional and the so-called topological action [15, VI.3] on S_θ^4 . Suppose \mathcal{E} is a finite projective module over $C^\infty(S_\theta^4)$ defined by a projection $p \in M_N(C^\infty(S_\theta^4))$. The topological action for \mathcal{E} is given by a pairing between the class of p in K-theory and the cyclic cohomology of $C^\infty(S_\theta^4)$. For computational purposes, we give the following definition in terms of the curvature of a connection on \mathcal{E}

Definition 19. *Let ∇ be a connection on \mathcal{E} with curvature F . The topological action is given by*

$$\text{Top}(\mathcal{E}) = -(F, *_\theta F)_2 = - \int *_\theta \text{tr}(F^2)$$

where the trace is taken over internal indices and in the second equality we have used the identity $*_\theta \circ *_\theta = \text{id}$.

Let us show that this does not depend on the choice of a connection on \mathcal{E} . Since two connections differ by an element α in $\text{End}_{C^\infty(S_\theta^4)}(\mathcal{E}, \mathcal{E} \otimes \Omega^1(S_\theta^4))$, we have to establish that $(F', *_\theta F')_2 = (F, *_\theta F)_2$ where $F' = F + t[\nabla, \alpha] + \mathcal{O}(t^2)$ is the curvature of $\nabla' := \nabla + t\alpha$, $t \in \mathbb{R}$. By definition of the inner product $(\cdot, \cdot)_2$ we then have

$$\begin{aligned} (F', *_\theta F')_2 - (F, *_\theta F)_2 &= t(F, *_\theta[\nabla, \alpha])_2 + t([\nabla, \alpha], *_\theta F)_2 + \mathcal{O}(t^2) \\ &= t(F, [\nabla^*, *_\theta \alpha])_2 + t([\nabla^*, *_\theta \alpha], F)_2 + \mathcal{O}(t^2), \end{aligned}$$

which vanishes due to the Bianchi identity $[\nabla, F] = 0$, Proposition 1.

The Hodge star operator $*_\theta$ splits $\Omega^2(S_\theta^4)$ into a self-dual and antiself-dual space,

$$\Omega^2(S_\theta^4) = \Omega_+^2(S_\theta^4) \oplus \Omega_-^2(S_\theta^4). \quad (4.31)$$

In fact, $\Omega_\pm^2(S_\theta^4) = L_\theta(\Omega_\pm^2(S^4))$. This decomposition is orthogonal with respect to the inner product $(\cdot, \cdot)_2$ which follows from the property $(\alpha, \beta)_2 = \overline{(\beta, \alpha)_2}$, so that we can write the Yang-Mills action functional as

$$\text{YM}(\nabla) = (F_+, F_+)_2 + (F_-, F_-)_2. \quad (4.32)$$

Comparing this with the topological action,

$$\text{Top}(\mathcal{E}) = -(F_+, F_+)_2 + (F_-, F_-)_2, \quad (4.33)$$

we see that $\text{YM}(\nabla) \geq |\text{Top}(\mathcal{E})|$, with equality holding iff

$$*_\theta F = \pm F. \quad (4.34)$$

Solutions of these equations are called instantons. We conclude that instantons are absolute minima of the Yang-Mills action functional.

5 Construction of $\text{SU}(2)$ -instantons on S_θ^4

In this section, we construct a set of charge -1 $\text{SU}(2)$ instantons on S_θ^4 , by acting with a twisted infinitesimal conformal symmetry on the basic instanton on S_θ^4 constructed in [19]. We will find a five-parameter family of such instantons. Then we prove that the ‘tangent space’ of the moduli space of irreducible instantons at the basic instanton is five-dimensional, proving that this set is complete. Here, one has to be careful with the notion of tangent space to the moduli space. As will be discussed elsewhere [30], it turns out that the moduli space is a noncommutative space given as the quantum quotient space of the deformed conformal group $\text{SL}_\theta(2, \mathbb{H})$ by the deformed gauge group $\text{Sp}_\theta(2)$. It turns out that the basic instanton of [19] is a ‘classical point’ in this moduli space of instantons. We perturb this connection ∇_0 linearly by sending $\nabla_0 \mapsto \nabla_0 + t\alpha$ where $t \in \mathbb{R}$ and $\alpha \in \text{End}(\mathcal{E}, \mathcal{E} \otimes \Omega^1(S_\theta^4))$. In order for this new connection still to be an instanton, we have to impose the self-duality equation on its curvature. After deriving this equation with respect to t , setting $t = 0$ afterwards, we obtain the linearized self-duality equation to be fulfilled by α . It is in this sense that we are considering the tangent space to the moduli space of instantons at the origin ∇_0 .

5.1 The basic instanton

Let us start with a technical lemma that simplifies the discussion.

Lemma 20. *There is the following isomorphism of right $C^\infty(S_\theta^4)$ -modules:*

$$\mathcal{E} \otimes_{C^\infty(S_\theta^4)} \Omega(S_\theta^4) \simeq \Omega(S_\theta^4) \otimes_{C^\infty(S_\theta^4)} \mathcal{E}.$$

Consequently, $\text{End}(\mathcal{E}, \mathcal{E} \otimes_{C^\infty(S_\theta^4)} \Omega(S_\theta^4)) \simeq \Omega(S_\theta^4) \otimes_{C^\infty(S_\theta^4)} \text{End}(\mathcal{E})$.

Proof. This follows from Corollary 8 and Remark 13, together with the observation that the $C^\infty(S^4)$ -module $\Omega(S^4)$ has a \mathbb{T}^2 -homogeneous basis. \square

Thus, we can unambiguously use the notation $\Omega(S_\theta^4, \mathcal{E})$ for the above right $C^\infty(S_\theta^4)$ -module.

We let $\nabla_0 = p \circ d$ be the canonical (Grassmann) connection on the projective module $\mathcal{E} = \Gamma(S_{\theta'}^7 \otimes_{SU(2)} \mathbb{C}^2) = p(C^\infty(S_\theta^4))^4$, with the projection $p = \Psi^\dagger \Psi$ of (4.8) and Ψ is the matrix (4.6). In the light of the isomorphism between this module and the collection of equivariant maps $C^\infty(S_{\theta'}^7) \boxtimes_\rho \mathbb{C}^2$ recalled in Example 16, we can write ∇_0 on equivariant maps as

$$\begin{aligned} \nabla_0 : \mathcal{E} &\rightarrow \mathcal{E} \otimes_{C^\infty(S_\theta^4)} \Omega^1(S_\theta^4), \\ (\nabla_0 f)_i &= df_i + \omega_{ij} \times_\theta f_j, \end{aligned} \tag{5.1}$$

where ω – referred to as the gauge potential – is given in terms of the matrix Ψ by

$$\omega = \Psi^\dagger d\Psi. \tag{5.2}$$

The above, is a 2×2 -matrix with entries in $\Omega^1(S_{\theta'}^7)$ satisfying $\overline{\omega_{ij}} = \omega_{ji}$ and $\sum_i \omega_{ii} = 0$. Note here that the entries ω_{ij} commute with all elements in $C^\infty(S_{\theta'}^7)$. Indeed, from (4.6) we see that the elements in ω_{ij} are \mathbb{T}^2 -invariant and hence central (as one forms) in $\Omega(S_{\theta'}^7)$. In other words $L_\theta(\omega) = \omega$, which shows that for an element $f \in \mathcal{E}$ as above, we have $\nabla_0(f)_i = df_i + \omega_{ij} \times_\theta f_j = df_i + \omega_{ij} f_j$ which coincides with the action of the classical connection $d + \omega$ on f .

The curvature $F_0 = \nabla_0^2 = d\omega + \omega^2$ of ∇_0 is an element of $\text{End}(\mathcal{E}) \otimes_{C^\infty(S_\theta^4)} \Omega^2(S_\theta^4)$ that satisfies [2, 18] the self-duality equation

$$*_\theta F_0 = F_0, \tag{5.3}$$

hence this connection is an instanton. At the classical value of the deformation parameter, $\theta = 0$, the connection (5.2) is nothing but the $SU(2)$ instanton of [9].

Its ‘topological charge’, i.e. the values of $\text{Top}(\mathcal{E})$ in Definition 19, was already computed in [19] (with the change in orientation mentioned in the introduction). Clearly it depends only on the class $[p]$ of the bundle and can be evaluated as the index

$$\text{Top}([p]) = \text{index}(D_p) = \int \gamma_5 \pi_D(\text{ch}_2(p)) \tag{5.4}$$

of the twisted Dirac operator

$$D_p = p(D \otimes \mathbb{I}_4)p.$$

The last equality in (5.4) follows from the vanishing of the class $\text{ch}_1(p)$ of the bundle¹. On the hand, one finds

$$\pi_D(\text{ch}_2(p)) = -3\gamma_5,$$

which, together with the fact that

$$\int 1 = \text{Tr}_\omega |D|^{-4} = \frac{1}{3},$$

¹The Chern character classes and their realization as operators are in the appendix

on S^4 (see for instance [26, 29]), gives the value $\text{Top}([p]) = -1$.

We aim at constructing all connections ∇ on \mathcal{E} whose curvature satisfies this self-duality equation and of topological charge equal to -1 . We can write any such connection in terms of the canonical connection as in eq. (2.9), i.e. $\nabla = \nabla_0 + \alpha$ with α a one-form valued endomorphism of \mathcal{E} . Clearly, this will not change the value of the topological charge. We are particularly interested in $\text{SU}(2)$ -instantons, so we impose that α is traceless and skew-hermitian. Here the trace is taken in the second leg of $\text{End}(\mathcal{E}) \simeq P \boxtimes_{\text{ad}} M_2(\mathbb{C})$. When complexified, this gives an element $\alpha \in \Omega^1(S_\theta^4) \otimes_{C^\infty(S_\theta^4)} \Gamma(\text{ad}(S_{\theta'}^7)) =: \Omega^1(\text{ad}(S_\theta^4))$ (cf. Example 16).

As usual, we impose an irreducibility condition on the instanton connections, a connection on \mathcal{E} being *irreducible* if it cannot be written as the sum of two other connections on \mathcal{E} . We are interested only in the irreducible instanton connections on the module \mathcal{E} .

Remark 21. In [31], we constructed projections $p_{(n)}$ for all modules $C^\infty(S_{\theta'}^7) \boxtimes_\rho \mathbb{C}^n$ over $C^\infty(S_\theta^4)$ associated to the irreducible representations \mathbb{C}^n of $\text{SU}(2)$. The induced Grassmann connections $\nabla_0^{(n)} := p_{(n)}d$, when acting on $C^\infty(S_{\theta'}^7) \boxtimes_\rho \mathbb{C}^n$, were written as $d + \omega_{(n)}$, with $\omega_{(n)}$ an $n \times n$ matrix with entries in $\Omega^1(S_{\theta'}^7)$. A similar argument as above then shows that all $\omega_{(n)}$ have entries that are central (as one forms) in $\Omega(S_{\theta'}^7)$; again, this means that $L_\theta(\omega_{(n)}) = \omega_{(n)}$. In particular, this holds for the adjoint bundle associated to the adjoint representation on $\mathfrak{su}(2)_\mathbb{C} \simeq \mathbb{C}^3$ (as complex representation spaces), from which we conclude that $\nabla_0^{(2)}$ coincides with $[\nabla_0, \cdot]$ (since this is the case if $\theta = 0$).

5.2 Twisted infinitesimal symmetries

The noncommutative sphere S_θ^4 can be realized as a quantum homogeneous space of the quantum orthogonal group $\text{SO}_\theta(5)$ [42, 18]. In other words, $\mathcal{A}(S_\theta^4)$ can be obtained as the subalgebra of $\mathcal{A}(\text{SO}_\theta(5))$ made of the elements that are coinvariant under the natural coaction of $\text{SO}_\theta(4)$ on $\text{SO}_\theta(5)$. For our purposes, it turns out to be more convenient to take a dual point of view and consider an *action* instead of a coaction [41]. We obtain a twisted symmetry action of the enveloping algebra $U_\theta(\mathfrak{so}(5))$ on S_θ^4 . This is lifted to $S_{\theta'}^7$ and the above basic instanton ∇_0 is invariant under this infinitesimal quantum symmetry.

Different instantons are obtained by a twisted symmetry action of $U_\theta(\mathfrak{so}(5, 1))$. Classically, $\mathfrak{so}(5, 1)$ is the conformal Lie algebra consisting of the infinitesimal diffeomorphisms leaving the conformal structure invariant. We construct the Hopf algebra $U_\theta(\mathfrak{so}(5, 1))$ by adding 5 generators to $U_\theta(\mathfrak{so}(5))$ and describe its action on S_θ^4 together with its lift to $S_{\theta'}^7$. The induced action of $U_\theta(\mathfrak{so}(5, 1))$ on forms leaves the conformal structure invariant. The action of $U_\theta(\mathfrak{so}(5, 1))$ on ∇_0 eventually results in a five-parameter family of instantons.

Let us start with the construction of the twisted symmetry $U_\theta(\mathfrak{so}(5))$. The eight roots of the Lie algebra $\mathfrak{so}(5)$ are two-component vectors $r = (r_1, r_2)$ of the form $r = (\pm 1, \pm 1)$ and $r = (0, \pm 1), r = (\pm 1, 0)$. There are corresponding generators E_r of $\mathfrak{so}(5)$ together with two mutually commuting generators H_1, H_2 of the Cartan subalgebra. The Lie brackets are

$$\begin{aligned} [H_1, H_2] &= 0, & [H_j, E_r] &= r_j E_r, \\ [E_{-r}, E_r] &= r_1 H_1 + r_2 H_2, & [E_r, E_{r'}] &= N_{r,r'} E_{r+r'}, \end{aligned} \tag{5.5}$$

with $N_{r,r'} = 0$ if $r + r'$ is not a root. The universal enveloping algebra $U(\mathfrak{so}(5))$ is the algebra generated by elements $\{H_j, E_r\}$ modulo relations given by the previous Lie brackets². It is a Hopf algebra with the (primitive) coproduct

$$\begin{aligned}\Delta_0 : U(\mathfrak{so}(5)) &\rightarrow U(\mathfrak{so}(5)) \otimes U(\mathfrak{so}(5)), \\ X &\mapsto \Delta_0(X) = X \otimes \mathbb{1} + \mathbb{1} \otimes X,\end{aligned}$$

the counit $\varepsilon(X) = 0$ and the antipode $S(X) = -X$.

The twisted universal enveloping algebra $U_\theta(\mathfrak{so}(5))$ is generated as above (i.e. one does not change the algebra structure) but is endowed with a twisted coproduct,

$$\begin{aligned}\Delta_\theta : U_\theta(\mathfrak{so}(5)) &\rightarrow U_\theta(\mathfrak{so}(5)) \otimes U_\theta(\mathfrak{so}(5)), \\ X &\mapsto \Delta_\theta(X) = \mathcal{F}\Delta_0(X)\mathcal{F}^{-1}.\end{aligned}$$

For the symmetries studied in the present paper the twist \mathcal{F} is given explicitly [41] by

$$\mathcal{F} = \lambda^{-H_1 \otimes H_2}. \quad (5.6)$$

On the generators E_r, H_j , the twisted coproduct reads

$$\begin{aligned}\Delta_\theta(E_r) &= E_r \otimes \lambda^{-r_1 H_2} + \lambda^{-r_2 H_1} \otimes E_r, \\ \Delta_\theta(H_j) &= H_j \otimes \mathbb{1} + \mathbb{1} \otimes H_j.\end{aligned} \quad (5.7)$$

This coproduct allows one to represent $U_\theta(\mathfrak{so}(5))$ as an algebra of twisted derivations on both S_θ^4 and S_θ^7 , as we shall see below. With counit and antipode given by

$$\begin{aligned}\varepsilon(E_r) &= \varepsilon(H_j) = 0, \\ S(E_r) &= -\lambda^{r_2 H_1} E_r \lambda^{r_1 H_2}, \quad S(H_j) = -H_j,\end{aligned} \quad (5.8)$$

the algebra $U_\theta(\mathfrak{so}(5))$ becomes a Hopf algebra [11]. At the classical value of the deformation parameter, $\theta = 0$, one recovers the Hopf algebra structure of $U(\mathfrak{so}(5))$.

We are ready for the representation of $U_\theta(\mathfrak{so}(5))$ on S_θ^4 . For convenience, we introduce ‘‘partial derivatives’’, ∂_μ and ∂_μ^* with the usual action on the generators of the algebra $\mathcal{A}(S_\theta^4)$ i.e. $\partial_\mu(z_\nu) = \delta_{\mu\nu}$, $\partial_\mu(z_\nu^*) = 0$, and $\partial_\mu^*(z_\nu^*) = \delta_{\mu\nu}$, $\partial_\mu^*(z_\nu) = 0$. Then, the action of $U_\theta(\mathfrak{so}(5))$ on $\mathcal{A}(S_\theta^4)$ is given by the following operators,

$$\begin{aligned}H_1 &= z_1 \partial_1 - z_1^* \partial_1^*, & H_2 &= z_2 \partial_2 - z_2^* \partial_2^* \\ E_{+1,+1} &= z_2 \partial_1^* - z_1 \partial_2^*, & E_{+1,-1} &= z_2^* \partial_1^* - z_1 \partial_2, \\ E_{+1,0} &= \frac{1}{\sqrt{2}}(2z_0 \partial_1^* - z_1 \partial_0), & E_{0,+1} &= \frac{1}{\sqrt{2}}(2z_0 \partial_2^* - z_2 \partial_0),\end{aligned} \quad (5.9)$$

and $E_{-r} = (E_r)^*$, with the obvious meaning of the adjoint.

Remark 22. *Note that the operators H_1 and H_2 are the infinitesimal generators of the action of \mathbb{T}^2 on S_θ^4 as given in eq. (4.13).*

²There are additional Serre relations; they generate an ideal that needs to be quotiented out. This is not problematic and we shall not dwell upon this point here.

These operators (not the partial derivatives!) are extended to the whole of $\mathcal{A}(S_\theta^4)$ as twisted derivations via the coproduct (5.7),

$$\begin{aligned} E_r(ab) &= \Delta_\theta(E_r)(a \otimes b) = E_r(a)\lambda^{-r_1 H_2}(b) + \lambda^{-r_2 H_1}(a)E_r(b), \\ H_j(ab) &= \Delta_\theta(H_j)(a \otimes b) = H_j(a)b + aH_j(b), \end{aligned} \quad (5.10)$$

for any two elements $a, b \in \mathcal{A}(S_\theta^4)$. With these twisted rules, one readily checks compatibility with the commutation relations (4.1) of $\mathcal{A}(S_\theta^4)$.

We can also write the twisted action of the Hopf algebra $U_\theta(so(5))$ on $\mathcal{A}(S_\theta^4)$ by using the quantization map L_θ introduced in Section 3. For $L_\theta(a) \in \mathcal{A}(S_\theta^4)$ and $T \in U_\theta(so(5))$ we define a twisted action by

$$T \cdot L_\theta(a) = L_\theta(t \cdot a) \quad (5.11)$$

where t is the classical limit ($\theta = 0$) of T and $t \cdot a$ is the classical action of $U(so(5))$ on $\mathcal{A}(S^4)$. One checks that both of these definitions of the twisted action coincide (cf. [41]).

The representation of $U_\theta(so(5))$ on S_θ^4 given in (5.9) is the fundamental vector representation. When lifted to S_θ^7 one gets the fundamental spinor representation: as we see from the quadratic relations among corresponding generators, as given in (4.9), the lifting amounts to take the ‘‘square root’’ representation. The action on $U_\theta(so(5))$ on $\mathcal{A}(S_\theta^7)$ is constructed by requiring twisted derivation properties via the coproduct (5.10) so as to reduce to the action (5.9) on $\mathcal{A}(S_\theta^4)$ when using the defining quadratic relations (4.9). The action on $\mathcal{A}(S_\theta^7)$ can be given as the action of matrices Γ 's on the ψ 's,

$$\psi_a \mapsto \sum_b \Gamma_{ab} \psi_b; \quad \psi_a^* \mapsto \sum_b \tilde{\Gamma}_{ab} \psi_b^* \quad (5.12)$$

with the matrices $\Gamma = \{H_j, E_r\}$ given explicitly by,

$$\begin{aligned} H_1 &= \frac{1}{2} \begin{pmatrix} 1 & & & \\ & -1 & & \\ & & -1 & \\ & & & 1 \end{pmatrix}, & H_2 &= \frac{1}{2} \begin{pmatrix} -1 & & & \\ & 1 & & \\ & & -1 & \\ & & & 1 \end{pmatrix}, \\ E_{+1,+1} &= \begin{pmatrix} 0 & 0 & & \\ 0 & 0 & -1 & \\ & 0 & 0 & \\ & & & 0 \end{pmatrix}, & E_{+1,-1} &= \begin{pmatrix} 0 & 0 & 0 & \\ -\mu & 0 & 0 & \\ 0 & & 0 & \\ & & & 0 \end{pmatrix}, \\ E_{+1,0} &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & \\ \mu & 0 & 0 & \\ 0 & 0 & 0 & \\ & & & 0 \end{pmatrix}, & E_{0,+1} &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & \bar{\mu} & \\ 0 & 1 & 0 & \\ 0 & 0 & 0 & \\ & & & 0 \end{pmatrix}, \end{aligned} \quad (5.13)$$

and $\tilde{\Gamma} := \sigma \Gamma \sigma^{-1}$ with

$$\sigma := \begin{pmatrix} 0 & -1 & & \\ 1 & 0 & 0 & \\ & 0 & & \\ & 0 & -1 & \\ & & & 0 \end{pmatrix}. \quad (5.14)$$

Furthermore, $E_{-r} = (E_r)^*$. With the twisted rules (5.10) for the action on products, one checks compatibility of the above action with the commutation relations (4.4) of $\mathcal{A}(S_\theta^7)$.

Remark 23. Compare the form of the matrices H_1 and H_2 above with the lifted action $\tilde{\sigma}$ of $\tilde{\mathbb{T}}^2$ on S_θ^7 as defined in (4.14). One checks that

$$\tilde{\sigma}_s = e^{\pi i((s_1+s_2)H_1 + (-s_1+s_2)H_2)},$$

when acting on the spinor (ψ_1, \dots, ψ_4) .

Notice that $\tilde{\Gamma} = -\Gamma^t$ at $\theta = 0$. There is a beautiful correspondence between these matrices and the twisted Dirac matrices introduced in (4.11),

$$\begin{aligned}\frac{1}{4}[\gamma_1^*, \gamma_1] &= 2H_1 & \frac{1}{4}[\gamma_2^*, \gamma_2] &= 2H_2 \\ \frac{1}{4}[\gamma_1, \gamma_2] &= (\mu + \bar{\mu})E_{+1,+1} & \frac{1}{4}[\gamma_1, \gamma_2^*] &= (\mu + \bar{\mu})E_{+1,-1} \\ \frac{1}{4}[\gamma_1, \gamma_0] &= \sqrt{2}E_{+1,0} & \frac{1}{4}[\gamma_2, \gamma_0] &= \sqrt{2}\bar{\mu}E_{0,+1}\end{aligned}\quad (5.15)$$

Remark 24. *The twisted Dirac matrices satisfy the following relations under conjugation by σ :*

$$(\sigma\gamma_0\sigma^{-1})^t = \gamma_0; \quad (\sigma\gamma_1\sigma^{-1})^t = \gamma_1\lambda^{H_2}; \quad (\sigma\gamma_2\sigma^{-1})^t = \gamma_2\lambda^{H_1}. \quad (5.16)$$

Due to the form of $\tilde{\Gamma}$ and the property $\Psi_{a2} = \sigma_{ab}\psi_b^*$ for the second column of the matrix Ψ in (4.6), we have also that $U_\theta(so(5))$ acts on Ψ by left matrix multiplication by Γ , and by right matrix multiplication on Ψ^* by the matrix transpose $\tilde{\Gamma}^t$ as follows

$$\Psi_{ai} \mapsto \sum_b \Gamma_{ab}\Psi_{bi}, \quad \Psi_{ia}^* \mapsto \sum_a \Psi_{ib}^* \tilde{\Gamma}_{ab}^t. \quad (5.17)$$

These are used in the following

Proposition 25. *The instanton gauge potential ω is invariant under the twisted action of $U_\theta(so(5))$.*

Proof. From the above observations, the gauge potential transforms as:

$$\omega = \Psi^* d\Psi \mapsto \Psi^* (\tilde{\Gamma}^t \lambda^{-r_1 H_2} + \lambda^{r_2 H_1} \Gamma) d\Psi.$$

where $\lambda^{-r_i H_j}$ is understood in its representation (5.13) on $\mathcal{A}(S^4)$. Direct computation for $\Gamma = \{H_j, E_r\}$ shows that $\tilde{\Gamma}^t \lambda^{-r_1 H_2} + \lambda^{r_2 H_1} \Gamma = 0$, which finishes the proof. \square

5.3 Twisted conformal transformations

In order to have new instantonic configurations we need to use conformal transformations. The conformal Lie algebra $so(5, 1)$ consists of the generators of $so(5)$ together with the dilation and the so-called special conformal transformations. On \mathbb{R}^4 with coordinates $\{x_\mu, \mu = 1, \dots, 4\}$ they are given by the operators $H_0 = \sum_\mu x_\mu \partial / \partial x_\mu$ and $G_\mu = 2x_\mu \sum_\nu x_\nu \partial / \partial x_\nu - \sum_\nu x_\nu^2 (\partial / \partial x_\mu)$, respectively [32].

In the definition of the enveloping algebra $U_\theta(so(5, 1))$ we do not change the algebra structure, i.e. we take the relations of $U(so(5, 1))$, as we did in the case of $U_\theta(so(5))$. We thus define $U_\theta(so(5, 1))$ as the algebra $U_\theta(so(5))$ with five extra generators adjoined, $H_0, G_r, r = (\pm 1, 0), (0, \pm 1)$, subject to the relations of $U_\theta(so(5))$ of eq. (5.5) together with the (undeformed) relations,

$$\begin{aligned}[H_0, H_i] &= 0, & [H_j, G_r] &= r_j G_r, \\ [H_0, G_r] &= \sqrt{2}E_r, & [H_0, E_r] &= (\sqrt{2})^{-1}G_r,\end{aligned}\quad (5.18)$$

whenever $r = (\pm 1, 0), (0, \pm 1)$, and

$$\begin{aligned}[G_{-r}, G_r] &= 2r_1 H_1 + 2r_2 H_2, & [G_r, G_{r'}] &= N_{r,r'} E_{r+r'}, \\ [E_r, G_{r'}] &= \tilde{N}_{r,r'} G_{r+r'}, & [E_{-r}, G_r] &= \sqrt{2}H_0,\end{aligned}\quad (5.19)$$

with as before $N_{r,r'} = 0$ if $r + r'$ is not of $so(5)$ and $\tilde{N}_{r,r'} = 0$ if $r + r' \notin \{(\pm 1, 0), (0, \pm 1)\}$. Although the algebra structure is unchanged, again the Hopf algebra structure of $U_\theta(so(5, 1))$ gets twisted. The twisted structures are given by equations (5.7) and (5.8) together with analogous ones for the extra generators,

$$\begin{aligned}\Delta_\theta(G_r) &= G_r \otimes \lambda^{-r_1 H_2} + \lambda^{-r_2 H_1} \otimes G_r, & \Delta_\theta(H_0) &= H_0 \otimes 1 + 1 \otimes H_0, \\ S(G_r) &= -\lambda^{r_2 H_1} G_r \lambda^{r_1 H_2}, & S(H_0) &= -H_0, \\ \varepsilon(G_r) &= 0, & \varepsilon(H_0) &= 0\end{aligned}\tag{5.20}$$

These structures make $U_\theta(so(5, 1))$ a Hopf algebra [11].

The action of $U_\theta(so(5, 1))$ on $\mathcal{A}(S_\theta^4)$ is given by the operators (5.9) together with

$$\begin{aligned}H_0 &= \partial_0 - z_0(z_0 \partial_0 + z_1 \partial_1 + z_1^* \partial_1^* + z_2 \partial_2 + z_2^* \partial_2^*), \\ G_{1,0} &= 2\partial_1^* - z_1(z_0 \partial_0 + z_1 \partial_1 + z_1^* \partial_1^* + \bar{\lambda} z_2 \partial_2 + \lambda z_2^* \partial_2^*), \\ G_{0,1} &= 2\partial_2^* - z_2(z_0 \partial_0 + z_1 \partial_1 + z_1^* \partial_1^* + z_2 \partial_2 + z_2^* \partial_2^*),\end{aligned}\tag{5.21}$$

and $G_{-r} = (G_r)^*$. Note that the introduction of the extra λ 's in $G_{1,0}$ (and $G_{-1,0}$) are necessary for the algebra structure of $U_\theta(so(5, 1))$, as dictated by the above Lie brackets in (5.18), (5.19) to be preserved. Since the operators H_0 and G_r are quadratic in the z 's, one has to be careful when deriving the above Lie brackets and use the twisted rules (5.10). For instance, on the generator z_2 , we have

$$\begin{aligned}[E_{-1,-1}, G_{1,0}](z_2) &= E_{-1,-1}(-\bar{\lambda} z_1 z_2) + G_{1,0}(z_1^*) \\ &= -\bar{\lambda}(E_{-1,-1}(z_1) \lambda^{H_2}(z_2) + \lambda^{H_1}(z_1) E_{-1,-1}(z_2)) + G_{1,0}(z_1^*) \\ &= -z_2^* z_2 + z_1 z_1^* + 2 - z_1 z_1^* = G_{0,-1}(z_2)\end{aligned}$$

Again, the operators H_0, G_r in (5.21) are extended to the whole of $\mathcal{A}(S_\theta^4)$ by analogues of (5.10) using the twisted coproduct,

$$\begin{aligned}G_r(ab) &:= \Delta_\theta G_r = G_r(a) \lambda^{-r_1 H_2}(b) + \lambda^{-r_2 H_1}(a) G_r(b), \\ H_0(ab) &:= \Delta_\theta H_0 = H_0(a)b + aH_0(b), \quad \forall a, b \in \mathcal{A}(S_\theta^4).\end{aligned}\tag{5.22}$$

Equivalently, the Hopf algebra $U_\theta(so(5, 1))$ could be defined to act on $\mathcal{A}(S_\theta^4)$ by

$$T \cdot L_\theta(a) = L_\theta(t \cdot a),\tag{5.23}$$

for $T \in U_\theta(so(5, 1))$ deforming $t \in U(so(5, 1))$ and $L_\theta(a) \in \mathcal{A}(S_\theta^4)$ deforming $a \in \mathcal{A}(S^4)$.

The Hopf algebra $U_\theta(so(5, 1))$ consists of the infinitesimal twisted conformal transformations on S_θ^4 due to the following lemma.

Lemma 26. *1. The twisted action of the Hopf algebra $U_\theta(so(5, 1))$ on $\mathcal{A}(S_\theta^4)$ can be extended to the differential calculus $(\Omega(S_\theta^4), d)$ by requiring it to commute with the exterior derivative,*

$$T \cdot d\omega := d(T \cdot \omega).$$

for $T \in U_\theta(so(5, 1))$, $\omega \in \Omega(S_\theta^4)$.

2. Under this twisted action, the Hopf algebra $U_\theta(\mathfrak{so}(5, 1))$ leaves the Hodge $*_\theta$ -structure of $\Omega(S_\theta^4)$ invariant,

$$T \cdot (*_\theta \omega) = *_\theta(T \cdot \omega),$$

Proof. 1. is Lemma 3 in [41] and 2. follows from the fact that $T(L_\theta(a)) = L_\theta(t \cdot a)$ for $a \in \mathcal{A}(S^4)$ and $t \in U(\mathfrak{so}(5, 1))$ the classical limit ($\theta = 0$) of $T \in U_\theta(\mathfrak{so}(5, 1))$. Then, since $U(\mathfrak{so}(5, 1))$ leaves the Hodge $*$ -structure of $\Omega(S^4)$ invariant and the differential d commutes with the action of $U_\theta(\mathfrak{so}(5, 1))$, it follows that the latter algebra leaves the Hodge $*_\theta$ -structure of $\Omega(S_\theta^4)$ invariant as well. \square

In the same manner as for $U_\theta(\mathfrak{so}(5))$, the action of the Hopf algebra $U_\theta(\mathfrak{so}(5, 1))$ on S_θ^4 can be lifted to an action on S_θ^7 . Again, the latter action can be written as in (5.12) in terms of matrices Γ 's acting on the ψ 's,

$$\psi_a \mapsto \sum_b \Gamma_{ab} \psi_b, \quad \psi_a^* \mapsto \sum_b \tilde{\Gamma}_{ab} \psi_b^*, \quad (5.24)$$

where in addition to (5.13) we have the additional matrices $\Gamma = \{H_0, G_r\}$,

$$\begin{aligned} H_0 &= \frac{1}{2}(-z_0 \mathbb{I}_4 + \gamma_0), \\ G_{1,0} &= \frac{1}{2}(-z_1 \lambda^{-H_2} + \gamma_1), \quad G_{0,1} = \frac{1}{2}(-z_2 + \lambda^{-H_1} \gamma_2), \end{aligned} \quad (5.25)$$

$G_{-r} = (G_r)^*$ and $\tilde{\Gamma} = \sigma \Gamma \sigma^{-1}$. Notice the reappearance of the twisted Dirac matrices γ_μ, γ_μ^* of (4.11) in the above expressions.

As for $U_\theta(\mathfrak{so}(5))$, the action of $U_\theta(\mathfrak{so}(5, 1))$ on the matrix Ψ is found to be by left matrix multiplication by Γ and on Ψ^* by $\tilde{\Gamma}$,

$$\Psi_{ai} \mapsto \sum_b \Gamma_{ab} \Psi_{bi}, \quad \Psi_{ia}^* \mapsto \sum_a \tilde{\Gamma}_{ab} \Psi_{ib}^*. \quad (5.26)$$

Here we have to be careful with the ordering between $\tilde{\Gamma}$ and Ψ^* in the second term since the $\tilde{\Gamma}$'s involve the (not-central) z 's. There are the following useful commutation relations between the z 's and Ψ :

$$\begin{aligned} z_1 \Psi_{ai} &= (\lambda^{-H_2})_{ab} \Psi_{bi} z_1, & z_2 \Psi_{ai} &= (\lambda^{-H_1})_{ab} \Psi_{bi} z_2, \\ z_1 \Psi_{ia}^* &= \Psi_{ib}^* (\lambda^{-H_2})_{ba} z_1, & z_2 \Psi_{ia}^* &= \Psi_{ib}^* (\lambda^{-H_1})_{ba} z_2. \end{aligned} \quad (5.27)$$

with λ^{-H_j} understood as 4×4 matrices in the suitable representation.

Proposition 27. *The instanton gauge potential $\omega = \Psi^* d\Psi$ transforms under the action of the Hopf algebra $U_\theta(\mathfrak{so}(5, 1))$ as $\omega \mapsto \omega + \delta\omega_i$, where*

$$\begin{aligned} \delta\omega_0 &:= H_0(\omega) = -z_0 \omega - \frac{1}{2} dz_0 \mathbb{I}_2 + \Psi^* \gamma_0 d\Psi, \\ \delta\omega_1 &:= G_{+1,0}(\omega) = -z_1 \omega - \frac{1}{2} dz_1 \mathbb{I}_2 + \Psi^* \gamma_1 d\Psi, \\ \delta\omega_2 &:= G_{0,+1}(\omega) = -z_2 \omega - \frac{1}{2} dz_2 \mathbb{I}_2 + \Psi^* \gamma_2 d\Psi, \\ \delta\omega_3 &:= G_{-1,0}(\omega) = -\omega \bar{z}_1 - \frac{1}{2} d\bar{z}_1 \mathbb{I}_2 + \Psi^* \gamma_1^* d\Psi, \\ \delta\omega_4 &:= G_{0,-1}(\omega) = -\omega \bar{z}_2 - \frac{1}{2} d\bar{z}_2 \mathbb{I}_2 + \Psi^* \gamma_2^* d\Psi, \end{aligned}$$

with γ_μ, γ_μ^* the twisted 4×4 Dirac matrices defined in (4.11).

Proof. The action of H_0 on the instanton gauge potential $\omega = \Psi^*d\Psi$ takes the form

$$H_0(\omega) = H_0(\Psi^*)d\Psi + \Psi^*d(H_0(\Psi)) = \Psi^*(-z_0\mathbb{I}_4 + \gamma_0)d\Psi - \frac{1}{2}dz_0\Psi^*\Psi,$$

since z_0 is central. Direct computation results in the above expression for $\delta\omega_0$. Instead, the twisted action of G_r on ω takes the form,

$$G_r : \omega_{ij} \mapsto \sum_{a,b,c} \tilde{\Gamma}_{ab} \Psi_{ib}^* (\lambda^{-r_1 H_2})_{ac} d\Psi_{cj} + (\lambda^{r_2 H_1})_{ab} \Psi_{ib}^* \Gamma_{ac} d\Psi_{cj} + (\lambda^{r_2 H_1})_{ab} \Psi_{ib}^* (d\Gamma_{ac}) \Psi_{cj},$$

where we used the fact that $\tilde{H}_j = \sigma H_j \sigma^{-1} = -H_j$. Let us consider the case $r = (+1, 0)$. Firstly, note that the complex numbers $(\lambda^{-H_2})_{ac}$ commute with Ψ_{ib}^* so that from the definition of Γ and $\tilde{\Gamma}$, and using (5.27), we obtain for the first two terms,

$$-z_1(\Psi^*d\Psi)_{ij} + \frac{1}{2}\Psi_{ib}^*(\sigma\gamma_1\sigma^{-1})_{cb}(\lambda^{-H_2})_{cd}d\Psi_{dj} + \frac{1}{2}\Psi_{ib}^*(\gamma_1)_{bc}d\Psi_{cj}.$$

The first term forms the matrix $-z_1\omega$ whereas the second two terms combine to give $\Psi^*\gamma_1d\Psi$, due to relation (5.16). Finally, using eq. (5.27) the term $\Psi_{ib}^*(d\Gamma_{ac})\Psi_{cj}$ reduces to $-\frac{1}{2}dz_1\Psi_{ib}^*\Psi_{bj} = -\frac{1}{2}dz_1\mathbb{I}_2$. The formulae for $r = (-1, 0)$ and $r = (0, \pm 1)$ are established in likewise manner. \square

Remark 28. *At first sight, the infinitesimal gauge potentials $\delta\omega_j$ given above do not seem to be $su(2)$ -gauge potentials, in that they do not satisfy $\overline{(\delta\omega_j)_{kl}} = (\delta\omega_j)_{lk}$ and $\sum_k (\delta\omega_j)_{kk} = 0$. This is only due to the fact that the generators G_r and H_0 are the deformed analogues of the generators of the complexified Lie algebra $so(5, 1) \otimes_{\mathbb{R}} \mathbb{C}$. One recovers $su(2)$ -gauge potentials by acting with the real generators $\frac{1}{2}(G_r + G_r^*)$, $\frac{1}{2i}(G_r - F_r^*)$ and H_0 . The resulting gauge potentials, $\delta\omega_0$, $\frac{1}{2}(\delta\omega_1 + \delta\omega_3)$, $\frac{1}{2i}(\delta\omega_1 - \delta\omega_3)$, $\frac{1}{2}(\delta\omega_2 + \delta\omega_4)$ and $\frac{1}{2i}(\delta\omega_2 - \delta\omega_4)$, are traceless skew-hermitian matrices with entries in $\Omega^1(S_{\theta'}^7)$.*

The transformations in Proposition 27 of the gauge potential ω under the twisted symmetry $U_{\theta}(so(5, 1))$ induce natural transformations of the canonical connection ∇_0 in (5.1) to $\nabla_{t,i} := \nabla_0 + t\delta\omega_i + \mathcal{O}(t^2)$. We shall presently see explicitly that these new connections are (infinitesimal) instantons, i.e. their curvatures are self-dual. In fact, this also follows from Lemma 26 which states that $U_{\theta}(so(5, 1))$ acts by conformal transformation therefore leaving the self-duality equation $*_{\theta}F_0 = F_0$ for the basic instanton ∇_0 invariant.

We start by writing $\nabla_{t,i}$ in terms of the canonical connection on $\mathcal{E} \simeq p(\mathcal{A}(S_{\theta}^4))^4$. Using the explicit isomorphism, described in Example 16, between this module and the module of equivariant maps $\mathcal{A}(S_{\theta}^7) \boxtimes_{\rho} \mathbb{C}^2$, we find that $\nabla_{t,i} = pd + t\delta\alpha_i + \mathcal{O}(t^2)$ with explicit expressions

$$\begin{aligned} \delta\alpha_0 &= p\gamma_0(dp)p - \frac{1}{2}\Psi dz_0\Psi^*, \\ \delta\alpha_1 &= p\gamma_1(dp)p - \frac{1}{2}\Psi dz_1\Psi^*, & \delta\alpha_3 &= p\gamma_1^*(dp)p - \frac{1}{2}\Psi dz_1^*\Psi^*, \\ \delta\alpha_2 &= p\gamma_2(dp)p - \frac{1}{2}\Psi dz_2\Psi^*, & \delta\alpha_4 &= p\gamma_2^*(dp)p - \frac{1}{2}\Psi dz_2^*\Psi^*, \end{aligned} \quad (5.28)$$

The $\delta\alpha'_i$'s are 4×4 matrices with entries in the one-forms $\Omega^1(S_{\theta}^4)$ and satisfying $p\delta\alpha_i = \delta\alpha_i p = p\delta\alpha_i p = \delta\alpha_i$, as expected from the general theory on connections on modules in Section 2. Indeed, using relations (5.27) one can move the dz 's to the left of Ψ at the cost of some μ 's, so getting expression like $dz_i p \in M_4(\Omega^1(S_{\theta}^4))$.

From eq. (2.10), the curvature $F_{t,i}$ of the connection $\nabla_{t,i}$ is given by

$$F_{t,i} = F_0 + tpd(\delta\alpha_i) + \mathcal{O}(t^2). \quad (5.29)$$

In order to check self-duality (modulo t^2) of this curvature, we will express it in terms of the projection p and consider $F_{t,i}$ as a two-form valued endomorphism on $\mathcal{E} \simeq p(\mathcal{A}(S_\theta^4))^4$.

Proposition 29. *The curvatures $F_{t,i}$ of the connections $\nabla_{t,i}$, $i = 0, \dots, 4$, are given by $F_{t,i} = F_0 + t\delta F_i + \mathcal{O}(t^2)$, where $F_0 = pdpd p$ and*

$$\begin{aligned} \delta F_0 &= -2z_0 F_0, \\ \delta F_1 &= -2z_1 \lambda^{H_2} F_0, & \delta F_3 &= -2z_1^* \lambda^{-H_2} F_0, \\ \delta F_2 &= -2z_2 \lambda^{H_1} F_0; & \delta F_4 &= -2z_2^* \lambda^{-H_1} F_0. \end{aligned} \quad (5.30)$$

Proof. A small computation yields for $\delta F_i = pd(\delta\alpha_i)$, thought of as an $\Omega^2(S_\theta^4)$ -valued endomorphism on \mathcal{E} the expression, $\delta F_i = p(dp)\gamma_i(dp)p - p\gamma_i(dp)(dp)p$, with the notation $\gamma_3 = \gamma_1^*$ and $\gamma_4 = \gamma_2^*$, and using $p(dp)p = 0$. Then, the crucial property $p(dp\gamma_i + \gamma_i dp)(dp)p = 0$ all $i = 0, \dots, 4$ yields $\delta F_i = -2p\gamma_i dpdp$. This is expressed as $\delta F_i = -2p\gamma_i pdpd p$ by using $dp = (dp)p + pdp$. Finally, $p\gamma_i p = \Psi(\Psi^*\gamma_i\Psi)\Psi^*$, so that the result follows from the definition of the z 's in terms of the Dirac matrices given in eq. (4.10), together with the commutation relations between them and the matrix Ψ in eq. (5.27) \square

Proposition 30. *The connections $\nabla_{t,i}$ are (infinitesimal) instantons, i.e.*

$$*_\theta F_{t,i} = F_{t,i} \quad \text{mod } t^2. \quad (5.31)$$

Moreover, the connections $\nabla_{t,i}$ are not gauge equivalent to ∇_0 .

Proof. The first point follows directly from the above expressions for δF_i and the self-duality of F_0 .

To establish the not gauge equivalence of the connections $\nabla_{t,i}$ with ∇_0 , recall that an infinitesimal gauge transformation is given by $\nabla_0 \mapsto \nabla_0 + t[\nabla_0, X]$ for $X \in \Gamma(\text{ad}(S_{\theta'}^7))$. We need to show that $\delta\omega_i$ is orthogonal to $[\nabla_0, X]$ for any such X , i.e. that

$$([\nabla_0, X], \delta\omega_i)_2 = 0,$$

with the natural inner product on $\Omega^1(\text{ad}(S_{\theta'}^7)) := \Omega^1(S_\theta^4) \otimes_{C^\infty(S_\theta^4)} \Gamma(\text{ad}(S_{\theta'}^7))$. From Remark 21, it follows that

$$(\nabla_0^{(2)}(X), \delta\omega_i)_2 = (X, (\nabla_0^{(2)})^*(\delta\omega_i))_2,$$

which then should vanish for all X . From equation (5.23), we see that $\delta\omega_i = T_i(\omega)$ coincides with $L_\theta(t_i \cdot \omega^{(0)})$ with t_i and $\omega^{(0)}$ the classical counterparts of T_i and ω , respectively. In the undeformed case, the infinitesimal gauge potentials generated by acting with elements in $so(5, 1) - so(5)$ on the basic instanton gauge potential $\omega^{(0)}$ satisfy $(\nabla_0^{(2)})^*(\delta\omega_i^{(0)}) = 0$ as shown in [6]. The result then follows from the observation that $\nabla_0^{(2)}$ commutes with the quantization map L_θ (cf. Remark 21). \square

5.4 Local expressions

In this section, we obtain “local expressions” for the instantons on S_θ^4 constructed in the previous section; that is we map then to a noncommutative \mathbb{R}_θ^4 obtained by ‘removing a point’ from S_θ^4 . The algebra $\mathcal{A}(\mathbb{R}_\theta^4)$ of polynomial functions on the 4-plane \mathbb{R}_θ^4 is defined to be the $*$ -algebra generated by ζ_1, ζ_2 satisfying

$$\zeta_1 \zeta_2 = \lambda \zeta_2 \zeta_1; \quad \zeta_1 \zeta_2^* = \bar{\lambda} \zeta_2^* \zeta_1. \quad (5.32)$$

with $\lambda = e^{2\pi i \theta}$ as above. At $\theta = 0$ one recovers the $*$ -algebra of polynomial functions on the usual 4-plane \mathbb{R}^4 . Again, one can define this algebra as the fixed point algebra $(\mathcal{A}(\mathbb{R}^4) \otimes \mathcal{A}(\mathbb{T}_\theta^2))^{\sigma \otimes \tau^{-1}}$; and the torus \mathbb{T}^2 acts as $\sigma_s : \zeta_i \mapsto e^{2\pi i s_i} \zeta_i$. In particular, one can define the algebra $C^\infty(\mathbb{R}_\theta^4)$ of smooth functions on \mathbb{R}_θ^4 . In this smooth algebra, there belongs the central element $(1 + |\zeta|^2)^{-1}$ with $|\zeta| := \zeta_1^* \zeta_1 + \zeta_2^* \zeta_2$. Then, one defines elements $\tilde{z}_\mu, \mu = 1, 2, 3$ as

$$\tilde{z}_0 = (1 - |\zeta|^2)(1 + |\zeta|^2)^{-1}, \quad \tilde{z}_j = 2\zeta_j(1 + |\zeta|^2)^{-1} \quad j = 1, 2, \quad (5.33)$$

and checks that they satisfy the same relations as in (4.1) of the generators z_μ of $\mathcal{A}(S_\theta^4)$. The difference is that the classical point $z_0 = -1, z_j = z_j^* = 0$ of S_θ^4 is not in the spectrum of \tilde{z}_μ . We interpret the noncommutative plane \mathbb{R}_θ^4 as a “chart” of the noncommutative 4-sphere S_θ^4 and the eq. (5.33) as the (inverse) stereographical projection from S_θ^4 to \mathbb{R}_θ^4 . In fact, one can cover S_θ^4 by two such charts with domain \mathbb{R}_θ^4 , and transition functions on $\mathbb{R}_\theta^4 \setminus \{0\}$, where $\{0\}$ is the classical point $\zeta_j = \zeta_j^* = 0$ of \mathbb{R}_θ^4 .

A differential calculus $(\Omega(\mathbb{R}_\theta^4), d)$ on \mathbb{R}_θ^4 is obtained from the general procedure described in Section 3. Explicitly, $\Omega(\mathbb{R}_\theta^4)$ is the graded $*$ -algebra generated by the elements ζ^μ of degree 0 and $d\zeta^\mu$ of degree 1 with relations,

$$\begin{aligned} d\zeta^\mu d\zeta^\nu + \lambda^{\mu\nu} d\zeta^\nu d\zeta^\mu &= 0, & d\bar{\zeta}^\mu d\zeta^\nu + \lambda^{\nu\mu} d\zeta^\nu d\bar{\zeta}^\mu &= 0, \\ \zeta^\mu d\zeta^\nu &= \lambda^{\mu\nu} d\zeta^\nu \zeta^\mu, & \bar{\zeta}^\mu d\zeta^\nu &= \lambda^{\nu\mu} d\zeta^\nu \bar{\zeta}^\mu. \end{aligned} \quad (5.34)$$

There is a unique differential d on $\Omega(\mathbb{R}_\theta^4)$ such that $d : \zeta^\mu \mapsto d\zeta^\mu$ and a Hodge star operator $*_\theta : \Omega^p(\mathbb{R}_\theta^4) \rightarrow \Omega^{4-p}(\mathbb{R}_\theta^4)$, obtained from the classical Hodge star operator as before. In terms of the standard Riemannian metric on \mathbb{R}^4 , on two-forms we have,

$$*_\theta d\zeta_1 d\zeta_2 = -d\zeta_1 d\zeta_2, \quad *_\theta d\zeta_1 d\zeta_1^* = -d\zeta_2 d\zeta_2^*, \quad *_\theta d\zeta_1 d\zeta_2^* = d\zeta_1 d\zeta_2^*, \quad (5.35)$$

and $*_\theta^2 = \text{id}$. These are the same formulae as the ones for the undeformed Hodge $*$ on \mathbb{R}^4 – since the metric is not changed in an isospectral deformation.

The stereographical projection from S^4 onto \mathbb{R}^4 is a conformal map commuting with the action of \mathbb{T}^2 ; thus it makes sense to investigate the form of the instanton connections on S_θ^4 obtained in Proposition 27 on the local chart \mathbb{R}_θ^4 . As in [28], we first introduce a “local section” of the principal bundle $S_\theta^7 \rightarrow S_\theta^4$ on the local chart of S_θ^4 defined in (5.33). Let $u = (u_1, u_2)$ be a complex spinor of modulus one, $u_1^* u_1 + u_2^* u_2 = 1$, and define

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = \rho \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}, \quad \begin{pmatrix} \psi_3 \\ \psi_4 \end{pmatrix} = \rho \begin{pmatrix} \zeta_1^* & \zeta_2^* \\ -\mu \zeta_2 & \bar{\mu} \zeta_1 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}. \quad (5.36)$$

Here ρ is a central element in $C^\infty(\mathbb{R}_\theta^4)$ such that $\rho^2 = (1 + |\zeta|^2)^{-1}$ and the commutations rules of the u_j 's with the ζ_k 's are dictated by those of the ψ_j ,

$$u_1\zeta_j = \mu\zeta_j u_1, \quad u_2\zeta_j = \bar{\mu}\zeta_j u_2, \quad j = 1, 2. \quad (5.37)$$

The right action of $SU(2)$ rotates the vector u while mapping to the ‘‘same point’’ of S_θ^4 , which, using the definition (4.9), from the choice in (5.36) is found to be

$$2(\psi_1\psi_3^* + \psi_2^*\psi_4) = \tilde{z}_1, \quad 2(-\psi_1^*\psi_4 + \psi_2\psi_3^*) = \tilde{z}_2, \quad 2(\psi_1^*\psi_1 + \psi_2^*\psi_2) - 1 = \tilde{z}_0, \quad (5.38)$$

and which is in the local chart (5.33), as expected.

By writing the unit vector u as an $SU(2)$ matrix, $u = \begin{pmatrix} u_1 & -u_2^* \\ u_2 & u_1^* \end{pmatrix}$, we have

$$\Psi = \rho \begin{pmatrix} \mathbb{I}_2 & 0 \\ 0 & \mathcal{Z} \end{pmatrix} \begin{pmatrix} u \\ u \end{pmatrix}, \quad \text{with } \mathcal{Z} = \begin{pmatrix} \zeta_1^* & \zeta_2^* \\ -\mu\zeta_2 & \bar{\mu}\zeta_1 \end{pmatrix}. \quad (5.39)$$

Then, by direct computation the gauge potential $\omega = \Psi^*d\Psi$ takes the form

$$\begin{aligned} u\omega u^* &= \rho^{-1}d\rho + \rho^2\mathcal{Z}^*d\mathcal{Z} + (du)u^* \\ &= \frac{1}{(1 + |\zeta|^2)} \begin{pmatrix} \sum_i \zeta_i d\zeta_i^* - d\zeta_i \zeta_i^* & 2(\zeta_1 d\zeta_2^* - d\zeta_1 \zeta_2^*) \\ 2(\zeta_2 d\zeta_1^* - d\zeta_2 \zeta_1^*) & \sum_i \zeta_i^* d\zeta_i - d\zeta_i^* \zeta_i \end{pmatrix} + (du)u^*, \end{aligned} \quad (5.40)$$

while its curvature $F = d\omega + \omega^2$ is,

$$uFu^* = \rho^4 d\mathcal{Z}^*d\mathcal{Z} = \frac{1}{(1 + |\zeta|^2)^2} \begin{pmatrix} d\zeta_1 d\zeta_1^* - d\zeta_2 d\zeta_2^* & 2d\zeta_1 d\zeta_2^* \\ 2d\zeta_2 d\zeta_1^* & -d\zeta_1 d\zeta_1^* - d\zeta_2 d\zeta_2^* \end{pmatrix}. \quad (5.41)$$

From (5.35) one checks that this curvature is self-dual: $*_\theta(uFu^*) = uFu^*$, as expected.

The explicit local expressions for the transformed gauge potentials and their curvature can be obtained in a similar manner. As an example, let us work out the local expression for $\delta\omega_0$, being the most transparent one. Given the expression for $\delta\omega_0$ in Proposition 27, a direct computation shows that

$$\delta\omega_0 = -2\rho d\rho u^*u - 2\rho^4 u^* \mathcal{Z}^* d\mathcal{Z}u, \quad (5.42)$$

which gives for the transformed curvature,

$$F_{t,0} = F_0 + 2t(1 - 2\rho^2)F_0 + \mathcal{O}(t^2). \quad (5.43)$$

It is clear that this rescaled curvature still satisfies the self-duality equation; this is also in concordance with Proposition 29, being $\tilde{z}_0 = 2\rho^2 - 1$.

5.5 Moduli space of instantons

We will closely follow the infinitesimal construction in [6] of instantons for the undeformed case. This will eventually result in the computation of the dimension of the ‘tangent space’ to the moduli space of instantons on S_θ^4 by index methods. It will turn out that

the five-parameter family of instantons constructed in the previous section is indeed the complete set of infinitesimal instantons on S_θ^4 .

Let us start by considering the following family of connections on S_θ^4 ,

$$\nabla_t = \nabla_0 + t\alpha \quad (5.44)$$

where $\alpha \in \Omega^1(\text{ad}(S_\theta^7))$. For ∇_t to be an instanton, we have to impose the self-duality equation $*_\theta F_t = F_t$ on the curvature $F_t = F_0^2 + t[\nabla, \alpha] + \mathcal{O}(t^2)$ of ∇_t . This leads, when differentiated with respect to t , setting $t = 0$ afterwards, to the following *linearized self-duality equation*

$$P_-[\nabla_0, \alpha] = 0, \quad (5.45)$$

with $P_- := \frac{1}{2}(1 - *_\theta)$ the projection onto the antiself-dual 2-forms. Here $[\nabla_0, \alpha]$ is an element in $\Omega^2(\Gamma(\text{ad}(S_\theta^7)))$,

$$[\nabla_0, \alpha]_{ij} = d\alpha_{ij} + \omega_{ik}\alpha_{kj} - \alpha_{ik}\omega_{kj} \quad (5.46)$$

and has vanishing trace, due to the fact that $\omega_{ik}\alpha_{kj} = \alpha_{kj}\omega_{ik}$ (cf. equations (5.1), (5.2) and the related discussion).

If the family were obtained from an infinitesimal gauge transformation, we would have had $\alpha = [\nabla_0, X]$, for some $X \in \Gamma(\text{ad}(S_\theta^7))$. Indeed, $[\nabla_0, X]$ is an element in $\Omega^1(\text{ad}(S_\theta^7))$ and $P_-[\nabla_0, [\nabla_0, X]] = [P_-F_0, X] = 0$, since F_0 is self-dual. Hence, we have defined an element in the first cohomology group H^1 of the so-called *self-dual complex*:

$$0 \rightarrow \Omega^0(\text{ad}(S_\theta^7)) \xrightarrow{d_0} \Omega^1(\text{ad}(S_\theta^7)) \xrightarrow{d_1} \Omega^2_-(\text{ad}(S_\theta^7)) \rightarrow 0 \quad (5.47)$$

where $\Omega^0(\text{ad}(S_\theta^7)) = \Gamma(\text{ad}(S_\theta^7))$ and $d_0 = [\nabla_0, \cdot]$ and $d_1 := P_-[\nabla_0, \cdot]$. Note that these operators are Fredholm operators, so that the cohomology groups of the complex are finite dimensional. As usual, the complex can be replaced by a single Fredholm operator

$$d_0^* + d_1 : \Omega^1(\text{ad}(S_\theta^7)) \rightarrow \Omega^0(\text{ad}(S_\theta^7)) \oplus \Omega^2_-(\text{ad}(S_\theta^7)), \quad (5.48)$$

where d_0^* is the adjoint of d_0 with respect to the inner product (3.28).

Our goal is to compute $h^1 = \dim H^1$ – the number of “true” gauge instantons. This is achieved by calculating the alternating sum $h^0 - h^1 + h^2$ of Betti numbers from the index of this Fredholm operator,

$$\text{index}(d_0^* + d_1) = -h^0 + h^1 - h^2, \quad (5.49)$$

while showing that $h^0 = h^2 = 0$.

By definition, H^0 consists of the covariant constant elements in $\Gamma(\text{ad}(S_\theta^7))$. Since $[\nabla_0, \cdot]$ commutes with the action of \mathbb{T}^2 and coincides with $\nabla_0^{(2)}$ on $\Gamma(\text{ad}(S_\theta^7))$ (cf. Remark 21), being covariantly constant means

$$[\nabla_0, X] = \nabla_0^{(2)}(X) = 0. \quad (5.50)$$

If we write once more $X = L_\theta(X^{(0)})$ in terms of its classical counterpart, we find that this condition entails

$$\nabla_0^{(2)}(L_\theta(X^{(0)})) = L_\theta(\nabla_0^{(2)}(X^{(0)})) = 0 \quad (5.51)$$

since $\nabla_0^{(2)}$ commutes with L_θ (cf. Remark 21). Since for the undeformed case, there are no covariant constant elements in $\Gamma(\text{ad}(S^7))$ for an irreducible self-dual connection on \mathcal{E} , we conclude that $h^0 = 0$. A completely analogous argument for the kernel of the operator d_1^* shows that also $h^2 = 0$.

5.6 Dirac operator associated to the complex

The next step consists in computing the index of the Fredholm operator $d_0^* + d_1$ defined in (5.48). Firstly, it can be replaced by a Dirac operator on the spinor bundle \mathcal{S} with coefficients in the adjoint bundle. For this, we need the following lemma, which is a straightforward modification of its classical analogue [6]. Recall that the \mathbb{Z}^2 -grading γ_5 induces a decomposition of the spinor bundle $\mathcal{S} = \mathcal{S}^+ \oplus \mathcal{S}^-$. Note that \mathcal{S}^- coincides classically with the charge -1 instanton bundle. Indeed, the Levi-Civita connection – when lifted to the spinor bundle and restricted to negative chirality spinors – has self-dual curvature. Similarly, \mathcal{S}^+ coincides with the charge $+1$ instanton bundle. Note that then Remark 13 implies that the $C^\infty(S^4)$ -modules $\Gamma(S^4, \mathcal{S}^\pm)$ have the homogeneous decomposition property of Definition 6. We conclude from \mathbb{T}^2 -equivariance that also $\Gamma(S_\theta^4, \mathcal{S}^-)$ is isomorphic to the charge -1 instanton bundle $\Gamma(S_{\theta'}^7 \times_{\text{SU}(2)} \mathbb{C}^2)$ on S_θ^4 . Similarly $\Gamma(S_\theta^4, \mathcal{S}^+)$ is isomorphic to the charge $+1$ instanton bundle.

Lemma 31. *There are the following isomorphisms of right $C^\infty(S_\theta^4)$ -modules:*

$$\begin{aligned} \Omega^1(S_\theta^4) &\simeq \Gamma(S_\theta^4, \mathcal{S}^+ \otimes \mathcal{S}^-) \simeq \Gamma(S_\theta^4, \mathcal{S}^+) \otimes_{C^\infty(S_\theta^4)} \Gamma(S_\theta^4, \mathcal{S}^-) \\ \Omega^0(S_\theta^4) \oplus \Omega_-^2(S_\theta^4) &\simeq \Gamma(S_\theta^4, \mathcal{S}^- \otimes \mathcal{S}^-) \simeq \Gamma(S_\theta^4, \mathcal{S}^-) \otimes_{C^\infty(S_\theta^4)} \Gamma(S_\theta^4, \mathcal{S}^-) \end{aligned}$$

Proof. Since this is true in the undeformed case, we have to establish equivariance of the isomorphisms under the action of \mathbb{T}^2 . The result then follows from Lemma 7. By definition, the action of \mathbb{T}^2 on S^4 is lifted to an action of $\widetilde{\mathbb{T}}^2$ on the spinor bundle \mathcal{S} such that it coincides on the endomorphism bundle $\text{End}(\mathcal{S}) \simeq \mathcal{S}^+ \otimes \mathcal{S}^-$ with the underlying action of \mathbb{T}^2 on the cotangent bundle. A analogous statement holds for the second isomorphism. \square

Let us forget for the moment the adjoint bundle $\text{ad}(S_{\theta'}^7)$. Since $\Omega(S_\theta^4) \simeq \Omega(S^4)$ as vector spaces and both d and $*$ commute with the action of \mathbb{T}^2 , the operator $d^* + P_- d$ can be understood as a map from $\Omega^1(S^4) \rightarrow \Omega^0(S^4) \oplus \Omega_-^2(S^4)$, (see Section 3.4). Under the isomorphisms of the above Lemma, this operator is replaced [6] by the Dirac operator with coefficients in \mathcal{S}^- , i.e.

$$D' : \Gamma(S_\theta^4, \mathcal{S}^+ \otimes \mathcal{S}^-) \rightarrow \Gamma(S_\theta^4, \mathcal{S}^- \otimes \mathcal{S}^-). \quad (5.52)$$

Twisting by the adjoint bundle, merely results into a composition with the projection $p_{(2)}$ defining the adjoint bundle $\text{ad}(S_{\theta'}^7)$. Hence, the operator $d_0^* + d_1$ is replaced by the Dirac operator

$$\mathcal{D} : \Gamma(S_\theta^4, \mathcal{S}^+ \otimes \mathcal{S}^- \otimes \text{ad}(S_{\theta'}^7)) \rightarrow \Gamma(S_\theta^4, \mathcal{S}^- \otimes \mathcal{S}^- \otimes \text{ad}(S_{\theta'}^7)), \quad (5.53)$$

with coefficients in the vector bundle $\mathcal{S}^- \otimes \text{ad}(S_{\theta'}^7)$ on S_θ^4 .

We have finally arrived to the computation of the index of this Dirac operator by means of the Connes-Moscovici local index formula. It is given by the pairing,

$$\text{index}(\mathcal{D}) = \langle \phi^*, \text{ch}(\mathcal{S}^- \otimes \text{ad}(S_{\theta'}^7)) \rangle = \langle \phi^*, \text{ch}(\mathcal{S}^-) \cdot \text{ch}(\text{ad}(S_{\theta'}^7)) \rangle. \quad (5.54)$$

In the Appendix we recall the expression for both the cyclic cocycle ϕ^* and the Chern characters, as well as their realization as operators $\pi_D(\text{ch}(\mathcal{E}))$ on the Hilbert space \mathcal{H} .

In [31], we computed these operators for all modules associated to the noncommutative principal bundle $S_{\theta'}^7 \rightarrow S_\theta^4$. In particular, for the adjoint bundle we found that

$$\pi_D(\text{ch}_0(\text{ad}(S_{\theta'}^7))) = 3, \quad \pi_D(\text{ch}_1(\text{ad}(S_{\theta'}^7))) = 0, \quad \pi_D(\text{ch}_2(\text{ad}(S_{\theta'}^7))) = 4(3\gamma_5).$$

To compute the Chern character of the spinor bundle \mathcal{S}^- we use its mentioned identification with the charge -1 instanton bundle $\Gamma(S_{\theta'}^7 \times_{\text{SU}(2)} \mathbb{C}^2)$ on S_θ^4 . It then follows from [19] (cf. also [31]) that

$$\pi_D(\text{ch}_0(\mathcal{S}^-)) = 2, \quad \pi_D(\text{ch}_1(\mathcal{S}^-)) = 0, \quad \pi_D(\text{ch}_2(\mathcal{S}^-)) = -3\gamma_5.$$

Combining both Chern characters and using the local index formula on S_θ^4 , we find that

$$\text{index}(\mathcal{D}) = 6 \underset{z=0}{\text{Res}} z^{-1} \text{tr}(\gamma_5 |D|^{-2z}) + 0 + \frac{1}{2}(2 \cdot 4 - 3 \cdot 1) \underset{z=0}{\text{Res}} \text{tr}(3\gamma_5^2 |D|^{-4-2z}). \quad (5.55)$$

with D which is the classical Dirac operator on S^4 (recall that we do not change it in the isospectral deformation). Now, the first term vanishes due to the fact that $\text{index}(D) = 0$. On the other hand $\gamma_5^2 = \mathbb{I}_4$, and

$$3 \underset{z=0}{\text{Res}} \text{tr}(|D|^{-4-2z}) = 6 \text{Tr}_\omega(|D|^{-4}) = 2,$$

since the Dixmier trace of $|D|^{-m}$ on the m -sphere equals $8/m!$ (cf. for instance [26, 29]). We conclude that $\text{index}(\mathcal{D}) = 5$ and for the moduli space of instantons on S_θ^4 , we have the following

Theorem 32. *The tangent space at the base point ∇_0 to the moduli space of (irreducible) $\text{SU}(2)$ -instantons on S_θ^4 is five-dimensional.*

6 Towards Yang-Mills theory on M_θ

In this final Section, we shall briefly describe how the just constructed Yang-Mills theory on S_θ^4 can be generalized to any four-dimensional toric noncommutative manifold M_θ .

With G a semisimple Lie group, let $P \rightarrow M$ be a principal G bundle on M . We take M to be a four-dimensional Riemannian manifold equipped with an isometrical action σ of the torus \mathbb{T}^2 . For the construction to work, we assume that this action can be lifted to an action $\tilde{\sigma}$ of a cover $\tilde{\mathbb{T}}^2$ on P , while it commutes with the action of G . As in Section 3, we define the noncommutative algebras $C^\infty(P_\theta)$ and $C^\infty(M_\theta)$ as the vector spaces $C^\infty(P)$ and $C^\infty(M)$ with star products defined like in (3.2) with respect to the action of $\tilde{\mathbb{T}}^2$ and \mathbb{T}^2 respectively; or, equivalently as the images of $C^\infty(P)$ and $C^\infty(M)$ under the corresponding quantization map L_θ . Since the action of $\tilde{\mathbb{T}}^2$ is taken to commute with the action of G on P , the corresponding action α of G on the algebra $C^\infty(P)$ by

$$\alpha_g(f)(p) = f(g^{-1} \cdot p) \quad (6.1)$$

induces an action of G by automorphisms on the algebra $C^\infty(P_\theta)$. This means that also the inclusion $C^\infty(M) \subset C^\infty(P)$ as G -invariant elements in $C^\infty(P)$ extends to an inclusion $C^\infty(M_\theta) \subset C^\infty(P_\theta)$ of G -invariant element in $C^\infty(P_\theta)$. Clearly, the action of G translates trivially into a coaction of the Hopf algebra $C^\infty(G)$ on $C^\infty(P_\theta)$.

Proposition 33. *The inclusion $C^\infty(M_\theta) \hookrightarrow C^\infty(P_\theta)$ is a (principal) Hopf-Galois $C^\infty(G)$ extension.*

Proof. As in [31], it is enough to establish surjectivity of the canonical map

$$\begin{aligned} \chi : C^\infty(P_\theta) \otimes_{C^\infty(M_\theta)} C^\infty(P_\theta) &\rightarrow C^\infty(P_\theta) \otimes C^\infty(G), \\ f' \otimes_{C^\infty(M_\theta)} f &\mapsto f' \Delta_R(f) = f' f_{(0)} \otimes f_{(1)} \end{aligned}$$

All additional nice properties would then follow from the cosemisimplicity of the Hopf algebra $C^\infty(G)$. Note that in the classical case, the bijectivity of the canonical map $\chi^{(0)} : C^\infty(P) \otimes_{C^\infty(M)} C^\infty(P) \rightarrow C^\infty(P) \otimes C^\infty(G)$ follows by the very definition of a principal bundle. Furthermore, there is an isomorphism of vector spaces:

$$\begin{aligned} T : C^\infty(P_\theta) \otimes_{C^\infty(M_\theta)} C^\infty(P_\theta) &\rightarrow C^\infty(P) \otimes_{C^\infty(M_\theta)} C^\infty(P_\theta) \\ f' \otimes_{C^\infty(M_\theta)} f &\mapsto \sum_r f'_r \otimes_{C^\infty(M)} \tilde{\sigma}_{r\theta}(f) \end{aligned}$$

where $f' = \sum_r f'_r$ is the homogeneous decomposition of f' under the action of $\tilde{\mathbb{T}}^2$ – compare with the proof of Lemma 7. We claim that the canonical map is given as the composition $\chi = \chi^{(0)} \circ T$, hence, it is bijective. Indeed,

$$\begin{aligned} \chi^{(0)} \circ T(f' \otimes_{C^\infty(M_\theta)} f) &= \sum_r f'_r \tilde{\sigma}_{r\theta}(f_{(0)}) \otimes f_{(1)} \\ &= f' \times_\theta f_{(0)} \otimes f_{(1)} \\ &= \chi(f' \otimes_{C^\infty(M_\theta)} f) \end{aligned}$$

since the action of $\tilde{\mathbb{T}}^2$ on $C^\infty(P_\theta)$ commutes with the coaction of $C^\infty(G)$. □

Noncommutative associated bundles are defined as before by setting

$$\mathcal{E} = C^\infty(P_\theta) \boxtimes_\rho V := \{f \in C^\infty(P_\theta) \otimes V \mid (\alpha_g \otimes \text{id})(f) = (\text{id} \otimes \rho(g)^{-1})(f)\}$$

for a representation ρ of G on V . These $C^\infty(M_\theta)$ bimodules are finite projective since they are of the form of the modules defined in Section 3.3 (cf. Remark 13). Moreover, the corresponding classical $C^\infty(M)$ -modules $C^\infty(P) \boxtimes_\rho V$ have the homogeneous decomposition property. Indeed, let $f \in C^\infty(P) \boxtimes_\rho V$ and write in a basis of V as

$$f = (f^1, \dots, f^n), \tag{6.2}$$

so that $f^i \in C^\infty(P)$ satisfies $\alpha_g(f^i) = \rho(g)_{ij} f^j$. Suppose now that $f^i = \sum_r f_r^i$ as a sum of $\tilde{\mathbb{T}}^2$ -homogeneous elements in $C^\infty(P)$: $\sigma_t(f_r^i) = e^{2\pi i t \cdot r} f_r^i$ for $t \in \tilde{\mathbb{T}}^2$. Since $\alpha_g \circ \sigma_t = \sigma_t \circ \alpha_g$, we find that

$$\alpha_g\left(\sum_r e^{2\pi i t \cdot r} f_r^i\right) = \sum_r \rho(g)_{ij} e^{2\pi i t \cdot r} f_r^j. \tag{6.3}$$

for $t \in \tilde{\mathbb{T}}^2$ and $g \in G$. By linear independence of the exponentials, we derive that $\alpha_g(f_r^i) = \rho(g)_{ij} f_r^j$, and conclude that $C^\infty(P) \boxtimes_\rho V$ has the homogeneous decomposition property.

Moreover, Proposition 14 generalizes and reading $\text{End}(\mathcal{E}) \simeq C^\infty(P_\theta) \boxtimes_{\text{ad}} L(V)$, where ad is the adjoint representation of G on $L(V)$. Also, one identifies the adjoint bundle as the module coming from the adjoint representation of G on $\mathfrak{g} \subset L(V)$, namely $\Gamma(\text{ad}(P_\theta)) := C^\infty(P_\theta) \boxtimes_{\text{ad}} \mathfrak{g}$.

For a (right) finite projective $C^\infty(M_\theta)$ -module \mathcal{E} we define an inner product $(\cdot, \cdot)_2$ on $\text{End}_{C^\infty(M_\theta)}(\mathcal{E}, \mathcal{E} \otimes_{C^\infty(M_\theta)} \Omega(M_\theta))$ as in Section 4.3. The Yang-Mills action functional for a connection ∇ on \mathcal{E} in terms of its curvature F is then defined by

$$\text{YM}(\nabla) = (F, F)_2. \tag{6.4}$$

This is a gauge invariant, positive and quartic functional. The derivation of the Yang-Mills equations (4.30) on S_θ^4 does not rely on the specific properties of S_θ^4 and continues to hold on M_θ . The same is true for the topological action, and $\text{YM}(\nabla) \geq \text{Top}(\mathcal{E})$ with equality iff $*_\theta F = \pm F$. In other words, instanton connections are minima of the Yang-Mills action.

The explicit construction of instanton connections on S_θ^4 carried over in Section 5 can of course not be generalized to the manifold M_θ . However, local expressions could in principle be obtained on a ‘‘local chart’’ \mathbb{R}_θ^4 of M_θ , if \mathbb{T}^2 acts on the corresponding local chart \mathbb{R}^4 of M . On the other hand, the infinitesimal construction of instantons on M_θ giving the dimension of the ‘tangent of the moduli space’ can be generalized to any toric noncommutative manifold M_θ , again following closely [6]. Essential here is the existence of a ‘base point’, i.e. an instanton connection that can be linearly perturbed to obtain a family of infinitesimal instantons.

7 Final remarks

Twisting of algebras and coalgebras has been known for some time [23, 25] with interesting consequences, e.g. for the nonassociativity of differential calculi [8]. The use of a twist to implement symmetries of the toric noncommutative manifolds used in the present paper was made explicit in [41]. For symmetries of the usual noncommutative planes and their use for quantum field theories on it, one has the approach of [43, 37, 44]. More recently [10, 45], a twist was used to implement Poincaré symmetry on the Moyal plane while conformal transformations are twisted in [35]. Twisting of infinitesimal diffeomorphisms and their use for gravity theories are in [1, 3] and infinite dimensional (infinitesimal) conformal symmetries on a two dimensional Moyal plane are twisted in [33]. There are also studies of spin and statistics and their relations in the context of these twisted symmetries of the Moyal plane [7].

A Local index formula

Suppose in general that $(\mathcal{A}, \mathcal{H}, D, \gamma)$ is an even p -summable spectral triple with discrete simple dimension spectrum. For a projection $e \in M_N(\mathcal{A})$, the operator

$$D_e = e(D \otimes \mathbb{I}_N)e$$

is a Fredholm operator, thought of as the Dirac operator with coefficient in the module determined by e . Then the local index formula of Connes and Moscovici [20] provides a method to compute its index via the pairing of suitable cyclic cycles and cocycles. We shall recall the ‘even’ case since it is the one that is relevant for the present paper.

Let $C_*(\mathcal{A})$ be the complex consisting of cycles over the algebra \mathcal{A} , that is in degree n , $C_n(\mathcal{A}) := \mathcal{A}^{\otimes(n+1)}$. On this complex there are defined the Hochschild operator $b : C_n(\mathcal{A}) \rightarrow C_{n-1}(\mathcal{A})$ and the boundary operator $B : C_n(\mathcal{A}) \rightarrow C_{n+1}(\mathcal{A})$, satisfying $b^2 = 0, B^2 = 0, bB + Bb = 0$; thus $(b+B)^2 = 0$. From general homological theory, one defines a bicomplex $CC_*(\mathcal{A})$ by $CC_{(n,m)}(\mathcal{A}) := CC_{n-m}(\mathcal{A})$ in bi-degree (n, m) . Dually, one defines $CC^*(\mathcal{A})$ as functionals on $CC_*(\mathcal{A})$, equipped with the dual Hochschild operator b and coboundary operator B (we refer to [15] and [34] for more details on this).

Theorem 34 (Connes-Moscovici [20]).

(a) An even cocycle $\phi^* = \sum_{k \geq 0} \phi^k$ in $CC^*(\mathcal{A})$, $(b+B)\phi^* = 0$, defined by the following formulæ. For $k = 0$,

$$\phi^0(a) := \operatorname{Res}_{z=0} z^{-1} \operatorname{tr}(\gamma a |D|^{-2z}); \quad (\text{A.1})$$

whereas for $k \neq 0$

$$\phi^{2k}(a^0, \dots, a^{2k}) := \sum_{\alpha} c_{k,\alpha} \operatorname{Res}_{z=0} \operatorname{tr}(\gamma a^0 [D, a^1]^{(\alpha_1)} \dots [D, a^{2k}]^{(\alpha_{2k})} |D|^{-2(|\alpha|+k+z)}) \quad (\text{A.2})$$

where

$$c_{k,\alpha} = (-1)^{|\alpha|} \Gamma(k + |\alpha|) (\alpha! (\alpha_1 + 1)(\alpha_1 + \alpha_2 + 2) \dots (\alpha_1 + \dots + \alpha_{2k} + 2k))^{-1}$$

and $T^{(j)}$ denotes the j 'th iteration of the derivation $T \mapsto [D^2, T]$.

(b) For $e \in K_0(\mathcal{A})$, the Chern character $\operatorname{ch}_*(e) = \sum_{k \geq 0} \operatorname{ch}_k(e)$ is the even cycle in $CC_*(\mathcal{A})$, $(b+B)\operatorname{ch}_*(e) = 0$, defined by the following formulæ. For $k = 0$,

$$\operatorname{ch}_0(e) := \operatorname{tr}(e); \quad (\text{A.3})$$

whereas for $k \neq 0$

$$\operatorname{ch}_k(e) := (-1)^k \frac{(2k)!}{k!} \sum (e_{i_0 i_1} - \frac{1}{2} \delta_{i_0 i_1}) \otimes e_{i_1 i_2} \otimes e_{i_1 i_2} \otimes \dots \otimes e_{i_{2k} i_0}. \quad (\text{A.4})$$

(c) The index is given by the natural pairing between cycles and cocycles

$$\operatorname{index} D_e = \langle \phi^*, \operatorname{ch}_*(e) \rangle. \quad (\text{A.5})$$

The components of the Chern character are represented as operators on the Hilbert space \mathcal{H} by explicit formulæ,

$$\pi_D(\text{ch}_k(e)) = (-1)^k \frac{(2k)!}{k!} \sum (\pi(e_{i_0 i_1}) - \frac{1}{2} \delta_{i_0 i_1}) [D, \pi(e_{i_1 i_2})] [D, \pi(e_{i_2 i_3})] \cdots [D, \pi(e_{i_{2k} i_0})]. \quad (\text{A.6})$$

In the case of the toric noncommutative manifolds introduced in [19] (cf. Section 3), the local index formula simplifies drastically [31].

Theorem 35. *For a projection $p \in M_N(C^\infty(M_\theta))$, we have*

$$\text{index } D_p = \text{Res}_{z=0} z^{-1} \text{tr} \left(\gamma p |D|^{-2z} \right) + \sum_{k \geq 1} c_k \text{Res}_{z=0} \text{tr} \left(\gamma \left(p - \frac{1}{2} \right) [D, p]^{2k} |D|^{-2(k+z)} \right)$$

where $c_k = (k-1)!/(2k)!$.

Proof. As noted before, the twist L_θ commutes with the action α_s of $\widetilde{\mathbb{T}}^n$ on an operator T . In fact, if T is homogeneous of degree r , then $L_\theta(T)$ is of degree r :

$$\alpha_s(L_\theta(T)) = U(s) T U(r') U(s)^{-1} = U(s) T U(s)^{-1} U(r') = e^{2\pi i s_\mu r_\mu} L_\theta(T).$$

with $r'_\nu = r_\mu \theta_{\mu\nu}$ so that $r' \in \widetilde{\mathbb{T}}^n$.

We write the cocycles ϕ^{2k} that define the local index formula in (A.2) in terms of the twist L_θ :

$$\begin{aligned} \phi^{2k}(L_\theta(f^0), L_\theta(f^1), \dots, L_\theta(f^{2k})) = \\ \text{Res}_{z=0} \text{tr} \left(\gamma L_\theta(f^0 \times_\theta [D, f^1]^{(\alpha_1)} \cdots \times_\theta [D, f^{2k}]^{(\alpha_{2k})}) |D|^{-2(|\alpha|+k+z)} \right) \end{aligned} \quad (\text{A.7})$$

where we extended the \times_θ -product to $C^\infty(M_\theta) \cup [D, C^\infty(M_\theta)]$ which can be done unambiguously since D is of degree 0. Suppose now that $f^0, \dots, f^{2k} \in C^\infty(M)$ are homogeneous of degree r^0, \dots, r^{2k} , respectively, under the action of \mathbb{T}^n , so that the operator $f^0 \times_\theta [D, f^1] \cdots \times_\theta [D, f^{2k}]$ is a homogeneous element of degree r (which can be expressed in terms of the r^i). It is in fact a multiple of $f^0 [D, f^1] \cdots [D, f^{2k}]$ by working out the \times_θ -product. Forgetting about this factor – which is a power of λ – we obtain from (3.6)

$$L_\theta(f^0 [D, f^1] \cdots [D, f^{2k}]) = f^0 [D, f^1] \cdots [D, f^{2k}] U(r_\mu \theta_{\mu 1}, \dots, r_\mu \theta_{\mu n}). \quad (\text{A.8})$$

Each term in the local index formula for $(C^\infty(M_\theta), \mathcal{H}, D)$ then takes the form

$$\text{Res}_{z=0} \text{tr} \left(\gamma f^0 [D, f^1]^{(\alpha_1)} \cdots [D, f^{2k}]^{(\alpha_{2k})} |D|^{-2(|\alpha|+k+z)} U(s) \right)$$

for $s_\nu = r_\mu \theta_{\mu\nu}$ so that $s \in \mathbb{T}^n$. The appearance of $U(s)$ here, is a consequence of the close relation with the index formula for a \mathbb{T}^n -equivariant Dirac spectral triple on M . In [12], Chern and Hu considered an even dimensional compact spin manifold M on which a (connected compact) Lie group G acts by isometries. The equivariant Chern character was defined as an equivariant version of the JLO-cocycle, the latter being an element in equivariant entire cyclic cohomology. The essential point is that they obtained an explicit formula for the above residues. In the case of the previous \mathbb{T}^n -action on M , one gets

$$\begin{aligned} \text{Res}_{z=0} \text{tr} \left(\gamma f^0 [D, f^1]^{(\alpha_1)} \cdots [D, f^{2k}]^{(\alpha_{2k})} |D|^{-2(|\alpha|+k+z)} U(s) \right) \\ = \Gamma(|\alpha| + k) \lim_{t \rightarrow 0} t^{|\alpha|+k} \text{tr} \left(\gamma f^0 [D, f^1]^{(\alpha_1)} \cdots [D, f^{2k}]^{(\alpha_{2k})} e^{-tD^2} U(s) \right) \end{aligned} \quad (\text{A.9})$$

for every $s \in \mathbb{T}^n$; moreover, this limit vanishes when $|\alpha| \neq 0$ (Thm 2 in [12]). \square

References

- [1] P. Aschieri, C. Blohmann, M. Dimitrijevic, F. Meyer, P. Schupp, J. Wess. A Gravity Theory on Noncommutative Spaces. *Class. Quant. Grav.* 22 (2005) 3511–3532.
- [2] P. Aschieri and F. Bonechi. On the noncommutative geometry of twisted spheres. *Lett. Math. Phys.* 59 (2002) 133–156.
- [3] P. Aschieri, M. Dimitrijevic, F. Meyer, J. Wess. Noncommutative Geometry and Gravity. hep-th/0510059.
- [4] M. F. Atiyah. *The Geometry of Yang-Mills Fields*. Fermi Lectures, Scuola Normale Pisa, 1979.
- [5] M. F. Atiyah, N. J. Hitchin, V. G. Drinfel'd and Yu. I. Manin. Construction of instantons. *Phys. Lett.* A65 (1978) 185–187.
- [6] M. F. Atiyah, N. J. Hitchin, and I. M. Singer. Self-duality in four-dimensional Riemannian geometry. *Proc. Roy. Soc. Lond.* A362 (1978) 425–461.
- [7] A.P. Balachandran, G. Mangano, A. Pinzul, S. Vaidya. Spin and Statistics on the Groenewold-Moyal Plane: Pauli-Forbidden Levels and Transitions. hep-th/0508002.
- [8] E.J. Beggs, S. Majid. Quantization by cochain twists and nonassociative differentials. math.QA/0506450.
- [9] A. Belavin, A. Polyakov, A. Schwarz, Y. Tyupkin. Pseudoparticle Solutions of the Yang-Mills Equations, *Phys. Lett.* 59B (1975) 85–87.
- [10] M. Chaichian, P. Kulish, K. Nishijima, A. Tureanu. On a Lorentz-Invariant Interpretation of Noncommutative Space-Time and Its Implications on Noncommutative QFT. *Phys. Lett.* B604 (2004) 98–102.
- [11] V. Chari and A. Pressley. *A guide to quantum groups*. Cambridge University Press, Cambridge, 1994.
- [12] S. Chern and X. Hu. Equivariant Chern character for the invariant Dirac operator. *Michigan Math. J.* 44 (1997) 451–473.
- [13] A. Connes. C^* -algèbres et géométrie différentielle. *C.R. Acad. Sci. Paris Ser. A-B*, 290 (1980) A599–A604.
- [14] A. Connes. Noncommutative differential geometry. *IHES Sci. Publ. Math.* 62 (1985) 257–360.
- [15] A. Connes. *Noncommutative Geometry*. Academic Press, San Diego, 1994.
- [16] A. Connes. Gravity coupled with matter and the foundation of noncommutative geometry. *Commun. Math. Phys.* 182 (1996) 155–176.
- [17] A. Connes. Noncommutative geometry and reality. *J. Math. Phys.* 36 (1995) 6194–6231.

- [18] A. Connes and M. Dubois-Violette. Noncommutative finite-dimensional manifolds. I. Spherical manifolds and related examples. *Commun. Math. Phys.* 230 (2002) 539–579.
- [19] A. Connes and G. Landi. Noncommutative manifolds: The instanton algebra and isospectral deformations. *Commun. Math. Phys.* 221 (2001) 141–159.
- [20] A. Connes and H. Moscovici. The local index formula in noncommutative geometry. *Geom. Funct. Anal.* 5 (1995) 174–243.
- [21] J. Cuntz and D. Quillen. Algebra extension and nonsingularity. *J. Amer. Math. Soc.* 8 (1995) 251–289.
- [22] J. Dixmier. Existence de traces non normales. *C.R. Acad. Sci. Paris Sér A-B*, 262 (1966) A1107–A1108.
- [23] V.G. Drinfel’d. Quasi-Hopf algebras. *Leningrad. Math. J.* 1 (1990) 1419–1457.
- [24] V. Gayral, B. Iochum, and J. C. Várilly. Dixmier traces on noncompact isospectral deformations. [hep-th/0507206](#).
- [25] A. Giaquinto, J.J.Zhang. Bialgebra actions, twists and universal deformation formulas. [hep-th/9411140](#).
- [26] J. M. Gracia-Bondía, J. C. Várilly, and H. Figueroa. *Elements of Noncommutative Geometry*. Birkhäuser, Boston, 2001.
- [27] P. Julg. K-théorie équivariante et produits croisés. *C.R. Acad. Sci. Paris*, 292 (1981) 629–632.
- [28] G. Landi. Spin-Hall effect with quantum group symmetry. [hep-th/0504092](#). *Lett. Math. Phys.* in press.
- [29] G. Landi. *An Introduction to Noncommutative Spaces and their Geometry*. Springer-Verlag, 1997.
- [30] G. Landi, C. Pagani, C. Reina, and W. van Suijlekom. work in progress.
- [31] G. Landi and W. van Suijlekom. Principal fibrations from noncommutative spheres. *Commun. Math. Phys.* 260 (2005) 203–225.
- [32] S. Lie. *Theorie der Transformationsgruppen*. Chelsea, New York, 1970.
- [33] F. Lizzi, S. Vaidya, P. Vitale. Twisted Conformal Symmetry in Noncommutative Two-Dimensional Quantum Field Theory. [hep-th/0601056](#).
- [34] J.-L. Loday. *Cyclic Homology*. Springer-Verlag, Berlin, 1992.
- [35] P. Matlock, Non-Commutative Geometry and Twisted Conformal Symmetry. *Phys. Rev. D* 71 (2005) 126007.

- [36] N. Nekrasov, A. Schwarz. Instantons on noncommutative R^4 , and (2,0) superconformal six dimensional theory. Commun. Math. Phys. 198 (1998) 689–703.
- [37] R. Oeckl. Untwisting noncommutative \mathbb{R}^g and the equivalence of quantum field theories. Nucl. Phys. B581 (2000) 559–574.
- [38] M. A. Rieffel. Non-commutative tori - A case study of non-commutative differentiable manifolds. Contemp. Math. 105 (1990) 191–212.
- [39] M.A. Rieffel. *Deformation Quantization for Actions of \mathbb{R}^d* . Memoirs of the Amer. Math. Soc. 506, Providence, RI, 1993.
- [40] M. A. Rieffel. K -groups of C^* -algebras deformed by actions of R^d . J. Funct. Anal. 116 (1993) 199–214.
- [41] A. Sitarz. Twists and spectral triples for isospectral deformations. Lett. Math. Phys. 58 (2001) 69–79.
- [42] J. C. Várilly. Quantum symmetry groups of noncommutative spheres. Commun. Math. Phys. 221 (2001) 511–523.
- [43] P. Watts. Noncommutative string theory, the R-matrix, and Hopf algebras. Phys. Lett. B474 (2000) 295–302.
- [44] P. Watts. Derivative and the role of the Drinfel'd twist in noncommutative string theory. hep-th/0003234.
- [45] J. Wess. Deformed coordinate spaces: derivatives. hep-th/0408080.