2000]Primary 11N25, Secondary 11Y50

ODD PERFECT NUMBERS HAVE AT LEAST NINE DISTINCT PRIME FACTORS

PACE P. NIELSEN

ABSTRACT. An odd perfect number, N, is shown to have at least nine distinct prime factors. If $3 \nmid N$ then N must have at least twelve distinct prime divisors. The proof ultimately avoids previous computational results for odd perfect numbers.

1. INTRODUCTION

A perfect number is one where $\sigma(N) = 2N$. In other words, the sum of the divisors of N is twice N. These numbers have been studied since antiquity. It is known that N is an even perfect number if and only if $N = (2^p - 1)2^{p-1}$ with $2^p - 1$ prime. Sufficiency was proven by Euclid and necessity by Euler. A prime number of the form $2^p - 1$ is called a Mersenne prime, and there are currently 43 known. There is an ongoing, online, distributed search for such primes at http://www.mersenne.org.

The search for odd perfect numbers has not been as successful. Currently there are none known, making their existence the oldest unanswered question in mathematics. However there are a great number of necessary conditions, which go through periodic improvements. The list of conditions given here is the same list as in [25], but with recent improvements included.

Let N be an odd perfect number (if such exists). Write $N = \prod_{i=1}^{k} p_i^{a_i}$ where each p_i is prime, $p_1 < p_2 < \ldots < p_k$, and $k = \omega(N)$ is the number of distinct prime factors. The factors $p_i^{a_i}$ are called the *prime components* of N. Then:

- Eulerian Form: We have $N = \pi^{\alpha} m^2$ for some integers $\pi, \alpha, m \in \mathbb{Z}_+$, with $\pi \equiv \alpha \equiv 1 \pmod{4}$ and π prime. The prime π is called the *special* prime of N.
- Lower Bound: Brent, Cohen, and te Riele [3] using a computer search found that $N > 10^{300}$. William Lipp, using the same techniques, is close to pushing the bound to $N > 10^{500}$, and plans to start a distributed search at the website http://www.oddperfect.org.
- Upper Bound: Dickson [7] proved that there are finitely many odd perfect numbers with a fixed number of distinct prime factors. Pomerance [24] gave an effective bound in terms of k. This was improved in succession by Heath-Brown [12], Cook [6], and finally Nielsen [22] to 2^{4^k} .
- Large Factors: Jenkins [15] proved that $p_k > 10^7$, and Iannucci [13],[14] proved $p_{k-1} > 10^4$ and $p_{k-2} > 10^2$.
- Small Factors: The smallest prime factor satisfies $p_1 < \frac{2}{3}k + 2$ as proved by Grün [8]. For $2 \leq i \leq 6$, Kishore [17] showed that $p_i < 2^{2^{i-1}}(k-i+1)$, and this has been slightly improved by Cohen and Sorli [5].

¹⁹⁹¹ Mathematics Subject Classification. [.

Key words and phrases. abundant, deficient, odd perfect.

- Number of Total Prime Factors: Hare [11] proved that the total number of (not necessarily distinct) prime factors of N must be at least 47. In unpublished work he has improved this to 75.
- Number of Distinct Prime Factors: Chein [4] and Hagis [9] independently proved that $\omega(N) \ge 8$. Hagis [10] and Kishore [18] showed that if $3 \nmid N$ then $\omega(N) \ge 11$. This paper improves both of these bounds by 1.
- The Exponents: For the non-special primes, p_i , write $a_i = 2b_i$. If $d = \gcd(2b_i + 1)$ then $d \neq 0 \pmod{3}$ by a result of McDaniel [19].

With such a number of conditions, it might seem that an odd perfect number could not exist. Pomerance has given an interesting heuristic, available at Lipp's website, suggesting that odd perfect numbers are very unlikely.

I would like to thank William Lipp for providing some of the factorizations given in the lemmas. Also, much of the terminology, notation, and lemmas of the early sections of this paper match those found in [25] (which in turn match those in [13], and earlier work like [23]) in an effort to establish a sense of both continuity and improvement, and I'd like to thank John Voight for some interesting conversations on topics related to his paper.

2. Fixed Notations and Conventions

Let N be an odd perfect number and $k = \omega(N)$. We will write π for the special prime. By N we mean the non-negative integers and by \mathbb{Z}_+ the positive integers. We will use an algorithm to prove our results. At each stage in the algorithm there will be prime divisors of N that are known and some that are unknown, meaning that the prime divisors are either specified by the algorithm or they are not, respectively. This set of known primes will change at every stage of the algorithm as it runs through different cases, and so the known and unknown primes are constantly changing. A more formal definition will be given in a later section. We let k_1 be the number of known, distinct prime divisors of N (at any given stage), and let $k_2 = k - k_1$ be the number of unknown, distinct prime divisors. Among the known prime divisors of N, some of the prime components are also known (again, known being a technical term meaning specified by the algorithm). In other words, if p is a known prime divisor of N and if our algorithm yields some $a \in \mathbb{Z}_+$ so that $p^a ||N$, we say p^a is a known prime component. We let ℓ_1 be the number of known prime components of N, and let $\ell_2 = k - \ell_1$ be the number of unknown prime components.

A word of warning: In some theorems we will assume p is a prime with $p^a||N$, but this doesn't even mean that p is a known prime, let alone that the prime component is known. Further, even if p is a known prime, the component may still be unknown (in this technical sense) even though we (from our position) are given that $p^a||N$. In other words, we are given the hypothesis $p^a||N$ but the algorithm may not have specified p or a. We will, throughout, only use the words "known" and "unknown" in the sense of known to us *through our algorithm*, rather than by hypothesis.

3. Cyclotomic Integers

The equation $\sigma(N) = 2N$ can be (trivially) rewritten as $\sigma(N)/N = 2$ which tells us that each odd prime divisor of $\sigma(N)$ must somehow divide N, and vice versa.

Thus, we want to study the prime factorization of

$$\sigma(N) = \prod_{i=1}^{k} \sigma(p_i^{a_i}) = \prod_{i=1}^{k} \frac{p_i^{a_i+1} - 1}{p_i - 1}.$$

Letting $\Phi_n(x)$ be the *n*th cyclotomic polynomial (i.e. the minimal polynomial over \mathbb{Q} for a primitive *n*th root of unity), we have the partial factorization

$$p^n - 1 = \prod_{d|n} \Phi_d(p)$$

and so

(1)
$$\sigma(p^{n-1}) = \frac{p^n - 1}{p - 1} = \prod_{d|n, d > 1} \Phi_d(p).$$

We are further interested in the factorization of $\Phi_d(p)$. If c and d are integers with d > 1 and gcd(c, d) = 1 we write $o_d(c)$ for the multiplicative order of c modulo d. If p is prime, we write v_p for the valuation associated to p. In other words, for $n \in \mathbb{Z}_+$ we have $p^{v_p(n)} || n$. The following results of Nagell [21] are fundamental and are often left as exercises in modern abstract algebra books.

Lemma 1 ([25, Lemma 3]). Let m > 1 be an integer, and let q be prime. Write $m = q^b n$ with gcd(q, n) = 1.

If b = 0 then

$$\Phi_m(x) \equiv 0 \pmod{q}$$

is solvable if and only if $q \equiv 1 \pmod{m}$. The solutions are those x with $o_q(x) = m$. Further, $v_q(\Phi_m(x)) = v_q(x^m - 1)$ for such solutions.

If $b \neq 0$ then

$$\Phi_m(x) \equiv 0 \pmod{q}$$

is solvable if and only if $q \equiv 1 \pmod{n}$. The solutions are those x with $o_q(x) = n$. Further, if m > 2 then $v_q(\Phi_m(x)) = 1$ for such solutions.

We have immediately from Lemma 1 and Equation 1:

Lemma 2 ([13, Equation 4],[25, Lemma 4]). Let p and q be primes, $q \ge 3$, and $a \in \mathbb{Z}_+$. Then

$$v_q(\sigma(p^a)) = \begin{cases} v_q(p^{o_q(p)} - 1) + v_q(a+1) & \text{if } o_q(p) | (a+1) \text{ and } o_q(p) \neq 1, \\ v_q(a+1) & \text{if } o_q(p) = 1, \\ 0 & \text{otherwise.} \end{cases}$$

It turns out that the first case of Lemma 1 (when b = 0) is the more common means of obtaining factors for odd perfect numbers. However, to make sure we can always reduce to that case we need the following result, usually attributed to Bang [1], but given other proofs such as in [2].

Lemma 3. Let $m, x \in \mathbb{Z}_+$ with $x \ge 2$. Then $\Phi_m(x)$ is divisible by a prime q with $o_q(x) = m$, except if x = 2 and m = 1 or 6, or if $x = 2^i - 1$ (for some $i \in \mathbb{Z}_+$) and m = 2.

Note that we are only interested in this lemma when x = p is a prime dividing an odd perfect number N. Thus, the case x = 2 never happens. Also, if m = 2this corresponds to the special prime (in the Eulerian form) so $x = \pi \equiv 1 \pmod{4}$ and hence cannot be of the form $2^i - 1$. Thus, both exceptions in the lemma do not affect our work.

Define $\sigma_i(n) = \sum_{d|n} d^i$, for $i \in \mathbb{Z}$ and $n \in \mathbb{Z}_+$. It is clear that each of these functions is multiplicative, $\sigma_1 = \sigma$ is the usual sum of divisors function, and σ_0 is the number of divisors function. The following is immediate:

Lemma 4. Let N be an odd perfect number. If $p^a || N$, where p is prime, then for each d|(a + 1) the number $\Phi_d(p)$ is divisible by a prime q with $o_q(p) = d$ and $q \equiv 1 \pmod{d}$. In particular, $\sigma(p^a)$ has at least $\sigma_0(a + 1) - 1$ distinct prime factors in common with N.

4. Fermat Primes

A prime q is called a *Fermat prime* if it is of the form $q = 2^j + 1$ for some $j \in \mathbb{Z}_+$. One can show it is necessary that $j = 2^i$ for some $i \in \mathbb{N}$. It is easily seen that if i = 0, 1, 2, 3, 4 then $2^{2^i} + 1$ is prime (i.e. q = 3, 5, 17, 257, 65537), but no other Fermat primes are known. These primes play a special role in the study of odd perfect numbers. This is because the prime factorization of q - 1 is exactly a power of 2.

The first thing we can do is restate Lemma 2 in terms of divisors of N, and Fermat primes.

Lemma 5 ([23],[25, Lemma 5]). Let N be an odd perfect number, $p^a || N$ with p prime, and let q be a Fermat prime. Then:

$$v_q(\sigma(p^a)) = \begin{cases} v_q(p+1) + v_q(a+1) & \text{if } \pi = p \equiv -1 \pmod{q}, \\ v_q(a+1) & \text{if } p \equiv 1 \pmod{q}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. First suppose that $p \equiv 1 \pmod{q}$. In this case $o_q(p) = 1$ and so we just use Lemma 2.

Next suppose $p \not\equiv 1 \pmod{q}$. Note $o_q(p)$ must be a divisor of $q - 1 = 2^{2^k}$. However, from the Eulerian form, we see that a + 1 is divisible by 2 if and only if $p = \pi$. Further, in this case $a \equiv 1 \pmod{4}$ and hence 2||(a+1). Thus $v_q(\sigma(p^a)) = 0$ unless $p = \pi$ and $o_q(p) = 2$. But this last equality is equivalent to $p \equiv -1 \pmod{q}$. Lemma 2 gives us the needed valuation in this case.

Lemma 5 tells us that there are only a few sources in $\sigma(N)$ for copies of q. In particular, if $q^n | N$ for some large n, we can force the size of the special prime to be large. To prove this we first need another lemma.

Lemma 6 ([25, Proposition 9]). Let N be an odd perfect number, let q be a Fermat prime, and suppose $p^a || N$ with p prime.

- (i) If $p \neq \pi$ and $q^b | \sigma(p^a)$ then $\sigma(p^a)$ is divisible by b distinct primes r_1, r_2, \ldots, r_b with $r_i \equiv 1 \pmod{q^i}$.
- (ii) If $p = \pi$, $p \equiv -1 \pmod{q}$, and $q^c | (a+1)$, then $\sigma(p^a)$ is divisible by 2c distinct primes $r_1, r'_1, \ldots, r_c, r'_c$ with $r_i \equiv r'_i \equiv 1 \pmod{q^i}$.

Proof. For part (i), by Lemma 5 we have $q^b|(a+1)$. So we take r_i to be the divisor of $\Phi_{q^i}(p)$ specified by Lemma 4. Part (ii) follows from the same lemmas, noticing that since $p = \pi$ is special we have 2|(a+1), and hence we can take r_i and r'_i to be the factors specified by Lemma 4 of $\Phi_{q^i}(p)$ and $\Phi_{2q^i}(p)$ respectively.

Proposition 7 (c.f. [25, Proposition 10]). Let N be an odd perfect number, and let q be a Fermat prime with $q^n|N$. Suppose k, k_1 , k_2 , ℓ_1 , and ℓ_2 have their usual meanings. Further suppose $q^b||\sigma(known prime components of N)$. Finally, let k'_1 (respectively ℓ'_1) be the number of distinct prime factors among the k_1 known prime divisors (respectively, among the ℓ_1 known prime components) which are congruent to 1 (mod q).

If

$$\tau = n - b - (k_1' - \ell_1' + k_2)(k_1' + k_2 - 1) > 0$$

then $\pi \equiv -1 \pmod{q}$ and π is among the unknown prime components. If, further, each known prime, p, with unknown component and with $p \equiv 1 \pmod{4}$ satisfies $p \not\equiv -1 \pmod{q^{\tau}}$, then π is among the unknown primes, and

$$v_q(\pi+1) \ge n - b - (k_1' - \ell_1' + k_2 - 1)(k_1' + k_2 - 2) - \lfloor (k_1' + k_2 - 1)/2 \rfloor = \tau'.$$

Proof. First, we will show the contrapositive of the initial statement. Suppose π is among the known components, or if it isn't then $\pi \not\equiv -1 \pmod{q}$. Let p^a be an unknown component of N. Lemma 5, combined with what we've just said, implies $v_q(\sigma(p^a)) = v_q(a+1)$ if $p \equiv 1 \pmod{q}$, and equals 0 otherwise. Using the equation $\sigma(N)/N = 2$, there are n - b copies of q that must be accounted for by σ of the unknown components. Note that there are at most $k'_1 + k_2$ distinct prime factors of N that are congruent to 1 (mod q). Thus, by Lemma 6, at most $k'_1 + k_2 - 1$ copies of q divide a+1 if $p \equiv 1 \pmod{q}$ (since p itself cannot divide $\sigma(p^a)$), otherwise we end up with too many distinct prime divisors of N which are congruent to 1 (mod q). There are at most $k'_1 - \ell'_1 + k_2$ primes p with unknown component satisfying $p \equiv 1$ (mod q). Hence at most $(k'_1 - \ell'_1 + k_2)(k'_1 + k_2 - 1)$ copies of q can be accounted for by σ of the unknown components. Therefore

$$\tau = n - b - (k_1' - \ell_1' + k_2)(k_1' + k_2 - 1) \leqslant 0$$

or we will have left over copies of q not accounted for in $\sigma(N)$.

To get the last statement, now suppose $\tau > 0$, $\pi \equiv -1 \pmod{q}$, and π is among the unknown components. Lemma 5 tells us that the only place the extra τ copies of q can be accounted for is $v_q(\pi + 1)$, and hence $q^{\tau}|(\pi + 1)$. Therefore $\pi \equiv -1$ (mod q^{τ}) and so π is an unknown prime.

This means that when we counted the maximum number of possible (unknown) prime divisors $\equiv 1 \pmod{q}$ we included one too many. Thus there are at most $k'_1 + k_2 - 1$ primes dividing N that are congruent to 1 (mod q), and hence at most $(k'_1 - \ell'_1 + k_2 - 1)(k'_1 + k_2 - 2)$ copies of q can be accounted for by σ of the unknown, non-special components. For the special component π^{α} , by Lemma 6 part (ii) we see that at most $(k'_1 + k_2 - 1)/2$ copies of q can divide $\alpha + 1$, else we obtain too many factors of N congruent to 1 (mod q). Since, in fact, an integer number of copies of q divides $\alpha + 1$, we can take the floor. Putting this all together, τ' copies of q must divide $\pi + 1$.

Thus, if a large power of a Fermat prime divides N, we see that $\pi + 1$, and hence π , must be large. This isn't quite good enough to simplify our search to

manageable cases. We need a way to find another large prime divisor of N. The trick is to consider divisors of $\sigma(q^n)$. Suppose q is a Fermat prime, with $q^n || N$, and n very large. Then using the previous lemma, we can force π to be very large. One might wonder if $\pi |\sigma(q^n)$. The following result speaks to this issue.

Lemma 8 ([3, Lemma 1]). If p and q are odd primes with $p|\sigma(q^k)$ and $q^m|(p+1)$ then $k \ge 3m$.

So, using the terminology of Proposition 7, if $\tau > 0$, $q^{\tau} \nmid (p+1)$ for the known primes, and also $3\tau' > n$, then $\pi \nmid \sigma(q^n)$ by Lemma 8. But since n is big we would expect a large prime divisor of N which divides $\sigma(q^n)$. The following work clarifies how large a divisor we can find for $\sigma(q^n)$.

Lemma 9. Let p be an odd prime and let $q \equiv 3 \text{ or } 5$. If $q^{p-1} \equiv 1 \pmod{p^2}$ then either (q, p) = (3, 11) or $q^{o_p(q)} - 1$ has a prime divisor greater than 10^{13} . If q = 17then either (q, p) = (17, 3) or $q^{o_p(q)} - 1$ has a prime divisor greater than 10^{11} .

Proof. The papers [20] and [16] give a list of (q, p) for which $q^{p-1} \equiv 1 \pmod{p^2}$ and $p < 10^{13}$ with q = 3 or 5 (or $p < 10^{11}$ with q = 17). In the cases $(q, p) \neq (3, 11), (17, 3)$ the following table gives the requisite factor of $q^{o_p(q)} - 1$.

q	p	Large factor of $q^{o_p(q)} - 1$
3	1006003	154680726732318637
5	20771	625552508473588471
5	40487	625552508473588471
5	53471161	50493456782731
5	164533507	52082118058261
5	6692367337	8930008316757509
5	188748146801	40093613041379
17	46021	1365581260423071390161
17	48947	63895279579889
	•	

For use shortly, we make the following definition. Letting p,q be odd primes, $p\neq q,$ we set

$$o'_{q}(p) = \begin{cases} \text{if } 2 \nmid o_{q}(p), \\ \text{or } 4 \nmid o_{q}(p) \text{ and either} \\ o_{q}(p) & (\text{i}) \ p = \pi \text{ or} \\ (\text{ii}) \ p \equiv 1 \pmod{4} \text{ and } \pi \text{ is not} \\ \text{among the known components} \\ 0 & \text{otherwise.} \end{cases}$$

In other words, $o'_q(p)$ is the usual order function, unless it is impossible for both $p^a || N$ and $o_q(p) | (a + 1)$ to hold, due to consideration of the Eulerian form.

Proposition 10. Let N be an odd perfect number with k, k_1 , and k_2 having their usual meanings. Suppose q = 3 or 5 is a known prime divisor of N, $q^n || N, q \neq \pi$ and $\pi \nmid \sigma(q^n)$. Suppose p_1, \ldots, p_{k_1-1} are the other known factors of N, besides q.

For each $i \in [1, k_1 - 1]$ define

$$\epsilon_{i} = \begin{cases} 0 & \text{if } o_{p_{i}}'(q) = 0\\ \max(s+t-1,1) & \text{if } o_{p_{i}}'(q) \neq 0, \, s = v_{p_{i}}(\sigma(q^{o_{p_{i}}(q)-1})),\\ & \text{and } t \in \mathbb{Z}_{+} \text{ minimal so that } p_{i}^{t} > 100. \end{cases}$$

Set $V = \prod_{i=1}^{k_1-1} p_i^{\epsilon_i}$. Suppose π is among the k_2 unknown prime factors, and $k_2 > 1$. Finally, assume that all unknown prime factors are greater than 100. If

$$\min\left(10^{13}, \left(\frac{\sigma(q^n)}{V}\right)^{\frac{1}{k_2-1}}, \left(\frac{\sigma(q^{100})}{V}\right)^{\frac{1}{k_2-1}}\right) > 1,$$

then $\sigma(q^n)$ has a prime divisor among the unknown primes at least as big as the above minimum. If q = 17 one can replace 10^{13} with 10^{11} , and then the result still holds.

Proof. We do the case when q = 3, since the other cases are similar. First suppose $\sigma(q^n) = (q^{n+1} - 1)/(q - 1)$ is at most divisible by $p_i^{\epsilon_i}$ for the known primes, and square-free for the unknown primes. Then since $\pi, q \nmid \sigma(q^n)$, the largest unknown divisor of $\sigma(q^n)$ is at least

$$\left(\frac{\sigma(q^n)}{V}\right)^{\frac{1}{k_2-1}},$$

unless this quantity is ≤ 1 (in which case there might be no unknown factors).

So we may assume there is some prime $p|N, o'_p(q) \neq 0$, so that $\sigma(q^n)$ is divisible by p^2 if p is unknown, or $p^{\epsilon+1}$ if p is known (and ϵ is the corresponding ϵ_i), with pmaximal among such primes. By Lemma 9 we may also assume that if $p^2 \nmid (q^{p-1}-1)$ and p is unknown then p|(n+1). (This is where 10^{13} comes into the minimum.)

Thus in either case, $p^t|(n+1)$, where $p^t > 100$ (taking t = 1 if p is an unknown prime). Then we have

$$\sigma(q^{p^{\tau}-1})|\sigma(q^n).$$

Thus it suffices to find a large divisor of $(q^{p^t} - 1)/(q - 1)$. By Lemma 1, $(q^{p^t} - 1)/(q - 1)$ is only divisible by primes larger than p, or p itself to the first power. (In this case, q being Fermat means the quantity isn't divisible by p, and we could replace $\max(s + t - 1, 1)$ by s + t - 1 in the definition of ϵ_i . But to keep similar notations later when we take q to be an arbitrary prime, we don't use this fact.) But then, by the maximality condition on p, $(q^{p^t} - 1)/(q - 1)$ is not divisible by more than $p_i^{\epsilon_i}$ for known primes and the first power for all the unknown primes. So the analysis we used in the first paragraph goes through by only changing n + 1 to p^t . Finally, note that $p^t > 100$, so we have the appropriate bound.

The most useful case when we will use Proposition 10 is when n is very large and k_1 is close to k. So, in practice, we will usually end up with 10^{13} as the lower bound on a divisor of $\sigma(q^n)$. In fact, a lot of work could be saved if the bounds given in [16] were improved, or there was some means to work around square divisors.

5. Non-Fermat Primes

Sometimes our odd perfect number will not be divisible by a large power of a Fermat prime, but rather by a large power of some arbitrary prime, q. Unfortunately, we don't have Lemma 5, and so we can't put all of the "extra" factors of qinto $\pi + 1$. This causes two problems. First, we have to spread the extra factors of q among the unknown primes, thus reducing the number of extra factors we have at hand, exponentially. Second, if $p \not\equiv 1 \pmod{q}$ is one of our unknown primes, we cannot reduce to the case $q|\Phi_2(p)$ but rather $q|\Phi_d(p)$ for some (arbitrary) d > 1, d|(q-1). Thus, we need a way of bounding the size of p for which $q^n|\Phi_d(p)$. This second problem is easily dealt with.

Lemma 11 ([16, Theorem 2]). Let q be an odd prime. Let a_1 be a primitive root modulo q. For $r \ge 2$, define $a_r = a_1^{q^{r-1}} \pmod{q^r}$. Then $\{a_r^m \pmod{q^r} \mid m = 0, \ldots, q-2\}$ gives a complete set of incongruent solutions to $a^{q-1} \equiv 1 \pmod{q^r}$.

Lemma 12. Let q < 1000 be an odd prime. If $p^{q-1} \equiv 1 \pmod{q^n}$ for some $n \in \mathbb{Z}_+$ and some odd prime p, then $p \ge \min(q^{n-2}, 10^{50})$ except when

(p,q) = (40663372766570611389846294355914421,7).

Proof. Let q < 1000 be an odd prime. Fix $m \in \mathbb{N}$ so that $q^{m-2} > 10^{50}$ and m is minimal. A computer search using Lemma 11 demonstrates that every positive solution to $x^{q-1} \equiv 1 \pmod{q^m}$ with $x \neq 1$ satisfies $x > 10^{50}$. Hence the same is true if we replace m with a larger integer. Thus, it suffices to search for prime solutions to $x^{q-1} \equiv 1 \pmod{q^n}$ for $n \leq m$. Again, a computer search yields the stated result.

The choice of 1000 in Lemma 12 is easily improved, but is large enough for our needs. Also, the exceptional (p,q) in the lemma is irrelevant to our work, since in this case we find $o_q(p) = 6$ and $\sigma(p^5)$ gives rise to at least 12 additional prime factors of N besides p and q.

Lemma 13. Let N be an odd perfect number, let q be an odd prime, and suppose $p^a || N$ with p prime.

- (i) If $p \equiv 1 \pmod{q}$ and $q^b | \sigma(p^a)$ then $\sigma(p^a)$ is divisible by b distinct primes r_1, r_2, \ldots, r_b with $r_i \equiv 1 \pmod{q^i}$.
- (ii) If $p \not\equiv 1 \pmod{q}$, $o_q(p)|(a+1)$, and $q^c|(a+1)$, then $\sigma(p^a)$ is divisible by $c \cdot \sigma_0(o_q(p))$ distinct primes divisors $r_{i,j}$, $i \in [1,c]$, $j \in [1,\sigma_0(o_q(p))]$, with $r_{i,j} \equiv 1 \pmod{q^i}$.

Proof. Analogous to Lemma 6.

Proposition 14. Let N be an odd perfect number, and let q < 1000 be a prime divisor of N with $q^n | N$. Suppose b, k, k_1 , k_2 , ℓ_1 , ℓ_2 , k'_1 , and ℓ'_1 , have the same meanings as in Proposition 7. Suppose further that the exceptional case of Lemma 12 doesn't hold. Let T be the set of known primes with unknown component, different from q, and $\neq 1 \pmod{q}$. Let

$$\tau = n - b - \sum_{p \in T, \, o'_q(p) \neq 0} \left(v_q(p^{o_q(p)} - 1) + \left\lfloor \frac{k'_1 + k_2}{\sigma_0(o_q(p))} \right\rfloor \right) - (k'_1 - \ell'_1 + k_2)(k'_1 + k_2 - 1).$$

If $\tau > 0$ then one of the unknown primes is not congruent to 1 (mod q). Further, in this case, one of the unknown primes is at least as large as $\min(q^{\tau'-2}, 10^{50})$ where

г /

$$\tau' = \min_{1 \le m \le k_2} \left| \left(n - b - \sum_{p \in T, \, o'_q(p) \neq 0} \left(v_q(p^{o_q(p)} - 1) + \left\lfloor \frac{k'_1 + k_2 - m}{\sigma_0(o_q(p))} \right\rfloor \right) - (k'_1 - \ell'_1 + k_2 - m)(k'_1 + k_2 - m - 1) - m \left\lfloor \frac{k'_1 + k_2 - m}{2} \right\rfloor \right) / m \right|.$$

Proof. The proof is similar to Proposition 7. From the equation $\sigma(N)/N = 2$ we know that $q^n | \sigma(N)$, and so we try to account for as many copies of q in $\sigma(N)$ as we can. The quantity τ is exactly how many copies of q are unaccounted for, if all of the unknown primes are $\equiv 1 \pmod{q}$, and we try to account for as many copies of q as possible from the known primes using Lemmas 2 and 13. If $\tau > 0$ this means we actually have left over copies of q, which yields a contradiction, and hence not all of the unknown primes are $\equiv 1 \pmod{q}$.

In this case, let *m* be the number of unknown primes $\neq 1 \pmod{q}$. Lemma 2 tells us that there are two sources of copies of *q* in $\sigma(N)$; the exponents of the primes, and $p^{o_q(p)} - 1$. The quantity

(2)
$$n-b-\sum_{p\in T, \, o'_q(p)\neq 0} \left(v_q(p^{o_q(p)}-1) + \left\lfloor \frac{k'_1+k_2-m}{\sigma_0(o_q(p))} \right\rfloor \right) - (k'_1-\ell'_1+k_2-m)(k'_1+k_2-m-1) - m \left\lfloor \frac{k'_1+k_2-m}{2} \right\rfloor$$

is the number of copies of q in $\sigma(N)$ not yet accounted for, after we account for as many copies of q as we can (again using Lemmas 2 and 13) except those copies of q which come from $\sigma(p^{o_q(p)-1})$ for the unknown primes p with $p \not\equiv 1 \pmod{q}$. Thus, if we divide Equation 2 by m, and take the ceiling, we have a number of copies of q that must be accounted for by $p^{o_q(p)} - 1$ for an unknown prime p with $o_q(p) \neq 1$. Since m is unspecified by the hypotheses, we take the minimum over all possibilities. Finally, we apply Lemma 12.

Next we want to prove a result analogous to Proposition 10, except for arbitrary primes. However, there have to be a few differences. First, we need something to play the role of Lemma 8. It turns out that it doesn't hurt much to just assume that the large prime, call it p, coming from Proposition 14 may in fact divide $\sigma(q^n)$. So we can write $q^{n+1} - 1 = (q-1)p^c m$ for some $c \in \mathbb{N}$ and $m \in \mathbb{Z}_+$. Powering this equation to the (q-1)st power, since $p^{q-1} \equiv 1 \pmod{q^{\tau'}}$ and $\tau' \leq n$, we have

$$((q-1)m)^{q-1} \equiv ((q-1)mp^c)^{q-1} = (q^{n+1}-1)^{q-1} \equiv 1 \pmod{q^{\tau'}}$$

and hence we can bound (q-1)m using an analogue of Lemma 12 (where we look for solutions to the equation $x^{q-1} \equiv 1 \pmod{q^n}$ which are divisible by q-1, rather than prime solutions). The following lemma provides this result:

Lemma 15. Let q < 1000 be an odd prime. Suppose $a^{q-1} \equiv 1 \pmod{q^n}$ for some $n \in \mathbb{Z}_+$ and some integer a with (q-1)|a. If $q \ge 11$ then $a \ge \min(q^{n-2}, 10^{50})$. If q = 7, then $a \ge \min(q^{n-3}, 10^{50})$.

Proof. An easy computer search as in Lemma 12.

Next we need to work with possible square factors of $\sigma(q^n)$, similar to Lemma 9.

Lemma 16. Let p and q be primes with $p \in (10^2, 10^{11})$ and q = 7, 11 or 13. If $q^{p-1} \equiv 1 \pmod{p^2}$ then $\sigma(q^{o_p(q)-1})$ is divisible by two primes greater than 10^{11} .

Proof. By [16], there are only 3 pairs (p,q) satisfying the conditions, namely

(p,q) = (491531,7), (863,13), (1747591,13).

In the first case, since $65|o_p(q)$, just factor $\sigma(7^{64})$ to find two distinct primes greater than 10^{11} . In the second case $o_p(q) = 862$. One factor, 16002623839393, is easily found and another is

$$\frac{13^{431} + 1}{14 \cdot 863^3 \cdot 68099}$$

In the last case, since $195|o_p(q)$, one factors $\sigma(13^{38})$ to find one prime greater than 10^{11} and factors $\sigma(13^{64})$ to find the other one. (Or, to save time, use the online factorizations on Richard Brent's extensions of the Cunningham tables for these last two.)

We are now ready to prove:

Proposition 17. Let N be an odd perfect number and let q = 11 or 13 be a known prime divisor of N, with $q^n || N$. Let τ, τ' be as in Proposition 14, suppose all the hypotheses of that proposition are met, and let p be the guaranteed unknown prime. Let p_1, \ldots, p_{k_1-1} be the known primes different from q. Let ϵ_i be defined as before, and put $V = \prod_{i=1}^{k_1-1} p_i^{\epsilon_i}$. Suppose $k_2 > 1$. Finally, assume that all unknown prime factors are greater than 100.

If

$$\min\left(10^{11}, \left(\frac{10^{50}}{(q-1)V}\right)^{\frac{1}{k_2-1}}, \left(\frac{q^{\tau'-2}}{(q-1)V}\right)^{\frac{1}{k_2-1}}\right) > 1$$

then $\sigma(q^n)$ has a prime divisor, different from p, among the unknown primes, at least as big as the above minimum. If q = 7 the same result holds if we replace $q^{\tau'-2}$ by $q^{\tau'-3}$.

Proof. We do the case q = 11 or 13, the other being similar. So assume the above minimum is > 1. First note that if d|(n+1) then $\sigma(q^{d-1})|\sigma(q^n)$ and so it suffices to show that $\sigma(q^{d-1})$ has a prime divisor larger than the above minimum, different from p, for some d|(n+1).

By the method of proof given in Proposition 10, we may assume that at most ϵ_i copies of p_i divide $\sigma(q^{d-1})$, for some d either greater than 100 or equal to n + 1. Further, because 10^{11} occurs in the above minimum, we may assume that the only unknown prime that may divide $\sigma(q^n)$ which may be greater than 10^{11} is p. Then by Lemma 16 and the fact that the unknown primes are greater than 100, we may assume $\sigma(q^n)$ is square-free for unknown primes, except possibly p.

From the proof for Proposition 14, we have $p^{q-1} \equiv 1 \pmod{q^{\tau'}}$. Write $q^d - 1 = (q-1)mp^c$ with $c \in \mathbb{N}$, $m \in \mathbb{Z}_+$, and gcd(p,m) = 1. Powering this equation to the (q-1)st power, we have

$$((q-1)m)^{q-1} \equiv ((q-1)mp^c)^{q-1} = (q^d-1)^{q-1} \equiv 1 \pmod{q^{\min(\tau',100)}}.$$

By Lemma 15,
$$(q-1)m \ge \min(q^{\tau'-2}, q^{98}, 10^{50}) = \min(q^{\tau'-2}, 10^{50})$$
. Thus
$$\frac{m}{V} \ge \min\left(\frac{q^{\tau'-2}}{(q-1)V}, \frac{10^{50}}{(q-1)V}\right).$$

Since m/V is at least as big as the part of $\sigma(q^{d-1})$ made up from the unknown primes, different from p, if we take the $k_2 - 1$ root of the minimum we have the appropriate lower bound.

6. Abundance and Deficiency

Let $n \in \mathbb{Z}_+$. Recall the multiplicative function $\sigma_{-1}(n) = \sum_{d|n} d^{-1}$ we introduced earlier. This function can alternatively be written using the formula $\sigma_{-1}(n) = \sigma(n)/n$, and so $\sigma(n)/n = 2$ if and only if $\sigma_{-1}(n) = 2$. A number *n* is called *abundant* when $\sigma_{-1}(n) > 2$ and *deficient* when $\sigma_{-1}(n) < 2$. We can use abundance and deficiency computations to limit choices on possible prime factors of an odd perfect number *N*. First, we extend the definition of σ_{-1} by setting

$$\sigma_{-1}(p^{\infty}) = \lim_{a \to \infty} \sigma_{-1}(p^a) = \frac{p}{p-1}$$

Lemma 18 ([25, Proposition 2]). Let p and q be odd primes. If $1 \le a < b \le \infty$ then $1 < \sigma_{-1}(p^a) < \sigma_{-1}(p^b)$. If $1 \le a, b \le \infty$ and p < q then $\sigma_{-1}(q^b) < \sigma_{-1}(p^a)$.

Lemma 19. Let N be an odd perfect number. Suppose p_1, \ldots, p_{k_1} are the known prime factors of N, $p_i^{a_i}|N$, and $k_1 < k = \omega(N)$. If $\Pi = \prod_{i=1}^{k_1} \sigma_{-1}(p_i^{a_i}) < 2$, then the smallest unknown prime is

$$p_{k_1+1} \geqslant \frac{\Pi}{2-\Pi}$$

Proof. We find

$$2 = \sigma_{-1}(N) \ge \left(\prod_{i=1}^{k_1} \sigma_{-1}(p_i^{a_i})\right) \sigma_{-1}(p_{k_1+1}) = \Pi \cdot \frac{p_{k_1+1}+1}{p_{k_1+1}}.$$

where the inequality in the middle follows from Lemma 18. Noting $\Pi \ge 1$, we obtain $\frac{2}{\Pi} \ge 1 + \frac{1}{p_{k_1+1}}$. Therefore $\frac{2-\Pi}{\Pi} \ge \frac{1}{p_{k_1+1}}$ and taking reciprocals gives us the result, since $2 - \Pi > 0$.

Note that in the lemma if $\Pi > 2$ then $\prod_{i=1}^{k_1} p_i^{a_i}$ is abundant, hence N is abundant. If $\Pi = 2$, then $\prod_{i=1}^{k_1} p_i^{a_i}$ is already an odd perfect number.

The following lemma is the true key to our search for odd perfect numbers, as simple as the proof is (after wading through the hypotheses). This is because we built up machinery in the last few sections to find bounds for large prime divisors of N.

Lemma 20 (c.f. [5, Lemma 2.2]). Let N be an odd perfect number. Let p_1, \ldots, p_k be the prime divisors of N, and let a_i be such that $p_i^{a_i} || N$. Fix the numbering on the indices so that p_1, \ldots, p_{ℓ_1} are the primes with known prime component, $p_{\ell+1}, \ldots, p_{k_1}$ are the other known primes, and $p_{k_1+1} < \ldots < p_k$ are the unknown primes. Suppose among the unknown primes we have bounds $p_k > P_1, \ldots, p_{k-\nu+1} > P_\nu$, with $\nu < k_2$ and $P_i > 1$ for each $i \in [1, \nu]$. For each $u \in [0, \nu]$, set

$$\Delta_u = \left(\prod_{i=1}^{\ell_1} \sigma_{-1}(p_i^{a_i})\right) \left(\prod_{i=\ell_1+1}^{k_1} \frac{p_i}{p_i-1}\right) \left(\prod_{i=1}^{u} \frac{P_i}{P_i-1}\right).$$

Finally, suppose $k_2 > 0$.

If $\Delta_u < 2$ then the smallest unknown prime is

$$p_{k_1+1} \leqslant \frac{\Delta_u(k_2-u)}{2-\Delta_u} + 1.$$

Therefore,

$$p_{k_1+1} \leq \min_{u \in [0,v], \, \Delta_u < 2} \left(\frac{\Delta_u(k_2 - u)}{2 - \Delta_u} + 1 \right).$$

Proof. We compute

$$2 = \sigma_{-1}(N) = \prod_{i=1}^{k} \sigma_{-1}(p_{i}^{a_{i}})$$

$$\leq \left(\prod_{i=1}^{\ell_{1}} \sigma_{-1}(p_{i}^{a_{i}})\right) \left(\prod_{i=\ell_{1}+1}^{k-u} \sigma_{-1}(p_{i}^{\infty})\right) \left(\prod_{i=k-u+1}^{k} \sigma_{-1}(P_{k-i+1}^{\infty})\right)$$

$$= \Delta_{u} \prod_{i=k_{1}+1}^{k-u} \sigma_{-1}(p_{i}^{\infty}) \leq \Delta_{u} \prod_{i=0}^{k-u-k_{1}-1} \sigma_{-1}((p_{k_{1}+1}+i)^{\infty})$$

$$= \Delta_{u} \frac{p_{k_{1}+1}+k-u-k_{1}-1}{p_{k_{1}+1}-1} = \Delta_{u} \left(1 + \frac{k_{2}-u}{p_{k_{1}+1}-1}\right).$$

Now, recall that $u \leq v < k_2$ which implies $k_2 - u \geq 1$. Also $0 < \Delta_u < 2$, so we solve the main inequality as we did in the previous lemma, finding

$$p_{k_1+1} \leqslant \frac{\Delta_u(k_2-u)}{2-\Delta_u} + 1.$$

The last statement follows.

One major difference between Lemma 20 and Lemma 19 is that if $\Delta_u > 2$ then that doesn't necessarily imply N is abundant. (It is true that if $k_1 = k$ and $\Delta_0 < 2$ then N is deficient, however.) This means that we might end up with $\Delta_0 > 2$, and hence we have no upper bound on p_{k_1+1} .

7. The Algorithm

To verify that an odd perfect number has at least 9 distinct prime factors, we use a factor chain algorithm to check all possible cases for odd perfect numbers with exactly 8 distinct prime factors, and find contradictions in each case, similar to the algorithms described in both [5] and [25]. This section will describe the algorithm, and the next section will explain my implementation.

We start by knowing 3|N, so we consider each of the cases $3^2||N, 3^4||N$, and so forth. We think of each of these cases as branches on a tree. On the branch $3^2||N$ we know that $\sigma(3^2)|2N$ and so 13|N. Hence the case $3^2||N$ branches further and we have to consider each of the subcases $13||N, 13^2||N, 13^4||N$, and so forth. We continue finding new factors, and the tree continues branching out.

Some cases do not result in new primes. To continue the factor chain in these cases we use the bounds of Section 6 to find an interval for the smallest unknown prime. Then we consider all the new primes in the given interval. We only stop branching when we reach a contradiction (such as having too many factors).

As it stands there are still an infinite number of branches. To get around this problem, we can combine all the cases $p^n || N$ for large n together into one case.

More precisely, let B be a large integer (around the size 10^{50}) which we fix at the beginning of the algorithm. Then, once we reach the branch p^n , with n large enough so that $p^n > B$, we no longer consider the case $p^n || N$ but rather we just assume $p^a |N$ for some $a \ge n$, and thus consider all the remaining cases together. (In particular, p is still a known prime divisor of N, but the component is unknown, although we are given a lower bound of B on the size of the component.) We then label this composite branch by p^{∞} .

So, for example, if $B = 10^2$ then since $3^4 < B < 3^6$ the first level of our tree would consist of the branches 3^2 , 3^4 , and 3^∞ . When we are on a branch with p^∞ we say p is an *infinite prime* (not to be confused with the infinite primes of algebraic number theory). Notice that infinite primes do not provide more primes for the factor chain, and so we have to rely on on the interval bounds of Section 6. If we set B too low then the primes become infinite too quickly and we may have the case the there is no upper bound for an interval. (This corresponds to the case when $\Delta_0 > 2$ in Lemma 20.) If we make B large enough this never happens (for a proof see [5]), and the algorithm will only have to consider a finite number of cases.

We are now able to expressly define what we mean by *known* primes and components. Suppose we are on the branch $3^{\infty}5^4$. Then the known primes are 3, 5, 11, 71 (the primes 11 and 71 come from $\sigma(5^4)|2N$), and the only known component is 5^4 . In this case we say that 3 and 5 are *on* while 11 and 71 are *off*. In other words, the *on* primes are exactly the known primes for which we have started the branching process. Note that $k_1 - \ell_1$ is exactly the number of (known) primes which are infinite or off.

In our example, since 11 is the smallest off prime we continue the branching process on this prime, rather than 71. When there are no off primes we use the interval bounds to arrive at another prime, as explained earlier. Whenever we reach a contradiction, we go to the next available branch.

The following is a possible, initial print-out in the case k = 5, B = 50. (Note: When calculating intervals this output only used Δ_0 in Lemma 20.)

The letter \mathbb{N} means that there are no primes in the interval, which is a contradiction. Adding more contradictions, or using the full power of Lemma 20, can further simplify the output.

8. An Implementation

There are three main differences between our implementation of this algorithm, and the implementations in [5] and [25]:

First, the bound B is not allowed to increase within the algorithm. Allowing B to vary fully automates the algorithm at the expense of unnecessary complexity. The number B is fixed at the outset, and only increased manually if needed.

Second, the use of Lemma 20 allows for stronger upper bounds on intervals. In the terminology of that lemma, we always have $v \leq 3$. We take $P_1 = 10^7$, $P_2 = 10^4$, and $P_3 = 10^2$, unless we already have known prime divisors larger than these bounds, or unless these bounds are superseded by the work in Sections 4 and 5. (So, for example, if the largest known prime is > 10⁷, and the next largest known prime is < 10², then we could take $P_1 = 10^4$ and $P_2 = 10^2$. If large powers of small primes divide N then we can use the work in previous sections to increase P_1 and P_2 .)

Third, the contradictions used are different. Here is a complete list of the contradictions in our implementation:

MT There are too many total factors.

- MS There are too many copies of a single prime with known component.
 - S There is an off prime smaller than an on prime coming from interval computations.
 - A The number is abundant.
 - D There are k known primes, and $\Delta_0 < 2$, hence N is deficient.
 - F The special prime π belongs to a known component, but the hypotheses of Proposition 7 hold showing π must be in an unknown component due to a Fermat prime.
 - N There are no primes in the interval given by Lemmas 19 and 20, or all the known primes are on and the only primes in the interval are already known.
- SF1 There are k-1 known primes, and the interval formula gives an upper bound of $p_k < C$, but we know from the fact that a large power of a small Fermat prime divides N that some unknown prime is larger than C, by Proposition 7.
- SF2 Similar to SF1 except we have a contradiction between the interval formula and Proposition 10.
- SNF1 Similar to SF1, except we have a contradiction from a small non-Fermat prime, using Proposition 14.
- SNF2 Similar, using Proposition 17.
- P1Int There are k 1 known primes all smaller than 10⁷, and the interval formula gives an upper bound $p_k < 10^7$.
- P2Int There are k 2 known primes all smaller than 10^4 , or k 1 known primes with the largest $> 10^7$ and none other $> 10^4$; and the interval formula says the next prime is $< 10^4$.
- P3Int There are k 3 known primes all smaller than 10^2 , or k 2 known primes only one greater than 10^4 and none other $> 10^2$, or k - 1 known primes with one larger than 10^6 and another larger that 10^4 and none other $> 10^2$; and the interval formula says the next prime is $< 10^2$.

Of course, before the algorithm checks for any contradictions it always checks if we have an odd perfect number. There were other contradictions we might have included. For example, if there are k known primes, and none of them can be the special prime, this means N cannot be an odd perfect number. However, the strength of our method lies in the fact that we rarely have exactly k known primes, and in those cases one of the other contradictions will do.

9. Points of Improvement and the Results

The algorithm was implemented in Mathematica on a Pentium 4 personal computer. The factorizations of $\sigma(p^a)$ were carried out using a only a probable primality

test, good for integers $< 10^{16}$. Thus, if a factor of $\sigma(p^a)$ was larger than this bound, Mathematica's primality proving routine was also run. After running the algorithm it became clear that certain modifications would speed up the process. First, the bound *B* was too uniform. So, *B* was replaced by two bounds B_1 and B_2 , where a prime p < 1000 became infinite when $p^a | N$ and $p^a > B_1$, while a prime p > 1000became infinite when $p^a | S_2$.

With $B_1 = 10^{50}$ and $B_2 = 10^{18}$, the algorithm didn't terminate until we reached the case

$$3^{\infty}5^{\infty}17^{\infty}257^{\infty}65537^{\circ}$$

because B_2 wasn't large enough. After increasing it to $B_2 = 10^{30}$ the algorithm finished the case $\omega(N) = 8$ without finding an odd perfect number, running for a total of less than 3 days, after numerous stops and starts.

One of the longest cases (besides the one mentioned above) was $3^47^{\infty}5^1$. This is because the bound given in Proposition 17 depends on taking $(k_2 - 1)$ st roots. So, when $k_2 = 3$ the bound is really around the square-root of where it should be. This led to formulating the improved result:

Proposition 21. Assume all of the hypotheses of Proposition 10 hold. Further, suppose that we can bound $p_{k_1+1} < P$. Finally assume $k_2 > 2$. If

$$\min\left(10^{13}, \left(\frac{\sigma(q^n)}{VP}\right)^{\frac{1}{k_2-2}}, \left(\frac{\sigma(q^{100})}{VP}\right)^{\frac{1}{k_2-2}}\right) > 1,$$

(replacing 10^{13} by 10^{11} if q = 17) then $\sigma(q^n)$ has a prime divisor among the unknown primes at least as big as the above minimum.

Proof. One reduces to the case that $\sigma(q^n)$ is square-free for unknown primes, just as in Proposition 10. Then the result is obvious.

Of course, Proposition 17 is similarly improved. Once these improvements were put into place, the case $3^47^{\infty}5^1$ took less than an hour. The case $\omega(N) = 8$ took less than half a day of non-stop running, with less than 800,000 lines of output. We thus have:

Theorem 22. An odd perfect number has at least 9 distinct prime divisors.

Running the algorithm with $B_1 = 10^{50}$, $B_2 = 10^{30}$, k = 11, and forcing $3 \nmid N$, also terminated without finding any odd perfect numbers after about one million lines of output. Thus:

Theorem 23. An odd perfect number N with $3 \nmid N$ has at least 12 distinct prime divisors.

At this point, the results seemed to depend on a lot of previous computation. To avoid this, the algorithm was rerun with the following changes:

- All uses of the P1Int, P2Int, and P3Int contradictions were removed.
- The results of [13], [14], and [15] were not used in interval calculations.
- It wasn't assumed that 3|N. In fact, Grün's result wasn't even used, but rather Lemma 20, to obtain $p_1 \leq k+1$.
- It wasn't assumed that $\omega(N) \ge 8$. So each $k \in [1, 8]$ was checked.

With these changes, the only results (outside this paper) which were used were (i) some of the easy to prove non-computational lemmas (such as the Eulerian form) and (ii) the main result of [16] for primes $a \leq 17$. The algorithm again terminated

after about one million lines of output, verifying Theorem 22. A similar run verified Theorem 23.

10. FUTURE RESULTS

To do the case $\omega(N) = 9$ with these techniques, in a reasonable amount of time, it is necessary to find bounds for p_{k-2} , or increase the lower bound on p_{k-1} . For example, consider the case of

$3^{\infty}5^{\infty}17^{\infty}257^{\infty}65537^{\infty}.$

We can make the powers on these primes so large that we may, for all intents and purposes, assume p_k is as big as we like (except not quite big enough to use [22]). However, the best we can do with our tools for p_{k-1} is 10^{13} , which is much smaller than the lower bound already given by the interval formula. The interval for p_{k-1} has a length of about 10^{18} , which is much too large to check one prime at a time. Further, it is currently computationally unfeasible to push the bound in [16] up to 10^{19} .

Another possible line of attack would be to suppose that two primes q_1, q_2 divide N, each to a high power. Then one might be able to prove that $\sigma(q_1^{n_1})$ and $\sigma(q_2^{n_2})$ must each have large divisors, different from one another, once n_1 and n_2 are large enough. Another alternative would be to show that the square-free part of $\sigma(q^n)$ becomes large as n does.

References

- 1. A. Bang, Taltheoretiske Undersøgelser, Tidsskrift Math. 5 IV (1886), 70-80, 130-137.
- 2. Geo. D. Birkhoff and H. S. Vandiver, On the integral divisors of $a^n b^n$, Ann. of Math. (2)
- (1904), no. 4, 173–180. MR 1503541
 R. P. Brent, G. L. Cohen, and H. J. J. te Riele, *Improved techniques for lower bounds for odd perfect numbers*, Math. Comp. 57 (1991), no. 196, 857–868. MR 1094940 (92c:11004)
- 4. Joseph E. Z. Chein, An odd perfect number has at least 8 prime factors, Ph.D. thesis, Penn-sylvania State University, 1979.
- Graeme L. Cohen and Ronald M. Sorli, On the number of distinct prime factors of an odd perfect number, J. Discrete Algorithms 1 (2003), no. 1, 21–35, Combinatorial algorithms. MR 2016472 (2004h:11003)
- R. J. Cook, Bounds for odd perfect numbers, Number theory (Ottawa, ON, 1996), CRM Proc. Lecture Notes, vol. 19, Amer. Math. Soc., Providence, RI, 1999, pp. 67–71. MR 1684591 (2000d:11010)
- Leonard Eugene Dickson, Finiteness of the Odd Perfect and Primitive Abundant Numbers with n Distinct Prime Factors, Amer. J. Math. 35 (1913), no. 4, 413–422. MR 1506194
- Otto Grün, Über ungerade vollkommene Zahlen, Math. Z. 55 (1952), 353–354. MR 0053123 (14,724g)
- Peter Hagis, Jr., Outline of a proof that every odd perfect number has at least eight prime factors, Math. Comp. 35 (1980), no. 151, 1027–1032. MR 572873 (81k:10004)
- 10. _____, Sketch of a proof that an odd perfect number relatively prime to 3 has at least eleven prime factors, Math. Comp. 40 (1983), no. 161, 399–404. MR 679455 (85b:11004)
- Kevin G. Hare, More on the total number of prime factors of an odd perfect number, Math. Comp. 74 (2005), no. 250, 1003–1008 (electronic). MR 2114661 (2005h:11010)
- D. R. Heath-Brown, Odd perfect numbers, Math. Proc. Cambridge Philos. Soc. 115 (1994), no. 2, 191–196. MR 1277055 (96b:11130)
- Douglas E. Iannucci, The second largest prime divisor of an odd perfect number exceeds ten thousand, Math. Comp. 68 (1999), no. 228, 1749–1760. MR 1651761 (2000i:11200)
- The third largest prime divisor of an odd perfect number exceeds one hundred, Math. Comp. 69 (2000), no. 230, 867–879. MR 1651762 (2000i:11201)
- Paul M. Jenkins, Odd perfect numbers have a prime factor exceeding 10⁷, Math. Comp. 72 (2003), no. 243, 1549–1554 (electronic). MR 1972752 (2004a:11002)

- 16. Wilfrid Keller and Jörg Richstein, Solutions of the congruence $a^{p-1} \equiv 1 \pmod{p^r}$, Math. Comp. 74 (2005), no. 250, 927–936 (electronic). MR 2114655 (2005i:11004)
- Masao Kishore, On odd perfect, quasiperfect, and odd almost perfect numbers, Math. Comp. 36 (1981), no. 154, 583–586. MR 606516 (82h:10006)
- _____, Odd perfect numbers not divisible by 3. II, Math. Comp. 40 (1983), no. 161, 405–411. MR 679456 (84d:10009)
- Wayne L. McDaniel, The non-existence of odd perfect numbers of a certain form, Arch. Math. (Basel) 21 (1970), 52–53. MR 0258723 (41 #3369)
- 20. Peter L. Montgomery, New solutions of $a^{p-1} \equiv 1 \pmod{p^2}$, Math. Comp. **61** (1993), no. 203, 361–363. MR **1182246 (94d:**11003)
- Trygve Nagell, Introduction to Number Theory, John Wiley & Sons Inc., New York, 1951. MR 0043111 (13,207b)
- Pace P. Nielsen, An upper bound for odd perfect numbers, Integers 3 (2003), A14, 9 pp. (electronic). MR 2036480 (2004k:11009)
- Carl Pomerance, Odd perfect numbers are divisible by at least seven distinct primes, Acta Arith. 25 (1973/74), 265–300. MR 0340169 (49 #4925)
- 24. _____, Multiply perfect numbers, Mersenne primes, and effective computability, Math. Ann. **226** (1977), no. 3, 195–206. MR 0439730 (55 #12616)
- John Voight, On the nonexistence of odd perfect numbers, MASS selecta, Amer. Math. Soc., Providence, RI, 2003, pp. 293–300. MR 2027187 (2004j:11006)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720 *E-mail address*: pace@math.berkeley.edu