The time evolution of permutations under random stirring

Bálint Vető

Abstract. We consider permutations of $\{1, \ldots, n\}$ obtained by $\lfloor \sqrt{nt} \rfloor$ independent applications of random stirring. In each step the same marked stirring element is transposed with probability 1/n with any one of the *n* elements. Normalizing by \sqrt{n} we describe the asymptotic distribution of the cycle structure of these permutations, for all $t \geq 0$, as $n \to \infty$.

1 Introduction

We consider the following random stirring mechanism: n numbered balls are given in the beginning on their corresponding numbered places. In each step, independently, the first ball, which is referred to as the *stirring particle* or *stirring element*, changes place with one of the n balls or stays unchanged with probability 1/n. We investigate that permutation which brings the balls from their initial place to their place after i steps.

Formally, let $\pi^{(n)}(i) = T_i^{(n)} \circ T_{i-1}^{(n)} \circ \cdots \circ T_1^{(n)}$ be a permutation acting on the set $[n] := \{1, \ldots, n\}$. The permutations $(T_i^{(n)})_{i=1}^{\infty}$ are chosen independently with uniform distribution from the n-1 transpositions moving the stirring particle and the identity permutation.

Let σ be a permutation of a finite set S, i.e. an $S \to S$ bijective function. The cycles (orbits) of σ are the sets of form $\{v, \sigma(v), \sigma^2(v), \dots\} \subseteq S$ for some $v \in S$. The set S is the disjoint union of its cycles. The cycle structure of σ is the sequence of the cardinalities of the different cycles in non-increasing order.

In our case one of the cycles can be distinguished from the others (namely the cycle of the stirring element), which will be called the *active cycle*. For the total description it is enough to determine the distribution of the cycle structure of the permutation $\pi^{(n)}(i)$ (regarding the active cycle separately). This gives the distribution of the conjugacy class of $\pi^{(n)}(i)$ restricting ourself to the conjugation with permutations fixing the stirring particle. The distribution of $\pi^{(n)}(i)$ is uniform within a fixed conjugacy class.

We encode the permutation $\pi^{(n)}(i)$ with the vector $\mathbf{C}^{(n)}(i) := (C_0^{(n)}(i), C_1^{(n)}(i), C_2^{(n)}(i), \ldots)$ where $C_0^{(n)}(i)$ denotes the length of the active cycle, $C_1^{(n)}(i), C_2^{(n)}(i), \ldots$ the lengths of those cycles in non-increasing order which are already moved by one of the transpositions $(T_j^{(n)})_{j=1}^i$. Other $C_j^{(n)}(i)$ -s are 0. $(\mathbf{C}^{(n)}(i))_{i=0}^{\infty}$ is a process on



Figure 1: The metric on \mathbf{S}

the state space

$$\mathbf{S} := \{(s_0, s_1, s_2, \dots) : s_n \in \mathbb{R}, \quad s_n \ge 0 \quad n = 0, 1, 2, \dots, \\ s_1 \ge s_2 \ge \dots \ge s_n \ge \dots \text{ and } s_j > 0 \text{ for finitely many } j\}$$
(1)

with the distance

$$d(A,B) := \sup\left\{ \left| \sum_{j=0}^{k} A_j - \sum_{j=0}^{k} B_j \right| : k = 0, 1, 2, \dots \right\}$$
(2)

where $A = (A_0, A_1, A_2, ...)$ and $B = (B_0, B_1, B_2, ...)$ are elements of **S**. (See Figure 1.) The ranking is not a natural part of the problem, but it facilitates studying the model.

At each step after applying a random transposition two types of changes may happen in the cycle structure: merging of two distinct cycles or splitting of a cycle in two. While different transpositions $(T_j^{(n)})_{j=1}^i$ are applied (meaning that the stirring particle chooses a new element in each step until *i*), the cycle decomposition of $\pi^{(n)}(i)$ contains only fixed points and the active cycle, which increases by one in each step: $\mathbf{C}^{(n)}(i) = (i+1,0,0,\ldots)$. If a transposition recurs, then the cycle splits in two, one of which will be the new active cycle. If there are already more than one non-trivial cycles in the decomposition, then the active cycle can merge another cycle. (See Figure 2.) The model realizes a coagulation-fragmentation process.

A reduction of the problem is to study the coagulation and fragmentation events of the cycles together, because both of these events happen when the stirring element steps to a place already visited. We investigate this simpler question first. Then we introduce a continuous time process on \mathbf{S} , which turns out to be the limit process. The convergence is proved by coupling. In Section 4. we show that the stationary distribution of the underlying split-and-merge transformation is the adequate modification of the Poisson-Dirichlet distribution. (See the definition later.)

A similar model is studied by Schramm in [11]. He chooses $(T_i)_{i=1}^{\infty}$ to be independent random transpositions with uniform distribution from all possible transpositions of the set [n]. The limit distribution of the proportions of the giant cycles in the permutation $\Pi(t) = T_t \circ \cdots \circ T_1$ after t = cn steps as $n \to \infty$ is identified (where c > 1/2 is a constant).

This result is in accordance with the classical theory of the random graphs derived from Erdős [6]. Let us consider the random graph G(t) on the vertex set



Figure 2: Coagulation and fragmentation of cycles

[n] where $\{u, v\}$ is an edge in it if and only if the transposition (u, v) appears in $\{T_1, \ldots, T_t\}$. By the Erdős-Rényi Theorem [7] the graph G(t) has a giant connected component only in the case t/n = c > 1/2 similarly to the condition on the random permutations. (For random graphs and random graph processes see [8] and [12]).

In Schramm's paper the vector of the cycle sizes of $\Pi(t)$ in non-increasing order normalized by the magnitude of the giant connected component of G(t) converges in distribution to Poisson-Dirichlet distribution with parameter 1 after t = cn steps (c > 1/2) as $n \to \infty$. That is the limit distribution of the relative cycle sizes in a random permutation chosen uniformly from all permutations of [n] as $n \to \infty$. Thus for large n the permutation $\Pi(t)$ behaves on the giant connected component of the Erdős-Rényi graph G(t) as a uniform permutation.

Our paper is motivated by Tóth in connection with the quantum-physical applications of the problem [13]. Angel analysed Tóth's random walk model on regular trees in [1]. For similar random stirring models see also [2] and [5].

2. Return times of the stirring particle

The movement of the stirring particle is a random walk $(B_i^{(n)})_{i=0}^{\infty}$ on the set [n], which is homogeneous in space and time. Let

$$V_i^{(n)} := \#\{k : k \le i, \exists j < k : B_j^{(n)} = B_k^{(n)}\}$$
(3)

be the number of the *returns* until the *i*th step to places already visited by the random walk $(B_j^{(n)})_{j=0}^{\infty}$. We also include those steps when the stirring particle keeps its place.

After the *i*th step the stirring element has already visited exactly $i + 1 - V_i^{(n)}$ places (including the starting point), so the transition probabilities of the Markovchain $(V_i^{(n)})_{i=0}^{\infty}$ are

$$\mathbb{P}\left(V_{i+1}^{(n)} - V_i^{(n)} = 1|V_i^{(n)}\right) = 1 - \mathbb{P}\left(V_{i+1}^{(n)} - V_i^{(n)} = 0|V_i^{(n)}\right) = \frac{i+1-V_i^{(n)}}{n}.$$
 (4)

In order to get a non-trivial limit distribution the time of the processes should be accelerated. As opposed to Schramm [11], in Theorem 1 the scaling is \sqrt{n} . This means that we describe the beginning of the evolution, because after $\mathcal{O}(\sqrt{n})$ steps the bulk of the elements is still unchanged. Simultaneously we normalize the cycle sizes with \sqrt{n} and we let $n \to \infty$.

From now on we investigate the limit of the vectors $\left(\frac{\mathbf{C}^{(n)}(\lfloor\sqrt{nt}\rfloor)}{\sqrt{n}}\right)_{t\geq 0}$ as $n \to \infty$, where the division is meant coordinatewise, namely $\frac{\mathbf{C}^{(n)}(\lfloor\sqrt{nt}\rfloor)}{\sqrt{n}} := \left(\frac{C_0^{(n)}(\lfloor\sqrt{nt}\rfloor)}{\sqrt{n}}\right)$, $\frac{C_1^{(n)}(\lfloor\sqrt{nt}\rfloor)}{\sqrt{n}},\ldots$). Elementary calculations, similar to the classical birthday problem, give the following limit distribution of the returns. For limit theorems related to generalizations of the birthday problem see also [3].

Proposition 1. Let $(V_t)_{t\geq 0}$ be an inhomogeneous Poisson point process with intensity $\rho(t) = t$. Then

$$(V_{\lfloor\sqrt{n}t\rfloor}^{(n)})_{t\geq 0} \stackrel{\mathrm{d}}{\Rightarrow} (V_t)_{t\geq 0} \qquad (n\to\infty)$$
(5)

in terms of the finite dimensional marginal distributions.

3. Coupling

Much more can be stated for the above model. Not only $(V_{\lfloor\sqrt{n}t\rfloor}^{(n)})_{t\geq 0}$, but the sequence of the processes $\left(\frac{\mathbf{C}^{(n)}(\lfloor\sqrt{n}t\rfloor)}{\sqrt{n}}\right)_{t\geq 0}$ converges. Moreover by means of coupling a stronger type of convergence is realized.

The limit process is a natural continuous extension of the discrete processes $(\mathbf{C}^{(n)}(i))_{i=0}^{\infty}$. For large *n* the active coordinate $C_0^{(n)}(i)$ increases in the bulk of the steps (when no split or merge occurs). In the times of jumps of $(V_i^{(n)})_{i=0}^{\infty}$ a split or a merge happens depending on the proportions of the cycle sizes as follows. The probability of a split in the *i*th step, conditionally given that the stirring particle returns to a place already visited, is

$$\frac{C_0^{(n)}(i-1)}{\sum_{m=0}^{\infty} C_m^{(n)}(i-1)}.$$
(6)

The conditional probability of the merge of the jth cycle and the active one is

$$\frac{C_j^{(n)}(i-1)}{\sum_{m=0}^{\infty} C_m^{(n)}(i-1)}.$$
(7)

We define an **S** valued continuous time stochastic process $\mathbf{C}(t) = (C_0(t), C_1(t), C_2(t), \ldots)$ with càdlàg paths, which imitates the above process. It is built on a Poisson point process $(V_t)_{t\geq 0}$ with intensity $\rho(t) = t$. Similarly to the discrete processes $(\mathbf{C}^{(n)}(i))_{i=0}^{\infty}$ at the times of jumps of $(V_t)_{t\geq 0}$ a split or a merge event occurs with probability proportional to the coordinates of **C**.

The initial state is $\mathbf{C}(0) := (0, 0, 0, ...)$. The evolution of the process is the following: the coordinate $C_0(t)$ increases with constant speed 1 between the jumps of $(V_t)_{t\geq 0}$. Let τ_k be the *k*th time of jump of $(V_t)_{t\geq 0}$, in other words $V_{\tau_k} = k$ and $V_{\tau_k-\varepsilon} = \lim_{\varepsilon \downarrow 0} V_{\tau_k-\varepsilon} = k - 1$. Let $(U_k)_{k=1}^{\infty}$ be i.i.d. random variables with uniform distribution on [0, 1] independent of $(V_t)_{t\geq 0}$. One of the next two actions occurs at time τ_k .

1. Split: If

$$U_k \le \frac{C_0(\tau_k -)}{\sum_{m=0}^{\infty} C_m(\tau_k -)},$$
(8)

then let $C_0(\tau_k) := U_k \sum_{m=0}^{\infty} C_m(\tau_k-)$, and the sequence $(C_m(\tau_k))_{m=1}^{\infty}$ will be the collection of $(C_m(\tau_k-))_{m=1}^{\infty}$ and $C_0(\tau_k-) - U_k \sum_{m=0}^{\infty} C_m(\tau_k-)$ rearranged in decreasing order.

2. Merge: Otherwise a unique index $j \ge 1$ can be chosen a.s. via

$$\frac{\sum_{m=0}^{j-1} C_m(\tau_k -)}{\sum_{m=0}^{\infty} C_m(\tau_k -)} < U_k \le \frac{\sum_{m=0}^{j} C_m(\tau_k -)}{\sum_{m=0}^{\infty} C_m(\tau_k -)}.$$
(9)

Let $C_0(\tau_k) := C_0(\tau_k -) + C_j(\tau_k -)$, and $C_m(\tau_k) := C_m(\tau_k -)$ if $1 \le m < j$, and $C_m(\tau_k) := C_{m+1}(\tau_k -)$ if $m \ge j$ restoring the decreasing order.

Observe that $\sum_{m=0}^{\infty} C_m(t) = t$, but we did not use it to simplify the formulas (8) and (9) in the above definition because the analogous discrete assertion is not true, compare with (6) and (7).

The main result of this paper is that the normalized discrete processes converge in probability to $(\mathbf{C}(t))_{t\geq 0}$ in the following uniform sense in terms of the distance defined by (2).

Theorem 1. There exists a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, on which the discrete processes $(\mathbf{C}^{(n)}(i))_{i=0}^{\infty}$ n = 1, 2, ... and the continuous time process $(\mathbf{C}(t))_{t\geq 0}$ can be jointly realized so that if T > 0 is fixed and f(n) is any function tending to infinity with n, then

$$\mathbb{P}\left(\sup_{0 \le t \le T} d\left(\mathbf{C}(t), \frac{\mathbf{C}^{(n)}(\lfloor\sqrt{n}t\rfloor)}{\sqrt{n}}\right) < \frac{f(n)}{\sqrt{n}}\right) \to 1 \quad as \ n \to \infty.$$
(10)

3.1. The convergence of the return process

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be such a probability space where a Poisson point process $(V_t)_{t\geq 0}$ with intensity $\rho(t) = t$ and the i.i.d. random variables $(U_k)_{k=1}^{\infty}$ and $(Z_i^{(n)})_{i,n=1}^{\infty}$ with uniform distribution on [0, 1] are given independently of each other.

We have constructed the process $(\mathbf{C}(t))_{t\geq 0}$ from $(V_t)_{t\geq 0}$ and $(U_k)_{k=1}^{\infty}$ earlier. We first re-create the processes $(V_i^{(n)})_{i=0}^{\infty}$ with the appropriate distributions on the new probability space $(\Omega, \mathcal{F}, \mathbb{P})$. The main idea of the construction is that we observe the process $(V_t)_{t\geq 0}$ in $\frac{1}{\sqrt{n}}$ long time intervals.

Let $X_i^{(n)} := \mathbb{1}\left(V_{\frac{i}{\sqrt{n}}} - V_{\frac{i-1}{\sqrt{n}}} \ge 1\right)$ i = 1, 2... n = 1, 2, ... be the indicators of the increase of the process $(V_t)_{t \ge 0}$, which are Bernoulli random variables with respective parameters

$$p_i^{(n)} = 1 - \exp\left(-\frac{2i-1}{2n}\right) = \frac{i}{n} + \mathcal{O}\left(\frac{i^2}{n^2}\right). \tag{11}$$

The required parameter for the increase of $V_i^{(n)}$ is

$$q_i^{(n)} = \frac{i}{n} - \frac{V_{i-1}^{(n)}}{n}.$$
(12)

We define the values of $V_i^{(n)}$ for fixed n with induction on i. Let $V_0^{(n)} := 0$ $n = 1, 2, \ldots$ and

$$Y_{i}^{(n)} := X_{i}^{(n)} - \mathbb{1}\left(p_{i}^{(n)} > q_{i}^{(n)}\right) \mathbb{1}\left(X_{i}^{(n)} = 1\right) \mathbb{1}\left(Z_{i}^{(n)} > \frac{q_{i}^{(n)}}{p_{i}^{(n)}}\right) \\ + \mathbb{1}\left(p_{i}^{(n)} < q_{i}^{(n)}\right) \mathbb{1}\left(X_{i}^{(n)} = 0\right) \mathbb{1}\left(Z_{i}^{(n)} < \frac{q_{i}^{(n)} - p_{i}^{(n)}}{1 - p_{i}^{(n)}}\right).$$
(13)

We define $V_i^{(n)} := V_{i-1}^{(n)} + Y_i^{(n)}$.

It is easy to see that the distribution of the new $(V_i^{(n)})_{i=0}^{\infty}$ is in accordance with (4). Later on we say that a *correction* happens if the products of the indicators in (13) do not disappear. We will see that the total probability that a correction ever occurs is small if n is large enough. This gives an alternative proof of Proposition 1.

Lemma 1. Let T > 0 be fixed and denote $0 = \tau_0, \tau_1, \ldots, \tau_{\kappa}$ the random times of jumps of the process $(V_t)_{0 \le t \le T}$ and denote $0 = \tau_0^{(n)}, \tau_1^{(n)}, \ldots, \tau_{\kappa^{(n)}}^{(n)}$ that of the discrete process $(V_{\lfloor\sqrt{nt}\rfloor}^{(n)})_{0 \le t \le T}$ defined above. Then for sufficiently large n with probability close to 1 the number of the jumps are equal: $\kappa = \kappa^{(n)}$. Furthermore, there exists a bijection between the jumps of the processes in such a way that

$$|\tau_k - \tau_k^{(n)}| \le \frac{1}{\sqrt{n}} \quad k = 1, \dots, \kappa \tag{14}$$

holds with large probability.

For technical convenience we introduce the following events for fixed $\varepsilon, \delta > 0$:

$$E_{\varepsilon} := \{ V_{\lfloor \sqrt{n}T \rfloor}^{(n)} \le K_{\varepsilon} \quad n = N_{\varepsilon}, N_{\varepsilon} + 1, \dots \},$$
(15)

where K_{ε} is a sufficiently large constant and N_{ε} is a threshold satisfying $\mathbb{P}(E_{\varepsilon}) \geq 1 - \varepsilon$. It makes sense by Proposition 1. Let

$$M_{\delta} := \{ \min_{k:\tau_k \le T} \{ \tau_k - \tau_{k-1} \} > \delta \} \cap \{ V_T - V_{T-\delta} = 0 \}$$
(16)

where τ_k is the time of the *k*th jump of the process $(V_t)_{0 \le t \le T}$ and $\tau_0 = 0$. It is elementary that $\lim_{\varepsilon \downarrow 0} \mathbb{P}(E_{\varepsilon}) = \lim_{\delta \downarrow 0} \mathbb{P}(M_{\delta}) = 1$.

Proof of Lemma 1: By (16), on the event M_{δ} the increment of the process $(V_t)_{0 \le t \le T}$ on any interval $\left[\frac{i}{\sqrt{n}}, \frac{i+1}{\sqrt{n}}\right]$ does not exceed 1 if $n > \frac{1}{\delta^2}$, hence $V_{\frac{i}{\sqrt{n}}} - V_{\frac{i-1}{\sqrt{n}}} = X_i^{(n)}$. Since $V_{\frac{i-1}{\sqrt{n}}}^{(n)} - V_{\frac{i-1}{\sqrt{n}}}^{(n)} = Y_i^{(n)}$, it is enough to prove that

$$\mathbb{P}(\{\exists i \leq \lfloor \sqrt{n}T \rfloor : X_i^{(n)} \neq Y_i^{(n)}\} \cap E_{\varepsilon} \cap M_{\delta}) \to 0 \quad (n \to \infty)$$
(17)

for all fixed $\varepsilon, \delta > 0$.

On the event E_{ε} , $X_i^{(n)} = 1$ can be true for at most K_{ε} many indices *i*. So the probability of the correction in the cases $p_i^{(n)} > q_i^{(n)}$ satisfies

$$1 - \frac{q_i^{(n)}}{p_i^{(n)}} = \mathcal{O}\left(\frac{1}{\sqrt{n}}\right) \quad (n \to \infty)$$
(18)

using the power series of the exponential function and the equations (11) and (12) estimating $p_i^{(n)}$ and $q_i^{(n)}$. If we add this at most K_{ε} many times, then the sum still goes to 0 as $n \to \infty$. A similar calculation shows that for an *i*, for which $p_i^{(n)} < q_i^{(n)}$ holds, the probability of the correction is at most

$$\frac{q_i^{(n)} - p_i^{(n)}}{1 - p_i^{(n)}} = \mathcal{O}\left(\frac{1}{n}\right) \quad (n \to \infty).$$

$$\tag{19}$$

Summing up for $i = 1, ..., \lfloor \sqrt{nT} \rfloor$ the total probability still tends to 0, as required.

3.2. Splits and merges

With the processes $(V_i^{(n)})_{i=0}^{\infty}$ we have determined when a split or a merge occurs, our task is now to define how it should happen. Similarly to the definition of the limit process $(\mathbf{C}^{(n)}(i))_{i\geq 0}^{\infty}$ we can prescribe the evolution of the discrete processes $(\mathbf{C}^{(n)}(i))_{i=0}^{\infty}$ with the use of the same independent uniform random variables $(U_k)_{k=1}^{\infty}$ as follows. Let $C_0^{(n)}(0) := 1$, $C_m^{(n)}(0) := 0$ $m = 1, 2, \ldots$ The evolution of the process $\mathbf{C}^{(n)}$ in the steps $i = 1, 2, \ldots$ is described below:

- if $V_i^{(n)} V_{i-1}^{(n)} = 0$, then $C_0^{(n)}(i) := C_0^{(n)}(i-1) + 1$ and other coordinates unchanged,
- if $V_i^{(n)} V_{i-1}^{(n)} = 1$ and $V_i^{(n)} = k$, then the uniform random variable U_k determines a unique index j with probability 1 as in (9) via

$$\frac{\sum_{m=0}^{j-1} C_m^{(n)}(i-1)}{\sum_{m=0}^{\infty} C_m^{(n)}(i-1)} < U_k \le \frac{\sum_{m=0}^{j} C_m^{(n)}(i-1)}{\sum_{m=0}^{\infty} C_m^{(n)}(i-1)}.$$
(20)

Similarly to the definition of the limit process

- 1. j = 0: split. If $U_k \sum_{m=0}^{\infty} C_m^{(n)}(i-1) < 1$, then let everything be unchanged: $\mathbf{C}^{(n)}(i) := \mathbf{C}^{(n)}(i-1)$, let us call this case fictive split (corresponding to the event that the stirring particle keeps its place). Otherwise $C_0^{(n)}(i) := [U_k \sum_{m=0}^{\infty} C_m^{(n)}(i-1)]$, let the broken fragment $C_0^{(n)}(i-1) - [U_k \sum_{m=0}^{\infty} C_m^{(n)}(i-1)]$ add to the collection of nonactive pieces $(C_m^{(n)}(i-1))_{m=1}^{\infty}$ to form the new ranked sequence $(C_m^{(n)}(i))_{m=1}^{\infty}$.
- 2. j > 0: merge. Let $C_0^{(n)}(i) := C_0^{(n)}(i-1) + C_j^{(n)}(i-1)$ and for the re-ranking $C_m^{(n)}(i) := C_m^{(n)}(i-1)$ if 0 < m < j, and $C_m^{(n)}(i) := C_{m+1}^{(n)}(i-1)$ if $m \ge j$.

It is easy to show that this new definition of $(\mathbf{C}^{(n)}(i))_{i=0}^{\infty}$ provides the same distribution as in the model generated by transpositions, so we prove the convergence for these processes.

Proof of Theorem 1: Let $\varepsilon, \delta > 0$ be fixed. Let A_n denote the event that the assertion of Lemma 1 holds for $(V_{\lfloor \sqrt{n}t \rfloor}^{(n)})_{0 \le t \le T}$. We restrict ourselves to the events $E_{\varepsilon} \cap M_{\delta} \cap A_n$. Let us define a measure (which is not a probability measure) on the sets $B \in \mathcal{F}$:

$$\mathbb{P}_{\varepsilon,\delta,n}(B) := \mathbb{P}(B \cap E_{\varepsilon} \cap M_{\delta} \cap A_n).$$
(21)

By Lemma 1 it is enough to show that for fixed $\varepsilon, \delta > 0$ the processes $\mathbf{C}(t)$ and $\frac{\mathbf{C}^{(n)}(\lfloor \sqrt{nt} \rfloor)}{\sqrt{n}}$ are sufficiently close to each other for large n except a set with $\mathbb{P}_{\varepsilon,\delta,n}$ -measure tending to 0 as $n \to \infty$. The proof consists of the following steps:

- 1. We estimate the increase of the distance between $\mathbf{C}(t)$ and $\frac{\mathbf{C}^{(n)}(\lfloor\sqrt{n}t\rfloor)}{\sqrt{n}}$ between two successive split or merge events.
- 2. We introduce those events when the distance under discussion cannot be estimated: the awkward events (defined later) and the fictive splits. We show that they have small probability.
- 3. On the complementer event, which has probability tending to 1 as $n \to \infty$, we show that a merge does not increase the distance between $\mathbf{C}(t)$ and $\frac{\mathbf{C}^{(n)}(\lfloor\sqrt{n}t\rfloor)}{\sqrt{n}}$ very much.

- 4. We do this also for the splits.
- 5. We summarize the estimates.

Step 1. Let

$$d_k^- := d\left(\mathbf{C}(\tau_k -), \frac{\mathbf{C}^{(n)}(\lfloor \sqrt{n\tau_k^{(n)}} \rfloor - 1)}{\sqrt{n}}\right), \quad d_k^+ := d\left(\mathbf{C}(\tau_k), \frac{\mathbf{C}^{(n)}(\lfloor \sqrt{n\tau_k^{(n)}} \rfloor)}{\sqrt{n}}\right)$$
(22)

denote the distance between the discrete and continuous processes before and after the time of the kth split or merge. (Recall that τ_k is the time of the kth jump of $(V_t)_{0 \le t \le T}$ and $\tau_k^{(n)}$ is that of $(V_{\lfloor \sqrt{nt} \rfloor}^{(n)})_{0 \le t \le T}$, which are close $\mathbb{P}_{\varepsilon,\delta,n}$ -almost surely by Lemma 1.)

While no split or merge occurs, the distance between the processes does not increase very much. From Lemma 1 the difference between τ_k and $\tau_k^{(n)}$ can be at most $\frac{1}{\sqrt{n}}$. The discrete processes $\left(\frac{\mathbf{C}^{(n)}(\lfloor\sqrt{n}t\rfloor)}{\sqrt{n}}\right)_{t\geq 0}$ change only in the times which are multiples of $\frac{1}{\sqrt{n}}$. Thus $\mathbb{P}_{\varepsilon,\delta,n}$ -almost surely

$$d_k^- \le d_{k-1}^+ + \frac{2}{\sqrt{n}}.$$
(23)

STEP 2. From now on we investigate only the split or merge points of the processes. At the kth time of jump of $(V_t)_{0 \le t \le T}$ and $(V_{\lfloor \sqrt{nt} \rfloor}^{(n)})_{0 \le t \le T}$ we choose with the help of U_k one of the components of $\mathbf{C}(\tau_k-)$ and $\mathbf{C}^{(n)}(\lfloor \sqrt{n\tau_k^{(n)}} \rfloor - 1)$ via (9) and (20). Let us call the possibility that these components are of different indices an awkward event. If an awkward event or a fictive split (meaning that $U_k \sum_{m=0}^{\infty} C_m^{(n)}(i-1) < 1$) occurs, then we cannot estimate $d(\mathbf{C}, \mathbf{C}^{(n)}/\sqrt{n})$. We will see that these events have probability tending to 0 as $n \to \infty$.

We can choose the components of **C** and $\mathbf{C}^{(n)}$ as follows. We set the coordinates of the vector $\mathbf{C}(\tau_k-)/\sum_{m=0}^{\infty} C_m(\tau_k-)$ to the real line from the origin one after another, which gives a partition of the unit interval [0, 1]. We do this also with the coordinates of

$$\frac{\mathbf{C}^{(n)}(\lfloor\sqrt{n}\tau_k^{(n)}\rfloor-1)}{\sum_{m=0}^{\infty}C_m^{(n)}(\lfloor\sqrt{n}\tau_k^{(n)}\rfloor-1)}.$$
(24)

Let W_k denote the set of those points in [0, 1] which are covered by the coordinates of C and $C^{(n)}$ of different indices. The probability of the awkward events (which is an upper estimate for their $\mathbb{P}_{\varepsilon,\delta,n}$ -measure) is exactly the Lebesgue measure of W_k . We know that $\sum_{m=0}^{\infty} C_m(t) = t$ for all $t \ge 0$. From the construction

$$\frac{\lfloor \sqrt{n}t \rfloor - K_{\varepsilon}}{\sqrt{n}} \le \frac{\sum_{m=0}^{\infty} C_m^{(n)}(\lfloor \sqrt{n}t \rfloor)}{\sqrt{n}} \le t \quad \text{if} \quad t \in [0, T],$$
(25)

because at the split or merge points (occurring at most K_{ε} many times) the total length of the discrete process does not increase. From this

$$\left|\sum_{m} C_{m}(\tau_{k}-) - \frac{\sum_{m} C_{m}^{(n)}(\lfloor \sqrt{n}\tau_{k}^{(n)} \rfloor - 1)}{\sqrt{n}}\right|$$
$$\leq \left|\tau_{k} - \frac{\lfloor \sqrt{n}\tau_{k}^{(n)} \rfloor - 1 - K_{\varepsilon}}{\sqrt{n}}\right| \leq \frac{K_{\varepsilon}}{\sqrt{n}} + \frac{3}{\sqrt{n}}$$
(26)

follows using Lemma 1.

From the above it is an elementary exercise to show that the distance between the corresponding dividing points of the partitions of [0, 1] generated by $\mathbf{C}(\tau_k -)/\tau_k$ and by the vector (24) can be, respectively, at most

$$\frac{d_k^- + \frac{K_\varepsilon + 3}{\sqrt{n}}}{\tau_k}.$$

Since the number of coordinates is at most K_{ε} , this provides the following upper bound:

$$\operatorname{Leb}(W_k) \le \frac{d_k^- + \frac{K_\varepsilon + 3}{\sqrt{n}}}{\tau_k} K_\varepsilon \le \frac{d_k^- + \frac{K_\varepsilon + 3}{\sqrt{n}}}{\delta} K_\varepsilon, \tag{27}$$

where we used the fact that $\tau_k = \sum_{m=0}^{\infty} C_m(\tau_k) \ge \delta$ holds for $k = 1, 2, \ldots$ on the event M_{δ} . This yields

$$\mathbb{P}_{\varepsilon,\delta,n}(\text{awkward event at }\tau_k) \le \operatorname{Leb}(W_k) \le \frac{d_k^- + \frac{K_\varepsilon + 3}{\sqrt{n}}}{\delta} K_\varepsilon.$$
(28)

Furthermore

$$\mathbb{P}_{\varepsilon,\delta,n}(\text{fictive split at } \tau_k^{(n)}) \le \frac{\frac{1}{\sqrt{n}}}{\sum_{m=0}^{\infty} C_m^{(n)} (\lfloor \sqrt{n} \tau_k^{(n)} \rfloor - 1) / \sqrt{n}} \le \frac{2}{\delta\sqrt{n}}, \qquad (29)$$

if n is large enough by (25). So we conclude that

 $\mathbb{P}_{\varepsilon,\delta,n}$ (awkward event or fictive split at the kth split or merge point)

$$\leq \frac{K_{\varepsilon}}{\delta} d_k^- + \frac{K_{\varepsilon}^2 + 3K_{\varepsilon} + 2}{\delta\sqrt{n}}.$$
(30)

STEP 3. In the case when the random variable U_k chooses the same components of **C** and **C**⁽ⁿ⁾ and it is not the active coordinate, i.e. there is a merge in both processes (see Figure 3), then

$$d_k^+ \le 3d_k^-. \tag{31}$$



Figure 3: The piece C_2 merges C_0 parallel with the $C_2^{(n)} - C_0^{(n)}$ coagulation



Figure 4: Split: the broken pieces from the coordinate 0 are X and X' which have to be moved to places A' and B

STEP 4. If a (non-fictive) split occurs in the discrete and continuous processes, then using inequality (26) we have

$$\left| C_0(\tau_k) - \frac{C_0^{(n)}(\lfloor \sqrt{n}\tau_k^{(n)} \rfloor)}{\sqrt{n}} \right|$$

$$\leq \left| U_k \sum_{m=0}^{\infty} C_m(\tau_k -) - \frac{\lfloor U_k \sum_{m=0}^{\infty} C_m^{(n)}(\lfloor \sqrt{n}\tau_k^{(n)} \rfloor - 1) \rfloor}{\sqrt{n}} \right| \leq \frac{K_{\varepsilon} + 3}{\sqrt{n}} + \frac{1}{\sqrt{n}}. \quad (32)$$

This is why the broken pieces from $C_0(\tau_k-)$ and $C_0^{(n)}(\lfloor\sqrt{n}\tau_k^{(n)}\rfloor-1)/\sqrt{n}$ (denoted by X and X' on Figure 4) can differ at most $d_k^- + \frac{K_k+4}{\sqrt{n}}$: the difference can be $\frac{K_k+4}{\sqrt{n}}$ between the left end points and at most d_k^- between the right end points.

It is possible that the two broken pieces do not come to the same place in the decreasing order of the coordinates. This case is shown on Figure 4. Then we move first both X and X' to the closer of the final places of them in the decreasing order (to the places A and A' on the figure). Because $|A - A'| \leq d_k^-$, the result is two vectors (the modifications of $\mathbf{C}(\tau_k-)$ and $\mathbf{C}^{(n)}(\lfloor\sqrt{n}\tau_k^{(n)}\rfloor-1)/\sqrt{n}$, but one of them is not necessarily in decreasing order), which have $d(\cdot, \cdot)$ -distance at most $2d_k^- + \frac{K_k+4}{\sqrt{n}}$ more then before this modification.

In the second step we move X from A to B (see Figure 4). The lengths of the parts between A and B are at least |X| and at most $|X'| + 2d_k^- \leq |X| + 3d_k^- + \frac{K_{\varepsilon}+4}{\sqrt{n}}$. So any two of these parts have lengths differing at most $3d_k^- + \frac{K_{\varepsilon}+4}{\sqrt{n}}$. Swapping X always with its right neighbour until hitting place B, the number of the swaps is at most K_{ε} , and at each swap the distance can increase at most $3d_k^- + \frac{K_{\varepsilon}+4}{\sqrt{n}}$, so we

have

$$d_k^+ \le 2d_k^- + \frac{K_\varepsilon + 4}{\sqrt{n}} + K_\varepsilon \left(3d_k^- + \frac{K_\varepsilon + 4}{\sqrt{n}}\right). \tag{33}$$

STEP 5. Summing up the estimates (23), (31) and (33) we get easily the following recursive bound:

$$d_{k}^{+} \leq \max(3, 2+3K_{\varepsilon})d_{k-1}^{+} + \frac{K_{\varepsilon}^{2} + 5K_{\varepsilon} + 4 + 2\max(3, 2+3K_{\varepsilon})}{\sqrt{n}} =: ad_{k-1}^{+} + \frac{b}{\sqrt{n}}.$$
(34)

Hence $\sup_{0 \le k \le K_{\varepsilon}} d_k^{\pm} \le \left(\sum_{i=0}^{K_{\varepsilon}} ba^i\right) \frac{1}{\sqrt{n}}$. Considering the results of steps 1 and 2 the assertion of the theorem follows.

4. Stationary distribution and generalizations

It is a natural question to identify the stationary distribution of our stirring process. This means that we look at the asymptotic behaviour of the process $(\mathbf{C}^{(n)}(\lfloor nt \rfloor)/n)_{t\geq 0}$. Observe that the time scale is of order n, i.e. the time scale when the stirring element has already visited the bulk of the n places. This setup is the same as that of the problem studied by Schramm in [11], but different from the phenomenon described by Theorem 1.

In this section we consider the following split-and-merge transformation corresponding to the stirring generated by random transpositions. Let $\mathbf{C} = (C_0, C_1, C_2, \ldots) \in \mathbf{S}$ be a random probability distribution, i.e. $\sum_m C_m = 1$ almost surely. C_0 is the active component. Let U be a random variable with uniform distribution of [0, 1] which is independent of \mathbf{C} . If $U \leq C_0$, then the C_0 splits, i.e. the new active component will be U and $(C_0 - U, C_1, C_2, \ldots)$ will be the remaining components after restoring the decreasing order. If $\sum_{m=0}^{j-1} C_m < U \leq \sum_{m=0}^{j} C_m$, then C_0 merges with C_j similarly to (6-9) because $\sum_m C_m = 1$.

In limit theorems of random partitions and permutations the following distribution appears often. Let the random variables W_1, W_2, \ldots be independent with uniform distribution on [0, 1]. Let (Q_1, Q_2, \ldots) be the decreasing rearrangement of the random variables

$$(P_1, P_2, \dots) := (W_1, (1 - W_1)W_2, (1 - W_1)(1 - W_2)W_3, \dots).$$

Then the random sequence $(P_1, P_2, ...)$ has GEM(1) distribution after Griffiths, Engen and McCloskey. $(Q_1, Q_2, ...)$ has *Poisson – Dirichlet* distribution with parameter 1, abbreviated PD(1). For more about this family of distributions see [9].

Let $(p_1, p_2, ...)$ be a random probability distribution. We construct its size biased permutation. Let $U_1, U_2, ...$ be i.i.d. uniform random variables on [0, 1] independently of $(p_1, p_2, ...)$. Let I_j be the unique index for which $\sum_{i=1}^{I_j-1} p_i \leq U_j < \sum_{i=1}^{I_j} p_i$. Let J_k denote the kth smallest integer m satisfying $I_m \notin \{I_1, I_2, ..., I_{m-1}\}$. Then the vector $(p_{J_1}, p_{J_2}, ...)$ is called the size biased permutation of $(p_1, p_2, ...)$. It is well known that the size biased permutation of a random partition with PD(1) distribution has GEM(1) distribution. See also [10].

Consider the following probability distribution on **S**. Let $(Q_1, Q_2, ...)$ have PD(1) distribution. Let C_0 be a size biased part from $(Q_1, Q_2, ...)$ (i.e. the first component of the size biased permutation of $(Q_1, Q_2, ...)$) corresponding to the active cycle and the rest $(C_1, C_2, ...)$ is the vector of the remaining Q_j -s in non-increasing order. We denote by μ the distribution of $\mathbf{C} = (C_0, C_1, C_2, ...)$.

Theorem 2. The distribution μ is invariant under the above split-and-merge transformation.

Proof: By definition a random partition \mathbf{C} with distribution μ can be considered as follows. Let W_1, W_2, \ldots be i.i.d. uniform random variables on [0, 1] as in the definition of PD(1). Because the size biased permutation of PD(1) is GEM(1), we can suppose that for the active component $C_0 = W_1$ holds and (C_1, C_2, \ldots) is the decreasing rearrangement of $((1 - W_1)W_2, (1 - W_1)(1 - W_2)W_3, \ldots)$. Let ν be the distribution of the random partition obtained by the application of a stirring step to \mathbf{C} .

If $U < W_1$ for the [0,1]-uniform random variable U, then the new non-active components are $(W_1 - U, (1 - W_1)W_2, (1 - W_1)(1 - W_2)W_3, ...)$ in decreasing order. Conditionally on $\{U < W_1\}$ and on U, the variable W_1 is uniform on [U, 1], thus the vector of the non-active components has PD(1) distribution scaled by (1 - U). It yields that ν conditioned on $\{U < W_1\}$ and on U is the same as μ conditioned on the active component having size U.

If $U > W_1$, then a coagulation occurs. Conditioned on $\{U > W_1\}$ and on the value of W_1 , the size of the component which merges C_0 has uniform distribution on $[0, 1 - W_1]$, because it is a size biased component. We get the same distribution, if we choose this component merging C_0 to be of length $U - W_1$. Conditionally on $\{U > W_1\}$ and on U the rest has PD(1) distribution scaled by (1 - U). Thus, a sample from ν conditioned on $\{U > W_1\}$ and on U has an active coordinate of size U and the remaining components with a scaled PD(1) distribution.

Hence, a vector with distribution ν can be obtained by sampling U uniformly on [0, 1], taking the active coordinate of length U and taking a scaled PD(1) distribution on the rest. It shows that $\nu = \mu$, as required.

Theorem 2 proves that μ is a stationary measure for our process, but it is not at all clear if this is the *unique* stationary measure. The proof of this would be the analogue of Schramm's result in [11].

A possible generalization of the model studied in this paper is the *multiple stir*ring. It means that we consider more than one stirring particles. For a fixed number k of stirring elements an analogous limit theorem can be proved with a coupling similarly to Theorem 1. The case, if the number of the stirring elements depends on the size of the set [n], might also be worth studying (for example with $k(n) = n^{\alpha}$ where $0 < \alpha < 1$). Of course, we need different scaling of time and space in this case. An open question is for our original model to establish after how much time a permutation can be regarded as a random permutation chosen with uniform distribution, if it can be regarded at all. The solution of the problem in this simply describable model is not obvious in the least. For more about this problem in similar models see [4].

Acknowledgement. I thank Bálint Tóth and Benedek Valkó for initiating these investigations and for their permanent support and useful comments while writing this paper. I am grateful to the referee for pointing out the proof of Theorem 2, and I thank to Sándor Csörgő for his helpful remarks on this paper.

References

- O. ANGEL, Random infinite permutations and the cyclic time random walk, in C. Banderier and C. Krattenthaler, editors, *Random Walks and Discrete Potential Theory*, Discrete Mathematics and Theoretical Computer Science, pages 9-16, 2003.
- [2] N. BERESTYCKI, R. DURRETT, A phase transition in the random transposition random walk, To appear in *Probability theory and related fields*, 2005.
- [3] M. CAMARRI, J. PITMAN, Limit distributions and random trees derived from the birthday problem with unequal probabilities, *Electronic Journal of Probability*, vol. 5, paper no. 2, pages 1-18, 2000.
- [4] P. DIACONIS, Group Representations in Probability an Statistics, IMS Lecture notes-Monograph Series, vol. 11, Institute of Mathematical Statistics, Hayward, California, 1988.
- [5] E. MAYER-WOLF, O. ZEITOUNI, M. ZERNER, Asymptotics of certain coagulation-fragmentation processes and invariant Poisson-Dirichlet measures, *Electronic Journal of Probability*, vol. 7, pages 1-25, 2002.
- [6] P. ERDŐS, Some remarks on the theory of graphs, Bull. Amer. Math. Soc. 53, 292-294, 1947.
- [7] P. ERDŐS, A. RÉNYI, On random graphs I. Publ. Math. Debrecen 6, 290-297, 1959.
- [8] S. JANSON, T. ŁUCZAK, A. RUCIŃSKI, Random Graphs, Wiley-Interscience Series in Discrete Mathematics and Optimization, New York, Wiley-Interscience, 2000.
- [9] J. PITMAN, Combinatorial Stochastic Processes, Technical Report no. 621, Dept. Statistics, U. C. Berkeley, Lecture notes for St. Flour course, 2002.
- [10] J. PITMAN, Poisson-Dirichlet and GEM invariant distributions for split-andmerge transformations of an interval partition, *Combinatorics, Probability and Computing* 11, 501-514, 2002.

- [11] O. SCHRAMM, Compositions of random transpositions. Israel J. Math. vol. 147, 221-244, 2005.
- [12] V. E. STEPANOV, The probability of connectedness of a random graph $\mathcal{G}_m(t)$, *Teor. Veroyatnost. i Primenen.* **15**, 55-68 (Russian); English transl. *Theor. Probab. Appl.* **15**, 55-67, 1970.
- [13] B. TÓTH, Improved lower bound on the thermodynamic pressure of the spin 1/2 Heisenberg ferromagnet. Lett. Math. Phys., 28(1):75-84, 1993.

B. VETŐ, Institute of Mathematics, Technical University Budapest, Egry J. u. 1, 1111 Budapest, Hungary; *e-mail:* vetob@math.bme.hu