Motivic interpretation of Milnor K-groups attached to Jacobian varieties

Satoshi Mochizuki *

mochi@ms.u-tokyo.ac.jp

Abstract

In the paper [Som90] p.105, Somekawa conjectures that his Milnor Kgroup $K(k, G_1, \ldots, G_r)$ attached to semi-abelian varieties G_1, \ldots, G_r over a field k is isomorphic to $\operatorname{Ext}_{\mathcal{M}_k}^r(\mathbb{Z}, G_1[-1] \otimes \ldots \otimes G_r[-1])$ where \mathcal{M}_k is a certain category of motives over k. The purpose of this note is to give remarks on this conjecture, when we take \mathcal{M}_k as Voevodsky's category of motives $\operatorname{DM}^{\operatorname{eff}}(k)$.

Key words: motivic cohomology, 1-motives, Milnor K-groups, Weil reciprocity law

Contents

0	Inti	roduction	2	
1	Mil	nor K-groups attached to semi-abelian varieties	3	
	1.1	Extension of valuations and tame symbols	3	
	1.2	Definition of the Milnor K -groups attached to semi-abelian varietie	s 4	
2	Tria	angulated categories of motives	5	
	2.1	Triangulated category of effective geometric motives	5	
	2.2	Triangulated category of effective motivic complexes	6	
	2.3	Motives with compact support	9	
	2.4	Tate object	10	
	2.5	The triangulated category of geometric motives	11	
3	Var	ious morphisms between motives	12	
	3.1	Transpose for finite equidimensional morphisms	12	
	3.2	Pull back for flat equidimensional morphisms	12	
	3.3	Motives with closed support	13	
	3.4	Thom isomorphism	14	
	3.5	Normal cone deformation	16	
	3.6	Gysin triangles	16	

 $^{^{*}}$ This research is supported by the 21 century COE program at Graduate School of Mathematical Sciences, the University of Tokyo.

		c cohomology groups attached to pointed smooth curves 1	18	
	4.1 Def	finition \ldots \ldots \ldots \ldots \ldots 18	3	
	4.2 Fur	ndamental properties 19	9	
5	Calcula	tion of motivic cohomology groups attached to pointed sm	mooth curves 20)
5		tion of motivic cohomology groups attached to pointed sm p-motives)
5	5.1 Pro))

0 Introduction

To unify the Moore exact sequence and the Bloch exact sequence, K. Kato defined the generalized Milnor K-groups attached to finite family of semiabelian varieties over a base field k in [Som90]. (See also [Akh00], [Kah92].) That is, for semi-abelian varieties G_1, \ldots, G_r over k, he associated the group $K(k, G_1, \ldots, G_r)$. (For precise definition, see 1.4) This group is a generalization of the Milnor K-group as the following example shows.

Example 0.1. In the notation above, if $G_1 = G_2 = \ldots = G_r = \mathbb{G}_m$, the following equality holds.

$$K(k, \mathbb{G}_{\mathrm{m}}, \dots, \mathbb{G}_{\mathrm{m}}) = K_r^M(k)$$

On the other hand this group is also a generalization of the Bloch group V.

Example 0.2. Let C be a projective smooth curve over k such that $C(k) \neq \phi$. We have the following equality

$$K(k, \operatorname{Jac} C, \mathbb{G}_{\mathrm{m}}) = V(C).$$

where V(C) is defined by S. Bloch (c.f. [Blo81]) as the following way

$$V(C) = \frac{\operatorname{Ker}(\bigoplus_{x \in C^1} k(x)^{\times \sum N_k(x)/k} k^{\times})}{\operatorname{Im}(K_2(K(C)) \stackrel{\oplus}{\to} \stackrel{x}{\to} \bigoplus_{x \in C^1} k(x)^{\times})}.$$

- ----

As is explained in [Som90], there is the generalized Bloch-Moore exact sequence.

0.3. Let k be a number field and A a semi-abelian variety over k. We write $G = \operatorname{Gal}(\overline{k}/k)$ and $G_v = \operatorname{Gal}(\overline{k_v}/k_v)$ for a place v. Let T(A) be the Tate module of A. S is a finite set of places including all Archimedian and places where A has bad reduction. Then $T(A)_G$ is a finite group by owing to [KL81]. Let m be a nonzero integer divisible by the order of $T(A)_G$. Somekawa proves the following generalized Moore-Bloch exact sequence (c.f. [Som90] Theorem 4.1):

$$K(k, A, \mathbb{G}_{\mathrm{m}}) \to \bigoplus_{v \notin S} T(A)_{G_v} \oplus \bigoplus_{v \in S} K(k_v, A_v, \mathbb{G}_{\mathrm{m}})/m \to T(A)_G \to 0$$

In the case of $A = \mathbb{G}_{m}$, the above exact sequence is proved by Moore (c.f. [Moo69])

$$K_2(k) \to \bigoplus_{v:\text{not complex}} \mu(k_v) \to \mu(k) \to 0.$$

In the case of A = Jac C in the notation in 0.2, the above exact sequence is proved by S. Bloch, K. Kato and S. Saito (c.f. [Blo81], [KS83])

$$V(C) \to \bigoplus_{v \notin S} T(\operatorname{Jac} C)_{G_v} \oplus \bigoplus_{v \in S} V(C \times_k k_v) / m \to T(\operatorname{Jac} C)_G \to 0.$$

In [Som90], Somekawa conjectures that the Somekawa K-groups should be motivic cohomology groups attached to semi-abelian varieties. More precisely

Conjecture 0.4. (Somekawa conjecture)

Let G_1, \ldots, G_r be semi-abelian varieties over k, then $K(k, G_1, \ldots, G_r)$ is isomorphic to $\operatorname{Ext}^r_{\mathcal{M}_k}(\mathbb{Z}, G_1[-1] \otimes \ldots \otimes G_r[-1])$, where \mathcal{M}_k is a certain category of motives over k and $G_i[-1]$ means 1-motif (c.f [Del74]).

In this paper we will examine this conjecture, if we take \mathcal{M}_k as Voevodsky's category of motives $\mathrm{DM}^{\mathrm{eff}}_{-}(k)$.

Main Theorem 0.5. (Somekawa conjecture for Jacobian varieties) Let $(C_1, a_1), \ldots, (C_n, a_n)$ be pointed projective smooth curves over perfect field k which admits resolution of singularities. Then

 $K(k, \operatorname{Jac} C_1, \ldots, \operatorname{Jac} C_n) \xrightarrow{\sim} \operatorname{Hom}_{\operatorname{DM}^{\operatorname{eff}}_{e}(k)}(\operatorname{M}_{\operatorname{gm}}(\operatorname{Spec} k), \mathbb{Z}(\bigwedge_{i=1}^n (C_i, a_1))[n]).$

0.6. In this paper, let k be a perfect field which admits resolution of singularity.

1 Milnor K-groups attached to semi-abelian varieties

1.1 Extension of valuations and tame symbols

1.1. Suppose k is a field and G is a semi-abelian variety defined over k, that is, there is an exact sequence of group schemes (viewed as sheaves in the flat topology) over k:

$$0 \to T \to G \to A \to 0$$

where T is a torus and A is an abelian variety.

1.2. In the notation above, let K/k be an algebraic function field and v a place of K/k. Let L/K_v be a finite unramified Galois extension such that $T \times_k F \xrightarrow{\sim} \mathbb{G}_m^n$ for the residue field F of L and some n; let w be the unique extension of v of L. We obtain the following commutative diagram of exact sequences defining a

map $r_w = (r_w^1, \ldots, r_w^n);$



1.3. In the notation above, we are going to construct a map

$$\partial_v : G(K_v) \otimes K_v^{\times} \to G(k(v)).$$

Fix $g \in G(K_v)$ and $h \in K_v^{\times}$. For each i = 1, ..., n, we define $h_i \in T(L)$ to be the *n*-th tuple having h in the i-th coordinate and 1 elsewhere. Then set

$$\varepsilon(g,h) = ((-1)^{\operatorname{ord}_w(h)r_w^1(g)}, \dots, (-1)^{\operatorname{ord}_w(h)r_w^n(g)}) \in T(\mathcal{O}_w) \subset G(\mathcal{O}_w)$$

and

$$\tilde{\partial_v}(g,h) = \varepsilon(g,h) g^{\operatorname{ord}_w(h)} \prod_{i=1}^n h_i^{-r_w^i(g)} \in G(\mathcal{O}_w).$$

We define the "extended tame symbol" $\partial_v(g,h)$ to be the image of $\tilde{\partial}_v(g,h)$ under the canonical map $G(\mathcal{O}_w) \to G(F)$; Then $\partial_v(g,h)$ is invariant under the action of $\operatorname{Gal}(F/k(v))$, so that it belongs to G(k(v)). This definition of ∂_v is independent of the choice of L and of the isomorphic from the torus to $\mathbb{G}_m^{\oplus n}$.

1.2 Definition of the Milnor *K*-groups attached to semiabelian varieties

1.4. Let k be a field and G_1, \ldots, G_r a finite (possibly empty) family of semiabelian varieties defined over k. We define Milnor K-groups attached to semiabelian varieties $K(k, G_1, \ldots, G_r)$ as follows. If r = 0, we write $K(k, \phi)$ for our groups and set $K(k, \phi) = \mathbb{Z}$.

For $r \geq 1$, we define

$$K(k, G_1, \dots, G_r) = F/R$$

where

$$F = \bigoplus_{E/k: \text{finite}} G_1(E) \otimes \ldots \otimes G_r(E)$$

and $R \subset F$ is the subgroup generated by the relation **R1-R2** below.

R1 For any finite extensions $k \hookrightarrow E_1 \stackrel{\phi}{\hookrightarrow} E_2$, let $g_{i_0} \in G_{i_0}(E_2)$ and $g_i \in G_i(E_1)$ for $i \neq i_0$, the relation

$$(\phi^*(g_1)\otimes\ldots\otimes g_{i_0}\otimes\ldots\otimes\phi^*(g_r))_{E_2}-(g_1\otimes\ldots\otimes N_{E_2/E_1}(g_{i_0})\otimes\ldots\otimes g_r)_{E_1}$$

(Here N_{E_2/E_1} denotes the norm map on the group scheme G_{i_0})

R2 For every algebraic function field K/k and all choices $g_i \in G_i(K)$, $h \in K^{\times}$ such that for each place v of K/k, there exists i(v) such that $g_i \in G_i(\mathcal{O}_v)$ for all $i \neq i(v)$, the relation

$$\sum_{\text{place of } K/k} (g_1(v) \otimes \ldots \otimes \partial_v(g_{i(v)}, h) \otimes \ldots \otimes g_r(v))_{k(v)/k}$$

Here $g_i(v) \in G_i(k(v))$ $(i \neq i(v))$ denotes the reduction of $g_i \in \mathcal{O}_v$ modulo m_v and $\partial_v(g_{i(v)}, h)$ is the extended tame symbol as defined in 1.3

The class in F/R of an element $a_1 \otimes \ldots \otimes a_r \in G_1(E) \otimes \ldots \otimes G_r(E)$ will be denoted $\{a_1, \ldots, a_r\}_{E/k}$.

Remark 1.5. By the relation **R1**, if ϕ is a k-isomorphism $E_1 \xrightarrow{\sim} E_2$, then $\{g_1, \ldots, g_r\}_{E_1/k} = \{\phi^*(g_1), \ldots, \phi^*(g_r)\}_{E_2/k}$. This shows that symbols form a set.

2 Triangulated categories of motives

In this section, we will briefly review the definition of the triangulated categories of motives. (c.f. [TriCa]).

2.1 Triangulated category of effective geometric motives

First we will review the construction of the category of geometric motives.

2.1. 1. Let Sm/k be the category of schemes which are separated smooth, and of finite type over k.

2. Recall the definition of the category $\operatorname{SmCor}(k)$: its objects are those of Sm/k . The set of morphism from Y to X is given by the group $\operatorname{Cor}(Y, X)$ of finite correspondences from Y to X, defined as the free abelian group on the symbol (Z), where Z runs through the integral closed subschemes of $Y \times_k X$ which are finite over Y and surjective over a connected component of Y. We will denote the object of $\operatorname{SmCor}(k)$ which corresponds to a smooth scheme X by [X].

2.2. The category $\operatorname{SmCor}(k)$ is an additive category. Consider the homotopy category $\mathcal{H}^b(\operatorname{SmCor}(k))$ of bounded complexes over $\operatorname{SmCor}(k)$. Let \mathcal{T} be the class of complexes of the following two forms:

1. For any smooth scheme X over k the complex

$$[X \times \mathbb{A}^1_k] \stackrel{[pr_1]}{\to} [X]$$

belongs to \mathcal{T} .

2. For any smooth scheme X over k and an open covering $X = U \cup V$ of X the complex

$$[U \cap V] \stackrel{[j_U] \oplus [j_V]}{\to} [U] \oplus [V] \stackrel{[i_U] \oplus (-[i_V])}{\to} [X]$$

belongs to \mathcal{T} . (here j_U , j_V , i_U , i_V are the obvious open embeddings.) Denote by $\overline{\mathcal{T}}$ the minimal thick subcategory of $\mathcal{H}^b(\mathrm{SmCor}(k))$ which contains \mathcal{T} .

The triangulated category $\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k)$ of effective geometric motives over k is the pseudo-Abelian envelope of the localization of $\mathcal{H}^{b}(\mathrm{SmCor}(k))$ with respect to the thick subcategory $\bar{\mathcal{T}}$. We denote the obvious functor $Sm/k \to \mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k)$ by M_{gm} .

2.3. For a pair of smooth schemes X, Y over k, we set

$$[X] \otimes [Y] := [X \times Y].$$

For any smooth schemes X_1 , Y_1 , X_2 , Y_2 the external product of cycles defines a homomorphism:

$$\operatorname{Cor}(X_1, Y_1) \otimes \operatorname{Cor}(X_2, Y_2) \to \operatorname{Cor}(X_1 \times X_2, Y_1 \times Y_2)$$

which gives us a definition of tensor product of morphisms in $\operatorname{SmCor}(k)$. This structure defines in the usual way a tensor category structure on $\mathcal{H}^b(\operatorname{SmCor}(k))$ which can be descended to the category $\operatorname{DM}_{\operatorname{gm}}^{\operatorname{eff}}(k)$ by the universal property of localization.

Note that the unit object our tensor structure is $M_{gm}(\operatorname{Spec} k)$. We will denote it by \mathbb{Z} .

Example 2.4. Let x, y: Spec $k \to \mathbb{P}^1_k$ be two k-rational points. Then $M_{gm}(x) = M_{gm}(y) : M_{gm}(\operatorname{Spec} k) \to M_{gm}(\mathbb{P}^1_k)$.

Proof. We take an affine open set $\mathbb{A}_k^1 \xrightarrow{j} \mathbb{P}_k^1$ which contains x and y. That is, there are \tilde{x}, \tilde{y} : Spec $k \to \mathbb{A}_k^1$ such that $x = j \circ \tilde{x}$ and $y = j \circ \tilde{y}$. Then we have $M_{gm}(x) = M_{gm}(j) \circ M_{gm}(\tilde{x}) = M_{gm}(j) \circ M_{gm}(p)^{-1} = M_{gm}(j) \circ M_{gm}(\tilde{y}) = M_{gm}(y)$, where $p : \mathbb{A}_k^1 \to \text{Spec } k$ is the structure morphism.

2.2 Triangulated category of effective motivic complexes

To study the fundamental property of $\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k)$, Voevodsky uses sheaf theoretic method in [TriCa]. More precisely, he constructs another category $\mathrm{DM}_{-}^{\mathrm{eff}}(k)$ using a sheaf category and he proves $\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k)$ admits a natural full embedding as a tensor category and a triangulated category to the category $\mathrm{DM}_{-}^{\mathrm{eff}}(k)$. We will review the construction of $\mathrm{DM}_{-}^{\mathrm{eff}}(k)$.

2.5. 1. A presheaf with transfers on Sm/k is an additive contravariant functor from the category SmCor(k) to the category of abelian groups. We denote by PST(k) the category of presheaf with transfers on Sm/k.

2. A presheaf with transfers on Sm /k is called a Nisnevich sheaf with transfers if the corresponding presheaf of abelian groups on Sm/k is a sheaf in the Nisnevich topology. We denoted by $\text{Shv}_{Nis}(\text{SmCor}(k))$ the category of Nisnevich sheaves with transfers.

Example 2.6. For any smooth scheme X over k, a presheaf $\mathbb{Z}_{tr}(X) := Cor(?, X)$ is a Nisnevich sheaf with transfers on Sm /k. (c.f. [TriCa] Lemma 3.1.2). For a k-rational point $x : \operatorname{Spec} k \to X$, we put

$$\mathbb{Z}_{\rm tr}(X,x) := \operatorname{Coker}(\mathbb{Z}_{\rm tr}(\operatorname{Spec} k) \xrightarrow{\mathbb{Z}_{\rm tr}(x)} \mathbb{Z}_{\rm tr}(X)).$$

2.7. 1. A presheaf with transfers F is called homotopy invariant if for any smooth scheme X over k the projection $X \times \mathbb{A}^1_k \to X$ induces the isomorphism $F(X) \to F(X \times \mathbb{A}^1_k)$.

2. A Nisnevich sheaf with transfers is called homotopy invariant if it is homotopy invariant as a presheaf with transfers.

2.8. $\operatorname{Shv}_{\operatorname{Nis}}(\operatorname{SmCor}(k))$ is an abelian category. (c.f. [TriCa] Theorem 3.1.4) Inside the derived category $D^{-}(\operatorname{Shv}_{\operatorname{Nis}}(\operatorname{SmCor}(k)))$ of complexes bounded from above, one defines the full subcategory $\operatorname{DM}_{-}^{\operatorname{eff}}(k)$ of effective motivic complexes over k as the one consisting of objects whose cohomology sheaves are homotopy invariant. This subcategory is triangulated. (Need the assumption of perfectness of k.) (c.f. [TriCa] Proposition 3.1.13).

2.9. 1. Let F be a presheaf with transfers. There is a canonical surjection of presheaves

$$\bigoplus_{(X,x\in F(X))} \mathbb{Z}_{\mathrm{tr}}(X) \to F$$

Iterating this construction we get a canonical left resolution $\mathcal{L}(F)$ of F which consists of direct sums of presheaves of the form $\mathbb{Z}_{tr}(X)$ for smooth schemes X over k.

2. We set for two smooth schemes X, Y:

$$\mathbb{Z}_{\rm tr}(X) \otimes \mathbb{Z}_{\rm tr}(Y) := \mathbb{Z}_{\rm tr}(X \times Y)$$

and for two presheaves with transfers F, G:

$$F \otimes G := \underline{\mathrm{H}}_{0}(\mathcal{L}(F) \otimes \mathcal{L}(G)).$$

3. This construction provides us with a tensor structure on the derived category $D^{-}(\text{Shv}_{Nis}(\text{SmCor}(k)))$.

To define the tensor structure on $DM_{-}^{eff}(k)$, we will need an alternative description of $DM_{-}^{eff}(k)$.

2.10. Let Δ^{\bullet} be the standard cosimplicial object in Sm/k. For any presheaf with transfers F on Sm/k let $C_*(F)$ be the complex of presheaves on Sm/k of the form $C_n(F) = \underline{\text{Hom}}(\Delta^n, F)$ with differentials given by alternated sums of morphisms which correspond to the boundary morphisms of Δ^{\bullet} . This complex is called the singular simplicial complex of F. The following properties are fundamental.

1. If F is a presheaf with transfers (resp. a Nisnevich sheaf with transfers) then $C_*(F)$ is a complex of presheaves with transfers (resp. Nisnevich sheaves with transfers).

2. For any presheaf with transfers F over k, the cohomology presheaf $h_i(F)$ of the complex $C_*(F)$ and its Nisnevich sheafication $h_i^{Nis}(F)$ are homotopy invariant. (Need the assumption of perfectness of k.) (c.f. [TriCa] Lemma 3.2.1).

3. In view of 1. and 2., $C_*(?)$ is a functor from the category of Nisnevich sheaves with transfers on Sm/k to $DM_{-}^{eff}(k)$.

Proposition 2.11. (c.f. [TriCa] Proposition 3.2.3) The functor $C_*(?)$ can be extended to a functor

 $\mathcal{R}C: D^{-}(\operatorname{Shv}_{\operatorname{Nis}}(\operatorname{SmCor}(k)) \to \operatorname{DM}^{\operatorname{eff}}_{-}(k)$

which is left adjoint to the natural embedding. The functor $\mathcal{R}C$ identifies $DM_{-}^{eff}(k)$ with localization of $D^{-}(Shv_{Nis}(SmCor(k)))$ with respect to the localizing subcategory \mathcal{A} generated by complexes of the form

$$\mathbb{Z}_{\rm tr}(X \times \mathbb{A}^1_{\rm k}) \stackrel{\mathbb{Z}_{\rm tr}(pr_1)}{\to} \mathbb{Z}_{\rm tr}(X)$$

for smooth schemes X over k.

2.12. 1. In the notation above, \mathcal{A} is a \otimes -ideal, that is, for any object T of $D^{-}(\operatorname{Shv}_{\operatorname{Nis}}(\operatorname{SmCor}(k)))$ and an object S of \mathcal{A} the object $T \otimes S$ belongs to \mathcal{A} . (c.f. [TriCa] Lemma 3.2.4).

2. We define tensor structure on $\text{DM}_{-}^{\text{eff}}(k)$ as the descent of the tensor structure on $D^{-}(\text{Shv}_{\text{Nis}}(\text{SmCor}(k)))$ with respect to the projector $\mathcal{R}C$. Note that such a descent exists by the universal property of localization and 1.

Theorem 2.13. (c.f. [TriCa] Theorem 3.2.6)

There is a commutative diagram of functors of the form such that the following conditions hold:

$$\mathcal{H}^{b}(\operatorname{SmCor}(k)) \xrightarrow{L} D^{-}(\operatorname{Shv}_{\operatorname{Nis}}(\operatorname{SmCor}(k))) \\ \downarrow \qquad \qquad \downarrow \mathcal{R}C \\ \operatorname{DM}_{\operatorname{gm}}^{\operatorname{eff}}(k) \xrightarrow{i} \operatorname{DM}_{-}^{\operatorname{eff}}(k)$$

1. The functor i is a full embedding with a dense image.

2. For any smooth scheme X over k the object $\mathcal{RC}(L(X))$ is canonically isomorphic to the $C_*(\mathbb{Z}_{tr}(X))$.

3. All functors preserve tensor and triangulated structures.

Example 2.14. Let $x : \operatorname{Spec} k \to X$ be a k-rational point of a smooth scheme X. Then we have an identification

$$C_*(\mathbb{Z}_{\mathrm{tr}}(X, x)) \xrightarrow{\sim} \mathrm{Cone}(\mathrm{M}_{\mathrm{gm}}(\mathrm{Spec}\,k) \xrightarrow{\mathrm{M}_{\mathrm{gm}}(x)} \mathrm{M}_{\mathrm{gm}}(X))$$

defined by

$$\begin{bmatrix} \mathbb{Z}_{\mathrm{tr}}(\operatorname{Spec} k) \\ \mathbb{Z}_{\mathrm{tr}}(x) \\ \mathbb{Z}_{\mathrm{tr}}(X) \end{bmatrix} \xrightarrow{\rightarrow} \begin{bmatrix} 0 \\ \downarrow \\ \mathbb{Z}_{\mathrm{tr}}(X, x) \end{bmatrix}.$$

2.3 Motives with compact support

In this subsection, we will briefly review the notation and fundamental result of [FV00], [RelCy] and [TriCa].

2.15. For any scheme of finite type X over k and any $r \ge 0$ we denote by $\mathcal{Z}_{\text{equi}}(X, r)$ the presheaf on the category of smooth schemes over k which takes a smooth scheme Y to free abelian groups generated by closed integral subschemes Z of $X \times Y$ which are equidimensional of relative dimension r over Y.

This presheaf has the following property.

1. $\mathcal{Z}_{equi}(X, r)$ is a sheaf in the Nisnevich topology.

2. It has a canonical structure of a presheaf with transfers.

3. The presheaf with transfers $\mathcal{Z}_{equi}(X, r)$ is covariantly functorial with respect to proper morphisms of X by means of the usual proper push-forward of cycles. 4. It is contravariantly functorial with an appropriate dimension shift with respect to flat equidimensional morphisms.

5. There is a pairing

$$\times : \mathcal{Z}_{\text{equi}}(X, r) \otimes \mathcal{Z}_{\text{equi}}(X', r') \to \mathcal{Z}_{\text{equi}}(X \times X', r + r')$$

of presheaves.:

Let U be a smooth scheme over k. For any pair of integral closed subschemes $Z \subset X \times U, Z' \subset X' \times U$ equidimensional over U. Sending Z, Z' to the cycle associated to the subscheme $Z \times_U Z' \subset X \times X' \times U$ determines a pairing.

6. (Need the assumption of 0.6.) The flat pull-back morphism induces a quasiisomorphism

$$C_*(\mathcal{Z}_{\text{equi}}(X,0)) \to C_*(\mathcal{Z}_{\text{equi}}(X \times \mathbb{A}^n, n)).$$

2.16. For any scheme of finite type over k the object $C_*(\mathcal{Z}_{\text{equi}}(X,0))$ belongs to $\text{DM}_{\text{-}}^{\text{eff}}(k)$. Moreover it belongs to $\text{DM}_{\text{gm}}^{\text{eff}}(k)$. (Using the assumption 0.6) (c.f. [TriCa], Corollary 4.1.6). We will denote $\text{M}_{\text{gm}}^{\text{c}}(X) := C_*(\mathcal{Z}_{\text{equi}}(X,0))$ and call it a motivic complex of X with compact support.

Since $\mathbb{Z}_{tr}(X)$ is a subsheaf of $\mathcal{Z}_{equi}(X,0)$, the inclusion morphism induces the natural morphism $M_{gm}(X) \to M_{gm}^c(X)$.

The following properties are fundamental. (c.f. [TriCa], Proposition 4.1.5, Proposition 4.1.7)

1. If X is proper then the canonical morphism $M_{gm}(X) \to M_{gm}^{c}(X)$ is the isomorphism.

2. (Need the assumption 0.6.) Let Z be a closed subscheme of X. Then there is a canonical distinguished triangle of the form

$$\mathrm{M}^{\mathrm{c}}_{\mathrm{gm}}(Z) \to \mathrm{M}^{\mathrm{c}}_{\mathrm{gm}}(X) \to \mathrm{M}^{\mathrm{c}}_{\mathrm{gm}}(X-Z) \to \mathrm{M}^{\mathrm{c}}_{\mathrm{gm}}(Z)[1].$$

3. For X, Y of finite type over k, the pairing $\mathcal{Z}_{equi}(X,0) \otimes \mathcal{Z}_{equi}(Y,0) \rightarrow \mathcal{Z}_{equi}(X \times_k Y,0)$ induces an isomorphism

$$\mathrm{M}^{\mathrm{c}}_{\mathrm{gm}}(X) \otimes \mathrm{M}^{\mathrm{c}}_{\mathrm{gm}}(Y) \xrightarrow{\sim} \mathrm{M}^{\mathrm{c}}_{\mathrm{gm}}(X \times_{k} Y)$$

2.4 Tate object

2.17. For any smooth scheme X over k, the morphism $X \to \operatorname{Spec} k$ gives us a morphism in $\operatorname{DM}_{\operatorname{gm}}^{\operatorname{eff}}(k)$ of the form $\operatorname{M}_{\operatorname{gm}}(X) \to \mathbb{Z}$. There is a canonical split distinguished triangle

$$\widetilde{\mathrm{M}_{\mathrm{gm}}}(X) \to \mathrm{M}_{\mathrm{gm}}(X) \to \mathbb{Z} \to \widetilde{\mathrm{M}_{\mathrm{gm}}}(X)[1]$$

where $\widetilde{\mathrm{M}_{\mathrm{gm}}}(X)$ is the reduced motif of X represented in $\mathcal{H}^{b}(\mathrm{SmCor}(k))$ by the complex $[X] \to [\mathrm{Spec} \ k]$.

Example 2.18. In the notation above, for any k-rational point $x : \operatorname{Spec} k \to X$, we have the canonical identification $\operatorname{M_{gm}}(X, x) \xrightarrow{\sim} \widetilde{\operatorname{M_{gm}}}(X)$ as the following way.

$\left[\operatorname{Spec} k\right]$	\longrightarrow	
	id – M (gon)	¥ [X]
	$\stackrel{\mathrm{id}}{\longrightarrow} \stackrel{\mathrm{M}_{\mathrm{gm}}(x \circ p)}{\longrightarrow}$	
	\longrightarrow	$\left[\operatorname{Spec} k\right]$

where $p: X \to \operatorname{Spec} k$ is the structure morphism. $x: \operatorname{Spec} k \to X$ defines splitting $\operatorname{M}_{\operatorname{gm}}(X) \xrightarrow{\sim} \operatorname{M}_{\operatorname{gm}}(X, x) \oplus \mathbb{Z}$.

2.19. We define the Tate object $\mathbb{Z}(1)$ of $\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k)$ as $\widetilde{\mathrm{M}_{\mathrm{gm}}}(\mathbb{P}^1)[-2]$. We further define $\mathbb{Z}(n)$ to be the *n*-th tensor power of $\mathbb{Z}(1)$. For any object A of $\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k)$ we put

$$A(n) = A \otimes \mathbb{Z}(n)$$
$$A\{n\} = A \otimes \mathbb{Z}(n)[n]$$
$$A((n)) = A \otimes \mathbb{Z}(n)[2n]$$

By Example 2.18, for any $x : \operatorname{Spec} k \to \mathbb{P}^1_k$, we have the canonical isomorphism $\operatorname{M}_{\operatorname{gm}}(\mathbb{P}^1_k, x) \xrightarrow{\sim} \mathbb{Z}((1)).$

2.20. Let $x : \operatorname{Spec} k \to \mathbb{P}^1_k$ be a k-rational point. Comparing the following split distinguished triangles

where $M_{gm}(\mathbb{P}^1_k) \to \mathbb{Z}((1))$ is defined by

$$\mathrm{M}_{\mathrm{gm}}(\mathbb{P}^1_k) \xrightarrow[]{}_{\mathrm{can}} \mathrm{M}_{\mathrm{gm}}(\mathbb{P}^1, x) \xrightarrow{\sim} \mathbb{Z}((1)),$$

we know that there is a natural isomorphism $\mathbb{Z}((1)) \xrightarrow{\sim} M^{c}_{gm}(\mathbb{A}^{1})$. It does not depend on the choice of a k-rational point by Example 2.4. Similarly using Mayer-Vietoris sequence for canonical covering of \mathbb{P}^{1}_{k} , we know also that there is a natural isomorphism $\mathbb{Z}\{1\} \xrightarrow{\sim} \widetilde{M}_{gm}(\mathbb{A}^{1} - \{0\})$.

2.5 The triangulated category of geometric motives

In this subsection, we will define the triangulated category $DM_{gm}(k)$ of geometric motives over k.

2.21. 1. We define the category $DM_{gm}(k)$: its objects are pairs of the form (A, n) where A is an object of $DM_{gm}^{\text{eff}}(k)$ and $n \in \mathbb{Z}$ and morphisms are defined by the following formula

$$\operatorname{Hom}_{\operatorname{DM}_{\operatorname{gm}}(k)}((A,n),(B,m)) := \lim_{k \ge -n,-m} \operatorname{Hom}_{\operatorname{DM}_{\operatorname{gm}}^{\operatorname{eff}}(k)}(A(k+n),B(k+m)).$$

2. The category $DM_{gm}(k)$ with the obvious shift functor and class of distinguished triangles is a triangulated category.

3. The permutation involution on $\mathbb{Z}(1) \otimes \mathbb{Z}(1)$ is identity in $\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k)$. (c.f. [TriCa] Corollary 2.1.5)

4. Using the fact of 3. and general theory, $\mathrm{DM}_{\mathrm{gm}}(k)$ has a natural tensor structure.

Theorem 2.22. (c.f. [Voe02] The cancellation theorem) (Need the assumption of perfectness of k.) For objects A, B in $DM_{gm}^{eff}(k)$ the natural map

 $? \otimes \mathrm{id}_{\mathbb{Z}(1)} : \mathrm{Hom}_{\mathrm{DM}_{\mathrm{eff}}^{\mathrm{eff}}(k)}(A, B) \to \mathrm{Hom}_{\mathrm{DM}_{\mathrm{eff}}^{\mathrm{eff}}(k)}(A(1), B(1))$

is an isomorphism. Thus the canonical functor

$$\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k) \to \mathrm{DM}_{\mathrm{gm}}(k)$$

is a full embedding.

3 Various morphisms between motives

In this section, we will briefly review [TriCa] and [MotGe]

3.1 Transpose for finite equidimensional morphisms

3.1. Let X, Y be smooth schemes over k and $f: X \to Y$ a finite equidimensional morphism. Then we have the transpose of $f, {}^tf: Y \to X$ in $\operatorname{SmCor}(k)$. That is $s(\Gamma_f) \in \operatorname{Hom}_{\operatorname{SmCor}(k)}(Y, X)$, where $s: X \times Y \to Y \times X$ is the switch morphism.

Example 3.2. Let L/K/k be finite field extensions. Then we have a canonical morphism i: Spec $L \to$ Spec K. Hence we get $M_{gm}(^ti)$: $M_{gm}(Spec K) \to M_{gm}(Spec L)$ in $DM_{gm}(k)$. We shall write this map as $N_{L/K}$ because there is the following commutative diagram

(c.f. [BKcon] Lemma 3.4.4. See also Theorem 5.16)

3.2 Pull back for flat equidimensional morphisms

3.3. Let X, Y be smooth schemes and $f : X \to Y$ a flat equidimensional morphism of relative dimension n. Then one can define a morphism f^* : $M^{c}_{gm}(Y)((n)) \to M^{c}_{gm}(X)$ as follows. (This is slightly different from the definition in [TriCa] Corollary 4.2.4).

$$\begin{aligned}
\mathbf{M}^{\mathbf{c}}_{\mathrm{gm}}(Y)((n)) &= & C_{*}(\mathcal{Z}_{\mathrm{equi}}(Y \times \mathbb{A}^{n}, 0)) \\
& \stackrel{C_{*}(f \times \mathrm{id}_{\mathbb{A}^{n}})^{*}}{\to} & C_{*}(\mathcal{Z}_{\mathrm{equi}}(X \times \mathbb{A}^{n}, n)) \\
& \stackrel{\leftarrow}{\underset{\mathrm{qis}}{\leftarrow}} & C_{*}(\mathcal{Z}_{\mathrm{equi}}(X, 0)) = \mathbf{M}^{\mathbf{c}}_{\mathrm{gm}}(X)
\end{aligned}$$

where every quasi-isomorphisms are induced from flat pull backs.

3.4. The following properties are easily proved.

1 In the notation above, if f is an open immersion, this morphism coincides with the canonical morphism $\mathrm{M}^{\mathrm{c}}_{\mathrm{gm}}(Y) \to \mathrm{M}^{\mathrm{c}}_{\mathrm{gm}}(X)$.

2 In the notation above, if X and Y are proper over Spec k and f is flat finite equidimensional, then $M_{gm}({}^tf) = f^*$. (c.f. [MotGe] Lemma 1.1.2)

3 Let X, Y and Z be smooth schemes and $f : X \to Y$, $g : Y \to Z$ flat equidimensional morphisms of relative dimension n and m respectively. Then we have $f^* \circ g^*((m)) = (g \circ f)^*$.

Lemma 3.5.

Let $p : \mathbb{A}^1_k \to \operatorname{Spec} k$ be the structure morphism. Then $p^* : \operatorname{M}^c_{\operatorname{gm}}(\operatorname{Spec} k)((1)) = \operatorname{M}^c_{\operatorname{gm}}(\mathbb{A}^1_k) \to \operatorname{M}^c_{\operatorname{gm}}(\mathbb{A}^1_k)$ coincides with $\operatorname{id}_{\operatorname{M}^c_{\operatorname{gm}}(\mathbb{A}^1_k)}$.

Proof. By definition (See 3.3),

$$p^* = C_*(\mathcal{Z}_{\text{equi}}(\mathbb{A}^1_{\mathbf{k}}, 0)) \stackrel{C_*(pr_1^*)}{\to} C_*(\mathcal{Z}_{\text{equi}}(\mathbb{A}^2_{\mathbf{k}}, 1)) \stackrel{(C_*(pr_2^*))^{-1}}{\to} C_*(\mathcal{Z}_{\text{equi}}(\mathbb{A}^1_{\mathbf{k}}, 0))$$

where pr_1, pr_2 are two projections $pr_1, pr_2 : \mathbb{A}^2_k \to \mathbb{A}^1_k$. We assert that two projections pr_1, pr_2 induce the same morphism $pr_1^* = pr_2^* : C_*(\mathcal{Z}_{\text{equi}}(\mathbb{A}^1_k, 0)) \to C_*(\mathcal{Z}_{\text{equi}}(\mathbb{A}^2_k, 1))$ in DM₋^{eff}(k). Since

$$pr_1 = \begin{pmatrix} 0 & 1\\ 1 & 0 \end{pmatrix} \circ pr_2,$$

it suffices to prove that the action of $\operatorname{GL}_2(k)$ on \mathbb{A}^2_k induces trivial action on $C_*(\mathcal{Z}_{\operatorname{equi}}(\mathbb{A}^2, 1))$ in $\operatorname{DM}_{-}^{\operatorname{eff}}(k)$ by flat pull back. On the other hand $\operatorname{GL}_2(k)$ is generated by the elements of conjugate of $\begin{pmatrix} 1 & 0 \\ * & * \end{pmatrix}$. For $A = \begin{pmatrix} 1 & 0 \\ c & d \end{pmatrix} \in \operatorname{GL}_2(k)$, considering the following diagram

$$C_*(\mathcal{Z}_{\text{equi}}(\mathbb{A}^2_k, 1)) \xrightarrow{A^*} C_*(\mathcal{Z}_{\text{equi}}(\mathbb{A}^2_k, 1))$$

$$pr_1^* \downarrow \land \qquad \land \uparrow pr_1^*$$

$$C_*(\mathcal{Z}_{\text{equi}}(\mathbb{A}^1_k, 0)) \xrightarrow{\text{id}} C_*(\mathcal{Z}_{\text{equi}}(\mathbb{A}^1_k, 0)),$$

we get the result.

3.3 Motives with closed support

3.6. We call (X, Z) a closed pair if X is a smooth scheme over k and Z is a closed subscheme. If Z is smooth over k, we call (X, Z) a smooth pair. We call a pair of morphisms of scheme $(f,g): (Y,T) \to (X,Z)$ a morphism of closed pair if a commutative square $T \xrightarrow{} X$ is a Cartesian square as underlying topolog- $\begin{array}{c} g \\ g \\ Z \xrightarrow{} Y \end{array}$

ical spaces. Such a morphisms called Cartesian (resp. excisive) if the diagram above is a Cartesian square (resp. f is étale and g_{red} is an isomorphism.)

3.7. Let X be a smooth scheme and U its open subset. Then we define

$$M_{gm}(X/U) := C^*(\operatorname{Coker}(\mathbb{Z}_{\operatorname{tr}}(U) \to \mathbb{Z}_{\operatorname{tr}}(X))).$$

For any closed pair (X, Z), we define relative motif associated (X, Z) by $M_Z(X) := M_{gm}(X/X - Z)$. By definition there is a canonical distinguished triangle of the form

$$\mathrm{M}_{\mathrm{gm}}(X-Z) \xrightarrow{\mathrm{M}_{\mathrm{gm}}(j)} \mathrm{M}_{\mathrm{gm}}(X) \xrightarrow{i^{\sharp}} \mathrm{M}_{Z}(X) \to \mathrm{M}_{\mathrm{gm}}(X-Z)[1].$$

where $i:Z \hookrightarrow X$ is a closed immersion and $j:X-Z \hookrightarrow X$ is an open immersion.

3.8. For any morphisms of closed pair $(f,g): (Y,T) \to (X,Z)$, we associate a morphism $(f,g)_*: M_T(Y) \to M_Z(X)$ which makes the following diagram commute

where $h: Y - T \to X - Z$ is a induced morphism from f.

3.9. In the notation above, if f is finite equidimensional, we associate a morphism $(f,g)^* : M_T(Y) \to M_Z(X)$ which makes the following diagram commute

$$0 \longrightarrow M_{gm}(Y - T) \longrightarrow M_{gm}(Y) \longrightarrow M_{T}(Y) \longrightarrow 0$$
$$M_{gm}({}^{t}h) \bigwedge M_{gm}({}^{t}f) \bigwedge (f,g)^{*}$$
$$0 \longrightarrow M_{gm}(X - Z) \longrightarrow M_{gm}(X) \longrightarrow M_{Z}(X) \longrightarrow 0.$$

Proposition 3.10. (c.f. [IntMo] Proposition 2.3)

(Red) Reduction: If (X, Z) is a closed pair, the canonical morphism $(X, Z_{red}) \rightarrow (X, Z)$ induces identity map $M_{Z_{red}}(X) \rightarrow M_Z(X)$.

(Add) Additivity: Let X be a smooth scheme, and Z, W disjoint closed subschemes of X. Then induced morphism $M_{Z \coprod W}(X) \to M_Z(X) \oplus M_W(X)$ is an isomorphism.

(Exc) Excision: Any excisive morphism $(Y,T) \to (X,Z)$ induces an isomorphism $M_T(Y) \to M_Z(X)$.

(MV) Mayer-Vietoris: Let X be a smooth scheme over k, U and V two open subsets of X such that $X = U \cup V$, and Z a closed subscheme of X. Then we have a distinguished triangle of the form

$$M_{Z \cap U \cap V}(U \cap V) \to M_{Z \cap U}(U) \oplus M_{Z \cap V}(V) \to M_Z(X) \xrightarrow{+1}$$

(*Htp*) Homotopy invariance: A Cartesian morphism $\pi : (\mathbb{A}^1_X, \mathbb{A}^1_Z) \to (X, Z)$ induced from the canonical projection induces an isomorphism

$$\mathrm{M}_{\mathrm{gm}}(\pi) : \mathrm{M}_{\mathbb{A}^1_Z}(\mathbb{A}^1_X) \to \mathrm{M}_Z(X).$$

3.4 Thom isomorphism

3.11. Let X be a scheme and E/X a vector bundle. We consider X as a closed subscheme of E by zero section. We define the Thom motif of E/X by $M_{gm}(ThE) := M_X(E)$.

3.12. In the notation above, if rank of E is n, there is the Thom isomorphism $\theta(E) : \operatorname{M}_{\operatorname{gm}}(ThE) \xrightarrow{\sim} \operatorname{M}_{\operatorname{gm}}(X)((n))$. We will briefly review the construction of this isomorphism.

3.13. Let X be a smooth scheme and $\Delta_X : X \to X \times_k X$ the diagonal immersion. Let \mathcal{M}, \mathcal{N} be objects of $\mathrm{DM}^{\mathrm{eff}}_{-}(k)$ and

$$\alpha : \mathrm{M}_{\mathrm{gm}}(X) \to \mathcal{M}$$

 $\beta : \mathrm{M}_{\mathrm{gm}}(X) \to \mathcal{N}$

are morphisms in $DM_{-}^{eff}(k)$.

We define external cup product of α and β over X is composition of the following morphisms

$$\mathrm{M}_{\mathrm{gm}}(X) \xrightarrow{\mathrm{M}_{\mathrm{gm}}(\Delta_X)} \mathrm{M}_{\mathrm{gm}}(X) \otimes \mathrm{M}_{\mathrm{gm}}(X) \xrightarrow{\alpha \otimes \beta} \mathcal{M} \otimes \mathcal{N}.$$

We denote this morphism by $\alpha \boxtimes_X \beta$ or $\alpha \boxtimes \beta$

3.14. In the notation above, if $\mathcal{M} = \mathbb{Z}((m))$ and $\mathcal{N} = \mathbb{Z}((n))$, then we have the canonical isomorphism $\varepsilon : \mathbb{Z}((m)) \otimes \mathbb{Z}((n)) \xrightarrow{\sim} \mathbb{Z}((m+n))$.

We put $\alpha \cup \beta := \varepsilon \circ \alpha \boxtimes \beta$ and call it internal cup product of α and β .

3.15. There is a natural isomorphism as a presheaf with transfers (c.f. [MotGe] Corollary 2.2.7)

$$c_1 : \operatorname{Pic}(?) \to \operatorname{Hom}_{\operatorname{DM}_{\operatorname{gm}}(k)}(\operatorname{M}_{\operatorname{gm}}(?), \mathbb{Z}((1))).$$

For any smooth scheme X and $\mathcal{L} \in \operatorname{Pic}(X)$, we will call $c_1(\mathcal{L})$ a motivic Chern class of \mathcal{L} .

3.16. Let X be a smooth scheme, E a vector bundle over X of rank n, λ_E the canonical invertible sheaf on $\mathbb{P}(E)$ and $p: \mathbb{P}(E) \to X$ the canonical projection. We define the r-th motivic Lefschetz projector of E by

$$l_r(E) := c_1(\lambda_E)^{\cup r} \boxtimes \mathrm{M}_{\mathrm{gm}}(p) : \mathrm{M}_{\mathrm{gm}}(\mathbb{P}(E)) \to \mathrm{M}_{\mathrm{gm}}(X)((r))$$

We put the motivic Lefschetz operator as

$$l(E) := \sum_{r=0}^{n-1} l_r(E).$$

Proposition 3.17. (c.f. [TriCa] Proposition 3.5.1) In the notation above, the morphism

$$l(E): \mathrm{M}_{\mathrm{gm}}(\mathbb{P}(E)) \to \bigoplus_{r=0}^{n-1} \mathrm{M}_{\mathrm{gm}}(X)((r))$$

is an isomorphism.

3.18. Let X be a smooth scheme and E/X a vector bundle of rank n. Put $\hat{E} := E \times_X \mathbb{A}^1_X$. Then we have the canonical isomorphisms

$$\mathrm{M}_{\mathrm{gm}}(ThE) \xrightarrow{1} \mathrm{M}_X(\mathbb{P}(\hat{E})) \xrightarrow{2} \mathrm{M}_{\mathrm{gm}}(\mathbb{P}(\hat{E})/\mathbb{P}(E)).$$

The first morphism is induced from an open immersion $E \to \mathbb{P}(\hat{E})$ which is an isomorphism by **(Exc)**. The second morphism is induced from the projection $\mathbb{P}(\hat{E}) - X \to \mathbb{P}(E)$ which is an isomorphism by **(MV)** and **(Htp)**. From this isomorphisms, we get the following distinguished triangle

$$\mathrm{M}_{\mathrm{gm}}(\mathbb{P}(E)) \to \mathrm{M}_{\mathrm{gm}}(\mathbb{P}(\hat{E})) \xrightarrow{\pi_E} \mathrm{M}_{\mathrm{gm}}(ThE) \xrightarrow{+1}$$
.

Using this distinguished triangle and Proposition 3.17, we get the following isomorphism

$$\mathrm{M}_{\mathrm{gm}}(X)((n)) \to \bigoplus_{r=0}^{n} \mathrm{M}_{\mathrm{gm}}(X)((r)) \stackrel{l(\hat{E})^{-1}}{\to} \mathrm{M}_{\mathrm{gm}}(\mathbb{P}(\hat{E})) \stackrel{\pi_{E}}{\to} \mathrm{M}_{\mathrm{gm}}(ThE).$$

We call the inverse of this isomorphism the Thom isomorphism and denote it by $\theta(E)$.

3.5 Normal cone deformation

3.19. Let (X, Z) be a smooth pair of pure codimension c over k such that dimension of X is n. We will write $B_Z X$ by the blow up of X in Z. Put $D_Z X := B_{0 \times Z}(\mathbb{A}^1 \times X) - B_Z X$. There are canonical isomorphisms

$$M_Z(X) \to M_{\mathbb{A}^1_r}(D_Z X) \leftarrow M_{\mathrm{gm}}(ThN_Z X).$$

Hence we get the isomorphism $M_Z(X) \to M_{gm}(ThN_Z X)$.

3.6 Gysin triangles

3.20. Let (X, Z) be a smooth pair of pure codimension c over k and $i : Z \to X$ a closed immersion. In [IntMo], Dèglise constructs the following functorial Gysin triangle in $DM_{gm}(k)$

$$\mathrm{M}_{\mathrm{gm}}(X-Z) \to \mathrm{M}_{\mathrm{gm}}(X) \xrightarrow{i^*} \mathrm{M}_{\mathrm{gm}}(Z)((c)) \xrightarrow{\partial_{X,Z}[1]} \mathrm{M}_{\mathrm{gm}}(X-Z)[1].$$

This is constructed from the following distinguished triangle

$$M_{gm}(X - Z) \to M_{gm}(X) \to M_Z(X) \to M_{gm}(X - Z)[1]$$

and the following isomorphisms

$$M_Z(X) \xrightarrow{\sim} M_{gm}(Th(N_Z(X))) \xrightarrow{\theta(N_Z(X))} M_{gm}(Z)((c))$$

where the first isomorphism is induced from the normal cone deformation.

3.21. Let $(f,g): (Y,T) \to (X,Y)$ be a morphism of closed pairs. We assume Z (resp. T) is connected and smooth over k of codimension n in X (resp. m in Y).

Then we define Gysin morphism associated to (f,g), denote (f,g)! by the following commutative diagram:

$$\begin{array}{cccc}
 & M_{T}(Y) & \xrightarrow{(1)} M_{gm}(Th(N_{T}Y)) \stackrel{\theta(N_{T}Y)}{\sim} M_{gm}(T)((m)) \\
 & (f,g)_{*} & & & \downarrow \\
 & & \downarrow & & \downarrow \\
 & M_{Z}(X) & \xrightarrow{(1)} M_{gm}(Th(N_{Z}X)) \stackrel{\theta(N_{Z}X)}{\sim} M_{gm}(Z)((n))
\end{array}$$

where morphisms (1) are isomorphisms induced from the morphisms of normal cone deformations.

3.22. In the notation above, consider $i: Z \to X, j: X - Z \to X, k: T \to Y$ and $l: Y - T \to T$ the canonical immersions. The following diagram is commutative:

$$\begin{split} \mathbf{M}_{\mathrm{gm}}(Y-T) & \stackrel{\mathbf{M}_{\mathrm{gm}}(l)}{\longrightarrow} \mathbf{M}_{\mathrm{gm}}(Y) \xrightarrow{k^{*}} \mathbf{M}_{\mathrm{gm}}(T)((m)) \xrightarrow{\partial_{Y,T}[1]} \mathbf{M}_{\mathrm{gm}}(Y-T)[1] \\ & \underset{\mathbf{M}_{\mathrm{gm}}(h)}{\longrightarrow} \mathbf{M}_{\mathrm{gm}}(f) & (f,g)_{!} \downarrow & \downarrow \mathbf{M}_{\mathrm{gm}}(h)[1] \\ & \underset{\mathbf{M}_{\mathrm{gm}}(X-Z)}{\longrightarrow} \mathbf{M}_{\mathrm{gm}}(X) \xrightarrow{i^{*}} \mathbf{M}_{\mathrm{gm}}(Z)((n)) \xrightarrow{\partial_{X,Z}[1]} \mathbf{M}_{\mathrm{gm}}(X-Z)[1] \end{split}$$

Lemma 3.23. Let $x : \operatorname{Spec} k \to \mathbb{A}^1_k \xrightarrow{j} \mathbb{P}^1$ be a k-rational point. Then the following diagram is commutative.

$$\begin{split} \mathbf{M}_{\mathrm{gm}}(\mathbb{P}^{1}) & \xrightarrow{x^{*}} \mathbf{M}_{\mathrm{gm}}(\mathrm{Spec}\,k)((1)) \\ & & \downarrow^{\flat} \\ \mathbf{M}_{\mathrm{gm}}^{\mathrm{c}}(\mathbb{P}^{1}) & \xrightarrow{j^{*}} \mathbf{M}_{\mathrm{gm}}^{\mathrm{c}}(\mathbb{A}_{\mathrm{k}}^{1}) \end{split}$$

where the vertical isomorphism $M_{gm}(\operatorname{Spec} k)((1)) \xrightarrow{\sim} M_{gm}^{c}(\mathbb{A}_{k}^{1})$ is defined in 2.20. Proof. It is just a matter of considering two split distinguished triangles below

where the commutativity of 1 follows from Example 2.4.

3.24. In the notation above, if f is finite equidimensional, then we define (f,g)[!] by the following commutative diagram:

$$\begin{array}{c|c} \mathbf{M}_{T}(Y) & \xrightarrow{(1)} \mathbf{M}_{\mathrm{gm}}(Th(N_{T}Y)) \xrightarrow{\theta(N_{T}Y)} \mathbf{M}_{\mathrm{gm}}(T)((m)) \\ & \xrightarrow{(f,g)^{*}} & \downarrow & \downarrow \\ & \downarrow & \downarrow \\ \mathbf{M}_{Z}(X) & \xrightarrow{(1)} \mathbf{M}_{\mathrm{gm}}(Th(N_{Z}X)) \xrightarrow{\theta(N_{Z}X)} \mathbf{M}_{\mathrm{gm}}(Z)((n)) \end{array}$$

where the morphisms (1) are the isomorphisms induced from the morphisms of normal cone deformations.

The following diagram is commutative:

$$\begin{array}{c|c} \mathrm{M}_{\mathrm{gm}}(Y-T) \xrightarrow{\mathrm{M}_{\mathrm{gm}}(l)} & \mathrm{M}_{\mathrm{gm}}(Y) \xrightarrow{k^{*}} \mathrm{M}_{\mathrm{gm}}(T)((m)) \xrightarrow{\partial_{Y,T}[1]} \mathrm{M}_{\mathrm{gm}}(Y-T)[1] \\ & & & & & \\ \mathrm{M}_{\mathrm{gm}}(^{t}h) \bigvee & & & & \\ \mathrm{M}_{\mathrm{gm}}(^{t}h) & & & & & \\ \mathrm{M}_{\mathrm{gm}}(^{t}f) & & & & & \\ \mathrm{M}_{\mathrm{gm}}(X-Z) \xrightarrow{}_{\mathrm{M}_{\mathrm{gm}}(j)} \mathrm{M}_{\mathrm{gm}}(X) \xrightarrow{}_{i^{*}} \mathrm{M}_{\mathrm{gm}}(Z)((n)) \xrightarrow{}_{\partial_{X,Z}[1]} \mathrm{M}_{\mathrm{gm}}(X-Z)[1] \end{array}$$

Proposition 3.25. ([MotGe] Proposition 2.5.2) In the notation above, if (f, g) is Cartesian, then

$$(f,g)^! = \operatorname{M_{gm}}({}^tg)((m)).$$

Next we cite the some proposition in [MotGe]. This is needed to prove that the motivic reciprocity law implies the Weil reciprocity law for Milnor K-groups.

3.26. By 2.20, we have the following distinguished triangle

$$\mathbb{Z} \to \mathrm{M}_{\mathrm{gm}}(\mathbb{G}_{\mathrm{m}}) \xrightarrow{\rho_1} \mathbb{Z}\{1\} \xrightarrow{+1}$$

where $\mathbb{Z} \to M_{gm}(\mathbb{G}_m)$ is induced from the unit morphism Spec $k \xrightarrow{1}{\to} \mathbb{G}_m$.

Proposition 3.27. (c.f. [MotGe] Proposition 2.6.6)

Let (X, Z) be a smooth closed pair of codimension 1. We denote $i: Z \to X$ a closed immersion and $j: X - Z \to X$ a canonical open immersion.

Suppose there is a regular function $\pi : X \to \mathbb{A}^1_k$ which parameterizes Z. Hence we have a morphism $\pi = \pi|_{X-Z} : X - Z \to \mathbb{G}_m$. Then the following diagram is commutative

4 Motivic cohomology groups attached to pointed smooth curves

4.1 Definition

4.1. For pointed smooth curves $(C_1, x_1), \ldots, (C_r, x_r)$ over field k, we define a motivic complex $\mathbb{Z}((C_1, x_1) \land \ldots \land (C_r, x_r))$, or $\mathbb{Z}(C_1 \land \ldots \land C_r)$ in $\mathrm{DM}_{-}^{\mathrm{eff}}(k)$ as follows

$$\mathbb{Z}(C_1 \wedge \ldots \wedge C_r) = C^*(\mathbb{Z}_{tr}(C_1, x_1) \otimes \ldots \otimes \mathbb{Z}_{tr}(C_r, x_r))[-r]$$

4.2. The restriction $\mathbb{Z}(\bigwedge_{i=1}^{r}(C_{i}, x_{i}))|_{X}$ of $\mathbb{Z}(\bigwedge_{i=1}^{r}(C_{i}, x_{i}))$ to the Zariski site of $X \in Sm/k$ is a complex of sheaves in Zariski topology and the motivic cohomology groups $\operatorname{H}^{n}_{\mathcal{M}}(X, \bigwedge_{i=1}^{r}(C_{i}, x_{i}))$, or $\operatorname{H}^{n}_{\mathcal{M}}(X, \bigwedge_{i=1}^{r}C_{r})$ are defined to be the hyper cohomology of the motivic complexes $\mathbb{Z}(\bigwedge_{i=1}^{r}(C_{i}, x_{i}))$ with respect to Zariski topology:

$$\operatorname{H}^{n}_{\mathcal{M}}(X, \bigwedge_{i=1}^{r}(C_{i}, x_{i})) = \operatorname{H}^{n}_{Zar}(X, \operatorname{\mathbb{Z}}(\bigwedge_{i=1}^{r}(C_{i}, x_{i}))|_{X}).$$

4.3. As in [TriCa] we have

$$\mathbb{H}^{n}_{Nis}(X, \mathbb{Z}(\bigwedge_{i=1}^{r} (C_{i}, x_{i}))|_{X}) = \operatorname{Hom}_{\operatorname{DM}^{\operatorname{eff}}_{*}(k)}(\operatorname{M}_{\operatorname{gm}}(X), \mathbb{Z}(\bigwedge_{i=1}^{r} (C_{i}, x_{i}))[n])$$

Moreover if k is a perfect field as in [CohTh] Proposition 3.1.11, we have

$$\mathbb{H}^n_{Zar}(X,\mathbb{Z}(\bigwedge_{i=1}^r(C_i,x_i))|_X) = \mathbb{H}^n_{Nis}(X,\mathbb{Z}(\bigwedge_{i=1}^r(C_i,x_i))|_X).$$

4.2 Fundamental properties

4.4. (Product structure) Let $(C_1, a_1), \ldots, (C_s, a_s), (D_1, b_1), \ldots, (D_t, b_t)$ be pointed smooth curves over field k. As usual motivic complexes, we have canonical morphism

$$\mathbb{Z}(\bigwedge_{i=1}^{s}(C_{i},a_{i}))\otimes\mathbb{Z}(\bigwedge_{j=1}^{t}(D_{j},b_{j}))\to\mathbb{Z}(\bigwedge_{i=1}^{s}(C_{i},a_{i})\wedge\bigwedge_{j=1}^{t}(D_{j},b_{j})).$$

Hence we get for any $X \in Sm/k$ the pairing

$$\mathrm{H}^{p}_{\mathcal{M}}(X, \bigwedge_{i=1}^{s}(C_{i}, a_{i})) \otimes \mathrm{H}^{q}_{\mathcal{M}}(X, \bigwedge_{j=1}^{t}(D_{j}, b_{j})) \to \mathrm{H}^{p+q}_{\mathcal{M}}(X, \bigwedge_{i=1}^{s}(C_{i}, a_{i}) \wedge \bigwedge_{j=1}^{t}(D_{j}, b_{j})).$$

4.5. Let $(C_1, a_1), \ldots, (C_r, a_r)$ be pointed smooth curves over field k. For any field extension L/k, we abbreviate $\operatorname{H}^p_{\mathcal{M}}(\operatorname{Spec} L, \bigwedge_{i=1}^r (C_i \times_k L, a_i \times_k \operatorname{id}_L)$ as $\operatorname{H}^p_{\mathcal{M}}(L, \bigwedge_{i=1}^r C_i)$. By definition, we have

$$\mathrm{H}^{p}_{\mathcal{M}}(L,\bigwedge_{i=1}^{r}C_{i}) = \mathrm{H}_{i-p}(C_{*}(\bigotimes_{i=1}^{r}\mathbb{Z}_{\mathrm{tr}}(C_{i}\times_{k}L,a_{i}\times_{k}\mathrm{id}_{L}))(\mathrm{Spec}\,L)).$$

4.6. (Norm map) In 4.5, if we assume L/k is a finite field extension, then using the description of 4.5 and the proper push-forward of cycles induces a map

$$N_{L/k}: H^p_{\mathcal{M}}(L, \bigwedge_{i=1}^r C_i) \to H^p_{\mathcal{M}}(k, \bigwedge_{i=1}^r C_i).$$

From the corresponding properties of proper push-forward, the following properties are immediately verified.

For finite field extension $k \subset L \subset M$ and $x \in \operatorname{H}^p_{\mathcal{M}}(M, \bigwedge_{i=1}^r C_i)$ and $y \in \operatorname{H}^p_{\mathcal{M}}(L, \bigwedge_{i=1}^r C_i)$

then we have

(1) $\mathcal{N}_{M/L}(y_M \cdot x) = y \cdot \mathcal{N}_{M/L}(x)$ and $\mathcal{N}_{M/L}(x \cdot y_M) = \mathcal{N}_{M/L}(x) \cdot y$

(2) $N_{M/k}(x) = N_{M/L}(N_{L/k}(x))$

(3) If M/k is a normal extension, we have

$$N_{L/k}(x)_M = [L:k]_{insep} \sum_{i:M \hookrightarrow L} j^*(x).$$

Example 4.7. Let (C, x) be a pointed projective smooth curve over k. Then we have

$$\mathrm{H}^{1}_{\mathcal{M}}(k, (C, x)) = \mathrm{Ker}(\mathrm{CH}_{0}(C) \xrightarrow{\mathrm{deg}} \mathbb{Z}).$$

Example 4.8. Let X be a smooth curve over k. A good compactification of X is a pair (\bar{X}, X_{∞}) such that there is an open embedding $X \stackrel{j}{\to} \bar{X}, \bar{X}$ is proper non-singular curve over k and $X_{\infty} = \bar{X} - X$ has an affine open neighborhood in \bar{X} .

Let (C, x) be a pointed smooth affine curve with a good compactification (\bar{X}, X_{∞}) , then

$$\mathrm{H}^{1}_{\mathcal{M}}(k, (C, x)) = \mathrm{Ker}(\mathrm{Pic}(\bar{X}, X_{\infty}) \stackrel{\mathrm{deg}}{\to} \mathbb{Z})$$

where $\operatorname{Pic}(\bar{X}, X_{\infty})$ is the relative Picard group. The elements of $\operatorname{Pic}(\bar{X}, X_{\infty})$ are the isomorphism classes (\mathcal{L}, t) of line bundle \mathcal{L} on \bar{X} with a trivialization t on X_{∞} .

4.9. In 4.7, for finite field extension L/k, using the property 4.6 (3), we know through isomorphisms in 4.7, norm maps in 4.6 and classical one are compatible.

5 Calculation of motivic cohomology groups attached to pointed smooth curves

5.1 Pro-motives

In this subsection, we will briefly review the result of [MotGe].

5.1. Let \mathcal{A} be a tensor triangulated category. We consider Pro- \mathcal{A} the procategory of \mathcal{A} . Then the following facts are fundamental.

1. Pro- \mathcal{A} is additive.

2. The shift functor of \mathcal{A} induces an auto-functor of Pro- \mathcal{A} .

3. There is a unique tensor structure over Pro- \mathcal{A} such that \otimes commutes projective limits.

5.2. In the notation above, we call any triangle in Pro- \mathcal{A} isomorphic to formal projective limit of distinguished triangles of \mathcal{A} a pro-distinguished triangle. Let $H: \mathcal{A}^{op} \to \mathcal{A}b$ be a cohomological functor. Then the functor

$$\overline{H}: (\operatorname{Pro-}\mathcal{A})^{op} \ni (X_i)_{i \in I} \mapsto \inf_{i \in I^{op}} H(X_i) \in \mathcal{A}b$$

sends pro-distinguished triangles to long exact sequences.

5.3. Let \mathcal{O} be a k-algebra. We say \mathcal{O} is local smooth over k iff there is an formally smooth of finite type k-algebra A, and a prime ideal x of A and an isomorphism $\mathcal{O} \xrightarrow{\sim} A_x$.

Since k is perfect, \mathcal{O} is local smooth iff it is regular and essentially of finite type. 5.4. Let \mathcal{O} be a local smooth k-algebra. A model of \mathcal{O}/k is a pair (X, x) consist of a smooth scheme X and a morphism $x : \operatorname{Spec} \mathcal{O} \to X$ such that, if we write the image of closed point of $\operatorname{Spec} \mathcal{O}$ to x, induced morphism $x^{\sharp} : \mathcal{O}_{X,x} \to \mathcal{O}$ is an isomorphism.

We put

$$\mathcal{M}^{\mathrm{smo}}(\mathcal{O}/k) := \{ A \subset \mathcal{O}; (\operatorname{Spec} A, \operatorname{Spec} \mathcal{O} \xrightarrow{\operatorname{obvious}} \operatorname{Spec} A) \text{ is a model of } \mathcal{O}/k \}.$$

 $\mathcal{M}^{\mathrm{smo}}(\mathcal{O}/k)$ is not empty and filtrant for inclusion. (c.f. [MotGe] Lemma 3.1.5). 5.5. 1. Let \mathcal{O} be a local smooth k-algebra. We consider a pro-object of Sm /k

$$(\mathcal{O}) := \{\operatorname{Spec} A\}_{A \in \mathcal{M}^{\operatorname{smo}}(\mathcal{O}/k)}.$$

2. Let X be a smooth scheme and $x \in X$, we define localization of X in x as a pro-object of Sm /k.

$$X_x := \{U\}_{x \in U \subset X}$$

where U runs through the open neighborhood of x.

5.6. (c.f. [MotGe] Lemma 3.1.8) Let \mathcal{O} be a local smooth k-algebra, and (X, x) a model of \mathcal{O} . Then x induces a canonical isomorphism

$$(\mathcal{O}) \to X_x$$

5.7. Let \mathcal{O} be a local smooth k-algebra and $n, m \in \mathbb{Z}$. We consider a pro-object of $\operatorname{Pro-DM}_{\operatorname{gm}}(k)$

$$\mathcal{M}_{\mathrm{gm}}(\operatorname{Spec}\mathcal{O})(n)[m] := \{\mathcal{M}_{\mathrm{gm}}(\operatorname{Spec}A)(n)[m]\}_{A \in \mathcal{M}^{\mathrm{smo}}(\mathcal{O}/k)}.$$

Next we define a residue morphism associated to a discrete valuation.

5.8. Let E/k be a field extension of finite type, v a valuation of E/k, \mathcal{O}_v a valuation ring of v and (X,t) a k-model of \mathcal{O}_v . We say a special point of (X,t) for image of closed point of Spec \mathcal{O}_v for t and denote by s.

We say that (X, t) is a strict k-model of \mathcal{O}_v iff closure $\{s\}$ in X is a smooth scheme.

Any discrete valuation ring \mathcal{O}_v essentially of finite type over k admits a strict k-model. (c.f. [MotGe] Lemma 4.5.3).

5.9. Let E/k be a field extension of finite type, v a valuation of E/k, \mathcal{O}_v a valuation ring of v and (X, t) a strict k-model of \mathcal{O}_v . Put $Z := \{\bar{s}\}$. Since (X, Z) is a smooth closed pair of codimension 1. We have a distinguished triangle of the form

$$\mathrm{M}_{\mathrm{gm}}(Z)\{1\} \xrightarrow{\partial_{X,Z}} \mathrm{M}_{\mathrm{gm}}(X-Z) \xrightarrow{j_*} \mathrm{M}_{\mathrm{gm}}(X) \xrightarrow{+1} .$$

Since this triangle is natural for inclusions of open sets in X. (c.f. 3.22) Considering a cofiltrant system of open neighborhoods of s in X, we get a prodistinguished triangle

$$\mathrm{M}_{\mathrm{gm}}(Z_s)\{1\} \stackrel{\partial_{X_s,Z_s}}{\to} \mathrm{M}_{\mathrm{gm}}(X_s - Z_s) \stackrel{j_{s*}}{\to} \mathrm{M}_{\mathrm{gm}}(X_s) \stackrel{+1}{\to} .$$

Since (X, s) is a k-model of \mathcal{O}_v , a morphism s: Spec $\mathcal{O}_v \to X$ induces an isomorphism of pro-object $(\mathcal{O}_v) \to X_s$; Hence we get a pro-distinguished triangle isomorphic to the form

$$\operatorname{M}_{\operatorname{gm}}(\operatorname{Spec} k(v))\{1\} \xrightarrow{\partial_{X_s, Z_s}} \operatorname{M}_{\operatorname{gm}}(\operatorname{Spec} E) \xrightarrow{i^{\sharp}} \operatorname{M}_{\operatorname{gm}}(\operatorname{Spec} \mathcal{O}_v) \xrightarrow{+1}$$

where E (resp. k(v)) is a fraction field (resp. residue field) of $v, i : \mathcal{O}_v \to E$ is a canonical inclusion.

Lemma 5.10. (c.f. [MotGe] Lemma 4.5.5.)

Let E/k be a field extension of finite type, v a discrete valuation of E/k, \mathcal{O}_v a valuation ring of v and k(v) is this residue field.

Then adopting the notation above, if (X, s) and (Y, t) are two strict k-models of \mathcal{O}_v , put $Z := \{\bar{s}\}$ closure in X and $T := \{\bar{t}\}$ closure in Y. Then we have

$$\partial_{X_s,Z_s} = \partial_{Y_t,T_t}$$

5.11. Let E/k be a field extension of finite type, v a discrete valuation of E/k. We define a residue morphism associated to v, denoted by ∂_v defined by $\partial_v := \partial_{(X_s,Z_s)}$ where (X,s) is a strict k-model of valuation ring of v. By Lemma 5.10 this does not depend on a choice of a strict k-model of valuation ring of v. So we have the following pro-distinguished triangle of the form

$$\operatorname{M}_{\operatorname{gm}}(\operatorname{Spec} k(v))\{1\} \xrightarrow{\partial_v} \operatorname{M}_{\operatorname{gm}}(\operatorname{Spec} E) \xrightarrow{i^{\sharp}} \operatorname{M}_{\operatorname{gm}}(\operatorname{Spec} \mathcal{O}_v) \xrightarrow{+1}$$

Having defined residue morphisms, we explain the connection of Milnor K-groups and Hom sets in $\text{Pro-DM}_{\text{gm}}(k)$.

5.12. Using the distinguished triangle in 3.26 and the definition of tensor structure in $\text{DM}_{-}^{\text{eff}}(k)$, for any $n \in \mathbb{N}$, there is a distinguished triangle of the form

$$\bigoplus_{i=1}^{n} \mathrm{M}_{\mathrm{gm}}(\mathbb{G}_{\mathrm{m}}^{n-1}) \to \mathrm{M}_{\mathrm{gm}}(\mathbb{G}_{\mathrm{m}}^{n}) \xrightarrow{\rho_{n}} \mathbb{Z}\{n\} \xrightarrow{+1}.$$

where the first morphism is induced from sum of n closed immersions

$$\iota_i := \mathrm{id} \times \mathrm{id} \times \ldots \times \overset{\imath}{1} \times \ldots \times \mathrm{id} \times \mathrm{id} \,.$$

5.13. Let E/k be a field extension of finite type. $f : M_{gm}(\operatorname{Spec} E) \to \mathcal{M}$ and $g : M_{gm}(\operatorname{Spec} E) \to \mathcal{N}$ are morphisms in Pro-DM_{gm}(k). Then we can extend the definition of external cup product $f \boxtimes g : M_{gm}(\operatorname{Spec} E) \to \mathcal{M} \otimes \mathcal{N}$.

If $\mathcal{M} = \mathbb{Z}\{p\}$ and $\mathcal{N} = \mathbb{Z}\{q\}$, we have a canonical isomorphism $\mathbb{Z}\{p\} \otimes \mathbb{Z}\{q\} \xrightarrow{\sim} \mathbb{Z}\{p+q\}$. Then we have the following identity

$$\alpha \boxtimes \beta = -\beta \boxtimes \alpha.$$

(c.f. [MotGe] Remarque 4.4.2.)

5.14. Let E/k be a field extension of finite type. Then we have a morphism

$$(E^{\times})^{n} \xrightarrow{\sim} \operatorname{Hom}(\operatorname{Spec} E, \mathbb{G}_{m}^{n})$$

$$\xrightarrow{\operatorname{Mgm}} \operatorname{Hom}_{\operatorname{Pro-DM}_{gm}(k)}(\operatorname{Mgm}(\operatorname{Spec} E), \operatorname{Mgm}(\mathbb{G}_{m}^{n}))$$

$$\xrightarrow{\operatorname{Hom}(\operatorname{Mgm}(\operatorname{Spec} E), \rho_{n})} \operatorname{Hom}_{\operatorname{Pro-DM}_{gm}(k)}(\operatorname{Mgm}(\operatorname{Spec} E), \mathbb{Z}\{n\}).$$

This map induces a morphism

$$\alpha: K_n^M(E) \to \operatorname{Hom}_{\operatorname{Pro-DM}_{\operatorname{gm}}(k)}(\operatorname{M}_{\operatorname{gm}}(\operatorname{Spec} E), \mathbb{Z}\{n\}).$$

5.15. In the notation above, for any $x \in (E^{\times})^{\otimes n}$ and $y \in (E^{\times})^{\otimes m}$, we have

$$\alpha(x \otimes y) = \alpha(x) \boxtimes \alpha(y).$$

Theorem 5.16. (c.f. [MotGe] Theorem 4.4.4) (Need the assumption of perfectness of k.) In the notation above,

 $\alpha: K^M_*(E) \to \operatorname{Hom}_{\operatorname{Pro-DM}_{\operatorname{gm}}(k)}(\operatorname{M}_{\operatorname{gm}}(\operatorname{Spec} E), \mathbb{Z}\{*\})$

is an algebra isomorphism.

5.2 Motivic reciprocity law

The classical theorems "Weil reciprocity law" and "residue formula" are unified using Milnor K-groups. More precisely, the following statement is known. (c.f. [Sus82])

5.17. (Reciprocity law for Milnor K-groups)

Let K be an algebraic function field over a field k. Then the following composition are the zero maps for all non-negative integers n.

$$K_{n+1}^M(K) \stackrel{\bigoplus \partial_v}{\to} \bigoplus_v K_n^M(k(v)) \stackrel{\sum N_{k(v)/k}}{\to} K_n^M(k)$$

In this subsection, we will prove more fundamental style of the following reciprocity law.

Theorem 5.18. (Motivic reciprocity law) The following composition

$$\mathrm{M}_{\mathrm{gm}}(\mathrm{Spec}\,k)\{1\} \xrightarrow{\Sigma \operatorname{N}_{k(v)/k}\{1\}} \prod_{v}^{\sim} \mathrm{M}_{\mathrm{gm}}(\mathrm{Spec}\,k(v))\{1\} \xrightarrow{\widetilde{\Pi}\partial_{v}} \mathrm{M}_{\mathrm{gm}}(\mathrm{Spec}\,K)$$

is the zero map in $\operatorname{Pro-DM}_{gm}(k)$.

5.19. Let K/k be a field extension of transcendental degree one. Let C/k be a projective nonsingular curve such that K(C) = K. As in the previous subsection, we can construct the following pro-distinguished triangle in Pro-DM_{gm}(k).

$$\begin{split} \mathrm{M}_{\mathrm{gm}}(\operatorname{Spec} K) &\to \mathrm{M}_{\mathrm{gm}}(C) \\ &\to \prod_{x \in C: \mathrm{closed \ points}}^{\sim} \mathrm{M}_{\mathrm{gm}}(\operatorname{Spec} k(x))((1)) \stackrel{\widetilde{\Pi} \partial_x[1]}{\to} \mathrm{M}_{\mathrm{gm}}(\operatorname{Spec} K)[1] \end{split}$$

This is constructed as follows: For any closed set $Z \subset C$, there is the Gysin triangle

$$M_{gm}(C-Z) \to M_{gm}(C) \to \bigoplus_{x \in Z} M_{gm}(\operatorname{Spec} k(x))((1)) \xrightarrow{\partial_{C,Z}[1]} M_{gm}(C-Z)[1]$$

and we consider $\mathcal{M}_{gm}(\operatorname{Spec} K) = {\mathcal{M}_{gm}(C-Z)}_{Z \subset C: \text{closed subsets}} \in \operatorname{Pro-DM}_{gm}(k)$

Lemma 5.20.

In the notation above, for any closed point $x \in C$, the diagram of structure morphisms



induces the following commutative diagram:

$$\operatorname{M_{gm}(\operatorname{Spec} k(x))((1))}_{N_{k(x)/k}((1))} \underbrace{\operatorname{M_{gm}(C)}}_{p^*} p^*$$

Proof. First choose a finite equidimensional morphism $C \xrightarrow{\pi} \mathbb{P}^1$ which is unramified at every points over $\pi(x)$. (This can be done by using Bertini theorem.) Next using 3.4 and Proposition 3.25, we may assume $C = \mathbb{P}^1$. Replacing \mathbb{P}^1 by $\mathbb{P}^1_{k(x)}$ and using Proposition 3.25 again, we may assume k(x) = k. In this case, $i^* \circ p^* = \text{id}$ by Lemma 3.5 and 3.23.

5.21. Hence we get the following diagram:

$$M_{gm}(\operatorname{Spec} k)((1))$$

$$\downarrow^{p^{*}} \qquad \qquad \downarrow^{\Sigma N_{k(x)/k}((1))}$$

$$M_{gm}(C) \xrightarrow{p^{*}} \bigoplus_{x \in Z} M_{gm}(\operatorname{Spec} k(x))((1)) \longrightarrow M_{gm}(C - Z)[1]$$

Taking a limit with respect to Z, we get the motivic reciprocity law.

Next we prove that the motivic reciprocity law implies the Weil reciprocity law for Milnor K-groups.

Lemma 5.22.

In the notation above, let v be a valuation of K/k, \mathcal{O}_v a valuation ring of v, π a uniformizer element of \mathcal{O}_v . Then the following diagram is commutative.

Proof. Take a strict k-model of Spec \mathcal{O}_v . Denote it by (X, s). If we take X sufficiently small, $\pi \in \mathcal{O}_v \xrightarrow{\sim} \mathcal{O}_{X,s}$ determines a regular function $X \to \mathbb{A}^1_k$ which parameterizes Z. Using Proposition 3.27 and considering a cofiltrant system of open neighborhoods of s in X, we get the following commutative diagram of pro-motives

Hence we get the result.

Example 5.23. In the notation above, for any discrete valuation v of K/k, there is a commutative diagram

Proof. For $u_1, \ldots, u_{n+1} \in \mathcal{O}_v^{\times}$ and a uniformizer π , it is enough to check the following two conditions.

1 Hom $(\partial_v, \mathbb{Z}\{n+1\})(\alpha(\{u_1, \dots, u_{n+1}\})) = 0$

 $2 \operatorname{Hom}(\partial_v, \mathbb{Z}\{n+1\})(\alpha(\{u_1, \dots, u_n, \pi\})) = (-1)^n \alpha(\{\overline{u_1}, \dots, \overline{u_n}\})$

To prove 1: Notice that there is a pro-distinguished triangle as follows (c.f. 5.11)

$$\mathrm{M}_{\mathrm{gm}}(\mathrm{Spec}\,k(v))\{1\} \xrightarrow{\mathcal{O}_v} \mathrm{M}_{\mathrm{gm}}(\mathrm{Spec}\,K) \to \mathrm{M}_{\mathrm{gm}}(\mathrm{Spec}\,\mathcal{O}_v) \xrightarrow{+1}$$

To prove 2: Anti-commutativity of \boxtimes (c.f. 5.13) and Lemma 5.22.

Example 5.24. Let K/k be a field extension of transcendental degree 1, (C, x) be a pointed smooth curve and v a place of K/k. There is a tame symbol ∂_v : Jac $C(K_v) \otimes K_v^{\times} \to \text{Jac } C(k(v))$. Then the following diagram is commutative.

This is proved in the same way as Example 5.23.

Corollary 5.25.

The motivic reciprocity law implies the Weil reciprocity law for Milnor K-groups.

Proof. Take Hom(?, $\mathbb{Z}\{n+1\}$) and use Theorem 2.22, Theorem 5.16 and notice Example 3.2 and Example 5.23.

5.3 Main result

5.26. In this section, let k be a perfect field which admits resolution of singularities and $(C_1, a_1), \ldots, (C_n, a_n)$ pointed projective smooth curves over k.

5.27. Let $p: Z \to \mathbb{A}^1_k$ be a finite surjective morphism and suppose that Z is integral. Let $f_i \in \text{Hom}(Z, C_i)$ and

$$p^{-1}(j) = \prod n_i^j z_i^j \quad (j = 0, 1)$$

where n_i^j are the multiplicities of points $z_i^j = \operatorname{Spec} L_i^j$. Define:

$$\phi_j = \sum n_i^j \{f_1, \dots, f_n\}_{L_i^j/k}$$

then we have

 $\phi_0 = \phi_1$

in $K(k, \operatorname{Jac} C_1, \ldots, \operatorname{Jac} C_n)$.

The proof is similar to [MVW02], p.45 Corollary 5.5.

5.28. As $\bigotimes_{i=1}^{n} \mathbb{Z}_{tr}(C_i, a_i)(\operatorname{Spec} k))$ is a quotient of the free abelian groups generated by the closed points of $C_1 \times \ldots \times C_n$ modulo the subgroup generated by all points of the form $(x_1, \ldots, a_i, \ldots, x_n)$ where the a_i 's can be any position. If x is a closed point of $C_1 \times \ldots \times C_n$ with residue field L then x is defined by a

canonical sequence $(x_1, \ldots, x_n) \in \operatorname{Jac} C_1(L) \times \ldots \times \operatorname{Jac} C_n(L)$. Since

$$\mathrm{H}^{n}_{\mathcal{M}}(k,\bigwedge_{i=1}^{n}C_{i}) = \mathrm{Coker}(\bigotimes_{i=1}^{n}\mathbb{Z}_{\mathrm{tr}}(C_{i},a_{i})(\mathbb{A}^{1}_{\mathrm{k}}) \xrightarrow{\partial_{0}-\partial_{1}} \bigotimes_{i=1}^{n}\mathbb{Z}_{\mathrm{tr}}(C_{i},a_{i})(\mathrm{Spec}\,k))$$

using 5.27, we have a natural map $\operatorname{H}^{n}_{\mathcal{M}}(k, \bigwedge_{i=1}^{n} C_{i}) \to K(k, \operatorname{Jac} C_{1}, \ldots, \operatorname{Jac} C_{n}).$ 5.29. Using 4.7, for every finite field extension L/k we have an isomorphism

$$\bigotimes_{i=1}^{n} \operatorname{Jac} C_{i}(L) \xrightarrow{\sim} \bigotimes_{i=1}^{n} \operatorname{H}^{1}_{\mathcal{M}}(L, C_{i}).$$

Combining a natural pairing in 4.4

$$\bigotimes_{i=1}^{n} \operatorname{H}^{1}_{\mathcal{M}}(L, C_{i}) \to \operatorname{H}^{n}_{\mathcal{M}}(L, \bigwedge_{i=1}^{n} C_{i})$$

and a norm map (c.f 4.6)

$$N_{L/k}: H^n_{\mathcal{M}}(L, \bigwedge_{i=1}^n C_i) \to H^n_{\mathcal{M}}(k, \bigwedge_{i=1}^n C_i)$$

we get a canonical map

$$\bigoplus_{L/k:\text{finite extension}} \bigotimes_{i=1}^{n} \operatorname{Jac} C_{i}(L) \to \operatorname{H}^{n}_{\mathcal{M}}(k, \bigwedge_{i=1}^{n} C_{i}).$$

If we use 4.6 (1), Theorem 4.9, Theorem 5.18 and Example 5.24, this map should factor through the map $K(k, \operatorname{Jac} C_1, \ldots, \operatorname{Jac} C_n) \to \operatorname{H}^n_{\mathcal{M}}(k, \bigwedge_{i=1}^n C_i).$

5.30. Obviously the morphisms above are inverse to each other. Hence we get the following result.

Theorem 5.31. (Somekawa conjecture for Jacobian varieties) Let $(C_1, a_1), \ldots, (C_n, a_n)$ be pointed projective smooth curves over perfect field k which admits resolution of singularities. Then

 $K(k, \operatorname{Jac} C_1, \ldots, \operatorname{Jac} C_n) \xrightarrow{\sim} \operatorname{Hom}_{\operatorname{DM}^{\operatorname{eff}}_{*}(k)}(\operatorname{M}(\operatorname{Spec} k), \mathbb{Z}(\bigwedge_{i=1}^{n} C_i)[n]).$

Acknowledgment The author is greatful for Professors Takeshi Saito, Shuji Saito, Kazuya Kato, Takao Yamazaki and Kenichiro Kimura.

References

[Akh00] Reza Akhtar, Milnor K-theory and zero-cycles on algebraic varieties, thesis.

- [Akh02] Reza Akhtar, Milnor K-theory of smooth quasi projective varieties, preprint.
- [BT73] H. Bass and J. Tate, The Milnor ring of global field, Springer Lecture Notes in Math. 342 (1973), p. 349-446.
- [Blo76] S. Bloch, Some elementary theorems about algebraic cycles on abelian varieties, Inventions math. 37, (1976), p. 215-228.
- [Blo81] S. Bloch, Algebraic K-theory and class field theory for arithmetic surface, Ann. of Math. 114, (1981), p. 229-266.
- [IntMo] F. Déglise, Interprétation motivique de la formule d'excès d'intersection, C. R. Acad. Sci. Paris, Ser. I 338 (2004) p. 41-46.
- [MotGe] F. Déglise, Motifs génériques, preprint.
- [Del74] P. Deligne, Théorie de Hodge III, Publ. Math. I.H.E.S. 44 (1974), p.5-78.
- [FV00] Eric M. Friedlander and V. Voevodsky, Bivariant cycle cohomology, in Cycles, transfers, and motivic homology theories, Annals of Mathematics Studies, vol 143, Princeton University press, (2000), p. 138-187.
- [Gil81] H. Gillet, Riemann-Roch theorems for higher algebraic K-theory, Adv. in Math. 40 (1981), no. 3, p. 203-289.
- [Kah92] B. Kahn, Nullité de certains groupes attachés aux variétés semiabéliennes sur corps fini, C. R. Acad. Sci. Paris Sér. I Math. 314 (1992), no. 13, p. 1039-1042.
- [KS83] K. Kato and S. Saito, Unramified class field theory of arithmetic surfaces, Ann. of Math. 118 (1983), p. 241-275.
- [KL81] N. Katz and S. Lang, Finiteness theorems in geometric class field theory, l'Enseignement Mathématique 27 (1981) p. 185-314.
- [Lan59] S. Lang, Abelian variety, New York: Interscience-Wiley (1959)
- [MVW02] C. Mazza, V. Voevodsky, C. Weibel Notes on Motivic Cohomology, preprint.
- [Mil70] J. Milnor, Algebraic K theory and quadratic forms, Inventions Math. (1970), p. 318-344.
- [Moo69] C. Moore, Group extension of p-adic and adelic linear groups, Publ. Math. I.H.E.S. 35 (1969), p. 251-281.
- [Som90] M. Somekawa, On Milnor K-groups attached at semi-Abelian varieties, K-theory, 4 (1990), p. 105-119.

- [Sus82] A. Suslin, Menicke symbols and their applications in the K-theory of fields, Proceedings of a Conference held at Oberwolfach, June 16-20, 1980, Springer-Verlag, Berlin, 1982, p. 334-356.
- [RelCy] A. Suslin and V. Voevodsky, *Relative cycles and Chow sheaves*, in *Cycles, transfers, and motivic homology theories*, Annals of Mathematics Studies, vol 143, Princeton University press, (2000), p. 10-86.
- [BKcon] A. Suslin and V. Voevodsky, Bloch-Kato conjecture and motivic cohomology with finite coefficients, The Arithmetic and Geometry of Algebraic Cycles, Nato ASI series C, vol. 548, Kluwer, (2000), p. 117-189.
- [CohTh] V. Voevodsky, Cohomological theory of presheaves with transfers, in Cycles, transfers, and motivic homology theories, Annals of Mathematics Studies, vol 143, Princeton University press, (2000), p. 87-137.
- [TriCa] V. Voevodsky, Triangulated categories of motives over field, in Cycles, transfers, and motivic homology theories, Annals of Mathematics Studies, vol 143, Princeton University press, (2000), p. 188-254.
- [Voe02] V. Voevodsky, Cancellation theorem, preprint.
- [Wei67] A. Weil, Basic Number Theory, Springer-Verlag (1967)

Satoshi Mochizuki Graduate school of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro-ku Tokyo 153-8914, JAPAN E-mail:mochi@ms.u-tokyo.ac.jp