### THE GAUSS MAP OF PSEUDO-ALGEBRAIC MINIMAL SURFACES IN $\mathbb{R}^4$

YU KAWAKAMI

ABSTRACT. In this paper, we prove effective estimates for the number of exceptional values and the totally ramified value number for the Gauss map of pseudo-algebraic minimal surfaces in Euclidean four-space and give a kind of unicity theorem.

### 1. INTRODUCTION

The Gauss map of complete regular minimal surfaces has many properties which have similarities to results in value distribution theory of meromorphic functions on the complex plane **C**. Actually, by using the techniques of value distribution theory, Fujimoto [3] proved that the Gauss map of nonflat complete regular minimal surfaces immersed in Euclidean 3-space  $\mathbb{R}^3$ can omit at most four values. He also obtained the result on the number of exceptional values of the Gauss map of complete minimal surfaces in Euclidean 4-space  $\mathbb{R}^4$ .

On the other hand, Osserman [16] proved that the Gauss map of nonflat algebraic minimal surfaces in  $\mathbb{R}^3$  can omit at most three values. By an algebraic minimal surface, we mean a complete minimal surface with finite total curvature. And Hoffman and Osserman [9] extended the result to algebraic minimal surfaces in  $\mathbb{R}^4$ . However, for the results of  $\mathbb{R}^4$ , we did not know the relationship between Fujimoto's result and Hoffman-Osserman's.

Recently, the author [12] found non-flat algebraic minimal surfaces with totally ramified value number  $\nu_g = 2.5$ . Moreover, the author, Kobayashi and Miyaoka defined the class of pseudo-algebraic minimal surfaces in  $\mathbf{R}^3$  which contains algebraic minimal surfaces as a proper subclass and obtained estimates for the number of exceptional values and the totally ramified value

<sup>2000</sup> Mathematics Subject Classification. Primary 53A10; Secondary 30D35, 30F10.

Key words and phrases. minimal surface, Gauss map, totally ramified value number. The author is partially supported by OCAMI (Osaka City University Advanced Math-

number of the Gauss map for this class (see [13]). These are the best possible estimates for pseudo-algebraic minimal surfaces and some special cases of algebraic minimal surfaces. Therefore, for  $\mathbf{R}^3$ , we can understand the relationship between Fujimoto's result and Osserman's.

In this paper, we first define pseudo-algebraic minimal surfaces in  $\mathbb{R}^4$ . Next, by using some results of complex algebraic geometry, we give estimates for the number of exceptional values and the totally ramified value number of the Gauss map of pseudo-algebraic minimal surfaces and examples which show that these estimates are the best possible. By these estimates, for  $\mathbb{R}^4$ , we can reveal the relationship between Fujimoto's result and Hoffman-Osserman's. Finally we give a kind of unicity theorem and show that the theorem is the best possible.

### 2. Preliminaries

Let  $x = (x^1, x^2, x^3, x^4) \colon M \to \mathbf{R}^4$  be a (connected oriented) regular minimal surface in  $\mathbf{R}^4$ . There canonically exists a system of positive isothermal local coordinate system (u, v) defining the system of holomorphic local coordinates  $z = u + \sqrt{-1}v$  of M. We may thus regard M as a Riemann surface. It is well-known that the set of all oriented 2-planes in  $\mathbf{R}^4$  is canonically identified with the quadric

$$\mathbf{Q}^{2}(\mathbf{C}) = \{(w^{1}: w^{2}: w^{3}: w^{4}) | (w^{1})^{2} + \dots + (w^{4})^{2} = 0\}$$

in  $\mathbf{P}^3(\mathbf{C})$ . By definition, the Gauss map  $g: M \to \mathbf{Q}^2(\mathbf{C})$  is the map which maps each point z of M to the point of  $\mathbf{Q}^2(\mathbf{C})$  corresponding to the oriented tangent plane of M at z. By the assumption of minimality of M, g is a holomorphic map. On the other hand, the quadric  $\mathbf{Q}^2(\mathbf{C})$  is biholomorphic to  $\widehat{\mathbf{C}} \times \widehat{\mathbf{C}}$ , where we denote by  $\widehat{\mathbf{C}}$  the Riemann sphere. Thus we may regard g as a pair of meromorphic functions  $g = (g_1, g_2)$  on M. For  $i = 1, \ldots, 4$ , set  $\phi_i = \partial x = (\partial x/\partial z) dz$  where  $\partial/\partial z = \partial/\partial u - \sqrt{-1}\partial/\partial v$  following Osserman [17]. Then these satisfy

(C)  $\sum \phi_i^2 = 0$ : conformality condition (R)  $\sum |\phi_i|^2 > 0$ : regularity condition (P) For any cycle  $\gamma \in H_1(M, \mathbb{Z}), \Re \int_{\gamma} \phi_i = 0$ : period condition We recover x by the real Abel-Jacobi map (called the Weierstrass-Enneper representation formula)

$$x(z) = \Re \int_{z_0}^{z} (\phi_1, \phi_2, \phi_3, \phi_4)$$

up to translation. If we put

$$hdz = \phi_1 - \sqrt{-1}\phi_2, \quad g_1 = \frac{\phi_3 + \sqrt{-1}\phi_4}{\phi_1 - \sqrt{-1}\phi_2}, \quad g_2 = \frac{-\phi_3 + \sqrt{-1}\phi_4}{\phi_1 - \sqrt{-1}\phi_2}$$

then hdz is a holomorphic differential and both  $g_1$  and  $g_2$  are meromorphic functions on M. Geometrically,  $g = (g_1, g_2)$  is the Gauss map of M. We call  $(hdz, g_1, g_2)$  the Weierstrass data. This is related to  $\phi_j$ 's in one to one way

$$\begin{cases} \phi_1 = \frac{1}{2}(1+g_1g_2)hdz ,\\ \phi_2 = \frac{\sqrt{-1}}{2}(1-g_1g_2)hdz ,\\ \phi_3 = \frac{1}{2}(g_1-g_2)hdz ,\\ \phi_4 = -\frac{\sqrt{-1}}{2}(g_1+g_2)hdz \end{cases}$$

If we are given a holomorphic differential hdz and two meromorphic functions  $g_1$  and  $g_2$  on M, we get  $\phi_j$ 's by this formula. They satisfy (C) automatically, and the condition (R) is interpreted as the zeros of hdz of order k coincide exactly with the poles of  $g_1$  or  $g_2$  of order k, because the induced metric on M is given by

$$ds^{2} = \frac{1}{4} |h|^{2} (1 + |g_{1}|^{2})(1 + |g_{2}|^{2})|dz|^{2} .$$

A minimal surface is complete if all divergent paths have infinite length with respect to this metric. In general, for given two meromorphic functions  $g_1$ and  $g_2$  on M, it is not so hard to find a holomorphic differential hdz satisfying (R). However the period condition (P) always causes trouble. When (P) is not satisfied, we anyway obtain a minimal surface on the universal covering surface of M.

Now the Gauss curvature K of M is given by

$$K = -\frac{2}{|h|^2 (1+|g_1|^2)(1+|g_2|^2)} \left(\frac{|g_1'|^2}{(1+|g_1|^2)^2} + \frac{|g_2'|^2}{(1+|g_2|^2)^2}\right)$$

and the total curvature by

$$\tau(M) = \int_M K dA = -\int_M \left( \frac{2|g_1'|^2}{(1+|g_1|^2)^2} + \frac{2|g_2'|^2}{(1+|g_2|^2)^2} \right) |dz|^2$$

4

where dA is the surface element of M. When the total curvature of a complete minimal surface is finite, the surface is called *an algebraic minimal surface*.

THEOREM 2.1 (Huber-Osserman). An algebraic minimal surface  $x: M \to \mathbb{R}^4$  satisfies:

- (i) M is conformally equivalent to  $\overline{M} \setminus \{p_1, \ldots, p_k\}$  where  $\overline{M}$  is a compact Riemann surface, and  $p_j \in \overline{M}$   $(j = 1, \ldots, k)$ .
- (ii) The Weierstrass data  $(hdz, g_1, g_2)$  extend meromorphically to  $\overline{M}$  ([16]).

DEFINITION. We call a complete minimal surface in  $\mathbb{R}^4$  pseudo-algebraic, if the following conditions are satisfied:

- (i) The Weierstrass data  $(hdz, g_1, g_2)$  is defined on a Riemann surface  $M = \overline{M} \setminus \{p_1, \ldots, p_k\}$  where  $\overline{M}$  is a compact Riemann surface, and  $p_j \in \overline{M} \ (j = 1, \ldots, k)$ .
- (ii) The Weierstrass data  $(hdz, g_1, g_2)$  can extend meromorphically to  $\overline{M}$ .

We call M the basic domain of the pseudo-algebraic minimal surface.

Since we do not assume the period condition on M, a pseudo-algebraic minimal surface is defined on some covering surface of M, in the worst case, on the universal covering.

Algebraic minimal surfaces are certainly pseudo-algebraic. The following examples are pseudo-algebraic.

EXAMPLE 2.2 (Mo-Osserman [14]). We consider the Weierstrass data

$$(hdz, g_1, g_2) = \left(\frac{dz}{\prod_{i=1}^3 (z - a_i)}, z, z\right).$$

on  $M = \mathbf{C} \setminus \{a_1, a_2, a_3\}$  for distinct  $a_1, a_2, a_3 \in \mathbf{C}$ . As this data does not satisfy period condition, we get a minimal surface  $x \colon \mathbf{D} \to \mathbf{R}^4$  on the universal covering disk  $\mathbf{D}$  of M. In particular, it has infinite total curvature. We can see that the surface is complete and both  $g_1$  and  $g_2$  omit four values  $a_1, a_2, a_3, \infty$ . This surface does not lie fully in  $\mathbf{R}^4$  because the component function  $x_3$  is equal to 0. Thus it is one-degenerate. (For details, see [9])

EXAMPLE 2.3 (Mo-Osserman [14]). We cosider the Weierstrass data

$$(hdz, g_1, g_2) = \left(\frac{dz}{\prod_{i=1}^2 (z - a_i)}, z, 0\right).$$

on  $M = \mathbb{C} \setminus \{a_1, a_2\}$  for distinct  $a_1, a_2 \in \mathbb{C}$ . As this data does not satisfy period condition, we get a complete minimal surface with infinite total curvature  $x : \mathbb{D} \to \mathbb{R}^4$  on the universal covering disk  $\mathbb{D}$  of M. We can see that  $g_1$  omits three values  $a_1, a_2, \infty$ . This surface is a complex curve in  $\mathbb{C}^2 \simeq \mathbb{R}^4$ because the Gauss map  $g_2$  is constant.

# 3. On the totally ramified value number of the Gauss map of a pseudo-algebraic minimal surface in ${f R}^4$

We recall the definition of the totally ramified value number.

DEFINITION (Nevanlinna [15]). Let M be a Riemann surface and f a meromorphic function on M. We call  $b \in \hat{\mathbf{C}}$  a totally ramified value of f when at all the inverse image points of b, f branches. We regard exceptional values also as totally ramified values. Let  $\{a_1, \ldots, a_{r_0}, b_1, \ldots, b_{l_0}\} \in \hat{\mathbf{C}}$  be the set of all totally ramified values of f, where  $a_j$   $(j = 1, \ldots, r_0)$  are exceptional values. For each  $a_j$ , put  $\nu_j = \infty$ , and for each  $b_j$ , define  $\nu_j$  to be the minimum of the multiplicities of f at points  $f^{-1}(b_j)$ . Then we have  $\nu_j \geq 2$ . We call

$$\nu_f = \sum_{a_j, b_j} \left( 1 - \frac{1}{\nu_j} \right) = r_0 + \sum_{j=1}^{l_0} \left( 1 - \frac{1}{\nu_j} \right)$$

the totally ramified value number of f.

We obtain upper bound estimates for the totally ramified value number of the Gauss map of pseudo-algebraic minimal surfaces in  $\mathbb{R}^4$ .

THEOREM 3.1. Consider a non-flat pseudo-algebraic minimal surface in  $\mathbf{R}^4$  with the basic domain  $M = \overline{M} \setminus \{p_1, \ldots, p_k\}$ . Let G be the genus of  $\overline{M}$ ,  $d_i$  be the degree of  $g_i$  considered as a map  $\overline{M}$  and  $\nu_{g_i}$  be the totally ramified value number of  $g_i$ .

(i) If  $g_1 \not\equiv constant$  and  $g_2 \not\equiv constant$ , then  $\nu_{g_1} \leq 2$ , or  $\nu_{g_2} \leq 2$ , or

(1) 
$$\frac{1}{\nu_{g_1} - 2} + \frac{1}{\nu_{g_2} - 2} \ge R_1 + R_2 \ge 1, \quad R_i = \frac{d_i}{2G - 2 + k} \ (i = 1, 2)$$

and for an algebraic minimal surface,  $R_1 + R_2 > 1$ .

(ii) If one of  $g_1$  and  $g_2$  is costant, say  $g_2 \equiv constant$ , then

(2) 
$$\nu_{g_1} \le 2 + \frac{1}{R_1}, \quad \frac{1}{R_1} = \frac{2G - 2 + k}{d_1} \le 1$$

and for an algebraic minimal surface,  $1/R_1 < 1$ .

REMARK 3.2. The ratios  $R_1$  and  $R_2$  have the geometrical meaning. For details, see [13, Section 6].

PROOF. By a suitable rotation of the surface, we may assume that both  $g_1$  and  $g_2$  are no pole at  $p_j$ , and have only simple poles. By completeness, hdz has poles of order  $\mu_j \geq 1$  at  $p_j$ . The period condition implies  $\mu_j \geq 2$ , however here we do not assume this. Let  $\alpha_s$  be (simple) poles of  $g_1$ ,  $\beta_t$  (simple) poles of  $g_2$ . The following table shows the relationship between zeros and poles of  $g_1$ ,  $g_2$  and hdz. The upper index means the order.

z	$\alpha_s$	$\beta_t$	$p_j$
$g_1$	$\infty^1$		
$g_2$		$\infty^1$	
hdz	$0^{1}$	$0^{1}$	$\infty^{\mu_i}$

Applying the Riemann-Roch formula to the meromorphic differential hdz on  $\overline{M}$ , we get

$$d_1 + d_2 - \sum_{i=1}^k \mu_i = 2G - 2$$
.

Note that this equality depends on above setting of poles of  $g_1$  and  $g_2$ , though  $d_1$  and  $d_2$  are invariant. Thus we obtain

(3) 
$$d_1 + d_2 = 2G - 2 + \sum_{i=1}^k \mu_i \ge 2G - 2 + k$$

and

(4) 
$$R_1 + R_2 = \frac{d_1 + d_2}{2G - 2 + k} \ge 1$$

When M is an algebraic minimal surface, we have  $\mu_j \ge 2$  and so  $R_1 + R_2 > 1$ .

Now we prove (i). Assume  $g_i$  is not constant and omits  $r_{i0}$  values. Let  $n_{i0}$  be the sum of the branching orders of  $g_i$  at the inverse image of these exceptional values. We see

$$(5) k \ge d_i r_{i0} - n_{i0}.$$

Let  $b_{i1}, \ldots, b_{il_0}$  be the totally ramified values which are not exceptional values. Let  $n_{ir}$  be the sum of branching orders of  $g_i$  at the inverse image of these values. For each  $b_{ij}$ , we denote

$$\nu_{ij} = \min_{g^{-1}(b_{ij})} \{ \text{multiplicity of } g(z) = b_{ij} \} ,$$

then the number of points in the inverse image  $g_i^{-1}(b_{ij})$  is less than or equal to  $d_i/\nu_{ij}$ . Thus we obtain

(6) 
$$d_i l_0 - n_{ir} \le \sum_{j=1}^{l_0} \frac{d_i}{\nu_{ij}}$$

This implies

(7) 
$$l_0 - \sum_{j=1}^{l_0} \frac{1}{\nu_{ij}} \le \frac{n_{ir}}{d_i} \; .$$

Let  $n_{i1}$  be the total branching order of  $g_i$ . Then applying Riemann-Hurwitz's theorem to the meromorphic function  $g_i$  on  $\overline{M}$ , we obtain

(8) 
$$n_{i1} = 2(d_i + G - 1)$$
.

Thus we get

(9) 
$$\nu_{g_i} = r_{i0} + \sum_{j=1}^{l_0} \left( 1 - \frac{1}{\nu_{ij}} \right) \le \frac{n_{i0} + k}{d_i} + \frac{n_{ir}}{d_i} \le \frac{n_{i1} + k}{d_i} = 2 + \frac{1}{R_i}.$$

If  $\nu_{g_1} > 2$  and  $\nu_{g_2} > 2$ , then

$$\frac{1}{\nu_{g_i} - 2} \ge R_i \quad (i = 1, 2) \; .$$

Thus we get

$$\frac{1}{\nu_{g_1} - 2} + \frac{1}{\nu_{g_2} - 2} \ge R_1 + R_2 \; .$$

Next, we prove (ii). Then the simple poles of  $g_1$  coincides exactly with the simple zeros of hdz and hdz has a pole of order  $\mu_j$  at each  $p_j$ . Applying the Riemann-Roch formula to the meromorphic differential hdz on  $\overline{M}$ , we obtain

$$d_1 - \sum_{i=1}^k \mu_i = 2G - 2$$
.

Thus we get

(10) 
$$d_1 = 2G - 2 + \sum_{i=1}^k \mu_i \ge 2G - 2 + k$$

and

(11) 
$$\frac{1}{R_1} = \frac{2G - 2 + k}{d_1} \le 1 \; .$$

When M is an algebraic minimal surface, we have  $\mu_j \ge 2$  and so  $R_1 > 1$ . By (9), we get

$$\nu_{g_1} \le 2 + \frac{1}{R_1} \; .$$

Thus, we complete the proof of this theorem.

We have the following result as an immediate consequence of Theorem 3.1.

COROLLARY 3.3 (Fujimoto[3], Hoffman-Osserman[9]). Let  $x: M \to \mathbb{R}^4$ be a pseudo-algebraic minimal surface,  $g = (g_1, g_2)$  be its Gauss map.

- (i) If both g<sub>1</sub> and g<sub>2</sub> omit more than four values, then M must be a plane. In particular, if M is an algebraic minimal surface and if both g<sub>1</sub> and g<sub>2</sub> omit more than three values, then M must be a plane.
- (ii) If one of g<sub>1</sub> and g<sub>2</sub> is constant, say g<sub>2</sub> ≡ constant and the other g<sub>1</sub> omits more than three values, then M must be a plane. In particular, if M is an algebraic minimal surface and if g<sub>1</sub> omits more than two values, then M must be a plane.

Example 2.2 and Example 2.3 show Corollary 3.3 is the best possible for pseudo-algebraic minimal surfaces. The following example shows Corollary 3.3 (ii) is the best possible for algebraic minimal surfaces.

EXAMPLE 3.4. We consider the Weierstrass data on  $M = \widehat{\mathbf{C}} \setminus \{0, \infty\}$ , by

$$(hdz,g_1,g_2) = \left(\frac{dz}{z^3},z,c\right),$$

where c is a complex number. This data satisfy conformality condition, period condition on M and the surface is complete. Thus we get an algebraic minimal surface  $x: M \to \mathbb{R}^4$  whose Gauss map  $g_1$  omits two values  $0, \infty$ .

However we do not know whether Corollary 3.3 (i) is the best possible for algebraic minimal surfaces.

REMARK 3.5. S.S. Chern and Osserman [2], Fujimoto [7], Ru [18] and Jin and Ru [11] obtain the results on the number of exceptinal values (hyperplanes) and the totally ramified value (hyperplane) number for the generalized Gauss map  $g: M \to \mathbf{P}^{n-1}(\mathbf{C})$  of a complete minimal surface in  $\mathbf{R}^n$ . However these results do not contain Theorem 3.1 and Corollary 3.3 because corresponding hyperplanes in  $\mathbf{P}^3(\mathbf{C})$  are not necessary located in general position. (For details, see [14, p353].)

# 4. Unicity theorem for the Gauss map of a pseudo-algebraic minimal surface in $\mathbf{R}^4$

We give an application of Theorem 3.1. It is a unicity theorem for the Gauss map of a pseudo-algebraic minimal surface in  $\mathbb{R}^4$ .

THEOREM 4.1. Consider two non-flat pseudo-algebraic minimal surfaces in  $\mathbb{R}^4$   $M_A$ ,  $M_B$  with the same basic domain  $M = \overline{M} \setminus \{p_1, \ldots, p_k\}$ . Let Gbe the genus of  $\overline{M}$ , and  $g_A = (g_{A1}, g_{A2})$ ,  $g_B = (g_{B1}, g_{B2})$  be the Gauss map of  $M_A$  and  $M_B$  respectively. For each i (i = 1, 2), assume that  $g_{Ai}$  and  $g_{Bi}$ have the same degree  $d_i$  when considered as a map on  $\overline{M}$ .

- (i) Let  $a_1, \ldots, a_p \in \widehat{\mathbf{C}}, b_1, \ldots, b_q \in \widehat{\mathbf{C}}$  be distinct points such that  $g_{A1}^{-1}(a_j) \cap M = g_{B1}^{-1}(a_j) \cap M$  for  $1 \leq j \leq p$ ,  $g_{A2}^{-1}(b_k) \cap M = g_{B2}^{-1}(b_k) \cap M$  for  $1 \leq k \leq q$  respectively. If  $g_{A1} \neq g_{B1}, g_{A2} \neq g_{B2}, p > 4$  and q > 4, then we have
- (12)  $\frac{1}{p-4} + \frac{1}{q-4} \ge R_1 + R_2 \ge 1, \quad R_i = \frac{d_i}{2G-2+k} \ (i=1,2) \ .$

In particular, if  $p \ge 7$  and  $q \ge 7$  then  $g_A \equiv g_B$ .

(ii) Let  $a_1, \ldots, a_p \in \widehat{\mathbf{C}}$  be distinct points such that  $g_{A1}^{-1}(a_j) \cap M = g_{B1}^{-1}(a_j) \cap M$ M for  $1 \leq j \leq p$ . If  $g_{A1} \neq g_{B1}$  and  $g_{A2} \equiv g_{B2} \equiv constant$ , then we have

(13) 
$$p \le 4 + \frac{1}{R_1}, \quad \frac{1}{R_1} = \frac{2G - 2 + k}{d_1} \le 1.$$

In particular, if  $p \ge 6$  then  $g_A \equiv g_B$ .

PROOF. Put

$$\delta_j = \sharp(g_{A1}^{-1}(a_j) \cap M) = \sharp(g_{B1}^{-1}(a_j) \cap M)$$

where  $\sharp$  denotes the number of points. Then we have

(14) 
$$pd_1 \le \sum_{j=1}^p \delta_j + n_{11} + k$$

using the same notation as in proof of Theorem 3.1. Consider a meromorphic function  $\varphi = 1/(g_{A1} - g_{B1})$  on M.  $\varphi$  has a pole, while the total number of the poles of  $\varphi$  is at most  $2d_1$ , we get

(15) 
$$\sum_{j=1}^{p} \delta_j \le 2d_1 \; .$$

Thus we obtain

$$pd_1 \le 2d_1 + n_{11} + k$$

and

(16) 
$$p \le \frac{2d_1 + n_{11} + k}{d_1} = 4 + \frac{1}{R_1} \ .$$

In the same way, we obtain

$$(17) q \le 4 + \frac{1}{R_2}$$

Thus we get (12) and (13) immediately.

We give an example which shows (p,q) = (7,7) in Theorem 4.1 (i) is the best possible for pseudo-algebraic minimal surfaces.

EXAMPLE 4.2. Taking a complex number  $\alpha$  with  $\alpha \neq 0, \pm 1$ , we consider the Weierstrass data

$$(hdz, g_1, g_2) = \left(\frac{dz}{z(z-\alpha)(\alpha z - 1)}, z, z\right)$$

and the universal covering surface M of  $\mathbb{C}\setminus\{0, \alpha, 1/\alpha\}$ . Then we can construct a pseudo-algebraic minimal surface on M. On the other hand, we can also construct a pseudo-algebraic minimal surface on M whose Weierstrass data is

$$(hdz, \widehat{g_1}, \widehat{g_2}) = \left(\frac{dz}{z(z-\alpha)(\alpha z-1)}, \frac{1}{z}, \frac{1}{z}\right).$$

For the maps  $g_i$  and  $\widehat{g}_i$ , we have  $g_i \neq \widehat{g}_i$  and  $g_i^{-1}(\alpha_j) = \widehat{g}_i^{-1}(\alpha_j)$  for six values

$$\alpha_1 := 0, \ \alpha_2 := \infty, \ \alpha_3 := \alpha, \ \alpha_4 := \frac{1}{\alpha}, \ \alpha_5 := 1, \ \alpha_6 := -1.$$

We also give an example which shows p = 6 in Theorem 4.1 (ii) is the best possible for pseudo-algebraic minimal surfaces.

EXAMPLE 4.3. Taking a complex number  $\alpha$  with  $\alpha \neq 0, \pm 1$ , we consider the Weierstrass data

$$(hdz, g_1, g_2) = \left(\frac{dz}{z(z-\alpha)}, z, 0\right)$$

and the universal covering surface M of  $\mathbb{C}\setminus\{0,\alpha\}$ . Then we can construct a pseudo-algebraic minimal surface on M. On the other hand, we can also construct a pseudo-algebraic minimal surface on M whose Weierstrass data is

$$(hdz, \widehat{g_1}, \widehat{g_2}) = \left(\frac{dz}{z(z-\alpha)}, \frac{1}{z}, 0\right) .$$

10

For the maps  $g_1$  and  $\hat{g}_1$ , we have  $g_1 \not\equiv \hat{g}_1$  and  $g_1^{-1}(\alpha_j) = \hat{g}_1^{-1}(\alpha_j)$  for five values

$$\alpha_1 := 0, \ \alpha_2 := \infty, \ \alpha_3 := \alpha, \ \alpha_4 := 1, \ \alpha_5 := -1.$$

### Acknowledgement.

The author expresses gratitude to Profs Ryoichi Kobayashi, Shin Nayatani and Reiko Miyaoka for many helpful comments.

### References

- C.C. Chen, On the image of generalized Gauss map of a complete minimal surface in R<sup>4</sup>, Pacific J. Math., **102** (1982), 9 – 14.
- [2] S.S. Chern and R. Osserman, Complete minimal surface in Euclidean n-space, J. Analyse. Math., 19 (1967), 15 – 34.
- [3] H. Fujimoto, On the number of exceptional values of the Gauss map of minimal surfaces, J. Math. Soc. Japan, 40 (1988), 235 – 247.
- [4] H. Fujimoto, On the Gauss curvature of minimal surfaces, J. Math. Soc. Japan, 44 (1992), 427–439.
- [5] H. Fujimoto, Unicity theorems for the Gauss maps of complete minimal surfaces, J. Math. Soc. Japan, 45 (1993), 481 – 487.
- [6] H. Fujimoto, Unicity theorems for the Gauss maps of complete minimal surfaces, II, Kodai. Math. J., 16 (1993), 335 – 354.
- [7] H. Fujimoto, Value Distribution Theory of the Gauss Map of Minimal Surfaces in *R<sup>m</sup>*, Braunschweig/Wiesbaden, Vieweg, 1993.
- [8] A. Huber, On subharmonic functions and differential geometry in the large, Comment. Math. Helv., 32 (1957), 13 – 72.
- [9] D.A. Hoffman and R.Osserman, The geometry of the generalized Gauss map, Memoirs. Amer. Math. Soc., 236 (1980).
- [10] D.A. Hoffman and R.Osserman, The Gauss map of surfaces in R<sup>3</sup> and R<sup>4</sup>, Proc. London Math. Soc., 50 (1985)
- [11] L. Jin and M. Ru, Algebraic curves and the Gauss map of algebraic minimal surfaces, Differential Geometry and its Applications, 25 (2007), 701 – 712.
- [12] Y. Kawakami, On the totally ramified value number of the Gauss map of minimal surfaces, Proc. Japan Acad. 82, Ser. A (2006), 1 – 3.
- [13] Y. Kawakami, R. Kobayashi and R. Miyaoka, The Gauss map of pseudo-algebraic minimal surfaces, to appear in Forum Mathematicum, math.DG/0511543.
- [14] X. Mo and R. Osserman, On the Gauss map and total curvature of complete minimal surfaces and an extension of Fujimoto's theorem, J. Differential Geometry, **31** (1990), 343–355.
- [15] R. Nevanlinna, Analytic Function, Translated from the second German edition by Phillip Emig. Die Grundlehren der mathematischen Wissenschaften, New York, Springer, 1970.
- [16] R. Osserman, Global properties of minimal surfaces in  $\mathbf{E}^3$  and  $\mathbf{E}^n$ , Ann. of Math., 80 (1964), 340–364.

- [17] R. Osserman, A Survey of Minimal Surfaces, Second edition, New York, Dover, 1986.
- [18] M. Ru, Gauss map of minimal surfaces with ramification, Trans. A.M.S., **339** (1993), 751 764.

Graduate school of Mathematics, Nagoya University, Nagoya, 464-8602 /Japan

*E-mail address*: m02008w@math.nagoya-u.ac.jp