

# INVARIANT METRICS AND DISTANCES ON GENERALIZED NEIL PARABOLAS

NIKOLAI NIKOLOV AND PETER PFLUG

ABSTRACT. We present the Carathéodory-Reiffen metric and the inner Carathéodory distance on generalized parabolas. It turns out that on such parabolas the Carathéodory distance is not inner.

## 1. INTRODUCTION AND RESULTS

In the survey paper [3] the authors had asked for an effective formula for the Carathéodory distance  $c_{A_{2,3}}$  on the Neil parabola  $A_{2,3}$  (in the bidisc). In a recent paper, such a formula was presented by G. Knese. To repeat the main result of [4] recall that the Neil parabola is given by  $A_{2,3} := \{(z, w) \in \mathbb{D}^2 : z^2 = w^3\}$ , where  $\mathbb{D}$  denotes the open unit disc in the complex plane. Then there is the natural parametrization  $p_{2,3} : \mathbb{D} \rightarrow A_{2,3}$ ,  $p_{2,3}(\lambda) := (\lambda^3, \lambda^2)$ . Moreover, let  $\rho$  denote the Poincaré distance of the unit disc. Recall that  $\rho(\lambda, \mu) := \frac{1}{2} \log \frac{1+m_{\mathbb{D}}(\lambda, \mu)}{1-m_{\mathbb{D}}(\lambda, \mu)}$ , where  $m_{\mathbb{D}}(\lambda, \mu) := \left| \frac{\lambda - \mu}{1 - \overline{\lambda}\mu} \right|$ ,  $\lambda, \mu \in \mathbb{D}$ .

Let  $\lambda, \mu \in \mathbb{D}$ . Then Knese's result is the following one:

$$c_{A_{2,3}}(p_{2,3}(\lambda), p_{2,3}(\mu)) = \begin{cases} \rho(\lambda^2, \mu^2) & \text{if } |\alpha_0| \geq 1 \\ \rho\left(\lambda^2 \frac{\alpha_0 - \lambda}{1 - \overline{\alpha_0}\lambda}, \mu^2 \frac{\alpha_0 - \mu}{1 - \overline{\alpha_0}\mu}\right) & \text{if } |\alpha_0| < 1 \end{cases},$$

where  $\alpha_0 := \alpha_0(\lambda, \mu) := \frac{1}{2}(\lambda + \frac{1}{\overline{\lambda}} + \mu + \frac{1}{\overline{\mu}})$ . In the case when  $\lambda\mu = 0$  the formula should be read as in the case  $|\alpha_0| \geq 1$ .

Observe that if  $\lambda$  and  $\mu$  have a non-obtuse angle, i.e.,  $\operatorname{Re}(\lambda\overline{\mu}) \geq 0$ , then  $|\alpha_0(\lambda, \mu)| > 1$  (compare with Corollary 2).

---

2000 *Mathematics Subject Classification.* 32F45.

*Key words and phrases.* generalized Neil parabola, Carathéodory and Kobayashi pseudodistance, Carathéodory-Reiffen pseudometric, Kobayashi-Royden pseudometric.

This note was written during the stay of the first named author at the Universität Oldenburg supported by a grant from the DFG (January – March 2006). He likes to thank both institutions for their support.

Moreover, in [4] the formula for the Carathéodory-Reiffen pseudometric  $\gamma_{A_{2,3}}$  is given as:

$$\gamma_{A_{2,3}}((a, b); X) = \begin{cases} |X_2| & \text{if } a = b = 0, |X_2| \geq 2|X_1| \\ |X_1| & \text{if } a = b = 0, |X_2| < 2|X_1| \\ \frac{2|\lambda b|}{1-|b|^2} & \text{if } (a, b) \neq (0, 0), X = \lambda(3a, 2b), \lambda \in \mathbb{C} \end{cases},$$

where  $(a, b) \in A_{2,3}$  and  $X \in T_{(a,b)}A_{2,3} :=$  the tangent space in  $(a, b)$  at  $A_{2,3}$ .

We point out that these are the first effective formulas for the Carathéodory distance and the Carathéodory-Reiffen pseudodistance of a non-trivial complex space.

In this paper we will discuss more general Neil parabolas, namely the spaces

$$A_{m,n} := \{(z, w) \in \mathbb{D}^2 : z^m = w^n\}, \quad m, n \in \mathbb{N}, \quad m \leq n, \quad \text{relatively prime.}$$

For short, we will call  $A_{m,n}$  the  $(m,n)$ -parabola. As in the case of the classical Neil parabola we have the following globally bijective holomorphic parametrization of  $A_{m,n}$ , namely

$$p_{m,n} : \mathbb{D} \longrightarrow A_{m,n}, \quad p_{m,n}(\lambda) := (\lambda^n, \lambda^m), \quad \lambda \in \mathbb{D}.$$

Observe that  $q_{m,n} := p_{m,n}^{-1} : A_{m,n} \longrightarrow \mathbb{D}$  is given outside of the origin by  $q_{m,n}(z, w) = z^k w^l$  where  $k, l \in \mathbb{Z}$  are such that  $kn + lm = 1$ ; moreover,  $q_{m,n}(0, 0) = 0$ . It is clear that  $q_{m,n}$  is continuous on  $A_{m,n}$  and holomorphic outside of the origin.

We will study the Carathéodory and the Kobayashi distances and also the Carathéodory-Reiffen and the Kobayashi-Royden pseudometrics of  $A_{m,n}$ . So let us recall the objects we will deal with in this paper:

$$m_{A_{m,n}}(\zeta, \eta) := \sup\{m_{\mathbb{D}}(f(\zeta), f(\eta)) : f \in \mathcal{O}(A_{m,n}, \mathbb{D})\}, \quad \zeta, \eta \in A_{m,n},$$

where  $\mathcal{O}(A_{m,n}, \mathbb{D})$  denotes the family of holomorphic functions on  $A_{m,n}$ , i.e., the family of those functions on  $A_{m,n}$  that are locally restriction of holomorphic functions on an open set in  $\mathbb{C}^2$ .

Observe that the Carathéodory distance  $c_{A_{m,n}}$  is given by  $c_{A_{m,n}}(\zeta, \eta) = \tanh^{-1} m_{A_{m,n}}(\zeta, \eta)$ ; moreover,  $c_{\mathbb{D}} = \rho$ .

So, we have to study holomorphic function on the  $(m,n)$ -parabola. Recall that there is the following bijection of  $\mathcal{O}(A_{m,n}, \mathbb{D})$  and a part  $\mathcal{O}_{m,n}(\mathbb{D})$  of  $\mathcal{O}(\mathbb{D}, \mathbb{D})$ , where

$$\mathcal{O}_{m,n}(\mathbb{D}) := \{h \in \mathcal{O}(\mathbb{D}, \mathbb{D}) : h^{(s)}(0) = 0, s \in S_{m,n}\}$$

and  $S_{m,n} := \{s \in \mathbb{N} : s \notin \mathbb{Z}_+ m + \mathbb{Z}_+ n\}$  (recall that  $S_{1,n} = \emptyset$  and if  $m \geq 2$ , then  $\max_{s \in S_{m,n}} s = nm - m - n$ ). To be precise, if  $f \in \mathcal{O}(A_{m,n}, \mathbb{D})$

then  $f \circ p_{m,n} \in \mathcal{O}_{m,n}(\mathbb{D})$ , and conversely, if  $h \in \mathcal{O}_{m,n}(\mathbb{D})$  then  $h \circ q_{m,n} \in \mathcal{O}(A_{m,n}, \mathbb{D})$ .

From this consideration it follows that there is the following description of the Carathéodory distance on  $A_{m,n}$ :

$$\begin{aligned} m_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu)) &= \max\{m_{\mathbb{D}}(h(\lambda), h(\mu)) : h \in \mathcal{O}_{m,n}(\mathbb{D})\} \\ &= \max\{m_{\mathbb{D}}(h(\lambda), h(\mu)) : h \in \mathcal{O}_{m,n}(\mathbb{D}), h(0) = 0\} \\ &= \max\{m_{\mathbb{D}}(\lambda^m h(\lambda), \mu^m h(\mu)) : h \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}}), \\ &\quad h^{(j)}(0) = 0, j + m \in S_{m,n}\}, \quad \lambda, \mu \in \mathbb{D}. \end{aligned}$$

We like to mention that the calculation of the Carathéodory distance of a generalized Neil parabola may be read as the following interpolation problem for holomorphic functions on the unit disc. Let  $\lambda, \mu$  be as above and let  $\zeta, \eta \in \mathbb{D}$ . Then there exists an  $h \in \mathcal{O}_{m,n}(\mathbb{D})$  with  $h(\lambda) = \zeta, h(\mu) = \eta$  if and only if  $m_{\mathbb{D}}(\zeta, \eta) \leq m_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu))$ . Note that  $m_{A_{1,n}}(p_{1,n}(\lambda), p_{1,n}(\mu)) = m_{\mathbb{D}}(\lambda, \mu)$ .

From the case of domains in  $\mathbb{C}^n$  it is well known that the Carathéodory distance need not to be an inner distance (see [2]). In the case of a generalized Neil parabola it turns out that the Carathéodory distance is an inner distance if and only if  $m = 1$ .

Recall that the associated inner distance is given by

$$c_{A_{m,n}}^i(\zeta, \eta) := \inf\{L_{c_{A_{m,n}}}( \alpha) : \alpha \text{ is a } \|\cdot\| - \text{rectifiable curve in } A_{m,n} \text{ connecting } \zeta, \eta\}, \zeta, \eta \in A_{m,n},$$

where  $L_{c_{A_{m,n}}}$  denotes the  $c_{A_{m,n}}$ -length. Obviously,  $c_{A_{m,n}} \leq c_{A_{m,n}}^i$ . Then we have the following result for the inner distance.

**Theorem 1.** *Let  $\lambda, \mu \in \mathbb{D}$ . Then*

$$\begin{aligned} c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) &= \begin{cases} c_{\mathbb{D}}(\lambda^m, \mu^m) & \text{if } \operatorname{Re}(\lambda\bar{\mu}) \geq \cos(\pi/m)|\lambda\mu| \\ c_{\mathbb{D}}(\lambda^m, 0) + c_{\mathbb{D}}(0, \mu^m) & \text{if otherwise} \end{cases}. \end{aligned}$$

Moreover, there is the following comparison result between the Carathéodory distance and its associated inner one.

**Corollary 2.** *Let  $\lambda, \mu \in \mathbb{D}$ .*

(a) *If  $\operatorname{Re}(\lambda\bar{\mu}) \geq \cos(\pi/m)|\lambda\mu|$ , then*

$$c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) = c_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu)).$$

(b) *If  $\operatorname{Re}(\lambda\bar{\mu}) < \cos(\pi/m)|\lambda\mu|$ , then*

$$c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) = c_{A_{m,n}}((p_{m,n}(\lambda), p_{m,n}(\mu)) \text{ iff } (\lambda\bar{\mu})^m < 0.$$

Thus, the following conditions are equivalent.

- $c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) = c_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu));$
- $c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) = c_{\mathbb{D}}(\lambda^m, \mu^m);$
- $\operatorname{Re}(\lambda\bar{\mu}) \geq \cos(\pi/m)|\lambda\mu|$  or  $(\lambda\bar{\mu})^m < 0.$

In particular,  $c_{A_{m,n}}$  is not inner if  $m > 1.$

Observe that the condition  $\operatorname{Re}(\lambda\bar{\mu}) \geq \cos(\pi/m)|\lambda\mu|$  in these results means geometrically that  $\mu$  lies inside an angular sector around  $\lambda$  of opening angle equal  $\pi/m$  (compare with Knese's result from above). Moreover, opposite to the  $A_{2,3}$ -case the new area  $(\lambda\bar{\mu})^m < 0$  (i.e., the "rays" on which the angle between  $\lambda$  and  $\mu$  equals to  $\frac{(2j-1)\pi}{m}$ ,  $j = 2, \dots, m-1$ ) appears for  $A_{m,n}$  with  $m > 2.$

In order to prove Theorem 1, we have to calculate the Carathéodory-Reiffen metric  $\gamma_{A_{m,n}}$  outside of the origin.

First, let us recall its definition

$$\gamma_{A_{m,n}}((z, w); X) := \max\{|f'(z, w)X| : f \in \mathcal{O}(A_{m,n}, \mathbb{D})\},$$

where  $(z, w) \in A_{m,n}$  and  $X$  a tangent vector in  $(z, w)$  at  $A_{m,n}$ . Recall that if  $(z, w) = \zeta = p_{m,n}(\lambda)$ ,  $\lambda \in \mathbb{D} \setminus \{0\}$ , then the tangent space  $T_{\zeta}(A_{m,n})$  at  $\zeta$  is spanned by the vector  $p'_{m,n}(\lambda)$  whereas the tangent space at the origin equals  $\mathbb{C}^2$  if  $m \geq 2.$

Using the above description of  $\mathcal{O}(A_{m,n}, \mathbb{D})$  we may reformulate this definition for  $\lambda \in \mathbb{D}$ ,  $\lambda \neq 0$ , in the following appropriate form which will be used here:

$$\gamma_{A_{m,n}}(p_{m,n}(\lambda); p'_{m,n}(\lambda)) = \sup\left\{\frac{|h'(\lambda)|}{1 - |h(\lambda)|^2} : h \in \mathcal{O}_{m,n}(\mathbb{D})\right\}.$$

Then we have the following result.

**Theorem 3.** *Let  $\lambda \in \mathbb{D}$ ,  $\lambda \neq 0$ . Then*

$$\gamma_{A_{m,n}}(p_{m,n}(\lambda); p'_{m,n}(\lambda)) = \frac{m|\lambda|^{m-1}}{1 - |\lambda|^{2m}}.$$

On the other hand, note that

$$\gamma_{A_{m,n}}(0; X) = \max\{|f'(z, w)X| : f \in \mathcal{O}(A_{m,n}, \mathbb{D}), f(0) = 0\}.$$

Then for such  $f$  we have  $f \circ p_{m,n}(\zeta) = \zeta^m h(\zeta)$ ,  $\zeta \in \mathbb{D}$ , where  $h \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}})$ . Observe that  $\frac{\partial f}{\partial z}(0) = \frac{h^{(n-m)}(0)}{(n-m)!}$  and  $\frac{\partial f}{\partial w}(0) = h(0)$ . Thus, if  $m \geq 2$  and  $X = (X_1, X_2) \in \mathbb{C}^2$ , then

$$\begin{aligned} \gamma_{A_{m,n}}(0; X) &= \max\left\{|X_1 \frac{h^{(n-m)}(0)}{(n-m)!} + X_2 h(0)| : \right. \\ &\quad \left. h \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}}), h^{(j)}(0) = 0, j + m \in S_{m,n}\right\}; \end{aligned}$$

in particular,  $\gamma_{A_{m,n}}(0; X) = \|X\|$  if  $X_1 X_2 = 0$ .

It turns out that the general calculation of the Carathéodory-Reiffen metric at the origin becomes much more difficult. The next theorem may give some flavor of the nature of this formulas.

**Proposition 4.** *Let  $X = (X_1, X_2) \in \mathbb{C}^2$ . Then*

$$\gamma_{A_{3,4}}(0; X) = \begin{cases} |X_1| & \text{if } |X_1| \geq 2|X_2| \\ |X_2| & \text{if } |X_2| \geq \sqrt{2}|X_1| \\ |X_1|^{\frac{c^3 - 18c + (c^2 + 24)^{3/2}}{108}} & \text{if } 1 < c := 2\frac{|X_2|}{|X_1|} < 2\sqrt{2} \end{cases}.$$

It seems rather difficult to calculate an effective formula of the Carathéodory distance of  $A_{m,n}$ . However, we have its value at pairs of “opposite” points; to be more precise the following is true.

**Proposition 5.** *Let  $\lambda \in \mathbb{D}$ ,  $\lambda \neq 0$ . Then*

$$m_{A_{2,2k+1}}(p_{2,2k+1}(\lambda), p_{2,2k+1}(-\lambda)) = \frac{2|\lambda|^{2k+1}}{1 + |\lambda|^{4k+2}}.$$

Observe that now, opposite to the cases before, the number  $n = 2k + 1$  appears in the formula.

Finally, the discussion of the Kobayashi distance and the Kobayashi-Royden metric on  $A_{m,n}$  becomes comparably much simpler. Let us first recall the definitions of the Lempert function  $\tilde{k}_{A_{m,n}}$ , the Kobayashi distance  $k_{A_{m,n}}$  and the Kobayashi-Royden metric  $\kappa_{A_{m,n}}$ .

- $\tilde{k}_{A_{m,n}}(\zeta, \eta) := \inf\{\rho(\lambda, \mu) : \lambda, \mu \in \mathbb{D} \exists \varphi \in \mathcal{O}(\mathbb{D}, A_{m,n}) : \varphi(\lambda) = \zeta, \varphi(\mu) = \eta\}, \quad \zeta, \eta \in A_{m,n};$
- $k_{A_{m,n}} :=$  the largest distance on  $A_{m,n}$  below of  $\tilde{k}_{A_{m,n}}$ ;
- $\kappa_{A_{m,n}}(\zeta; X) := \inf\{\alpha \in \mathbb{R}_+ : \exists \varphi \in \mathcal{O}(\mathbb{D}, A_{m,n}) : \varphi(0) = \zeta, \alpha\varphi'(0) = X\}, \quad \zeta \in A_{m,n}, X \in T_\zeta(A_{m,n}).$

We set  $\tilde{k}_{A_{m,n}}(\zeta, \eta) := \infty$  or  $\kappa_{A_{m,n}}(\zeta; X) := \infty$  if there are no respective discs  $\varphi$ .

Since  $\mathcal{O}(\mathbb{D}, A_{m,n}) = \{p_{m,n} \circ \psi : \psi \in \mathcal{O}(\mathbb{D}, \mathbb{D})\}$ , then we have the following formulas (see also [3, 4]).

**Proposition 6.** *Let  $\lambda, \mu \in \mathbb{D}$ . Then*

$$k_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu)) = \tilde{k}_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu)) = \rho(\lambda, \mu).$$

$$\text{If } \lambda \neq 0, \text{ then } \kappa_{A_{m,n}}(p_{m,n}(\lambda); p'_{m,n}(\lambda)) = \frac{1}{1 - |\lambda|^2}.$$

$$\text{Let } m \geq 2 \text{ and } X \in \mathbb{C}^2 \setminus \{0\}. \text{ Then } \kappa_{A_{m,n}}(0; X) = \infty.$$

At the end of the paper a simple reducible variety is also discussed.

## 2. PROOFS AND ADDITIONAL REMARKS

We start with the proof of Theorem 3 which will serve as the basic information for Theorem 1.

*Proof of Theorem 3.* Recall that

$$\gamma_{A_{m,n}}(p_{m,n}(\lambda); p'_{m,n}(\lambda)) = \max\left\{\frac{|h'(\lambda)|}{1 - |h(\lambda)|^2} : h \in \mathcal{O}_{m,n}(\mathbb{D})\right\}.$$

Observe that if  $\alpha \in \mathbb{D}$  and  $\Phi_\alpha(\zeta) = \frac{\alpha - \zeta}{1 - \bar{\alpha}\zeta}$ , then  $h_\alpha = \Phi_\alpha \circ h \in \mathcal{O}_{m,n}(\mathbb{D})$  (use, for example, the Faà di Bruno formula) and

$$\frac{|h'_\alpha(\lambda)|}{1 - |h_\alpha(\lambda)|^2} = \frac{|h'(\lambda)|}{1 - |h(\lambda)|^2}.$$

Then

$$\begin{aligned} \gamma_{A_{m,n}}(p_{m,n}(\lambda); p'_{m,n}(\lambda)) &= \max\left\{\frac{|h'(\lambda)|}{1 - |h(\lambda)|^2} : h \in \mathcal{O}_{m,n}(\mathbb{D}), h(0) = 0\right\} \\ &= \max\left\{\frac{|(\lambda^m \tilde{h}(\lambda))'|}{1 - |\lambda^m \tilde{h}(\lambda)|^2} : \tilde{h} \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}}), \tilde{h}^{(j)}(0) = 0, j + m \in S_{m,n}\right\} \\ &= |\lambda|^{m-1} \max\left\{\frac{|mh(\lambda) + \lambda h'(\lambda)|}{1 - |\lambda^m h(\lambda)|^2} : \right. \\ &\quad \left. h \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}}), h^{(j)}(0) = 0, j + m \in S_{m,n}\right\} = \frac{m|\lambda|^{m-1}}{1 - |\lambda|^{2m}}. \end{aligned}$$

The last equality is a consequence of the fact that the unimodular constants are the only extremal functions for

$$\max\left\{\frac{|mh(\lambda) + \lambda h'(\lambda)|}{1 - |\lambda^m h(\lambda)|^2} : h \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}})\right\}.$$

To prove this fact, observe that  $(h(\lambda), h'(\lambda))$  varies on all pairs  $(a, b)$  satisfying  $|b| \leq \frac{1 - |a|^2}{1 - |\lambda|^2}$ . Thus, we have to show that if  $0 \leq c, s < 1$  and  $0 \leq t \leq t_s := \frac{1 - s^2}{1 - c^2}$ , then  $F(s, t) < F(1, 0)$ , where  $F(s, t) = \frac{ms + ct}{1 - c^{2m}s^2}$ . Since  $F(s, t) \leq F(s, t_s)$ , the problem may be reduced to the inequality

$$\frac{m(1 - c^2)s + c(1 - s^2)}{1 - c^{2m}s^2} < \frac{m(1 - c^2)}{1 - c^{2m}} \iff \frac{c(1 - c^{2m})}{m(1 - c^2)} < \frac{1 + c^{2m}s}{1 + s}.$$

Using the inequality  $\frac{1+c^{2m}}{2} < \frac{1+c^{2m}s}{1+s}$ , one has to see that

$$\frac{c(1-c^{2m})}{m(1-c^2)} < \frac{1+c^{2m}}{2} \iff 2c \sum_{j=0}^{m-1} c^{2j} < m(1+c^{2m}).$$

Finally, by summing up the inequalities  $1 - c^{2j+1} > c^{2m-2j-1}(1 - c^{2j+1})$  for  $j = 0, \dots, m-1$ , the last inequality follows.  $\square$

Now, we are in the position to prove Theorem 1.

*Proof of Theorem 1.* Set  $A_{\lambda,m} = \{\zeta \in \mathbb{D} : \operatorname{Re}(\lambda\bar{\zeta}) \geq \cos(\pi/m)|\lambda\zeta|\}$ ,  $\lambda \in \mathbb{D}$ ,  $m \in \mathbb{N}$ . Recall again that  $A_{\lambda,m}$  is an angular sector around  $\lambda$ .

In a first step we shall prove that if  $\lambda \in \mathbb{D}$  and  $\mu \in A_{\lambda,m}$ , then

$$c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) = c_{\mathbb{D}}(\lambda^m, \mu^m).$$

Since

$$(1) \quad c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) \geq c_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu)) \geq c_{\mathbb{D}}(\lambda^m, \mu^m),$$

we have only to prove the opposite inequality. After rotation, we may assume that  $\lambda \in [0, 1)$ . By continuity, we may also assume that  $\lambda, \mu \neq 0$  and  $\arg(\mu) \in (-\pi/m, \pi/m)$ . Then the geodesic for  $c_{\mathbb{D}}^i(\lambda^m, \mu^m)$  does not intersect the segment  $(-1, 0]$ . Denote by  $\alpha$  this geodesic and by  $\alpha_m$  its  $m$ -th root ( $\sqrt[m]{1} = 1$ ). Observe that if  $\zeta, \eta \in A_{m,n}^* := A_{m,n} \setminus \{0\}$ , then

$$c_{A_{m,n}}^i(\zeta, \eta) = \inf \left\{ \int_0^1 \gamma_{A_{m,n}}(\alpha(t); \alpha'(t)) dt : \alpha : [0, 1] \rightarrow A_{m,n}^* \right. \\ \left. \text{is a } C^1\text{-curve connecting } \zeta, \eta \right\}$$

(see Theorem 4.2.7 in [5]).

It follows by Theorem 3 that

$$\begin{aligned} c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) &\leq \int_0^1 \gamma_{A_{m,n}}(p_{m,n} \circ \alpha_m(t); (p_{m,n} \circ \alpha_m)'(t)) dt \\ &= \int_0^1 \frac{m|\alpha_m(t)|^{m-1} \alpha_m'(t)}{1 - |\alpha_m(t)|^{2m}} dt = \int_0^1 \frac{|\alpha'(t)|}{1 - |\alpha(t)|^2} dt \\ &= c_{\mathbb{D}}^i(\lambda^m, \mu^m) = c_{\mathbb{D}}(\lambda^m, \mu^m). \end{aligned}$$

It remains to prove that if  $\mu \notin A_{\alpha,m}$ , then

$$c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) = c_{A_{m,n}}^i(p_{m,n}(\lambda), 0) + c_{A_{m,n}}^i(0, p_{m,n}(\mu)).$$

By the triangle inequality, we only have to prove that

$$(2) \quad c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) \geq c_{A_{m,n}}^i(p_{m,n}(\lambda), 0) + c_{A_{m,n}}^i(0, p_{m,n}(\mu)).$$

Take an arbitrary  $C^1$ -curve  $\alpha : [0, 1] \rightarrow A_{m,n}^*$  with  $\alpha(0) = p_{m,n}(\lambda)$  and  $\alpha(1) = p_{m,n}(\mu)$ . Let  $t_0 \in (0, 1)$  be the smallest numbers such that  $\alpha(t_0) \in \partial A_{\alpha,m}$ . If  $\alpha(t_0) = p(\lambda_0)$ , then

$$\begin{aligned}
& \int_0^1 \gamma_{A_{m,n}}(\alpha(t); \alpha'(t)) dt \\
&= \int_0^{t_0} \gamma_{A_{m,n}}(\alpha(t); \alpha'(t)) dt + \int_{t_0}^1 \gamma_{A_{m,n}}(\alpha(t); \alpha'(t)) dt \\
&\geq c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\lambda_0)) + c_{A_{m,n}}^i(p_{m,n}(\lambda_0), p_{m,n}(\mu)) \\
&\geq c_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\lambda_0)) + c_{A_{m,n}}(p_{m,n}(\lambda_0), p_{m,n}(\mu)) \\
&\geq c_{\mathbb{D}}(\lambda^m, \lambda_0^m) + c_{\mathbb{D}}(\lambda_0^m, \mu^m) \\
&= c_{\mathbb{D}}(\lambda^m, 0) + c_{\mathbb{D}}(0, \lambda_0^m) + c_{\mathbb{D}}(\lambda_0^m, \mu^m) \quad (\text{since } \lambda_0^m \in (-1, 0)) \\
&\geq c_{\mathbb{D}}(\lambda^m, 0) + c_{\mathbb{D}}(0, \mu^m).
\end{aligned}$$

Now, (2) follows by taking the infimum over all curves under consideration.  $\square$

Next, the proof of Corollary 2 will be given.

*Proof of Corollary 2.* (a) Follows by Theorem 1 and the inequality (1).

(b) The inequalities

$$\begin{aligned}
c_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu)) &\leq \max\{c_{\mathbb{D}}(\lambda^m f(\lambda), \mu^m f(\mu)) : f \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}})\} \\
&\leq \max\{c_{\mathbb{D}}(\lambda^m f(\lambda), 0) + c_{\mathbb{D}}(0, \mu^m + f(\mu)) : f \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}})\} \\
&\leq c_{\mathbb{D}}(\lambda^m, 0) + c_{\mathbb{D}}(0, \mu^m) = c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu))
\end{aligned}$$

show that

$$c_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu)) = c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu))$$

if and only if  $\lambda^m f(\lambda)$  and  $\mu^m f(\mu)$  lie on opposite rays and  $|f(\lambda)| = |f(\mu)| = 1$  for some  $f \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}})$ , i.e.,  $f$  is a unimodular constant, and  $(\lambda \overline{\mu})^m < 0$ .

The remaining part of Corollary 2 follows by the fact that  $c_{\mathbb{D}}(z, 0) + c_{\mathbb{D}}(0, w) = c_{\mathbb{D}}(z, w)$  if and only if  $z\overline{w} \leq 0$ .  $\square$

**Remarks.** (a) For  $m \in \mathbb{N}$ , consider the following distance on  $\mathbb{D}$ :

$$\rho^{(m)}(\lambda, \mu) := \max\{\rho_{\mathbb{D}}(\lambda^m h(\lambda), \mu^m h(\mu)) : h \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}})\}.$$

Note that

$$\begin{aligned}
\lim_{\varepsilon \rightarrow 0, \varepsilon \neq 0} \frac{\rho^{(m)}(\lambda, \lambda + \varepsilon)}{|\varepsilon|} &= |\lambda|^{m-1} \max\left\{\frac{|mh(\lambda) + \lambda h'(\lambda)|}{1 - |\lambda^m h(\lambda)|^2} : h \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}})\right\} \\
&= \gamma_{A_{m,n}}(p_{m,n}(\lambda); p'_{m,n}(\lambda))
\end{aligned}$$



by the proof of Theorem 3. So it follows that the associated inner distance of  $\rho^{(m)}$  equals  $c_{A_{m,n}}^i(p_{m,n}(\cdot), p_{m,n}(\cdot))$ . Then

$$\begin{aligned} c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) &\geq \rho^{(m)}(\lambda, \mu) \\ &\geq c_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu)) \geq \rho(\lambda^m, \mu^m). \end{aligned}$$

Moreover, the proof of Corollary 2 shows that the following conditions are equivalent:

- $c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) = \rho^{(m)}(\lambda, \mu)$ ;
- $c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) = c_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu))$ ;
- $c_{A_{m,n}}^i(p_{m,n}(\lambda), p_{m,n}(\mu)) = \rho(\lambda^m, \mu^m)$ ;
- $\operatorname{Re}(\lambda\bar{\mu}) \geq \cos(\pi/m)|\lambda\mu|$  or  $(\lambda\bar{\mu})^m < 0$ .

As an application of these observations we obtain a simple proof (without calculations) of Lemma 14 in [6]:

*If  $a, b \in [0, 1]$ ,  $s \in (0, 1]$  and  $\theta \in [-\pi, \pi]$ , then  $\rho(a, be^{i\theta}) \leq \rho(a^s, b^s e^{is\theta})$ .*

In fact, we may assume that  $s \in \mathbb{Q}$ . If  $s = \frac{p}{q}$  ( $1 \leq p \leq q$ ),  $\lambda = \sqrt[q]{a}$ ,  $\mu = \sqrt[q]{b}e^{i\theta/q}$ , then we have to prove that  $\rho(\lambda^q, \mu^q) \leq \rho(\lambda^p, \mu^p)$ . But the angle between  $\lambda$  and  $\mu$  does not exceed  $\frac{\pi}{q} \leq \frac{\pi}{p}$  and hence

$$\rho(\lambda^p, \mu^p) = \rho^{(p)}(\lambda, \mu) \geq \rho(\lambda^q, \mu^q)$$

(the last inequality holds for any  $\lambda, \mu \in \mathbb{D}$  and  $q \geq p$ ).

(b) Recall that

$$\begin{aligned} c_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu)) &= \max\{\rho_{\mathbb{D}}(\lambda^m h(\lambda), \mu^m h(\mu)) : \\ &\quad h \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}}), h^{(j)}(0) = 0, j + m \in S_{m,n}\}. \end{aligned}$$

If  $m = 1$  or  $(m, n) = (2, 3)$ , then  $\rho^{(m)}(\lambda, \mu) = c_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu))$ , since  $S_{1,n} = \emptyset$  and  $S_{2,3} = \{1\}$ .

On the other hand, if  $m \neq 1$  and  $m \neq n - 1$ , then the following conditions are equivalent:

- $\rho^{(m)}(\lambda, \mu) = \rho(\lambda^m, \mu^m)$ ;
- $\rho^{(m)}(\lambda, \mu) = c_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu))$ .

It is clear that the first condition implies the second one. For the converse, observe that as  $h$  varies over  $\mathcal{O}(\mathbb{D}, \mathbb{D})$ , the pair  $(h(\lambda), h(\mu))$  varies over all  $(z, w) \in \mathbb{D}^2$  with  $m_{\mathbb{D}}(z, w) \leq m_{\mathbb{D}}(\lambda, \mu)$ . Thus,

$$\begin{aligned} \rho^{(m)}(\lambda, \mu) &= \max\{\rho_{\mathbb{D}}(\lambda^m z, \mu^m w) : z, w \in \mathbb{D} \text{ with} \\ &\quad m_{\mathbb{D}}(z, w) \leq m_{\mathbb{D}}(\lambda, \mu) \text{ or } z = w \in \partial D.\} \end{aligned}$$

It follows by the maximum principle for the continuous plurisubharmonic function  $m_{\mathbb{D}}(\lambda^m \cdot, \mu^m w)$  that if  $\rho^{(m)}(\lambda, \mu) = \rho_{\mathbb{D}}(\lambda^m z, \mu^m w)$ , then either  $z = w \in \partial D$ , or  $m_{\mathbb{D}}(z, w) = m_{\mathbb{D}}(\lambda, \mu)$ . Assuming that

$\rho^{(m)}(\lambda, \mu) \neq \rho(\lambda^m, \mu^m)$  excludes the first possibility. Then any extremal function  $h$  for  $\rho^{(m)}(\lambda, \mu)$  satisfies  $m_{\mathbb{D}}(h(\lambda), h(\mu)) = m_{\mathbb{D}}(\lambda, \mu)$ , i.e.,  $h \in \text{Aut}(\mathbb{D})$ . Since any such function should be also extremal for  $c_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu))$ , it follows that either  $h^{(j)} \neq 0$  for any  $j \in \mathbb{N}$ , or  $h$  is a rotation. In particular,  $m+1 \notin S_{m,n}$ , i.e.,  $m = 1$  or  $m = n-1$ , a contradiction.

Let  $m \geq 3$ . Then  $m+2 \notin S_{m,m+1}$  and hence  $h$  must be a rotation. Thus, the following conditions are equivalent:

- $\rho^{(m)}(\lambda, \mu) = \max\{\rho(\lambda^m, \mu^m), \rho(\lambda^{m+1}, \mu^{m+1})\}$ ;
- $\rho^{(m)}(\lambda, \mu) = c_{A_{m,m+1}}(p_{m,n}(\lambda), p_{m,n}(\mu))$ .

Concerning the first condition, we point out that

$$\rho^{(2m)}(\lambda, -\lambda) = \rho(\lambda^{2m+1}, -\lambda^{2m+1}),$$

since

$$\begin{aligned} m_{\mathbb{D}}(\lambda^{2m}\Phi_{\alpha}(\lambda), \lambda^{2m}\Phi_{\alpha}(-\lambda)) &= \frac{2(1-|\alpha|^2)|\lambda|^{2m+1}}{|1+|\lambda|^{4m+2}-|\alpha|^2(|\lambda|^2+|\lambda|^{4m})+(1-|\lambda|^{4m})(\alpha\bar{\lambda}-\bar{\alpha}\lambda)|}} \\ &\leq \frac{2(1-|\alpha|^2)|\lambda|^{2m+1}}{1+|\lambda|^{4m+2}-|\alpha|^2(|\lambda|^2+|\lambda|^{4m})} \leq \frac{2|\lambda|^{2m+1}}{1+|\lambda|^{4m+2}}, \end{aligned}$$

(use that  $1+|\lambda|^{4m+2} > |\lambda|^2+|\lambda|^{4m}$ ).

On the other hand, for  $m \geq 5$  we may choose  $\lambda \in \mathbb{D}$  such that  $2 \cos \frac{2\pi}{m} > |\lambda|^2+|\lambda|^{2m+2}$ . If  $\mu = e^{2\pi i/m}\lambda$ , then

$$\rho^{(m)}(\lambda, \mu) > \rho(\lambda^{m+1}, \mu^{m+1}) > \rho(\lambda^m, \mu^m),$$

since

$$\begin{aligned} \tanh(\rho^{(m)}(\lambda, \mu)) &\geq m_{\mathbb{D}}(\lambda^m\Phi_{\mu}(\lambda), \mu^m\Phi_{\mu}(\mu)) = \frac{|\lambda|^m|\lambda-\mu|}{|1-|\lambda|^{2m}e^{2\pi i/m}|}} \\ &> \frac{|\lambda|^m|\lambda-\mu|}{|1-|\lambda|^{2m+2}e^{2\pi i/m}|}} = m_{\mathbb{D}}(\lambda^{m+1}, \mu^{m+1}) > m_{\mathbb{D}}(\lambda^m, \mu^m) = 0. \end{aligned}$$

Observe also that for  $\lambda_0 = \sqrt{\frac{2}{3}}$  and  $\mu_0 = \sqrt{\frac{1}{3}}$  one has that

$$\begin{aligned} m_{\mathbb{D}}(\lambda_0^4\Phi_{\mu_0}(i\lambda_0), \mu_0^4\Phi_{\mu_0}(\mu_0)) &= \lambda_0^4 \sqrt{\frac{\lambda_0^2 + \mu_0^2}{1 + \lambda_0^2\mu_0^2}} > \frac{\lambda_0^4 - \mu_0^4}{1 - \lambda_0^4\mu_0^4} \\ &> \sqrt{\frac{\lambda_0^6 + \mu_0^6}{1 + \lambda_0^6\mu_0^6}} > \sqrt{\frac{\lambda_0^{10} + \mu_0^{10}}{1 + \lambda_0^{10}\mu_0^{10}}} \end{aligned}$$

and hence

$$\rho^{(3)}(i\lambda_0, \mu_0) \geq \rho^{(4)}(i\lambda_0, \mu_0) > \rho(\lambda_0^4, \mu_0^4) > \rho(-i\lambda_0^3, \mu_0^3) > \rho(i\lambda_0^5, \mu_0^5).$$

Now we turn to the discussion of the Carathéodory-Reiffen pseudo-metric on the  $(3, 4)$ -parabola.

*Proof of Proposition 4.* Recall that

$$\gamma_{A_{3,4}}(0; X) = \max\{|X_1 h'(0) + X_2 h(0)| : h \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}}), h''(0) = 0\}.$$

So, we have to describe the pairs  $(a_0, a_1) \in \mathbb{C}^2$  for which there is a function  $h \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}})$  of the form  $h(\zeta) = a_0 + a_1 \zeta + o(\zeta^2)$ . Let  $I_3$  be the  $3 \times 3$  unit matrix and

$$M = \begin{bmatrix} a_0 & a_1 & 0 \\ 0 & a_0 & a_1 \\ 0 & 0 & a_0 \end{bmatrix}.$$

It follows by Schur's theorem (cf. [1]) that such an  $h$  exists if only if  $I_3 - M^* M$  is a semipositive matrix. It is easy to check that the last conditions just means that the pair  $(|a_0|^2, |a_1|^2)$  belongs to the set

$$C := \{(a, b) \in \mathbb{R}_+^2 : a + \sqrt{b} \leq 1, ab(1 - a) \leq ((1 - a)^2 - b)(1 - a - b)\}.$$

The second inequality can be written as

$$b \leq (1 - a)(1 - \sqrt{a}) \quad \text{or} \quad b \geq (1 - a)(1 + \sqrt{a}).$$

Hence  $C = \{(a, b) \in \mathbb{R}_+^2 : b \leq (1 - a)(1 - \sqrt{a}), a \leq 1\}$ . Thus,

$$\begin{aligned} \gamma_{A_{3,4}}(0; X) &= \max\{|X_1| \sqrt{b} + |X_2| \sqrt{a} : (a, b) \in C\} \\ &= \max\{t \in [0; 1] : |X_1|(1 - t)\sqrt{1 + t} + |X_2|t\}. \end{aligned}$$

Straightforward calculations show that the last maximum is equal to

$$\begin{cases} |X_1| & \text{if } |X_1| \geq 2|X_2| \\ |X_2| & \text{if } |X_2| \geq \sqrt{2}|X_1| \\ |X_1|^{\frac{c^3 - 18c + (c^2 + 24)^{3/2}}{108}} & \text{if } 1 < c := 2\frac{|X_2|}{|X_1|} < 2\sqrt{2} \end{cases}.$$

□

Now, we shall go to prove Proposition 5.

*Proof of Proposition 5.* Recall that

$$\begin{aligned} &m_{A_{2,2k+1}}(p_{2,2k+1}(\lambda), p_{2,2k+1}(\mu)) \\ &= \max\{m_{\mathbb{D}}(f(\lambda), f(\mu)) : f \in \mathcal{O}(\mathbb{D}, \mathbb{D}), f^{(2j-1)}(0) = 0, j = 1, \dots, k\} \\ &= \max\{m_{\mathbb{D}}(\lambda^2 h(\lambda), \mu^2 h(\mu)) : \\ &\quad h \in \mathcal{O}(\mathbb{D}, \overline{\mathbb{D}}), h^{(2j-1)}(0) = 0, j = 1, \dots, k - 1\}. \end{aligned}$$

It follows that

$$m_{A_{2,2k+1}}(p_{2,2k+1}(\lambda), p_{2,2k+1}(\mu))$$

$$= \sup\{m_{\mathbb{D}}(\lambda^2 z, \mu^2 w) : m_{\mathbb{D}}(z, w) \leq m_{A_{2,2k-1}}(p_{2,2k-1}(\lambda), p_{2,2k-1}(\mu))\}.$$

Then Proposition 5 will follow by induction on  $n \in \mathbb{Z}_+$  if we show that

$$m_{\mathbb{D}}(z, w) \leq \frac{2|\lambda|^{2k-1}}{1 + |\lambda|^{4k-2}} \implies m_{\mathbb{D}}(\lambda^2 z, \lambda^2 w) \leq \frac{2|\lambda|^{2k+1}}{1 + |\lambda|^{4k+2}}.$$

Since  $\frac{2|\lambda|^{2k-1}}{1 + |\lambda|^{4k-2}} = m_{\mathbb{D}}(\lambda^{2k-1}, -\lambda^{2k-1})$ , we may assume as in Remark (b) that  $z = \Phi_{\alpha}(\lambda^{2k-1})$  and  $w = \Phi_{\alpha}(-\lambda^{2k-1})$  for some  $\alpha \in \mathbb{D}$ . Then

$$\begin{aligned} & m_{\mathbb{D}}(\lambda^2 z, \lambda^2 w) \\ &= \frac{2(1 - |\alpha|^2)|\lambda|^{2k+1}}{|1 + |\lambda|^{4k+2} - |\alpha|^2(|\lambda|^4 + |\lambda|^{4k-2}) + (1 - |\lambda|^4)(\alpha\bar{\lambda}^{2k-1} - \bar{\alpha}\lambda^{2k-1})|} \\ &\leq \frac{2(1 - |\alpha|^2)|\lambda|^{2k+1}}{1 + |\lambda|^{4k+2} - |\alpha|^2(|\lambda|^4 + |\lambda|^{4k-2})} \leq \frac{2|\lambda|^{2k+1}}{1 + |\lambda|^{4k+2}}, \end{aligned}$$

since  $1 + |\lambda|^{4k+2} > |\lambda|^4 + |\lambda|^{4k-2}$ .  $\square$

**Remark.** From the result above one may conclude the following interpolation result. Namely, for given  $k \in \mathbb{N}$ ,  $\lambda, \eta, \zeta \in \mathbb{D}$  the following conditions are equivalent:

- (i)  $m_{\mathbb{D}}(\eta, \zeta) \leq m_{\mathbb{D}}(\lambda^{2k+1}, -\lambda^{2k-1})$ ;
- (i)  $\exists f \in \mathcal{O}(\mathbb{D}, \mathbb{D}) : f(\lambda^{2k+1}) = \eta, f(-\lambda^{2k+1}) = \zeta$ ;
- (iii)  $\exists f \in \mathcal{O}(\mathbb{D}, \mathbb{D}) : f(\lambda) = \eta, f(-\lambda) = \zeta, f^{(j)}(0) = 0, j = 1, \dots, 2k$ ;
- (iv)  $\exists f \in \mathcal{O}(\mathbb{D}, \mathbb{D}) : f(\lambda) = \eta, f(-\lambda) = \zeta, f^{(2j-1)}(0) = 0, j = 1, \dots, k$ .

Indeed, it is trivial that (i)  $\implies$  (ii)  $\implies$  (iii)  $\implies$  (iv), and the implication (iv)  $\implies$  (i) follows by the equalities

$$m_{A_{2,2k+1}}(p_{2,2k+1}(\lambda), p_{2,2k+1}(\mu)) = \frac{2|\lambda|^{2k+1}}{1 + |\lambda|^{4k+2}} = m_{\mathbb{D}}(\lambda^{2k+1}, -\lambda^{2k-1}).$$

Finally, we discuss the proof for the Kobayashi distance and metric.

*Proof of Proposition 6.* The proof of the formula for  $\tilde{k}_{A_{m,n}}$  follows the one for the case  $(m, n) = (2, 3)$  (see [4]). For convenience of the reader we include it.

First,  $\tilde{k}_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu)) \leq \rho(\lambda, \mu)$  because  $p_{m,n}$  is holomorphic. Second, since  $m$  and  $n$  are relatively prime, it is easy to see that  $\mathcal{O}(\mathbb{D}, A_{m,n}) = \{p_{m,n} \circ \psi : \psi \in \mathcal{O}(\mathbb{D}, \mathbb{D})\}$ . Then any  $\varphi \in \mathcal{O}(\mathbb{D}, A_{m,n})$  with  $\varphi(\tilde{\lambda}) = p_{m,n}(\lambda)$  and  $\varphi(\tilde{\mu}) = p_{m,n}(\mu)$  corresponds to some  $\psi \in \mathcal{O}(\mathbb{D}, \mathbb{D})$  with  $\psi(\tilde{\lambda}) = \lambda$  and  $\psi(\tilde{\mu}) = \mu$ . Thus,  $\rho(\lambda, \mu) \leq \rho(\tilde{\lambda}, \tilde{\mu})$  and hence  $\rho(\lambda, \mu) \leq \tilde{k}_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu))$ . So,  $\tilde{k}_{A_{m,n}}(p_{m,n}(\lambda), p_{m,n}(\mu)) = \rho(\lambda, \mu)$ ; in particular,  $\tilde{k}_{A_{m,n}}$  is a distance and therefore  $\tilde{k}_{A_{2,3}} = k_{A_{2,3}}$ .

The formulas for  $\kappa_{A_{m,n}}$  can be proven in a similar way and we omit the details.  $\square$

We conclude this paper by mentioning the simplest example of a reducible variety.

**Remark.** Put  $A_{2,2} := \{(z, w) \in \mathbb{D}^2 : z^2 = w^2\}$ ;  $A_{2,2}$  is reducible. Obviously,  $A_{2,2}$  is biholomorphically equivalent to the coordinate cross  $V := \{(z, w) \in \mathbb{D}^2 : zw = 0\}$ . Therefore, we discuss  $V$  instead of  $A_{2,2}$ .

It is clear that  $c_V((z_1, 0), (z_2, 0)) = \tilde{k}_V((z_1, 0), (z_2, 0)) = \rho(z_1, z_2)$ ,

$$\tilde{k}_V((z, 0), (0, w)) = \infty \quad (zw \neq 0)$$

and

$$k_V((z, 0), (0, w)) = \tilde{k}_V((z, 0), (0, 0)) + \tilde{k}_V((0, 0), (0, w)) = \rho(|z|, -|w|).$$

Moreover,  $\gamma_V((z, 0); (1, 0)) = \kappa_V((z, 0); (1, 0)) = \frac{1}{1 - |z|^2}$  and

$$\kappa_V(0; X) = \begin{cases} |X| & \text{if } X_1 X_2 = 0 \\ \infty & \text{if otherwise} \end{cases}.$$

Recall now that

$$\begin{aligned} \mathcal{O}(V, \mathbb{D}) = & \{f + g - f(0) : \\ & f \in \mathcal{O}(\mathbb{D} \times \{0\}, \mathbb{D}), g \in \mathcal{O}(\{0\} \times \mathbb{D}, \mathbb{D}), f(0) = g(0)\}. \end{aligned}$$

Then obviously  $\gamma_V(0; X) = |X_1| + |X_2|$ .

Finally, since  $z + w \in \mathcal{O}(V, \mathbb{D})$ , it follows that

$$c_V((z, 0), (0, w)) = c_V((|z|, 0), (-|w|, 0)) \geq \rho(|z|, -|w|).$$

Thus,  $c_V = k_V$ ; in particular,  $c_V = c_V^i$ .

## REFERENCES

- [1] J. B. Garnett, *Bounded analytic functions*, Pure and Appl. Math. 96, Academic Press, New York, 1981.
- [2] M. Jarnicki, P. Pflug, *Invariant distances and metrics in complex analysis*, de Gruyter Exp. Math. 9, de Gruyter, Berlin, New York, 1993.
- [3] M. Jarnicki, P. Pflug, *Invariant distances and metrics in complex analysis – revisited*, Diss. Math. 430 (2005), 1–192.
- [4] G. Kneese, *Function theory on the Neile parabola*, Preprint (2005) (arXiv:math.CV/0512473).
- [5] S. Kobayashi, *Hyperbolic complex spaces*, Grundlehren Math. Wiss. 318, Springer, Berlin, 1998.
- [6] P. Pflug, W. Zwonek *Effective formulas for invariant functions – case of elementary Reinhardt domains*, Ann. Polon. Math. 69 (1998), 175–196 (erratum: ibid., 301).

INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, ACAD. G. BONCHEV 8, 1113 SOFIA, BULGARIA

*E-mail address:* nik@math.bas.bg

CARL VON OSSIETZKY UNIVERSITÄT OLDENBURG, INSTITUT FÜR MATHE-  
MATIK, FAKULTÄT V, POSTFACH 2503, D-26111 OLDENBURG, GERMANY  
*E-mail address:* `pflug@mathematik.uni-oldenburg.de`