SPLITTING THE AUTOMORPHISM GROUP OF AN ABELIAN P-GROUP

MARIA ALICIA AVIÑO

ABSTRACT. Let G be an abelian p-group sum of finite homocyclic groups G_i . Here, we determine in which cases the automorphism group of G splits over ker σ , where σ :Aut $(G) \rightarrow \prod_i \operatorname{Aut}(G_i/pG_i)$ is the natural epimorphism.

1. Preliminaries

Throughout this paper, p is an arbitrary prime and r is a fixed ordinal number. Let $G = \bigoplus_{i \leq r} G_i$ be an abelian p-group, such that G_i is an homocyclic group of exponent p^{n_i} and finite p-rank r_i , with $n_i < n_{i+1}$ for all i. It is known that \mathcal{E} the endomorphism ring of G is isomorphic to the ring E(G)of all row finite $r \times r$ -matrices (A_{ij}) where $A_{ij} \in Hom(G_i, G_j)$. We denote by $\mathcal{A}(H)$, the automorphism group of a group H and consider $\mathcal{A}(H)$ as the group of units of the endomorphism ring E(H).

Let σ be the natural epimorphism of $\mathcal{A}(G)$ onto the product of the $\mathcal{A}(G_i/pG_i)$. We have the following exact sequence (see [2], page 256)

(1)
$$1 \to ker\sigma \to \mathcal{A}(G) \to \prod_i \mathcal{A}(G_i/pG_i) \cong \prod_i GL_{r_i}(\mathbf{Z}_p) \to 1,$$

where $GL_{r_i}(\mathbf{Z}_p)$ is the general linear group of $r_i \times r_i$ -matrices over the field \mathbf{Z}_p .

In this paper we prove a Theorem which together with Theorem 1.1 proved in [1] and [3], give a necessary and sufficient condition for the decomposition of $\mathcal{A}(G)$ as a semidirect product of ker $\sigma = \Delta(G)$ by $\Pi(G) = \prod_i GL_{r_i}(\mathbf{Z}_p)$, whenever $p \geq 5$. For the cases p = 2, 3 we give sufficient and necessary conditions for such decomposition in case $n_i + 1 < n_{i+1}$ for all *i*.

Because the p-rank of G_i is finite, $Hom(G_i, G_j) \cong p^{n_j - n_i} M_{r_i \times r_j}(\mathbf{Z}_{p^{n_i}})$ for i < j, where $M_{r_i \times r_j}(\mathbf{Z}_{p^{n_i}})$ denotes the additive group of $r_i \times r_j$ -matrices over the integers modulo p^{n_i} .

Evidently $\Delta(G) = 1 + I$ where

$$I = \{ (A_{ij})_{r \times r} \in E(G) | A_{ii} \equiv 0 \pmod{p}, \text{ for all } i \}.$$

¹⁹⁹¹ Mathematics Subject Classification. Primary: 20K10, 20K30, 20F28.

Key words and phrases. abelian p-groups, endomorphism rings, automorphism group.

If r is a natural number, then G is finite and because $GL_{r_i}(\mathbf{Z}_p)$ does not have any normal p-subgroup, $\Delta(G)$ is the maximal normal p-subgroup of $\mathcal{A}(G)$, denoted $O_p(\mathcal{A}(G))$.

Consider the exact sequences

(2i)
$$1 \to ker\lambda_i \to \mathcal{A}(G_i) \to GL_{r_i}(\mathbf{Z}_p) \to 1$$

where $GL_{r_i}(\mathbf{Z}_p)$ is the general linear group of $r_i \times r_i$ -matrices over the field \mathbf{Z}_p . It follows that $ker\lambda_i = J(\text{End } G_i) + 1_i = O_p(\mathcal{A}G_i) = \Delta(G_i)$.

By Theorem I, II.2 and III.4 in [1] and the Theorem in [3] we have the following.

Theorem 1.1. 1.1.1) For $p \ge 5$, the exact sequence (2i) splits if and only if G_i is of type either (p^{n_i}) or (p, ..., p) with $n_i > 1$.

1.1.2) For p = 3, the exact sequence (2i) splits if and only if G_i is of type $(3^{n_i}), (3^{n_i}, 3^{n_i})$ or (3, ..., 3) with $n_i > 1$.

1.1.3) For p = 2 the exact sequence (2i) splits if and only if G_i is of type $(2^{n_i}), (2^{n_i}, 2^{n_i}), (2^{n_i}, 2^{n_i}, 2^{n_i})$ or (2, ..., 2) with $n_i > 1$.

2.
$$\mathcal{A}(G)$$
 as semidirect product of $\Delta(G)$ and $\Pi(G)$

This section is devoted to the proof of Theorem 2.1.

Theorem 2.1. 2.1.1) For $p \ge 5$, the exact sequence (1) splits if and only if the exact sequences (2i) split for all i.

2.1.2) If $n_{i+1} - n_i > 1$ for all *i*, the exact sequence (1) splits if and only if the exact sequences (2*i*) split for all *i*.

This Theorem is a consequence of the following lemmas and propositions.

Proposition 2.2. If the exact sequences (2i) split for all *i*, then the exact sequence (1) splits.

Proof. Because the exact sequences (2_i) split for all i, there exists a family of morphisms $\{\theta_i : GL_{r_i}(\mathbf{Z}_p) \to \mathcal{A}(G_i)\}_r$ such that $\theta_i \lambda_i = 1_i$, where 1_i is the identity of $\mathcal{A}(G_i)$. So we have the diagram

where ι is the canonical monomorphism, p_i , δ_i are the canonical epimorphisms and $\sigma|_{\Pi}$ is the restriction of σ to $\prod_r \mathcal{A}(G_i)$. Now consider the morphisms $f_i = \delta_i \theta_i$, for all *i*.

There exists $\theta : \prod_i GL_{r_i}(\mathbb{Z}_p) \to \prod_r \mathcal{A}(G_i)$ such that $\theta p_i = f_i$ for all i, and the diagram commutes. It is clear that $\theta \sigma|_{\Pi} = 1_{\Pi}$ where 1_{Π} is the identity map of $\prod_i GL_{r_i}(\mathbb{Z}_p)$, then $\theta \sigma = 1_{\Pi}$ so the Proposition holds. In the following lemma we consider the exact sequence (1) for the group pG.

(3)
$$1 \to \Delta(pG) \to \mathcal{A}(pG) \xrightarrow{\sigma'} \Pi(pG) \to 1$$

Lemma 2.3. If the exact sequence (1) splits then the exact sequence (3) splits.

Proof. Set $\alpha : G \to pG$ such that $g\alpha = pg$. But $ker\alpha = G[p]$ is the socle of G. This morphism induces a morphism $\tau : \mathcal{A}(G) \to \mathcal{A}(pG)$. Because $\Delta(G)$ is a characteristic subgroup $\Delta(G)\tau \subset \Delta(pG)$ so we have the following exact commutative diagram

and two cases

a) If $n_1 > 1$, then $G[p] \subset pG$, so $\Pi(G) = \Pi(pG)$ and $\underline{\tau}$ is the identity. Thus if θ splits (1) then $\theta \tau$ splits (3).

b) If $n_1 = 1$, then G_1 is of type (p, p, ..., p) and $G = G_1 \oplus \overline{G}_2$ where $\overline{G}_2 = \bigoplus_{i \ge 2} G_i$. Therefore $\Pi(pG) = \Pi(\overline{G}_2)$. Thus, if ι is the canonical monomorphism of $\Pi(pG)$ to $\Pi(G)$ we have that if θ splits (1), then the morphism $\iota\theta\tau$ splits (3) and the Lemma holds.

Lemma 2.4. Set $\overline{G}_2 = \bigoplus_{i \ge 2} G_i$. If $n_1 = 1$, then (1) splits if and only if the exact sequence $1 \to \Delta(\overline{G}_2) \to \mathcal{A}(\overline{G}_2) \xrightarrow{\sigma_2} \Pi(\overline{G}_2) \to 1$ splits.

Proof. Let $g \in G$ and $g = g_1 + g_2$, $g_1 \in G_1$, $g_2 \in \overline{G}_2$. Then $\gamma : G \to \overline{G}_2$ such that $g\gamma = g_2$ is a morphism with $ker\gamma = G_1 \subseteq G[p]$ the socle of G. This morphism induces a morphism $\overline{\gamma} : \mathcal{A}(G) \to \mathcal{A}(\overline{G}_2)$.

Here ι_2 is the natural inclusion and $\overline{p}(A_{ii})_r = (A_{ii})_{i>1}$. Then the following diagram commutes

$$\begin{array}{ccc} \mathcal{A}(G) & \stackrel{\sigma}{\longrightarrow} & \Pi(G) \\ \stackrel{\iota_2 \uparrow \downarrow \overline{\gamma}}{\longleftarrow} & \stackrel{\iota_{\uparrow \downarrow} \overline{p}}{\longleftarrow} \\ \mathcal{A}(\overline{G}_2) & \stackrel{\sigma_2}{\longrightarrow} & \Pi(\overline{G}_2) \end{array}$$

and our claim holds.

Lemma 2.5. Let $n_1 = 2$ and $n_2 \ge 4$. If (1) splits then (2₁) splits.

Proof. Set $A = (A_{ij}) \in \mathcal{A}(G)$, we can define a map $\mu : \mathcal{A}(G) \to \mathcal{A}(G_1)$ so that $\mu(A) = A_{11}$. In this case, the map μ is a morphism. In fact, if $B = (B_{ij}) \in \mathcal{A}(G)$ then $(BA)_{11} = B_{11}A_{11}$ since $A_{i1} \equiv 0 \pmod{p^{n_i - n_1}}$ and $n_i - n_1 \ge 2$ for i > 1.

Now, consider the diagram

$$\begin{array}{ccc} \mathcal{A}(G) & \stackrel{\theta}{\rightarrow} & \Pi(G) \\ \downarrow \mu & \iota_1 \uparrow \downarrow \mu' \\ \mathcal{A}(G_1) & \stackrel{\lambda_1}{\rightarrow} & \Pi(G_1) \end{array}$$

where μ' is the morphism induced by μ , and ι_1 is the canonical monomorphism. If θ splits (1) then $\iota_1 \theta \mu$ splits (2₁).

Proposition 2.6. If $n_{i+1} - n_i > 1$ for all *i* and the exact sequence (1) splits then (2_i) split for all *i*.

Proof. Suppose there exists θ such that (1) splits, but there exists *i* such that (2_i) does not split. Let *t* be the least *i* such that (2_i) does not split. We have two cases

a) t > 1, so $n_t > 2$. Applying the Lemma 2.3 we can obtain that the exact sequence

$$1 \to \Delta(\underline{G}) \to \mathcal{A}(\underline{G}) \to \Pi(\underline{G}) \to 1$$

splits where $\underline{G} = p^{n_t-2}G = \bigoplus_{i \ge t} \underline{G}_i$, with $\underline{G}_i = p^{n_t-2}G_i$ for $i \ge t$. But the exponent of \underline{G}_t is p^2 and the exponent of \overline{G}_{t+1} is $\ge p^4$.

Then using Lemma 2.5 we obtain that the exact sequence

$$1 \to \Delta(\underline{G}_t) \to \mathcal{A}(\underline{G}_t) \to \Pi(\underline{G}_t) \to 1$$

splits. But this is impossible because the rank of \underline{G}_t is the same as that of G_t , that is $r_t \ge 2$ if $p \ge 5$, $r_t \ge 3$ if p = 3 and $r_t \ge 4$ if p = 2, contradicting Theorem 1.1.

b) If t = 1, but $n_1 \ge 2$, we proceed as in a). If t = 1 but $n_1 = 1$, G_1 is an elementary abelian group and by Theorem 1.1, (2_1) splits, which is not the case.

Proposition 2.7. If $p \ge 5$ and the exact sequence (1) splits then the exact sequences (2_i) split, for all *i*.

Proof. Suppose that there exists t such that (2_t) does not split. Then by Theorem 1.1, G_t is a subgroup of exponent p^{n_t} , $n_t > 1$ and p-rank $r_t > 1$. Using Lemmas 2.3 and 2.4, we may assume t = 1 and $n_t = 2$. Thus we may assume that (1) splits and (2_1) does not split with $n_1 = 2$ and $r_1 \ge 2$.

Let $\overline{A} = 1_{\Pi} + (E_{1r_1}, 0, ..., 0) \in \Pi(G)$, where $E_{1r_1} = (e_{uv}) \in GL_{r_1}(\mathbb{Z}_p)$ such that $e_{1r_1} = 1$ and $e_{uv} = 0$ for $(u, v) \neq (1, r_1)$, and $A = 1 + B = 1 + (B_{ij}) \in \mathcal{A}(G)$, where $B_{11} = E_{1r-1}$ and $B_{ij} = 0$ for $(i, j) \neq (1, 1)$.

If θ is the monomorphism such that $\theta \sigma = 1_{\Pi}$, then $\theta(\overline{A}) \equiv A \pmod{\Delta(G)}$. So $\theta(\overline{A}) = A(1 + \overline{C})$ with $1 + \overline{C} \in \Delta(G) = 1 + I$. Set $A\overline{C} = C = (c_{uv}) \in I$, then we have that $\overline{A}\theta = A + C$.

Because the order of \overline{A} is $p, 1 = (A + C)^p$. Therefore

$$1 = [1 + (B + C)]^{p} = 1 + p(B + C) + {\binom{p}{2}}(B + C)^{2} + {\binom{p}{3}}(B + C)^{3} + \dots + {\binom{p}{p-1}}(B + C)^{p-1} + (B + C)^{p}$$

Since $(B+C)^2 \in I$, we have that the block (1,1) of $(B+C)^t$ is 0 for $t \ge 4$ and for the matrices $(B+C)^2$, $(B+C)^3$ the block (1,1) is congruent to pmodulo p^2 . For $p \ge 5$, we have in the place $(1, r_1)$

$$0 \equiv p(1 + pc_{1r_1}) \pmod{p^2},$$

4

which is impossible and the proposition holds.

Corollary 2.8. 2.8.1) For $p \geq 5$, $\mathcal{A}(G)$ is a semidirect product of $\Delta(G)$ by $\Pi(G)$ if and only if $\mathcal{A}(G_i)$ is a semidirect product of $O_p(\mathcal{A}(G_i))$ by $GL_{r_i}(\mathbb{Z}_p)$.

2.8.2) For p = 3, p = 2 and $n_{i+1} - n_i > 1$, i $\mathcal{A}(G)$ is a semidirect product of $\Delta(G)$ by $\Pi(G)$ if and only if $\mathcal{A}(G_i)$ is a semidirect product of $O_p(\mathcal{A}(G_i))$ by $GL_{r_i}(\mathbb{Z}_p)$.

Corollary 2.9. $\mathcal{A}(G)$ is a semidirect product of $\Delta(G)$ by $\Pi(G)$ if and only if

2.9.1) For $p \ge 5$, G_i has rank 1 for i > 1 and G_1 either is elementary abelian or has rank 1 if $n_1 > 1$.

2.9.2) For p = 3 and $n_{i+1} - n_i > 1$, for all i, G_i has rank ≤ 2 for i > 1and G_1 either is elementary abelian or has rank ≤ 2 if $n_1 > 1$.

2.9.3) For p = 2 and $n_{i+1} - n_i > 1$ for all i, G_i has rank ≤ 3 for i > 1and G_1 either is elementary abelian or has rank ≤ 3 if $n_1 > 1$.

References

- Beletskii, P and Morgado, E. On the automorphism group of a finite abelian p-group. Ukrain. Mat. Zh. 33 No 5, 589-596.
- [2] Fuchs, L. Infinite Abelian Groups. Academic Press Vol II (1973).
- [3] Morgado, E. On the automorphism group of a finite abelian p-group. Ukrain. Mat. Zh. 32 No 5, 617-622.

(Aviño) DEPARTMENT OF MATHEMATICAL SCIENCES, NEW MEXICO STATE UNIVER-SITY,, LAS CRUCES, 88003,

E-mail address: mavino@nmsu.edu