HUREWICZ-SERRE THEOREM IN EXTENSION THEORY

M. CENCELJ, J. DYDAK, A. MITRA, AND A. VAVPETIČ

ABSTRACT. The paper is devoted to generalizations of Cencelj-Dranishnikov theorems relating extension properties of nilpotent CW complexes to its homology groups. Here are the main results of the paper:

Theorem 0.1. Suppose L is a nilpotent CW complex and F is the homotopy fiber of the inclusion i of L into its infinite symmetric product SP(L). If X is a metrizable space such that $X\tau K(H_k(L), k)$ for all $k \ge 1$, then $X\tau K(\pi_k(F), k)$ and $X\tau K(\pi_k(L), k)$ for all $k \ge 2$.

Theorem 0.2. Let X be a metrizable space such that $dim(X) < \infty$ or $X \in ANR$. Suppose L is a nilpotent CW complex and SP(L) is its infinite symmetric product. If $X\tau SP(L)$, then $X\tau L$ in the following cases:

a. $H_1(L)$ is finitely generated.

b. $H_1(L)$ is a torsion group.

Date: January 21, 2006.

¹⁹⁹¹ Mathematics Subject Classification. Primary: 54F45; Secondary: 55M10, 54C65.

Key words and phrases. Extension dimension, cohomological dimension, absolute extensor, nilpotent groups.

Supported in part by the Slovenian-USA research grant BI–US/05-06/002 and the ARRS research project No. J1–6128–0101–04. The second-named author was partially supported by Grant No.2004047 from the United States-Israel Binational Science Foundation (BSF), Jerusalem, Israel.

CONTENTS

1.	Introduction	2
2.	Properties of the homotopy fiber of $L \to SP(L)$	3
3.	Homotopy groups with coefficients	4
4.	Main results	6
5.	Appendix	8
References		9

1. INTRODUCTION

Recall that $X\tau L$ and $L \in AE(X)$ are shortcuts to the statement: L is an absolute extensor of X. The geometric result being repeatedly used in the paper is the consequence of Theorem 1.9 in [10]: if F is the homotopy fiber of $L \to K$ and $X\tau F$, then $X\tau L$ is equivalent to $X\tau K$.

Given a metrizable space X and a connected CW complex L consider the following conditions:

(1) $X\tau L$.

(2) $X \tau K(\pi_n(L), n)$ for all $n \ge 1$.

(3) $X \tau K(H_n(L), n)$ for all $n \ge 1$.

It is well-known that (1) implies (3) as proved by Dranishnikov [7] for X compact and for arbitrary X in [[12], Theorem 3.4]. The difficulty in generalizing results in cohomological dimension theory from compact spaces to arbitrary metrizable spaces usually lies in the fact that the First Bockstein Theorem does not hold for metric spaces.

By a Hurewicz-Serre Theorem in Extension Theory we mean any result showing (3) implies (2). However, in practice we are really interested in arriving at (1).

Here is the main problem we are interested in:

Problem 1.1. Suppose X is a metrizable space such that $X \tau K(H_n(L), n)$ for all $n \ge 1$. If L is nilpotent, does $X \tau K(\pi_n(L), n)$ hold for all $n \ge 1$?

Even a specialized version of 1.1 is open:

Problem 1.2. Suppose L is a nilpotent CW complex. If X is a metrizable space such that $X \tau K(H_k(L), k)$ for all $k \ge 1$, does $X \tau K(\pi_1(L), 1)$ hold?

Notice that it is not sufficient to assume $X \tau K(H_1(L), 1)$ in 1.2. Namely, take the group G from [22] whose abelianization is $\mathbb{Q} \oplus \mathbb{Q}$ and whose commutator group is \mathbb{Z}/p^{∞} . Pick a compactum X so that $\dim_{\mathbb{Q}}(X) = 1$ and $\dim_{\mathbb{Z}/p^{\infty}}(X) = 2$. The complex L = K(G, 1) is nilpotent and $X \tau K(H_1(L), 1)$ but $X \tau L$ does not hold. Indeed, as $\pi_1(L)$ is not \bar{p} -local and $H_1(L; \mathbb{Z}/p^{\infty}) = 0$, 5.4 says $H_2(L; \mathbb{Z}/p^{\infty}) \neq 0$ which means $H_2(L)/\operatorname{Tor}(H_2(L))$ is not p-divisible. If $X \tau L$, then $X \tau K(H_2(L), 2)$ and $\dim_{\mathbb{Z}_{(p)}}(X) \leq 2$ and that, in combination with $\dim_{\mathbb{Q}}(X) = 1$, implies $\dim_{\mathbb{Z}/p^{\infty}}(X) \leq 1$, a contradiction. See more in [3] about Bockstein First Theorem for nilpotent groups.

However, if $H_1(L)$ is a torsion group, then the answer to 1.2 is positive.

Lemma 1.3. Suppose N is a nilpotent group. If $X \tau K(Ab(N), 1)$ for some metrizable space X and Ab(N) is a torsion group, then $X \tau K(N, 1)$.

 $\mathbf{2}$

Proof. We will prove 1.3 by induction on the nilpotency class n of N. Let $\Gamma^n N = \Gamma^n$. Notice N/Γ^n is a nilpotent group of class (n-1) whose abelianization is an image of Ab(N). Thus $X \tau K(N/\Gamma^n, 1)$. The epimorphism

$$\otimes^n \operatorname{Ab} N \longrightarrow \Gamma^n N = \Gamma^n$$

implies $X \tau K(\Gamma^n, 1)$, so the fact that N is a central extension

$$1 \to \Gamma^n \to N \to N/\Gamma^n \to 1$$

concludes the proof.

For the sake of completeness let us show (2) is always stronger than (3).

Proposition 1.4. Suppose X is a metrizable space and L is a connected CW complex. If $X \tau K(\pi_n(L), n)$ hold for all $n \ge 1$, then $X \tau K(H_n(L), n)$ for all $n \ge 1$.

Proof. Let L_n be the CW complex obtained from L by killing all homotopy groups higher than n. Since L_n is obtained from L by attaching k-cells for k > n+1, $H_n(L_n) = H_n(L)$. Also, one has $X\tau L_n$ as $X\tau K(\pi_i(L_n), i)$ holds for all i and only finitely many homotopy groups of L_n are non-trivial (see Theorem G of [11]). Therefore $X\tau K(H_n(L_n), n)$.

Also, (1) is always stronger than (2) provided the issue of the fundamental group is avoided.

Proposition 1.5. Suppose X is a metrizable space and L is a connected CW complex. If $X \tau K(\pi_1(L), 1)$ and $X \tau L$, then $X \tau K(\pi_n(L), n)$ hold for all $n \ge 1$.

Proof. Notice that the homotopy fiber of the covering projection $\tilde{L} \to L$ is $K(\pi_1(L), 1)$. Therefore $X\tau\tilde{L}$ and $X\tau K(H_n(\tilde{L}), n)$ for all n. By Theorem F of [11] (see also 4.2 of this paper) one has $X\tau K(\pi_n(\tilde{L}), n) = K(\pi_n(L), n)$ for all $n \ge 2$.

Definition 1.6. X is called a *Knoxville space* if it is metrizable and for any connected CW complex L the conditions $X\tau K(\pi_n(L), n)$ for all $n \ge 1$ imply $X\tau L$.

Problem 1.7. Characterize Knoxville spaces.

It follows from Theorem G of [11] that any finitely dimensional X or any $X \in ANR$ is a Knoxville space. Also, it is easy to see that any countable union of closed Knoxville subspaces is a Knoxville subspace.

2. Properties of the homotopy fiber of $L \to SP(L)$

Notice that condition (3) of Section 1 is equivalent to $X\tau SP(L)$ as SP(L) is the weak product of $K(H_n(L), n)$ for all $n \ge 1$ according to the famous theorem of Dold and Thom [6]. Since we are interested in deriving $X\tau L$ it makes sense to ponder the stronger condition $X\tau F$, where F is the homotopy fiber of the inclusion $L \to SP(L)$. That is the main idea of the whole paper and in this section we concentrate on basic properties of F and its homotopy groups.

Proposition 2.1. Suppose L is a CW complex and F is the homotopy fiber of the inclusion i of L into its infinite symmetric product SP(L). If L is nilpotent, then F is nilpotent.

Proof. The homotopy sequence

$$\cdots \to \pi_n(F) \xrightarrow{j_*} \pi_n(L) \xrightarrow{i_*} \pi_n(SP(L)) \xrightarrow{\partial} \pi_{n-1}(F) \to \cdots$$

of the fibration $F \xrightarrow{j} L \xrightarrow{i} SP(L)$ is sequence of $\pi_1(L)$ -modules [17, Proposition $8^{bis}.2$]. For the action of $\pi_1(L)$ on $\pi_n(F)$ which is described in the proof of [17, Proposition $8^{bis}.2$] holds $g \cdot \alpha = j_*(g) \cdot \alpha$ for $\alpha \in \pi_n(F)$ and $g \in \pi_1(F)$.

Let I_F and I_L be the augmentation ideals of group rings $\mathbb{Z}[\pi_1(F)]$ and $\mathbb{Z}[\pi_1(L)]$, respectively. Because L is a nilpotent, there is an integer c, such that $(I_L)^c \pi_n(L) = 0$. Let $\eta \in (I_F)^c$ and $\alpha \in \pi_n(F)$. Then $j_*(\eta \alpha) = j_*(\eta)j_*(\alpha) = 0$, because $j_*(\eta) \in (I_L)^c$. Thus there exists $\beta \in \pi_{n+1}(SP(L))$, such that $\partial \beta = \eta \alpha$. Let $g \in \pi_1(F)$. Then $(j_*(g) - 1) \in I_L$ and

$$\partial((j_*(g) - 1)\beta) = (j_*(g) - 1)\partial\beta = (j_*(g) - 1)\eta\alpha = (g - 1)\eta\alpha.$$

The action of $\pi_1(L)$ on $\pi_n(SP(L))$ is defined as $l\gamma = i_*(l)\gamma$ for $l \in \pi_1(L)$ and $\gamma \in \pi_n(SP(L))$. Hence

$$(j_*(g) - 1)\beta = (i_*j_*(g) - 1)\beta = (1 - 1)\beta = 0,$$

therefore $(g-1)\eta\alpha = 0$. This shows that $(I_F)^{c+1}\pi_n(F) = 0$, so the space F is nilpotent.

Proposition 2.2. Suppose L is a nilpotent CW complex and F is the homotopy fiber of the inclusion i of L into its infinite symmetric product SP(L). If \mathcal{P} is a set of primes such that $H_k(L)$ is a \mathcal{P} -torsion group for all $k \leq n$, where $n \geq 1$ is given, then $\pi_k(F)$ is a \mathcal{P} -torsion group for all $1 \leq k \leq n+1$.

Proof. Let \mathcal{P}' be the complement of \mathcal{P} in the set of all primes. Consider the localization $L_{(\mathcal{P}')}$ of L at \mathcal{P}' . Notice that $L_{(\mathcal{P}')}$ is n-connected, so the Hurewicz homomorphism $\phi_k : \pi_k(L_{(\mathcal{P}')}) \to H_k(L_{(\mathcal{P}')})$ is an isomorphism for k = n + 1 and an epimorphism for k = n + 2. Let us split the exact sequence $\ldots \to \pi_k(F) \to \pi_k(L) \to H_k(L) \to \ldots$ into $\ldots \to \pi_2(F) \to \pi_2(L) \to H_2(L) \to A \to 0$ and $1 \to A \to \pi_1(F) \to B \to 1$, where B is the commutator subgroup of $\pi_1(L)$. Localizing the first sequence at \mathcal{P}' yields A being a \mathcal{P} -torsion group and $\pi_k(F)$ being \mathcal{P} -torsion for $2 \le k \le n + 1$. Since B is \mathcal{P} -torsion, 2.2 follows.

Corollary 2.3. Suppose L is a nilpotent CW complex and F is the homotopy fiber of the inclusion i of L into its infinite symmetric product SP(L). If n > 1 is a number such that $H_k(L)$ is a torsion group for all k < n, then for any metrizable space X the conditions $X \tau K(H_k(L), k)$ for all $k \le n$ imply $X \tau K(\pi_k(F), k)$ for all $1 \le k \le n$.

Proof. The case k = 1 is taken care of by 1.3. If *p*-torsion of $\pi_k(F)$ is not trivial, then 2.2 implies that *p*-torsion of $H_m(L)$ is not trivial for some m < k. Therefore $X\tau K(\mathbb{Z}/p^{\infty}, m)$ and $X\tau K(\mathbb{Z}/p, m+1)$. This implies $X\tau K(G, k)$ for all G in the Bockstein basis of $\pi_k(F)$ resulting in $X\tau K(\pi_k(F), k)$.

3. Homotopy groups with coefficients

Given a countable Abelian group G consider a pointed compactum $P_n(G)$ such that its integral cohomology is concentrated in dimension n and equals G. The n-th homotopy group $\pi_n(L; G)$ of a pointed CW complex L is defined in [19] to be the set $[P_n(G), L]$ of pointed homotopy classes from $P_n(G)$ to L. If $P_{n-1}(G)$ exists (that is always true if n > 2 or G is torsion free and $n \ge 2$), then $P_n(G)$ could be taken as the suspension $\Sigma P_{n-1}(G)$ of $P_{n-1}(G)$ with the resulting group structure on $\pi_n(L, G)$.

If one puts $D = P_2(G)$ (or $D = P_1(G)$ if G is torsion-free), then one can analyze homotopy groups of $L^D = \operatorname{Map}(D, L)$ and realize that $\pi_n(L^D) = \pi_{n+2}(L;G)$ (respectively, $\pi_n(L^D) = \pi_{n+1}(L;G)$). Therefore, given a Hurewicz fibration $F \to E \to B$, one concludes there is a long exact sequence $\ldots \to \pi_n(F;G) \to \pi_n(E;G) \to \pi_n(G;G) \to \pi_{n-1}(F;G) \to \ldots$ (see [19] for the special case of $G = \mathbb{Z}/m$) because $F^D \to E^D \to B^D$ is a Serre fibration.

In the special case of $G = \mathbb{Z}/m$ one can pick the Moore space $D = M(\mathbb{Z}/m, 1)$ for $P_2(G)$. In that case one has a Serre fibration (that follows from the Homotopy Extension Theorem) $\operatorname{Map}(S^2, L) \to \operatorname{Map}(D, L) \to \operatorname{Map}(S^1, L)$ where S^1 is the 1-skeleton of D and $S^2 = D/S^1$. The map $\operatorname{Map}(D, L) \to \operatorname{Map}(S^1, L)$ is simply restriction induced. Since the boundary homomorphism $\pi_{n+1}(B) \to \pi_n(F)$ in that case amounts to multiplication by m from $\pi_{n+1}(\operatorname{Map}(S^1, L)) = \pi_{n+2}(L)$ to $\pi_n(\operatorname{Map}(S^2, L)) = \pi_{n+2}(L)$, one concludes the following (see [19] for another way of deriving an equivalent result):

Lemma 3.1. Let $D = M(\mathbb{Z}/m, 1)$ for some $m \ge 2$. For each pointed CW complex L and each $n \ge 0$ one has a natural exact sequence

$$0 \to \pi_{n+2}(L) \otimes \mathbb{Z}/m \to \pi_n(L^D) \to \pi_{n+1}(L) * \mathbb{Z}/m \to 0,$$

where $\pi_1(L) * \mathbb{Z}/m$ is $\{x \in \pi_1(L) | x^m = 1\}$.

We are interested in homotopy groups with coefficients in \mathbb{Z}/p^{∞} , the direct limit of $\mathbb{Z}/p \to \mathbb{Z}/p^2 \to \ldots$ Notice that one can construct $P_2(\mathbb{Z}/p^{\infty})$ as the inverse limit of $\ldots \to M(\mathbb{Z}/p^{n+1}, 1) \to M(\mathbb{Z}/p^n, 1) \to \ldots \to M(\mathbb{Z}/p, 1)$ which can be viewed as $M(\hat{Z}_p, 1)$, the Moore space for the *p*-adic integers \hat{Z}_p in terms of Steenrod homology. In that case 3.1 becomes

Lemma 3.2. Let p be prime. For each pointed CW complex L and each $n \ge 0$ one has a natural exact sequence

 $0 \to \pi_{n+2}(L) \otimes \mathbb{Z}/p^{\infty} \to \pi_{n+2}(L; \mathbb{Z}/p^{\infty}) \to \pi_{n+1}(L) * \mathbb{Z}/p^{\infty} \to 0,$

where $\pi_1(L) * \mathbb{Z}/p^{\infty}$ is $\{x \in \pi_1(L) | x^{p^k} = 1 \text{ for some } k \ge 1\}.$

As a consequence of 3.1, 3.2, and Dold-Thom Theorem [6] $(\pi_n(SP(L) = H_n(L)))$ one can get that $\pi_n(SP(L); G) = H_n(L; G)$ for all n and $G = \mathbb{Z}/p$ or $G = \mathbb{Z}/p^{\infty}$.

Proposition 3.3. Suppose L is a nilpotent CW complex whose fundamental group is \bar{p} -local for some prime p. Let F be the homotopy fiber of the inclusion $i : L \to$ SP(L) of L into its infinite symmetric product. If $H_k(L, \mathbb{Z}/p) = 0$ for $k \leq n$, where $n \geq 1$, then $\pi_k(F, \mathbb{Z}/p) = 0$ for $2 \leq k \leq n + 1$ and $\pi_k(L, \mathbb{Z}/p) = 0$ for $2 \leq k \leq n$.

Proof. Let \tilde{L} be the universal cover of L and let $\pi : \tilde{L} \to L$ be the covering projection. Recall that every nilpotent CW complex L has the p-completion L_p with the map $L \to L_p$ inducing isomorphisms of all homology groups with coefficients in \mathbb{Z}/p such that one has a natural exact sequence

$$0 \to \operatorname{Ext}(\mathbb{Z}/p^{\infty}, \pi_n(L)) \to \pi_n(L_p) \to \operatorname{Hom}(\mathbb{Z}/p^{\infty}, \pi_{n-1}(L)) \to 0$$

for all $n \ge 1$ (see [[13], Theorem 3.7 on p.416]). By 5.2 and 5.3 one has $\operatorname{Ext}(\mathbb{Z}/p^{\infty}, \pi_1(L)) = \operatorname{Hom}(\mathbb{Z}/p^{\infty}, \pi_1(L)) = 0$, so the induced map $\hat{L}_p \to \hat{L}_p$ is a homotopy equivalence. Therefore it induces isomorphism of homology mod p and π induces isomorphism of homology mod p. However, $H_2(\tilde{L}; \mathbb{Z}/p) = \pi_2(\tilde{L}; \mathbb{Z}/p) = \pi_2(L; \mathbb{Z}/p)$ and $\pi_3(L) \to H_3(\tilde{L})$ is an epimorphism resulting in $\pi_3(L; \mathbb{Z}/p) \to H_3(L; \mathbb{Z}/p)$ being an

epimorphism. By exactness of mod p groups of a fibration one gets $\pi_2(F, \mathbb{Z}/p) = 0$. That proves 3.3 for n = 1.

If n > 1 we apply mod p Hurewicz Theorem of [19] to \tilde{L} to conclude $\pi_{n+1}(\tilde{L}; \mathbb{Z}/p) \to H_{n+1}(\tilde{L}; \mathbb{Z}/p)$ is an isomorphism and $\pi_{n+2}(\tilde{L}; \mathbb{Z}/p) \to H_{n+2}(\tilde{L}; \mathbb{Z}/p)$ is an epimorphism. Consequently, $\pi_{n+1}(L; \mathbb{Z}/p) \to H_{n+1}(L; \mathbb{Z}/p)$ is an isomorphism and $\pi_{n+2}(L; \mathbb{Z}/p) \to H_{n+2}(L; \mathbb{Z}/p) \to H_{n+2}(L; \mathbb{Z}/p)$ is an epimorphism. Thus $\pi_{n+1}(F; \mathbb{Z}/p) = 0$.

Corollary 3.4. Suppose L is a nilpotent CW complex whose fundamental group is \bar{p} -local for some prime p. Let F be the homotopy fiber of the inclusion $i: L \to SP(L)$ of L into its infinite symmetric product. If $H_k(L, \mathbb{Z}/p^{\infty}) = 0$ for $k \leq n$, where $n \geq 2$, then $\pi_k(L, \mathbb{Z}/p^{\infty}) = \pi_k(F, \mathbb{Z}/p^{\infty}) = 0$ for $2 \leq k \leq n$.

Proof. Case 1: n > 2. Notice $H_k(L, \mathbb{Z}/p) = 0$ for $k \le n - 1$ resulting in $\pi_k(F, \mathbb{Z}/p) = 0$ for $2 \le k \le n$. Hence $\pi_k(F, \mathbb{Z}/p^\infty) = 0$ for $2 \le k \le n$ and from an exact sequence we get $\pi_k(L, \mathbb{Z}/p^\infty) = 0$ for $2 \le k \le n$.

Case 2: n = 2. Let \tilde{L} be the universal cover of L and let $\pi : \tilde{L} \to L$ be the covering projection. Notice that the induced map $\hat{L}_p \to \hat{L}_p$ is a homotopy equivalence. Therefore it induces isomorphism of homology mod p and π induces isomorphism of homology mod p. However, $H_2(\tilde{L}; \mathbb{Z}/p^{\infty}) = \pi_2(L; \mathbb{Z}/p^{\infty})$ and $\pi_3(L) \to H_3(\tilde{L})$ is an epimorphism resulting in $\pi_3(L; \mathbb{Z}/p^{\infty}) \to H_3(L; \mathbb{Z}/p^{\infty})$ being an epimorphism. By exactness of mod p groups of a fibration one gets $\pi_2(F, \mathbb{Z}/p^{\infty}) = \pi_2(L, \mathbb{Z}/p^{\infty}) = 0$.

Proposition 3.5. Suppose L is a nilpotent CW complex, F is the homotopy fiber of the inclusion $i: L \to SP(L)$ of L into its infinite symmetric product, and $n \ge 2$. If $H_k(L, \mathbb{Z}_{(\mathcal{P})}) = 0$ for $k \le n$, then $\pi_k(F, \mathbb{Z}_{(\mathcal{P})}) = 0$ for $2 \le k \le n + 1$ and $\pi_k(L, \mathbb{Z}_{(\mathcal{P})}) = 0$ for $2 \le k \le n$.

Proof. Let \mathcal{P}' be the complement of \mathcal{P} in the set of all primes. If $H_k(L, \mathbb{Z}_{(\mathcal{P})}) = 0$ for $k \leq n$, then $H_1(L)$ is a \mathcal{P}' -torsion group resulting in $\pi_1(L)$ being a \mathcal{P}' -torsion group. Let $L_{(\mathcal{P})}$ be the \mathcal{P} -localization of L. It is *n*-connected, so by the classical Hurewicz Theorem $\pi_k(L_{(\mathcal{P})}) \to H_k(L_{(\mathcal{P})})$ is an isomorphism for $k \leq n + 1$ and an epimorphism for k = n + 2. That corresponds to $\pi_k(L; \mathbb{Z}_{(\mathcal{P})}) \to H_k(L; \mathbb{Z}_{(\mathcal{P})})$ being an isomorphism for $k \leq n + 1$ and an epimorphism for k = n + 2. In view of exactness of $\ldots \to \pi_k(F; \mathbb{Z}_{(\mathcal{P})}) \to \pi_k(L; \mathbb{Z}_{(\mathcal{P})}) \to H_k(L; \mathbb{Z}_{(\mathcal{P})}) \to \ldots$, 3.5 follows.

4. Main results

Lemma 4.1. Suppose X is a metrizable space, G is an Abelian group, and $n \ge 1$. If $\dim_G(X) > n$, then one of the following conditions holds:

- a. $\dim_{\mathbb{O}}(X) \ge n+1$ and G is not a torsion group.
- b. There is a prime p such that $G \otimes \mathbb{Z}/p^{\infty} \neq 0$ and $\dim_{\mathbb{Q}}(X) \leq n$, $\dim_{\mathbb{Z}/p^{\infty}}(X) \geq n$.
- c. There is a prime p such that G is p-divisible, $Tor_p(G) \neq 0$, and $\dim_{\mathbb{Z}/p^{\infty}}(X) \geq n+1$.
- d. There is a prime p such that $Tor_p(G)$ is not p-divisible and $\dim_{\mathbb{Z}/p}(X) \ge n+1$.

Proof. Let $\dim_G(X) = m$. Suppose none of (a)-(d) holds. According to Part (b) of Theorem B of [11] one has $m = \dim_{G/\operatorname{Tor}(G)}(X)$ or $m = \dim_{\operatorname{Tor}(G)}(X)$. If $\dim_{\operatorname{Tor}(G)}(X) \ge n+1$, then, according to Part (a) of Theorem B of [11], there is a

prime p such that either $\operatorname{Tor}(G)$ is p-divisible, $\operatorname{Tor}_p(G) \neq 0$, and $\dim_{\operatorname{Tor}(G)}(X) = \dim_{\mathbb{Z}/p^{\infty}}(X)$ (in which case (c) holds) or $\operatorname{Tor}(G)$ is not p-divisible, $\operatorname{Tor}_p(G) \neq 0$, and $\dim_{\operatorname{Tor}(G)}(X) = \dim_{\mathbb{Z}/p}(X)$ (in which case (d) holds). Therefore $m = \dim_{G/\operatorname{Tor}(G)}(X)$ and $\dim_{\operatorname{Tor}(G)}(X) \leq n$. In particular G is not a torsion group, so $\dim_{\mathbb{Q}}(X) \leq n$ as (a) fails to hold.

Consider $\mathcal{P} = \{p | G \otimes \mathbb{Z}/p^{\infty} \neq 0\}$, the set of primes p such that $G/\operatorname{Tor}(G)$ is not p-divisible. It is shown in [11] (Part (f) of Theorem B) that $\dim_{\mathbb{Z}(\mathcal{P})}(X) \geq \dim_{G/\operatorname{Tor}(G)}(X)$, so $\dim_{\mathbb{Z}(\mathcal{P})}(X) \geq m$. As (b) does not hold, one has $\dim_{\mathbb{Z}/p^{\infty}}(X) \leq n-1$ for all $p \in \mathcal{P}$. From the exact sequence

$$0 \to \mathbb{Z}_{(\mathcal{P})} \to \mathbb{Q} \to \bigoplus_{p \in \mathcal{P}} \mathbb{Z}/p^{\infty} \to 0$$

one concludes that the homotopy fiber of $K(\mathbb{Z}_{(\mathcal{P})}, m-1) \to K(\mathbb{Q}, m-1)$ is $K(\bigoplus_{p\in\mathcal{P}} \mathbb{Z}/p^{\infty}, m-2)$. Since $m-2 \ge n-1$, $X\tau K(\bigoplus_{p\in\mathcal{P}} \mathbb{Z}/p^{\infty}, m-2)$ which implies $X\tau K(\mathbb{Z}_{(\mathcal{P})}, m-1)$ as $X\tau K(\mathbb{Q}, m-1)$ is true. Thus $\dim_{\mathbb{Z}_{(\mathcal{P})}}(X) \le m-1$, a contradiction.

Theorem 4.2. Suppose L is a nilpotent CW complex and F is the homotopy fiber of the inclusion i of L into its infinite symmetric product SP(L). If X is a metrizable space such that $X\tau K(H_k(L), k)$ for all $k \ge 1$, then $X\tau K(\pi_k(F), k)$ and $X\tau K(\pi_k(L), k)$ for all $k \ge 2$.

Proof. Suppose $n \ge 2$ is the smallest natural number such that $X\tau K(\pi_k(F), k)$ fails (similar argument in case $X\tau K(\pi_k(L), k)$ fails). By 4.1 one of the following cases holds for $G = \pi_n(F)$:

- a. dim_{\mathbb{O}} $(X) \ge n+1$ and G is not a torsion group.
- b. There is a prime p such that $G \otimes \mathbb{Z}/p^{\infty} \neq 0$ and $\dim_{\mathbb{Q}}(X) \leq n$, $\dim_{\mathbb{Z}/p^{\infty}}(X) \geq n$.
- c. There is a prime p such that G is p-divisible, $\operatorname{Tor}_p(G) \neq 0$, and $\dim_{\mathbb{Z}/p^{\infty}}(X) \geq n+1$.
- d. There is a prime p such that $\operatorname{Tor}_p(G)$ is not p-divisible and $\dim_{\mathbb{Z}/p}(X) \ge n+1$.

Case 1: $\dim_{\mathbb{Q}}(X) \leq n-1$. Now only (b)-(d) are possible. Let p be the prime from one of those cases. Notice $H_k(L)$ is \bar{p} -local for $k \leq n-1$ as otherwise $\dim_{\mathbb{Z}/p^{\infty}}(X) \leq n-1$ and $\dim_{\mathbb{Z}/p}(X) \leq n$ so none of (b)-(d) would be valid. Another observation is $H_n(L) \otimes \mathbb{Z}/p^{\infty} = 0$. Indeed, $H_n(L) \otimes \mathbb{Z}/p^{\infty} \neq 0$ leads to $H_n(L)/\operatorname{Tor}(H_n(L))$ not being p-divisible in which case [[11], Part (d) of Theorem B] implies $\dim_{\mathbb{Z}_p}(X) \leq n$ as $\dim_{H_n(L)}(X) \leq n$. Therefore $\dim_{\mathbb{Z}_{(p)}}(X) \leq n$ (see Part (e) of Theorem B in [11]) and $\dim_{\mathbb{Z}/p^{\infty}}(X) \leq \max(\dim_{\mathbb{Q}}(X), \dim_{\mathbb{Z}_{(p)}}(X) - 1) \leq n-1$, a contradiction.

 $\pi_1(L)$ is \bar{p} -local by 5.4 and $G \otimes \mathbb{Z}/p^{\infty} = 0$ by 3.4. That means (b) is not possible. If $H_n(L)$ is not p-divisible, then $\dim_{\mathbb{Z}/p}(X) \leq n$ and neither (c) nor (d) would be possible. Thus $H_k(L;\mathbb{Z}/p) = 0$ for $k \leq n$ resulting in G being p-divisible by 3.3. That means only (c) is possible. In addition, $\operatorname{Tor}_p(H_n(L)) = 0$. Also $H_{n+1}(L) \otimes \mathbb{Z}/p^{\infty} = 0$ (otherwise $\dim_{\mathbb{Z}(p)}(X) \leq n+1$ and $\dim_{\mathbb{Z}/p^{\infty}}(X) \leq \max(\dim_{\mathbb{Q}}(X), \dim_{\mathbb{Z}(p)}(X) - 1) \leq n$). Thus $H_k(L;\mathbb{Z}/p^{\infty}) = 0$ for $k \leq n+1$. By 3.4 $\operatorname{Tor}_p(G) = 0$, a contradiction.

Case 2: $\dim_{\mathbb{Q}}(X) > n-1$. By 2.2 the group G is \mathcal{P} -torsion such that $\dim_{\mathbb{Z}/p^{\infty}}(X) \leq n-1$ for all $p \in \mathcal{P}$ which implies $\dim_G(X) \leq n$, a contradiction.

Corollary 4.3. Suppose L is a nilpotent CW complex such that $\pi_n(L) = \pi_{n+1}(L) = 0$ for some $n \ge 1$. If $X \tau SP(L)$ for some metrizable space X, then $X \tau K(H_{n+1}(L), n)$.

Proof. If n = 1, then $H_{n+1}(L) = 0$, so assume $n \ge 2$. Notice that $\pi_n(F) = H_{n+1}(L)$, where F is the homotopy fiber of $i: L \to SP(L)$.

Lemma 4.4. Suppose L is a nilpotent CW complex and F is the homotopy fiber of the inclusion i of L into its infinite symmetric product SP(L). If $X\tau K(H_1(L), 1)$ for some metrizable space X and $H_1(L)$ is finitely generated, then $X\tau K(\pi_1(F), 1)$

Proof. If $H_1(L)$ is finitely generated and non-torsion, then X is at most 1dimensional in which case $X \tau L$ for all connected CW complexes. Therefore assume $H_1(L)$ is a torsion group and (see 2.2) there is an exact sequence $1 \to A \to \pi_1(F) \to B \to 1$ such that A and B are \mathcal{P} -torsion groups, where $\mathcal{P} = \{p | \text{Tor}_p(H_1(L)) \neq 0\}$. Notice $H_1(L)$ does not contain \mathbb{Z}/p^{∞} for any prime p, so $X \tau K(A, 1)$ and $X \tau K(B, 1)$ which implies $X \tau K(\pi_1(F), 1)$.

Theorem 4.5. Let X be a metrizable space such that $dim(X) < \infty$ or $X \in ANR$. Suppose L is a nilpotent CW complex and SP(L) is its infinite symmetric product. If $X\tau SP(L)$, then $X\tau L$ in the following cases:

- a. $H_1(L)$ is finitely generated.
- b. $H_1(L)$ is a torsion group.

Proof. a. By 4.2 and 4.4 one concludes $X\tau K(\pi_n(F), n)$ for all $n \ge 1$. Theorem G of [11] gives $X\tau F$ which implies $X\tau L$.

b. By 4.2 and 1.3 one concludes $X\tau K(\pi_n(L), n)$ for all $n \ge 1$. Theorem G of [11] yields $X\tau L$.

5. Appendix

Lemma 5.1. Suppose N is a nilpotent group and p is a prime. Ab(N) is p-divisible if and only if N is p-divisible.

Proof. If N is p-divisible clearly so is its abelianization. We will prove the converse by induction on the nilpotency class n of N. Let $\Gamma^n N = \Gamma^n$. Notice N/Γ^n is a nilpotent group of class (n-1) whose abelianization is p-divisible. Thus it is p-divisible. The epimorphism

$$\otimes^n \operatorname{Ab} N \longrightarrow \Gamma^n N = \Gamma^n$$

implies Γ^n is *p*-divisible, so the fact that N is a central extension

$$1 \to \Gamma^n \to N \to N/\Gamma^n \to 1$$

concludes the proof.

Lemma 5.2. Suppose N is a nilpotent group and p is a prime. The following conditions are equivalent:

- a. $Ext(\mathbb{Z}/p^{\infty}, N) = 0$,
- b. $Ext(\mathbb{Z}/p^{\infty}, N)$ is p-divisible,
- c. N is p-divisible.

Proof. (a) \Longrightarrow (b) is obvious. For (c) \Longrightarrow (a) let N be p-divisible. Then so is its abelianization and Proposition 3 of [4] implies E = 0.

(b) \implies (c) If N is not p-divisible neither is its abelianization by 5.1. Therefore, by Proposition 3 of [4] Ext($\mathbb{Z}_{p^{\infty}}$, AbN) is not p-divisible. Then the six-term exact

sequence of Hom and Ext [[2], p.170] implies that $\text{Ext}(\mathbb{Z}/p^{\infty}, N)$ is not p-divisible.

Lemma 5.3. Suppose G is a nilpotent group and p is a prime. The following conditions are equivalent:

- a. $Hom(\mathbb{Z}/p^{\infty}, G) = 0$,
- b. $Hom(\mathbb{Z}/p^{\infty}, G)$ is p-divisible,
- c. $Hom(\mathbb{Z}/p^{\infty}, G) \otimes \mathbb{Z}/p^{\infty} = 0$
- d. G does not contain \mathbb{Z}/p^{∞} .

Proof. Note that albeit Bousfield and Kan [2] defined *Hom* as a space, they showed that it is also the set of the respective homomorphisms.

(a) \Longrightarrow (b) and (b) \Longrightarrow (c) are obvious.

(c) \implies (b). Notice the *p*-torsion of $\operatorname{Hom}(\mathbb{Z}/p^{\infty}, G)$ is trivial. Indeed, if $i : \mathbb{Z}/p^{\infty} \to G$ and $i^p = 1$, then for any $a \in \mathbb{Z}/p^{\infty}$ we find $b \in \mathbb{Z}/p^{\infty}$ satisfying $b^p = a$. Now, $i(a) = i(b^p) = i^p(b) = 1$. If an Abelian group A has no *p*-torsion and $A \otimes \mathbb{Z}/p^{\infty} = 0$, then A is *p*-divisible.

(b) \implies (d) Suppose $i : \mathbb{Z}/p^{\infty} \to G$ is a monomorphism. Given $a \in \mathbb{Z}/p^{\infty}$ find $k \ge 1$ such that $a^{p^k} = 1$ and choose $\phi : \mathbb{Z}/p^{\infty} \to G$ so that $i = \phi^{p^k}$. Now $i(a) = \phi^{p^k}(a) = (\phi(a))^{p^k} = \phi(a^{p^k}) = \phi(1) = 1$, a contradiction.

(d) \Longrightarrow (a). Given a non-trivial $i : \mathbb{Z}/p^{\infty} \to G$ its image is a direct sum of copies of \mathbb{Z}/p^{∞} , a contradiction.

Lemma 5.4. Suppose L is a nilpotent CW complex and p is a prime. If $H_1(L; \mathbb{Z}/p^{\infty}) = H_2(L; \mathbb{Z}/p^{\infty}) = 0$, then $\pi_1(L)$ is \bar{p} -local.

Proof. In view of $H_2(L; \mathbb{Z}/p^{\infty}) = 0$, $H_1(L)$ has trivial *p*-torsion and $H_1(L; \mathbb{Z}/p^{\infty}) = 0$ implies $H_1(L)$ is *p*-divisible. So is $\pi_1(L)$ (see 5.1). Consider the *p*-completion \hat{L}_p of *L*. As $\pi_1(\hat{L}_p) = \text{Ext}(\mathbb{Z}/p^{\infty}, \pi_1(L)) = 0$ and $H_2(\hat{L}_p; \mathbb{Z}/p^{\infty}) = H_2(L; \mathbb{Z}/p^{\infty}) = 0$ one gets $\pi_2(\hat{L}_p) \otimes \mathbb{Z}/p^{\infty} = 0$ by the Hurewicz Theorem. The exact sequence

$$0 \to \operatorname{Ext}(\mathbb{Z}/p^{\infty}, \pi_2(L)) \to \pi_2(L_p) \to \operatorname{Hom}(\mathbb{Z}/p^{\infty}, \pi_1(L)) \to 0$$

implies $\operatorname{Hom}(\mathbb{Z}/p^{\infty}, \pi_1(L)) \otimes \mathbb{Z}/p^{\infty} = 0$. By 5.3 $\pi_1(L)$ is \bar{p} -local.

References

- A.K.Bousfield, Localization and periodicity in unstable homotopy theory, J. Amer. Math. Soc. 7 (1994), no. 4, 831–873.
- [2] A.K.Bousfield and D.M.Kan, Homotopy limits completions and localizations, Springer Lecture Notes in Math., Vol.304 (2nd corrected printing), Springer-Verlag, Berlin-Heidelberg-New York, 1987.
- [3] M.Cencelj, J.Dydak, A.Mitra, and A.Vavpetič, Bockstein Theorem for nilpotent groups, preprint
- M.Cencelj and A.N.Dranishnikov, Extension of maps to nilpotent spaces II, Topology Appl. 124 (2002), no. 1, 77–83.
- [5] M.Cencelj and A.N.Dranishnikov, Extension of maps to nilpotent spaces III, Topology Appl. 153 (2005) 208-212.
- [6] A.Dold and R.Thom, Quasifaserungen und Unendliche Symmetrische Produkte, Annals of Math. 67 (1958),239–281.
- [7] A.N.Dranishnikov, Extension of mappings into CW complexes, Mat. USSR Sbornik, 74 (1993), 47–56.
- [8] A.N.Dranishnikov, Cohomological Dimension Theory of Compact Metric Spaces, Topology Atlas (1999).

- [9] A. N. Dranishnikov and J. Dydak, Extension dimension and extension types, Proc. Steklov Math. Inst. 212 (1996), 55–88.
- [10] A.Dranishnikov and J.Dydak, Extension theory of separable metrizable spaces with applications to dimension theory, Transactions of the American Math.Soc. 353 (2000), 133–156.
- J.Dydak, Cohomological dimension and metrizable spaces, Transactions of the Amer.Math.Soc. 337 (1993),219–234.
- [12] J.Dydak, Cohomological dimension and metrizable spaces, Transactions of the Amer.Math.Soc. 348 (1996), 1647–1661.
- [13] P.G.Goerss and J.F.Jardine, Simplicial Homotopy Theory, Birkhauser, Basel-Boston-Berlin, 1999.
- [14] A.Hatcher, Algebraic Topology, Cambridge University Press, Cambridge, 2002.
- [15] P. Hilton, G. Mislin, J. Roitberg, *Localization of Nilpotent groups and spaces*, North-Holland Publishing Co. , Amsterdam-Oxford; American Elsevier Publishing Co. , Inc. , New York, 1975.
- [16] W.I.Kuzminov, Homological dimension theory, Russian Math. Surveys 23 (1968), 1–45.
- [17] J. MacCleary, A User's Guide to Spectral Sequence, Cambridge studies in advanced mathematics, second edition, 2
- [18] S.Mardešić and J.Segal, Shape theory, North-Holland Publ.Co. (1982).
- [19] J.A.Neisendorfer, Primary homotopy theory, Memoirs.Amer.Math.Soc 232 (1980).
- [20] D. J. S. Robinson, A course in the theory of groups, Springer-Verlag, New York, 1993.
- [21] D.Sullivan, Genetics of homotopy theory and the Adams conjecture, Ann. of Math. (2) 100 (1974), 1–79.
- [22] R.B.Warfield, Nilpotent groups, Springer Lecture Notes in Math., Vol.513, Springer-Verlag, Berlin-Heidelberg-New York, 1976.

Fakulteta za Matematiko in Fiziko, Univerza v Ljubljani, Jadranska ulica 19, SI-1111 Ljubljana, Slovenija

E-mail address: matija.cencelj@guest.arnes.si

UNIVERSITY OF TENNESSEE, KNOXVILLE, TN 37996, USA *E-mail address*: dydak@math.utk.edu

UNIVERSITY OF TENNESSEE, KNOXVILLE, TN 37996, USA *E-mail address*: ajmitra@math.utk.edu

Fakulteta za Matematiko in Fiziko, Univerza v Ljubljani, Jadranska ulica 19, SI-1111 Ljubljana, Slovenija

E-mail address: ales.vavpetic@fmf.uni-lj.si