

Bethe Eigenvectors of Higher Transfer Matrices

E. Mukhin^{*}, V. Tarasov^{*}, and A. Varchenko[◇]

^{*,*}*Department of Mathematical Sciences
Indiana University Purdue University at Indianapolis
Indianapolis, IN 46202, USA*

^{*}*St. Petersburg Branch of Steklov Mathematical Institute
Fontanka 27, St. Petersburg 191023, Russia*

[◇]*Department of Mathematics, University of North Carolina
Chapel Hill, NC 27599, USA*

April 2006

Abstract. We consider the XXX-type and Gaudin quantum integrable models associated with the Lie algebra \mathfrak{gl}_N . The models are defined on a tensor product $M_1 \otimes \dots \otimes M_n$ of irreducible \mathfrak{gl}_N -modules. For each model, there exist N one-parameter families of commuting operators on $M_1 \otimes \dots \otimes M_n$, called the transfer matrices. We show that the Bethe vectors for these models, given by the algebraic nested Bethe ansatz, are eigenvectors of higher transfer matrices and compute the corresponding eigenvalues.

1. Introduction

In this paper we consider quantum integrable models associated with the Lie algebra \mathfrak{gl}_N , the XXX-type and Gaudin models. The models are defined on a tensor product $M_1 \otimes \dots \otimes M_n$ of irreducible \mathfrak{gl}_N -modules. For each model, there exist N one-parameter families of mutually commuting operators on $M_1 \otimes \dots \otimes M_n$, called the transfer matrices. In this paper the transfer matrices for the XXX-type model are denoted by $\mathcal{T}_{1,Q}^{M_1 \otimes \dots \otimes M_n}(u; z), \dots, \mathcal{T}_{N,Q}^{M_1 \otimes \dots \otimes M_n}(u; z)$, see Section 6, and the transfer matrices for the Gaudin model are denoted by $\mathcal{G}_{1,K}^{M_1 \otimes \dots \otimes M_n}(u; z), \dots, \mathcal{G}_{N,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$, see Section 9.

The important problem is to find eigenvectors and eigenvalues of the transfer matrices. If M_1, \dots, M_n are highest weight modules, this can be done by the algebraic Bethe ansatz method. The $N = 2$ case is fairly well understood, for example, see [G], [KBI]. However, the $N > 2$ case is much more technically involved and less studied.

^{*}*E-mail: mukhin@math.iupui.edu*

^{*}Supported in part by RFFI grant 05-01-00922
E-mail: vt@math.iupui.edu, vt@pdmi.ras.ru

[◇]Supported in part by NSF grant DMS-0244579
E-mail: anv@email.unc.edu

The purpose of this paper is to obtain eigenvectors and eigenvalues of the transfer matrices for $N > 2$ using the algebraic nested Bethe ansatz method. For the transfer matrix $\mathcal{T}_{1,Q}^{M_1 \otimes \dots \otimes M_n}(u; z)$ this was done in [KR1]. The transfer matrix $\mathcal{T}_{N,Q}^{M_1 \otimes \dots \otimes M_n}(u; z)$ comes from the central elements of the Yangian $Y(\mathfrak{gl}_N)$ and is proportional to the identity operator on the whole tensor product $M_1 \otimes \dots \otimes M_n$ [KS]. In this paper we extend the technique developed in [KR1] to show that the Bethe vectors are eigenvectors of the higher transfer matrices $\mathcal{T}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u)$, $k > 1$, and to compute the eigenvalues, see Theorems 5.4 and 6.1. The formula for eigenvalues of the higher transfer matrices on Bethe vectors were conjectured in [KR2]. However, no proof can be found in literature.

For the Gaudin model, the transfer matrix $\mathcal{G}_{1,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$ comes from the central elements of the current algebra $\mathfrak{gl}_N[x]$ and is proportional to the identity operator on the whole tensor product $M_1 \otimes \dots \otimes M_n$. The transfer matrix $\mathcal{G}_{2,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$ is the generating function of the famous Gaudin Hamiltonians which also appear as the operators in the Knizhnik-Zamolodchikov equations. Eigenvectors and eigenvalues of $\mathcal{G}_{2,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$ can be obtained from those of $\mathcal{T}_{1,Q}^{M_1 \otimes \dots \otimes M_n}(u; z)$ by taking a suitable limit. For more straightforward approach see [Ju]. For a construction of eigenvectors and eigenvalues of the Gaudin Hamiltonians as a quasiclassical limit of solutions to the Knizhnik-Zamolodchikov equations see [RV].

The transfer matrices for the XXX-type model were constructed in [KS] more than 20 years ago. Suprisingly, an explicit form of the higher transfer matrices for the Gaudin model became known only recently. In this paper we use the approach suggested in [T] that gives the transfer matrices of the Gaudin model as a suitable limit of proper linear combinations of the transfer matrices of the XXX-type model. We trace this limit in the proof of the Theorem 5.4. This allows us to show that the Bethe vectors of the Gaudin model are eigenvectors of the higher transfer matrices and to compute the eigenvalues; see Theorems 8.6 and 9.2.

There is another approach to higher transfer matrices of the Gaudin model and the Bethe ansatz, based on the representation theory of affine Lie algebras at the critical level [FFR]. It is interesting and important to identify the transfer matrices $\mathcal{G}_{1,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$, $\dots, \mathcal{G}_{N,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$ among the higher Gaudin Hamiltonians constructed in [FFR].

Following [T], it is convenient to combine the transfer matrices of the XXX-type model into a single difference operator

$$\mathfrak{D}_{N,Q}^{M_1 \otimes \dots \otimes M_n}(u; \partial_u; z) = 1 - \mathcal{T}_{1,Q}^{M_1 \otimes \dots \otimes M_n}(u; z) e^{-\partial_u} + \dots + (-1)^N \mathcal{T}_{N,Q}^{M_1 \otimes \dots \otimes M_n}(u; z) e^{-N\partial_u},$$

where $\partial_u = d/du$, cf. (A.1), (4.16). The operator $\mathfrak{D}_{N,Q}^{M_1 \otimes \dots \otimes M_n}(u; \partial_u; z)$ acts on functions of u with values in $M_1 \otimes \dots \otimes M_n$. Then any common eigenvector $v \in M_1 \otimes \dots \otimes M_n$ of the transfer matrices,

$$\mathcal{T}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u; z) v = \mathcal{A}_k^v(u) v, \quad k = 1, \dots, N,$$

where $\mathcal{A}_1^v(u), \dots, \mathcal{A}_N^v(u)$ are suitable scalar functions, defines the scalar difference operator

$$\mathfrak{M}^v(u; \partial_u) = 1 - \mathcal{A}_1^v(u) e^{-\partial_u} + \dots + (-1)^N \mathcal{A}_N^v(u) e^{-N\partial_u}.$$

If $v = \mathbb{B}_\xi^{v_1 \otimes \dots \otimes v_n}(t; z)$ is the Bethe vector of the XXX-type model, cf. (6.2), (5.2), then the operator $\mathfrak{M}^v(u; \partial_u)$ coincides with the operator $\mathfrak{M}_{\xi, Q}(u, \partial_u; t; z; \Lambda)$, cf. (6.6), which is written as a product,

$$(1.1) \quad \mathfrak{M}_{\xi, Q}(u, \partial_u; t; z; \Lambda) = (1 - \mathcal{X}_{\xi, Q}^1(u; t; z; \Lambda) e^{-\partial_u}) \dots (1 - \mathcal{X}_{\xi, Q}^N(u; t; z; \Lambda) e^{-\partial_u}),$$

where the functions $\mathcal{X}_{\xi, Q}^a(u; t; z; \Lambda)$, see (6.4), are explicitly defined in terms of the solution t of the Bethe ansatz equations (6.3). This gives a formula for the corresponding eigenvalues $\mathcal{A}_1^v(u; z), \dots, \mathcal{A}_N^v(u)$.

It is also convenient to combine the transfer matrices of the Gaudin model into a single differential operator

$$\mathcal{D}_K^{M_1 \otimes \dots \otimes M_n}(u; \partial_u; z) = \partial_u^N - \mathcal{G}_{1, K}^{M_1 \otimes \dots \otimes M_n}(u; z) \partial_u^{N-1} + \dots + (-1)^N \mathcal{G}_{N, K}^{M_1 \otimes \dots \otimes M_n}(u; z),$$

cf. (A.2), (8.3). Then any common eigenvector $v \in M_1 \otimes \dots \otimes M_n$ of the transfer matrices,

$$\mathcal{G}_{k, K}^{M_1 \otimes \dots \otimes M_n}(u; z) v = \mathcal{Z}_k^v(u) v, \quad k = 1, \dots, N,$$

where $\mathcal{Z}_1^v(u), \dots, \mathcal{Z}_N^v(u)$ are suitable scalar functions, defines the scalar differential operator

$$\mathcal{M}^v(u; \partial_u) = \partial_u^N - \mathcal{Z}_1^v(u) \partial_u^{N-1} + \dots + (-1)^N \mathcal{Z}_N^v(u).$$

If $v = \mathbb{F}_\xi^{v_1 \otimes \dots \otimes v_n}(t; z)$ is the Bethe vector of the Gaudin model, cf. (9.6), (8.7), then the operator $\mathcal{M}^v(u; \partial_u)$ coincides with the operator $\mathcal{M}_{\xi, K}(u, \partial_u; t; z; \Lambda)$, cf. (9.4), which is written as a product,

$$(1.2) \quad \mathcal{M}_{\xi, K}(u, \partial_u; t; z; \Lambda) = (\partial_u - \mathcal{Y}^1(u; t; z; \Lambda)) \dots (\partial_u - \mathcal{Y}^N(u; t; z; \Lambda)),$$

where the functions $\mathcal{Y}^a(u; t; z; \Lambda)$ are explicitly defined in terms of the solution t of the Bethe ansatz equations (9.3). This gives a formula for the eigenvalues $\mathcal{Z}_1^v(u), \dots, \mathcal{Z}_N^v(u)$.

The Gaudin model can be obtained from the XXX-type model in the limit as u, z tend to infinity. In this limit, the difference operators $\mathfrak{D}_{N, Q}^{M_1 \otimes \dots \otimes M_n}(u; \partial_u; z)$ and $\mathfrak{M}_{\xi, Q}(u, \partial_u; t; z; \Lambda)$ turn into the differential operators $\mathcal{D}_K^{M_1 \otimes \dots \otimes M_n}(u; \partial_u; z)$ and $\mathcal{M}_{\xi, K}(u, \partial_u; t; z; \Lambda)$,

$$\mathfrak{D}_{N, Q}^{M_1 \otimes \dots \otimes M_n}(\varepsilon^{-1}u, \varepsilon \partial_u; \varepsilon^{-1}z) = \varepsilon^N \mathcal{D}_K^{M_1 \otimes \dots \otimes M_n}(u; \partial_u; z) + O(\varepsilon^{N+1}),$$

$$\mathfrak{M}_{N, Q}^{M_1 \otimes \dots \otimes M_n}(\varepsilon^{-1}u, \varepsilon \partial_u; \varepsilon^{-1}t; \varepsilon^{-1}z; \Lambda) = \varepsilon^N \mathcal{M}_K^{M_1 \otimes \dots \otimes M_n}(u; \partial_u; t; z; \Lambda) + O(\varepsilon^{N+1}),$$

as $Q = 1 + \varepsilon K$ and $\varepsilon \rightarrow 0$. Moreover, the Bethe ansatz equations (6.3) of the XXX-type model become in this limit the Bethe ansatz equations (9.3) of the Gaudin model, and the Bethe vector $\mathbb{B}_\xi^{v_1 \otimes \dots \otimes v_n}(t; z)$ turns into the Bethe vector $\mathbb{F}_\xi^{v_1 \otimes \dots \otimes v_n}(t; z)$, cf. Theorem A.1.

The operators (1.1) and (1.2) of the XXX-type and Gaudin models appeared in [MV2] and [MV1], respectively. Those operators were called in [MV1], [MV2] the fundamental

operators of the corresponding solutions of the Bethe ansatz equations. It was shown therein that for finite-dimensional \mathfrak{gl}_N -modules M_1, \dots, M_n , the kernel of a fundamental operator is generated by polynomials only and the singular points of the fundamental operator were described. These facts and Schubert calculus imply that the number of solutions of the corresponding Bethe ansatz equations (properly counted) does not exceed the dimension of $M_1 \otimes \dots \otimes M_n$ [MV1], [MV2]. That was a step to the proof of the completeness of the Bethe ansatz conjecture, which says that the Bethe vectors form a basis of $M_1 \otimes \dots \otimes M_n$ under certain conditions.

An important observation for applications is the fact that the transfer matrices are symmetric with respect to a suitable symmetric bilinear form on $M_1 \otimes \dots \otimes M_n$, see Theorems 9.1 and 7.6. Under certain assumptions, this fact allows us to conclude that the eigenvalues of the transfer matrices take real values for real u . In particular, this implies that the polynomials generating the kernel of the fundamental operator of a solution of the Bethe ansatz equations can be chosen to have real coefficients only, see applications to real algebraic geometry in [MTV].

2. Basic notation

We will be using the standard superscript notation for embeddings of tensor factors into tensor products. If $\mathcal{A}_1, \dots, \mathcal{A}_k$ are unital associative algebras, and $a \in \mathcal{A}_i$, then

$$a^{(i)} = 1^{\otimes(i-1)} \otimes a \otimes 1^{\otimes(k-i)} \in \mathcal{A}_1 \otimes \dots \otimes \mathcal{A}_k.$$

If $a \in \mathcal{A}_i$ and $b \in \mathcal{A}_j$, then $(a \otimes b)^{(ij)} = a^{(i)} b^{(j)}$, etc.

Example. Let $k = 2$. Let $\mathcal{A}_1, \mathcal{A}_2$ be two copies of the same algebra \mathcal{A} . Then for any $a, b \in \mathcal{A}$ we have $a^{(1)} = a \otimes 1$, $b^{(2)} = 1 \otimes b$, $(a \otimes b)^{(12)} = a \otimes b$ and $(a \otimes b)^{(21)} = b \otimes a$.

Fix a positive integer N . All over the paper we use the convention $V = \mathbb{C}^N$, and we identify elements of $\text{End}(V)$ with $N \times N$ matrices using the standard basis of \mathbb{C}^N .

Let e_{ab} , $a, b = 1, \dots, N$, be the standard generators of the Lie algebra \mathfrak{gl}_N :

$$[e_{ab}, e_{cd}] = \delta_{bc} e_{ad} - \delta_{ad} e_{cb}.$$

A vector v in a \mathfrak{gl}_N -module is a weight vector of weight $(\Lambda^1, \dots, \Lambda^N)$ if $e_{aa} v = \Lambda^a v$ for any $a = 1, \dots, N$. A vector v is called a *singular vector* if $e_{ab} v = 0$ for any $1 \leq a < b \leq N$.

Let $\mathbb{A}_{\{k\}} \in \text{End}(V^{\otimes k})$ be the skew-symmetrization projector:

$$\mathbb{A}_{\{k\}} v_1 \otimes \dots \otimes v_k = \frac{1}{k!} \sum_{\sigma \in S_k} \text{sign}(\sigma) v_{\sigma_1} \otimes \dots \otimes v_{\sigma_k}.$$

We denote by $V^{\wedge k}$ the image of $\mathbb{A}_{\{k\}}$, and for any $Q \in \text{End}(V)$ set $Q^{\wedge k} = Q^{\otimes k}|_{V^{\wedge k}}$. In particular, $Q^{\wedge N} = \det Q$.

The space V is considered as a \mathfrak{gl}_N -module with the natural action: $e_{ab} \mapsto E_{ab}$, where $E_{ab} \in \text{End}(V)$ is the matrix with the only nonzero entry equal to 1 at the intersection of the a -th row and b -th column. The \mathfrak{gl}_N -module V is called the *vector representation*. The space $V^{\wedge k}$ is a \mathfrak{gl}_N -module with the action $e_{ab} \mapsto (E_{ab}^{(1)} + \dots + E_{ab}^{(k)})|_{V^{\wedge k}}$.

We will use the following notation for products of noncommuting factors:

$$\begin{aligned} \overrightarrow{\prod}_{1 \leq i \leq k} X_i &= X_1 \dots X_k, & \overleftarrow{\prod}_{1 \leq i \leq k} X_i &= X_k \dots X_1, \\ \overrightarrow{\prod}_{1 \leq i < j \leq k} &= \overrightarrow{\prod}_{1 \leq i \leq k} \overrightarrow{\prod}_{i < j \leq k} & \overleftarrow{\prod}_{1 \leq i < j \leq k} &= \overleftarrow{\prod}_{1 \leq j \leq k} \overleftarrow{\prod}_{1 \leq i < j} \end{aligned}$$

3. R -matrices

Let $P = \sum_{a,b=1}^N E_{ab} \otimes E_{ba}$. It is the flip map: $P(x \otimes y) = y \otimes x$ for any $x, y \in V$, and $A^{(21)} = PAP$ for any $A \in \text{End}(V^{\otimes 2})$.

The *rational R -matrix* is $R(u) = u + P \in \text{End}(V^{\otimes 2})$. It satisfies the inversion relation

$$(3.1) \quad R(u) R^{(21)}(-u) = 1 - u^2$$

and the Yang-Baxter equation

$$(3.2) \quad R^{(12)}(u-v) R^{(13)}(u) R^{(23)}(v) = R^{(23)}(v) R^{(13)}(u) R^{(12)}(u-v).$$

Remark. The R -matrix $R(u)$ is symmetric: $R(u) = R^{(21)}(u)$. However, we will not use this property and will write all relations, for instance (3.1), in the form that naturally extends to more general R -matrices.

Below we use the well known *fusion procedure* to define R -matrices $R^{\wedge k, \wedge l}(u)$ acting in the tensor products $V^{\wedge k} \otimes V^{\wedge l}$, and describe their properties. The proofs can be found in [MNO, Sections 1, 2], [N, Sections 2, 3]. Notice that our notation differs from those in [MNO], [N], instead of $R(u)$ the R -matrix $1 - P/u$ is used therein.

Lemma 3.1.
$$\overrightarrow{\prod}_{1 \leq i < j \leq k} R^{(ij)}(i-j) = \overleftarrow{\prod}_{1 \leq i < j \leq k} R^{(ij)}(i-j) = \mathbb{A}_{\{k\}} (-1)^k \prod_{j=1}^k (-j)^{k-j+1}.$$

The Yang-Baxter equation (3.2) and Lemma 3.1 imply that

$$(3.3) \quad \mathbb{A}_{\{k\}}^{(1\dots k)} \mathbb{A}_{\{l\}}^{(k+1\dots k+l)} \overrightarrow{\prod}_{1 \leq i \leq k} \overleftarrow{\prod}_{1 \leq j \leq l} R^{(i,j+k)}(u+i-j-k+l) = \\ = \left(\overleftarrow{\prod}_{1 \leq i \leq k} \overrightarrow{\prod}_{1 \leq j \leq l} R^{(i,j+k)}(u+i-j-k+l) \right) \mathbb{A}_{\{k\}}^{(1\dots k)} \mathbb{A}_{\{l\}}^{(k+1\dots k+l)}.$$

Set

$$R^{\wedge k, \wedge l}(u) = \overleftarrow{\prod}_{1 \leq i \leq k} \overrightarrow{\prod}_{1 \leq j \leq l} R^{(i,j+k)}(u+i-j-k+l)|_{V^{\wedge k} \otimes V^{\wedge l}} \in \text{End}(V^{\wedge k}) \otimes \text{End}(V^{\wedge l}).$$

The introduced R -matrices satisfy the inversion relation

$$R^{\wedge k, \wedge l}(u) (R^{\wedge l, \wedge k}(-u))^{(21)} = \prod_{i=1}^k \prod_{j=1}^l (1 - (u - i + j)^2),$$

and the Yang-Baxter equation

$$\begin{aligned} (R^{\wedge k, \wedge l}(u - v))^{(12)} (R^{\wedge k, \wedge m}(u))^{(13)} (R^{\wedge l, \wedge m}(v))^{(23)} = \\ = (R^{\wedge l, \wedge m}(v))^{(23)} (R^{\wedge k, \wedge m}(u))^{(13)} (R^{\wedge k, \wedge l}(u - v))^{(12)}. \end{aligned}$$

In addition, for any $Q \in \text{End}(V)$,

$$(3.4) \quad [R^{\wedge k, \wedge l}(u), Q^{\wedge k} \otimes Q^{\wedge l}] = 0.$$

Set

$$R_{\wedge k, \wedge 1}(u) = u + \sum_{a,b=1}^N (E_{ab}^{(1)} + \dots + E_{ab}^{(k)})|_{V^{\wedge k}} \otimes E_{ba} \in \text{End}(V^{\wedge k}) \otimes \text{End}(V),$$

and

$$R_{\wedge 1, \wedge k}(u) = u + k - 1 + \sum_{a,b=1}^N E_{ab} \otimes (E_{ba}^{(1)} + \dots + E_{ba}^{(k)})|_{V^{\wedge k}} \in \text{End}(V) \otimes \text{End}(V^{\wedge k}).$$

In particular, $R_{\wedge N, \wedge 1}(u) = u + 1$ and $R_{\wedge 1, \wedge N}(u) = u + N$.

Lemma 3.2. $R^{\wedge k, \wedge 1}(u) = R_{\wedge k, \wedge 1}(u) \prod_{i=1}^{k-1} (u - i), \quad R^{\wedge 1, \wedge k}(u) = R_{\wedge 1, \wedge k}(u) \prod_{i=0}^{k-2} (u + i).$

Corollary 3.3. $R_{\wedge k, \wedge 1}(u) (R_{\wedge 1, \wedge k}(-u))^{(21)} = (u + 1)(k - u).$

4. Yangian $Y(\mathfrak{gl}_N)$

In this section we collect some known facts from the representation theory of the Yangian $Y(\mathfrak{gl}_N)$ that will be used in the paper. We refer the reader to reviews [MNO] and [Mo] for proofs and details. Notice that the series $T_{ab}(u)$ in (4.1) corresponds to the series $T_{ba}(u)$ in [MNO], [Mo].

The Yangian $Y(\mathfrak{gl}_N)$ is the unital associative algebra with generators $T_{ab}^{\{s\}}$, $a, b = 1, \dots, N$ and $s = 1, 2, \dots$. Organize them into generating series:

$$(4.1) \quad T_{ab}(u) = \delta_{ab} + \sum_{s=1}^{\infty} T_{ab}^{\{s\}} u^{-s}, \quad a, b = 1, \dots, N.$$

The defining relations in $Y(\mathfrak{gl}_N)$ have the form

$$(4.2) \quad (u - v) [T_{ab}(u), T_{cd}(v)] = T_{cb}(v) T_{ad}(u) - T_{cb}(u) T_{ad}(v),$$

for all $a, b, c, d = 1, \dots, N$.

Combine all series (4.1) together into a series $T(u) = \sum_{a,b=1}^N E_{ab} \otimes T_{ab}(u)$ with coefficients in $\text{End}(\mathbb{C}^N) \otimes Y(\mathfrak{gl}_N)$. Relations (4.2) can be written as the following equality for series with coefficients in $\text{End}(\mathbb{C}^N) \otimes \text{End}(\mathbb{C}^N) \otimes Y(\mathfrak{gl}_N)$:

$$(4.3) \quad R^{(12)}(u-v)T^{(13)}(u)T^{(23)}(v) = T^{(23)}(v)T^{(13)}(u)R^{(12)}(u-v).$$

The Yangian $Y(\mathfrak{gl}_N)$ is a Hopf algebra. In terms of generating series (4.1), the coproduct $\Delta : Y(\mathfrak{gl}_N) \rightarrow Y(\mathfrak{gl}_N) \otimes Y(\mathfrak{gl}_N)$ reads as follows:

$$(4.4) \quad \Delta(T_{ab}(u)) = \sum_{c=1}^N T_{cb}(u) \otimes T_{ac}(u), \quad a, b = 1, \dots, N.$$

This formula amounts to an equality

$$(4.5) \quad (\text{id} \otimes \Delta)(T(u)) = T^{(13)}(u)T^{(12)}(u)$$

for series with coefficients in $\text{End}(V) \otimes Y(\mathfrak{gl}_N) \otimes Y(\mathfrak{gl}_N)$.

There is a one-parameter family of automorphisms $\rho_x : Y(\mathfrak{gl}_N) \rightarrow Y(\mathfrak{gl}_N)$ that is defined in terms of the series $T(u)$ by the rule $\rho_x(T(u)) = T(u-x)$; in the right hand side, $(u-x)^{-1}$ has to be expanded as a power series in u^{-1} .

The Yangian $Y(\mathfrak{gl}_N)$ contains the universal enveloping algebra $U(\mathfrak{gl}_N)$ as a Hopf subalgebra. The embedding is given by $e_{ab} \mapsto T_{ba}^{\{1\}}$ for any $a, b = 1, \dots, N$. We identify $U(\mathfrak{gl}_N)$ with its image in $Y(\mathfrak{gl}_N)$ under this embedding. It is clear from relations (4.2) that for any $a, b = 1, \dots, N$,

$$(4.6) \quad [E_{ab} \otimes 1 + 1 \otimes e_{ab}, T(u)] = 0.$$

The *evaluation homomorphism* $\epsilon : Y(\mathfrak{gl}_N) \rightarrow U(\mathfrak{gl}_N)$ is given by the rule $\epsilon : T_{ab}^{\{1\}} \mapsto e_{ba}$ for any $a, b = 1, \dots, N$, and $\epsilon : T_{ab}^{\{s\}} \mapsto 0$ for any $s > 1$ and all a, b . Both the automorphisms ρ_x and the homomorphism ϵ restricted to the subalgebra $U(\mathfrak{gl}_N)$ are the identity maps.

For a \mathfrak{gl}_N -module M denote by $M(x)$ the $Y(\mathfrak{gl}_N)$ -module induced from M by the homomorphism $\epsilon \circ \rho_x$. The module $M(x)$ is called an *evaluation module* over $Y(\mathfrak{gl}_N)$ with the *evaluation point* x .

Denote by $Y_+(\mathfrak{gl}_N)$ the left ideal in $Y(\mathfrak{gl}_N)$ generated by the coefficients of the series $T_{ab}(u)$ for $1 \leq b < a \leq N$. If $A, B \in Y(\mathfrak{gl}_N)$ and $A - B \in Y_+(\mathfrak{gl}_N)$, then we will write $A \simeq B$. Relations (4.2) imply the following lemma.

Lemma 4.1.

- a) For any $a, b = 1, \dots, N$, the coefficients of the series $[T_{aa}(u), T_{bb}(v)]$ belong to $Y_+(\mathfrak{gl}_N)$.
- b) For any $Z \in Y_+(\mathfrak{gl}_N)$ and any $a = 1, \dots, N$, the coefficients of the series $ZT_{aa}(u)$ belong to $Y_+(\mathfrak{gl}_N)$.

Therefore, for any $a = 1, \dots, N$, the coefficients of the series $T_{aa}(u)$ act on the space $Y(\mathfrak{gl}_N)/Y_+(\mathfrak{gl}_N)$ by multiplication from the right, and those actions commute.

Lemma 4.2. *The coefficients of the series $\Delta(T_{aa}(u)) - T_{aa}(u) \otimes T_{aa}(u)$, $a = 1, \dots, N$, belong to $Y(\mathfrak{gl}_N) \otimes Y_+(\mathfrak{gl}_N) + Y_+(\mathfrak{gl}_N) \otimes Y(\mathfrak{gl}_N)$.*

The proof is straightforward, cf. (4.4).

A vector v in a $Y(\mathfrak{gl}_N)$ -module is called *singular* with respect to the action of $Y(\mathfrak{gl}_N)$ if $Y_+(\mathfrak{gl}_N)v = 0$. A singular vector v that is an eigenvector for the action of $T_{11}(u), \dots, T_{NN}(u)$ is called a *weight singular vector*.

Example. Let M be a \mathfrak{gl}_N -module and $v \in M$ a singular vector of weight $(\Lambda^1, \dots, \Lambda^N)$. Then for any $x \in \mathbb{C}$, the vector v is a weight singular vector with respect to the action of $Y(\mathfrak{gl}_N)$ in the evaluation module $M(x)$.

There is a Hopf algebra anti-involution $\varpi : Y(\mathfrak{gl}_N) \rightarrow Y(\mathfrak{gl}_N)$ that is defined in terms of the series $T_{ab}(u)$ by the rule

$$(4.7) \quad \varpi(T_{ab}(u)) = T_{ba}(u), \quad a, b = 1, \dots, N.$$

In particular, for any $X \in Y(\mathfrak{gl}_N)$ we have

$$(4.8) \quad \Delta(\varpi(X)) = ((\varpi \otimes \varpi)(\Delta(X)))^{(21)}.$$

Consider the series $T^{(k, k+1)}(u) \dots T^{(1, k+1)}(u - k + 1)$ with coefficients in $\text{End}(V^{\otimes k}) \otimes Y(\mathfrak{gl}_N)$. Relations (4.3) and Lemma 3.1 imply that

$$(4.9) \quad \mathbb{A}_{\{k\}}^{(1 \dots k)} T^{(1, k+1)}(u - k + 1) \dots T^{(k, k+1)}(u) = \\ = T^{(k, k+1)}(u) \dots T^{(1, k+1)}(u - k + 1) \mathbb{A}_{\{k\}}^{(1 \dots k)}.$$

Therefore, the coefficients of the series $T^{(k, k+1)}(u) \dots T^{(1, k+1)}(u - k + 1)$ preserve $V^{\wedge k}$. Define the series $T^{\wedge k}(u)$ with coefficients in $\text{End}(V^{\wedge k}) \otimes Y(\mathfrak{gl}_N)$:

$$(4.10) \quad T^{\wedge k}(u) = T^{(k, k+1)}(u) \dots T^{(1, k+1)}(u - k + 1) \Big|_{V^{\wedge k}}^{(1 \dots k)}.$$

In particular, $T^{\wedge 1}(u) = T(u)$. Formulae (4.3) and (3.3) imply that the introduced series satisfy the following relations:

$$(4.11) \quad (R^{\wedge k, \wedge l}(u - v))^{(12)} (T^{\wedge k}(u))^{(13)} (T^{\wedge l}(v))^{(23)} = \\ = (T^{\wedge l}(v))^{(23)} (T^{\wedge k}(u))^{(13)} (R^{\wedge k, \wedge l}(u - v))^{(12)}.$$

Lemma 4.3. $(\text{id} \otimes \Delta)(T^{\wedge k}(u)) = (T^{\wedge k}(u))^{(13)} (T^{\wedge k}(u))^{(12)}.$

Proof. The statement follows from the coproduct formula (4.5) and the definition (4.10) of $T^{\wedge k}(u)$. \square

Since the space $V^{\wedge N}$ is one-dimensional, the coefficients of the series $T^{\wedge N}(u)$ belong to the Yangian $Y(\mathfrak{gl}_N)$. The series $T^{\wedge N}(u)$ is called the *quantum determinant* and is denoted by $\text{qdet } T(u)$. Explicit formulae for the quantum determinant that follow from (4.9) and (4.10) are

$$(4.12) \quad \begin{aligned} \text{qdet } T(u) &= \text{sign}(\sigma) \sum_{\tau \in S_N} \text{sign}(\tau) T_{\sigma_1, \tau_1}(u) \dots T_{\sigma_N, \tau_N}(u - N + 1) \\ &= \text{sign}(\sigma) \sum_{\mathbf{b} \in S_N} \text{sign}(\tau) T_{\tau_N, \sigma_N}(u - N + 1) \dots T_{\tau_1, \sigma_1}(u), \end{aligned}$$

where the permutation $\sigma \in S_N$ is arbitrary. Relations (4.11) and Lemma 3.2 imply that the coefficients of $\text{qdet } T(u)$ are central in $Y(\mathfrak{gl}_N)$.

Theorem 4.4. [MNO] *The coefficients of $\text{qdet } T(u)$ are free generators of the center of the Yangian $Y(\mathfrak{gl}_N)$.*

Let $\text{tr}_W : \text{End}(W) \rightarrow \mathbb{C}$ be the trace map. For any $Q \in \text{End}(V)$ define the series

$$(4.13) \quad \mathcal{T}_{k,Q}(u) = (\text{tr}_{V^{\wedge k}} \otimes \text{id})(Q^{\wedge k} T^{\wedge k}(u)),$$

$k = 1, \dots, N$, with coefficients in $Y(\mathfrak{gl}_N)$. We set $\mathcal{T}_{0,Q}(u) = 1$ by definition. It is easy to see that $\mathcal{T}_{N,Q}(u) = \det Q \text{qdet } T(u)$. The series $\mathcal{T}_{1,Q}(u), \dots, \mathcal{T}_{N,Q}(u)$ are called the *transfer matrices*. They have been introduced in [KS].

Proposition 4.5. [KS] *We have $[\mathcal{T}_{k,Q}(u), \mathcal{T}_{l,Q}(v)] = 0$.*

Proof. The statement follows from formulae (4.11) and (3.4):

$$\begin{aligned} \mathcal{T}_{k,Q}(u) \mathcal{T}_{l,Q}(v) &= (\text{tr}_{V^{\wedge k}} \otimes \text{tr}_{V^{\wedge l}} \otimes \text{id}) \left((Q^{\wedge k} T^{\wedge k}(u))^{(13)} (Q^{\wedge l} T^{\wedge l}(v))^{(23)} \right) = \\ &= (\text{tr}_{V^{\wedge k}} \otimes \text{tr}_{V^{\wedge l}} \otimes \text{id}) \left((Q^{\wedge k})^{(1)} (Q^{\wedge l})^{(2)} (T^{\wedge k}(u))^{(13)} (T^{\wedge l}(v))^{(23)} \right) = \\ &= (\text{tr}_{V^{\wedge k}} \otimes \text{tr}_{V^{\wedge l}} \otimes \text{id}) \left((Q^{\wedge k})^{(1)} (Q^{\wedge l})^{(2)} \left((R^{\wedge k, \wedge l}(u - v))^{(12)} \right)^{-1} \times \right. \\ &\quad \left. \times (T^{\wedge k}(v))^{(23)} (T^{\wedge l}(u))^{(13)} (R^{\wedge k, \wedge l}(u - v))^{(12)} \right) = \\ &= (\text{tr}_{V^{\wedge k}} \otimes \text{tr}_{V^{\wedge l}} \otimes \text{id}) \left(\left((R^{\wedge k, \wedge l}(u - v))^{(12)} \right)^{-1} (Q^{\wedge k})^{(1)} (Q^{\wedge l})^{(2)} \times \right. \\ &\quad \left. \times (T^{\wedge k}(v))^{(23)} (T^{\wedge l}(u))^{(13)} (R^{\wedge k, \wedge l}(u - v))^{(12)} \right) = \\ &= (\text{tr}_{V^{\wedge k}} \otimes \text{tr}_{V^{\wedge l}} \otimes \text{id}) \left((Q^{\wedge k})^{(1)} (Q^{\wedge l})^{(2)} (T^{\wedge l}(v))^{(23)} (T^{\wedge k}(u))^{(13)} \right) = \\ &= (\text{tr}_{V^{\wedge k}} \otimes \text{tr}_{V^{\wedge l}} \otimes \text{id}) \left((Q^{\wedge l} T^{\wedge l}(v))^{(23)} (Q^{\wedge k} T^{\wedge k}(u))^{(13)} \right) = \\ &= \mathcal{T}_{l,Q}(v) \mathcal{T}_{k,Q}(u). \end{aligned}$$

□

Theorem 4.6. [NO] *If the matrix $Q \in \text{End}(V)$ has a simple spectrum, then the coefficients of the series $\mathcal{T}_{1,Q}(u), \dots, \mathcal{T}_{N-1,Q}(u)$ and $\text{qdet } T(u)$ generate a maximal commutative subalgebra in $Y(\mathfrak{gl}_N)$.*

Recall that the Yangian $Y(\mathfrak{gl}_N)$ contains $U(\mathfrak{gl}_N)$ as a subalgebra, the embedding being given by $e_{ab} \mapsto T_{ba}^{\{1\}}$.

Proposition 4.7. [KS] *Let Q be the identity matrix. Then for any $k = 1, \dots, N$, the coefficients of $\mathcal{T}_{k,Q}(u)$ commute with the subalgebra $U(\mathfrak{gl}_N) \subset Y(\mathfrak{gl}_N)$.*

Proof. Similarly to (4.6), for any $a, b = 1, \dots, N$, we have

$$[e_{ab}|_{V^{\wedge k}} \otimes 1 + 1 \otimes e_{ab}, T^{\wedge k}(u)] = 0,$$

which implies that for the identity matrix Q ,

$$[e_{ab}, \mathcal{T}_{k,Q}(u)] = -(\text{tr}_{V^{\wedge k}} \otimes \text{id})\left([e_{ab}|_{V^{\wedge k}} \otimes 1, T^{\wedge k}(u)]\right) = 0$$

due to the cyclic property of the trace. □

Lemma 4.8. *For any $m = k, \dots, N$, and any distinct $i_1, \dots, i_k \in \{1, \dots, m\}$, we have*

$$\begin{aligned} \mathcal{T}_{k,Q}(u) &= \\ &= \frac{m!(N-m)!}{k!(N-k)!} (\text{tr}_{V^{\otimes m}} \otimes \text{id})(Q^{(i_1)} \dots Q^{(i_k)} T^{(i_1, m+1)}(u) \dots T^{(i_k, m+1)}(u - k + 1) \mathbb{A}_{\{m\}}^{(1 \dots m)}). \end{aligned}$$

Proof. If $m = k$ and $i_j = j$ for all $j = 1, \dots, k$, the claim is equivalent to formula (4.13). In general, we are also using the relations $P^{(ij)} Q^{(i)} T^{(i, m+1)}(u) = T^{(j, m+1)}(u) Q^{(j)} P^{(ij)}$ and $P^{(ij)} \mathbb{A}_{\{m\}} = -\mathbb{A}_{\{m\}} = \mathbb{A}_{\{m\}} P^{(ij)}$, $1 \leq i < j \leq m$, for the flip maps, and the formula

$$(\text{id}^{\otimes k} \otimes \text{tr}_{V^{\otimes(m-k)}}) \mathbb{A}_{\{m\}} = \frac{m!(N-m)!}{k!(N-k)!} \mathbb{A}_{\{k\}}. \quad \square$$

Consider the algebra $Y(\mathfrak{gl}_N)[[u^{-1}, \partial_u]]$ of formal differential operators; the indeterminates u^{-1} and ∂_u do not commute obeying the following relation $\partial_u \cdot u^{-s} = u^{-s} \partial_u - s u^{-s-1}$ instead. Following [T], for any $m = 1, \dots, N$, set

$$\begin{aligned} (4.14) \quad \mathfrak{D}_{m,Q}(u, \partial_u) &= \\ &= (\text{tr}_{V^{\otimes m}} \otimes \text{id}) \left((1 - Q^{(1)} T^{(1, m+1)}(u) e^{-\partial_u}) \dots (1 - Q^{(m)} T^{(m, m+1)}(u) e^{-\partial_u}) \mathbb{A}_{\{m\}}^{(1 \dots m)} \right) \end{aligned}$$

and $\mathfrak{D}_{0,Q}(u, \partial_u) = 1$.

Proposition 4.9. [T] *For any $m = 1, \dots, N$, we have*

$$(4.15) \quad \mathfrak{D}_{m,Q}(u, \partial_u) = \frac{1}{(N-m)!} \sum_{k=0}^m (-1)^k \frac{(N-k)!}{(m-k)!} \mathcal{T}_{k,Q}(u) e^{-k \partial_u}.$$

Proof. Expand the product under the trace in formula (4.14),

$$\begin{aligned} \mathfrak{D}_{m,Q}(u, \partial_u) &= \sum_{k=0}^m \sum_{1 \leq i_1 < \dots < i_k \leq m} (-1)^k \times \\ &\times (\text{tr}_{V^{\otimes m}} \otimes \text{id})(Q^{(i_1)} T^{(i_1, m+1)}(u) \dots Q^{(i_k)} T^{(i_k, m+1)}(u - k + 1) \mathbb{A}_{\{m\}}^{(1\dots m)}) e^{-k\partial_u}. \end{aligned}$$

Then Lemma 4.8 yields the claim. \square

Remark. Formula (4.15) for $m = N$ shows that the expression $\mathfrak{D}_{N,Q}(u, \partial_u)$ serves as a generating function of the transfer matrices $\mathcal{T}_{1,Q}(u), \dots, \mathcal{T}_{N,Q}(u)$:

$$(4.16) \quad \mathfrak{D}_{N,Q}(u, \partial_u) = \sum_{k=0}^N (-1)^k \mathcal{T}_{k,Q}(u) e^{-k\partial_u}.$$

Remark. Taking into account formula (4.9) and the equality $(\mathbb{A}_{\{m\}})^2 = \mathbb{A}_{\{m\}}$, one can show that

$$\begin{aligned} (4.17) \quad &(1 - Q^{(1)} T^{(1, m+1)}(u) e^{-\partial_u}) \dots (1 - Q^{(m)} T^{(m, m+1)}(u) e^{-\partial_u}) \mathbb{A}_{\{m\}}^{(1\dots m)} = \\ &= \mathbb{A}_{\{m\}}^{(1\dots m)} (1 - Q^{(1)} T^{(1, m+1)}(u) e^{-\partial_u}) \dots (1 - Q^{(m)} T^{(m, m+1)}(u) e^{-\partial_u}) \mathbb{A}_{\{m\}}^{(1\dots m)}. \end{aligned}$$

Since $\mathbb{A}_{\{N\}}$ is a one-dimensional projector, formula (4.17) implies the following proposition.

Proposition 4.10. *We have*

$$\begin{aligned} &(\text{id}^{\otimes N} \otimes \mathfrak{D}_{N,Q}(u, \partial_u)) \mathbb{A}_{\{N\}}^{(1\dots N)} = \\ &= (1 - Q^{(1)} T^{(1, N+1)}(u) e^{-\partial_u}) \dots (1 - Q^{(N)} T^{(N, N+1)}(u) e^{-\partial_u}) \mathbb{A}_{\{N\}}^{(1\dots N)}. \end{aligned}$$

Let Q^\dagger be the matrix transposed to Q ; for example, $E_{ab}^\dagger = E_{ba}$.

Proposition 4.11. *We have $\varpi(\mathcal{T}_{k,Q}(u)) = \mathcal{T}_{k,Q^\dagger}(u)$, where ϖ is given by (4.7).*

Proof. The claim follows from Lemma 4.8 for $\mathcal{T}_{k,Q}(u)$ with $m = k$ and $i_j = k - j + 1$, formula (4.9), two more observations, $\varpi(T(u)) = (T(u))^\dagger$ and $(\mathbb{A}_{\{k\}}^{(1\dots k)})^\dagger = \mathbb{A}_{\{k\}}^{(1\dots k)}$, and the standard properties of the trace:

$$\begin{aligned} \varpi(\mathcal{T}_{k,Q}(u)) &= \\ &= \frac{1}{k!} (\text{tr}_{V^{\otimes k}} \otimes \varpi)(Q^{(k)} \dots Q^{(1)} T^{(k, k+1)}(u) \dots T^{(1, k+1)}(u - k + 1) \mathbb{A}_{\{k\}}^{(1\dots k)}) = \\ &= \frac{1}{k!} (\text{tr}_{V^{\otimes k}} \otimes \varpi)(Q^{(k)} \dots Q^{(1)} \mathbb{A}_{\{k\}}^{(1\dots k)} T^{(1, k+1)}(u - k + 1) \dots T^{(k, k+1)}(u)) = \\ &= \frac{1}{k!} (\text{tr}_{V^{\otimes k}} \otimes \text{id})(Q^{(k)} \dots Q^{(1)} \mathbb{A}_{\{k\}}^{(1\dots k)} (T^{(k, k+1)}(u))^\dagger \dots (T^{(1, k+1)}(u - k + 1))^\dagger) = \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{k!} (\text{tr}_{V^{\otimes k}} \otimes \text{id}) \left((Q^{(k)} \dots Q^{(1)} \mathbb{A}_{\{k\}}^{(1\dots k)} (T^{(k,k+1)}(u))^\dagger \dots (T^{(1,k+1)}(u-k+1))^\dagger)^\dagger \right) = \\
&= \frac{1}{k!} (\text{tr}_{V^{\otimes k}} \otimes \text{id}) \left(T^{(k,k+1)}(u) \dots T^{(1,k+1)}(u-k+1) \mathbb{A}_{\{k\}}^{(1\dots k)} (Q^{(k)})^\dagger \dots (Q^{(1)})^\dagger \right) = \\
&= \frac{1}{k!} (\text{tr}_{V^{\otimes k}} \otimes \text{id}) \left((Q^{(k)})^\dagger \dots (Q^{(1)})^\dagger T^{(k,k+1)}(u) \dots T^{(1,k+1)}(u-k+1) \mathbb{A}_{\{k\}}^{(1\dots k)} \right) = \\
&= \mathcal{T}_{k,Q^\dagger}(u). \quad \square
\end{aligned}$$

5. The Bethe ansatz

In this section we formulate the main technical result of this paper, Theorem 5.4. Equality (5.8) is, in a sense, a universal formulation of the eigenvalue problem for Bethe eigenvectors of the transfermatrix $\mathcal{T}_{k,Q}(u)$. Given a $Y(\mathfrak{gl}_N)$ -module M and a weight singular vector $v \in M$, we will apply formula (5.8) to v and obtain the Bethe eigenvector for the action of $\mathcal{T}_{k,Q}(u)$ in M and the formula for the corresponding eigenvalue. In Section 6 we will do it in detail for M being a tensor product of highest weight evaluation modules and $v \in M$ being the tensor product of highest weight vectors.

Let $\xi = (\xi^1, \dots, \xi^{N-1})$ be a collection of nonnegative integers. Set $\xi^{<a} = \xi^1 + \dots + \xi^{a-1}$, $a = 1, \dots, N$, and $|\xi| = \xi^1 + \dots + \xi^{N-1} = \xi^{<N}$. Consider a series in $|\xi|$ variables $t_1^1, \dots, t_{\xi^1}^1, \dots, t_1^{N-1}, \dots, t_{\xi^{N-1}}^{N-1}$ with coefficients in $Y(\mathfrak{gl}_N)$:

$$\begin{aligned}
(5.1) \quad \widehat{\mathbb{B}}_\xi(t_1^1, \dots, t_{\xi^{N-1}}^{N-1}) &= (\text{tr}_{V^{\otimes |\xi|}} \otimes \text{id}) \left(T^{(1,|\xi|+1)}(t_1^1) \dots T^{(|\xi|,|\xi|+1)}(t_{\xi^{N-1}}^{N-1}) \times \right. \\
&\quad \times \left. \prod_{(a,i) < (b,j)}^{\rightarrow} R^{(\xi^{<b}+j, \xi^{<a}+i)}(t_j^b - t_i^a) E_{21}^{\otimes \xi^1} \otimes \dots \otimes E_{N,N-1}^{\otimes \xi^{N-1}} \otimes 1 \right),
\end{aligned}$$

where the pairs are ordered lexicographically, $(a,i) < (b,j)$ if $a < b$, or $a = b$ and $i < j$, and the product is taken over all two element subsets of the set $\{(c,k) \mid c = 1, \dots, N-1, k = 1, \dots, \xi^c\}$. Further on, we will abbreviate, $t = (t_1^1, \dots, t_{\xi^{N-1}}^{N-1})$.

Lemma 5.1. *The series $\widehat{\mathbb{B}}_\xi(t)$ is divisible by $\prod_{a=1}^N \prod_{1 \leq i < j \leq \xi^a} (t_j^a - t_i^a + 1)$ in $Y(\mathfrak{gl}_N)[t_1^1, \dots, t_{\xi^{N-1}}^{N-1}][[(t_1^1)^{-1}, \dots, (t_{\xi^{N-1}}^{N-1})^{-1}]]$.*

Proof. Using the Yang-Baxter equation (3.2) we can rearrange the factors in the product of R -matrices in formula (5.1) such that the product $\prod_{1 \leq a < N}^{\rightarrow} \prod_{1 \leq i < j \leq \xi^a}^{\rightarrow} R^{(\xi^{<a}+j, \xi^{<a}+i)}(t_j^a - t_i^a)$ appears on the right. Then formula $R(u)E_{bc} \otimes E_{bc} = (u+1)E_{bc} \otimes E_{bc}$ yields the claim. \square

Let $\theta : Y(\mathfrak{gl}_N) \rightarrow Y(\mathfrak{gl}_N)/Y_+(\mathfrak{gl}_N)$ be the canonical projection.

Lemma 5.2. *The series $\theta(\widehat{\mathbb{B}}_\xi(t))$ is divisible by $\prod_{a=1}^{N-2} \prod_{b=a+2}^{N-1} \prod_{i=1}^{\xi^a} \prod_{j=1}^{\xi^b} (t_j^b - t_i^a)$ in $(Y(\mathfrak{gl}_N)/Y_+(\mathfrak{gl}_N))[t_1^1, \dots, t_{\xi^{N-1}}^{N-1}][[(t_1^1)^{-1}, \dots, (t_{\xi^{N-1}}^{N-1})^{-1}]]$.*

The proof is given in Section 11 after Proposition 11.3.

Set

$$(5.2) \quad \mathbb{B}_\xi(t) = \widehat{\mathbb{B}}_\xi(t) \prod_{a=1}^{N-1} \prod_{1 \leq i < j \leq \xi^a} \frac{1}{t_j^a - t_i^a + 1} \prod_{1 \leq a < b < N} \prod_{i=1}^{\xi^a} \prod_{j=1}^{\xi^b} \frac{1}{t_j^b - t_i^a},$$

cf. (5.1). To indicate the dependence on N , if necessary, we will write $\mathbb{B}_\xi^{(N)}(t)$.

Remark. The expression $\mathbb{B}_\xi(t)$ serves as a creation operator to the universal weight function and Bethe vectors in tensor products of evaluation modules. See more details in Section 6.

Example. Let $N = 2$ and $\xi = (\xi^1)$. Then $\mathbb{B}_\xi(t) = T_{12}(t_1^1) \dots T_{12}(t_{\xi^1}^1)$.

Example. Let $N = 3$ and $\xi = (1, 1)$. Then

$$\mathbb{B}_\xi(t) = T_{12}(t_1^1) T_{23}(t_1^2) + \frac{1}{t_1^2 - t_1^1} T_{13}(t_1^1) T_{22}(t_1^2).$$

Example. Let $N = 4$ and $\xi = (1, 1, 1)$. Then

$$\begin{aligned} \mathbb{B}_\xi(t) = & T_{12}(t_1^1) T_{23}(t_1^2) T_{34}(t_1^3) + \\ & + \frac{1}{t_1^2 - t_1^1} T_{13}(t_1^1) T_{22}(t_1^2) T_{34}(t_1^3) + \frac{1}{t_1^3 - t_1^2} T_{12}(t_1^1) T_{24}(t_1^2) T_{33}(t_1^3) + \\ & + \frac{1}{(t_1^2 - t_1^1)(t_1^3 - t_1^1)} (T_{14}(t_1^1) T_{22}(t_1^2) T_{33}(t_1^3) + T_{13}(t_1^1) T_{24}(t_1^2) T_{32}(t_1^3)) + \\ & + \frac{(t_1^2 - t_1^1)(t_1^3 - t_1^1) + 1}{(t_1^2 - t_1^1)(t_1^3 - t_1^1)(t_1^3 - t_1^2)} T_{14}(t_1^1) T_{23}(t_1^2) T_{32}(t_1^3). \end{aligned}$$

In general, $\mathbb{B}_\xi(t)$ is a sum of the products

$$(5.3) \quad T_{a_1, b_1}(t_1^1) \dots T_{a_{|\xi|}, b_{|\xi|}}(t_{\xi^{N-1}}^{N-1}) p(t_1^1, \dots, t_{\xi^{N-1}}^{N-1}) \prod_{1 \leq a < b < N} \prod_{i=1}^{\xi^a} \prod_{j=1}^{\xi^b} \frac{1}{t_j^b - t_i^a}$$

with various $a_1, \dots, a_{|\xi|}$, $b_1, \dots, b_{|\xi|}$, and polynomials $p(t_1^1, \dots, t_{\xi^{N-1}}^{N-1})$.

The direct product of the symmetric groups $S_{\xi^1} \times \dots \times S_{\xi^{N-1}}$ acts on expressions in $|\xi|$ variables, permuting the variables with the same superscript:

$$(5.4) \quad \sigma^1 \times \dots \times \sigma^{N-1} : f(t_1^1, \dots, t_{\xi^{N-1}}^{N-1}) \mapsto f(t_{\sigma_1^1}^1, \dots, t_{\sigma_{\xi^1}^1}^1; \dots; t_{\sigma_1^{N-1}}^{N-1}, \dots, t_{\sigma_{\xi^{N-1}}^{N-1}}^{N-1}),$$

where $\sigma^a \in S_{\xi^a}$, $a = 1, \dots, N-1$.

Lemma 5.3. [TV1, Theorem 3.3.4] *The expression $\mathbb{B}_\xi(t)$ is invariant under the action (5.4) of the group $S_{\xi^1} \times \dots \times S_{\xi^{N-1}}$.*

Fix a diagonal matrix $Q = \sum_{a=1}^N Q_a E_{aa}$. Introduce the series

$$\begin{aligned} (5.5) \quad \mathcal{J}_{\xi, Q}^{a, i}(t) = & Q_a T_{aa}(t_i^a) \prod_{j=1}^{\xi^{a-1}} (t_i^a - t_j^{a-1} + 1) \prod_{j=1}^{\xi^a} (t_i^a - t_j^a - 1) \prod_{j=1}^{\xi^{a+1}} (t_i^a - t_j^{a+1}) + \\ & + Q_{a+1} T_{a+1, a+1}(t_i^a) \prod_{j=1}^{\xi^{a-1}} (t_i^a - t_j^{a-1}) \prod_{j=1}^{\xi^a} (t_i^a - t_j^a + 1) \prod_{j=1}^{\xi^{a+1}} (t_i^a - t_j^{a+1} - 1), \end{aligned}$$

$a = 1, \dots, N-1$, $i = 1, \dots, \xi^a$. Here we set $\xi^0 = \xi^N = 0$.

Given the following data: integers $a_1, \dots, a_{|\xi|+k-1}, b_1, \dots, b_{|\xi|+k-1}, c \in \{1, \dots, N\}$ and $i \in \{1, \dots, \xi^c\}$, a sequence $s_1, \dots, s_{|\xi|+k-1}$, which is a permutation of the sequence $u, \dots, u-k+1, t_1^1, \dots, \widehat{t_i^c}, \dots, t_{\xi^{N-1}}^{N-1}$, where the variable t_i^c is omitted, and a polynomial $p(u; t_1^1, \dots, t_{\xi^{N-1}}^{N-1})$, consider the product

$$(5.6) \quad T_{a_1, b_1}(s_1) \dots T_{a_{|\xi|+k-1}, b_{|\xi|+k-1}}(s_{|\xi|+k-1}) \mathcal{J}_{\xi, Q}^{c, i}(t) \times \\ \times p(u; t_1^1, \dots, t_{\xi^{N-1}}^{N-1}) \prod_{a=1}^{N-1} \prod_{j=1}^{\xi^a} \left(\frac{1}{u - t_j^a} \prod_{1 \leq j < l \leq \xi^a} \frac{1}{t_j^a - t_l^a} \right) \prod_{a=1}^{N-2} \prod_{j=1}^{\xi^a} \prod_{l=1}^{\xi^{a+1}} \frac{1}{(t_l^{a+1} - t_j^a)^2}$$

where the factors $(u - t_j^a)^{-1}$ in (5.6) have to be expanded as power series in u^{-1} . We denote by $I_{\xi, k, Q}$ the span over \mathbb{C} of all products (5.6) with various $a_1, \dots, a_{|\xi|+k-1}$, $b_1, \dots, b_{|\xi|+k-1}$, c, i , $s_1, \dots, s_{|\xi|+k-1}$, and $p(u; t_1^1, \dots, t_{\xi^{N-1}}^{N-1})$.

Set

$$(5.7) \quad \mathcal{X}_{\xi, Q}^a(u; t) = Q_a T_{aa}(u) \prod_{i=1}^{\xi^{a-1}} \frac{u - t_i^{a-1} + 1}{u - t_i^{a-1}} \prod_{i=1}^{\xi^a} \frac{u - t_i^a - 1}{u - t_i^a},$$

$a = 1, \dots, N$. Expressions (5.7) are regarded as power series in u^{-1} with coefficients in $Y(\mathfrak{gl}_N)[t_1^1, \dots, t_{\xi^{N-1}}^{N-1}]$.

Theorem 5.4. *Let Q be a diagonal matrix. For any $k = 1, \dots, N$, we have*

$$(5.8) \quad \mathcal{T}_{k, Q}(u) \mathbb{B}_{\xi}(t) \simeq \mathbb{B}_{\xi}(t) \sum_{\mathbf{a}} \prod_{r=1}^k \mathcal{X}_{\xi, Q}^{a_r}(u - r + 1; t) + \mathcal{U}_{\xi, k, Q}(u; t),$$

where the sum is taken over all k -tuples $\mathbf{a} = (a_1, \dots, a_k)$ such that $1 \leq a_1 < \dots < a_k \leq N$, and $\mathcal{U}_{\xi, k, Q}(u; t)$ belongs to $I_{\xi, k, Q}$, cf. (5.6). By Lemma 4.1, the order of the product of elements $\mathcal{X}_{\xi, Q}^{a_r}$ is irrelevant.

The theorem is proved in Section 11.

Remark. Similarly to formula (4.16), the expressions $\sum_{\mathbf{a}} \prod_{r=1}^k \mathcal{X}_{\xi, Q}^{a_r}(u - r + 1; t)$ can be obtained as coefficients of a suitable formal differential operator:

$$(5.9) \quad \mathfrak{M}_{\xi, Q}(u, \partial_u) = (1 - \mathcal{X}_{\xi, Q}^1(u; t) e^{-\partial_u}) \dots (1 - \mathcal{X}_{\xi, Q}^N(u; t) e^{-\partial_u}) = \\ = \sum_{k=0}^N (-1)^k \sum_{1 \leq a_1 < \dots < a_k \leq N} \prod_{r=1}^k \mathcal{X}_{\xi, Q}^{a_r}(u - r + 1; t) e^{-k\partial_u}.$$

Remark. Since the coefficients of the series $\text{qdet } T(u)$ are central in $Y(\mathfrak{gl}_N)$, the first formula of (4.12) for the identity permutation σ implies that

$$\text{qdet } T(u) \mathbb{B}_{\xi}(t) \simeq \mathbb{B}_{\xi}(t) \prod_{a=1}^N T_{aa}(u - a + 1).$$

Together with the relation $\mathcal{T}_{N, Q}(u) = \det Q \text{ qdet } T(u)$, this proves Theorem 5.4 for $k = N$.

6. Tensor products of evaluation modules

In this section we consider the action of the transfer matrices $\mathcal{T}_{1,Q}(u), \dots, \mathcal{T}_{N,Q}(u)$ in a tensor product of evaluation modules over the Yangian $Y(\mathfrak{gl}_N)$. We will apply Theorem 5.4 to get eigenvectors and eigenvalues of the transfer matrices. In Section 7 we will also show that the operators of the action are symmetric with respect to a certain symmetric bilinear form, see Theorem 7.6.

Let M_1, \dots, M_n be \mathfrak{gl}_N -modules. Consider the tensor product $M_1(z_1) \otimes \dots \otimes M_n(z_n)$ of evaluation $Y(\mathfrak{gl}_N)$ -modules. The operators $T_{ab}(u)|_{M_1(z_1) \otimes \dots \otimes M_n(z_n)}$, $a, b = 1, \dots, N$, are $\text{End}(M_1 \otimes \dots \otimes M_n)$ -valued rational functions in u, z_1, \dots, z_n with the denominator $\prod_{i=1}^n (u - z_i)$, and $T_{ab}(u)|_{M_1(z_1) \otimes \dots \otimes M_n(z_n)} = \delta_{ab} + O(u^{-1})$ as $u \rightarrow \infty$.

Further on, we will abbreviate, $z = (z_1, \dots, z_n)$. For any $k = 1, \dots, n$, the operator

$$(6.1) \quad \mathcal{T}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u; z) = \mathcal{T}_{k,Q}(u)|_{M_1(z_1) \otimes \dots \otimes M_n(z_n)}$$

is a rational function in u, z_1, \dots, z_n with the denominator $\prod_{i=1}^n \prod_{j=0}^{k-1} (u - j - z_i)$, and $\mathcal{T}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u; z) = \text{tr}_{V^{\wedge k}} Q^{\wedge k} + O(u^{-1})$ as $u \rightarrow \infty$. The operators $\mathcal{T}_{1,Q}^{M_1 \otimes \dots \otimes M_n}(u; z), \dots, \mathcal{T}_{N,Q}^{M_1 \otimes \dots \otimes M_n}(u; z)$ are called the *transfer matrices of the XXX-type model* associated with the Lie algebra \mathfrak{gl}_N .

Let M_1, \dots, M_n be highest weight \mathfrak{gl}_N -modules with highest weights $\Lambda_1, \dots, \Lambda_n$, $\Lambda_i = (\Lambda_i^1, \dots, \Lambda_i^N)$, and highest weight vectors v_1, \dots, v_n . Then for any z_1, \dots, z_n , the vector $v_1 \otimes \dots \otimes v_n$ is a weight singular vector with respect to the action of $Y(\mathfrak{gl}_N)$ in the module $M_1(z_1) \otimes \dots \otimes M_n(z_n)$ and

$$T_{aa}(u)v_1 \otimes \dots \otimes v_n = v_1 \otimes \dots \otimes v_n \prod_{i=1}^n \frac{u - z_i + \Lambda_i^a}{u - z_i}, \quad a = 1, \dots, N.$$

Let $\xi = (\xi^1, \dots, \xi^{N-1})$ be a collection of nonnegative integers. The function

$$(6.2) \quad \mathbb{B}_{\xi}^{v_1 \otimes \dots \otimes v_n}(t; z) = \mathbb{B}_{\xi}(t)v_1 \otimes \dots \otimes v_n \prod_{a=1}^{N-1} \prod_{i=1}^{\xi^a} \prod_{j=1}^n (t_i^a - z_j) \prod_{a=1}^{N-2} \prod_{i=1}^{\xi^a} \prod_{j=1}^{\xi^{a+1}} (t_j^{a+1} - t_i^a)$$

is a polynomial in $t_1^1, \dots, t_{\xi^{N-1}}^{N-1}$, z_1, \dots, z_n , cf. (5.3) and Lemma 5.2. We call it the *universal weight function* for the XXX-type model.

Let $Q = \sum_{a=1}^N Q_a E_{aa}$ be a diagonal matrix. The system of algebraic equations

$$(6.3) \quad Q_a \prod_{j=1}^n (t_i^a - z_j + \Lambda_j^a) \prod_{j=1}^{\xi^{a-1}} (t_i^a - t_j^{a-1} + 1) \prod_{\substack{j=1 \\ j \neq i}}^{\xi^a} (t_i^a - t_j^a - 1) \prod_{j=1}^{\xi^{a+1}} (t_i^a - t_j^{a+1}) = \\ = Q_{a+1} \prod_{j=1}^n (t_i^a - z_j + \Lambda_j^{a+1}) \prod_{j=1}^{\xi^{a-1}} (t_i^a - t_j^{a-1}) \prod_{\substack{j=1 \\ j \neq i}}^{\xi^a} (t_i^a - t_j^a + 1) \prod_{j=1}^{\xi^{a+1}} (t_i^a - t_j^{a+1} - 1),$$

$a = 1, \dots, N-1$, $i = 1, \dots, \xi^a$, $\xi^0 = \xi^N = 0$, is called the *Bethe ansatz equations*. Say that a solution $\tilde{t} = (\tilde{t}_1^1, \dots, \tilde{t}_{\xi^{N-1}}^{N-1})$ of system (6.3) is *off-diagonal* if $\tilde{t}_i^a \neq \tilde{t}_j^a$ for any $a = 1, \dots, N-1$, $1 \leq i < j \leq \xi^a$, and $\tilde{t}_i^a \neq \tilde{t}_j^{a+1}$ for any $a = 1, \dots, N-2$, $i = 1, \dots, \xi^a$, $j = 1, \dots, \xi^{a+1}$. For any $a = 1, \dots, N$, set

$$(6.4) \quad \mathcal{X}_{\xi, Q}^a(u; t; z; \Lambda) = Q_a \prod_{i=1}^n \frac{u - z_i + \Lambda_i^a}{u - z_i} \prod_{i=1}^{\xi^{a-1}} \frac{u - t_i^{a-1} + 1}{u - t_i^{a-1}} \prod_{i=1}^{\xi^a} \frac{u - t_i^a - 1}{u - t_i^a}.$$

Theorem 6.1. *Let M_1, \dots, M_n be highest weight \mathfrak{gl}_N -modules with highest weights $\Lambda_1, \dots, \Lambda_n$ and highest weight vectors v_1, \dots, v_n . Let Q be a diagonal matrix and $\tilde{t} = (\tilde{t}_1^1, \dots, \tilde{t}_{\xi^{N-1}}^{N-1})$ an off-diagonal solution of system (6.3). Then for any $k = 1, \dots, N$, we have*

$$(6.5) \quad \mathcal{T}_{k, Q}^{M_1 \otimes \dots \otimes M_n}(u; z) \mathbb{B}_{\xi}^{v_1 \otimes \dots \otimes v_n}(\tilde{t}; z) = \mathbb{B}_{\xi}^{v_1 \otimes \dots \otimes v_n}(\tilde{t}; z) \sum_{\mathbf{a}} \prod_{r=1}^k \mathcal{X}_{\xi, Q}^{a_r}(u - r + 1; \tilde{t}; z; \Lambda),$$

where the sum is taken over all k -tuples $\mathbf{a} = (a_1, \dots, a_k)$ such that $1 \leq a_1 < \dots < a_k \leq N$.

Proof. Theorem 5.4 and formula (5.6) imply that

$$\begin{aligned} & \mathcal{T}_{k, Q}^{M_1 \otimes \dots \otimes M_n}(u; z) \mathbb{B}_{\xi}^{v_1 \otimes \dots \otimes v_n}(t; z) = \\ & = \mathbb{B}_{\xi}^{v_1 \otimes \dots \otimes v_n}(t; z) \sum_{\mathbf{a}} \prod_{r=1}^k \mathcal{X}_{\xi, Q}^{a_r}(u - r + 1; t; z; \Lambda) + \mathcal{U}_{\xi, k, Q}^{v_1 \otimes \dots \otimes v_n}(u; t; z; \Lambda), \end{aligned}$$

where $\mathcal{U}_{\xi, k, Q}^{v_1 \otimes \dots \otimes v_n}(u; t; z; \Lambda)$ is a ratio of a polynomial with coefficients in $M_1 \otimes \dots \otimes M_n$ vanishing at solutions of system (6.3) and the product

$$\prod_{a=1}^{N-1} \prod_{1 \leq i < j \leq \xi^a} (t_i^a - t_j^a) \prod_{a=1}^{N-2} \prod_{i=1}^{\xi^a} \prod_{j=1}^{\xi^{a+1}} (t_j^{a+1} - t_i^a)$$

that does not vanish at off-diagonal points $(\tilde{t}_1^1, \dots, \tilde{t}_{\xi^{N-1}}^{N-1})$. This yields formula (6.5). \square

For an off-diagonal solution $\tilde{t} = (\tilde{t}_1^1, \dots, \tilde{t}_{\xi^{N-1}}^{N-1})$ of system (6.3), the vector $\mathbb{B}_{\xi}^{v_1 \otimes \dots \otimes v_n}(\tilde{t}; z)$ is called the *Bethe vector*.

Remark. Similarly to formula (5.9), the eigenvalues $\sum_{\mathbf{a}} \prod_{r=1}^k \mathcal{X}_{\xi, Q}^{a_r}(u - r + 1; t; z; \Lambda)$ of the transfer matrices $\mathcal{T}_{k, Q}^{M_1 \otimes \dots \otimes M_n}(u; z)$ can be obtained as coefficients of a suitable difference operator:

$$\begin{aligned} (6.6) \quad \mathfrak{M}_{\xi, Q}(u, \partial_u; t; z; \Lambda) &= (1 - \mathcal{X}_{\xi, Q}^1(u; t; z; \Lambda) e^{-\partial_u}) \dots (1 - \mathcal{X}_{\xi, Q}^N(u; t; z; \Lambda) e^{-\partial_u}) = \\ &= \sum_{k=0}^N (-1)^k \sum_{1 \leq a_1 < \dots < a_k \leq N} \prod_{r=1}^k \mathcal{X}_{\xi, Q}^{a_r}(u - r + 1; t; z; \Lambda) e^{-k \partial_u}. \end{aligned}$$

Remark. Since the coefficients of the series $\text{qdet } T(u)$ are central in $Y(\mathfrak{gl}_N)$, the first formula of (4.12) for the identity permutation σ implies that

$$\text{qdet } T(u) \big|_{M_1(z_1) \otimes \dots \otimes M_n(z_n)} = \prod_{i=1}^n \prod_{a=1}^N \frac{u - a + 1 - z_i + \Lambda_i^a}{u - a + 1 - z_i}.$$

Together with relation $\mathcal{T}_{N,Q}(u) = \det Q \text{qdet } T(u) \prod_{i=1}^n \prod_{a=1}^N (u - a + 1 - z_i)$, this proves Theorem 6.1 for $k = N$.

Proposition 6.2. [KR1] *Let Q be the identity matrix, that is, $Q_1 = \dots = Q_N = 1$, and $\tilde{t} = (\tilde{t}_1^1, \dots, \tilde{t}_{\xi^{N-1}}^{N-1})$ an off-diagonal solution of the Bethe ansatz equations (6.3). Then the Bethe vector $\mathbb{B}_\xi^{v_1 \otimes \dots \otimes v_n}(\tilde{t}; z)$ is a singular vector with respect to the \mathfrak{gl}_N -action in $M_1 \otimes \dots \otimes M_n$ of weight $(\sum_{i=1}^n \Lambda_i^1 - \xi^1, \sum_{i=1}^n \Lambda_i^2 + \xi^1 - \xi^2, \dots, \sum_{i=1}^n \Lambda_i^N + \xi^{N-1})$.*

Proof. Clearly, for any t the vector $\mathbb{B}_\xi^{v_1 \otimes \dots \otimes v_n}(t; z)$ has the indicated weight. The fact that the Bethe vector $\mathbb{B}_\xi^{v_1 \otimes \dots \otimes v_n}(\tilde{t}; z)$ is singular is proved in Section 11. \square

Remark. Let M_1, \dots, M_n be any \mathfrak{gl}_N -modules, not necessarily highest weight ones, and let $v \in M_1 \otimes \dots \otimes M_n$ be any weight singular vector with respect to the action of $Y(\mathfrak{gl}_N)$ in the module $M_1(z_1) \otimes \dots \otimes M_n(z_n)$. Define polynomials $q_1^v(u), \dots, q_N^v(u)$ by the rule

$$T_{aa}(u)v \prod_{i=1}^n (u - z_i) = q_a^v(u)v, \quad a = 1, \dots, N.$$

The function

$$\mathbb{B}_\xi^v(t; z) = \mathbb{B}_\xi(t; z)v \prod_{a=1}^{N-1} \prod_{j=1}^{\xi^a} \prod_{i=1}^n (t_i^a - z_j) \prod_{a=1}^{N-2} \prod_{i=1}^{\xi^a} \prod_{j=1}^{\xi^{a+1}} (t_j^{a+1} - t_i^a)$$

is a polynomial in $t_1^1, \dots, t_{\xi^{N-1}}^{N-1}$, cf. (5.3) and Lemma 5.2. For $a = 1, \dots, N$, set

$$\mathcal{X}_{\xi,Q}^{a,v}(u; t) = Q_a q_a^v(u) \prod_{i=1}^n \frac{1}{u - z_i} \prod_{i=1}^{\xi^{a-1}} \frac{u - t_i^{a-1} + 1}{u - t_i^{a-1}} \prod_{i=1}^{\xi^a} \frac{u - t_i^a - 1}{u - t_i^a}.$$

Similarly to Theorem 6.1, we can show that for any off-diagonal solution \tilde{t} of the system

$$\begin{aligned} & Q_a q_a^v(t_i^a) \prod_{j=1}^{\xi^{a-1}} (t_i^a - t_j^{a-1} + 1) \prod_{\substack{j=1 \\ j \neq i}}^{\xi^a} (t_i^a - t_j^a - 1) \prod_{j=1}^{\xi^{a+1}} (t_i^a - t_j^{a+1}) = \\ & = Q_{a+1} q_{a+1}^v(t_i^a) \prod_{j=1}^{\xi^{a-1}} (t_i^a - t_j^{a-1}) \prod_{\substack{j=1 \\ j \neq i}}^{\xi^a} (t_i^a - t_j^a + 1) \prod_{j=1}^{\xi^{a+1}} (t_i^a - t_j^{a+1} - 1), \end{aligned}$$

$a = 1, \dots, N-1$, $i = 1, \dots, \xi^a$, where $\xi^0 = \xi^N = 0$, we have

$$\mathcal{T}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u; z) \mathbb{B}_\xi^v(\tilde{t}; z) = \mathbb{B}_\xi^v(\tilde{t}; z) \sum_{\mathbf{a}} \prod_{r=1}^k \mathcal{X}_{\xi,Q}^{a_r, v}(u - r + 1; \tilde{t}).$$

Moreover, for any t ,

$$\text{qdet } T(u) \mathbb{B}_\xi^v(t; z) = \mathbb{B}_\xi^v(t; z) \prod_{j=0}^{N-1} \left(q_a^v(u - j) \prod_{i=1}^n \frac{1}{u - j - z_i} \right).$$

7. Shapovalov form

Define an anti-involution $\tau : U(\mathfrak{gl}_N) \rightarrow U(\mathfrak{gl}_N)$ by the rule $\tau(e_{ab}) = e_{ba}$, $a, b = 1, \dots, N$. Let M be a highest weight \mathfrak{gl}_N -module with highest weight vector v . The *Shapovalov form* $S_M : M \otimes M \rightarrow \mathbb{C}$ is the unique symmetric bilinear form such that $S_M(v, v) = 1$, and $S_M(Xw_1, w_2) = S_M(w_1, \tau(X)w_2)$ for any $X \in U(\mathfrak{gl}_N)$ and any $w_1, w_2 \in M$. For an irreducible module M , the form S_M is nondegenerate.

Lemma 7.1. *For any $X \in Y(\mathfrak{gl}_N)$, $w_1, w_2 \in M$, and $x \in \mathbb{C}$, we have*

$$S_M(X|_{M(x)} w_1, w_2) = S_M(w_1, \varpi(X)|_{M(x)} w_2).$$

The proof is straightforward.

Let M_1, \dots, M_n be highest weight \mathfrak{gl}_N -modules. Then formula (4.8) and Lemma 7.1 imply that for any $X \in Y(\mathfrak{gl}_N)$ and any $w_1, w_2 \in M_1 \otimes \dots \otimes M_n$, we have

$$\begin{aligned} (7.1) \quad S_{M_1} \otimes \dots \otimes S_{M_n} (X|_{M_1(z_1) \otimes \dots \otimes M_n(z_n)} w_1, w_2) &= \\ &= S_{M_1} \otimes \dots \otimes S_{M_n} \left(w_1, (\varpi(X)|_{M_n(z_n) \otimes \dots \otimes M_1(z_1)})^{(n \dots 1)} w_2 \right), \end{aligned}$$

where S_{M_i} is the Shapovalov form for the module M_i , $i = 1, \dots, n$.

Proposition 7.2. *For any $k = 1, \dots, N$, and any $w_1, w_2 \in M_1 \otimes \dots \otimes M_n$, we have*

$$\begin{aligned} S_{M_1} \otimes \dots \otimes S_{M_n} (\mathcal{T}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u; z_1, \dots, z_n) w_1, w_2) &= \\ &= S_{M_1} \otimes \dots \otimes S_{M_n} \left(w_1, (\mathcal{T}_{k,Q^\dagger}^{M_n \otimes \dots \otimes M_1}(u; z_n, \dots, z_1))^{(n \dots 1)} w_2 \right), \end{aligned}$$

where the superscript \dagger stands for the transposition of matrices.

Proof. The claim follows from formulae (6.1), (7.1) and Proposition 4.11. \square

We remind below some facts on R -matrices for tensor products of evaluation modules over the Yangian $Y(\mathfrak{gl}_N)$. Propositions 7.3–7.5 follow from Drinfeld's results on the universal R -matrices for Yangians [D].

Proposition 7.3. *Let L and M be highest weight \mathfrak{gl}_N -modules with the respective highest weight vectors v and w . For generic $x, y \in \mathbb{C}$ there exists a unique operator $R_{LM}(x - y) \in \text{End}(L \otimes M)$ such that*

$$(7.2) \quad R_{LM}(x - y) X|_{L(x) \otimes M(y)} = (X|_{M(y) \otimes L(x)})^{(21)} R_{LM}(x - y)$$

for any $X \in Y(\mathfrak{gl}_N)$, and

$$(7.3) \quad R_{LM}(u) v \otimes w = v \otimes w.$$

The operator $R_{LM}(u)$ is called the *rational R-matrix* for the tensor product $L \otimes M$.

Example. Let $L = V^{\wedge l}$ and $M = V^{\wedge m}$. Then

$$R_{LM}(u) = R^{\wedge l, \wedge m}(u) \frac{u + \max(m - l, 0)}{u + m} \prod_{i=0}^{l-1} \prod_{j=0}^{m-1} \frac{1}{u + j - i}.$$

Proposition 7.4. $R_{LM}(u) = ((R_{ML}(-u))^{(21)})^{-1}$.

Proposition 7.5. *Let M_1, M_2, M_3 be highest weight \mathfrak{gl}_N -modules. Then the operators $R_{M_i M_j}(u)$, $1 \leq i < j \leq 3$, satisfy the Yang-Baxter equation*

$$R_{M_1 M_2}^{(12)}(u_1 - u_2) R_{M_1 M_3}^{(13)}(u_1) R_{M_2 M_3}^{(23)}(u_2) = R_{M_2 M_3}^{(23)}(u_2) R_{M_1 M_3}^{(13)}(u_1) R_{M_1 M_2}^{(12)}(u_1 - u_2).$$

By the definition of evaluation modules over $Y(\mathfrak{gl}_N)$, formula (7.2) means that the operator $R_{LM}(u)$ commutes with the \mathfrak{gl}_N -action in $L \otimes M$ and for any $a, b = 1, \dots, N$,

$$R_{LM}(u) \left(u e_{ab} \otimes 1 + \sum_{c=1}^N e_{ac} \otimes e_{cb} \right) = \left(u e_{ab} \otimes 1 + \sum_{c=1}^N e_{cb} \otimes e_{ac} \right) R_{LM}(u).$$

The operator $R_{LM}(u)$ is uniquely determined by these properties together with the normalization condition (7.3), which implies that the restriction of $R_{LM}(u)$ to any weight subspace of the \mathfrak{gl}_N -module $L \otimes M$ is a rational function of u , and

$$(7.4) \quad R_{LM}(u) = 1 + O(u^{-1}), \quad u \rightarrow \infty.$$

In addition, for any $w_1, w_2 \in L \otimes M$, we have

$$(7.5) \quad S_L \otimes S_M (R_{LM}(u) w_1, w_2) = S_L \otimes S_M (w_1, R_{LM}(u) w_2).$$

Let M_1, \dots, M_n be highest weight \mathfrak{gl}_N -modules. Set

$$(7.6) \quad R_{M_1 \dots M_n}(z) = \prod_{1 \leq i < j \leq n}^{\rightarrow} R_{M_i M_j}^{(ij)}(z_i - z_j) \in \text{End}(M_1 \otimes \dots \otimes M_n).$$

By Proposition 7.4 we have

$$(R_{M_1 \dots M_n}(z))^{-1} = \prod_{1 \leq i < j \leq k}^{\leftarrow} R_{M_j M_i}^{(ji)}(z_j - z_i).$$

For any $w_1, w_2 \in M_1 \otimes \dots \otimes M_n$, formula (7.5) and Proposition 7.5 yield

$$(7.7) \quad S_{M_1} \otimes \dots \otimes S_{M_n} (R_{M_1 \dots M_n}(z) w_1, w_2) = S_{M_1} \otimes \dots \otimes S_{M_n} (w_1, R_{M_1 \dots M_n}(z) w_2),$$

and for any $X \in Y(\mathfrak{gl}_N)$, formula (7.2) implies

$$(7.8) \quad R_{M_1 \dots M_n}(z) X|_{M_1(z_1) \otimes \dots \otimes M_n(z_n)} = (X|_{M_n(z_n) \otimes \dots \otimes M_1(z_1)})^{(n \dots 1)} R_{M_1 \dots M_n}(z).$$

Define a bilinear form $S_{M_1 \otimes \dots \otimes M_n}^z$ on $M_1 \otimes \dots \otimes M_n$ by the rule

$$(7.9) \quad S_{M_1 \otimes \dots \otimes M_n}^z(w_1, w_2) = (S_{M_1} \otimes \dots \otimes S_{M_n})(w_1, R_{M_1 \dots M_n}(z) w_2).$$

For irreducible modules M_1, \dots, M_n and generic z , the form $S_{M_1 \otimes \dots \otimes M_n}^z$ is nondegenerate. By equality (7.7), the form $S_{M_1 \otimes \dots \otimes M_n}^z$ is symmetric.

Theorem 7.6. *For a symmetric matrix Q , $Q = Q^\dagger$, the operators $\mathcal{T}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u; z)$, $k = 1, \dots, N$, are symmetric with respect to the symmetric bilinear form $S_{M_1 \otimes \dots \otimes M_n}^z$.*

Proof. Proposition 7.2 and formula (7.8) yield

$$S_{M_1 \otimes \dots \otimes M_n}^z(\mathcal{T}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u; z) w_1, w_2) = S_{M_1 \otimes \dots \otimes M_n}^z(w_1, \mathcal{T}_{k,Q^\dagger}^{M_1 \otimes \dots \otimes M_n}(u; z) w_2)$$

for any $k = 1, \dots, N$. This proves the statement. \square

8. Current algebra $\mathfrak{gl}_N[x]$

Let $\mathfrak{gl}_N[x]$ be the Lie algebra of polynomials with coefficients in \mathfrak{gl}_N with the pointwise commutator. We call it the *current algebra*. In this section we construct, following [T], analogues of the transfer matrices $\mathcal{T}_{1,Q}(u), \dots, \mathcal{T}_{N,Q}(u)$ for the algebra $U(\mathfrak{gl}_N[x])$, and formulate the counterparts of Theorems 5.4 and 6.1.

We will be using the standard generators of $\mathfrak{gl}_N[x]$, $e_{ab}^{\{s\}} = e_{ab} x^s$, $a, b = 1, \dots, N$, and $s = 0, 1, \dots$, which commute as follows: $[e_{ab}^{\{r\}}, e_{cd}^{\{s\}}] = \delta_{bc} e_{ad}^{\{r+s\}} - \delta_{ad} e_{cb}^{\{r+s\}}$. We will identify the Lie algebra \mathfrak{gl}_N with the subalgebra in $\mathfrak{gl}_N[x]$ of constant polynomials, thus making no difference between the elements $e_{ab}^{\{0\}}$ and the generators e_{ab} of \mathfrak{gl}_N .

Let $\mathfrak{n}_+ = \bigoplus_{1 \leq a < b \leq N} \mathbb{C} e_{ab}$ and $\mathfrak{n}_- = \bigoplus_{1 \leq a < b \leq N} \mathbb{C} e_{ba}$ be the standard nilpotent subalgebras in \mathfrak{gl}_N , and let $\mathfrak{n}_\pm[x]$ be the corresponding subalgebras in $\mathfrak{gl}_N[x]$. If $A, B \in U(\mathfrak{gl}_N[x])$ and $A - B \in U(\mathfrak{gl}_N[x]) \mathfrak{n}_+[x]$, then we will write $A \simeq B$.

A vector v in a $\mathfrak{gl}_N[x]$ -module is called *singular* with respect to the action of $\mathfrak{gl}_N[x]$ if $\mathfrak{n}_+[x]v = 0$. A singular vector v that is an eigenvector for the action of $e_{aa}^{\{s\}}$, $a = 1, \dots, N$, $s = 0, 1, \dots$, is called a *weight singular vector*.

Let M be a \mathfrak{gl}_N -module. For any $y \in \mathbb{C}$ denote by $M(y)$ the corresponding evaluation $\mathfrak{gl}_N[x]$ -module, that is, $e_{ab}^{\{s\}}|_{M(y)} = y^s e_{ab}|_M$, $a, b = 1, \dots, N$, $s = 0, 1, \dots$.

We organize elements $e_{ab}^{\{s\}}$, $a, b = 1, \dots, N$, $s = 0, 1, \dots$, into generating series

$$(8.1) \quad L_{ab}(u) = \sum_{s=0}^{\infty} e_{ba}^{\{s\}} u^{-s-1}, \quad a, b = 1, \dots, N.$$

and combine those series together into a series $L(u) = \sum_{a,b=1}^N E_{ab} \otimes L_{ab}(u)$ with coefficients in $\text{End}(\mathbb{C}^N) \otimes U(\mathfrak{gl}_N[x])$. Notice the flip of subscripts of the generators in the sum (8.1); this is done to have better notational correspondence further with the Yangian case. The commutation relations in $U(\mathfrak{gl}_N[x])$ can be written in the following form:

$$(u-v)[L_{ab}(u), L_{cd}(v)] = \delta_{bc}(L_{ad}(u) - L_{ad}(v)) - \delta_{ad}(L_{cb}(u) - L_{cb}(v)),$$

that amounts to the equality for series with coefficients in $\text{End}(\mathbb{C}^N) \otimes \text{End}(\mathbb{C}^N) \otimes U(\mathfrak{gl}_N[x])$:

$$(8.2) \quad (u-v)[L^{(13)}(u), L^{(23)}(v)] = [L^{(13)}(u) + L^{(23)}(v), P^{(12)}].$$

Here $P \in \text{End}(\mathbb{C}^N) \otimes \text{End}(\mathbb{C}^N)$ is the flip map.

Fix a matrix $K \in \text{End}(V)$ and consider the formal differential operator

$$(8.3) \quad \mathcal{D}_K(u, \partial_u) = (\text{tr}_{V \otimes N} \otimes \text{id}) \left((\partial_u - K^{(1)} - L^{(1, N+1)}(u)) \dots (\partial_u - K^{(N)} - L^{(N, N+1)}(u)) \mathbb{A}_{\{N\}}^{(1 \dots N)} \right).$$

of order N . Let $\mathcal{G}_{0,K}(u), \dots, \mathcal{G}_{N,K}(u) \in U(\mathfrak{gl}_N[x])[u^{-1}]$ be its coefficients:

$$(8.4) \quad \mathcal{D}_K(u, \partial_u) = \sum_{k=0}^N (-1)^k \mathcal{G}_{k,K}(u) \partial_u^{N-k}.$$

The series $\mathcal{G}_{1,K}(u), \dots, \mathcal{G}_{N,K}(u)$ are called the *Gaudin transfer matrices*.

Example. Clearly, $\mathcal{G}_{0,K}(u) = 1$ because $\text{tr}_{V \otimes N} \mathbb{A}_{\{N\}} = 1$. Let $K = \sum_{a,b=1}^N K_{ab} E_{ab}$. We also have that

$$(8.5) \quad \begin{aligned} \mathcal{G}_{1,K}(u) &= (\text{tr}_V \otimes \text{id})(K^{(1)} + L^{(12)}(u)) = \sum_{a=1}^N (K_{aa} + L_{aa}(u)), \\ \mathcal{G}_{2,K}(u) &= (\text{tr}_{V \otimes 2} \otimes \text{id}) \left(((K^{(1)} + L^{(13)}(u))(K^{(2)} + L^{(23)}(u)) - \dot{L}^{(23)}(u)) \mathbb{A}_{\{2\}}^{(12)} \right) = \\ &= \frac{1}{2} \left((\mathcal{G}_{1,K}(u))^2 - (N-1) \dot{\mathcal{G}}_{1,K}(u) - \sum_{a,b=1}^N (K_{ab} + L_{ab}(u))(K_{ba} + L_{ba}(u)) \right), \end{aligned}$$

where $\dot{G} = dG/du$.

The next two propositions were obtained in [T] from the Yangian case. We will give their proofs at the end of Section 10.

Proposition 8.1. *We have*

$$\mathbb{A}_{\{N\}}^{(1\dots N)} \mathcal{D}_K^{(N+1)}(u, \partial_u) = (\partial_u - K^{(1)} - L^{(1, N+1)}(u)) \dots (\partial_u - K^{(N)} - L^{(N, N+1)}(u)) \mathbb{A}_{\{N\}}^{(1\dots N)}.$$

Proposition 8.2. *We have $[\mathcal{G}_{k,K}(u), \mathcal{G}_{l,K}(v)] = 0$.*

Proposition 8.3. *Let K be the zero matrix. Then for any $k = 1, \dots, N$, the coefficients of $\mathcal{G}_{k,K}(u)$ commute with the subalgebra $U(\mathfrak{gl}_N) \subset U(\mathfrak{gl}_N[x])$.*

Proof. Similarly to (4.6), for any $a, b = 1, \dots, N$, we have $[E_{ab} \otimes 1 + 1 \otimes e_{ab}, L(u)] = 0$. Hence, the claim follows from formulae (8.3), (8.4), and the cyclic property of the trace, because $[E_{ab}^{(1)} + \dots + E_{ab}^{(N)}, \mathbb{A}_{\{N\}}^{(1\dots N)}] = 0$. \square

We extend the anti-involution $\tau : U(\mathfrak{gl}_N) \rightarrow U(\mathfrak{gl}_N)$ to the algebra $U(\mathfrak{gl}_N[x])$ by the rule

$$(8.6) \quad \tau(e_{ab}^{\{s\}}) = e_{ba}^{\{s\}}, \quad a, b = 1, \dots, N, \quad s = 0, 1, \dots$$

The extension is consistent with the standard embedding $U(\mathfrak{gl}_N) \hookrightarrow U(\mathfrak{gl}_N[x])$.

Proposition 8.4. *We have $\tau(\mathcal{G}_{k,K}(u)) = \mathcal{G}_{k,K^\dagger}(u)$, where the superscript \dagger stands for the transposition of matrices.*

The statement is proved in Section 10.

Let $\xi = (\xi^1, \dots, \xi^{N-1})$ be a collection of nonnegative integers. Set $|\xi| = \xi^1 + \dots + \xi^{N-1}$. Consider an expression in $|\xi|$ variables $t_1^1, \dots, t_{\xi^1}^1, \dots, t_1^{N-1}, \dots, t_{\xi^{N-1}}^{N-1}$ with coefficients in $U(\mathfrak{n}_-[x])$:

$$(8.7) \quad \mathbb{F}_\xi(t_1^1, \dots, t_{\xi^{N-1}}^{N-1}) = \\ = \text{Sym}_{t_1^1, \dots, t_{\xi^1}^1} \dots \text{Sym}_{t_1^{N-1}, \dots, t_{\xi^{N-1}}^{N-1}} \left\{ \sum_{\mathbf{m}} \prod_{1 \leq a < b \leq N} \left(\frac{1}{m_{ab}!} \prod_{i=1}^{m_{ab}} L_{ab}(t_{\widehat{m}_{ab}^a+i}^a) \times \right. \right. \\ \left. \left. \times \prod_{c=a}^{b-2} \frac{1}{t_{\widehat{m}_{ab}^{c+1}+i}^{c+1} - t_{\widehat{m}_{ab}^c+i}^c} \right) \right\},$$

where the sum is taken over all arrays of nonnegative integers $\mathbf{m} = (m_{ab})_{1 \leq a < b \leq N}$ such that $\sum_{c=1}^a \sum_{b=a+1}^N m_{cb} = \xi^a$, and $\widehat{m}_{ab}^c = \sum_{\substack{(r,s) < (a,b) \\ s > c}} m_{rs}$.

Lemma 8.5. *The product $\mathbb{F}_\xi(t_1^1, \dots, t_{\xi^{N-1}}^{N-1}) \prod_{a=1}^{N-2} \prod_{i=1}^{\xi^a} \prod_{j=1}^{\xi^{a+1}} (t_j^{a+1} - t_i^a)$ is a power series in $(t_1^1)^{-1}, \dots, (t_{\xi^{N-1}}^{N-1})^{-1}$ with coefficients in $U(\mathfrak{n}_-[x])$.*

The proof is straightforward.

Let K be a diagonal matrix, $K = \sum_{a=1}^N K_a E_{aa}$. Introduce the expressions

$$(8.8) \quad \mathcal{J}_{\xi,K}^{a,i}(t) = K_a - K_{a+1} + L_{aa}(t_i^a) - L_{a+1,a+1}(t_i^a) + \\ + \sum_{j=1}^{\xi^{a-1}} \frac{1}{t_i^a - t_j^{a-1}} - 2 \sum_{\substack{j=1 \\ j \neq i}}^{\xi^a} \frac{1}{t_i^a - t_j^a} + \sum_{j=1}^{\xi^{a+1}} \frac{1}{t_i^a - t_j^{a+1}},$$

$a = 1, \dots, N-1$, $i = 1, \dots, \xi^a$. Here we set $\xi^0 = \xi^N = 0$.

Denote $L_{ab}^{[r]}(u) = \frac{d^r}{du^r} L_{ab}(u)$. Given the following data:

- i) an integer m not exceeding $|\xi| + k - 1$, integers $a_1, \dots, a_m, b_1, \dots, b_m, c \in \{1, \dots, N\}$ and $i \in \{1, \dots, \xi^c\}$,
- ii) a sequence s_1, \dots, s_m , where each s_j is either u or one of the variables $t_1^1, \dots, \widehat{t_i^c}, \dots, t_{\xi^{N-1}}^{N-1}$, where the variable t_i^c is excluded,
- iii) nonnegative integers r_1, \dots, r_m such that $r_j = 0$ if $s_j \neq u$, and $\sum_{s_j=u} r_j \leq k$,
- iv) a polynomial $p(u; t_1^1, \dots, t_{\xi^{N-1}}^{N-1})$,

consider the product

$$(8.9) \quad L_{a_1 b_1}^{[r_1]}(s_1) \dots L_{a_m b_m}^{[r_m]}(s_m) \mathcal{J}_{\xi,K}^{c,i}(t) \times \\ \times p(u; t_1^1, \dots, t_{\xi^{N-1}}^{N-1}) \prod_{a=1}^{N-1} \prod_{j=1}^{\xi^a} \frac{1}{(u - t_j^a)^k} \prod_{a=1}^{N-2} \prod_{j=1}^{\xi^a} \prod_{l=1}^{\xi^{a+1}} \frac{1}{t_l^{a+1} - t_j^a}$$

where the factors $(u - t_j^a)^{-k}$ in (8.9) have to be expanded as power series in u^{-1} . We denote by $J_{\xi,k,K}$ the span over \mathbb{C} of all products (8.9) with various m , a_1, \dots, a_m , b_1, \dots, b_m , r_1, \dots, r_m , c, i , s_1, \dots, s_m , and $p(u; t_1^1, \dots, t_{\xi^{N-1}}^{N-1})$.

Consider the formal differential operator

$$\mathcal{M}_{\xi,K}(u, \partial_u; t) = \overrightarrow{\prod}_{1 \leq a \leq N} \left(\partial_u - K_a - L_{aa}(u) - \sum_{i=1}^{\xi^{a-1}} \frac{1}{u - t_i^{a-1}} + \sum_{i=1}^{\xi^a} \frac{1}{u - t_i^a} \right).$$

Let $\mathcal{Z}_{\xi,1,K}(u; t), \dots, \mathcal{Z}_{\xi,N,K}(u; t)$ be its coefficients:

$$\mathcal{M}_{\xi,K}(u, \partial_u; t) = \sum_{k=0}^N (-1)^k \mathcal{Z}_{\xi,k,K}(u; t) \partial_u^{N-k}.$$

Expressions $\mathcal{Z}_{\xi,1,K}(u; t), \dots, \mathcal{Z}_{\xi,N,K}(u; t)$ are regarded as power series in u^{-1} with coefficients in $U(\mathfrak{gl}_N[x])[t_1^1, \dots, t_{\xi^{N-1}}^{N-1}]$.

Theorem 8.6. *Let K be a diagonal matrix. For any $k = 1, \dots, N$, we have*

$$(8.10) \quad \mathcal{G}_{k,K}(u) \mathbb{F}_{\xi}(t) \simeq \mathbb{F}_{\xi}(t) \mathcal{Z}_{\xi,k,K}(u; t) + \mathcal{W}_{\xi,k,K}(u; t),$$

where $\mathcal{W}_{\xi,k,K}(u; t)$ belongs to $J_{\xi,k,K}$.

The theorem is proved in Section 12.

Remark. It is easy to see that the coefficients of the series $\mathcal{G}_{1,K}(u)$ are central in $U(\mathfrak{gl}_N[x])$ and $\mathcal{Z}_{\xi,1,K}(u;t) = \mathcal{G}_{1,K}(u)$ for any ξ . This implies Theorem 8.6 for $k = 1$.

9. Tensor products of evaluation modules of the current algebra

Let M_1, \dots, M_n be \mathfrak{gl}_N -modules. Consider the tensor product $M_1(z_1) \otimes \dots \otimes M_n(z_n)$ of evaluation $\mathfrak{gl}_N[x]$ -modules. Then

$$(9.1) \quad L_{ab}(u) \Big|_{M_1(z_1) \otimes \dots \otimes M_n(z_n)} = \sum_{i=1}^n \frac{e_{ba}^{(i)}}{u - z_i}, \quad a = 1, \dots, N,$$

where we consider $e_{ba}^{(i)} = 1^{\otimes(i-1)} \otimes e_{ab} \otimes 1^{\otimes(n-i)}$ as an operator acting in $M_1 \otimes \dots \otimes M_n$, cf. (8.1). By formulae (8.3), (8.4), for any $k = 1, \dots, n$, the operator

$$(9.2) \quad \mathcal{G}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z) = \mathcal{G}_{k,K}(u) \Big|_{M_1(z_1) \otimes \dots \otimes M_n(z_n)}$$

is a rational function in u, z_1, \dots, z_n with the denominator $\prod_{i=1}^n (u - z_i)^k$ and

$$\mathcal{G}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z) = \text{tr } K^{\wedge k} + O(u^{-1}), \quad u \rightarrow \infty.$$

The operators $\mathcal{G}_{1,K}^{M_1 \otimes \dots \otimes M_n}(u; z), \dots, \mathcal{G}_{N,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$ are called the *transfer matrices of the Gaudin model* associated with the Lie algebra \mathfrak{gl}_N .

Let τ be the anti-involution (8.6) on $U(\mathfrak{gl}_N[x])$. Recall that for a highest weight \mathfrak{gl}_N module M we denote by S_M the Shapovalov form on M .

Let M_1, \dots, M_n be highest weight \mathfrak{gl}_N -modules. For any z_1, \dots, z_n , $X \in U(\mathfrak{gl}_N[x])$, and $w_1, w_2 \in M_1 \otimes \dots \otimes M_n$, we have

$$\begin{aligned} S_{M_1} \otimes \dots \otimes S_{M_n} (X \Big|_{M_1(z_1) \otimes \dots \otimes M_n(z_n)} w_1, w_2) &= \\ &= S_{M_1} \otimes \dots \otimes S_{M_n} (w_1, \tau(X) \Big|_{M_1(z_1) \otimes \dots \otimes M_n(z_n)} w_2). \end{aligned}$$

Then Proposition 8.4 and formula (9.2) yield that for any $k = 1, \dots, N$,

$$S_{M_1} \otimes \dots \otimes S_{M_n} (\mathcal{G}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z) w_1, w_2) = S_{M_1} \otimes \dots \otimes S_{M_n} (w_1, \mathcal{G}_{k,K^\dagger}^{M_1 \otimes \dots \otimes M_n}(u; z) w_2).$$

In particular, we have the following result.

Theorem 9.1. *For a symmetric matrix K , $K = K^\dagger$, the operators $\mathcal{G}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$, $k = 1, \dots, N$, are symmetric with respect to the tensor product $S_{M_1} \otimes \dots \otimes S_{M_n}$ of the Shapovalov forms.*

Let M_1, \dots, M_n be highest weight \mathfrak{gl}_N -modules with highest weights $\Lambda_1, \dots, \Lambda_n$, $\Lambda_i = (\Lambda_i^1, \dots, \Lambda_i^N)$, and highest weight vectors v_1, \dots, v_n . Then for any z_1, \dots, z_n , the vector

$v_1 \otimes \dots \otimes v_n$ is a weight singular vector with respect to the action of $\mathfrak{gl}_N[x]$ in the module $M_1(z_1) \otimes \dots \otimes M_n(z_n)$ and

$$L_{aa}(u) v_1 \otimes \dots \otimes v_n = v_1 \otimes \dots \otimes v_n \sum_{i=1}^n \frac{\Lambda_i^a}{u - z_i}, \quad a = 1, \dots, N.$$

Fix a diagonal matrix $K = \sum_{a=1}^N K_a E_{aa}$. Let $\xi = (\xi^1, \dots, \xi^{N-1})$ be a collection of non-negative integers. Set $\xi^0 = \xi^N = 0$. The system of equations

$$(9.3) \quad \sum_{j=1}^n \frac{\Lambda_j^a - \Lambda_j^{a+1}}{t_i^a - z_j} + \sum_{j=1}^{\xi^{a-1}} \frac{1}{t_i^a - t_j^{a-1}} - 2 \sum_{\substack{j=1 \\ j \neq i}}^{\xi^a} \frac{1}{t_i^a - t_j^a} + \sum_{j=1}^{\xi^{a+1}} \frac{1}{t_i^a - t_j^{a+1}} = K_{a+1} - K_a,$$

$a = 1, \dots, N-1$, $i = 1, \dots, \xi^a$, is called the *Bethe ansatz equations*. We always assume that for a solution $\tilde{t} = (\tilde{t}_1^1, \dots, \tilde{t}_{\xi^{N-1}}^{N-1})$ of system (9.3) any denominator in equations does not equal zero unless the corresponding numerator equals zero.

Consider the differential operator

$$(9.4) \quad \mathcal{M}_{\xi, K}(u, \partial_u; t; z; \Lambda) = \prod_{1 \leq a \leq N}^{\rightarrow} \left(\partial_u - K_a - \sum_{i=1}^n \frac{\Lambda_i^a}{u - z_i} - \sum_{i=1}^{\xi^{a-1}} \frac{1}{u - t_i^{a-1}} + \sum_{i=1}^{\xi^a} \frac{1}{u - t_i^a} \right).$$

Let $\mathcal{Z}_{\xi, 1, K}(u; t; z; \Lambda), \dots, \mathcal{Z}_{\xi, N, K}(u; t; z; \Lambda)$ be its coefficients:

$$(9.5) \quad \mathcal{M}_{\xi, K}(u, \partial_u; t; z; \Lambda) = \sum_{k=0}^N (-1)^k \mathcal{Z}_{\xi, k, K}(u; t; z; \Lambda) \partial_u^{N-k}.$$

The expression (8.7) defines a vector-valued rational function $\mathbb{F}_{\xi}^{v_1 \otimes \dots \otimes v_n}(t; z)$ of the variables $t_1^1, \dots, t_{\xi^{N-1}}^{N-1}, z_1, \dots, z_n$,

$$(9.6) \quad \mathbb{F}_{\xi}^{v_1 \otimes \dots \otimes v_n}(t; z) = \mathbb{F}_{\xi}(t) v_1 \otimes \dots \otimes v_n,$$

with the denominator $\prod_{i=1}^n \prod_{a=1}^{N-1} \prod_{j=1}^{\xi^a} (t_j^a - z_i) \prod_{a=1}^{N-2} \prod_{i=1}^{\xi^a} \prod_{j=1}^{\xi^{a+1}} (t_j^{a+1} - t_i^a)$, cf. Lemma 8.5. The function $\mathbb{F}_{\xi}^{v_1 \otimes \dots \otimes v_n}(t; z)$ coincides with the *universal weight function* used in integral formulae for hypergeometric solutions of the Knizhnik-Zamolodchikov equations, see [Ma], [SV], [RSV].

Theorem 9.2. *Let M_1, \dots, M_n be highest weight \mathfrak{gl}_N -modules with highest weights $\Lambda_1, \dots, \Lambda_n$ and highest weight vectors v_1, \dots, v_n . Let K be a diagonal matrix and $\tilde{t} = (\tilde{t}_1^1, \dots, \tilde{t}_{\xi^{N-1}}^{N-1})$ a solution of system (9.3). Then for any $k = 1, \dots, N$, we have*

$$(9.7) \quad \mathcal{G}_{k, K}^{M_1 \otimes \dots \otimes M_n}(u; z) \mathbb{F}_{\xi}^{v_1 \otimes \dots \otimes v_n}(\tilde{t}; z) = \mathcal{Z}_{\xi, k, K}(u; \tilde{t}; z; \Lambda) \mathbb{F}_{\xi}^{v_1 \otimes \dots \otimes v_n}(\tilde{t}; z).$$

Proof. The statement follows from Theorem 8.6 and formulae (9.2), (9.6). \square

Remark. Since the coefficients of the series $\mathcal{G}_{1,K}(u)$ are central in $U(\mathfrak{gl}_N[x])$, formulae (8.5) and (9.4) yield

$$\mathcal{G}_{1,K}^{M_1 \otimes \dots \otimes M_n}(u; z) = \sum_{a=1}^N \left(K_a + \sum_{i=1}^n \frac{\Lambda_i^a}{u - z_i} \right) = \mathcal{Z}_{\xi,1,K}(u; t; z; \Lambda),$$

which implies Theorem 8.6 for $k = 1$.

Proposition 9.3. [RV] *Let K be the zero matrix, that is, $K_1 = \dots = K_N = 0$, and $\tilde{t} = (\tilde{t}_1^1, \dots, \tilde{t}_{\xi^{N-1}}^{N-1})$ an off-diagonal solution of the Bethe ansatz equations (9.3). Then the Bethe vector $\mathbb{F}_{\xi}^{v_1 \otimes \dots \otimes v_n}(\tilde{t}; z)$ is a singular vector with respect to the \mathfrak{gl}_N -action in $M_1 \otimes \dots \otimes M_n$ of weight $(\sum_{i=1}^n \Lambda_i^1 - \xi^1, \sum_{i=1}^n \Lambda_i^2 + \xi^1 - \xi^2, \dots, \sum_{i=1}^n \Lambda_i^N + \xi^{N-1})$.*

Proof. The claim follows from identities for the universal weight function [RSV] and quasi-classical asymptotics of hypergeometric solutions of the Knizhnik-Zamolodchikov equations [RV]. \square

Remark. Let M_1, \dots, M_n be any \mathfrak{gl}_N -modules, not necessarily highest weight ones, and let $v \in M_1 \otimes \dots \otimes M_n$ be any weight singular vector with respect to the action of $\mathfrak{gl}_N[x]$ in the module $M_1(z_1) \otimes \dots \otimes M_n(z_n)$. Define rational functions $r_1^v(u), \dots, r_N^v(u)$ by the rule $L_{aa}(u)v = r_a^v(u)v$, $a = 1, \dots, N$, and consider the differential operator

$$\mathcal{M}_{\xi,K}^v(u, \partial_u; t) = \prod_{1 \leq a \leq N} \left(\partial_u - K_a - r_a^v(u) - \sum_{i=1}^{\xi^{a-1}} \frac{1}{u - t_i^{a-1}} + \sum_{i=1}^{\xi^a} \frac{1}{u - t_i^a} \right).$$

Let $\mathcal{Z}_{\xi,1,K}^v(u; t; z), \dots, \mathcal{Z}_{\xi,N,K}^v(u; t; z)$ be its coefficients:

$$\mathcal{M}_{\xi,K}^v(u, \partial_u; t; z; \Lambda) = \sum_{k=0}^N (-1)^k \mathcal{Z}_{\xi,k,K}^v(u; t; z) \partial_u^{N-k}.$$

Similarly to Theorem 9.2, for any solution \tilde{t} of the system

$$r_a^v(t_i^a) + \sum_{j=1}^{\xi^{a-1}} \frac{1}{t_i^a - t_j^{a-1}} - 2 \sum_{\substack{j=1 \\ j \neq i}}^{\xi^a} \frac{1}{t_i^a - t_j^a} + \sum_{j=1}^{\xi^{a+1}} \frac{1}{t_i^a - t_j^{a+1}} = K_{a+1} - K_a,$$

$a = 1, \dots, N-1$, $i = 1, \dots, \xi^a$, we have

$$\mathcal{G}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z) \mathbb{F}_{\xi}(\tilde{t})v = \mathcal{Z}_{\xi,k,K}^v(u; \tilde{t}; z) \mathbb{F}_{\xi}(\tilde{t})v.$$

Moreover, for any ξ and t ,

$$\mathcal{G}_{1,K}^{M_1 \otimes \dots \otimes M_n}(u; z) \mathbb{F}_{\xi}(t)v = \mathcal{Z}_{\xi,1,K}^v(u; t; z) \mathbb{F}_{\xi}(t)v = \sum_{a=1}^N (K_a + r_a^v(u)) \mathbb{F}_{\xi}(t)v.$$

10. Filtration on $Y(\mathfrak{gl}_N)$

In this section we are going to relate the series $\mathcal{T}_{1,Q}(u), \dots, \mathcal{T}_{N,Q}(u)$ with coefficients in the Yangian $Y(\mathfrak{gl}_N)$ and the series $\mathcal{G}_{1,K}(u), \dots, \mathcal{G}_{N,K}(u)$ with coefficients in the algebra $U(\mathfrak{gl}_N[x])$, and to prove Propositions 8.1–8.4. The results of this section are essentially borrowed from [T].

Given a filtered algebra \mathcal{A} with an ascending filtration $\dots \subset \mathcal{A}_{s-1} \subset \mathcal{A}_s \subset \dots \subset \mathcal{A}$, possibly double infinite, we denote by $\text{grad}_s^{\mathcal{A}} : \mathcal{A}_s \rightarrow \mathcal{A}_s/\mathcal{A}_{s-1}$ the canonical projection. We will identify the quotient spaces $\mathcal{A}_s/\mathcal{A}_{s-1}$ with the corresponding homogeneous subspaces in the algebra

$$\text{grad } \mathcal{A} = \bigoplus_{r \in \mathbb{Z}} \mathcal{A}_r/\mathcal{A}_{r-1}$$

and will regard $\text{grad}_s^{\mathcal{A}}$ as a map from \mathcal{A}_s to $\text{grad } \mathcal{A}$. Abusing notation, we will write just grad_s , dropping the superscript, if it can cause no confusion.

For a filtered algebra \mathcal{A} , we consider the algebra $\text{End}(V) \otimes \mathcal{A}$ to be filtered by the subspaces $(\text{End}(V) \otimes \mathcal{A})_s = \text{End}(V) \otimes \mathcal{A}_s$.

The Yangian $Y(\mathfrak{gl}_N)$ admits a degree function such that $\deg T_{ab}^{\{s\}} = s - 1$ for any $a, b = 1, \dots, N$, $s = 1, 2, \dots$, see Section 1.20 in [MNO]; the function \deg here coincides with the function \deg_2 in [MNO]. Then $Y(\mathfrak{gl}_N)$ is a filtered algebra with an ascending filtration $Y(\mathfrak{gl}_N)_0 \subset Y(\mathfrak{gl}_N)_1 \subset \dots \subset Y(\mathfrak{gl}_N)$, where the subspace $Y(\mathfrak{gl}_N)_s$ contains elements of degree not greater than s .

Theorem 10.1. [MNO, Theorem 1.26] *We have $\text{grad}(Y(\mathfrak{gl}_N)) = U(\mathfrak{gl}_N[x])$.*

Moreover, $\text{grad}_{s-1}(T_{ab}^{\{s\}}) = e_{ba}^{\{s\}}$, $a, b = 1, \dots, N$, $s = 1, 2, \dots$.

The anti-involution ϖ is compatible with the filtration on $Y(\mathfrak{gl}_N)$. For any $X \in Y(\mathfrak{gl}_N)$ we have $\deg \varpi(X) = \deg X$, and if $\deg X = r$, then

$$(10.1) \quad \text{grad}_r(\varpi(X)) = \tau(\text{grad}_r X).$$

We extend the filtration from the Yangian $Y(\mathfrak{gl}_N)$ to the algebra $Y(\mathfrak{gl}_N)[[u^{-1}, \partial_u]]$ by the rule $\deg u^{-1} = \deg \partial_u = -1$. Notice that the indeterminates u^{-1} and ∂_u do not commute obeying the following relation $\partial_u \cdot u^{-s} = u^{-s} \partial_u - s u^{-s-1}$ instead. Clearly,

$$\text{grad}(Y(\mathfrak{gl}_N)[[u^{-1}, \partial_u]]) = U(\mathfrak{gl}_N[x])[u^{-1}, \partial_u].$$

It is easy to see that the series $T_{ab}(u) - \delta_{ab} \in Y(\mathfrak{gl}_N)[[u^{-1}, \partial_u]]$ has degree -1 and

$$(10.2) \quad \text{grad}_{-1}(T_{ab}(u) - \delta_{ab}) = L_{ab}(u), \quad a, b = 1, \dots, N.$$

Further on we assume that in the definition (4.13) of the transfer matrices Q is a series from $\text{End}(V)[[\zeta]]$ instead of being an element of $\text{End}(V)$. Then the obtained transfer matrices $\mathcal{T}_{1,Q}(u), \dots, \mathcal{T}_{N,Q}(u)$ are power series in u^{-1} and ζ with coefficients in $Y(\mathfrak{gl}_N)$. We also regard them as elements of $Y(\mathfrak{gl}_N)[[u^{-1}, \partial_u, \zeta]]$. The results of Sections 4 and 5 naturally generalize to the described setting. We extend the filtration from $Y(\mathfrak{gl}_N)[[u^{-1}, \partial_u]]$

to $Y(\mathfrak{gl}_N)[[u^{-1}, \partial_u, \zeta]]$ by setting $\deg \zeta = -1$. Similarly, in formulae (8.3), (8.4) we will assume K being a series from $\text{End}(V)[[\zeta]]$.

Recall that $\mathcal{T}_{0,Q}(u) = 1$ by convention. For any $k = 0, \dots, N$, set

$$(10.3) \quad \mathcal{S}_{k,Q}(u) = \frac{1}{(N-k)!} \sum_{l=0}^k (-1)^{k-l} \frac{(N-l)!}{(k-l)!} \mathcal{T}_{l,Q}(u).$$

For example, $\mathcal{S}_{0,Q}(u) = 1$ and $\mathcal{S}_{1,Q}(u) = \mathcal{T}_{1,Q}(u) - N$. Formulae (10.3) for all $k = 0, \dots, N$ taken together are equivalent to the identity

$$(10.4) \quad \sum_{k=0}^N (-1)^k \mathcal{S}_{k,Q}(u) y^{N-k} = \sum_{l=0}^N (-1)^l \mathcal{T}_{l,Q}(u) (y+1)^{N-l}.$$

The series $\mathcal{S}_{1,Q}(u), \dots, \mathcal{S}_{N,Q}(u)$ are called the *modified transfer matrices*.

Theorem 10.2. [T] *Let $\deg(Q-1) \leq -1$. Then $\deg(\mathcal{S}_{k,Q}(u)) = -k$ for any $k = 0, \dots, N$. Moreover,*

$$(10.5) \quad \text{grad}_{-k}(\mathcal{S}_{k,Q}(u)) = \mathcal{G}_{k,K}(u)$$

where $K = \text{grad}_{-1}(Q-1)$.

Proof. Taking the $(N-m)$ -th derivative of formula (10.4) with respect to y and making there the substitution $y = e^{\partial_u} - 1$, we get

$$(10.6) \quad \sum_{k=0}^m (-1)^k \frac{(N-k)!}{(m-k)!} \mathcal{S}_{k,Q}(u) (e^{\partial_u} - 1)^{m-k} = \sum_{l=0}^m (-1)^l \frac{(N-l)!}{(m-l)!} \mathcal{T}_{l,Q}(u) e^{(m-l)\partial_u} = \\ = (N-m)! \mathfrak{D}_{m,Q}(u, \partial_u) e^{m\partial_u},$$

cf. (4.15). Exploiting the standard identity

$$\sum_{r=0}^s \frac{(-1)^r}{r! (s-r)!} = 0, \quad s \geq 1,$$

we obtain that

$$(10.7) \quad \mathcal{S}_{k,Q}(u) = \frac{1}{(N-k)!} \sum_{m=0}^k (-1)^m \frac{(N-m)!}{(k-m)!} \mathfrak{D}_{m,Q}(u, \partial_u) e^{m\partial_u} (e^{\partial_u} - 1)^{k-m}.$$

Since $\deg(T(u) - 1) = -1$, $\deg(e^{\partial_u} - 1) = -1$, and $\deg(Q-1) \leq -1$, formula (4.14) implies that $\deg(\mathfrak{D}_{m,Q}(u, \partial_u)) = -m$. Hence, $\deg(\mathcal{S}_{k,Q}(u)) = -k$ by formula (10.7).

Observe that $\text{grad}_{-1}(T(u) - 1) = L(u)$, cf. (10.2), and $\text{grad}_{-1}(e^{\partial_u} - 1) = \partial_u$. Then computing $\text{grad}_{-m} \mathfrak{D}_{m,Q}(u, \partial_u)$ in two ways using either formula (4.14) or formula (10.6),

we get

$$\begin{aligned} (\mathrm{tr}_{V^{\otimes m}} \otimes \mathrm{id}) \left((\partial_u - K^{(1)} - L^{(1,m+1)}(u)) \dots (\partial_u - K^{(m)} - L^{(m,m+1)}(u)) \mathbb{A}_{\{m\}}^{(1\dots m)} \right) = \\ = \frac{1}{(N-m)!} \sum_{k=0}^m (-1)^k \frac{(N-k)!}{(m-k)!} \mathrm{grad}_{-k}(\mathbb{S}_{k,Q}(u)) \partial_u^{m-k}. \end{aligned}$$

The last step is to compare the obtained formula for $m = N$ with formulae (8.3) and (8.4), which completes the proof. \square

Proof of Propositions 8.1–8.4. Let $Q = 1 + \zeta K$. Propositions 4.5, 4.11 and 4.10 together with formulae (10.3), (10.5), (10.1) and the equality $\mathrm{grad}_{-N}(\mathfrak{D}_{N,Q}(u, \partial_u)) = \mathcal{D}_{\zeta K}(u, \partial_u)$ imply respectively Propositions 8.2, 8.4 and 8.1 with the matrix K replaced by ζK . Since all the expressions $\mathcal{G}_{1,\zeta K}(u), \dots, \mathcal{G}_{N,\zeta K}(u)$ and $\mathcal{D}_{\zeta K}(u, \partial_u)$ are polynomials in ζ , we can evaluate them at $\zeta = 1$, which completes the proof. \square

11. Proof of Theorem 5.4

We will prove Theorem 5.4 by induction with respect to N . The key points are Proposition 11.3 and 11.5, and formula (11.25).

Set $W = \mathbb{C}^{N-1}$. Let $\mathbf{w}_1, \dots, \mathbf{w}_{N-1}$ be the standard basis of W and $\mathbf{v}_1, \dots, \mathbf{v}_N$ the standard basis of $V = \mathbb{C}^N$. We identify W with its image in V under the embedding $\mathbf{w}_a \mapsto \mathbf{v}_{a+1}$, $a = 1, \dots, N-1$.

Let $P^{\langle N-1 \rangle} \in \mathrm{End}(W^{\otimes 2})$ be the flip map and $R^{\langle N-1 \rangle}(u) = u + P^{\langle N-1 \rangle}$ the rational R -matrix used in the definition of the Yangian $Y(\mathfrak{gl}_{N-1})$. The R -matrix $R(u)$ preserves the subspace $W^{\otimes 2} \subset V^{\otimes 2}$ and the restriction of $R(u)$ on $W^{\otimes 2}$ coincides with $R^{\langle N-1 \rangle}(u)$.

We will use the embedding $\psi : Y(\mathfrak{gl}_{N-1}) \hookrightarrow Y(\mathfrak{gl}_N) : \psi(T_{ab}^{\langle N-1 \rangle}(u)) = T_{a+1,b+1}(u)$, $a, b = 1, \dots, N-1$, where $T_{ab}^{\langle N-1 \rangle}(u)$ are series (4.1) for the algebra $Y(\mathfrak{gl}_{N-1})$. It is clear that $\psi(Y_+(\mathfrak{gl}_{N-1})) \subset Y_+(\mathfrak{gl}_N)$.

Let $W(x)$ be the evaluation $Y(\mathfrak{gl}_{N-1})$ -module and $\pi(x) : Y(\mathfrak{gl}_{N-1}) \rightarrow \mathrm{End}(W)$ be the corresponding homomorphism. In other words

$$(11.1) \quad \pi(x) : T^{\langle N-1 \rangle}(u) \mapsto (u - x)^{-1} R^{\langle N-1 \rangle}(u - x).$$

Introduce a map $\psi(x_1, \dots, x_r) : Y(\mathfrak{gl}_{N-1}) \rightarrow \mathrm{End}(W^{\otimes r}) \otimes Y(\mathfrak{gl}_N)$,

$$(11.2) \quad \psi(x_1, \dots, x_r) = (\pi(x_1) \otimes \dots \otimes \pi(x_r) \otimes \psi) \circ (\Delta^{\langle N-1 \rangle})^{(r)},$$

where $(\Delta^{\langle N-1 \rangle})^{(r)} : Y(\mathfrak{gl}_{N-1}) \rightarrow (Y(\mathfrak{gl}_{N-1}))^{\otimes(r+1)}$ is the multiple coproduct. For any $X \in \mathrm{End}(W)$ set $\nu(X) = X \mathbf{w}_1$. Define a map $\tilde{\psi}(x_1, \dots, x_r) : Y(\mathfrak{gl}_{N-1}) \rightarrow W^{\otimes r} \otimes Y(\mathfrak{gl}_N)$,

$$\tilde{\psi}(x_1, \dots, x_r) = (\nu^{\otimes r} \otimes \mathrm{id}) \circ \psi(x_1, \dots, x_r).$$

Lemma 11.1. $\tilde{\psi}(x_1, \dots, x_r)(Y_+(\mathfrak{gl}_{N-1})) \subset W^{\otimes r} \otimes Y_+(\mathfrak{gl}_N).$

Proof. The claim follows from the facts that $Y_+(\mathfrak{gl}_{N-1})$ is a coideal in $Y(\mathfrak{gl}_{N-1})$, and the vector \mathbf{w}_1 is a singular vector with respect to the action of $Y(\mathfrak{gl}_{N-1})$. \square

Consider the embedding $\phi : Y(\mathfrak{gl}_{N-2}) \hookrightarrow Y(\mathfrak{gl}_{N-1}) : \phi(T_{ab}^{\langle N-2 \rangle}(u)) = T_{a+1, b+1}^{\langle N-1 \rangle}(u)$, $a, b = 1, \dots, N-2$, similar to the embedding ψ . Recall that $\theta : Y(\mathfrak{gl}_N) \rightarrow Y(\mathfrak{gl}_N)/Y_+(\mathfrak{gl}_N)$ is the canonical projection.

Lemma 11.2. We have $(\text{id}^{\otimes r} \otimes \theta) \circ \tilde{\psi}(x_1, \dots, x_r) \circ \phi = \mathbf{w}_1^{\otimes r} \otimes (\theta \circ \psi \circ \phi).$

For any element $g \in W^{\otimes r} \otimes Y(\mathfrak{gl}_N)$ we define its components g^{a_1, \dots, a_r} by the rule

$$g = \sum_{a_1, \dots, a_r=1}^{N-1} \mathbf{w}_{a_1} \otimes \dots \otimes \mathbf{w}_{a_r} \otimes g^{a_1, \dots, a_r}.$$

Set $\bar{\xi} = (\xi^2, \dots, \xi^{N-1})$ and $\bar{t} = (t_1^2, \dots, t_{\xi^2}^2; \dots; t_1^{N-1}, \dots, t_{\xi^{N-1}}^{N-1}).$

Proposition 11.3. [TV1, Theorem 3.4.2] We have

$$(11.3) \quad \mathbb{B}_\xi(t) = \sum_{a_1, \dots, a_{\xi^1}=1}^{N-1} T_{1, a_1+1}(t_1^1) \dots T_{1, a_{\xi^1}+1}(t_{\xi^1}^1) \left(\tilde{\psi}(t_1^1, \dots, t_{\xi^1}^1) (\mathbb{B}_{\bar{\xi}}^{\langle N-1 \rangle}(\bar{t})) \right)^{a_1, \dots, a_{\xi^1}}.$$

Proof of Lemma 5.2. Lemma 11.2, Proposition 11.3 and formulae (11.1), (11.2) imply that the denominator of $\theta(\mathbb{B}_\xi(t))$ is at most $\prod_{a=1}^{N-2} \prod_{i=1}^{\xi^a} \prod_{j=1}^{\xi^{a+1}} (t_j^{a+1} - t_i^a)$. Now the statement follows from formula (5.2). \square

Since $V = \mathbb{C}\mathbf{v}_1 \oplus W$ we have $V^{\wedge k} = (\mathbf{v}_1 \wedge W^{\wedge(k-1)}) \oplus W^{\wedge k}$, where the first summand is spanned by the vectors $\mathbf{v}_1 \wedge \mathbf{v}_{a_1} \wedge \dots \wedge \mathbf{v}_{a_{k-1}}$, $2 \leq a_1 < \dots < a_{k-1} \leq N$, and the second summand is spanned by the vectors $\mathbf{v}_{b_1} \wedge \dots \wedge \mathbf{v}_{b_k}$, $2 \leq b_1 < \dots < b_k \leq N$. Abusing notation, we will often identify $W^{\wedge(k-1)}$ and $\mathbf{v}_1 \wedge W^{\wedge(k-1)}$ taking a vector \mathbf{x} to $\mathbf{v}_1 \wedge \mathbf{x}$.

The R -matrix $R_{\wedge k, \wedge 1}(u)$ acting in $V^{\wedge k} \otimes V$ preserves the subspaces

$$(\mathbf{v}_1 \wedge W^{\wedge(k-1)}) \otimes \mathbb{C}\mathbf{v}_1, \quad ((\mathbf{v}_1 \wedge W^{\wedge(k-1)}) \otimes W) \oplus (W^{\wedge k} \otimes \mathbb{C}\mathbf{v}_1), \quad W^{\wedge k} \otimes W.$$

Its restrictions onto these subspaces are

$$(11.4) \quad R_{\wedge k, \wedge 1}(u)|_{(\mathbf{v}_1 \wedge W^{\wedge(k-1)}) \otimes \mathbb{C}\mathbf{v}_1} = u + 1, \quad R_{\wedge k, \wedge 1}(u)|_{W^{\wedge k} \otimes W} = R_{\wedge k, \wedge 1}^{\langle N-1 \rangle}(u),$$

$$R_{\wedge k, \wedge 1}(u)|_{((\mathbf{v}_1 \wedge W^{\wedge(k-1)}) \otimes W) \oplus (W^{\wedge k} \otimes \mathbb{C}\mathbf{v}_1)} = \begin{pmatrix} R_{\wedge(k-1), \wedge 1}^{\langle N-1 \rangle}(u) & \tilde{S}' \\ \tilde{S} & u \end{pmatrix},$$

where $\tilde{S}((\mathbf{v}_1 \wedge \mathbf{x}) \otimes \mathbf{w}) = (\mathbf{w} \wedge \mathbf{x}) \otimes \mathbf{v}_1$, and we will not use the explicit form of \tilde{S}' . A similar decomposition can be written for the R -matrix $R_{\wedge 1, \wedge k}(u)$.

In what follows we regard $T(u)$ and $T^{\wedge k}(u)$ as matrices over the algebra $Y(\mathfrak{gl}_N)$ and introduce their submatrices induced by the decompositions $V = \mathbb{C}\mathbf{v}_1 \oplus W$ and $V^{\wedge k} = (\mathbf{v}_1 \wedge W^{\wedge(k-1)}) \oplus W^{\wedge k}$:

$$(11.5) \quad T(u) = \begin{pmatrix} A(u) & B(u) \\ C(u) & D(u) \end{pmatrix}, \quad T^{\wedge k}(u) = \begin{pmatrix} \hat{A}(u) & \hat{B}(u) \\ \hat{C}(u) & \hat{D}(u) \end{pmatrix}.$$

For example, $A(u) = T_{11}(u)$, $B(u) = (T_{12}(u), \dots, T_{1N}(u))$, $D(u) = (T_{ij}(u))_{i,j=2}^N$.

Denote by $\text{HY}(L, M)$ the space $\text{Hom}(L, M) \otimes Y(\mathfrak{gl}_N)$ of matrices with noncommuting entries. We call L the *domain* of those matrices. The coefficients of the series $B, D, \hat{A}, \hat{B}, \hat{D}$ respectively belong to $\text{HY}(W, \mathbb{C})$, $\text{HY}(W, W)$, $\text{HY}(W^{\wedge(k-1)}, W^{\wedge(k-1)})$, $\text{HY}(W^{\wedge k}, W^{\wedge(k-1)})$, $\text{HY}(W^{\wedge k}, W^{\wedge k})$. For $k = 1$ we have $W^{\wedge(k-1)} = \mathbb{C}$, $\hat{A} = A$, $\hat{B} = B$, $\hat{D} = D$.

Set

$$\bar{R}(u) = \frac{1}{u} R^{\langle N-1 \rangle}(u), \quad \tilde{R}(u) = \frac{1}{u+1} R^{\langle N-1 \rangle}_{\wedge(k-1), \wedge 1}(u), \quad \hat{R}(u) = \frac{1}{u} R^{\langle N-1 \rangle}_{\wedge k, \wedge 1}(u).$$

Define a map $S : W^{\wedge(k-1)} \otimes W \rightarrow W^{\wedge k}$ by the rule

$$(11.6) \quad S(\mathbf{x} \otimes \mathbf{w}) = \mathbf{w} \wedge \mathbf{x}.$$

We will use the following commutation relations that are obtained from formulae (4.11), (11.4), Corollary 3.3 and the equality $S\tilde{R}(u-1) = (u-k)S$:

$$(11.7) \quad \hat{B}^{[1]}(u) B^{[2]}(v) = \frac{u-v}{u-v+1} B^{[2]}(v) \hat{B}^{[1]}(u) \hat{R}^{(12)}(u-v),$$

$$(11.8) \quad \hat{A}^{(1)}(u) B^{[2]}(v) = B^{[2]}(v) \hat{A}^{(1)}(u) \tilde{R}^{(12)}(u-v-1) + \frac{1}{u-v} \hat{B}(u) S^{[12]} A(v),$$

$$(11.9) \quad \hat{D}^{(1)}(u) B^{[2]}(v) = B^{[2]}(v) \hat{D}^{(1)}(u) \hat{R}^{(12)}(u-v) - \frac{1}{u-v} S \hat{B}^{[1]}(u) D^{(2)}(v).$$

The equalities hold respectively in $\text{HY}(W^{\wedge k} \otimes W, W^{\wedge(k-1)})$, $\text{HY}(W^{\wedge(k-1)} \otimes W, W^{\wedge(k-1)})$ and $\text{HY}(W^{\wedge k} \otimes W, W^{\wedge k})$, and the superscripts in brackets indicate what tensor factors are domains of the respective matrices. If $k = 1$, relations (11.7)–(11.9) become

$$(11.10) \quad B^{[1]}(u) B^{[2]}(v) = \frac{u-v}{u-v+1} B^{[2]}(v) B^{[1]}(u) \bar{R}^{(12)}(u-v),$$

$$(11.11) \quad A(u) B(v) = \frac{u-v-1}{u-v} B(v) A(u) + \frac{1}{u-v} B(u) A(v),$$

$$(11.12) \quad D^{(1)}(u) B^{[2]}(v) = B^{[2]}(v) D^{(1)}(u) \bar{R}^{(12)}(u-v) - \frac{1}{u-v} B^{[1]}(u) D^{(2)}(v).$$

Let $\check{R}(u) = (u+1)^{-1} P^{(N-1)} R^{(N-1)}(u)$. For an expression $f(u_1, \dots, u_r)$ with coefficients in matrices with the domain $W^{\otimes r}$ and a simple transposition $(i, i+1)$, $i = 1 \dots k-1$, set

$$(11.13) \quad {}^{(i, i+1)}f(u_1, \dots, u_r) = f(u_1, \dots, u_{i-1}, u_{i+1}, u_i, u_{i+2}, \dots, u_r) \check{R}^{(i, i+1)}(u_i - u_{i+1}).$$

The matrix $\check{R}(u)$ has the properties $\check{R}(u) \check{R}(-u) = 1$ and

$$\check{R}^{(12)}(u-v) \check{R}^{(23)}(u) \check{R}^{(12)}(v) = \check{R}^{(23)}(v) \check{R}^{(12)}(u) \check{R}^{(23)}(u-v),$$

cf. (3.2). This yields the following lemma.

Lemma 11.4. *Formula (11.13) extends to the action of the symmetric group S_r on expressions $f(u_1, \dots, u_r)$ with coefficients in matrices with the domain $W^{\otimes r}$: $f \mapsto {}^\sigma f$, $\sigma \in S_r$.*

By formula (11.10) the expression $B^{[1]}(u_1) \dots B^{[r]}(u_r)$ is invariant under the action (11.13) of the symmetric group S_r .

For an expression $f(u_1, \dots, u_r)$ with coefficients in matrices with the domain $W^{\otimes r}$, set

$$(11.14) \quad {}^R\text{Sym}_{u_1, \dots, u_r}^{(1 \dots r)} f(u_1, \dots, u_r) = \sum_{\sigma \in S_r} {}^\sigma f(u_1, \dots, u_r).$$

Proposition 11.5.

$$(11.15) \quad \begin{aligned} & \hat{A}^{(0)}(u) B^{[1]}(u_1) \dots B^{[r]}(u_r) = \\ & = B^{[1]}(u_1) \dots B^{[r]}(u_r) \hat{A}^{(0)}(u) \tilde{R}^{(0r)}(u - u_r - 1) \dots \tilde{R}^{(01)}(u - u_1 - 1) + \\ & + \frac{1}{(r-1)!} \hat{B}(u) {}^R\text{Sym}_{u_1, \dots, u_r}^{(1 \dots r)} \left(\frac{1}{u - u_1} \prod_{i=2}^r \frac{u_1 - u_i - 1}{u_1 - u_i} S^{[01]} B^{[2]}(u_2) \dots B^{[r]}(u_r) A(u_1) \right), \end{aligned}$$

$$(11.16) \quad \begin{aligned} & \hat{D}^{(0)}(u) B^{[1]}(u_1) \dots B^{[r]}(u_r) = \\ & = B^{[1]}(u_1) \dots B^{[r]}(u_r) \hat{D}^{(0)}(u) \hat{R}^{(0r)}(u - u_r) \dots \hat{R}^{(01)}(u - u_1) - \\ & - \frac{1}{(r-1)!} S \hat{B}^{[0]}(u) {}^R\text{Sym}_{u_1, \dots, u_r}^{(1 \dots r)} \left(\frac{1}{u - u_1} B^{[2]}(u_2) \dots B^{[r]}(u_r) D^{(1)}(u_1) \times \right. \\ & \quad \left. \times \bar{R}^{(1r)}(u_1 - u_r) \dots \bar{R}^{(12)}(u_1 - u_2) \right), \end{aligned}$$

where the tensor factors are counted by $0, 1, \dots, r$.

Proof. The statement follows from relations (11.7)–(11.12) by induction with respect to r . We apply formula (11.8) or (11.9) to the product of the first factors in the left hand side and then use the induction assumption. \square

Remark. Formulae (11.15) and (11.16) have the following structure. The first term in the right hand side comes from repeated usage of the first term in the right hand side of relation (11.8) or (11.9), respectively. The second term, involving symmetrization, is effectively

determined by the fact that the whole expression in the right hand side is regular at $u = u_i$ for any $i = 1, \dots, r$, and is invariant with respect to action (11.13) of the symmetric group S_r . The symmetrized expression is obtained by applying once the second term in the right hand side of the relevant relation (11.8) or (11.9) followed by repeated usage of the first term of the respective relation.

Let $Q = \sum_{a=1}^N Q_a E_{aa}^{(N)} \in \text{End}(V)$ be the diagonal matrix from Theorem 5.4. Set

$$(11.17) \quad \bar{Q} = \sum_{a=1}^{N-1} Q_{a+1} E_{aa}^{(N-1)} \in \text{End}(W).$$

Denote $\tilde{Q} = \bar{Q}^{\wedge(k-1)}$ and $\hat{Q} = \bar{Q}^{\wedge k}$. Then formula (4.13) implies that

$$(11.18) \quad \mathcal{T}_{k,Q}(u) = Q_1 \text{tr}_{W^{\wedge(k-1)}}(\tilde{Q} \hat{A}(u)) + \text{tr}_{W^{\wedge k}}(\hat{Q} \hat{D}(u)).$$

To prove formula (5.8) we will employ Propositions 11.3 and 11.5. In our present notation, relation (11.3) reads as follows:

$$(11.19) \quad \mathbb{B}_\xi(t) = B^{[1]}(t_1^1) \dots B^{[\xi_1]}(t_{\xi_1}^1) \tilde{\psi}(t_1^1, \dots, t_{\xi_1}^1) (\mathbb{B}_\xi^{(N-1)}(\bar{t})).$$

We substitute (11.18) and (11.19) into the left hand side of formula (5.8) and use relations (11.15), (11.16) to move $\hat{A}(u)$ and $\hat{D}(u)$ through the product $B^{[1]}(t_1^1) \dots B^{[\xi_1]}(t_{\xi_1}^1)$.

To simplify writing, let $r = \xi^1$ and $u_i = t_i^1$, $i = 1, \dots, r$. We have

$$(11.20) \quad \begin{aligned} & \mathcal{T}_{k,Q}(u) B^{[1]}(u_1) \dots B^{[r]}(u_r) = \\ & = B^{[1]}(u_1) \dots B^{[r]}(u_r) \times \\ & \times \left(Q_1 (\text{tr}_{W^{\wedge(k-1)}} \otimes \text{id}^{\otimes r}) (\tilde{Q}^{(0)} \hat{A}^{(0)}(u) \tilde{R}^{(0r)}(u - u_r - 1) \dots \tilde{R}^{(01)}(u - u_1 - 1)) + \right. \\ & \quad \left. + (\text{tr}_{W^{\wedge k}} \otimes \text{id}^{\otimes r}) (\hat{Q}^{(0)} \hat{D}^{(0)}(u) \hat{R}^{(0r)}(u - u_r) \dots \hat{R}^{(01)}(u - u_1)) \right) + \\ & + \frac{1}{(r-1)!} {}^R\text{Sym}_{u_1, \dots, u_r}^{(1 \dots r)} \left[\frac{1}{u - u_1} \mathcal{B}_Q^{[1]}(u) B^{[2]}(u_2) \dots B^{[r]}(u_r) \times \right. \\ & \quad \left. \times \left(Q_1 \prod_{i=2}^r \frac{u_1 - u_i - 1}{u_1 - u_i} A(u_1) - \bar{Q}^{(1)} D^{(1)}(u_1) \bar{R}^{(1r)}(u_1 - u_r) \dots \bar{R}^{(12)}(u_1 - u_2) \right) \right], \end{aligned}$$

where the tensor factors for the products under the traces are counted by $0, 1, \dots, r$. Here

$$(11.21) \quad \mathcal{B}_Q(u) = (\text{tr}_{W^{\wedge(k-1)}} \otimes \text{id}) (\bar{Q}^{\wedge(k-1)} \hat{B}(u) S)$$

is a series with coefficients in $\text{HY}(W, \mathbb{C})$. Besides this, we use the equality

$$(\text{tr}_{W^{\wedge k}} \otimes \text{id}) \left(\bar{Q}^{\wedge k} S(\hat{B}(u) \otimes \text{id}) \right) = \mathcal{B}_Q(u) \bar{Q}$$

that follows from the cyclic property of the trace and the formula $\bar{Q}^{\wedge k} S = S(\bar{Q}^{\wedge(k-1)} \otimes \bar{Q})$. Notice that $\mathcal{B}_Q(u) = B(u)$ if $k = 1$.

To proceed further, we use the next lemmas.

Lemma 11.6.

$$\begin{aligned} D^{(0)}(u) \bar{R}^{(0r)}(u - u_r) \dots \bar{R}^{(01)}(u - u_1) &= \psi(u_1, \dots, u_r) (T^{\langle N-1 \rangle}(u)), \\ \hat{D}^{(0)}(u) \hat{R}^{(0r)}(u - u_r) \dots \hat{R}^{(01)}(u - u_1) &= \psi(u_1, \dots, u_r) \left((T^{\langle N-1 \rangle}(u))^{\wedge k} \right). \end{aligned}$$

Proof. The equalities follow from formula (4.5), Lemma 4.3 and formula (11.2). \square

Lemma 11.7. For any $X \in Y(\mathfrak{gl}_{N-1})$,

$$(11.22) \quad A(u) \psi(u_1, \dots, u_r)(X) \simeq \psi(u_1, \dots, u_r)(X) A(u),$$

$$\begin{aligned} (11.23) \quad \hat{A}^{(0)}(u) \tilde{R}^{(0r)}(u - u_r - 1) \dots \tilde{R}^{(01)}(u - u_1 - 1) \psi(u_1, \dots, u_r)(X) &\simeq \\ &\simeq \prod_{i=1}^r \frac{u - u_i - 1}{u - u_i} A(u) \psi(u_1, \dots, u_r) \left((T^{\langle N-1 \rangle}(u - 1))^{\wedge(k-1)} X \right). \end{aligned}$$

Proof. Denote by $Y_{\times}(\mathfrak{gl}_N)$ the left ideal in $Y(\mathfrak{gl}_N)$ generated by the coefficients of the series $T_{a1}(u)$ for $2 \leq a \leq N$. It is a subideal of $Y_+(\mathfrak{gl}_N)$. Relations (4.2) imply that for any $Z \in Y(\mathfrak{gl}_{N-1})$ and $C \in Y_{\times}(\mathfrak{gl}_N)$ the coefficients of the series $[T_{11}^{\langle N \rangle}(u), \psi(Z)]$ and the product $C\psi(Z)$ belong to $Y_{\times}(\mathfrak{gl}_N)$. Then formula (11.22) holds because $A(u) = T_{11}^{\langle N \rangle}(u)$ and the coefficients of components of $\psi(u_1, \dots, u_r)(X)$ belong to $\psi(Y(\mathfrak{gl}_{N-1}))$.

To prove formula (11.23) we first observe that the coefficients of entries of the matrix $\hat{A}(u) - A(u) D^{\wedge(k-1)}(u - 1)$ belong to $Y_{\times}(\mathfrak{gl}_N)$; this fact follows from the formula for $\hat{A}(u)$ similar to the first formula of (4.12) for the identity permutation σ . Then we proceed like in Lemma 11.6 and use the fact that $\psi(u_1, \dots, u_r)$ is a homomorphism of algebras. The calculation goes as follows:

$$\begin{aligned} \hat{A}^{(0)}(u) \tilde{R}^{(0r)}(u - u_r - 1) \dots \tilde{R}^{(01)}(u - u_1 - 1) \psi(u_1, \dots, u_r)(X) &\simeq \\ &\simeq A(u) (D^{\wedge(k-1)}(u - 1))^{[0]} \tilde{R}^{(0r)}(u - u_r - 1) \dots \tilde{R}^{(01)}(u - u_1 - 1) \psi(u_1, \dots, u_r)(X) \simeq \\ &\simeq A(u) \psi(u_1, \dots, u_r) \left((T^{\langle N-1 \rangle}(u - 1))^{\wedge(k-1)} \right) \psi(u_1, \dots, u_r)(X) \prod_{i=1}^r \frac{u - u_i - 1}{u - u_i} \simeq \\ &\simeq A(u) \psi(u_1, \dots, u_r) \left((T^{\langle N-1 \rangle}(u - 1))^{\wedge(k-1)} X \right) \prod_{i=1}^r \frac{u - u_i - 1}{u - u_i}. \end{aligned} \quad \square$$

For an expression $f(v)$ set $\operatorname{res}_{v=w} f(u) = ((v-w)f(v))|_{v=w}$, if the substitution makes sense. Observe that

$$\begin{aligned}
(11.24) \quad & \bar{Q}^{(1)} D^{(1)}(u_1) \bar{R}^{(1r)}(u_1 - u_r) \dots \bar{R}^{(12)}(u_1 - u_2) = \\
& = \operatorname{res}_{v=u_1} \left((\operatorname{tr}_W \otimes \operatorname{id}^{\otimes r}) (\bar{Q}^{(0)} D^{(0)}(v) \bar{R}^{(0r)}(v - u_r) \dots \bar{R}^{(01)}(v - u_1)) \right) = \\
& = \operatorname{res}_{v=u_1} \left(\psi(u_1, \dots, u_r) (\mathcal{T}_{1, \bar{Q}}^{\langle N-1 \rangle}(v)) \right),
\end{aligned}$$

because $\operatorname{res}_{v=u_1} (\bar{R}(v - u_1)) = P^{\langle N-1 \rangle}$.

Altogether, for any $X \in Y(\mathfrak{gl}_{N-1})$ we have

$$\begin{aligned}
(11.25) \quad & \mathcal{T}_{k, Q}(u) B^{[1]}(u_1) \dots B^{[r]}(u_r) \psi(u_1, \dots, u_r)(X) \simeq \\
& \simeq \left\{ B^{[1]}(u_1) \dots B^{[r]}(u_r) \left(\psi(u_1, \dots, u_r) (\mathcal{T}_{k-1, \bar{Q}}^{\langle N-1 \rangle}(u-1) X) Q_1 A(u) \prod_{i=1}^r \frac{u - u_i - 1}{u - u_i} + \right. \right. \\
& \quad \left. \left. + \psi(u_1, \dots, u_r) (\mathcal{T}_{k, \bar{Q}}^{\langle N-1 \rangle}(u) X) \right) + \right. \\
& \quad \left. + \frac{1}{(r-1)!} {}^R\operatorname{Sym}_{u_1, \dots, u_r}^{(1 \dots r)} \left[\frac{1}{u - u_1} \mathcal{B}_Q^{[1]}(u) B^{[2]}(u_2) \dots B^{[r]}(u_r) \times \right. \right. \\
& \quad \left. \left. \times \left(\psi(u_1, \dots, u_r)(X) Q_1 A(u_1) \prod_{i=2}^r \frac{u_1 - u_i - 1}{u_1 - u_i} - \operatorname{res}_{v=u_1} \left(\psi(u_1, \dots, u_r) (\mathcal{T}_{1, \bar{Q}}^{\langle N-1 \rangle}(v) X) \right) \right) \right] \right\}.
\end{aligned}$$

The next step is to apply both sides of formula (11.25) to the vector $\mathbf{w}_1^{\otimes r}$. That amounts to replacing there the map $\psi(u_1, \dots, u_r)$ by $\tilde{\psi}(u_1, \dots, u_r)$, and changing the symmetrization ${}^R\operatorname{Sym}_{u_1, \dots, u_r}^{(1 \dots r)}$, cf. (11.14), to the ordinary symmetrization $\operatorname{Sym}_{u_1, \dots, u_r}$, because $\check{R}(u) \mathbf{w}_1 \otimes \mathbf{w}_1 = \mathbf{w}_1 \otimes \mathbf{w}_1$.

Now we replace X in formula (11.25) by $\mathbb{B}_{\xi}^{\langle N-1 \rangle}(\bar{t})$, cf. (11.19), and employ the induction assumption. We recall that $A(u) = T_{11}(u)$, $r = \xi^1$, and $u_i = t_i^1$, $i = 1, \dots, r$. We also use Lemma 4.2 and equalities

$$\begin{aligned}
\pi(x) (T_{11}^{\langle N-1 \rangle}(u)) \mathbf{w}_1 &= \frac{u - x + 1}{u - x} \mathbf{w}_1, \\
\pi(x) (T_{aa}^{\langle N-1 \rangle}(u)) \mathbf{w}_1 &= \mathbf{w}_1, \quad a = 2, \dots, N-1,
\end{aligned}$$

cf. (11.1), which imply that

$$\begin{aligned}
\tilde{\psi}(t_1^1, \dots, t_{\xi^1}^1) (T_{11}^{\langle N-1 \rangle}(u)) &\simeq \mathbf{w}_1^{\otimes r} \otimes T_{22}(u) \prod_{i=1}^{\xi^1} \frac{u - t_i^1 + 1}{u - t_i^1}, \\
\tilde{\psi}(t_1^1, \dots, t_{\xi^1}^1) (T_{aa}^{\langle N-1 \rangle}(u)) &\simeq \mathbf{w}_1^{\otimes r} \otimes T_{a+1, a+1}(u), \quad a = 2, \dots, N-1.
\end{aligned}$$

Besides this, we use Lemmas 4.1 and 11.1. As a result, we transform formula (11.25) to

$$\begin{aligned}
(11.26) \quad \mathcal{T}_{k,Q}(u) \mathbb{B}_\xi(t) &\simeq \\
&\simeq \mathbb{B}_\xi(t) \left(Q_1 T_{11}(u) \prod_{i=1}^{\xi^1} \frac{u - t_i^1 - 1}{u - t_i^1} \sum_{\mathbf{a}} \prod_{j=2}^k \mathcal{X}_{\xi,Q}^{a_j}(u - j + 1; t) + \right. \\
&\quad \left. + \sum_{\mathbf{b}} \prod_{j=1}^k \mathcal{X}_{\xi,Q}^{b_j}(u - j + 1; t) \right) + \mathcal{U}_{\xi,k,Q}(u; t),
\end{aligned}$$

cf. (5.7), where the first sum is taken over all $(k-1)$ -tuples $\mathbf{a} = (a_2, \dots, a_k)$ such that $2 \leq a_2 < \dots < a_k \leq N$, the second sum is taken over all k -tuples $\mathbf{b} = (b_1, \dots, b_k)$ such that $2 \leq b_1 < \dots < b_k \leq N$, and

$$\begin{aligned}
(11.27) \quad \mathcal{U}_{\xi,k,Q}(u; t) &\simeq \\
&\simeq B^{[1]}(t_1^1) \dots B^{[r]}(t_{\xi^1}^1) \left(\tilde{\psi}(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{U}_{\bar{\xi},k-1,\bar{Q}}^{\langle N-1 \rangle}(u-1; \bar{t})) Q_1 A(u) \prod_{i=1}^{\xi^1} \frac{u - t_i^1 - 1}{u - t_i^1} + \right. \\
&\quad \left. + \tilde{\psi}(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{U}_{\bar{\xi},k,\bar{Q}}^{\langle N-1 \rangle}(u; \bar{t})) \right) + \\
&+ \frac{1}{(\xi^1 - 1)!} \text{Sym}_{t_1^1, \dots, t_{\xi^1}^1} \left[\frac{1}{u - t_1^1} \mathcal{B}_Q^{[1]}(u) B^{[2]}(t_2^1) \dots B^{[r]}(t_{\xi^1}^1) \times \right. \\
&\quad \times \left(\tilde{\psi}(t_1^1, \dots, t_{\xi^1}^1) (\mathbb{B}_{\bar{\xi}}^{\langle N-1 \rangle}(\bar{t})) \times \right. \\
&\quad \times \left(Q_1 T_{11}(t_1^1) \prod_{i=2}^{\xi^1} \frac{t_1^1 - t_i^1 - 1}{t_1^1 - t_i^1} - Q_2 T_{22}(t_1^1) \prod_{i=2}^{\xi^1} \frac{t_1^1 - t_i^1 + 1}{t_1^1 - t_i^1} \prod_{j=1}^{\xi^2} \frac{t_1^1 - t_j^2 - 1}{t_1^1 - t_j^2} \right) - \\
&\quad \left. \left. - \text{res}_{v=t_1^1} \left(\tilde{\psi}(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{U}_{\bar{\xi},1,\bar{Q}}^{\langle N-1 \rangle}(v; \bar{t})) \right) \right) \right].
\end{aligned}$$

The last formula determines $\mathcal{U}_{\xi,k,Q}(u; t)$ modulo terms with coefficients belonging to $Y_+(\mathfrak{gl}_N)$. By the induction assumption the expressions $\mathcal{U}_{\bar{\xi},j,\bar{Q}}^{\langle N-1 \rangle}(v; \bar{t})$ belong to $I_{\bar{\xi},j,\bar{Q}}^{\langle N-1 \rangle}(v; \bar{t})$. Then together with Lemmas 3.2, 4.1, 4.2, 11.1, 11.2, and formulae (5.5), (11.1), (11.2), the induction assumption implies that there exists an expression $\mathcal{U}_{\xi,k,Q}(u; t)$ which belongs to $I_{\xi,k,Q}$ and obeys formula (11.27). Therefore, formula (11.26) coincides with formula (5.8). Theorem 5.4 is proved. \square

Proof of Proposition 6.2. We prove the statement by induction with respect to N using Proposition 11.3. It is straightforward to check that the induction assumption yields

$$e_{ab} \mathbb{B}_\xi^{v_1 \otimes \dots \otimes v_n}(\tilde{t}; z) = 0, \quad 2 \leq a < b \leq N.$$

Let $C(u)$ be the left bottom block in the decomposition (11.5) of $T(u)$. Consider the coefficients of the series $C(u)$, $C(u) = \sum_{s=1}^{\infty} C_s u^{-s}$. Then C_1 is a matrix from $\text{HY}(\mathbb{C}, W)$ whose entries are e_{12}, \dots, e_{1N} . Therefore, it remains to check that $C_1 \mathbb{B}_{\xi}^{v_1 \otimes \dots \otimes v_n}(\tilde{t}; z) = 0$.

Relations (4.2) implies that

$$C_1 B(u) - B(u) C_1 = D(u) - A(u).$$

Then, similarly to the proof of Proposition 11.5, we get

$$\begin{aligned} C_1 B^{[1]}(u_1) \dots B^{[r]}(u_r) &= B^{[1]}(u_1) \dots B^{[r]}(u_r) C_1 + \\ &+ \frac{1}{(r-1)!} {}^R\text{Sym}_{u_1, \dots, u_r}^{(1 \dots r)} \left(B^{[2]}(u_2) \dots B^{[r]}(u_r) \times \right. \\ &\times \left. \left(D^{[1]}(u_1) \bar{R}^{(1r)}(u_1 - u_r) \dots \bar{R}^{(12)}(u_1 - u_2) - \prod_{i=2}^r \frac{u_1 - u_i - 1}{u_1 - u_i} A(u_1) \right) \right). \end{aligned}$$

The rest of the proof of Proposition 6.2 is similar to considerations in the proof of Theorem 5.4 after Lemma 11.6 and in the proof of Theorem 6.1. \square

12. Proof of Theorem 8.6

We will prove Theorem 8.6 by induction with respect to N , suitably adapting the proof of Theorem 5.4.

We will continue to use notation introduced in the previous section. The maps ψ , $\psi(x_1, \dots, x_r)$ and $\tilde{\psi}(x_1, \dots, x_r)$ defined there for the Yangian $Y(\mathfrak{gl}_{N-1})$ have their counterparts for the current Lie algebra $\mathfrak{gl}_{N-1}[x]$. Abusing notation we will denote them by the same letters.

The embedding $\psi : U(\mathfrak{gl}_{N-1}[x]) \rightarrow U(\mathfrak{gl}_N[x])$ is defined by the rule $\psi(L_{ab}^{\langle N-1 \rangle}(u)) = L_{a+1, b+1}(u)$, $a, b = 1, \dots, N-1$, where $L_{ab}^{\langle N-1 \rangle}(u)$ is series (8.1) for $\mathfrak{gl}_{N-1}[x]$. The homomorphism $\psi(x_1, \dots, x_r) : U(\mathfrak{gl}_{N-1}[x]) \rightarrow \text{End}(W^{\otimes r}) \otimes U(\mathfrak{gl}_N[x])$ is given by

$$\psi(x_1, \dots, x_r)(L_{ab}^{\langle N-1 \rangle}(u)) = 1^{\otimes r} \otimes L_{a+1, b+1}(u) + \sum_{i=1}^r \frac{1^{\otimes(i-1)} \otimes E_{ba}^{\langle N-1 \rangle} \otimes 1^{\otimes(r-i)}}{u - x_i} \otimes 1,$$

$a, b = 1, \dots, N-1$. The map $\tilde{\psi}(x_1, \dots, x_r) : U(\mathfrak{gl}_{N-1}[x]) \rightarrow W^{\otimes r} \otimes U(\mathfrak{gl}_N[x])$ is such that

$$\tilde{\psi}(x_1, \dots, x_r)(L_{ab}^{\langle N-1 \rangle}(u)) = \mathbf{w}_1^{\otimes r} \otimes L_{a+1, b+1}(u) + \delta_{a1} \sum_{i=1}^r \frac{\mathbf{w}_1^{\otimes(i-1)} \otimes \mathbf{w}_b \otimes \mathbf{w}_1^{\otimes(r-i)}}{u - x_i} \otimes 1,$$

$a, b = 1, \dots, N-1$, and $\tilde{\psi}(x_1, \dots, x_r)(X_1 X_2) = \psi(x_1, \dots, x_r)(X_1) \tilde{\psi}(x_1, \dots, x_r)(X_2)$ for any $X_1, X_2 \in U(\mathfrak{gl}_{N-1}[x])$.

Lemma 12.1. *We have*

$$\tilde{\psi}(x_1, \dots, x_r) (U(\mathfrak{gl}_{N-1}[x]) \mathfrak{n}_+^{\langle N-1 \rangle}[x]) \subset W^{\otimes r} \otimes U(\mathfrak{gl}_{N-1}[x]) \mathfrak{n}_+^{\langle N-1 \rangle}[x].$$

The proof is straightforward.

Define the embedding $\phi : U(\mathfrak{gl}_{N-2}[x]) \hookrightarrow U(\mathfrak{gl}_{N-1}[x])$, $\phi(L_{ab}^{\langle N-2 \rangle}(u)) = L_{a+1, b+1}^{\langle N-1 \rangle}(u)$, $a, b = 1, \dots, N-2$, similar to the embedding ψ .

Lemma 12.2. *We have $\tilde{\psi}(x_1, \dots, x_r) \circ \phi = \mathbf{w}_1^{\otimes r} \otimes (\psi \circ \phi)$.*

The proof is straightforward.

For any element $g \in W^{\otimes r} \otimes U(\mathfrak{gl}_{N-1}[x])$ we consider its components g^{a_1, \dots, a_r} given by

$$g = \sum_{a_1, \dots, a_r=1}^{N-1} \mathbf{w}_{a_1} \otimes \dots \otimes \mathbf{w}_{a_r} \otimes g^{a_1, \dots, a_r}.$$

Recall that $\bar{\xi} = (\xi^2, \dots, \xi^{N-1})$ and $\bar{t} = (t_1^2, \dots, t_{\xi^2}^2; \dots; t_1^{N-1}, \dots, t_{\xi^{N-1}}^{N-1})$.

Proposition 12.3. *We have*

$$(12.1) \quad \mathbb{F}_{\xi}(t) = \sum_{a_1, \dots, a_{\xi^1}=1}^{N-1} L_{1, a_1+1}(t_1^1) \dots L_{1, a_{\xi^1}+1}(t_{\xi^1}^1) \left(\tilde{\psi}(t_1^1, \dots, t_{\xi^1}^1) (\mathbb{F}_{\bar{\xi}}^{\langle N-1 \rangle}(\bar{t})) \right)^{a_1, \dots, a_{\xi^1}}.$$

Proof. The statement follows from formula (8.7). \square

In Section 10 we introduced the degree function \deg on the algebra $Y(\mathfrak{gl}_N)[[u^{-1}, \partial_u, \zeta]]$ by the rule $\deg u^{-1} = \deg \partial_u = \deg \zeta = -1$. We define the degree to rational expressions in $t_1^1, \dots, t_{\xi^{N-1}}^{N-1}$ with coefficients in $Y(\mathfrak{gl}_N)[[u^{-1}, \partial_u, \zeta]]$, assuming in addition that $\deg t_i^a = 1$ and $\deg(t_i^a - t_j^b)^{-1} = -1$ for all possible a, b, i, j . We use similar definition of the degree for the Yangian $Y(\mathfrak{gl}_{N-1})$ and related rational expressions in $t_2^2, \dots, t_{\xi^{N-1}}^{N-1}$. Notice that the maps ψ , $\psi(t_1^1, \dots, t_{\xi^1}^1)$ and $\tilde{\psi}(t_1^1, \dots, t_{\xi^1}^1)$ agree with the introduced degree functions and the projections to the graded algebras. For example, if $X \in Y(\mathfrak{gl}_{N-1})$ and $\deg X = k$, then $\deg \psi(X) = k$ and $\psi(\text{grad}_k X) = \text{grad}_k(\psi(X))$.

Proposition 12.4. *We have $\deg(\mathbb{B}_{\xi}(t)) = -|\xi|$, and $\text{grad}_{-|\xi|}(\mathbb{B}_{\xi}(t)) = \mathbb{F}_{\xi}(t)$.*

Proof. The statement follows from formula (10.2) and Propositions 11.3, 12.3 by induction with respect to N . \square

Proof of Theorem 8.6. As in Section 10, set $Q = 1 + \zeta K$. We will employ Theorems 10.1, 10.2, and Proposition 12.4, and adapt arguments used in the proof of Theorem 5.4 to prove the statement of Theorem 8.6 by induction with respect to N .

Consider formula (11.25) and write there $\mathcal{B}_{k, Q}(u)$ instead of $\mathcal{B}_Q(u)$ to indicate the dependence on k . Recall that we have $r = \xi^1$, $u_i = t_i^1$, $i = 1, \dots, \xi^1$, and $A(u) = T_{11}(u)$.

Taking the sum of equalities (11.25) for $k = 1, \dots, l$, with coefficients $\frac{(-1)^{l-k}(N-k)!}{(N-l)!(l-k)!}$ and using formula (10.3), for any $X \in Y(\mathfrak{gl}_{N-1})$, we obtain

$$\begin{aligned}
(12.2) \quad & \mathcal{S}_{l,Q}(u) B^{[1]}(t_1^1) \dots B^{[\xi^1]}(t_{\xi^1}^1) \psi(t_1^1, \dots, t_{\xi^1}^1)(X) \simeq \\
& \simeq \left\{ B^{[1]}(t_1^1) \dots B^{[\xi^1]}(t_{\xi^1}^1) \times \right. \\
& \quad \times \left(\psi(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{S}_{l-1, \bar{Q}}^{\langle N-1 \rangle}(u-1) X) \left(Q_1 T_{11}(u) \prod_{i=1}^{\xi^1} \frac{u - t_i^1 - 1}{u - t_i^1} - 1 \right) + \right. \\
& \quad \left. \left. + \psi(t_1^1, \dots, t_{\xi^1}^1) \left((\mathcal{S}_{l-1, \bar{Q}}^{\langle N-1 \rangle}(u-1) - \mathcal{S}_{l-1, \bar{Q}}^{\langle N-1 \rangle}(u) + \mathcal{S}_{l, \bar{Q}}^{\langle N-1 \rangle}(u)) X \right) \right) + \right. \\
& \quad \left. + \frac{1}{(\xi^1 - 1)!} {}^R\text{Sym}_{t_1^1, \dots, t_{\xi^1}^1}^{(1 \dots \xi^1)} \left[\frac{1}{u - t_1^1} \mathcal{P}_{l, Q}^{[1]}(u) B^{[2]}(t_2^1) \dots B^{[\xi^1]}(t_{\xi^1}^1) \times \right. \right. \\
& \quad \times \left(\psi(t_1^1, \dots, t_{\xi^1}^1)(X) \left(Q_1 T_{11}(t_1^1) \prod_{i=2}^{\xi^1} \frac{t_1^1 - t_i^1 - 1}{t_1^1 - t_i^1} - 1 \right) + \right. \\
& \quad \left. \left. + \left(1 - \text{res}_{v=t_1^1} \left(\psi(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{T}_{1, \bar{Q}}^{\langle N-1 \rangle}(v)) \right) \right) \psi(t_1^1, \dots, t_{\xi^1}^1)(X) \right) \right] \Big\},
\end{aligned}$$

where

$$(12.3) \quad \mathcal{P}_{l, Q}(u) = \frac{1}{(N-l)!} \sum_{k=1}^l (-1)^{l-k} \frac{(N-k)!}{(l-k)!} \mathcal{B}_{k, Q}(u).$$

Notice that in formula (11.25) for $k = N$ we have $\mathcal{T}_{N, \bar{Q}}^{\langle N-1 \rangle}(u) = 0$ and $\mathcal{B}_{N, Q}(u) = 0$, and in formula (12.2) for $l = N$ we assume that $\mathcal{S}_{N, \bar{Q}}^{\langle N-1 \rangle}(u) = 0$. We remind that $\mathcal{T}_{0, Q}(u) = 1$ and $\mathcal{T}_{0, \bar{Q}}^{\langle N-1 \rangle}(u) = \mathcal{S}_{0, \bar{Q}}^{\langle N-1 \rangle}(u) = 1$ by convention.

Recall that $\mathcal{B}_{k, Q}(u)$ is an $(N-1) \times 1$ matrix whose entries are series in u^{-1} with coefficients in $Y(\mathfrak{gl}_N)$, and $\mathcal{P}_{l, Q}(u)$ is an $(N-1) \times 1$ matrix of the same kind,

$$\mathcal{B}_{k, Q}(u) = (\mathcal{B}_{k, Q; 1}(u), \dots, \mathcal{B}_{k, Q; N-1}(u)), \quad \mathcal{P}_{l, Q}(u) = (\mathcal{P}_{l, Q; 1}(u), \dots, \mathcal{P}_{l, Q; N-1}(u)).$$

Lemma 12.5. *For any $m = k, \dots, N$, distinct $i_1, \dots, i_k \in \{1, \dots, m\}$, and any $s = 1, \dots, k$, we have*

$$\begin{aligned}
\mathcal{B}_{k, Q; a}(u) &= \frac{m! (N-m)!}{(k-1)! (N-k)!} \times \\
&\times (\text{tr}_{V^{\otimes m}} \otimes \text{id})(E_{a+1, 1}^{(i_s)} Q^{(i_1)} \dots Q^{(i_k)} T^{(i_1, m+1)}(u) \dots T^{(i_k, m+1)}(u-k+1) \mathbb{A}_{\{m\}}^{(1 \dots m)}).
\end{aligned}$$

Proof. Formulae (11.21), (11.17), (11.5), and (11.6) give that

$$\mathcal{B}_{k, Q; a}(u) = k (\text{tr}_{V^{\otimes k}} \otimes \text{id})(E_{a+1, 1}^{(1)} Q^{(1)} \dots Q^{(k)} T^{(1, k+1)}(u) \dots T^{(k, k+1)}(u-k+1) \mathbb{A}_{\{k\}}^{(1 \dots k)}).$$

The remaining consideration is similar to the proof of Lemma 4.8. \square

Lemma 12.5 and formula (12.3) allow us to obtain $\mathcal{B}_{k,Q;a}(u)$ and $\mathcal{P}_{l,Q;a}(u)$ as coefficients of a formal difference operator similarly to formulae (4.14), (10.6),

$$\begin{aligned}
(12.4) \quad & \sum_{k=1}^m (-1)^{(k-1)} \frac{(N-k)!}{(m-k)!} \mathcal{B}_{k,Q;a}(u) e^{(m-k)\partial_u} = \\
& = \sum_{l=1}^m (-1)^{(l-1)} \frac{(N-l)!}{(m-l)!} \mathcal{P}_{l,Q;a}(u) (e^{\partial_u} - 1)^{m-l} = \\
& = m(N-m)! (\text{tr}_{V^{\otimes m}} \otimes \text{id}) \left(E_{a+1,1}^{(1)} Q^{(1)} T^{(1,m+1)}(u) e^{-\partial_u} \times \right. \\
& \quad \left. \times (1 - Q^{(2)} T^{(2,m+1)}(u) e^{-\partial_u}) \dots (1 - Q^{(m)} T^{(m,m+1)}(u) e^{-\partial_u}) \mathbb{A}_{\{m\}}^{(1\dots m)} \right) e^{m\partial_u}.
\end{aligned}$$

Introduce the following $(N-1) \times 1$ matrices

$$\mathcal{F}_{l,K}(u) = (\mathcal{F}_{l,K;1}(u), \dots, \mathcal{F}_{l,K;N-1}(u)), \quad l = 1, \dots, N,$$

whose entries are defined by the rule

$$\begin{aligned}
(12.5) \quad & \frac{1}{N} \sum_{l=1}^N (-1)^{(l-1)} \mathcal{F}_{l,K;a}(u) \partial_u^{N-l} = \\
& = (\text{tr}_{V^{\otimes N}} \otimes \text{id}) \left(E_{a+1,1}^{(1)} (\partial_u - K^{(2)} - L^{(2,N+1)}(u)) \dots (\partial_u - K^{(N)} - L^{(N,N+1)}(u)) \mathbb{A}_{\{N\}}^{(1\dots N)} \right).
\end{aligned}$$

It is easy to see that $\mathcal{F}_{1,K}(u) = 0$. Like in the proof of Theorem 10.2, formulae (12.4) and (12.5) imply that $\deg(\mathcal{P}_{l,Q}(u)) = 1-l$ for any $l = 1, \dots, N$, and

$$\text{grad}_{1-l}(\mathcal{P}_{l,Q}(u)) = \mathcal{F}_{l,\zeta K}(u).$$

Set $\bar{K} = \sum_{a=1}^{N-1} K_{a+1} E_{aa}^{\langle N-1 \rangle}$, so that $\bar{Q} = 1 + \zeta \bar{K}$.

Lemma 12.6. We have $\deg \left(1 - \text{res}_{v=t_1^1} \left(\psi(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{T}_{1,\bar{Q}}^{\langle N-1 \rangle}(v)) \right) \right) = -1$ and

$$\begin{aligned}
& \text{grad}_{-1} \left(1 - \text{res}_{v=t_1^1} \left(\psi(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{T}_{1,\bar{Q}}^{\langle N-1 \rangle}(v)) \right) \right) = \\
& = \left(\frac{1}{v - t_1^1} - \psi(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{G}_{1,\zeta \bar{K}}^{\langle N-1 \rangle}(v)) \right) \Big|_{v=t_1^1} - \text{res}_{v=t_1^1} \left(\psi(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{G}_{2,\zeta \bar{K}}^{\langle N-1 \rangle}(v)) \right).
\end{aligned}$$

Proof. The statement follows from formulae (11.24) and (8.5) by straightforward calculations. Notice that the denominator of the expression $(v - t_1^1)^{-1} - \psi(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{G}_{1,\bar{K}}^{\langle N-1 \rangle}(v))$ does not actually contain the factor $v - t_1^1$, which makes the substitution $v = t_1^1$ in the expression legal. \square

The Poincaré-Birkhoff-Witt theorem for the Yangian $Y(\mathfrak{gl}_N)$, see [MNO, Corollary 1.23], [Mo, Theorem 2.6], yields that if $X \in Y_+(\mathfrak{gl}_N)$, $\deg X = d$, then $\text{grad}_d X \in U(\mathfrak{gl}_N[x]) \mathfrak{n}_+[x]$.

Let $F(u) = (L_{12}(u), \dots, L_{1N}(u)) = \text{grad}_{-1}(B(u))$, and $d = \deg X$ in formula (12.2). We apply the map grad_{d-l} to both sides of (12.2), and then evaluate the result at $\zeta = 1$. Recall that we have

$$\text{grad}_{-l}(\mathcal{S}_{l,Q}(u)) = \mathcal{G}_{l,\zeta K}(u), \quad \text{grad}_{-s}(\mathcal{S}_{s,\bar{Q}}^{\langle N-1 \rangle}(u)) = \mathcal{G}_{s,\zeta K}^{\langle N-1 \rangle}(u).$$

Thus we obtain that for any $Z \in U(\mathfrak{gl}_{N-1}[x])$,

$$\begin{aligned} (12.6) \quad & \mathcal{G}_{l,K}(u) F^{[1]}(t_1^1) \dots F^{[\xi^1]}(t_{\xi^1}^1) \psi(t_1^1, \dots, t_{\xi^1}^1)(Z) \simeq \\ & \simeq \left\{ F^{[1]}(t_1^1) \dots F^{[\xi^1]}(t_{\xi^1}^1) \left(\psi(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{G}_{l-1,\bar{K}}^{\langle N-1 \rangle}(u) Z) \left(K_1 + L_{11}(u) - \sum_{i=1}^{\xi^1} \frac{1}{u - t_i^1} \right) + \right. \right. \\ & \quad \left. \left. + \psi(t_1^1, \dots, t_{\xi^1}^1) ((\mathcal{G}_{l,\bar{K}}^{\langle N-1 \rangle}(u) - \dot{\mathcal{G}}_{l-1,\bar{K}}^{\langle N-1 \rangle}(u)) Z) \right) \right\} + \\ & + \frac{1}{(\xi^1 - 1)!} \text{Sym}_{t_1^1, \dots, t_{\xi^1}^1} \left[\frac{1}{u - t_1^1} \mathcal{F}_{l,K}^{[1]}(u) F^{[2]}(t_2^1) \dots F^{[\xi^1]}(t_{\xi^1}^1) \times \right. \\ & \quad \times \left(\psi(t_1^1, \dots, t_{\xi^1}^1)(Z) \left(K_1 + L_{11}(t_1^1) - \sum_{i=2}^{\xi^1} \frac{1}{t_1^1 - t_i^1} \right) + \right. \\ & \quad \left. \left. + \left(\left(\frac{1}{v - t_1^1} - \psi(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{G}_{1,\bar{K}}^{\langle N-1 \rangle}(v)) \right) \right|_{v=t_1^1} - \text{res}_{v=t_1^1} \left(\psi(t_1^1, \dots, t_{\xi^1}^1) (\mathcal{G}_{2,\bar{K}}^{\langle N-1 \rangle}(v)) \right) \right) \right. \\ & \quad \left. \left. \times \psi(t_1^1, \dots, t_{\xi^1}^1)(Z) \right) \right] \Bigg\}. \end{aligned}$$

Here $\dot{\mathcal{G}}_{l-1,\bar{K}}^{\langle N-1 \rangle}(u) = \frac{d}{du} \mathcal{G}_{l-1,\bar{K}}^{\langle N-1 \rangle}(u)$.

The rest of the proof of Theorem 8.6 is similar to the proof of Theorem 5.4 after formula (11.25). We apply both sides of formula (12.6) to the vector $\mathbf{w}_1^{\otimes r}$, that amounts to replacing there the map $\psi(u_1, \dots, u_r)$ by $\tilde{\psi}(u_1, \dots, u_r)$, and then take $Z = \mathbb{F}_{\bar{\xi}}^{\langle N-1 \rangle}(\bar{t})$. Since formula (12.1) is equivalent to

$$\mathbb{F}_{\xi}(t) = F^{[1]}(t_1^1) \dots F^{[\xi^1]}(t_{\xi^1}^1) \tilde{\psi}(t_1^1, \dots, t_{\xi^1}^1) (\mathbb{F}_{\bar{\xi}}^{\langle N-1 \rangle}(\bar{t})),$$

the left hand side of (12.6) becomes $\mathcal{G}_{l,K}(u) \mathbb{F}_{\xi}(t)$, and to transform the right hand side to the required form, we employ the induction assumption. \square

Appendix A. The Gaudin model as a limit of the XXX-type model

Let M_1, \dots, M_n be \mathfrak{gl}_N -modules. In Sections 6 and 7, we describe the XXX-type model on the tensor product $M_1 \otimes \dots \otimes M_n$ considering the action of the Yangian $Y(\mathfrak{gl}_N)$ on the tensor product $M_1(z_1) \otimes \dots \otimes M_n(z_n)$ of evaluation modules. The objects associated with the XXX-type model are the operators

$$T_{ab}^{M_1 \otimes \dots \otimes M_n}(u; z) = T_{ab}(u) \Big|_{M_1(z_1) \otimes \dots \otimes M_n(z_n)}, \quad a, b = 1, \dots, N,$$

rationally depending on u and z_1, \dots, z_n ; the transfer matrices $\mathcal{T}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u; z)$, cf. (6.1); the modified transfer matrices

$$\begin{aligned} \mathcal{S}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u; z) &= \mathcal{S}_{k,Q}(u) \Big|_{M_1(z_1) \otimes \dots \otimes M_n(z_n)} = \\ &= \frac{1}{(N-k)!} \sum_{l=0}^k (-1)^{k-l} \frac{(N-l)!}{(k-l)!} \mathcal{T}_{l,Q}^{M_1 \otimes \dots \otimes M_n}(u), \end{aligned}$$

$k = 0, \dots, N$, cf. (10.3); the difference operator

$$\begin{aligned} (\text{A.1}) \quad \mathfrak{D}_{N,Q}^{M_1 \otimes \dots \otimes M_n}(u; \partial_u; z) &= \sum_{k=0}^N (-1)^k \mathcal{T}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u; z) e^{-k\partial_u} = \\ &= \sum_{k=0}^N (-1)^k \mathcal{S}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u; z) (e^{\partial_u} - 1)^{(N-k)\partial_u} e^{-N\partial_u}, \end{aligned}$$

the deformed Shapovalov form $S_{M_1 \otimes \dots \otimes M_n}^z$ on $M_1 \otimes \dots \otimes M_n$, see (7.9); the universal weight function $\mathbb{B}_{\xi}^{v_1 \otimes \dots \otimes v_n}(t; z)$, cf. (6.2); the rational functions

$$\begin{aligned} \mathcal{Q}_{\xi,Q}^{a,i}(t; z; \Lambda) &= \\ &= \frac{Q_a}{Q_{a+1}} \prod_{j=1}^n \frac{t_i^a - z_j + \Lambda_j^a}{t_i^a - z_j + \Lambda_j^{a+1}} \prod_{j=1}^{\xi^{a-1}} \frac{t_i^a - t_j^{a-1} + 1}{t_i^a - t_j^{a-1}} \prod_{\substack{j=1 \\ j \neq i}}^{\xi^a} \frac{t_i^a - t_j^a - 1}{t_i^a - t_j^a + 1} \prod_{j=1}^{\xi^{a+1}} \frac{t_i^a - t_j^{a+1}}{t_i^a - t_j^{a+1} - 1}, \end{aligned}$$

$a = 1, \dots, N-1$, $i = 1, \dots, \xi^a$, that define the Bethe ansatz equations (6.3); and the difference operator

$$\mathfrak{M}_{\xi,Q}(u, \partial_u; t; z; \Lambda) = (1 - \mathcal{X}_{\xi,Q}^1(u; t; z; \Lambda) e^{-\partial_u}) \dots (1 - \mathcal{X}_{\xi,Q}^N(u; t; z; \Lambda) e^{-\partial_u}),$$

cf. (6.6), (6.4), whose coefficients are eigenvalues of the transfer matrices.

In Section 9, we describe the Gaudin model on the tensor product $M_1 \otimes \dots \otimes M_n$ considering the action of the current algebra $\mathfrak{gl}_N[x]$ on the tensor product $M_1(z_1) \otimes \dots \otimes M_n(z_n)$

of evaluation modules over $\mathfrak{gl}_N[x]$. For the Gaudin model, the associated objects are the operators

$$L_{ab}^{M_1 \otimes \dots \otimes M_n}(u; z) = \sum_{i=1}^n \frac{e_{ba}^{(i)}}{u - z_i}, \quad a = 1, \dots, N,$$

rationally depending on u and z_1, \dots, z_n ; the transfer matrices $\mathcal{G}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$, cf. (9.1), (9.2); the differential operator

$$(A.2) \quad \mathcal{D}_K^{M_1 \otimes \dots \otimes M_n}(u, \partial_u; z) = \sum_{k=0}^N (-1)^k \mathcal{G}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z) \partial_u^{N-k}$$

with $\text{End}(M_1 \otimes \dots \otimes M_n)$ -valued rational coefficients, cf. (8.3); the Shapovalov form $S_{M_1 \otimes \dots \otimes M_n}$ on $M_1 \otimes \dots \otimes M_n$; the universal weight function $\mathbb{F}_\xi^{v_1 \otimes \dots \otimes v_n}(t; z)$, cf. (9.6); the rational functions

$$\begin{aligned} \mathcal{K}_{\xi,K}^{a,i}(t; z; \Lambda) = \\ = K_a - K_{a+1} + \sum_{j=1}^n \frac{\Lambda_j^a - \Lambda_j^{a+1}}{t_i^a - z_j} + \sum_{j=1}^{\xi^{a-1}} \frac{1}{t_i^a - t_j^{a-1}} - 2 \sum_{\substack{j=1 \\ j \neq i}}^{\xi^a} \frac{1}{t_i^a - t_j^a} + \sum_{j=1}^{\xi^{a+1}} \frac{1}{t_i^a - t_j^{a+1}}, \end{aligned}$$

$a = 1, \dots, N-1$, $i = 1, \dots, \xi^a$, that define the Bethe ansatz equations (9.3); and the differential operator

$$\mathcal{M}_{\xi,K}(u, \partial_u; t; z; \Lambda) = \prod_{1 \leq a \leq N} \left(\partial_u - K_a - \sum_{i=1}^n \frac{\Lambda_i^a}{u - z_i} - \sum_{i=1}^{\xi^{a-1}} \frac{1}{u - t_i^{a-1}} + \sum_{i=1}^{\xi^a} \frac{1}{u - t_i^a} \right).$$

cf. (9.4), whose coefficients are eigenvalues of the transfer matrices.

The objects associated with the Gaudin model can be obtained from the corresponding objects for the XXX-type model in the following limit.

Set $\varepsilon^{-1}z = (\varepsilon^{-1}z_1, \dots, \varepsilon^{-1}z_n)$ and $\varepsilon^{-1}t = (\varepsilon^{-1}t_1^1, \dots, \varepsilon^{-1}t_{\xi^{N-1}}^{N-1})$.

Theorem A.1. *Let $Q = 1 + \varepsilon K$. As $\varepsilon \rightarrow 0$, we have*

$$(A.3) \quad T_{ab}^{M_1 \otimes \dots \otimes M_n}(\varepsilon^{-1}u; \varepsilon^{-1}z) = \delta_{ab} + \varepsilon L_{ab}^{M_1 \otimes \dots \otimes M_n}(u; z) + O(\varepsilon^2),$$

$a, b = 1, \dots, N$,

$$(A.4) \quad \mathcal{S}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(\varepsilon^{-1}u; \varepsilon^{-1}z) = \varepsilon^k \mathcal{G}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z) + O(\varepsilon^{k+1}),$$

$k = 0, \dots, N$,

$$(A.5) \quad \mathcal{D}_{N,Q}^{M_1 \otimes \dots \otimes M_n}(\varepsilon^{-1}u, \varepsilon \partial_u; \varepsilon^{-1}z) = \varepsilon^N \mathcal{D}_K^{M_1 \otimes \dots \otimes M_n}(u; \partial_u; z) + O(\varepsilon^{N+1}),$$

$$(A.6) \quad S_{M_1 \otimes \dots \otimes M_n}^{\varepsilon^{-1}z} \rightarrow S_{M_1} \otimes \dots \otimes S_{M_n},$$

$$\begin{aligned}
(A.7) \quad \mathbb{B}_\xi^{v_1 \otimes \dots \otimes v_n}(\varepsilon^{-1}t; \varepsilon^{-1}z) & \prod_{i=1}^n \prod_{a=1}^{N-1} \prod_{j=1}^{\xi^a} \frac{\varepsilon}{t_j^a - z_i} \prod_{a=1}^{N-2} \prod_{i=1}^{\xi^a} \prod_{j=1}^{\xi^{a+1}} \frac{\varepsilon}{t_j^{a+1} - t_i^a} = \\
& = \varepsilon^{|\xi|} \mathbb{F}_\xi^{v_1 \otimes \dots \otimes v_n}(t; z) + O(\varepsilon^{|\xi|+1}),
\end{aligned}$$

$$(A.8) \quad \mathcal{Q}_{\xi, Q}^{a, i}(\varepsilon^{-1}t; \varepsilon^{-1}z; \Lambda) = 1 + \varepsilon \mathcal{K}_{\xi, K}^{a, i}(t; z; \Lambda) + O(\varepsilon^2),$$

$$(A.9) \quad \mathfrak{M}_{\xi, Q}(\varepsilon^{-1}u, \varepsilon \partial_u; \varepsilon^{-1}t; \varepsilon^{-1}z; \Lambda) = \varepsilon^N \mathcal{M}_{\xi, K}(u, \partial_u; t; z; \Lambda).$$

Proof. Formulae (A.3) and (A.8) are straightforward. The proof of formulae (A.4) and (A.5) is similar to the proof of Theorem 10.2. Formula (A.6) follows from formula (7.4). Formula (A.7) is similar to the claim of Proposition 12.4. \square

Remark. Using Theorem A.1 one can see that Theorem 7.6 implies Theorem 9.1, and Theorem 6.1 implies Theorem 9.2 for an isolated solution $\tilde{t} = (\tilde{t}_1^1, \dots, \tilde{t}_{\xi^{N-1}}^{N-1})$ of the Bethe ansatz equations (9.3).

Appendix B. Dynamical Hamiltonians as parts of transfer matrices

Let $C_1 = \sum_{a=1}^N e_{aa}$, and $C_2 = \sum_{1 \leq a < b \leq N} (e_{aa}e_{bb} - e_{ab}e_{ba} + e_{aa})$. They are central elements of the universal enveloping algebra $U(\mathfrak{gl}_N)$.

Let M_1, \dots, M_n be \mathfrak{gl}_N -modules. Let K be a diagonal matrix, $K = \sum_{a=1}^N K_a E_{aa}$. The Gaudin Hamiltonians $H_{1,K}(z), \dots, H_{n,K}(z)$, see [G], the rational dynamical Hamiltonians $G_{1,K}(z), \dots, G_{N,K}(z)$, see [FMTV], [TV2], and the trigonometric dynamical Hamiltonians $X_{1,K}(z), \dots, X_{N,K}(z)$, see [TV2], [TV3], acting in $M_1 \otimes \dots \otimes M_n$, are the following operators:

$$\begin{aligned}
H_{i,K}(z) &= \sum_{a=1}^N K_a e_{aa}^{(i)} + \sum_{a,b=1}^N \sum_{\substack{j=1 \\ j \neq i}}^n \frac{e_{ab}^{(i)} e_{ba}^{(j)}}{z_i - z_j}, \\
G_{a,K}(z) &= \sum_{i=1}^n z_i e_{aa}^{(i)} + \sum_{\substack{b=1 \\ b \neq a}}^N \frac{e_{ab} e_{ba} - e_{aa}}{K_a - K_b}.
\end{aligned}$$

$$X_{a,K}(z) = -\frac{(e_{aa})^2}{2} + \sum_{i=1}^n z_i e_{aa}^{(i)} + \sum_{b=1}^N \sum_{1 \leq i < j \leq n} e_{ab}^{(i)} e_{ba}^{(j)} + \sum_{\substack{b=1 \\ b \neq a}}^N \frac{K_b}{K_a - K_b} (e_{ab} e_{ba} - e_{aa}).$$

Recall that e_{ab} acts on $M_1 \otimes \dots \otimes M_n$ as $\sum_{i=1}^n e_{ab}^{(i)}$.

It is well known that the Gaudin Hamiltonians come from the residues of the transfer matrix $\mathcal{G}_{2,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$:

$$\begin{aligned} \mathcal{G}_{2,K}^{M_1 \otimes \dots \otimes M_n}(u; z) &= \\ &= \operatorname{tr} K^{\wedge 2} + \sum_{i=1}^n \frac{1}{u - z_i} \left[C_1^{(i)} \left(\operatorname{tr} K + \sum_{\substack{j=1 \\ j \neq i}}^n \frac{C_1^{(j)}}{z_i - z_j} \right) - H_{i,K}(z) \right] + \sum_{i=1}^n \frac{C_2^{(i)}}{(u - z_i)^2}. \end{aligned}$$

The rational and trigonometric dynamical Hamiltonians also can be recovered from the transfer matrices $\mathcal{G}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$ and $\mathcal{T}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$, respectively. To this end, consider the following $\operatorname{End}(M_1 \otimes \dots \otimes M_n)$ -valued rational function of u, x, z_1, \dots, z_n :

$$\begin{aligned} \widehat{\mathcal{G}}_K(u, x; z) &= \sum_{k=0}^N (-1)^k \mathcal{G}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z) x^{N-k} \prod_{a=1}^N \frac{1}{x - K_a}, \\ \widehat{\mathcal{T}}_K(u, x; z) &= \sum_{k=0}^N (-1)^k \mathcal{T}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z) x^{N-k} \prod_{a=1}^N \frac{1}{x - K_a}. \end{aligned}$$

Proposition B.1. *We have*

$$\begin{aligned} \widehat{\mathcal{G}}_K(u, x; z) &= 1 - u^{-1} \sum_{a=1}^N \frac{e_{aa}}{x - K_a} - \\ &\quad - u^{-2} \sum_{a=1}^N \frac{1}{x - K_a} \left(G_{a,K}(z) - \sum_{\substack{b=1 \\ b \neq a}}^N \frac{e_{aa} e_{bb}}{K_a - K_b} \right) + O(u^{-3}) \end{aligned}$$

and

$$\begin{aligned} \widehat{\mathcal{T}}_K(u, x; z) &= 1 - u^{-1} \sum_{a=1}^N \frac{K_a}{x - K_a} e_{aa} - \\ &\quad - u^{-2} \sum_{a=1}^N \frac{K_a}{x - K_a} \left(X_{a,K}(z) + \frac{(e_{aa})^2}{2} - \sum_{\substack{b=1 \\ b \neq a}}^N \frac{K_b}{K_a - K_b} e_{aa} e_{bb} \right) + O(u^{-3}). \end{aligned}$$

Proof. To get the first formula we compute the expansion of $\mathcal{G}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$ as $u \rightarrow \infty$ using formulae (8.3), (8.4), (9.1), (9.2), and Proposition 8.1. For the second formula we compute the expansion of $\mathcal{T}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$ using formulae (4.13), (6.1), and the definition of the module $M_1(z_1) \otimes \dots \otimes M_n(z_n)$ over the Yangian $Y(\mathfrak{gl}_N)$. \square

Appendix C. Eigenvalues of transfer matrices for real z_1, \dots, z_n

Denote by $\mathfrak{gl}_N^{\mathbb{R}}$ the real part of the Lie algebra \mathfrak{gl}_N generated by e_{ab} , $a, b = 1, \dots, N$. Let M_1, \dots, M_n be finite-dimensional irreducible \mathfrak{gl}_N -modules with highest weights $\Lambda_1, \dots, \Lambda_n$, and highest weight vectors v_1, \dots, v_n . Denote by $M_1^{\mathbb{R}}, \dots, M_n^{\mathbb{R}}$ be their real parts generated over $\mathfrak{gl}_N^{\mathbb{R}}$ by the vectors v_1, \dots, v_n , respectively. It is clear that for real u, z_1, \dots, z_n , real matrices Q, K , and any $k = 1, \dots, N$, the operators $\mathcal{T}_{k,Q}^{M_1 \otimes \dots \otimes M_n}(u; z)$ and $\mathcal{G}_{k,K}^{M_1 \otimes \dots \otimes M_n}(u; z)$ preserve the tensor product $M_1^{\mathbb{R}} \otimes_{\mathbb{R}} \dots \otimes_{\mathbb{R}} M_n^{\mathbb{R}}$.

Since M_1, \dots, M_n are finite-dimensional irreducible \mathfrak{gl}_N -modules, the tensor product $S_{M_1} \otimes \dots \otimes S_{M_n}$ of Shapovalov forms restricted to $M_1^{\mathbb{R}} \otimes_{\mathbb{R}} \dots \otimes_{\mathbb{R}} M_n^{\mathbb{R}}$ is a positive definite bilinear form. Hence, an operator, which is symmetric with respect to this form, has real eigenvalues. Therefore, Theorems 9.1 and 9.2 yield the following statement.

Theorem C.1. *Let K be a real diagonal matrix, z_1, \dots, z_n real numbers and $\tilde{t} = (\tilde{t}_1^1, \dots, \tilde{t}_{\xi^{N-1}}^{N-1})$ a solution of the Bethe ansatz equations (9.3). If $\mathbb{F}_{\xi}^{v_1 \otimes \dots \otimes v_n}(\tilde{t}; z) \neq 0$, then for any $k = 1, \dots, N$, the function $\mathcal{Z}_{\xi,k,K}(u; \tilde{t}; z; \Lambda)$, cf. (9.5), takes real values for real values of u .*

For a dominant weight $\Lambda = (\Lambda^1, \dots, \Lambda^N)$, set $\Lambda' = \Lambda^N + 1 - \max\{a \mid \Lambda^a > \Lambda^N\}$.

For finite-dimensional irreducible \mathfrak{gl}_N -modules L, M , the R -matrix $R_{LM}(u)$ is a rational function of u . Its poles and degeneracy points are known from the representation theory of the Yangian $Y(\mathfrak{gl}_N)$, see [Mo]. In particular, the following proposition holds.

Proposition C.2. *The R -matrix $R_{M_i M_j}(u)$ is well-defined and nondegenerate for real values of u such that $u > \Lambda_i^1 - \Lambda_j'$ or $u < \Lambda_i' - \Lambda_j^1$. Moreover, $R_{M_i M_j}(u)$ has a simple pole at $u = \Lambda_i' - \Lambda_j^1$ with the degenerate residue, and the value $R_{M_i M_j}(\Lambda_i^1 - \Lambda_j')$ is well-defined, non-zero, and degenerate.*

It is clear that for real z_1, \dots, z_n , the operator $R_{M_1 \dots M_n}(z)$, cf. (7.6), preserves the tensor product $M_1^{\mathbb{R}} \otimes_{\mathbb{R}} \dots \otimes_{\mathbb{R}} M_n^{\mathbb{R}}$.

Proposition C.3. *Let z_1, \dots, z_n be real numbers such that $z_i - z_j > \Lambda_i^1 - \Lambda_j'$ for any $i \neq j$, $i, j = 1, \dots, n$. Then the bilinear form $S_{M_1 \otimes \dots \otimes M_n}^z$, cf. (7.9), restricted to $M_1^{\mathbb{R}} \otimes_{\mathbb{R}} \dots \otimes_{\mathbb{R}} M_n^{\mathbb{R}}$ is positive definite.*

Proof. The statement follows from Proposition C.2 and formula (7.4). \square

Theorem C.4. *Let Q be a real diagonal matrix, z_1, \dots, z_n real numbers such that $z_i - z_j > \Lambda_i^1 - \Lambda_j'$ for any $i \neq j$, $i, j = 1, \dots, n$, and $\tilde{t} = (\tilde{t}_1^1, \dots, \tilde{t}_{\xi^{N-1}}^{N-1})$ a solution of the Bethe ansatz equations (6.3). If $\mathbb{B}_{\xi}^{v_1 \otimes \dots \otimes v_n}(\tilde{t}; z) \neq 0$, then for any $k = 1, \dots, N$, the function*

$$\sum_{1 \leq a_1 < \dots < a_k \leq N} \prod_{r=1}^k \mathcal{X}_{\xi,Q}^{a_r}(u - r + 1; \tilde{t}; z; \Lambda),$$

cf. (6.4), takes real values for real values of u .

Proof. The statement follows from Theorems 7.6 and 6.1, and Proposition C.3. \square

References

- [D] V.G.Drinfeld, *Quantum groups*, in Proceedings of the ICM, Berkeley, 1986, AMS, Providence RI, 1987, pp. 798–820.
- [FFR] B.Feigin, E.Frenkel, and N.Reshetikhin, *Gaudin model, Bethe ansatz and critical level*, Comm. Math. Phys. **166** (1994), no. 1, 27–62.
- [FMTV] G.Felder, Ya.Markov, V.Tarasov and A.Varchenko, *Differential equations compatible with KZ equations*, Math. Phys., Analysis and Geometry **3** (2000), 139–177.
- [G] M.Gaudin, *La fonction d’onde de Bethe*, Masson, Paris, 1983.
- [Ju] B.Jurčo, *Classical Yang-Baxter equations and quantum integrable systems*, J. Math. Phys. **30** (1989), 1289–1293.
- [KBI] V.E.Korepin, N.M.Bogoliubov and A.G.Izergin, *Quantum inverse scattering method and correlation functions*, Cambridge University Press, 1993.
- [KR1] P.P.Kulish and N.Yu.Reshetikhin, *Diagonalization of $GL(n)$ invariant transfer-matrices and quantum N -wave system (Lee model)*, J. Phys. A **15** (1983), L591–L596.
- [KR2] P.P.Kulish and N.Yu.Reshetikhin, *On GL_3 -invariant solutions of the Yang-Baxter equation and associated quantum systems*, J. Soviet Math. **34** (1986), 1948–1971.
- [KS] P.P.Kulish and E.K.Sklyanin, *Quantum spectral transform method. Recent developments*, Lect. Notes in Phys. **151** (1982), 61–119.
- [Ma] A.Matsuo, *An application of Aomoto-Gelfand hypergeometric functions to the $SU(n)$ Knizhnik-Zamolodchikov equation*, Comm. Math. Phys. **134** (1990), 65–77.
- [Mo] A.Molev, *Yangians and their applications*, Handbook of Algebra, vol. 3, North-Holland, Amsterdam, 2003, pp. 907–959.
- [MNO] A.Molev, M.Nazarov, and G.Olshanski, *Yangians and classical Lie algebras*, Russian Math. Surveys **51** (1996), 205–282.
- [MTV] E.Mukhin, V.Tarasov, and A.Varchenko, *The B. and M.Shapiro conjecture in real algebraic geometry and the Bethe ansatz*, Preprint (2005), 1–17, [math.AG/0512299](#).
- [MV1] E.Mukhin and A.Varchenko, *Critical points of master functions and flag varieties*, Commun. Contemp. Math. **6** (2004), no. 1, 111–163.
- [MV2] E.Mukhin and A.Varchenko, *Solutions to the XXX type Bethe ansatz equations and flag varieties*, Central Europ. J. Math. **1** (2003), 238–271.
- [N] M.Nazarov, *Yangians and Capelli identities*, Amer. Math. Society Transl., Ser. 2 **181** (1998), 139–163.
- [NO] M.Nazarov and G.Olshanski, *Bethe subalgebras in twisted Yangians*, Comm. Math. Phys. **178** (1996), no. 2, 483–506.
- [RSV] R.Rimányi, L.Stevens, and A.Varchenko, *Combinatorics of rational functions and Poincaré-Birchoff-Witt expansions of the canonical $U(\mathfrak{n}_-)$ -valued differential form*, Annals Combin. **9** (2005), no. 1, 57–74.
- [RV] N.Reshetikhin and A.Varchenko, *Quasiclassical asymptotics of solutions to the KZ equations*, Geometry, Topology, & Physics, Internat. Press, Cambridge, MA, 1995, pp. 293–322.
- [SV] V.V.Schechtman and A.N.Varchenko, *Arrangements of hyperplanes and Lie algebras homology*, Invent. Math. **106** (1991), 139–194.
- [T] D.Talalaev, *Quantization of the Gaudin system*, Preprint (2004), 1–19, [hep-th/0404153](#).

- [TV1] V.Tarasov and A.Varchenko, *Jackson integral representations for solutions to the quantized Knizhnik-Zamolodchikov equation*, St.Petersburg Math. J. **6** (1994), no. 2, 90–137.
- [TV2] V.Tarasov and A.Varchenko, *Duality for Knizhnik-Zamolodchikov and dynamical equations*, Acta Appl. Math. **73** (2002), no. 1-2, 141–154.
- [TV3] V.Tarasov and A.Varchenko, *Dynamical differential equations compatible with rational qKZ equations*, Lett. Math. Phys. **71** (2005), no. 2, 101–108.