

Relatively hyperbolic groups: geometry and quasi-isometric invariance

Cornelia Druţu

Abstract

In this paper it is proved that relative hyperbolicity is an invariant of quasi-isometry. As a byproduct we provide simplified definitions of relative hyperbolicity in terms of the geometry of a Cayley graph. In particular we obtain a definition very similar to the one of hyperbolicity, relying on the existence for every quasi-geodesic triangle of a central left coset of peripheral subgroup.

Contents

| | | |
|----------|---|-----------|
| 1 | Introduction | 1 |
| 1.1 | Rigidity result | 1 |
| 1.2 | Metric and algebraic relative hyperbolicity | 3 |
| 1.3 | New definitions of relative hyperbolicity | 5 |
| 1.4 | Organization of the paper | 6 |
| 2 | Preliminaries | 6 |
| 2.1 | Definitions and notations | 6 |
| 2.2 | Asymptotic cones of a metric space | 7 |
| 2.3 | Relatively hyperbolic groups | 9 |
| 3 | Tree-graded metric spaces | 10 |
| 3.1 | Definition and properties | 10 |
| 3.2 | Topological bigons and property (T_2) | 11 |
| 4 | Asymptotically tree-graded metric spaces | 13 |
| 4.1 | Definition and properties | 13 |
| 4.2 | Property (T_2) and polygons with limit edges | 16 |
| 4.3 | New definition, useful for the rigidity of relatively hyperbolic groups | 19 |
| 4.4 | New definition, closer to the definition of hyperbolicity | 23 |
| 5 | Quasi-isometric rigidity of relatively hyperbolic groups | 30 |

1 Introduction

1.1 Rigidity result

A finitely generated group G endowed with a word metric becomes a geometric object. In many cases G is related to other geometric structures as well, for instance if it acts properly

discontinuously and with compact quotient on a metric space (X, dist) . In the latter case, G is quasi-isometric to X (see § 2.1 for the definition of quasi-isometry).

In the general setting of a group G quasi-isometric to a metric space X , M. Gromov asked ([Gro87],[Gro93]) what are the structures on X that allow to obtain consistent information on G . In particular, X can be itself a finitely generated group with some property (P), and the question is whether this property transfers or not to G (or to a subgroup of finite index of G or to a quotient of G by a finite normal subgroup). In case it does, the property (P) is called *geometric*. In another terminology the class of groups with property (P) is called *rigid*, and the results of this type are called *rigidity results*. Recall that if a finite index subgroup of a group G has property (P) then G is said to *virtually* satisfy (P).

Examples of rigid/non-rigid classes of groups:

1. the class of virtually nilpotent groups is rigid [Gro81];
2. the class of virtually solvable groups is not rigid [Dyu00]; but smaller classes of virtually solvable groups are rigid ([FM98], [FM99], [EFW05]);
3. amenability is a geometric property [GdlH90];
4. property (T) is not geometric [Val04];
5. the class of fundamental groups of compact manifolds covered by a fixed symmetric space X of non-compact type is rigid (this statement includes many deep results of different authors: [Sul81], [Tuk86], [Pan89], [Cho96], [Tuk88], [Tuk94], [CJ94], [Gab92], [KL97b], [EF97]);
6. the same rigidity result holds for fundamental groups of finite volume non-compact manifolds ([Sch96b], [FS96], [Sch96a], [Esk98], [Dru00]);
7. hyperbolicity is a geometric property [Gro87].

The present paper focuses on the class of relatively hyperbolic groups.¹ This notion was introduced by M. Gromov in [Gro87] aiming to generalize the notion of hyperbolic group so as to include fundamental groups of non-compact finite volume hyperbolic manifolds. Other definitions of the same notion, as well as important developments of the theory of relatively hyperbolic groups can be found in [Bow97], [Far98], [Dah03b], [Yam04], [DS05b], [Osi06]. In § 2.3 we recall the definition of relative hyperbolicity due to B. Farb. We say that a group G is *properly* relatively hyperbolic if it is hyperbolic relative to finitely many *proper* subgroups of finite type.

Examples of relatively hyperbolic groups:

1. a hyperbolic group is hyperbolic relative to $\{1\}$;
2. $A *_F B$, where F is finite, is hyperbolic relative to A and B ; more generally, fundamental groups of finite graphs of groups with finite edge groups are hyperbolic relative to the vertex groups [Bow97];
3. fundamental groups of complete finite volume manifolds of pinched negative sectional curvature are hyperbolic relative to the fundamental groups of their cusps ([Bow97], [Far98]);

¹By relatively hyperbolic group we mean what is sometimes called in the literature *strongly relatively hyperbolic group*, in contrast with *weakly relatively hyperbolic group*.

4. fundamental groups of (non-geometric) Haken manifolds with at least one hyperbolic component are hyperbolic relative to fundamental groups of maximal graph-manifold components and to fundamental groups of tori and Klein bottles not contained in a graph-manifold component;
5. fully residually free groups, also known as limit groups, are hyperbolic relative to their maximal Abelian non-cyclic subgroups [Dah03a];
6. more generally, finitely generated groups acting freely on \mathbb{R}^n -trees are hyperbolic relative to their maximal Abelian non-cyclic subgroups [Gui04].

The main result of the paper is the following, answering Problem 1.15 in [DS05b].

Theorem 1.1 (relative hyperbolicity is geometric, Theorem 5.13). *Let G be a finitely generated group properly hyperbolic relative to a finite family of finitely generated subgroups $\mathcal{H} = \{H_1, \dots, H_n\}$. If a group G' is quasi-isometric to G then G' is properly hyperbolic relative to $\mathcal{H}' = \{H'_1, \dots, H'_m\}$, where each H'_i can be embedded quasi-isometrically in H_j for some $j = j(i) \in \{1, 2, \dots, n\}$.*

Rigidity has already been proved for smaller sub-classes of relatively hyperbolic groups: fundamental groups of non-geometric Haken manifolds with at least one hyperbolic component ([KL95] and [KL97a]), non-uniform lattices of isometries of rank one symmetric spaces different from $\mathbb{H}_{\mathbb{R}}^2$ [Sch96b], fundamental groups of graphs of groups with finite edge groups [PW02].

In [DS05b] the rigidity is proved in the particular case when all H_i are *unconstricted*, that is they have at least one asymptotic cone without (global) cut-points. Under this assumption it can be ensured that each H'_i is quasi-isometric to some H_j . The theorem has been improved in [BDM05], where it is only required that each H_i is *not properly relatively hyperbolic*.

This type of rigidity of relatively hyperbolic groups, also preserving the quasi-isometry classes of peripheral subgroups, can only be obtained under some extra-hypotheses on H_i , and not in the full generality assumed in Theorem 1.1. This can be seen in the case when $G = G' = A*B*C*D$, with G hyperbolic relative to $\{A*B, C*D\}$ and G' hyperbolic relative to $\{A, B, C, D\}$.

1.2 Metric and algebraic relative hyperbolicity

In order to study the rigidity of relatively hyperbolic groups it is necessary to have a definition only in terms of the Cayley graph of the group. Most of the definitions of relative hyperbolicity (except the ones in [DS05b] and in [Osi06]) use not only the Cayley graph of the group but also a metric space obtained from this graph by gluing to each left coset of a peripheral subgroup some geometric object (a hyperbolic horoball [Gro87], countably many edges with one common endpoint [Far98] etc).

In what follows, we recall the definition provided in [DS05b].

A complete geodesic metric space \mathbb{F} is *tree-graded with respect to a collection \mathcal{P} of closed geodesic subsets (called *pieces*)*, if the following two properties are satisfied:

- (T_1) two different pieces have at most one point in common;
- (T_2) any simple non-trivial geodesic triangle is contained in one piece.

A metric space X is *asymptotically tree-graded (ATG) with respect to a collection of subsets \mathcal{A}* if every asymptotic cone of X is tree-graded with respect to the collection of limit sets of sequences in \mathcal{A} .

Equivalently, X is ATG with respect to \mathcal{A} if a list of three geometric properties is satisfied, which goes approximately as follows (for details see Theorem 4.1 in [DS05b], Theorem 3.8 in this paper):

- (α_1) finite radius tubular neighborhoods of distinct elements in \mathcal{A} are either disjoint or intersect in sets of uniformly bounded diameter;
- (α_2) geodesics diverging slower than linearly from a set A in \mathcal{A} must intersect a finite radius tubular neighborhood of A ;
- (α_3) “fat” geodesic polygons must stay close to a set in \mathcal{A} (the meaning of “fat” here is the contrary of “thin” in its metric hyperbolic sense).

The metric space X is *properly* asymptotically tree-graded with respect to \mathcal{A} if moreover it is not contained in any tubular neighborhood of a subset in \mathcal{A} .

The purpose of the notion of ATG metric space is to generalize to metric spaces the concept of relative hyperbolicity (see also [BF01] for another metric version of the notion of relative hyperbolicity). The fact that the definition is coherent with the definition for groups is illustrated by the following result.

Theorem 1.2 ([DS05b], **Theorem 1.11**). *An infinite finitely generated group G is properly relatively hyperbolic with respect to a finite family of finitely generated subgroups $\mathcal{H} = \{H_1, \dots, H_m\}$ if and only if G is properly asymptotically tree-graded with respect to the collection of left cosets $\mathcal{L} = \{gH_i ; g \in G/H_i, i \in \{1, 2, \dots, m\}\}$.*

In [DS05b], the equivalence appears without the extra hypothesis that both structures are proper. Still, it is easy to prove that one structure is proper if and only if the other one is.

Indeed, if $G \subset \mathcal{N}_\tau(gH_i)$ for some $i \in \{1, 2, \dots, m\}$ and $\tau > 0$ then H_i is infinite. If $H_i \neq G$ then there exists $g'H_i \neq gH_i$, $g'H_i \subset \mathcal{N}_\tau(gH_i)$. This contradicts property (α_1).

The converse statement is immediate.

The equivalence in Theorem 1.2 suggests the following natural question.

Question 1.3 (see **Problem 1.16** in [DS05b]). *If a group is asymptotically tree-graded in a metric sense, that is with respect to a collection \mathcal{A} of subsets, does it follow that it is relatively hyperbolic with respect to some finite family of subgroups ?*

The implication in Question 1.3 was previously known to be true only under some restrictive metric conditions on \mathcal{A} (see [DS05b, Theorem 5.13] and its improvement in [BDM05]).

We prove the following.

Theorem 1.4 (**Theorem 5.1**). *Let G be an infinite finitely generated group properly asymptotically tree-graded with respect to a collection \mathcal{A} of subsets. Then G is either hyperbolic or it is properly relatively hyperbolic with respect to some finitely generated subgroups H_1, \dots, H_m , such that every H_i is contained in the finite radius tubular neighborhood of a set $A_i \in \mathcal{A}$.*

Remark 1.5. If G is hyperbolic then it is properly hyperbolic relative to $H = \{1\}$. Still, in this case one cannot state that H is contained in some $\mathcal{N}_\varepsilon(A)$ with $A \in \mathcal{A}$, because in the definition that we adopt of asymptotically tree-graded metric spaces the finite radius tubular neighborhoods of the sets $A \in \mathcal{A}$ do not cover the whole space (see Remark 4.10).

Theorem 1.4 implies Theorem 1.1. Indeed, a group quasi-isometric to a relatively hyperbolic group is asymptotically tree-graded as a metric space [DS05b, Theorem 5.1].

1.3 New definitions of relative hyperbolicity

If a group has an asymptotically tree-graded structure which is equivariant with respect to the action of the group on itself by left translations, then the group is relatively hyperbolic, by a standard argument (see Proposition 5.4). Thus, the main step in the proof of Theorem 1.4 is to construct an equivariant ATG structure on a group G from a given ATG structure \mathcal{A} on G . A natural idea is to consider all the translated ATG structures $g\mathcal{A} = \{gA ; A \in \mathcal{A}\}$ and to take intersections of the form $\bigcap_{g \in G} gA_g$, with $A_g \in \mathcal{A}$. To make such an argument work, we need a good behavior of the ATG properties with respect to intersection. This requires two things:

- that in (α_2) is specified the position of the point on the slowly diverging geodesic contained in the finite radius tubular neighborhood of A ;
- that in (α_3) it suffices to consider polygons with a fixed number of edges.

It is not difficult to prove that in an ATG metric space the following strengthening of (α_2) holds (see Theorem 4.21 for the precise formulation):

- (β_2) a geodesic diverging from a set A in \mathcal{A} at a speed slow enough compared to the linear one has its middle third contained in a finite radius tubular neighborhood of A .

Property (α_3) is required in order to ensure property (T_2) in every asymptotic cone, with respect to limit sets of sequences from \mathcal{A} . If (α_3) is known only for hexagons then (T_2) holds in any asymptotic cone for simple triangles whose edges are limits of sequences of geodesics (see Proposition 4.13). But generically a geodesic in an asymptotic cone of a group is not limit of a sequence of geodesics (see the example in the end of § 2.2).

Thus, in order to ensure (T_2) for an arbitrary geodesic triangle the argument in [DS05b] was to prove that such a triangle can be approximated by a geodesic triangle which is limit of a sequence of fat polygons with the same number k of edges, with k increasing as the constant of approximation decreases (see Lemma 4.14). This and (α_3) for an arbitrary number of edges implies (T_2) .

In this paper we show that if property (T_1) holds in every asymptotic cone, an inductive argument allows to deduce (T_2) from (α_3) reduced to the case of hexagons (Corollary 4.19). Thus we obtain the following version, essential for the rigidity result, of the three geometric properties defining an ATG metric space.

Theorem 1.6 (Theorem 4.21). *Let (X, dist) be a geodesic metric space and let \mathcal{A} be a collection of subsets of X . The space X is ATG with respect to \mathcal{A} if and only if properties (α_1) , (α_2) and (α_3) restricted to geodesic hexagons, are satisfied.*

In particular (α_2) can be replaced by its stronger version (β_2) .

Property (α_3) restricted to geodesic hexagons is denoted (β_3) in the text.

Asymptotically tree-graded metric spaces have a property that strongly reminds of hyperbolic metric spaces. Indeed, a metric space is hyperbolic if and only if the edges of every quasi-geodesic triangle intersect a ball of uniformly bounded radius [Gro87, §6]. A space X asymptotically tree-graded with respect to a collection of subsets \mathcal{A} has the following property, proved in [DS05b]:

- (*) the edges of any quasi-geodesic triangle in X either intersect a finite radius ball or a finite radius tubular neighborhood of a subset in \mathcal{A} . Moreover, in the latter case the distance between the entrance points of two edges starting from a common vertex into the tubular neighborhood is uniformly bounded.

See Section 4.3 in this paper for details.

If (X, \mathcal{A}) satisfy property $(*)$ then the space X is called $(*)$ -asymptotically tree-graded with respect to \mathcal{A} [DS05a]. This notion is weaker than the notion of ATG metric space (see Remark 4.34, (2)). A natural question to ask is the following.

Question 1.7. Under what additional conditions is a $(*)$ -asymptotically tree-graded metric space also asymptotically tree-graded ?

Recall that property $(*)$ was essential in the proof of the fact that the property of Rapid Decay transfers from the peripheral subgroups H_1, \dots, H_m of a relatively hyperbolic group to the group itself [DS05a].

The arguments used to prove Theorem 1.6 can be adapted to answer Question 1.7.

Theorem 1.8 (Theorem 4.35). *Let (X, dist) and \mathcal{A} be as in Theorem 1.6. The metric space X is asymptotically tree-graded with respect to \mathcal{A} if and only if (X, \mathcal{A}) satisfy properties (α_1) and (α_2) , and moreover X is $(*)$ -asymptotically tree-graded with respect to \mathcal{A} .*

1.4 Organization of the paper

Section 2 contains preliminaries on asymptotic cones and relatively hyperbolic groups, as well as notations used throughout the paper.

In Section 3 are recalled some basic facts about tree-graded spaces. Proposition 3.14 proved in the same section is very useful in different arguments deducing the general property (T_2) from (T_1) , and (T_2) restricted to some particular cases.

Section 4 begins with a short overview of the main properties of ATG metric spaces. In §4.2 an induction argument and Proposition 3.14 are used to show the following important result. Denote by (Π_3) the property (T_2) restricted to triangles with edges limits of sequences of geodesics. If in an asymptotic cone $\text{Con}_\omega(X)$ of a metric space X a collection \mathcal{A}_ω of closed subsets satisfies (T_1) and (Π_3) then \mathcal{A}_ω satisfies (T_2) in full generality (Corollary 4.18).

This statement is the main ingredient in the proof of Theorem 1.6, appearing in §4.3. It also plays a central part in the proof of Theorem 1.8 given in §4.4. Another main step in the proof of Theorem 1.8 is to deduce from properties $(*)$, (α_1) and (α_2) the fact that fat quadrilaterals are contained in finite radius tubular neighborhoods of subsets in \mathcal{A} (Lemma 4.39). Once this last statement proved, from it as well as from property $(*)$ and Proposition 3.10 can be deduced property (Π_3) . Corollary 4.18 allows to finish the argument.

In Section 5 first Theorem 1.4 is proved. The first step of the proof is to construct from a given ATG structure on a group an ATG structure that is equivariant with respect to the action of the group on itself by left translations. The subsets in the new ATG structure are indexed by equivalence classes of fat hexagons. A simple argument allows to finish the proof, by showing that the existence of an equivariant ATG structure implies that the group is relatively hyperbolic (Proposition 5.4).

Theorem 1.1 is deduced from Theorem 1.4 in the end of Section 5.

2 Preliminaries

2.1 Definitions and notations

Let Y be a subset in a metric space (X, dist) . We denote by $\mathcal{N}_\delta(Y)$ the set $\{x \in X \mid \text{dist}(x, Y) < \delta\}$, which we call the δ -tubular neighborhood of Y . We denote by $\overline{\mathcal{N}}_\delta(Y)$ the set $\{x \in X \mid \text{dist}(x, Y) \leq \delta\}$, called the δ -closed tubular neighborhood of Y .

When Y is a singleton y , we also use the notations $B(y, \delta)$ and respectively $\overline{B}(y, \delta)$.

The *Hausdorff distance* between two subsets Y and Z in X , denoted by $\text{dist}_H(Y, Z)$, is the minimal δ such that $Y \subset \overline{\mathcal{N}}_\delta(Z)$ and $Z \subset \overline{\mathcal{N}}_\delta(Y)$.

Definition 2.1. An action of a group G on a metric space X is called \mathcal{K} -*transitive*, where \mathcal{K} is a non-negative constant, if for every $x \in X$ the closed tubular neighborhood $\overline{\mathcal{N}}_{\mathcal{K}}(Gx)$ of the orbit of x coincides with X .

A 0-transitive action is transitive in the standard sense.

An (L, C) -*quasi-isometric embedding* of a metric space (X, dist_X) into a metric space (Y, dist_Y) is a map $\mathbf{q} : X \rightarrow Y$ such that for every $x_1, x_2 \in X$,

$$\frac{1}{L} \text{dist}_X(x_1, x_2) - C \leq \text{dist}_Y(\mathbf{q}(x_1), \mathbf{q}(x_2)) \leq L \text{dist}_X(x_1, x_2) + C, \quad (1)$$

for some constants $L \geq 1$ and $C \geq 0$.

If moreover Y is contained in the C -tubular neighborhood of $\mathbf{q}(X)$ then \mathbf{q} is called an (L, C) -*quasi-isometry*. In this case there exists an (L, C) -quasi-isometry $\bar{\mathbf{q}} : Y \rightarrow X$ such that $\bar{\mathbf{q}} \circ \mathbf{q}$ and $\mathbf{q} \circ \bar{\mathbf{q}}$ are at uniformly bounded distance from the respective identity maps (see [GdlH90] for a proof). We call $\bar{\mathbf{q}}$ *quasi-converse of \mathbf{q}* .

If $\mathbf{q} : [a, b] \rightarrow X$ is an (L, C) -quasi-isometric embedding then \mathbf{q} is called an (L, C) -*quasi-geodesic (segment)* in X . The same name is used for the image of \mathbf{q} .

If $\mathbf{q}_i : [0, \ell_i] \rightarrow X$, $i = 1, 2$, are two quasi-geodesic segments with $\mathbf{q}_1(\ell_1) = \mathbf{q}_2(0)$, then we denote by $\mathbf{q}_1 \sqcup \mathbf{q}_2$ the map $\mathbf{q} : [0, \ell_1 + \ell_2] \rightarrow X$ defined by $\mathbf{q}(t) = \mathbf{q}_1(t)$ for $t \in [0, \ell_1]$ and $\mathbf{q}(t) = \mathbf{q}_2(t - \ell_1)$ for $t \in [\ell_1, \ell_1 + \ell_2]$.

Definition 2.2 (almost geodesics). If an (L, C) -quasi-geodesic \mathbf{q} is L -Lipschitz then \mathbf{q} is called an (L, C) -*almost geodesic*.

Every (L, C) -quasi-geodesic in a geodesic metric space is at bounded (in terms of L, C) distance from an $(L + C, C)$ -almost geodesic with the same endpoints [BBI01, Proposition 8.3.4].

2.2 Asymptotic cones of a metric space

The notion of asymptotic cone of a metric space was introduced first informally in [Gro81], then more rigorously in [dDW84] and [Gro93]. It is meant to represent an image of a given metric space, seen from an infinite distance. To define such a limit image, one needs the notion of *non-principal ultrafilter*. This is a finitely additive measure ω defined on the set of all subsets of \mathbb{N} (or, more generally, of a countable set) and taking values in $\{0, 1\}$, such that $\omega(F) = 0$ for every finite subset F of \mathbb{N} .

Convention 2.3. Throughout the paper all ultrafilters are nonprincipal, therefore we will omit mentioning it each time.

Notation 2.4 (from [DS05b], Appendix). Let A_n and B_n be two sequences of objects and let \mathcal{R} be a relation that can be established between A_n and B_n for every $n \in \mathbb{N}$. We write $A_n \mathcal{R}_\omega B_n$ if and only if $A_n \mathcal{R} B_n$ ω -almost surely, that is

$$\omega(\{n \in \mathbb{N} \mid A_n \mathcal{R} B_n\}) = 1.$$

Examples: $=_\omega, <_\omega, \subset_\omega$.

Given an ultrafilter ω and a sequence (x_n) in a topological space X , an ω -limit $\lim_{\omega} x_n$ of the sequence (x_n) is an element $x \in X$ such that for every neighborhood \mathcal{N} of x , $x_n \in_{\omega} \mathcal{N}$.

In a Hausdorff separable space if the ω -limit of a sequence exists then it is unique.

The following property emphasizes the main interest of ultrafilters and ω -limits.

Proposition 2.5 ([Bou65]). *If (x_n) is contained in a compact space then it has an ω -limit.*

Given a space X one can define its *ultrapower* X^{ω} as the quotient $X^{\mathbb{N}} / \approx$, where $(x_n) \approx (y_n)$ if $x_n =_{\omega} y_n$.

Let now (X, dist) be a metric space, e a fixed element in its ultrapower X^{ω} , (e_n) a representative of e , and $d = (d_n)$ a sequence of numbers in $(0, +\infty)$ such that $\lim_{\omega} d_n = +\infty$.

Consider

$$\mathcal{S}_e(X) = \left\{ (x_n) \in X^{\mathbb{N}} ; \exists M_x \text{ such that } \text{dist}(x_n, e_n) \leq_{\omega} M_x d_n \right\}. \quad (2)$$

Define the equivalence relation

$$(x_n) \sim (y_n) \Leftrightarrow \lim_{\omega} \left(\frac{\text{dist}(x_n, y_n)}{d_n} \right) = 0.$$

The quotient space $\mathcal{S}_e(X) / \sim$ is denoted by $\text{Con}_{\omega}(X; e, d)$ and it is called *the asymptotic cone of X with respect to the ultrafilter ω , the scaling sequence d and the sequence of observation centers e* . It is endowed with the natural metric dist_{ω} defined by

$$\text{dist}_{\omega}(x, y) = \lim_{\omega} \left(\frac{\text{dist}(x_n, y_n)}{d_n} \right).$$

A sequence of subsets (A_n) in X gives rise to a *limit subset* in the cone, defined by

$$\lim_{\omega} (A_n) = \{ \lim_{\omega} (a_n) \mid a_n \in_{\omega} A_n \}.$$

If $\lim_{\omega} \frac{\text{dist}(e_n, A_n)}{d_n} = +\infty$ then $\lim_{\omega} (A_n) = \emptyset$.

Proposition 2.6 ([dDW84], [KL95]). (1) *Every asymptotic cone is a complete metric space.*

(2) *Every limit subset $\lim_{\omega} (A_n)$, if non-empty, is closed.*

If each set A_n is a geodesic \mathfrak{g}_n with length of order $O(d_n)$ and $\lim_{\omega} (\mathfrak{g}_n)$ is non-empty, then it is a geodesic in $\text{Con}_{\omega}(X; e, d)$. Therefore if X is a geodesic space then every asymptotic cone $\text{Con}_{\omega}(X; e, d)$ is geodesic.

Definition 2.7. We call a geodesic in $\text{Con}_{\omega}(X; e, d)$ which appears as $\lim_{\omega} (\mathfrak{g}_n)$ with \mathfrak{g}_n geodesics in X a *limit geodesic*.

Not every geodesic in $\text{Con}_{\omega}(X; e, d)$ is a limit geodesic, not even in the particular case when X is a group of finite type with a word metric.

Example of group with continuously many non-limit geodesics in an asymptotic cone:

On the two-dimensional unit sphere \mathbb{S}^2 consider a family of horizontal circles, and a family of vertical circles in parallel planes, such that two consecutive circles in each family are at spherical distance $\frac{\pi}{2^k}$, and such that the North and the South points are on one vertical circle, and are at distance $\frac{\pi}{2^k}$ from two respective horizontal circles.

The two families of circles compose a spherical grid Γ'_k . We have that $\Gamma'_k \subset \Gamma'_{k+1}$. Let Γ''_k be the graph obtained from Γ'_k by joining with spherical geodesics all vertices not on the same vertical or horizontal circle, and at distance at most $\frac{\pi}{\sqrt{2}^k}$. Let Γ_k be the graph obtained from Γ''_k by deleting all the vertical edges of length $\frac{\pi}{2^k}$ above the Equator, except the one having the East point $(1, 0, 0)$ as an endpoint, and replacing each of them by a path of double length $\frac{\pi}{2^{k-1}}$. Let dist_k be the shortest-path metric on Γ_k .

Proposition 7.26 from [DS05b] applied to the sequence of graphs $(\Gamma_k, \text{dist}_k)$, and Lemma 7.5 from the same paper imply that there exists a two-generated and recursively presented group G with one asymptotic cone tree-graded, with all pieces isometric to \mathbb{S}^2 . Moreover, from the construction it follows that in each of the pieces, for an appropriate choice of the North, South and East points, all geodesics joining North and South and not containing East are not limit geodesics.

The same argument as in [DS05b, §7] allows in fact to construct a two-generated and recursively presented group with continuously many non-homeomorphic asymptotic cones with the property that continuously many geodesics in each of them are not limit geodesics.

2.3 Relatively hyperbolic groups

We recall the definition of relative hyperbolicity given by B. Farb in [Far98], in its slightly modified version appearing in [Osi06]. Let G be a finitely generated group and let $\{H_1, \dots, H_m\}$ be a collection of finitely generated subgroups of G . We denote by \mathcal{EH} the set $\bigsqcup_{i=1}^m (H_i \setminus \{1\})$. Let S be a finite generating set of G invariant with respect to inversion. The Cayley graph of the group G with respect to S , $\text{Cayley}(G, S)$, is a subgraph of the Cayley graph $\text{Cayley}(G, S \cup \mathcal{EH})$, which has the same set of vertices but a larger set of edges, and is not locally finite. Let $\text{dist}_{S \cup \mathcal{EH}}$ and dist_S be the respective shortest path metrics in $\text{Cayley}(G, S \cup \mathcal{EH})$ and $\text{Cayley}(G, S)$.

Definition 2.8. Let \mathbf{p} be a path in $\text{Cayley}(G, S \cup \mathcal{EH})$. An \mathcal{EH} -component of \mathbf{p} is a maximal sub-path of \mathbf{p} contained in a left coset gH_i , $i \in \{1, 2, \dots, m\}$, $g \in G$.

The path \mathbf{p} is said to be *without backtracking* if it does not have two distinct \mathcal{EH} -components in the same left coset gH_i .

Definition 2.9. The group G is *weakly hyperbolic relative to* $\{H_1, \dots, H_m\}$ if and only if the graph $\text{Cayley}(G, S \cup \mathcal{EH})$ is hyperbolic.

Definition 2.10. The pair $(G, \{H_1, \dots, H_m\})$ satisfies the *Bounded Coset Penetration (BCP) property* if for every $\lambda \geq 1$ there exists $a = a(\lambda)$ such that the following holds. Let \mathbf{p} and \mathbf{q} be two λ -bi-Lipschitz paths without backtracking in $\text{Cayley}(G, S \cup \mathcal{EH})$ such that $\mathbf{p}_- = \mathbf{q}_-$ and $\text{dist}_S(\mathbf{p}_+, \mathbf{q}_+) \leq 1$.

- (1) Suppose that s is an \mathcal{EH} -component of \mathbf{p} such that $\text{dist}_S(s_-, s_+) \geq a$. Then \mathbf{q} has an \mathcal{EH} -component contained in the same left coset as s ;
- (2) Suppose that s and t are two \mathcal{EH} -components of \mathbf{p} and \mathbf{q} , respectively, contained in the same left coset. Then $\text{dist}_S(s_-, t_-) \leq a$ and $\text{dist}_S(s_+, t_+) \leq a$.

Definition 2.11. The group G is *(strongly) hyperbolic relative to* $\{H_1, \dots, H_m\}$ if it is weakly hyperbolic relative to $\{H_1, \dots, H_m\}$ and if $(G, \{H_1, \dots, H_m\})$ satisfies the BCP property.

Both in the case of weak and strong relative hyperbolicity, the subgroups H_1, \dots, H_m are called *peripheral subgroups*. A subgroup conjugate to some H_i , $i \in \{1, \dots, m\}$, is called a *maximal parabolic subgroup*. A subgroup contained in a maximal parabolic subgroup is called *parabolic*.

Definitions 2.12. A group is *elementary* if it has a cyclic subgroup of finite index.

A group G is *properly relatively hyperbolic* with respect to a finite family $\mathcal{H} = \{H_1, \dots, H_n\}$ of finitely generated subgroups, if G is relatively hyperbolic with respect to \mathcal{H} and $G \neq H_i$ for all $i \in \{1, 2, \dots, n\}$. In this case, if G is infinite then all H_i have infinite index in G .

We say that G is *non-relatively hyperbolic* when G is not properly relatively hyperbolic with respect to any finite family \mathcal{H} . In particular, a non-relatively hyperbolic group is not hyperbolic, hence it is non-elementary.

3 Tree-graded metric spaces

3.1 Definition and properties

The notion of tree-graded metric space has been introduced in [DS05b]. In this paper we use the following version of this notion. Recall that a subset A in a geodesic metric space X is called *geodesic* if every two points in A can be joined by a geodesic contained in A .

Definition 3.1. Let \mathbb{F} be a complete geodesic metric space and let \mathcal{P} be a collection of closed geodesic subsets, called *pieces*. Suppose that the following two properties are satisfied:

- (T_1) Every two different pieces have at most one point in common.
- (T_2) Every simple non-trivial geodesic triangle in \mathbb{F} is contained in one piece.

Then we say that the space \mathbb{F} is *tree-graded with respect to \mathcal{P}* .

Remarks 3.2 (pieces need not cover the space). (1) In [DS05b] trivial geodesic triangles are allowed in property (T_2). This is equivalent to asking that \mathbb{F} is covered by the pieces in \mathcal{P} . In the present paper we remove this convention. The reason is that a main purpose when introducing the notion of tree-graded space is to produce a convenient notion of relatively hyperbolic metric space (called asymptotically tree-graded metric space in [DS05b] and in this paper, see Definition 4.1). The condition that pieces cover \mathbb{F} produces some unnatural restrictions for a space to be asymptotically tree-graded (i.e. relatively hyperbolic) with respect to a list of subsets. See Remark 4.10 for details.

- (2) Possibly \mathcal{P} is empty, in which case \mathbb{F} is a real tree.
- (3) When a group G acts transitively on \mathbb{F} (for instance when \mathbb{F} is an asymptotic cone of a group) and G permutes the pieces, the condition that pieces cover \mathbb{F} is automatically satisfied.

All properties of tree-graded spaces in [DS05b, §2.1] hold with the new definition 3.1, as none of the proofs uses the property that pieces cover the space. In particular one has the following results.

Proposition 3.3 ([DS05b], Proposition 2.17). *Condition (T_2) in the definition of a tree-graded space can be replaced by any of the following two conditions:*

- (T'_2) *Let $\mathbf{c} : [0, d] \rightarrow \mathbb{F}$ be an arbitrary topological arc. For every $t \in [0, d]$, if $\mathbf{c}(t)$ is contained in a piece then take $\mathbf{c}[t - a, t + b]$ a maximal sub-arc of \mathbf{c} containing $\mathbf{c}(t)$ and contained in one piece; if not then take $t - a = t + b = t$.*

Every other topological arc with the same endpoints as \mathbf{c} must contain the points $\mathbf{c}(t - a)$ and $\mathbf{c}(t + b)$.

(T_2'') Every non-trivial simple loop in \mathbb{F} is contained in one piece.

Lemma 3.4 ([DS05b], §2.1). *Let x be an arbitrary point in \mathbb{F} and let T_x be the set of points $y \in \mathbb{F}$ which can be joined to x by a topological arc intersecting every piece in at most one point.*

- (1) *The subset T_x is a real tree.*
- (2) *For every $y \in T_x$, $T_y = T_x$.*
- (3) *Every topological arc joining two points in T_x is contained in T_x .*
- (4) *T_x is a closed subset of \mathbb{F} .*

Definition 3.5. A subset T_x as in Lemma 3.4 is called a *transversal tree* in \mathbb{F} .

Remark 3.6. One can ensure that pieces in a tree-graded space cover it by adding to the list of pieces the transversal trees.

Thus a tree-graded space \mathbb{F} with set of pieces \mathcal{P} in the sense of Definition 3.1 can be seen as tree-graded in the sense of Definition 2.1 in [DS05b] with respect to a set of pieces \mathcal{P}' such that $\mathcal{P}' \setminus \mathcal{P}$ is a collection of real trees.

Definition 3.7. Let \mathbb{F} be a tree-graded space with set of pieces \mathcal{P} . We say that $(\mathbb{F}, \mathcal{P})$ is a tree-graded space *without extra singletons* if no piece M is contained in a distinct piece M' . Since by (T_1) the piece M must be a singleton, this is equivalent to removing from the list of pieces every singleton contained in another piece.

One can define a partial order relation on the set of collections of pieces without extra singletons of a tree-graded space X . If \mathcal{P} and \mathcal{P}' are such, we write $\mathcal{P} \prec \mathcal{P}'$ if for every set $M \in \mathcal{P}$ there exists $M' \in \mathcal{P}'$ such that $M \subset M'$.

We adopt, as in [DS05b], the following

Convention 3.8. A singleton is assumed to be a set with cut-points.

The following result displays the relation between spaces with cut-points and tree-graded spaces.

Lemma 3.9 ([DS05b], Lemma 2.31). *Let X be a complete geodesic metric space containing at least two points and let \mathcal{C} be a non-empty set of global cut-points in X .*

There exists the largest in the sense of \prec collection \mathcal{P} of subsets of X such that

- *(X, \mathcal{P}) is tree-graded without extra singletons;*
- *any piece in \mathcal{P} is either a singleton or a set with no global cut-point from \mathcal{C} .*

Moreover the intersection of any two distinct pieces from \mathcal{P} is either empty or a point from \mathcal{C} .

3.2 Topological bigons and property (T_2)

Definition 3.10. We call *topological bigon* (τ -bigon, in short) formed by two topological arcs \mathbf{g}_1 and \mathbf{g}_2 a union between a sub-arc \mathbf{g}'_1 of \mathbf{g}_1 and a sub-arc \mathbf{g}'_2 of \mathbf{g}_2 such that \mathbf{g}'_1 and \mathbf{g}'_2 intersect only in their endpoints x and y . The *endpoints of the τ -bigon* are the points x and y . The *interior of the τ -bigon* is the set $\mathbf{g}'_1 \cup \mathbf{g}'_2 \setminus \{x, y\}$.

Note that a τ -bigon with non-empty interior cannot be trivial, i.e. reduced to a point.

The results in this section are useful in arguments aiming to prove property (T_2) . In several contexts it proves necessary to deduce from (T_1) , and (T_2) satisfied only for some special type of τ -bigons, the general property (T_2) .

Convention 3.11. When there is no possibility of confusion, we omit to mention that a bigon is topological (or τ -bigon). It will be the case throughout the rest of this section.

Lemma 3.12. *Let \mathfrak{g}_1 and \mathfrak{g}_2 be two topological arcs with common endpoints. Then every point $z \in \mathfrak{g}_1 \setminus \mathfrak{g}_2$ is in the interior of a simple bigon formed by \mathfrak{g}_1 and \mathfrak{g}_2 .*

Proof. For $i = 1, 2$, $\mathfrak{g}_i : [0, \ell_i] \rightarrow Y$ is a topological embedding. Let $t \in [0, \ell_1]$ be such that $\mathfrak{g}_1(t) = z$. The set $K = \mathfrak{g}_1^{-1}(\mathfrak{g}_2([0, \ell_2]))$ is a compact set not containing t . Let r be the maximal element of the compact $K \cap [0, t]$, and let s be the minimal element of the compact $K \cap [t, \ell_1]$. Then $\mathfrak{g}_1(r) = \mathfrak{g}_2(r')$ and $\mathfrak{g}_1(s) = \mathfrak{g}_2(s')$ for some $r', s' \in [0, \ell_2]$. The union of \mathfrak{g}_1 restricted to $[r, s]$ with \mathfrak{g}_2 restricted to $[r', s']$ is a simple bigon formed by \mathfrak{g}_1 and \mathfrak{g}_2 , containing z in its interior. \square

Lemma 3.13. *Let Y be a complete metric space and let \mathcal{B} be a collection of closed subsets of Y , \mathcal{B} with property (T_1) .*

Let \mathfrak{g}_1 and \mathfrak{g}_2 be two topological arcs with common endpoints and with the property that any non-trivial simple bigon formed by \mathfrak{g}_1 and \mathfrak{g}_2 is contained in a subset in \mathcal{B} .

If \mathfrak{g}_1 is contained in $B \in \mathcal{B}$ then \mathfrak{g}_2 is contained in B .

Proof. Take z an arbitrary point in $\mathfrak{g}_2 \setminus \mathfrak{g}_1$. By Lemma 3.12 the point z is in the interior of a simple bigon formed by \mathfrak{g}_1 and \mathfrak{g}_2 , of endpoints z_1, z_2 . By hypothesis this bigon is contained in a subset $B_z \in \mathcal{B}$. As $\{z_1, z_2\}$ is in $B \cap B_z$ it follows by (T_1) that $B_z = B$ and $z \in B$. \square




Figure 1: Step 1.

Proposition 3.14. *Let Y be a complete metric space and let \mathcal{B} be a collection of closed subsets of Y , \mathcal{B} with property (T_1) .*

Let \mathcal{L}_1 and \mathfrak{g}_1 be two topological arcs with common endpoints u, v . Let \mathcal{L}_2 and \mathfrak{g}_2 be two, possibly identical, topological arcs with common endpoints v, w . Assume that:

- (1) $\mathcal{L}_1 \cap \mathcal{L}_2 = \{v\}$;
- (2) $\mathfrak{g}_1 \cap \mathfrak{g}_2$ contains a point $a \neq v$;
- (3) all non-trivial simple bigons formed either by \mathfrak{g}_1 and \mathfrak{g}_2 , or by \mathfrak{g}_i and \mathcal{L}_i , $i = 1, 2$, are contained in a subset in \mathcal{B} .

Then the bigon formed by \mathfrak{g}_1 and \mathfrak{g}_2 with endpoints a and v is contained in a subset in \mathcal{B} .

Proof. STEP 1. We prove that there exists $b \in \mathfrak{g}_1 \cap \mathfrak{g}_2 \setminus \{a\}$, b between v and a on both \mathfrak{g}_i , $i = 1, 2$, such that the bigon formed by \mathfrak{g}_1 and \mathfrak{g}_2 of endpoints a, b is contained in some $B \in \mathcal{B}$.

Hypothesis (1) implies that either $a \notin \mathcal{L}_1$ or $a \notin \mathcal{L}_2$. Without loss of generality we may assume that $a \notin \mathcal{L}_1$. Then a is in the interior of a simple bigon formed by \mathcal{L}_1 and \mathfrak{g}_1 , of endpoints x and y (with y closer to v than x). Property (3) implies that this bigon is contained in a set $B_1 \in \mathcal{B}$.

If $y \in \mathfrak{g}_2$ then take $b = y$.

Assume that $y \notin \mathfrak{g}_2$. Then y is in the interior of a simple bigon formed by \mathfrak{g}_1 and \mathfrak{g}_2 , of endpoints y_1, y_2 (with y_2 closer to v than y_1 , and y_1 between y and a). By (3) this bigon is contained in some $B_2 \in \mathcal{B}$. The intersection $B_1 \cap B_2$ contains $\{y, y_1\}$ hence by (T_1) we have that $B_1 = B_2 = B$. Take $b = y_2$.

It remains to show that the sub-arc \mathfrak{g}'_2 of \mathfrak{g}_2 in between a and b is contained in B . The sub-arc \mathfrak{g}'_1 of \mathfrak{g}_1 with the same endpoints is known to be contained in B . By property (3) we can apply Lemma 3.13 to \mathfrak{g}'_1 and \mathfrak{g}'_2 and we obtain that $\mathfrak{g}'_2 \subset B$.

STEP 2. Consider the closest to v point b as in Step 1.

Assume that $b \neq v$. Step 1 applied with a replaced by b implies that there exists $c \in \mathfrak{g}_1 \cap \mathfrak{g}_2 \setminus \{b\}$, c closer to v than b such that the bigon formed by \mathfrak{g}_1 and \mathfrak{g}_2 of endpoints b, c is contained in some $B' \in \mathcal{B}$.

Since $b \neq v$ it cannot be contained simultaneously in \mathcal{L}_1 and in \mathcal{L}_2 . Assume that $b \notin \mathcal{L}_1$. Then b is in the interior of a simple bigon formed by \mathfrak{g}_1 and \mathcal{L}_1 . According to (3) this bigon is contained in some $B'' \in \mathcal{B}$. The intersections $B \cap B''$ and $B' \cap B''$ both contain non-trivial sub-arcs of \mathfrak{g}_1 , therefore $B = B'' = B'$. Thus the point c satisfies all the properties of b and it is closer to v , which contradicts the choice of b .

We conclude that $b = v$. □

4 Asymptotically tree-graded metric spaces

4.1 Definition and properties

Let (X, dist) be a geodesic metric space and let $\mathcal{A} = \{A_i \mid i \in I\}$ be a collection of subsets of X . In every asymptotic cone $\text{Con}_\omega(X; e, d)$, we consider the collection \mathcal{A}_ω of subsets

$$\left\{ \lim_{\omega} (A_{i_n}) ; i = (i_n)^\omega \in I^\omega \text{ such that } \exists M_i \text{ with the property } \text{dist}(e_n, A_{i_n}) \leq_\omega M_i d_n \right\}.$$

Definition 4.1. The metric space X is *asymptotically tree-graded (ATG)* with respect to \mathcal{A} if every asymptotic cone $\text{Con}_\omega(X; e, d)$ is tree-graded with respect to \mathcal{A}_ω .

The space X is *properly asymptotically tree-graded* with respect to \mathcal{A} if it is asymptotically tree-graded with respect to \mathcal{A} and if no $A \in \mathcal{A}$ contains X in a tubular neighborhood of it.

The notion of ATG metric space is meant to extend the notion of (strong) relative hyperbolicity from groups to metric spaces. Theorem 1.2 emphasizes that it is the correct definition to work with.

Remark 4.2. It is easy to see that if X is ATG with respect to \mathcal{A} and $\tau > 0$ then X is ATG with respect to $\{\mathcal{N}_\tau(A_i) ; i \in I\}$. See Lemma 4.12 below for a more general statement.

The notion of ATG metric space can also be defined by a list of geometric conditions, without involving asymptotic cones. First some notation and terminology.

Notation 4.3. For every quasi-geodesic \mathbf{p} in a metric space X , we denote the origin of \mathbf{p} by \mathbf{p}_- and the endpoint of \mathbf{p} by \mathbf{p}_+ .

Given $r > 0$ we denote by $\check{\mathbf{p}}_r$ the set $\mathbf{p} \setminus \mathcal{N}_r(\{\mathbf{p}_-, \mathbf{p}_+\})$.

We say that a metric space P is a *geodesic (quasi-geodesic) k -gonal line* if it is a union of k geodesics (quasi-geodesics) $\mathbf{q}_1, \dots, \mathbf{q}_k$ such that $(\mathbf{q}_i)_+ = (\mathbf{q}_{i+1})_-$ for $i = 1, \dots, k-1$. If moreover $(\mathbf{q}_k)_+ = (\mathbf{q}_1)_-$ then we say that P is a *geodesic (quasi-geodesic) k -gon*.

Let P be a quasi-geodesic polygon, with set of vertices \mathcal{V} . Points in $P \setminus \mathcal{V}$ are called *interior points* of P .

Notation 4.4. Given a vertex $x \in \mathcal{V}$ and \mathbf{q}, \mathbf{q}' the consecutive edges of P such that $x = \mathbf{q}_+ = \mathbf{q}'_-$, we denote the polygonal line $P \setminus (\mathbf{q} \cup \mathbf{q}')$ by $\mathcal{O}_x(P)$. When there is no possibility of confusion we simply denote it by \mathcal{O}_x .

Let $p \in P$. The *inscribed radius in p* with respect to P is either the distance from p to the set \mathcal{O}_p , if p is a vertex, or the distance from p to the set $P \setminus \mathbf{q}$ if p is contained in the interior of the edge \mathbf{q} (see Figure 2, taken from [DS05b]).




Figure 2: Properties (F_1) and (F_2) .

Definition 4.5 (fat polygons). Let $\vartheta > 0$, $\sigma \geq 1$ and $\nu \geq 4\sigma$. We call a k -gon P with quasi-geodesic edges (ϑ, σ, ν) -fat if the following properties hold:

(F_1) (**large inscribed radii in interior points, large comparison angles**) for every edge \mathfrak{q} we have, with the notation 4.3, that

$$\text{dist}(\check{\mathfrak{q}}_{\sigma\vartheta}, P \setminus \mathfrak{q}) \geq \vartheta;$$

(F_2) (**large inscribed radii in vertices, large edges**) for every vertex x we have

$$\text{dist}(x, \mathcal{O}_x) \geq \nu\vartheta.$$

When $\sigma = 2$ we say that P is (ϑ, ν) -fat.

Lemma 4.6. *Let P be a polygon (ϑ, σ, ν) -fat for some $\vartheta > 0$, $\sigma \geq 1$ and $\nu \geq 4\sigma$. Then any two edges of P without a common vertex are at distance at least ϑ from each other.*

Proof. Let \mathfrak{q} and \mathfrak{q}' be two edges without a common vertex. Assume that there exists a point $a \in \mathfrak{q}$ such that $\text{dist}(a, \mathfrak{q}') < \vartheta$. Property (F_1) implies that $a \in \mathcal{N}_{\sigma\vartheta}(\{x, y\})$, where x, y are the endpoints of \mathfrak{q} . Property (F_2) implies that $\text{dist}(\{x, y\}, \mathfrak{q}') \geq \nu\vartheta$. Therefore $\text{dist}(a, \mathfrak{q}') \geq (\nu - \sigma)\vartheta \geq 3\sigma\vartheta \geq 3\vartheta$. This contradicts the assumption that $\text{dist}(a, \mathfrak{q}') < \vartheta$. \square

Theorem 4.7 ([DS05b], **Theorem 4.1 and Remark 4.2, (3)**). *Let (X, dist) be a geodesic metric space and let $\mathcal{A} = \{A_i \mid i \in I\}$ be a collection of subsets of X . The metric space X is asymptotically tree-graded with respect to \mathcal{A} if and only if the following properties are satisfied:*

- (α_1) *For every $\delta > 0$ the diameters of the intersections $\mathcal{N}_\delta(A_i) \cap \mathcal{N}_\delta(A_j)$ are uniformly bounded for all $i \neq j$.*
- (α_2) *There exists ε in $[0, \frac{1}{2})$ and $M > 0$ such that for every geodesic \mathfrak{g} of length ℓ and every $A \in \mathcal{A}$ with $\mathfrak{g}(0), \mathfrak{g}(\ell) \in \mathcal{N}_{\varepsilon\ell}(A)$ we have $\mathfrak{g}([0, \ell]) \cap \mathcal{N}_M(A) \neq \emptyset$.*
- (α_3) *For every $k \geq 2$ there exist $\vartheta > 0$, $\nu \geq 8$ and $\chi > 0$ such that every k -gon P in X with geodesic edges which is (ϑ, ν) -fat satisfies $P \subset \mathcal{N}_\chi(A)$ for some $A \in \mathcal{A}$.*

Remarks 4.8 ([DS05b], **Theorem 4.1 and Remark 4.2**). (1) Property (α_2) from Theorem 4.7 is slightly modified, compared to the similar property in Theorem 4.1 in [DS05b]. Still, as stated it implies property (α_2^ε) from [DS05b, Remark 4.2, (3)], which accounts for the accuracy of the modified statement.

- (2) As a necessary condition, (α_2) can be strengthened to “for every ε from $[0, \frac{1}{2})$ there exists $M > 0$ such that etc.”

Notation 4.9. We denote by diam_δ an uniform bound provided by property (α_1) for an arbitrary $\delta \geq 0$.

Remarks 4.10 (on the condition that pieces cover the space). (1) If in property (T_2) of the definition of tree-graded spaces (Definition 3.1) we allow for trivial geodesic triangles, that is if we ask that pieces cover a tree-graded space, then in Theorem 4.7 the following condition has to be added:

- (α_0) there exists $\tau \geq 0$ such that $X = \bigcup_{A \in \mathcal{A}} \mathcal{N}_\tau(A)$.

If (X, \mathcal{A}) satisfy only the conditions (α_1) , (α_2) , (α_3) but not (α_0) then it suffices to add some singletons to \mathcal{A} in order to ensure (α_0) . Indeed, for some $\tau > 0$ consider in $X \setminus \bigcup_{A \in \mathcal{A}} \mathcal{N}_\tau(A)$ a maximal subset \wp with the property that $\text{dist}(p, p') \geq \tau$ for every $p, p' \in \wp$. The space X coincides with $\bigcup_{A \in \mathcal{A}} \mathcal{N}_\tau(A) \cup \bigcup_{p \in \wp} \mathcal{N}_\tau(\{p\})$. Properties (α_1) and (α_2) are obviously satisfied by singletons, whence X is ATG with respect to $\mathcal{A}' = \mathcal{A} \cup \{\{p\} ; p \in \wp\}$; moreover \mathcal{A}' also satisfies (α_0) .

- (2) Let \mathbb{H}^3 be the 3-dimensional real hyperbolic space and let $(Hbo_n)_{n \in \mathbb{N}}$ be a countable collection of pairwise disjoint open horoballs. The complementary set $X_0 = X \setminus \bigsqcup_{n \in \mathbb{N}} Hbo_n$ and the list of boundary horospheres $\mathcal{A} = \{\partial Hbo_n ; n \in \mathbb{N}\}$ is the typical example one has in mind when trying to define relative hyperbolicity for metric spaces. The pair (X_0, \mathcal{A}) does not in general satisfy (α_0) , one has to add singletons to \mathcal{A} to ensure that property. In order to remove this inconvenient, we give up the condition of pieces covering the space in Definition 3.1.

Remark 4.11. If X is a metric space ATG with respect to \mathcal{A} , and a group G acts \mathcal{K} -transitively (in the sense of Definition 2.1, with $\mathcal{K} \geq 0$) by isometries on X , G permuting the subsets in \mathcal{A} , then property (α_0) is satisfied with $M = \mathcal{K}$.

It is for instance the case when X is itself a group and \mathcal{A} is the collection of left cosets of a family of subgroups.

Lemma 4.12. *Let X be ATG with respect to \mathcal{A} . Let \mathcal{B} be a collection of subsets of X such that there exists a constant $K \geq 0$ and a bijection $\phi : \mathcal{A} \rightarrow \mathcal{B}$ verifying $\text{dist}_H(A, \phi(A)) \leq K$. Then X is ATG with respect to \mathcal{B} .*

Proof. Properties (α_1) and (α_3) for \mathcal{B} follow in an obvious way from the same properties for \mathcal{A} .

Let ε and M be the constants given by (α_2) for \mathcal{A} . We prove (α_2) for \mathcal{B} , with $\varepsilon' = \frac{\varepsilon}{2}$ and $M' = \max(\frac{K}{\varepsilon} + K, M + K)$.

Consider a geodesic \mathbf{g} of length ℓ and $B \in \mathcal{B}$ such that $\mathbf{g}(0), \mathbf{g}(\ell)$ are in $\mathcal{N}_{\varepsilon'\ell}(B) \subset \mathcal{N}_{\varepsilon'\ell+K}(\phi^{-1}(B))$.

If $K \geq \frac{\varepsilon}{2}\ell$ then $\mathbf{g} \subset \mathcal{N}_{\frac{K}{\varepsilon}}(\{\mathbf{g}(0), \mathbf{g}(\ell)\}) \subset \mathcal{N}_{\frac{K}{\varepsilon}+K}(B)$.

If $K < \frac{\varepsilon}{2}\ell$ then property (α_2) for \mathcal{A} implies that $\mathbf{g}([0, \ell])$ intersects $\mathcal{N}_M(\phi^{-1}(B))$, hence it intersects $\mathcal{N}_{M+K}(B)$. \square

4.2 Property (T_2) and polygons with limit edges

Property (α_3) in the definition of an ATG metric space X is used to prove property (T_2) in any of its asymptotic cones. If X is such that any geodesic in an asymptotic cone of it is a limit geodesic (for instance if X is a CAT(0) metric space) then it suffices to have (α_3) for $k = 6$, that is:

- (β_3) there exists $\vartheta > 0$, $\nu \geq 8$ and $\chi > 0$ such that any geodesic hexagon (ϑ, ν) -fat is contained in $\mathcal{N}_\chi(A)$, for some $A \in \mathcal{A}$.

This is due to the following general fact.

Proposition 4.13. *Let (X, dist) be a geodesic metric space and let $\vartheta > 0$ and $\nu \geq 8$ be two constants. In any asymptotic cone $\text{Con}_\omega(X; e, d)$, any simple non-trivial triangle whose edges are limit geodesics can be written as $\lim_\omega(H_n)$ where H_n are (ϑ, ν) -fat geodesic hexagons in X .*

Proof. Consider a non-trivial simple geodesic triangle Δ whose edges $[a, b]$, $[b, c]$ and $[c, a]$ appear as limits of sequences $[a_n, b'_n]$, $[b_n, c'_n]$ and $[c_n, a'_n]$ of geodesics in X . We have that $\text{dist}(a_n, a'_n)$, $\text{dist}(b_n, b'_n)$ and $\text{dist}(c_n, c'_n)$ are of order $o(d_n)$ ω -almost surely.

Define the positive number d_n^A to be the maximum between $\text{dist}([a_n, b'_n], [a'_n, c_n])$ and $\nu\vartheta$. Note that $d_n^A > 0$ and that $d_n^A =_\omega o(d_n)$. Take a_n^1 to be the farthest from a_n point on $[a_n, b'_n]$ at distance d_n^A from $[a'_n, c_n]$. Consider then a_n^2 the farthest from a'_n point on $[a'_n, c_n]$ at distance d_n^A from a_n^1 . Obviously $\text{dist}(a_n^1, a_n^2) = d_n^A$.

The pairs of points (b_n^1, b_n^2) in $[b_n, c'_n] \times [b'_n, a_n]$, and respectively (c_n^1, c_n^2) in $[c_n, a'_n] \times [c'_n, b_n]$ are chosen similarly. Because the limit triangle Δ is simple, it follows that the sets $\{a_n, a'_n, a_n^1, a_n^2\}$, $\{b_n, b'_n, b_n^1, b_n^2\}$ and $\{c_n, c'_n, c_n^1, c_n^2\}$ have ω -almost surely diameters of order $o(d_n)$. It follows that the sequence of geodesic hexagons H_n of vertices $a_n^1, b_n^2, b_n^1, c_n^2, c_n^1, a_n^2$ with edges $[a_n^1, b_n^2] \subset [a_n, b'_n]$, $[b_n^1, c_n^2] \subset [b_n, c'_n]$, $[c_n^1, a_n^2] \subset [c_n, a'_n]$, has the property that $\lim_\omega (H_n)$ is Δ . It remains to prove that H_n is ω -almost surely (ϑ, ν) -fat.

(F₁) The edge $[a_n^1, a_n^2]$ is at distance $O(d_n)$ from $[b_n^2, b_n^1] \cup [b_n^1, c_n^2] \cup [c_n^2, c_n^1]$. The existence of a point in $[a_n^1, a_n^2] \setminus \mathcal{N}_{2\vartheta}(\{a_n^1, a_n^2\})$ at distance less than ϑ either from $[a_n^1, b_n^2]$ or from $[a_n^2, c_n^1]$ would contradict the choice of a_n^1 or respectively the choice of a_n^2 .

In the same manner it can be shown that the edges $[b_n^1, b_n^2]$ and $[c_n^1, c_n^2]$ satisfy the property (F₁).

The edge $[a_n^1, b_n^2]$ is at distance $O(d_n)$ from $[c_n^1, c_n^2]$. The choice of a_n^1 and of the pair (b_n^1, b_n^2) implies that $[a_n^1, b_n^2]$ is at distance at least $\nu\vartheta$ from $[b_n^1, c_n^2] \cup [c_n^1, a_n^2]$.

If a point x in $[a_n^1, b_n^2] \setminus \mathcal{N}_{2\vartheta}(\{a_n^1, b_n^2\})$ is at distance $\leq \vartheta$ from a point $y \in [a_n^1, a_n^2]$ then $\text{dist}(a_n^1, y) \geq 2\vartheta - \vartheta \geq \text{dist}(x, y)$, hence $\text{dist}(x, a_n^2) \leq \text{dist}(x, y) + \text{dist}(y, a_n^2) \leq \text{dist}(a_n^1, a_n^2)$, contradicting the choice of a_n^1 . Likewise it is shown that the existence of a point in $[a_n^1, b_n^2] \setminus \mathcal{N}_{2\vartheta}(\{a_n^1, b_n^2\})$ at distance $\leq \vartheta$ from $[b_n^2, b_n^1]$ contradicts the choice of b_n^2 .

Similar arguments give that the edges $[b_n^1, c_n^2]$ and $[c_n^1, a_n^2]$ satisfy (F₁).

(F₂) The distance from a_n^1 to $[a_n^2, c_n^1]$ is at least $\nu\vartheta$ by construction, while the distance to $[b_n^2, b_n^1] \cup [b_n^1, c_n^2] \cup [c_n^2, c_n^1]$ is $O(d_n)$. The same kind of argument shows that (F₂) is satisfied ω -almost surely by all the vertices of H_n . \square

In general not every geodesic in an asymptotic cone is a limit geodesic (see the example in the end of Section 2.2). Thus, in order to ensure property (T₂) in every asymptotic cone, in [DS05b] property (α_3) in full generality is used, together with the following result.

Lemma 4.14 ([DS05b], Proposition 3.34). *Let Δ be an arbitrary simple geodesic triangle in $\text{Con}_\omega(X; e, d)$. For every $\varepsilon > 0$ sufficiently small there exists $k_0 = k_0(\varepsilon)$ and a simple geodesic triangle Δ_ε with the following properties:*

- (a) $\text{dist}_H(\Delta, \Delta_\varepsilon) \leq \varepsilon$;
- (b) Δ_ε contains the midpoints of the edges of Δ ;
- (c) for every $\vartheta > 0$ and $\nu \geq 8$ the triangle Δ_ε can be written as $\lim_\omega (P_n^\varepsilon)$, where ω -almost surely each P_n^ε is a geodesic k -gon in X , for some $k \leq k_0$, and P_n^ε is (ϑ, ν) -fat.

Remark 4.15. If Δ is non-trivial then the set of midpoints of edges of Δ has cardinal 3, hence the triangles Δ_ε are also non-trivial.

In this section we prove that if property (T₁) holds in every asymptotic cone, then (β_3) suffices to deduce (T₂). To this purpose, consider the following property in an asymptotic cone $\text{Con}_\omega(X; e, d)$:

(Π_k) every simple non-trivial k -gon with edges limit geodesics is contained in a subset from \mathcal{A}_ω .

Corollary 4.16. *Assume that in an asymptotic cone $\text{Con}_\omega(X; e, d)$ a collection \mathcal{A}_ω of closed subsets satisfies properties (T_1) and (Π_k) for every $k \in \mathbb{N}$, $k \geq 3$. Then \mathcal{A}_ω satisfies (T_2) .*

Proof. Consider a simple non-trivial geodesic triangle Δ in $\text{Con}_\omega(X; e, d)$. By Lemma 4.14 for every large enough $k \in \mathbb{N}$ there exists a simple non-trivial geodesic triangle Δ_k at Hausdorff distance $\leq \frac{1}{k}$ from Δ , containing the midpoints of its edges, moreover $\Delta_k = \lim_\omega \left(P_n^{(k)} \right)$, where $P_n^{(k)}$ is n - ω -almost surely a geodesic m -gon, $m = m(k)$. By property (Π_m) the triangle Δ_k is contained in some $A_k \in \mathcal{A}_\omega$. All A_k contain the midpoints of the edges of Δ . Property (T_1) implies that there exists $A \in \mathcal{A}_\omega$ such that $A_k = A$ for all k . All Δ_k are in A , Δ is the limit of Δ_k in the Hausdorff distance, and A is closed, therefore $\Delta \subset A$. \square

In view of Corollary 4.16 it suffices to prove (Π_k) for all $k \geq 3$ to obtain property (T_2) .

Obviously (Π_k) implies (Π_i) for every $i < k$. It turns out that with the additional assumption that (T_1) is satisfied, the converse implication also holds.

Lemma 4.17. *Assume that in an asymptotic cone $\text{Con}_\omega(X; e, d)$, the collection of subsets \mathcal{A}_ω satisfies the properties (T_1) and (Π_3) . Then for every $k \geq 3$ property (Π_k) holds.*

Proof. We prove property (Π_k) by induction on k . The cases $k = 2$ and $k = 3$ hold by hypothesis. Assume that the statement is true for every $k \leq m - 1$ and consider a simple non-trivial geodesic m -gon P in $\text{Con}_\omega(X; e, d)$, $m \geq 4$, with edges limit geodesics.

Let $[x, y]$ and $[y, z]$ be two consecutive edges of P , in clockwise order. Denote by \mathcal{L}_1 the union of the two edges $[x, y] \cup [y, z]$ of P , and by \mathcal{L} the union of the other $m - 2$ edges of P , in clockwise order.

Consider a limit geodesic \mathbf{g} joining x and z . If \mathbf{g} coincides with \mathcal{L}_1 or with \mathcal{L} then P is a simple geodesic polygon with at most $m - 1$ edges, all of them limit geodesics. By the inductive hypothesis P is contained in a subset A in \mathcal{A}_ω .

Assume that \mathbf{g} does not coincide either with \mathcal{L}_1 or with \mathcal{L} .




Figure 3: Step 1 in the proof of Lemma 4.17

STEP 1. We prove that $\mathbf{g} \cup \mathcal{L}$ is contained in some $A \in \mathcal{A}_\omega$.

Let $\alpha \in \mathcal{L} \setminus \mathbf{g}$. Lemma 3.12 implies that α is in the interior of a simple τ -bicon formed by \mathcal{L} and \mathbf{g} , of endpoints a and b , with a closer to x than b . This τ -bicon is a geodesic polygon with at most $m - 1$ edges which are limit geodesics, therefore by the inductive hypothesis it is contained in a subset $A \in \mathcal{A}_\omega$.

If $a = x$ and $b = z$ then $\mathbf{g} \cup \mathcal{L}$ is a simple τ -bicon and it is contained in some $A \in \mathcal{A}_\omega$ by the inductive hypothesis. Assume therefore that $(a, b) \neq (x, z)$. Without loss of generality we may assume that $a \neq x$.

We apply Proposition 3.14 to \mathcal{L}_1 , $\mathbf{g}_1 = \mathbf{g}$, and $\mathbf{g}_2 = \mathcal{L}_2$ the sub-arc of \mathcal{L} in between x and α . Property (3) is satisfied by the hypothesis of the induction. It follows that the τ -bigon formed by \mathbf{g} and \mathcal{L} of endpoints x and a is contained in some $A_1 \in \mathcal{A}_\omega$.

The point a is in $\mathcal{L} \setminus \mathcal{L}_1$, hence it is in $\mathbf{g} \setminus \mathcal{L}_1$. By Lemma 3.12, a is in the interior of some τ -bigon formed by \mathcal{L}_1 and \mathbf{g} , and by (Π_3) this τ -bigon is in a subset $A'_1 \in \mathcal{A}_\omega$. Since $A'_1 \cap A$ and $A'_1 \cap A_1$ contain non-trivial sub-arcs of \mathbf{g} property (T_1) implies that $A = A'_1 = A_1$.

If moreover $b \neq z$, a similar argument gives that the τ -bigon formed by \mathbf{g} and \mathcal{L} of endpoints z and b is contained in A (see Figure 3).

We conclude that $\mathbf{g} \cup \mathcal{L}$ is contained in A .

STEP 2. We prove that \mathcal{L}_1 is also contained in A . Property (Π_3) implies that any non-trivial simple topological bigon formed by \mathcal{L}_1 and \mathbf{g} is contained in a subset in \mathcal{A}_ω . We apply Lemma 3.13 to \mathcal{L}_1 and $\mathbf{g} \subset A$ and we conclude that $\mathcal{L}_1 \subset A$. Consequently A contains $\mathcal{L}_1 \cup \mathcal{L} = P$. \square

Corollary 4.18. *Assume that in an asymptotic cone $\text{Con}_\omega(X; e, d)$, the collection of closed subsets \mathcal{A}_ω satisfies properties (T_1) and (Π_3) . Then \mathcal{A}_ω satisfies property (T_2) .*

Corollary 4.19. *Let X be a geodesic metric space and \mathcal{A} a collection of subsets in X , such that (β_3) is satisfied and such that in any asymptotic cone $\text{Con}_\omega(X; e, d)$, the collection of limit subsets \mathcal{A}_ω satisfies property (T_1) . Then \mathcal{A}_ω satisfies property (T_2) .*

Note that the only thing missing in Corollary 4.19 to conclude that X is ATG with respect to \mathcal{A} is that \mathcal{A}_ω is composed of geodesic subsets.

Another useful consequence of Proposition 4.13 is the following.

Corollary 4.20. *Let (X, dist) be a geodesic metric space. Assume that for some $\vartheta > 0$ and $\nu \geq 8$ the set of (ϑ, ν) -fat geodesic hexagons is either empty or composed of hexagons of uniformly bounded diameter. Then X is hyperbolic.*

Proof. Proposition 4.13 implies that in any asymptotic cone of X any simple triangle with edges limit geodesics is trivial. This statement can be extended by induction to all polygons. Indeed, suppose that in any asymptotic cone of X for all $3 \leq k \leq m-1$ all simple k -gons with edges limit geodesics are trivial. Consider P a simple m -gon with edges limit geodesics in some $\text{Con}_\omega(X; e, d)$. Let $[x, y]$ and $[y, z]$ be two consecutive edges of P and let \mathbf{g} be a limit geodesic joining x and z . All τ -bigons formed by $[x, y] \cup [y, z]$ and \mathbf{g} must be trivial by the inductive hypothesis, thus $\mathbf{g} = [x, y] \cup [y, z]$. It follows that P is a simple $(m-1)$ -gon with edges limit geodesics, hence by the inductive hypothesis it is trivial.

Lemma 4.14 and Remark 4.15 imply that in any $\text{Con}_\omega(X; e, d)$ any simple geodesic triangle is trivial. It follows that $\text{Con}_\omega(X; e, d)$ is a real tree, and since this holds for all asymptotic cones we conclude that X is hyperbolic ([Gro93, §2.A], see also [Dru02, §3]). \square

4.3 New definition, useful for the rigidity of relatively hyperbolic groups

We need some variations of the definition of an ATG metric space that we state and prove in what follows. They will play an important part in the proof of the quasi-isometric invariance of relative hyperbolicity.

Theorem 4.21. *In Theorem 4.7 the following modifications in the list of properties can be made:*

(M₁) property (α_3) can be replaced by property (β_3) ;

(M₂) property (α_2) can be either maintained or replaced by one of the following two properties:

- (β_2) there exists $\epsilon > 0$ and $M \geq 0$ such that for any geodesic \mathbf{g} of length ℓ and any $A \in \mathcal{A}$ satisfying $\mathbf{g}(0), \mathbf{g}(\ell) \in \mathcal{N}_{\epsilon\ell}(A)$, the middle third $\mathbf{g}([\frac{\ell}{3}, \frac{2\ell}{3}])$ is contained in $\mathcal{N}_M(A)$;
- (Ω) (**uniform quasi-convexity of pieces**) there exists $t > 0$ and $M_0 \geq 0$ such that for every $A \in \mathcal{A}$, $M \geq M_0$ and $x, y \in \mathcal{N}_M(A)$, every geodesic joining x and y in X is contained in $\mathcal{N}_{tM}(A)$.

Proof. If X is ATG with respect to \mathcal{A} then (Ω) is satisfied according to [DS05b, Lemma 4.3].

Property (β_2) can be obtained for any $\epsilon < \frac{1}{6t}$, where t is the constant from (Ω), by the following argument. Given a geodesic \mathbf{g} and $A \in \mathcal{A}$ as in (β_2), by uniform quasi-convexity \mathbf{g} is contained in $\mathcal{N}_{t\epsilon\ell}(A)$. If $\theta = t\epsilon < \frac{1}{6}$ then by Theorem 4.7 and Remark 4.8 there exists $M = M(\theta)$ such that $\mathbf{g}([0, \frac{\ell}{3}])$ and $\mathbf{g}([\frac{2\ell}{3}, \ell])$ intersect $\mathcal{N}_M(A)$. Uniform convexity implies that $\mathbf{g}([\frac{\ell}{3}, \frac{2\ell}{3}])$ is contained in $\mathcal{N}_{tM}(A)$.

Property (β_3) follows from (α_3) for $k = 6$.

It remains to prove the converse statement, which is that any of the triples of properties (α_1)&(α_2)&(β_3), (α_1)&(β_2)&(β_3) or (α_1)&(Ω)&(β_3) imply that X is ATG with respect to \mathcal{A} .

Lemma 4.22 ([DS05b], Lemma 4.3). *Properties (α_1) and (α_2) imply (Ω).*

Lemma 4.23. *Properties (α_1) and (β_2) imply (Ω) with M_0 equal to the constant M in (β_2).*

Proof. Suppose by contradiction that for every $n \in \mathbb{N}^*$ there exists $A_n \in \mathcal{A}$, $M_n \geq M$ and $x_n, y_n \in \mathcal{N}_{M_n}(A_n)$ such that a geodesic $[x_n, y_n]$ is not contained in $\mathcal{N}_{M_n}(A_n)$. For each $n \in \mathbb{N}^*$ we define D_n to be the infimum over the distances $\text{dist}(x_n, y_n)$ between pairs of points satisfying the properties above for some set in \mathcal{A} . In what follows we assume that we chose x_n, y_n at distance $\delta_n \leq D_n + 1$ of each other. Since $[x_n, y_n]$ is in $\mathcal{N}_{\delta_n/2}(\{x_n, y_n\}) \subset \mathcal{N}_{\delta_n/2+M_n}(A_n)$ it follows that $\frac{1}{2n-1}\delta_n \geq M_n$. In particular for n large enough $M_n < \epsilon\delta_n$, where $\epsilon > 0$ is the constant in (β_2). It follows that the middle third $[a_n, b_n]$ of $[x_n, y_n]$ is contained in $\mathcal{N}_M(A_n)$. Since $M_n \geq M$, the fact that $[x_n, y_n] \not\subset \mathcal{N}_{M_n}(A_n)$ implies that either $[x_n, a_n]$ or $[b_n, y_n]$ is not contained in $\mathcal{N}_{M_n}(A_n)$. It follows that $D_n \leq \frac{\delta_n}{n} \leq \frac{D_n+1}{3}$, hence that the sequence (D_n) is uniformly bounded. This contradicts the fact that $D_n \geq (2n-1)M$. \square

Lemma 4.24. *Let P be a geodesic k -gon with two consecutive edges $[x, y]$ and $[y, z]$, such that $\text{dist}(x, [y, z]) = \text{dist}(x, y)$. If P is (ϑ, ν) -fat then the $(k+1)$ -gon P' obtained from P by adding as a vertex the point $v \in [x, y]$ with $\text{dist}(v, y) = \frac{\nu\vartheta}{2}$ is $(\vartheta, \frac{\nu}{2})$ -fat.*

Proof. Property (F_1) is preserved for the edges distinct from $[x, y]$. For $[x, v]$ and $[v, y]$ property (F_1) follows property (F_1) of $[x, y]$ in P and from the fact that $[x, v] \setminus \mathcal{N}_{2\vartheta}(\{x, v\})$ is at distance 2ϑ from $[v, y]$, and the same for $[v, y] \setminus \mathcal{N}_{2\vartheta}(\{v, y\})$ and $[x, v]$.

Property (F_2) holds for all the vertices different from x, v, y , by property (F_2) in P .

The polygonal line $\mathcal{O}_x(P') = \mathcal{O}_x(P) \cup [v, y]$ is in the $\frac{\nu\vartheta}{2}$ -tubular neighborhood of $\mathcal{O}_x(P)$, hence at distance at least $\frac{\nu\vartheta}{2}$ from x .

The polygonal line $\mathcal{O}_y(P')$ is equal to $\mathcal{O}_y(P) \cup [x, v]$. The line $\mathcal{O}_y(P)$ is at distance $\geq \nu\vartheta$ from y and $[x, v]$ is at distance $\nu\vartheta$ from y .

Finally, $\mathcal{O}_v(P') = \mathcal{O}_y(P) \cup [y, z]$. Since $\text{dist}(v, y) = \frac{\nu\vartheta}{2}$ it follows that $\text{dist}(v, \mathcal{O}_y) \geq \frac{\nu\vartheta}{2}$. If there exists $p \in [y, z]$ such that $\text{dist}(v, p) < \frac{\nu\vartheta}{2}$ then $\text{dist}(x, p) \leq \text{dist}(x, v) + \text{dist}(v, p) < \text{dist}(x, v) + \frac{\nu\vartheta}{2} = \text{dist}(x, y)$. This contradicts the hypothesis that $\text{dist}(x, [y, z]) = \text{dist}(x, y)$. \square

Lemma 4.25. *Properties (α_1), (Ω) and (β_3) imply (α_2) for small enough $\epsilon > 0$, and (β_2).*

Proof. Assume that (\mathfrak{Q}) and (β_3) are satisfied. Let $\mathbf{g} : [0, \ell] \rightarrow X$ be a geodesic with endpoints $x = \mathbf{g}(0)$ and $y = \mathbf{g}(\ell)$ contained in $\mathcal{N}_{\varepsilon\ell}(A)$ for some $A \in \mathcal{A}$. According to (\mathfrak{Q}) , the geodesic \mathbf{g} is contained in $\mathcal{N}_{t\varepsilon\ell}(A)$.

Notation 4.26. We denote $t\varepsilon$ by ϵ and we assume in what follows that $\epsilon < \frac{1}{8}$. We denote by D the maximum between $tM_0 + 4\nu\vartheta$, χ and diam_δ (with the notation 4.9) for $\delta = \max(\chi, tM_0)$. Here t and M_0 are the constants appearing in (\mathfrak{Q}) , while ν, ϑ, χ are the constants appearing in (β_3) .

Suppose by contradiction that \mathbf{g} does not intersect $\mathcal{N}_D(A)$. Note that since $\epsilon\ell \geq D$ it follows that $\ell > 8D$.

Consider x' and y' points in A such that $\text{dist}(x, x')$ and $\text{dist}(y, y')$ are at most $\varepsilon\ell$. By (\mathfrak{Q}) , a geodesic \mathbf{g}' joining x' and y' is contained in $\mathcal{N}_{tM_0}(A)$.

Let $c \in \mathbf{g}$ and $c' \in \mathbf{g}'$ be two points such that $\text{dist}(c, c') = \text{dist}(\mathbf{g}, \mathbf{g}')$. Without loss of generality we may suppose that $\text{dist}(x, c) \geq \frac{\ell}{2}$. We may also suppose that $\text{dist}(x, x') = \text{dist}(x, \mathbf{g}')$. In order to transform the 4-gon of vertices x, x', c, c' , into a fat polygon we make the following choices. Let x_1 be the point on \mathbf{g} between x and c which is farthest from x and at distance at most $2\nu\vartheta$ from $[x, x']$, where ν and ϑ are the constants from (β_3) . Obviously $\text{dist}(x_1, [x, x']) = 2\nu\vartheta$. Let x_2 be the farthest from x point on $[x, x']$ which is at distance $2\nu\vartheta$ from x_1 .

We prove in the sequel that the geodesic pentagon of vertices x_1, x_2, x', c', c is $(\vartheta, 2\nu)$ -fat. To simplify we shall denote its edges by $[v, w]$ if v, w are two consecutive vertices, keeping in mind that $[x_1, c] \subset \mathbf{g}$ and that $[x', c'] \subset \mathbf{g}'$.

(F₁) A point in $[c, c'] \setminus \mathcal{N}_{2\vartheta}(\{c, c'\})$ is at distance at least $(\frac{1}{2} - 2\epsilon)\ell$ from $[x, x']$, hence at distance at least $(\frac{1}{2} - 2\epsilon)\ell - 2\nu\vartheta$ of $[x_1, x_2]$. Since $\ell > 8D > 32\nu\vartheta$, it follows that $[c, c'] \setminus \mathcal{N}_{2\vartheta}(\{c, c'\})$ is at distance at least ϑ from $[x_1, x_2] \cup [x_2, x']$.

The choice of c, c' implies that all points in $[c, c'] \setminus \mathcal{N}_{2\vartheta}(\{c, c'\})$ are at distance at least 2ϑ from \mathbf{g} and from \mathbf{g}' .

The points in $[x_1, c] \setminus \mathcal{N}_{2\vartheta}(\{c, c'\})$ are at distance at least $D - tM_0$ from \mathbf{g}' .

If there exists $p \in [x_1, c] \setminus \mathcal{N}_{2\vartheta}(\{x_1, c\})$ and $p' \in [c, c']$ such that $\text{dist}(p, p') < \vartheta$, then by the triangle inequality $\text{dist}(c, p') \geq \text{dist}(c, p) - \text{dist}(p, p') > \vartheta$. Hence $\text{dist}(p, c') \leq \text{dist}(p, p') + \text{dist}(p', c) < \text{dist}(c, p') + \text{dist}(p', c)$, which contradicts the choice of c, c' . A similar argument shows that the existence of points $p \in [x_1, c] \setminus \mathcal{N}_{2\vartheta}(\{x_1, c\})$ and $p' \in [x_1, x_2]$ such that $\text{dist}(p, p') < \vartheta$ would contradict the choice of x_1 .

The distance between $[x_1, x_2]$ and $[c, c']$ is at least $(\frac{1}{2} - 2\epsilon)\ell - 2\nu\vartheta$, and the one between $[x_1, x_2]$ and $[x', c']$ is at least $D - tM_0 - 2\nu\vartheta$. Thus, it suffices to verify that the distance between $[x_1, x_2] \setminus \mathcal{N}_{2\vartheta}(\{x_1, x_2\})$ and $[c, x_1] \cup [x_2, x']$ is at least ϑ . According to the choices of x_1, x_2 this distance is 2ϑ .

The edge $[x_2, x']$ is at distance at least $(\frac{1}{2} - 2\epsilon)\ell$ from $[c, c']$ and at distance $2\nu\vartheta$ from $[x_1, c]$. The choice of x' implies that $[x_2, x'] \setminus \mathcal{N}_{2\vartheta}(\{x_2, x'\})$ is at distance 2ϑ from \mathbf{g}' . If a point $p \in [x_2, x'] \setminus \mathcal{N}_{2\vartheta}(\{x_2, x'\})$ and a point $p' \in [x_1, x_2]$ are such that $\text{dist}(p, p') < \vartheta$ then by the triangle inequality $\text{dist}(x_2, p') > \vartheta$. It follows that $\text{dist}(p, x_1) \leq \text{dist}(p, p') + \text{dist}(p', x_1) < \text{dist}(x_2, p') + \text{dist}(p', x_1)$, which contradicts the choice of x_2 .

The edge $[x', c']$ is at distance at least $D - tM_0 - 2\nu\vartheta$ from $\mathbf{g} \cup [x_1, x_2]$. If a point $p \in [x', c'] \setminus \mathcal{N}_{2\vartheta}(\{x', c'\})$ and a point $p' \in [x, x']$ satisfy $\text{dist}(p, p')$ then as above we obtain the contradictory inequality $\text{dist}(x, p) < \text{dist}(x, x')$. Likewise, if p' is in $[c, c']$ we obtain $\text{dist}(c, p) < \text{dist}(c, c')$.

(F₂) The vertex c' is at distance at least $D - tM_0 - 2\nu\vartheta$ from $[c, x_1] \cup [x_1, x_2]$ and at distance at least $(\frac{1}{2} - 2\epsilon)\ell$ from $[x_2, x']$.

The vertex c is at distance at least $D - tM_0$ from $[x', c']$ and at distance at least $(\frac{1}{2} - 2\epsilon)\ell - 2\nu\vartheta$ from $[x_1, x_2] \cup [x_2, x']$.

We have chosen x_1 at distance $2\nu\vartheta$ from $[x_2, x']$. The same vertex is at distance at least $D - tM_0$ from $[x', c']$ and at least $(\frac{1}{2} - 2\epsilon)\ell - 2\nu\vartheta$ from $[c, c']$.

Similarly, x_2 is at distance $2\nu\vartheta$ from $[x_1, c]$, at distance at least $(\frac{1}{2} - 2\epsilon)\ell - 2\nu\vartheta$ from $[c, c']$ and at least $D - tM_0 - 2\nu\vartheta$ from $[x', c']$.

The vertex c' is at distance at least $D - tM_0 - 2\nu\vartheta$ from $[x, c]$ and at least $(\frac{1}{2} - 2\epsilon)\ell - 2\nu\vartheta$ from $[x_1, x_2] \cup [x_2, x']$.

The pentagon of vertices x_1, x_2, x', c', c is $(\vartheta, 2\nu)$ -fat. Lemma 4.24 and the fact that $\text{dist}(c, c') = \text{dist}(c, [c', x'])$ implies that by adding a vertex on $[c, c']$ this pentagon becomes a hexagon (ϑ, ν) -fat. Therefore by (β_3) it is contained in $\mathcal{N}_\chi(A')$ for some $A' \in \mathcal{A}$. In particular the edge $[x', c']$ is contained in $\mathcal{N}_\chi(A') \cap \mathcal{N}_{tM_0}(A)$. This edge has length at least $\frac{\ell}{4} > 2D$ and D is at least diam_δ for $\delta = \max(\chi, tM_0)$. It follows that $A = A'$ and that $D < \chi$, which is a contradiction.

We conclude that property (α_2) is satisfied for $\varepsilon < \frac{1}{8t}$ and for D chosen above.

Property (β_2) is obtained as follows. If a geodesic $\mathbf{g} : [0, \ell] \rightarrow X$ joins two points in $\mathcal{N}_{\delta\ell}(A)$ then it is contained in $\mathcal{N}_{t\delta\ell}(A)$ by (\mathfrak{Q}) . If $\delta < \frac{\varepsilon}{3t}$ then by (α_2) the sub-geodesics $g([0, \frac{\ell}{3}])$ and $g([\frac{2\ell}{3}, \ell])$ intersect $\mathcal{N}_D(A)$. Then by (\mathfrak{Q}) , $g([\frac{\ell}{3}, \frac{2\ell}{3}])$ is contained in $\mathcal{N}_{tD}(A)$. \square

In view of Lemmata 4.22, 4.23 and 4.25, in order to finish the proof of Theorem 4.21 it suffices to prove the following.

Lemma 4.27. *Assume that (X, dist) and \mathcal{A} satisfy the properties (α_1) , (α_2) , (\mathfrak{Q}) and (β_3) . Then X is ATG with respect to \mathcal{A} .*

Proof. In an asymptotic cone $\text{Con}_\omega(X; e, d)$, the sets in \mathcal{A}_ω are closed, by Proposition 2.6, (2).

Lemma 4.28. *If (X, \mathcal{A}) satisfy property (\mathfrak{Q}) then any limit geodesic with endpoints in a set $A \in \mathcal{A}_\omega$ is contained in A . In particular if X is geodesic and (\mathfrak{Q}) is satisfied then all subsets in \mathcal{A}_ω are geodesic.*

Proof. Let $[x, y] = \lim_\omega([x_n, y_n])$ be a limit geodesic with its endpoints in a set $A = \lim_\omega(A_n)$, $A_n \in \mathcal{A}$. Then $\delta_n = \max[\text{dist}(x_n, A_n), \text{dist}(y_n, A_n)]$ satisfies $\delta_n =_\omega o(d_n)$. Property (\mathfrak{Q}) implies that $[x_n, y_n] \subset \mathcal{N}_{t\delta_n}(A_n)$. It follows that $[x, y] \subset A$. \square

Property (T_1) for \mathcal{A}_ω is deduced from (α_1) and (α_2) as in [DS05b, Lemma 4.5]. Properties $(T_1) \& (\beta_3)$ imply (T_2) by Corollary 4.19. \square

Proposition 4.29. *For any $\eta > 0$ property (β_3) can be replaced by the following:*

(β_3^η) *there exists $\vartheta > 0$, $\nu \geq 8$ and $\chi > 0$ such that any geodesic hexagon (ϑ, ν) -fat with diameter at least η is contained in $\mathcal{N}_\chi(A)$, for some $A \in \mathcal{A}$.*

Proof. Indeed, as a sufficient condition (β_3^η) is used to prove (Π_3) in any asymptotic cone, by means of Proposition 4.13. Given the sequence of hexagons H_n in Proposition 4.13, H_n has ω -almost surely diameter of order $O(d_n)$. Property (β_3^η) suffices therefore to obtain property (Π_3) .

Property (β_3) is also used in Lemma 4.25 to prove (α_2) . It suffices to take in that proof the constant D larger than $tM_0 + \eta$ to obtain that the geodesic pentagon with vertices x_1, x_2, x', c', c has diameter at least $\text{dist}(c, c') \geq D - tM_0 > \eta$. The same pentagon is $(\vartheta, 2\nu)$ -fat, hence by Lemma 4.24 it can be made into a hexagon (ϑ, ν) -fat of diameter larger than η ; therefore it is contained in $\mathcal{N}_\chi(A')$ for some $A' \in \mathcal{A}$. The rest of the argument is carried out similarly. \square




Figure 4: Case (P).

Proposition 4.30. *Let \mathcal{A}_{red}^η be the set of $A \in \mathcal{A}$ such that $\mathcal{N}_\chi(A)$ contains a (ϑ, ν) -fat geodesic hexagon with diameter at least η . Then the space X is ATG with respect to \mathcal{A}_{red}^η .*

Proof. Since $\mathcal{A}_{red}^\eta \subset \mathcal{A}$, properties (α_1) and (β_2) are satisfied. Property (β_3^η) is also satisfied by \mathcal{A}_{red}^η , hence by Proposition 4.29, X is ATG with respect to \mathcal{A}_{red}^η . \square

Corollary 4.31. *For every $\lambda > 0$ the space X is also ATG with respect to the subset \mathcal{A}_λ in \mathcal{A} composed of all the subsets of diameter at least λ in \mathcal{A} .*

Proof. Indeed $\mathcal{A}_\lambda \subset \mathcal{A}$ implies that properties (α_1) and (β_2) are still satisfied.

Let $\eta = \lambda + 2\chi$. Then $\mathcal{A}_{red}^\eta \subset \mathcal{A}_\lambda$, which implies that property (β_3^η) is satisfied by \mathcal{A}_λ . By Proposition 4.29, X is ATG with respect to \mathcal{A}_λ . \square

4.4 New definition, closer to the definition of hyperbolicity

In [DS05a] a version for groups of the following notion has been introduced.

Definition 4.32. Let X be a geodesic metric space and let \mathcal{A} be a collection of subsets of X . We say that X is $(*)$ -asymptotically tree-graded with respect to \mathcal{A} if for every $C \geq 0$ there exist two constants σ and δ such that every triangle xyz with $(1, C)$ -almost geodesic edges in X is in one of the following two cases:

- (C) there exists $a \in X$ such that $\overline{B}(a, \sigma)$ intersects each of the sides of the triangle;
- (P) there exists $A \in \mathcal{A}$ such that $\overline{\mathcal{N}}_\sigma(A)$ intersects each of the sides of the triangle, and the entrance (resp. exit) points x_1, y_1, z_1 (resp. y_2, z_2, x_2) of the sides $[x, y], [y, z], [z, x]$ in $\overline{\mathcal{N}}_\sigma(A)$ satisfy

$$\text{dist}(x_1, x_2) < \delta, \text{dist}(y_1, y_2) < \delta, \text{dist}(z_1, z_2) < \delta.$$

See Figure 4, taken from [DS05b].

Remark 4.33. If X is a geodesic metric space in which for some constant $\sigma > 0$ every geodesic triangle satisfies condition (C), then X is a hyperbolic space. Conversely, in a hyperbolic geodesic metric space for every $L \geq 1$ and $C \geq 0$ there exists $\sigma > 0$ such that every triangle with (L, C) -quasi-geodesic edges satisfies condition (C).

Remarks 4.34. (1) If a metric space X is ATG with respect to a collection of subsets \mathcal{A} then X is $(*)$ -ATG with respect to \mathcal{A} , by [DS05b, Corollary 8.14 and Lemma 8.19].

Moreover, according to [DS05b, Corollary 8.14] if a geodesic triangle is in case **(P)** then for every $\sigma' \geq \sigma$ there exists δ' such that the pairs of entrance points in $\overline{\mathcal{N}}_{\sigma'}(A)$ are at distance at most δ' .

- (2) The notion of $(*)$ -ATG space is weaker than the one of ATG space. For instance if X is a geodesic hyperbolic space and if \mathcal{A} is any collection of subsets covering X , then X is $(*)$ -ATG with respect to \mathcal{A} , and the collection \mathcal{A} needs not satisfy properties (α_1) or (\mathfrak{Q}) , for instance.

It turns out nevertheless that one can formulate an equivalent definition of ATG metric spaces using the $(*)$ -property.

Theorem 4.35. *Let (X, dist) be a geodesic metric space and let \mathcal{A} be a collection of subsets of X . The metric space X is ATG with respect to \mathcal{A} if and only if (X, \mathcal{A}) satisfy properties (α_1) and (α_2) , and moreover X is $(*)$ -ATG with respect to \mathcal{A} .*

Convention 4.36. In order to simplify some technical arguments of the equivalence we make the assumption that for all $C > 0$ the constant σ in the $(*)$ -property is larger than the constant M appearing in property (α_2) . By Remark 4.34, (1), if X is ATG then such a choice can be made.

Proof. The direct implication has already been discussed. Concerning the converse statement, as in Section 4.3 from (α_1) and (α_2) can be deduced property (\mathfrak{Q}) . By Proposition 2.6, (2), and Lemma 4.28, in any asymptotic cone $\text{Con}_\omega(X; e, d)$ the collection \mathcal{A}_ω is composed of closed geodesic subsets. Again (α_1) and (α_2) imply property (T_1) for \mathcal{A}_ω . According to Corollary 4.18, it remains to prove property (Π_3) .

Lemma 4.37. *Let (X, dist) be a geodesic metric space and let \mathcal{A} be a collection of subsets of X satisfying property (α_2) for some $\varepsilon \in [0, 1/2)$ and $M > 0$.*

Let $\mu \geq \nu \geq M$, and let \mathfrak{g} be a geodesic and A a subset in \mathcal{A} such that \mathfrak{g} intersects $\overline{\mathcal{N}}_\nu(A)$. If e_μ and e_ν are the entrance points of \mathfrak{g} in $\overline{\mathcal{N}}_\mu(A)$ and respectively $\overline{\mathcal{N}}_\nu(A)$ then $\text{dist}(e_\mu, e_\nu) \leq \frac{\mu}{\varepsilon}$.

Proof. If $\varepsilon \text{dist}(e_\mu, e_\nu) > \mu$ then by (α_2) the sub-arc of \mathfrak{g} between e_μ and e_ν intersects $\mathcal{N}_M(A) \subset \mathcal{N}_\nu(A)$, which contradicts the definition of e_ν . \square

Lemma 4.38. *If (X, dist) is a geodesic metric space $(*)$ -ATG with respect to a collection \mathcal{A} of subsets of X then for every $C \geq 0$ there exist $\kappa \geq 0$ and $\lambda \geq 0$ such that the following holds. For any two geodesics $\mathfrak{g}, \mathfrak{g}'$ with $\mathfrak{g}_- = \mathfrak{g}'_-$ and $\text{dist}(\mathfrak{g}_+, \mathfrak{g}'_+) \leq C$, any point z on \mathfrak{g}' is either contained in $\overline{\mathcal{N}}_\kappa(\mathfrak{g})$ or it is contained in $\overline{\mathcal{N}}_\kappa(A)$ for some $A \in \mathcal{A}$ such that $\overline{\mathcal{N}}_\kappa(A)$ intersects \mathfrak{g} . Moreover, if e and f are the entrance and respectively the exit point of \mathfrak{g}' from $\overline{\mathcal{N}}_\kappa(A)$, then $\{e, f\} \subset \overline{\mathcal{N}}_\lambda(\mathfrak{g})$.*

Proof. Let \mathfrak{p} be the path $\mathfrak{g} \sqcup [\mathfrak{g}_+, \mathfrak{g}'_+]$, where $[\mathfrak{g}_+, \mathfrak{g}'_+]$ is a geodesic segment joining \mathfrak{g}_+ and \mathfrak{g}'_+ . It is a $(1, 2C)$ -almost geodesic. Let σ and δ be the constants of property $(*)$ for $2C$, and let z be an arbitrary point on \mathfrak{g}' , dividing \mathfrak{g}' into two sub-arcs, \mathfrak{g}_1 and \mathfrak{g}_2 . The triangle Δ of edges $\mathfrak{g}_1, \mathfrak{g}_2$ and \mathfrak{p} is either in case **(C)** or in case **(P)**.

If it is in case **(C)** then there exist $a_1 \in \mathfrak{g}_1, a_2 \in \mathfrak{g}_2$ and $b \in \mathfrak{p}$ such that the set $\{a_1, a_2, b\}$ has diameter at most 2σ . The point z is on a geodesic joining a_1 and a_2 , hence it is at distance at most 3σ from b , thus it is contained in $\overline{\mathcal{N}}_{3\sigma+C}(\mathfrak{g})$.

If Δ is in case **(P)** then there exists $A \in \mathcal{A}$ with $\overline{\mathcal{N}}_\sigma(A)$ intersecting $\mathbf{g}_1, \mathbf{g}_2$ and \mathbf{p} . Let x_1, z_1, z_2, y_1 and x_2, y_2 be the entrance and exit points from $\overline{\mathcal{N}}_\sigma(A)$ of $\mathbf{g}_1, \mathbf{g}_2$ and \mathbf{p} respectively. Then $\text{dist}(x_1, x_2), \text{dist}(y_1, y_2)$ and $\text{dist}(z_1, z_2)$ are all at most δ . Since z is on a geodesic joining z_1 and z_2 , $z \in \overline{\mathcal{N}}_{\sigma+\delta/2}(A)$. Note that $\overline{\mathcal{N}}_\sigma(A)$ intersects \mathbf{p} , therefore $\overline{\mathcal{N}}_{\sigma+C}(A)$ intersects \mathbf{g} .

Take $\kappa = \max(3\sigma + C, \sigma + \frac{\delta}{2}, \sigma + C)$.

The points x_1 and y_1 are the entrance and respectively the exit point of \mathbf{g}' from $\overline{\mathcal{N}}_\sigma(A)$. If we consider e and f the entrance and exit points of \mathbf{g}' from $\overline{\mathcal{N}}_\kappa(A)$, Lemma 4.37 implies that $\text{dist}(x_1, e)$ and $\text{dist}(y_1, f)$ are at most $\frac{\kappa}{\varepsilon}$. Therefore $\text{dist}(e, \mathbf{g})$ and $\text{dist}(f, \mathbf{g})$ are both at most $\frac{\kappa}{\varepsilon} + \delta + C$. \square

Lemma 4.39. *Let (X, dist) be a geodesic metric space and let \mathcal{A} be a collection of subsets of X . Assume that X is $(*)$ -ATG with respect to \mathcal{A} and that (X, \mathcal{A}) satisfy properties (α_1) and (\mathfrak{Q}) . Then there exist $\vartheta > 0$, $\nu \geq 8$ and $\chi > 0$ such that any geodesic quadrilateral which is (ϑ, ν) -fat is contained in $\mathcal{N}_\chi(A)$ for some $A \in \mathcal{A}$.*

Proof. Let P be a (ϑ, ν) -fat quadrilateral with vertices in counterclockwise order x, y, z, w . Let $[x, z]$ be a geodesic joining the opposite vertices x and z .

CASE 1. Assume that both geodesic triangles xyz and xzw are in case **(C)**. Then there exists $a_1 \in [x, y]$, $a_2 \in [y, z]$ and $a_3 \in [x, z]$ such that the set $\{a_1, a_2, a_3\}$ has diameter at most 2σ . Likewise there exists $b_1 \in [z, w]$, $b_2 \in [w, x]$ and $b_3 \in [x, z]$ such that $\{b_1, b_2, b_3\}$ has diameter at most 2σ . If $\vartheta > 2\sigma$ then $a_1, a_2 \in B(y, 2\vartheta)$ and $b_1, b_2 \in B(w, 2\vartheta)$.




Figure 5: Case 1 in proof of Lemma 4.39.

Without loss of generality we may assume that $a_3 \in [x, b_3]$. Lemma 4.38 applied to $[x, b_2]$ and to $a_3 \in [x, b_3]$ implies that for some $\kappa = \kappa(2\sigma)$ either $a_3 \in \overline{\mathcal{N}}_\kappa([x, b_2])$, or $a_3 \in \overline{\mathcal{N}}_\kappa(A)$ such that the entrance respectively exit point, a_4, a_5 , of $[x, b_3]$ from $\overline{\mathcal{N}}_\kappa(A)$ are at distance at most λ from $[x, b_2]$, for some $\lambda = \lambda(2\sigma)$. If $\vartheta > 2\sigma + \kappa$ then by Lemma 4.6 the first case cannot occur.

In the second case we have that $\text{dist}(a_3, \{a_4, a_5\})$ is at least $\vartheta - 2\sigma - \lambda$. Lemma 4.38 applied to $[z, a_2]$ and to $a_5 \in [z, a_3]$ implies that either $a_5 \in \overline{\mathcal{N}}_\kappa([z, a_2])$ or that $a_5 \in \overline{\mathcal{N}}_\kappa(A')$ such that the entrance and the exit point a_6, a_7 , of $[a_3, z]$ from $\overline{\mathcal{N}}_\kappa(A)$ are at distance at most λ from $[z, a_2]$. The first case cannot occur if $\vartheta > \lambda + \kappa$. In the second case $\text{dist}(a_5, \{a_6, a_7\}) \geq \vartheta - 2\lambda$. The intersection $[a_3, a_5] \cap [a_6, a_7]$ has length at least $\vartheta - 2\sigma - 2\lambda$. By property (\mathfrak{Q}) this intersection is contained in $\overline{\mathcal{N}}_{t\kappa}(A) \cap \overline{\mathcal{N}}_{t\kappa}(A')$. If $\vartheta > 2\sigma + 2\lambda + \text{diam}_{t\kappa+1} + 1$ then $A = A'$.

The point a_4 is the entrance point of $[x, b_3]$ in $\overline{\mathcal{N}}_\kappa(A)$ while a_6 is the entrance point of $[a_3, z]$ in $\overline{\mathcal{N}}_\kappa(A)$. If $a_4 \in [a_3, b_3]$ then $a_4 = a_6$, and this point is at distance at most λ from both $[x, w]$ and $[y, z]$. If $\vartheta > 2\lambda$ then this cannot occur. Thus we may assume that $a_4 \in [x, a_3]$. Likewise we have that $a_7 \in [b_3, z]$ (see Figure 5).

We apply Lemma 4.38 to $[x, a_1]$ and to $a_4 \in [x, a_3]$. If we are in the second case of the conclusion then $a_4 \in \overline{\mathcal{N}}_\kappa(A'')$, and the entrance and exit point, a'_4, a''_4 , of $[x, a_3]$ from $\overline{\mathcal{N}}_\kappa(A'')$ are at distance at most λ from $[x, a_1]$. If $\text{dist}(a_4, a''_4) \geq \text{diam}_{t\kappa+1} + 1$ then $A'' = A$ and $a_4 = a'_4$. Thus, in all cases, a_4 is at distance at most $\kappa + \text{diam}_{t\kappa+1} + 1 + \lambda$ from $[x, a_1]$. Recall that a_4 is at distance at most λ from $[x, w]$. It follows that if $\vartheta > \kappa + \text{diam}_{t\kappa+1} + 1 + 2\lambda$ then $a_4 \in B(x, 2\vartheta + \lambda)$.

A similar argument gives that $a_7 \in B(z, 2\vartheta + \lambda)$.

We have thus that $\{x, z\} \subset \overline{\mathcal{N}}_{2\vartheta+\lambda+\kappa}(A)$. Also, since $\{a_3, b_3\} \subset [a_4, a_7] \subset \overline{\mathcal{N}}_{t\kappa}(A)$ it follows that $\{y, w\} \subset \overline{\mathcal{N}}_{2\vartheta+2\sigma+t\kappa}(A)$. By property (Ω) , $P \subset \overline{\mathcal{N}}_\chi(A)$ where $\chi = t(2\vartheta + \lambda + 2\sigma + t\kappa)$.

CASE 2. Assume that the triangle xyz is in case **(P)** while xzw is in case **(C)**. Then there exists $A \in \mathcal{A}$ such that $\overline{\mathcal{N}}_\sigma(A)$ intersects all the edges of xyz . Moreover if x_2, y_1 are the entrance and exit point of $[x, y]$ in $\overline{\mathcal{N}}_\sigma(A)$, while y_2, z_1 and z_2, x_1 are the entrance and exit points of $[y, z]$ and respectively $[z, x]$ in $\overline{\mathcal{N}}_\sigma(A)$ then $\text{dist}(x_1, x_2), \text{dist}(y_1, y_2)$ and $\text{dist}(z_1, z_2)$ are all at most δ .

Let also $b_1 \in [z, w]$, $b_2 \in [w, x]$ and $b_3 \in [x, z]$ be such that $\{b_1, b_2, b_3\}$ has diameter at most 2σ . If $\vartheta > 2\sigma$ then property (F_1) implies that $\{b_1, b_2\} \subset B(w, 2\vartheta)$.

CASE 2.A. Assume that $b_3 \in [x_1, z_2]$. Note that $\text{dist}(b_3, x_1) \geq \text{dist}(w, [x, y]) - 2\vartheta - 2\sigma - \delta \geq 6\vartheta - 2\sigma - \delta$. Same for $\text{dist}(b_3, z_2)$. Thus for ϑ large both $\text{dist}(b_3, x_1)$ and $\text{dist}(b_3, z_2)$ are large.

Lemma 4.38 applied to $x_1 \in [x, b_3]$ and to $[x, b_2]$ implies that either $x_1 \in \overline{\mathcal{N}}_\kappa([x, b_2])$ or $x_1 \in \overline{\mathcal{N}}_\kappa(A_1)$ such that the entrance and exit point x'_1, x''_1 of $[x, b_3]$ from $\overline{\mathcal{N}}_\kappa(A_1)$ are at distance at most λ from $[x, b_2]$. In the latter case if $\text{dist}(x_1, x''_1) > \text{diam}_\tau$ where $\tau = t \max(\sigma, \kappa) + 1$ then $A_1 = A$ and $x_1 = x'_1$. Thus in all cases $\text{dist}(x_1, [x, w]) = O(1)$. For ϑ large enough it follows that $x_2 \in B(x, 2\vartheta)$, hence $x \in \overline{\mathcal{N}}_{2\vartheta+\sigma}(A)$. A similar argument gives that $z \in \overline{\mathcal{N}}_{2\vartheta+\sigma}(A)$.

If $\vartheta > \delta$ then $\text{dist}(y, y_1) < 2\vartheta$ and $y \in \overline{\mathcal{N}}_{2\vartheta+\sigma}(A)$.

Also $b_3 \in [x_1, z_2] \subset \overline{\mathcal{N}}_{t\sigma}(A)$, hence $w \in \overline{\mathcal{N}}_{t\sigma+2\vartheta+2\sigma}(A)$. We conclude by property (Ω) that $P \subset \overline{\mathcal{N}}_\chi(A)$ for some $\chi = O(1)$.

CASE 2.B. Assume that $b_3 \notin [x_1, z_2]$. Without loss of generality we may assume that $[x_1, z_2] \subset [x, b_3]$.

Lemma 4.38 applied to $b_3 \in [z_2, z]$ and to $[z_1, z]$ implies that either $b_3 \in \overline{\mathcal{N}}_\kappa([z_1, z])$ or that $b_3 \in \overline{\mathcal{N}}_\kappa(B)$ such that if b_4, b_5 are the entrance and exit point of $[z_2, z]$ from $\overline{\mathcal{N}}_\kappa(B)$ then these points are at distance at most λ from $[z_1, z]$. For ϑ large enough the first case cannot occur. In the second case $\text{dist}(b_3, \{b_4, b_5\}) \geq \text{dist}(w, [y, z]) - \lambda - 2(\vartheta + \sigma) \geq 6\vartheta - \lambda - 2\sigma$.

Applying Lemma 4.38 now to $b_4 \in [x, b_3]$ and to $[x, b_2]$ gives that for ϑ large enough $b_4 \in \overline{\mathcal{N}}_\kappa(B')$ such that the entrance and exit points b_6, b_7 of $[x, b_3]$ from $\overline{\mathcal{N}}_\kappa(B')$ are at distance at most λ from $[x, b_2]$ (see Figure 6). Moreover $\text{dist}(b_4, \{b_6, b_7\}) \geq \vartheta - 2\lambda$. Thus $\overline{\mathcal{N}}_{t\kappa}(B) \cap \overline{\mathcal{N}}_{t\kappa}(B')$ contains $[b_4, b_7] \cap [b_4, b_3]$ of length $\min(6\vartheta - \lambda - 2\sigma, \vartheta - 2\lambda)$. For ϑ large we conclude that $B = B'$.

The point b_6 is the entrance point of $[x, b_3]$ into $\overline{\mathcal{N}}_\kappa(B)$ while b_4 is the entrance point of $[z_2, z]$ into $\overline{\mathcal{N}}_\kappa(B)$. If $b_6 \in [z_2, b_3]$ then $b_6 = b_4$ and $\text{dist}([y, z], [x, w]) \leq 2\lambda$. For ϑ large enough this case is impossible. Thus $b_6 \in [x, z_2]$.

The intersection $[x_1, z_2] \cap [b_6, z_2]$ is in $\overline{\mathcal{N}}_{t\sigma}(A) \cap \overline{\mathcal{N}}_{t\kappa}(B)$. Note that $\text{dist}(b_6, z_2) \geq \text{dist}([y, z], [x, w]) - \lambda - \delta$ thus for ϑ large we may assume that $\text{dist}(b_6, z_2) > \text{diam}_\tau + 1$ with $\tau = t \max(\sigma, \kappa) + 1$.

If $\text{dist}(x_1, z_2) \leq \text{diam}_\tau + 1$ then $\{x_2, x_1, z_1\}$ has diameter $O(1)$ and we are back in Case 1 with $a_1 = x_2, a_2 = z_1$ and $a_3 = x_1$ and with the constant σ possibly larger. We may then use the proof in Case 1 to finish the argument.




Figure 6: Case 2.B in proof of Lemma 4.39.

Assume now that $\text{dist}(x_1, z_2) > \text{diam}_\tau + 1$. Then $A = B$ and b_5 , the entrance point of $[z, z_2]$ into $\overline{\mathcal{N}}_\kappa(B)$, is also the entrance point of $[z, x]$ into $\overline{\mathcal{N}}_\kappa(A)$. As z_2 is the entrance point of $[z, x]$ into $\overline{\mathcal{N}}_\sigma(A)$, Lemma 4.37 implies that $\text{dist}(z_2, b_5) = O(1)$.

By construction $b_4 \in [z_2, b_5]$, hence $b_7 \in [b_4, b_3] \subset [z_2, b_5]$. Consequently $\text{dist}(z_2, b_7) = O(1)$. On the other hand $\text{dist}(z_2, [y, z]) < \delta$ and $\text{dist}(b_7, [x, w]) \leq \lambda$. Thus for ϑ large enough we obtain a contradiction.

CASE 3. Assume that both geodesic triangles xyz and xzw are in case **(P)**. Then there exists A_1 in \mathcal{A} such that $\overline{\mathcal{N}}_\sigma(A_1)$ intersects all the edges of xyz . Moreover the pairs of entrance points in it, (x_1, x_2) , (y_1, y_2) and (z_1, z_2) are all at respective distances less than δ . Likewise there exists A_2 in \mathcal{A} such that $\overline{\mathcal{N}}_\sigma(A_2)$ intersects all the edges of xzw , and the pairs of entrance points in $\overline{\mathcal{N}}_\sigma(A_2)$, (x'_1, x'_2) , (z'_1, z'_2) and (w_1, w_2) are all at distances less than δ .

If $\vartheta > \delta$ then $y_1, y_2 \in B(y, 2\vartheta)$ and $w_1, w_2 \in B(w, 2\vartheta)$.

If $A_1 = A_2 = A$ then $x_1 = x'_1$, $z_2 = z'_1$, hence $\text{dist}(x'_1, x_2)$ and $\text{dist}(z_1, z'_2)$ are less than 2δ . If $\vartheta > 2\delta$ then it follows that $x'_1, x_2 \in B(x, 2\vartheta)$ and that $z_1, z'_2 \in B(z, 2\vartheta)$. Thus $x, y, z, w \in \overline{\mathcal{N}}_{2\vartheta+\sigma}(A)$, which by (\mathfrak{Q}) implies that $P \subset \mathcal{N}_\chi(A)$ for $\chi > t(2\vartheta + \sigma)$.

Assume that $A_1 \neq A_2$. Then $[x_1, z_2] \subset [x, z]$ and $[x'_2, z'_1] \subset [x, z]$ intersect in a sub-geodesic of length at most $\text{diam}_{\sigma+1}$. If the intersection is non-empty then x'_1 and z_1 are at distance at most $2\delta + \text{diam}_{\sigma+1}$. If $\vartheta > 2\delta + \text{diam}_{\sigma+1}$ then this is impossible. It follows that the intersection is empty.

Without loss of generality we may assume that $[x_1, z_2] \subset [x, x'_2]$. Lemma 4.38 applied to the geodesic $[x, x'_1]$ and the point $z_2 \in [x, x'_2]$ implies that for some $\kappa = \kappa(\delta)$ either $z_2 \in \overline{\mathcal{N}}_\kappa([x, w])$ or $z_2 \in \overline{\mathcal{N}}_\kappa(A_3)$, and the entrance and exit points z_3, z_4 of $[x, x'_2]$ in $\overline{\mathcal{N}}_\kappa(A_3)$ are at distance at most λ from $[x, x'_1]$ (see Figure 7). If $\vartheta > \delta + \kappa$ the first case cannot occur. In the second case we have that $\text{dist}(z_2, \{z_3, z_4\}) > \vartheta - \delta - \lambda$. In particular we may assume that $\text{dist}(z_2, z_3) > \text{diam}_\tau + 1$ for $\tau = t \max(\sigma, \kappa) + 1$.

If $\text{dist}(x_1, z_2) \leq \text{diam}_\tau + 1$ then the set $\{x_2, x_1, z_1\}$ has diameter at most $2\delta + \text{diam}_\tau + 1$ and it contains one point on each edge of the triangle xyz . Therefore we find ourselves in Case 2.B, with the constant σ possibly larger. We can then finish the argument as in that case.

Assume that $\text{dist}(x_1, z_2) > \text{diam}_\tau + 1$. Then $[x_1, z_2] \cap [z_3, z_2]$ has diameter $> \text{diam}_\tau$ and




Figure 7: Case 3 in proof of Lemma 4.39.

it is contained in $\overline{\mathcal{N}}_{t\sigma}(A_1) \cap \overline{\mathcal{N}}_{t\kappa}(A_3)$. Therefore $A_1 = A_3$. In particular z_4 , the exit point of $[x, x'_2]$ from $\overline{\mathcal{N}}_{\kappa}(A_1)$, and z_2 , the exit point of $[x, z]$ (therefore also of $[x, x'_2]$) from $\overline{\mathcal{N}}_{\sigma}(A_1)$ are at distance $O(1)$ by Lemma 4.37. It follows that z_1 and $[x, w]$ are at distance $O(1)$, and if ϑ is large enough then this gives a contradiction. \square

Corollary 4.40. *Let (X, dist) and \mathcal{A} satisfy the hypotheses of Lemma 4.39. Then in any asymptotic cone of X property (Π_2) is satisfied.*

Proof. In an asymptotic cone $\text{Con}_{\omega}(X; e, d)$ consider a simple bigon of endpoints x, y whose edges are limit geodesics. Then there exist two sequences of geodesics $[x_n, y_n]$ and $[x'_n, y'_n]$ such that ω -almost surely $\text{dist}(x_n, x'_n)$ and $\text{dist}(y_n, y'_n)$ are of order $o(d_n)$, while $\text{dist}(x_n, y_n)$ and $\text{dist}(x'_n, y'_n)$ are of order $O(d_n)$. Let m_n and m'_n be the midpoints of $[x_n, y_n]$ and respectively of $[x'_n, y'_n]$. Let δ_n be the maximum between $\text{dist}([x_n, m_n], [x'_n, m'_n])$ and $\nu\vartheta$, where ϑ and ν are the constants provided by Lemma 4.39. Then $\delta_n =_{\omega} o(d_n)$. Similarly, $\delta'_n = \max\{\text{dist}([m_n, y_n], [m'_n, y'_n]), \nu\vartheta\}$ satisfies $\delta'_n =_{\omega} o(d_n)$. Let x_n^1 be the farthest from x_n point on $[x_n, m_n]$ at distance δ_n from $[x'_n, m'_n]$, and let x_n^2 be the farthest from x'_n point on $[x'_n, m'_n]$ at distance δ_n from x_n^1 . We choose in a similar manner $y_n^1 \in [y_n, m_n]$ and $y_n^2 \in [y'_n, m'_n]$. Since the limit bigon is simple, it follows that the sets $\{x_n, x'_n, x_n^1, x_n^2\}$ and $\{y_n, y'_n, y_n^1, y_n^2\}$ have diameters $o(d_n)$ ω -almost surely.

We prove that any quadrilateral having as two opposite edges $[x_n^1, y_n^1] \subset [x_n, y_n]$ and $[x_n^2, y_n^2] \subset [x'_n, y'_n]$ is (ϑ, ν) -fat. This suffices to finish the argument, by Lemma 4.39.

(F₁) By construction $\text{dist}([x_n^1, y_n^1], [x_n^2, y_n^2]) \geq \nu\vartheta$.

If the set $[x_n^1, y_n^1] \setminus \mathcal{N}_{2\vartheta}(\{x_n^1, y_n^1\})$ contains a point p at distance $< \vartheta$ from a point $q \in [x_n^1, x_n^2]$ then $\text{dist}(x_n^1, q) \geq \vartheta > \text{dist}(p, q)$, hence $\text{dist}(x_n^1, x_n^2) > \text{dist}(p, x_n^2)$, contradicting the choice of x_n^1 . Likewise it is shown that $[x_n^1, y_n^1] \setminus \mathcal{N}_{2\vartheta}(\{x_n^1, y_n^1\})$ is at distance at least ϑ from $[y_n^1, y_n^2]$; also that $[x_n^2, y_n^2] \setminus \mathcal{N}_{2\vartheta}(\{x_n^2, y_n^2\})$ is at distance at least ϑ from $[x_n^1, x_n^2] \cup [y_n^1, y_n^2]$.

By construction $[x_n^1, x_n^2] \setminus \mathcal{N}_{2\vartheta}(\{x_n^1, x_n^2\})$ is at distance 2ϑ from $[x_n^1, y_n^1] \cup [x_n^2, y_n^2]$. Same for $[y_n^1, y_n^2] \setminus \mathcal{N}_{2\vartheta}(\{y_n^1, y_n^2\})$. The edges $[x_n^1, x_n^2]$ and $[y_n^1, y_n^2]$ are at distance $O(d_n)$ from each other.

Property **(F₂)** follows immediately from the fact that $\text{dist}([x_n^1, y_n^1], [x_n^2, y_n^2]) \geq \nu\vartheta$ and that $\text{dist}([x_n^1, x_n^2], [y_n^1, y_n^2]) = O(d_n)$. \square

The following statement ends the proof of Theorem 4.35.

Proposition 4.41. *Let (X, dist) be a geodesic metric space $(*)$ -ATG with respect to a collection \mathcal{A} of subsets of it. If (X, \mathcal{A}) moreover satisfy properties (α_1) and (α_2) , then in any asymptotic cone of X property (Π_3) is satisfied.*

Proof. Let $\text{Con}_\omega(X; e, d)$ be an arbitrary asymptotic cone of X and let \mathcal{A}_ω be the collection of limit sets of sequences from \mathcal{A} . Since $(\alpha_1) \& (\alpha_2) \Rightarrow (\mathfrak{Q})$ it follows that the sets in \mathcal{A}_ω are geodesic, by Lemma 4.28. Also $(\alpha_1) \& (\alpha_2)$ imply that \mathcal{A}_ω satisfies (T_1) .

Let Δ be a non-trivial simple geodesic triangle in $\text{Con}_\omega(X; e, d)$, whose edges $[x, y]$, $[y, z]$ and $[z, x]$ appear as limits of sequences $[x_n, y'_n]$, $[y_n, z'_n]$ and $[z_n, x'_n]$ of geodesics in X . Then ω -almost surely $\text{dist}(x_n, x'_n)$, $\text{dist}(y_n, y'_n)$ and $\text{dist}(z_n, z'_n)$ are of order $o(d_n)$, while the lengths of $[x_n, y'_n]$, $[y_n, z'_n]$ and $[z_n, x'_n]$ are of order $O(d_n)$. Let T_n be a geodesic triangle with vertices x_n, y_n, z_n . We denote its edges by $[u, v]$, with $u, v \in \{x_n, y_n, z_n\}$. The three limit geodesics $\mathfrak{g}^x = \lim_\omega([y_n, z_n])$, $\mathfrak{g}^y = \lim_\omega([x_n, z_n])$ and $\mathfrak{g}^z = \lim_\omega([x_n, y_n])$ compose the limit triangle $T = \lim_\omega(T_n)$.

CASE 1. Assume that ω -almost surely T_n is in case **(C)**. Then there exists $a_n^1 \in [x_n, y_n]$, $a_n^2 \in [y_n, z_n]$ and $a_n^3 \in [z_n, x_n]$ such that the set $\{a_n^1, a_n^2, a_n^3\}$ has ω -almost surely diameter at most 2σ for some constant σ . It follows that $\lim_\omega(a_n^1) = \lim_\omega(a_n^2) = \lim_\omega(a_n^3) = a$. The point a is on the three edges of T .

Without loss of generality we may assume that $a \notin \{x, y\}$. The fact that $a \neq x$ implies that either $\mathfrak{g}^z \neq [x, y]$ or $\mathfrak{g}^y \neq [x, z]$. Property (Π_2) implies that we may apply Proposition 3.14 to $\mathcal{L}_1 = [x, y]$, $\mathcal{L}_2 = [x, z]$, $\mathfrak{g}_1 = \mathfrak{g}^z$, $\mathfrak{g}_2 = \mathfrak{g}^y$ and to the intersection point $a \in \mathfrak{g}^z \cap \mathfrak{g}^y$. We conclude that the bigon formed by \mathfrak{g}^z and \mathfrak{g}^y of endpoints a, x is contained in a subset $A_x \in \mathcal{A}_\omega$.

Similarly we deduce that the bigon formed by \mathfrak{g}^z and \mathfrak{g}^x of endpoints a, y is contained in a subset $A_y \in \mathcal{A}_\omega$.

If $a = z$ then $\mathfrak{g}_x \subset A_y$, $\mathfrak{g}_y \subset A_x$. Also $a \notin [x, y]$, which by Lemma 3.12 implies that a is contained in the interior of a simple bigon formed by $[x, y]$ and \mathfrak{g}^z . By property (Π_2) this bigon is contained in some $A \in \mathcal{A}_\omega$. The intersections $A \cap A_x$ and $A \cap A_y$ contain non-trivial sub-arcs of \mathfrak{g}^z therefore by (T_1) we conclude that $A = A_x = A_y$. The subset A contains also \mathfrak{g}^z .

Property (Π_2) allows to apply Lemma 3.13 to the pairs of arcs $(\mathfrak{g}^x, [y, z])$, $(\mathfrak{g}^y, [x, z])$ and $(\mathfrak{g}^z, [x, y])$ and deduce that $\Delta = [x, y] \cup [y, z] \cup [z, x]$ is contained in A .

If $a \neq z$ then again by Proposition 3.14 the bigon formed by \mathfrak{g}^x and \mathfrak{g}^y of endpoints a, z is contained in a subset $A_z \in \mathcal{A}_\omega$. Since a is not a vertex in Δ it is contained in at most one edge of Δ . Without loss of generality we assume that $a \notin [x, y] \cup [y, z]$.

The fact that $a \notin [x, y]$, Lemma 3.12, properties (Π_2) and (T_1) imply as above that $A_x = A_y$. Likewise from $a \notin [y, z]$ we deduce that $A_y = A_z$. Thus $A = A_x = A_y = A_z$ contains T . Property (Π_2) and Lemma 3.13 imply that Δ is also contained in A .

CASE 2. Assume that ω -almost surely T_n is in case **(P)**. Then there exist A_n in \mathcal{A} such that $\overline{\mathcal{N}}_\sigma(A_n)$ intersects all the edges of T_n . Moreover if (x_n^2, y_n^1) , (y_n^2, z_n^1) and (z_n^2, x_n^1) are the pairs of entrance and exit points from $\overline{\mathcal{N}}_\sigma(A)$ of $[x_n, y_n]$, $[y_n, z_n]$ and $[z_n, x_n]$ respectively, then $\text{dist}(x_n^1, x_n^2)$, $\text{dist}(y_n^1, y_n^2)$ and $\text{dist}(z_n^1, z_n^2)$ are less than δ .

Let $x' = \lim_\omega(x_n^1) = \lim_\omega(x_n^2)$, $y' = \lim_\omega(y_n^1) = \lim_\omega(y_n^2)$ and $z' = \lim_\omega(z_n^1) = \lim_\omega(z_n^2)$.

Assume that $\{x', y', z'\}$ has cardinal at most 2. Assume for instance that $x' = y'$. Then the point $a = x' = y'$ is in $\mathfrak{g}^x \cap \mathfrak{g}^y \cap \mathfrak{g}^z$. With the same argument as in Case 1 we deduce that both T and Δ are contained in some $A \in \mathcal{A}_\omega$.

Assume now that $\{x', y', z'\}$ has cardinal 3. The geodesic triangle T' of vertices x', y', z' and with edges contained in the edges of T is included in the piece $A = \lim_{\omega} (A_n)$.

Proposition 3.14 implies that the bigon of endpoints x, x' formed by \mathfrak{g}^z and \mathfrak{g}^y is either trivial or contained in some $A_x \in \mathcal{A}_{\omega}$. Similarly, the bigon of endpoints y, y' formed by \mathfrak{g}^z and \mathfrak{g}^x is either trivial or in some A_y , and the bigon of endpoints z, z' formed by \mathfrak{g}^x and \mathfrak{g}^y is either trivial or in some A_z .

If $x' \neq x$ then x' cannot be contained both in $[x, y]$ and in $[x, z]$. Suppose that $x' \notin [x, y]$. Then x' is in the interior of a non-trivial simple bigon formed by \mathfrak{g}^z and $[x, y]$. This bigon is contained in some $B_x \in \mathcal{A}_{\omega}$ by (Π_2) , and its intersection with A_x contains a non-trivial sub-arc of \mathfrak{g}^z . The same holds for its intersection with A , due to the fact that $x' \neq y'$. Thus $A_x = B_x = A$. A similar argument works for the case when $x' \notin [x, z]$, using the hypothesis that $x' \neq z'$. Thus the bigon of endpoints x, x' is either trivial or contained in A .

In the same way we obtain that the bigons of endpoints y, y' and z, z' are either trivial or contained in A . Thus in all cases $T \subset A$, which by Lemma 3.13 implies that $\Delta \subset A$. \square

5 Quasi-isometric rigidity of relatively hyperbolic groups

In this section we prove the following.

Theorem 5.1. *Let G be an infinite finitely generated group endowed with a word metric dist , (G, dist) properly asymptotically tree-graded with respect to a collection \mathcal{A} of subsets. Then G is either hyperbolic or it is properly relatively hyperbolic with respect to a finite family of finitely generated subgroups $\{H_1, \dots, H_m\}$ such that every H_i is contained in $\mathcal{N}_{\varkappa}(A_i)$ for some $A_i \in \mathcal{A}$, where \varkappa is the maximum between the constant M in (β_2) and the constant χ in (β_3) .*

An explanation of the fact that the case when G is hyperbolic is treated separately can be found in Remark 1.5.

Proof. The pair (G, \mathcal{A}) satisfies properties (α_1) , (β_2) and (β_3) .

By Corollary 4.20, if for $\vartheta > 0$ and $\nu \geq 8$ from (β_3) either there exists no (ϑ, ν) -fat geodesic hexagon, or the (ϑ, ν) -fat geodesic hexagons have uniformly bounded diameter, then G is hyperbolic.

Assume from now on that for every $\eta > 0$ there exists a (ϑ, ν) -fat geodesic hexagon with diameter at least η .

For \varkappa as defined in the statement of the theorem and diam_{\varkappa} given by property (α_1) of \mathcal{A} , consider the set

$$\Phi = \{P \text{ geodesic hexagon ; } P \text{ is } (\vartheta, \nu)\text{-fat , } \text{diameter}(P) \geq \text{diam}_{\varkappa} + 1\}.$$

Let $P \in \Phi$ and $g \in G$. Then $g^{-1}P$ is a (ϑ, ν) -fat geodesic hexagon, therefore by (β_3) it is contained in $\mathcal{N}_{\varkappa}(A)$ for some $A \in \mathcal{A}$. If $g^{-1}P$ is also contained in $\mathcal{N}_{\varkappa}(A')$ with $A' \in \mathcal{A}$ then $\mathcal{N}_{\varkappa}(A) \cap \mathcal{N}_{\varkappa}(A')$ has diameter at least the diameter of P , hence at least $\text{diam}_{\varkappa} + 1$, consequently $A = A'$. Thus P defines a map

$$\mathbf{A}_P : G \rightarrow \mathcal{A}, \mathbf{A}_P(g) = A \text{ such that } g^{-1}P \subset \mathcal{N}_{\varkappa}(A) \Leftrightarrow P \subset \mathcal{N}_{\varkappa}(gA).$$

We may then define

$$\mathbf{A} : \Phi \rightarrow \text{Map}(G, \mathcal{A}), \mathbf{A}(P) = \mathbf{A}_P,$$

where $\text{Map}(G, \mathcal{A})$ is the set of maps from G to \mathcal{A} . Consider the equivalence relation on Φ induced by \mathbf{A} , that is

$$P \sim P' \Leftrightarrow \mathbf{A}(P) = \mathbf{A}(P') \Leftrightarrow \forall g \in G, g^{-1}P \text{ and } g^{-1}P' \text{ are in the same } \mathcal{N}_\varkappa(A).$$

Let $[P]$ be the equivalence class of $P \in \Phi$. To it we associate the set

$$B[P] = \bigcap_{g \in G} \mathcal{N}_\varkappa(g \mathbf{A}_P(g))$$

Proposition 5.2. *The metric space (G, dist) is properly ATG with respect to*

$$\mathcal{B} = \{B[P] ; [P] \in \Phi / \sim\}.$$

Proof. According to Proposition 4.29 it suffices to prove (α_1) , (β_2) and (β_3^η) for some $\eta > 0$. The proof relies on the simple remark that for every $r > 0$,

$$\mathcal{N}_r(B[P]) \subset \bigcap_{g \in G} \mathcal{N}_{r+\varkappa}(g \mathbf{A}_P(g)).$$

(α_1) Let $[P] \neq [P']$, which is equivalent to the fact that there exists $g_0 \in G$ such that $P \subset \mathcal{N}_\varkappa(g_0 A)$ and $P' \subset \mathcal{N}_\varkappa(g_0 A')$ with $A \neq A'$. For every $\delta > 0$,

$$\mathcal{N}_\delta(B[P]) \cap \mathcal{N}_\delta(B[P']) \subset \mathcal{N}_{\delta+\varkappa}(g_0 A) \cap \mathcal{N}_{\delta+\varkappa}(g_0 A') = g_0 [\mathcal{N}_{\delta+\varkappa}(A) \cap \mathcal{N}_{\delta+\varkappa}(A')].$$

Property (α_1) for \mathcal{A} implies that the diameter of $\mathcal{N}_\delta(B[P]) \cap \mathcal{N}_\delta(B[P'])$ is uniformly bounded.

(β_2) Let ϵ be the constant appearing in (β_2) for \mathcal{A} . Take $\epsilon' = \frac{\epsilon}{2}$ and take $M' = \frac{\epsilon+1}{\epsilon}\varkappa$. We prove that (β_2) holds for \mathcal{B} with the constants ϵ' and M' .

Let \mathbf{g} be a geodesic of length ℓ and let $[P] \in \Phi / \sim$ such that $\mathbf{g}(0)$ and $\mathbf{g}(\ell)$ are in $\mathcal{N}_{\epsilon'\ell}(B[P])$. It follows that for every $g \in G$, $\mathbf{g}(0)$ and $\mathbf{g}(\ell)$ are in $\mathcal{N}_{\epsilon'\ell+\varkappa}(g \mathbf{A}_P(g))$.

If $\varkappa \geq \frac{\epsilon}{2}\ell \Leftrightarrow \ell \leq \frac{2\varkappa}{\epsilon}$ then $\mathbf{g} \subset \overline{\mathcal{N}_\varkappa}(\{\mathbf{g}(0), \mathbf{g}(\ell)\}) \subset \mathcal{N}_{\varkappa+\varkappa}(B[P]) = \mathcal{N}_{M'}(B[P])$.

Assume that $\varkappa < \frac{\epsilon}{2}\ell$. Then for every $g \in G$ the geodesic $g^{-1}\mathbf{g}$ of length ℓ has its end-points in $\mathcal{N}_{\epsilon\ell}(\mathbf{A}_P(g))$. Property (β_2) implies that $g^{-1}\mathbf{g}([\frac{\ell}{3}, \frac{2\ell}{3}])$ is contained in $\mathcal{N}_M(\mathbf{A}_P(g)) \subset \mathcal{N}_\varkappa(\mathbf{A}_P(g))$.

We have thus obtained that for every $g \in G$, $\mathbf{g}([\frac{\ell}{3}, \frac{2\ell}{3}])$ is contained in $\mathcal{N}_\varkappa(g \mathbf{A}_P(g))$. It follows that $\mathbf{g}([\frac{\ell}{3}, \frac{2\ell}{3}])$ is contained in $B[P]$.

Property (β_3^η) holds for ϑ and ν the same as in (β_3) for \mathcal{A} , $\chi = 0$, and $\eta = \text{diam}_\varkappa + 1$. Indeed every $P \in \Phi$ is contained in $B[P]$. \square

Lemma 5.3 (the group permutes the pieces). (1) *If $P \sim P'$ and $g \in G$ then $gP \sim gP'$.*

Consequently G acts on the left on Φ / \sim .

(2) *For every $g \in G$ and $P \in \Phi$, $gB[P] = B[gP]$.*

Proof. (1) The set $\mathbf{A}_P(\gamma)$ is defined by $P \subset \mathcal{N}_\varkappa(\gamma \mathbf{A}_P(\gamma))$. For every $g \in G$, $gP \subset \mathcal{N}_\varkappa(g\gamma \mathbf{A}_P(\gamma))$, hence $\mathbf{A}_{gP}(g\gamma) = \mathbf{A}_P(\gamma)$. From this can be deduced that $P \sim P' \Rightarrow gP \sim gP'$.

(2) The translate $gB[P] = \bigcap_{\gamma \in G} \mathcal{N}_\varkappa(g\gamma \mathbf{A}_P(\gamma))$ is equal to

$$\bigcap_{\gamma \in G} \mathcal{N}_\varkappa(g\gamma \mathbf{A}_{gP}(g\gamma)) = \bigcap_{\gamma' \in G} \mathcal{N}_\varkappa(\gamma' \mathbf{A}_{gP}(\gamma')) = B[gP].$$

\square

The following statement finishes the proof of Theorem 5.1.

Proposition 5.4 (equivariant ATG structure implies relative hyperbolicity). *Let G be a finitely generated group which endowed with a word metric is properly ATG with respect to a collection of subsets \mathcal{B} , such that G permutes the subsets in \mathcal{B} . Assume moreover that for every $D > 0$ the following set is non-empty:*

$$\mathcal{B}_D = \{B \in \mathcal{B} ; B \text{ of diameter at least } D\} .$$

Then G is properly hyperbolic relative to a finite family of finitely generated subgroups $\{H_1, \dots, H_m\}$ such that every H_i satisfies $H_i \subset B_i \subset \mathcal{N}_K(H_i)$ for a uniquely determined $B_i \in \mathcal{B}$, where K is a constant depending on G and \mathcal{B} .

Remark 5.5. In the proof of Theorem 5.1, the remaining case was the one when for every $\eta > 0$ there exists a (ϑ, ν) -fat geodesic hexagon of diameter at least η . This and property (β_3) for \mathcal{B} imply that in that case, for any $D > 0$ the set \mathcal{B}_D is non-empty.

The proof of Proposition 5.4 is done in several steps.

Lemma 5.6. *Finitely many subsets in \mathcal{B} contain 1.*

Proof. Property (Ω) applied to \mathcal{B} implies that there exists $\tau > 0$ such that for any x, y in some $B \in \mathcal{B}$ any geodesic $[x, y]$ is contained in $\mathcal{N}_\tau(B)$. Property (α_1) for \mathcal{B} implies that there exists D_τ such that for $B \neq B'$, $\mathcal{N}_\tau(B) \cap \mathcal{N}_\tau(B')$ has diameter at most D_τ .

Assume that $B \in \mathcal{B}$ contains 1 and has diameter at most $3D_\tau$. Then $B \subset \overline{B}(1, 3D_\tau)$. As $\overline{B}(1, 3D_\tau)$ is finite, only finitely many $B \in \mathcal{B}$ can be in this case.

Assume that B contains 1 and has diameter larger than $3D_\tau$. Then B contains some point x with $\text{dist}(1, x) > 3D_\tau$. The geodesic $[1, x]$ is contained in $\mathcal{N}_\tau(B)$ and it intersects $S(1, 2D_\tau)$. We define a map from the set $\{B \in \mathcal{B} ; 1 \in B, \text{diam } B > 3D_\tau\}$ to the set of subsets of $S(1, 2D_\tau)$, associating to each B the non-empty intersection $\mathcal{N}_\tau(B) \cap S(1, 2D_\tau)$. This map is injective, by (α_1) and the choice of D_τ , and since the set of subsets of $S(1, 2D_\tau)$ is finite, so is the considered subset of \mathcal{B} . \square

Let $\mathcal{F} = \{B_1, B_2, \dots, B_k\}$ be the set of $B \in \mathcal{B}$ containing 1. For every $i \in \{1, 2, \dots, k\}$ let

$$\mathcal{I}_i = \{j \in \{1, 2, \dots, k\} \mid \exists g \in G \text{ such that } gB_i = B_j\} .$$

For every $j \in \mathcal{I}_i$ we fix $g_j \in G$ such that $g_j B_i = B_j$.

Notation 5.7. Define the constants $\mathcal{K}_i = \max_{j \in \mathcal{I}_i} \text{dist}(1, g_j)$ and $\mathcal{K} = \max_{1 \leq i \leq k} \mathcal{K}_i$.

Lemma 5.8. *For every $B \in \mathcal{B}$ the stabilizer $\text{Stab}(B) = \{g \in G \mid gB = B\}$ is a subgroup of G acting \mathcal{K} -transitively on B .*

Proof. Let x and b be arbitrary points in B . Both sets $b^{-1}B$ and $x^{-1}B$ contain 1 and by Lemma 5.3 they both are in \mathcal{B} . It follows that $b^{-1}B = B_i$ for some $i \in \{1, 2, \dots, k\}$. Also $x^{-1}B = B_j$ and since $b^{-1}xB_j = B_i$ it follows that $j \in \mathcal{I}_i$ and that $B_j = g_j B_i$. The last equality can be re-written as $x^{-1}B = g_j b^{-1}B$ which implies that $xg_j b^{-1} \in \text{Stab}(B)$, hence that x is at distance at most $\text{dist}(1, g_j)$ from $\text{Stab}(B)b$. \square

Corollary 5.9. *For $i \in \{1, 2, \dots, k\}$, $\text{Stab}(B_i) \subset B_i \subset \mathcal{N}_K(\text{Stab}(B_i))$.*

Notation 5.10. Denote by B' the set $\mathcal{B}_{\text{diam}_{2\mathcal{K}}+1}$, where $\text{diam}_{2\mathcal{K}}$ is the uniform bound given by property (α_1) for (G, \mathcal{B}) and $\delta = 2\mathcal{K}$.

By hypothesis \mathcal{B}' is non-empty. Proposition 5.2 and Corollary 4.31 imply that G is ATG with respect to \mathcal{B}' . Obviously G also permutes the subsets in \mathcal{B}' .

Let $\mathcal{F}' = \mathcal{F} \cap \mathcal{B}'$. Let \mathcal{F}_0 be a subset of \mathcal{F}' such that for every $B \in \mathcal{F}'$, its orbit $G \cdot B$ intersects \mathcal{F}_0 in a unique element. Such a subset can be obtained for instance by considering one by one the elements B_i in \mathcal{F}' , and deleting from \mathcal{F}' all B_j with $j \in \mathcal{I}_i$, $j \neq i$.

It follows that for every $B \in \mathcal{B}'$, the orbit $G \cdot B$ intersects \mathcal{F}_0 in only one element.

Let $\bar{B}_1, \dots, \bar{B}_m$ be the elements of \mathcal{F}_0 .

Lemma 5.11. *For every $B \in \mathcal{B}'$ there exists a unique $j \in \{1, 2, \dots, m\}$ and a unique left coset $g\text{Stab}(\bar{B}_j)$ such that*

$$g\text{Stab}(\bar{B}_j) \subset B \subset \mathcal{N}_K(g\text{Stab}(\bar{B}_j)) . \quad (3)$$

Proof. Existence. Let $g \in B$. Then $g^{-1}B \in \mathcal{B}'$ and $1 \in g^{-1}B$. Therefore $g^{-1}B = \bar{B}_j$ for some $j \in \{1, 2, \dots, m\}$. Corollary 5.9 implies the double inclusion (3).

Unicity. Assume that $g\text{Stab}(\bar{B}_j)$ and $g'\text{Stab}(\bar{B}_l)$ both satisfy (3), for $j, l \in \{1, 2, \dots, m\}$. Then

$$g\bar{B}_j \subset \mathcal{N}_K(g\text{Stab}(\bar{B}_j)) \subset \mathcal{N}_K(B) \subset \mathcal{N}_{2K}(g'\text{Stab}(\bar{B}_l)) \subset \mathcal{N}_{2K}(g'\bar{B}_l) .$$

On the other hand, both $g\bar{B}_j$ and $g'\bar{B}_l$ are in \mathcal{B}' by Lemma 5.3, in particular $g\bar{B}_j$ has diameter at least $\text{diam}_{2K} + 1$ which by property (α_1) implies that $g\bar{B}_j = g'\bar{B}_l$. According to the definition of \mathcal{F}_0 this can only happen if $j = l$. Thus $g^{-1}g'$ is in $\text{Stab}(\bar{B}_j)$. \square

Lemma 5.12. *The group G is properly hyperbolic relative to $\{H_1, \dots, H_m\}$, where $H_j = \text{Stab}(\bar{B}_j)$.*

Proof. The fact that G is ATG with respect to \mathcal{B}' and Lemmata 5.11 and 4.12 imply that G is ATG with respect to $\{gH_j ; g \in G/H_j, j \in \{1, 2, \dots, m\}\}$. In particular by (\mathfrak{Q}) each H_j is quasi-convex in G , which implies that H_j is finitely generated. Theorem 1.2 implies that G is hyperbolic relative to H_1, \dots, H_m .

If $G = H_j = \text{Stab}(\bar{B}_j)$ then Corollary 5.9 implies that $G = \bar{B}_j$. \square

A consequence of Theorem 5.1 is the following.

Theorem 5.13. *Let G be a finitely generated group properly hyperbolic relative to a finite family of finitely generated subgroups $\mathcal{H} = \{H_1, \dots, H_n\}$. If a group G' is (L, C) -quasi-isometric to G then G' is properly hyperbolic relative to $\mathcal{H}' = \{H'_1, \dots, H'_m\}$, where each H'_i can be embedded (λ, κ) -quasi-isometrically in H_j for some $j = j(i) \in \{1, 2, \dots, n\}$, where (λ, κ) depend on (L, C) and on (G, \mathcal{H}) .*

Proof. If the group G is finite then the group G' is also finite. We assume henceforth that both groups are infinite.

Let \mathfrak{q} be an (L, C) -quasi-isometry from G to G' , and let $\bar{\mathfrak{q}}$ be its quasi-converse, such that $\text{dist}(\mathfrak{q} \circ \bar{\mathfrak{q}}, \text{id}_{G'}) \leq D$ and $\text{dist}(\bar{\mathfrak{q}} \circ \mathfrak{q}, \text{id}_G) \leq D$, where $D = D(L, C)$. By Theorem 1.2, G is ATG with respect to the collection of left cosets $\mathcal{A} = \{gH_i ; g \in G/H_i, i \in \{1, 2, \dots, n\}\}$. Theorem 5.1 in [DS05b] implies that G' is ATG with respect to $\mathfrak{q}(\mathcal{A}) = \{\mathfrak{q}(A) ; A \in \mathcal{A}\}$. Moreover all constants appearing in the properties (α_i) , $i = 1, 2, 3$, (β_j) , $j = 1, 2$, and (\mathfrak{Q}) for $(G', \mathfrak{q}(\mathcal{A}))$ can be expressed as functions of (L, C) and of the constants in the same properties for (G, \mathcal{A}) .

Theorem 5.1 implies that G' is either hyperbolic or relatively hyperbolic with respect to $\{H'_1, \dots, H'_m\}$; moreover each H'_i is contained in $\mathcal{N}_\varkappa(\mathfrak{q}(A_i))$ for some $A_i \in \mathcal{A}$, where \varkappa is a constant depending on (L, C) , on the constant M in (β_2) for (G, \mathcal{A}) , and on the constant χ in (β_3) for (G, \mathcal{A}) .

Let $\pi_1 : \mathcal{N}_\varkappa(\mathfrak{q}(A_i)) \rightarrow \mathfrak{q}(A_i)$ be a map such that $\text{dist}(x, \pi_1(x)) \leq \varkappa$. Then π_1 is a $(1, 2\varkappa)$ -quasi-isometric embedding. Let $\pi_2 : \mathcal{N}_D(A_i) \rightarrow A_i$ be an $(1, 2D)$ -quasi-isometric embedding constructed similarly. The restriction to H'_i of $\pi_2 \circ \bar{q} \circ \pi_1$ is a (λ, κ) -quasi-isometric embedding of H'_i into $A_i = gH_j$, for some $j \in \{1, 2, \dots, n\}$, with (λ, κ) depending on (L, C) , \varkappa and D .

If G' is hyperbolic then G' is relatively hyperbolic with respect to $\{1\} \neq \{G'\}$ and all the statements in the theorem hold.

If $G' = H'_i$ then $G' = \mathcal{N}_\varkappa(\mathfrak{q}(A_i))$, which implies that $G \subset \mathcal{N}_C(\bar{q}(G')) \subset \mathcal{N}_{L\varkappa+2C+D}(A_i)$. By Theorem 1.2, this contradicts the fact that G is properly hyperbolic relative to \mathcal{H} . \square

References

- [BBI01] D. Burago, Y. Burago, and S. Ivanov, *A course in metric geometry*, Graduate Studies in Math., vol. 33, American Mathematical Society, Providence, RI, 2001.
- [BDM05] J. Behrstock, C. Druţu, and L. Mosher, *Thick metric spaces, relative hyperbolicity, and quasi-isometric rigidity*, Preprint ARXIV:MATH.GT/0512592, 2005.
- [BF01] J. Brock and B. Farb, *Curvature and rank of Teichmüller space*, preprint, ARXIV:MATH.GT/0109045, 2001.
- [Bou65] Nicolas Bourbaki, *Topologie générale*, Hermann, Paris, 1965.
- [Bow97] B. Bowditch, *Relatively hyperbolic groups*, Preprint, University of Southampton <http://www.maths.soton.ac.uk/pure/preprints.phtml>, 1997.
- [Cho96] R. Chow, *Groups quasi-isometric to complex hyperbolic space*, Trans. AMS **348** (1996), no. 5, 1757–1769.
- [CJ94] A. Casson and D. Jungreis, *Convergence groups and Seifert fibered 3-manifolds*, Invent. Math. **118** (1994), 441–456.
- [Dah03a] F. Dahmani, *Combination of convergence groups*, Geometry and Topology **7** (2003), 933–963.
- [Dah03b] F. Dahmani, *Les groupes relativement hyperboliques et leurs bords*, Ph.D. thesis, University of Strasbourg, 2003.
- [dDW84] L. Van den Dries and A. Wilkie, *On Gromov’s theorem concerning groups of polynomial growth and elementary logic*, J. Algebra **89** (1984), 349–374.
- [Dru00] C. Druţu, *Quasi-isometric classification of non-uniform lattices in semisimple groups of higher rank*, Geom. Funct. Anal. **10** (2000), no. 2, 327–388.
- [Dru02] C. Druţu, *Quasi-isometry invariants and asymptotic cones*, Int. J. Algebra Comput. **12** (2002), 99–135.
- [DS05a] C. Druţu and M. Sapir, *Relatively hyperbolic groups with rapid decay property*, Int. Math. Res. Notices **19** (2005), 1181–1194.
- [DS05b] C. Druţu and M. Sapir, *Tree-graded spaces and asymptotic cones of groups*, Topology **44** (2005), 959–1058.

- [Dyu00] A. Dyubina, *Instability of the virtual solvability and the property of being virtually torsion-free for quasi-isometric groups*, Internat. Math. Res. Notices **21** (2000), 1097–1101.
- [EF97] A. Eskin and B. Farb, *Quasi-flats and rigidity in higher rank symmetric spaces*, Jour. AMS **10** (1997), no. 3, 653–692.
- [EFW05] A. Eskin, D. Fisher, and K. Whyte, *Quasi-isometries and rigidity of solvable groups*, Preprint ARXIV:MATH.GR/0511647, 2005.
- [Esk98] A. Eskin, *Quasi-isometric rigidity of nonuniform lattices in higher rank symmetric spaces*, Jour. AMS **11** (1998), no. 2, 321–361.
- [Far98] B. Farb, *Relatively hyperbolic groups*, Geom. Funct. Anal. **8** (1998), no. 5, 810–840.
- [FM98] B. Farb and L. Mosher, *A rigidity theorem for the solvable Baumslag-Solitar groups*, Invent. Math. **131** (1998), no. 2, 419–451, Appendix by D. Cooper.
- [FM99] B. Farb and L. Mosher, *Quasi-isometric rigidity for the solvable Baumslag-Solitar groups, II*, Invent. Math. **137** (1999), no. 3, 613–649.
- [FS96] B. Farb and R. Schwartz, *The large-scale geometry of Hilbert modular groups*, J. Diff. Geom. **44** (1996), no. 3, 435–478.
- [Gab92] D. Gabai, *Convergence groups are Fuchsian groups*, Ann. of Math. **136** (1992), 447–510.
- [GdlH90] E. Ghys and P. de la Harpe, *Sur les groupes hyperboliques d’apres Mikhael Gromov*, Progress in Mathematics, no. 83, Birkhauser, 1990.
- [Gro81] M. Gromov, *Groups of polynomial growth and expanding maps*, IHES Sci. Publ. Math. **53** (1981), 53–73.
- [Gro87] M. Gromov, *Hyperbolic groups*, Essays in group theory (S. Gersten, ed.), MSRI Publications, vol. 8, Springer, 1987.
- [Gro93] M. Gromov, *Asymptotic invariants of infinite groups*, Geometric Group Theory, Vol. 2 (Sussex, 1991) (G. Niblo and M. Roller, eds.), LMS Lecture Notes, vol. 182, Cambridge Univ. Press, 1993, pp. 1–295.
- [Gui04] V. Guirardel, *Limit groups and groups acting freely on \mathbb{R}^n -trees*, Geom. and Topol. **8** (2004), 1427–1470.
- [KL95] M. Kapovich and B. Leeb, *On asymptotic cones and quasi-isometry classes of fundamental groups of nonpositively curved manifolds*, Geom. Funct. Anal. **5** (1995), no. 3, 582–603.
- [KL97a] M. Kapovich and B. Leeb, *Quasi-isometries preserve the geometric decomposition of haken manifolds*, Invent. Math. **128** (1997), no. 2, 393–416.
- [KL97b] B. Kleiner and B. Leeb, *Rigidity of quasi-isometries for symmetric spaces and euclidean buildings*, IHES Publ. Math. **86** (1997), 115–197.
- [Osi06] D. Osin, *Relatively hyperbolic groups: intrinsic geometry, algebraic properties, and algorithmic problems*, Mem. AMS **179** (2006), no. 843, vi+100pp.

- [Pan89] P. Pansu, *Metriques de Carnot-Caratheodory et quasiisometries des espaces symmetriques de rang un*, Ann. of Math. **129** (1989), 1–60.
- [PW02] P. Papasoglu and K. Whyte, *Quasi-isometries between groups with infinitely many ends*, Comment. Math. Helv. **77** (2002), no. 1, 133–144.
- [Sch96a] R. Schwartz, *Quasi-isometric rigidity and Diophantine approximation*, Acta Math. **177** (1996), 75–112.
- [Sch96b] R. Schwartz, *The quasi-isometry classification of rank one lattices*, IHES Sci. Publ. Math. **82** (1996), 133–168.
- [Sul81] D. Sullivan, *On the ergodic theory at infinity of an arbitrary discrete group of hyperbolic motions*, Riemann surfaces and related topics, Proceedings of the 1978 Stony Brook Conference, Ann. Math. Studies, vol. 97, Princeton University Press, 1981, pp. 465–496.
- [Tuk86] P. Tukia, *On quasi-conformal groups*, J. D’Anal. Math. **46** (1986), 318–346.
- [Tuk88] P. Tukia, *Homeomorphic conjugates of fuchsian groups*, J. Reine Angew. Math. **391** (1988), 1–54.
- [Tuk94] P. Tukia, *Covergence groups and gromov’s metric hyperbolic spaces*, New Zealand J. Math. **23** (1994), 157–187.
- [Val04] A. Valette, *Nouvelles approches de la propriété (T) de Kazhdan*, Astérisque **294** (2004), 97–124.
- [Yam04] A. Yaman, *A topological characterisation of relatively hyperbolic groups*, J. Reine Angew. Math. (Crelle’s Journal) **566** (2004), 41–89.

Cornelia Druțu
 UFR de Mathématiques
 Université de Lille 1
 F-59655 Villeneuve d’Ascq
 FRANCE
 Cornelia.Drutu@math.univ-lille1.fr