# Gauss Decomposition of the Yangian $Y(\mathfrak{gl}_{m|n})$

Lucy Gow\*
University of Sydney

November 21, 2018

Here we describe a Gauss decomposition of the Yangian  $Y(\mathfrak{gl}_{m|n})$  of the general linear Lie superalgebra. This gives a connection between this Yangian and the Yangian of the classical Lie superalgebra Y(A(m-1,n-1)) defined and studied in papers by Stukopin. We also show that the coefficients of the quantum Berezinian generate the centre of the Yangian  $Y(\mathfrak{gl}_{m|n})$ . This is a conjecture of Nazarov dating back to 1991.

Such Gauss decompositions have already been studied for the Yangian  $Y(\mathfrak{gl}_n)$  in papers by Brundan and Kleshchev, and Crampé, as well as for the super-Yangian  $Y(\mathfrak{gl}_{1|1})$  in the work of Jin-fang Cai, Guo-xing Ju, Ke Wu and Shi-kun Wang.

#### 1 Introduction

The Yangian  $Y(\mathfrak{gl}_{m|n})$  is the  $\mathbb{Z}_2$ -graded associative algebra over  $\mathbb{C}$  with generators

$$\{t_{ij}^{(r)} \mid 1 \le i, j \le m + n; r \ge 1\}$$

and defining relations

$$[t_{ij}^{(r)}, t_{kl}^{(s)}] = (-1)^{\overline{i}\,\overline{j} + \overline{i}\,\overline{k} + \overline{j}\,\overline{k}} \sum_{p=0}^{\min(r,s)-1} (t_{kj}^{(p)} t_{il}^{(r+s-1-p)} - t_{kj}^{(r+s-1-p)} t_{il}^{(p)}). \tag{1.1}$$

where  $\overline{i}$  is the parity of the index i, and we write square brackets for the super-commutator. We take  $\overline{i} = 0$  for  $i \leq m$ ; and  $\overline{i} = 1$  for  $i \geq m + 1$ . We define the formal power series

$$t_{ij}(u) = \delta_{ij} + t_{ij}^{(1)}u^{-1} + t_{ij}^{(2)}u^{-2} + \dots$$

and a matrix

$$T(u) = \sum_{i,j=1}^{m+n} t_{ij}(u) \otimes E_{ij} (-1)^{\overline{j}(\overline{i}+1)}$$
(1.2)

<sup>\*</sup>lucyg@maths.usyd.edu.au

where  $E_{ij}$  is the standard elementary matrix. (Here we identify an operator  $\sum A_{ij} \otimes E_{ij} (-1)^{\overline{j}(\overline{i}+1)}$  in  $Y(\mathfrak{gl}_{m|n})[[u^{-1}]] \otimes \operatorname{End} \mathbb{C}^{m|n}$  with the matrix  $(A_{ij})_{i,j=1}^{m+n}$ . The extra sign ensures that the product of two matrices can still be calculated in the usual way). Then, as for the Yangian  $Y(\mathfrak{gl}_n)$  (see for example [2, 5, 14]), the defining relations may be expressed by the matrix product

$$R(u-v)T_1(u)T_2(v) = T_2(v)T_1(u)R(u-v)$$

where

$$R(u-v) = 1 - \frac{1}{(u-v)}P_{12}$$

and  $P_{12}$  is the permutation matrix:  $P_{12} = \sum_{i,j=1}^{m+n} E_{ij} \otimes E_{ji} (-1)^{\bar{j}}$ . We also have the following equivalent form of the defining relations:

$$[t_{ij}(u), t_{kl}(v)] = \frac{(-1)^{\overline{i}\,\overline{j} + \overline{i}\,\overline{k} + \overline{j}\,\overline{k}}}{(u - v)} (t_{kj}(u)t_{il}(v) - t_{kj}(v)t_{il}(u)). \tag{1.3}$$

The Yangian  $Y(\mathfrak{gl}_{m|n})$  is a Hopf algebra with comultiplication

$$\Delta: t_{ij}(u)(-1)^{\overline{j}(\overline{i}+1)} \mapsto \sum_{k=1}^{m+n} t_{ik}(u)(-1)^{\overline{k}(\overline{i}+1)} \otimes t_{kj}(u)(-1)^{\overline{j}(\overline{k}+1)}, \tag{1.4}$$

antipode  $S: T(u) \mapsto T(u)^{-1}$  and counit  $\epsilon: T(u) \mapsto 1$ . Throughout this article we observe the following notation for entries of the inverse of the matrix T(u):

$$T(u)^{-1} =: (t'_{ij}(u))_{i,j=1}^n.$$

A straightforward calculation yields the following relation in  $Y(\mathfrak{gl}_{m|n})$ :

$$[t_{ij}(u), t'_{kl}(v)] = \frac{(-1)^{\overline{i}\,\overline{j} + \overline{i}\,\overline{k} + \overline{j}\,\overline{k}}}{(u - v)} \cdot (\delta_{kj} \sum_{s=1}^{m+n} t_{is}(u)t'_{sl}(v) - \delta_{il} \sum_{s=1}^{m+n} t'_{ks}(v)t_{sj}(u)). \tag{1.5}$$

The Yangian  $Y(\mathfrak{gl}_{m|n})$  was introduced in [15]. It has applications in mathematical physics because it describes symmetry in integrable systems such as  $\mathfrak{gl}_{m|n}$ -color Calogero-Sutherland models [1, 12], superstrings in  $AdS_5 \times S^5$  [11], and in the hierarchy of a form of the non-linear super-Schrödinger equation with m bosons and n fermions [4]. Vladimir Stukopin [17, 18] has also introduced Yangians for classical simple Lie superalgebras.

In this article we provide a new presentation for the Yangian  $Y(\mathfrak{gl}_{m|n})$  that allows us to relate it to the Yangian  $Y(\mathfrak{sl}_{m|n})$  studied by Stukopin. This follows a similar treatment of the Yangians  $Y(\mathfrak{gl}_N)$  in [2, 6], and  $Y(\mathfrak{gl}_{1|1})$  in [3].

### 2 Filtrations

We may define two different filtrations on the Yangian  $Y(\mathfrak{gl}_{m|n})$ . These are defined by setting the degree of a generator as follows:

$$\deg_1(t_{ij}^{(r)}) = r; \qquad \deg_2(t_{ij}^{(r)}) = r - 1.$$

Let  $\operatorname{gr}_1 Y(\mathfrak{gl}_{m|n})$  and  $\operatorname{gr}_2 Y(\mathfrak{gl}_{m|n})$ , respectively, denote the corresponding graded algebras.

## 3 The Poincaré-Birkhoff-Witt Theorem for Super Yangians

In this section we prove the Poincaré-Birkhoff-Witt theorem for the Yangian  $Y(\mathfrak{gl}_{m|n})$ . The proof is based very closely on that of the corresponding theorem for  $Y(\mathfrak{gl}_N)$  given in [2].

The map

$$\pi: t_{ij}(u) \mapsto \delta_{ij} + E_{ij}(-1)^{\overline{i}} u^{-1}$$

defines a projection from the Yangian  $Y(\mathfrak{gl}_{m|n})$  onto the universal enveloping algebra  $U(\mathfrak{gl}_{m|n})$ . For each positive integer  $l \geq 1$ , we define a homomorphism

$$\kappa_l := (\pi \otimes \cdots \otimes \pi) \circ \Delta^{(l)} : Y(\mathfrak{gl}_{m|n}) \to U(\mathfrak{gl}_{m|n})^{\otimes l}$$

where  $\Delta^{(l)}: Y(\mathfrak{gl}_{m|n}) \to Y(\mathfrak{gl}_{m|n})^{\otimes l}$  is the coproduct iterated (l-1) times. Then

$$\kappa_l(t_{ij}^{(r)}) = \sum_{1 \le s_1 \le \dots \le s_r \le l} \sum_{1 \le i_1, \dots, i_{r-1} \le m+n} E_{ii_1}^{[s_1]} E_{i_1 i_2}^{[s_2]} \cdots E_{i_{r-1} j}^{[s_r]} (-1)^{\overline{i} \, \overline{i}_1 + \overline{i}_1 \overline{i}_2 + \dots + \overline{i}_{r-1} \overline{j} + \overline{i} + \overline{j}}$$

where  $E_{ij}^{[s]} = 1^{\otimes (s-1)} \otimes E_{ij} \otimes 1^{(l-s)}$ . For any  $r > l \ge 1$ , we have  $\kappa_l(t_{ij}^{(r)}) = 0$ .

**Theorem 1.** Suppose we have fixed some ordering on the generators  $t_{ij}^{(r)}$   $(1 \le i, j \le m+n; r \ge 1)$  for the Yangian  $Y(\mathfrak{gl}_{m|n})$ . Then the ordered products of these, containing no second or higher order powers of the odd generators, form a basis for  $Y(\mathfrak{gl}_{m|n})$ .

*Proof.* <sup>1</sup> By relation (1.1), the graded algebra  $\operatorname{gr}_1Y(\mathfrak{gl}_{m|n})$  is supercommutative, and thus the set of all ordered monomials in the generators  $t_{ij}^{(r)}$  (with no second and higher order powers of the odd generators) span the Yangian  $Y(\mathfrak{gl}_{m|n})$ . It remains to show that they are linearly independent.

We show that, for every  $l \geq 1$ , the corresponding monomials in  $\{\kappa_l(t_{ij}^{(r)}) \mid 1 \leq r \leq l\}$  are linearly independent in  $\kappa_l(Y(\mathfrak{gl}_{m|n}))$ . Consider the filtration

$$F_0U(\mathfrak{gl}_{m|n})^{\otimes l} \subseteq F_1U(\mathfrak{gl}_{m|n})^{\otimes l} \subseteq F_2U(\mathfrak{gl}_{m|n})^{\otimes l} \subseteq \dots$$

on  $U(\mathfrak{gl}_{m|n})^{\otimes l}$  defined be setting each generator  $E_{ij}^{[r]}$  to be of degree 1. Then the associated graded algebra gr $U(\mathfrak{gl}_{m|n})^{\otimes l}$  is the polynomial algebra on supersymmetric generators

$$x_{ij}^{[r]} := \operatorname{gr}_1 e_{ij}^{[r]},$$

where  $x_{ij}^{[r]}$  is even if  $\overline{i} + \overline{j} = \overline{0}$  and odd if  $\overline{i} + \overline{j} = \overline{1}$ . The map  $\kappa_l$  preserves the filtration on the Yangian given by setting  $\deg_1(t_{ij}^{(r)}) = r$ , and thus defines a homomorphism between the corresponding graded algebras. It is enough to show that the same monomials in the elements  $y_{ij}^{(r)} = \operatorname{gr}_r \kappa_l(t_{ij}^{(r)})$  in the graded algebra are linearly independent. But for this, it is enough to show that the superderivatives  $dy_{ij}^{(r)}$  are linearly independent at a point. We have:

$$y_{ij}^{(r)} = \sum_{1 \le s_1 < \dots < s_r < l} \sum_{1 \le i_1, \dots, i_{r-1} \le n} x_{ii_1}^{[s_1]} x_{i_1 i_2}^{[s_2]} \cdots x_{i_{r-1} j}^{[s_r]} (-1)^{\overline{i} \, \overline{i}_1 + \overline{i}_1 \overline{i}_2 + \dots + \overline{i}_{r-1} \overline{j} + \overline{i} + \overline{j}}.$$

<sup>&</sup>lt;sup>1</sup>This theorem was stated in [19] but the proof there is incomplete.

We will show that the matrix  $d\phi$  corresponding to the map  $\left(dx_{ij}^{[s]}\right) \mapsto \left(dy_{ij}^{(r)}\right)$  has non-zero determinant at a point. It suffices to show that the determinant of this matrix is nonzero even when the variables are specialized to  $x_{kl}^{(s)} = \delta_{kl} c_s (-1)^{\overline{k}}$  for some distinct  $c_s$   $(s \ge 1)$ . When the variables are specialized as described, we find:

$$dy_{ij}^{(r)} = \sum_{s=1}^{l} \sum_{\substack{1 \le s_1 < \dots < s_{r-1} \le l \\ s_i \ne s}} c_{s_1} c_{s_2} \cdots c_{s_{r-1}} (-1)^{\overline{i} + \overline{j} + \overline{i} \overline{j}} dx_{ij}^{[s]}.$$

Let J be the  $(m+n) \times (m \times n)$  matrix  $J = \left(\delta_{ij}(-1)^{\overline{i}+\overline{j}+\overline{i}\,\overline{j}}\right)$ . Then  $d\phi = J \otimes X_l$ , where

$$X_{l} = \begin{pmatrix} 1 & 1 & \dots & 1 \\ (c_{2} + c_{3} + \dots + c_{l}) & (c_{1} + c_{3} + \dots + c_{l}) & \dots & (c_{1} + c_{2} + \dots + c_{l-1}) \\ (\sum_{i,j\neq 1} c_{i}c_{j}) & (\sum_{i,j\neq 2} c_{i}c_{j}) & \dots & (\sum_{i,j\neq l} c_{i}c_{j}) \\ \vdots & & & \vdots \\ c_{2}c_{3}\cdots c_{l} & c_{1}c_{3}c_{4}\cdots c_{l} & \dots & c_{2}c_{3}\cdots c_{l-1} \end{pmatrix}.$$

We show by induction that  $\det X_l = \prod_{1 \leq i < j \leq l} (c_i - c_j) \neq 0$ , and hence  $\det d\phi \neq 0$ . Indeed, row-reducing  $X_l$  gives the following matrix:

$$\begin{pmatrix} 1 & \dots & 1 & 1 \\ (c_{l}-c_{1}) & \dots & (c_{l}-c_{l-1}) & 0 \\ (c_{l}-c_{1}) \sum_{\substack{i,j\neq 1 \\ i,j< l}} c_{i}c_{j} & \dots & (c_{l}-c_{l-1}) \sum_{\substack{i,j\neq l-1 \\ i,j< l}} c_{i}c_{j} & 0 \\ \vdots & & \vdots \\ (c_{l}-c_{1})c_{2}c_{3}\cdots c_{l-1} & \dots & (c_{l}-c_{l-1})c_{1}\cdots c_{l-2} & 0 \end{pmatrix},$$

which clearly has determinant  $(c_1 - c_l)(c_2 - c_l) \cdots (c_{l-1} - c_l) \det X_{l-1}$ .

Now, suppose we have some non-trivial linear combination P of the ordered monomials in  $t_{ij}^{(r)}$  (with no second or higher order powers of the odd generators) and take l to be any number greater than all the r that occur in P. Since the monomials in  $\kappa_l(t_{ij}^{(r)})$  are linearly independent in  $\kappa_l(Y(\mathfrak{gl}_{m|n}))$ , we must have  $\kappa_l(P) \neq 0$ . Therefore,  $P \neq 0$  in the Yangian.  $\square$ 

Now let  $\mathfrak{gl}_{m|n}[t]$  denote the algebra  $\mathfrak{gl}_{m|n} \otimes \mathbb{C}[t]$  with basis  $\{E_{ij}t^r\}_{1 \leq i,j \leq m+n,r \geq 0}$ .

Corollary 3.1. The graded algebra  $gr_2Y(\mathfrak{gl}_{m|n})$  is isomorphic to the algebra  $U(\mathfrak{gl}_{m|n}[x])$ , via the map

$$\operatorname{gr}_2 Y(\mathfrak{gl}_{m|n}) \to U(\mathfrak{gl}_{m|n}[x])$$
$$\operatorname{gr}_2^{r-1} t_{ij}^{(r)} \mapsto E_{ij} x^{r-1} (-1)^{\overline{i}} \quad (1 \le i, j \le m+n, r \ge 1).$$

## 4 Gauss Decomposition of T(u)

Here we describe a decomposition of the matrix T(u) in terms of the quasideterminants of Gelfand and Retakh [8].

**Definition 4.1.** Let X be a square matrix over a ring with identity such that its inverse matrix  $X^{-1}$  exists, and such that its (j,i)th entry is an invertible element of the ring. Then the (i,j)th quasideterminant of X is defined by the formula

$$|X|_{ij} = ((X^{-1})_{ji})^{-1} =: \begin{vmatrix} x_{11} & \cdots & x_{1j} & \cdots & x_{1n} \\ & \cdots & & \cdots & \\ x_{i1} & \cdots & \boxed{x_{ij}} & \cdots & x_{in} \\ & \cdots & & \cdots & \\ x_{n1} & \cdots & x_{nj} & \cdots & x_{nn} \end{vmatrix}.$$

By Theorem 4.96 in [8], the matrix T(u) defined in (1.2) has the following Gauss decomposition in terms of quasideterminants:

$$T(u) = F(u)D(u)E(u)$$

for unique matrices

$$D(u) = \begin{pmatrix} d_1(u) & \cdots & 0 \\ & d_2(u) & \vdots \\ \vdots & & \ddots \\ 0 & \cdots & d_{m+n}(u) \end{pmatrix},$$

$$E(u) = \begin{pmatrix} 1 & e_{12}(u) & \cdots & e_{1,m+n}(u) \\ & \ddots & & e_{2,m+n}(u) \\ & & \ddots & \vdots \\ 0 & & 1 \end{pmatrix}, F(u) = \begin{pmatrix} 1 & & \cdots & 0 \\ f_{21}(u) & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ f_{m+n,1}(u) & f_{m+n,2}(u) & \cdots & 1 \end{pmatrix},$$

where

$$d_{i}(u) = \begin{vmatrix} t_{11}(u) & \cdots & t_{1,i-1}(u) & t_{1i}(u) \\ \vdots & \ddots & & \vdots \\ t_{i1}(u) & \cdots & t_{i,i-1}(u) & t_{ii}(u) \end{vmatrix},$$

$$e_{ij}(u) = d_{i}(u)^{-1} \begin{vmatrix} t_{11}(u) & \cdots & t_{1,i-1}(u) & t_{1j}(u) \\ \vdots & \ddots & \vdots & & \vdots \\ t_{i-1,i}(u) & \cdots & t_{i-1,i-1}(u) & t_{i-1,j}(u) \\ t_{i1}(u) & \cdots & t_{i,i-1}(u) & t_{ii}(u) \end{vmatrix},$$

$$f_{ji}(u) = \begin{vmatrix} t_{11}(u) & \cdots & t_{1,i-1}(u) & t_{1i}(u) \\ \vdots & \ddots & \vdots & & \vdots \\ t_{i-1,1}(u) & \cdots & t_{i-1,i-1}(u) & t_{i-1,i}(u) \\ t_{ji}(u) & \cdots & t_{j,i-1}(u) & t_{ji}(u) \end{vmatrix} d_{i}(u)^{-1}.$$

We use the following notation for the coefficients:

$$d_{i}(u) = \sum_{r \geq 0} d_{i}^{(r)} u^{-r}; \quad (d_{i}(u))^{-1} = \sum_{r \geq 0} d_{i}^{\prime(r)} u^{-r};$$
  

$$e_{ij}(v) = \sum_{r \geq 1} e_{ij}^{(r)} v^{-r}; \quad f_{ji}(v) = \sum_{r \geq 1} f_{ji}^{(r)} v^{-r}.$$

It is easy to recover each generating series  $t_{ij}(u)$  by multiplying together and taking commutators of the series  $d_i(u)$ ,  $e_j(u) := e_{j,j+1}(u)$ , and  $f_j(v) := f_{j+1,j}(u)$  for  $1 \le i \le m+n$ ,  $1 \le j \le m+n-1$ . Indeed, for each pair i, j such that  $1 < i+1 < j \le m+n-1$ , we have:

$$e_{ij}^{(r)} = (-1)^{\overline{j-1}} [e_{i,j-1}^{(r)}, e_{j-1}^{(1)}]; \quad f_{ji}^{(r)} = (-1)^{\overline{j-1}} [f_{j-1}^{(1)}, f_{i,j-1}^{(r)}]. \tag{4.1}$$

Thus the Yangian  $Y(\mathfrak{gl}_{m|n})$  is generated by the coefficients of the series

$$\{d_i(u), e_j(u), f_j(u) \mid 1 \le i \le m+n; 1 \le j \le m+n-1\}.$$

## 5 Maps Between Yangians

For Yangians  $Y(\mathfrak{gl}_{m|n})$  with small m, n, such as  $Y(\mathfrak{gl}_{1|1})$  and  $Y(\mathfrak{gl}_{2|1})$ , it is feasible to use this matrix relationship T(u) = F(u)D(u)E(u) to translate the defining relations (1.3) into relations between the generating series  $d_i(u)$ ,  $e_j(u)$  and  $f_j(u)$ . In order to transfer these results to the general case of  $Y(\mathfrak{gl}_{m|n})$  we define various homomorphisms between Yangians.

**Lemma 5.1.** The map  $\rho_{m|n}: Y(\mathfrak{gl}_{m|n}) \to Y(\mathfrak{gl}_{n|m})$  defined by

$$\rho_{m|n}(t_{ij}(u)) = t_{m+n+1-i,m+n+1-j}(-u).$$

is an associative algebra isomorphism. (Note that we have swapped m and n in the above).

*Proof.* We check that the map  $\rho_{m|n}$  preserves the defining relation (1.3).

**Proposition 5.2.** Let  $\zeta_{m|n}: Y(\mathfrak{gl}_{m|n}) \to Y(\mathfrak{gl}_{n|m})$  be the associative algebra isomorphism given by  $\zeta_{m|n} = \rho_{m|n} \circ \omega_{m|n}$ , where  $\omega_{m|n}$  is the  $Y(\mathfrak{gl}_{m|n})$  automorphism given by  $\omega_{m|n}: T(u) \mapsto T(-u)^{-1}$ . That is,

$$\zeta_{m|n}: t_{ij}(u) \mapsto t'_{m+n+1-i,m+n+1-j}(u).$$

Then:

$$\zeta_{m|n} : \begin{cases}
d_i(u) & \mapsto (d_{m+n-i+1}(u))^{-1}, \\
e_k(u) & \mapsto -f_{m+n-k}(u), \\
f_k(u) & \mapsto -e_{m+n-k}(u),
\end{cases}$$
(5.1)

for  $1 \le i \le m + n \ and \ 1 \le k \le m + n - 1$ .

*Proof.* We multiply out the matrix products T(u) = F(u)D(u)E(u) and  $T(u)^{-1} = E(u)^{-1}D(u)^{-1}F(u)^{-1}$ . This shows that for all  $1 \le i < j \le m + n$ ,

$$t_{ii}(u) = d_i(u) + \sum_{k < i} f_{ik}(u) d_k(u) e_{ki}(u),$$

$$t_{ij}(u) = d_i(u) e_{ij}(u) + \sum_{k < i} f_{ik}(u) d_k(u) e_{kj}(u),$$

$$t_{ji}(u) = f_{ji}(u) d_i(u) + \sum_{k < i} f_{jk}(u) d_k(u) e_{ki}(u),$$

and

$$t'_{ii}(u) = d_i(u)^{-1} + \sum_{k>i} e'_{ik}(u) d_k(u)^{-1} f'_{ki}(u),$$

$$t'_{ij}(u) = e'_{ij}(u) d_j(u)^{-1} + \sum_{k>i} e'_{ik}(u) d_k(u)^{-1} f'_{kj}(u),$$

$$t'_{ji}(u) = d_j(u)^{-1} f'_{ji}(u) + \sum_{k>i} e'_{jk}(u) d_k(u)^{-1} f'_{ki}(u),$$

where

$$e'_{ij}(u) = \sum_{i=i_0 < i_1 < \dots < i_s = j} (-1)^s e_{i_0 i_1}(u) e_{i_1 i_2}(u) \cdots e_{i_{s-1} i_s}(u)$$

and

$$f'_{ji}(u) = \sum_{i=i_0 < i_1 < \dots < i_s = j} (-1)^s f_{i_s i_{s-1}}(u) \cdots f_{i_2 i_1}(u) f_{i_1 i_0}(u).$$

Then immediately we have  $\zeta_{m|n}(d_1(u)) = d_{m+n,m+n}(u)^{-1}$ ,  $\zeta_{m|n}(e_{1j}(u)) = f'_{m+n,m+n+1-j}(u)$ , and  $\zeta_{m|n}(f_{j1}(u)) = e'_{m+n+1-j,m+n}(u)$ . By induction on i, we derive:

$$\zeta_{m|n}(d_i(u)) = (d_{m+n+1-i}(u))^{-1}, 
\zeta_{m|n}(e_{ij}(u)) = f'_{m+n+1-i,m+n+1-j}(u), 
\zeta_{m|n}(f_{ji}(u)) = e'_{m+n+1-j,m+n+1-i}(u).$$

The result stated in the proposition is the special case of this where j = i + 1.

Finally, let  $\varphi_{m|n}: Y(\mathfrak{gl}_{m|n}) \hookrightarrow Y(\mathfrak{gl}_{m+k|n})$  be the inclusion which sends each  $t_{ij}^{(r)} \in Y(\mathfrak{gl}_{m|n})$  to the generator  $t_{k+i,k+j}^{(r)} \in Y(\mathfrak{gl}_{m+k|n})$ ; and let  $\psi_k: Y(\mathfrak{gl}_{m|n}) \to Y(\mathfrak{gl}_{m+k|n})$  be the injective homomorphism defined by

$$\psi_k = \omega_{m+k|n} \circ \varphi_{m|n} \circ \omega_{m|n}. \tag{5.2}$$

Then, for any  $1 \le i, j \le m + n$  (see Lemma 4.2 of [2]) we have:

$$\psi_k(t_{ij}(u)) = \begin{vmatrix} t_{11}(u) & \cdots & t_{1k}(u) & t_{1,k+j}(u) \\ \vdots & \ddots & \vdots & \vdots \\ t_{k1}(u) & \cdots & t_{kk}(u) & t_{k,k+j}(u) \\ t_{k+i,1}(u) & \cdots & t_{k+i,k}(u) & t_{k+i,k+j}(u) \end{vmatrix}.$$

As an immediate consequence we have the following lemma.

**Lemma 5.3.** For  $k, l \ge 1$ , we have

$$\psi_k(d_l(u)) = d_{k+l}(u),$$
  

$$\psi_k(e_l(u)) = e_{k+l}(u), and$$
  

$$\psi_k(f_l(u)) = f_{k+l}(u).$$

Notice that the map  $\psi_k$  sends  $t_{ij}^{\prime(r)} \in Y(\mathfrak{gl}_{m|n})$  to the element  $t_{k+i,k+j}^{\prime(r)}$  in  $Y(\mathfrak{gl}_{m+k|n})$ . Thus the subalgebra  $\psi_k(Y(\mathfrak{gl}_{m|n}))$  is generated by the elements  $\{t_{k+s,k+t}^{\prime(r)}\}_{s,t=1}^n$ . Then, by (1.5), all elements of this subalgebra commute with those of the subalgebra generated by the elements  $\{t_{ij}^{(r)}\}_{i,j=1}^k$ . This implies in particular that for any  $i,j\geq 1$ , the quasideterminants  $d_i(u)$  and  $d_j(v)$  commute.

#### 6 Gauss Decomposition of $Y(\mathfrak{gl}_{2|1})$

We begin by defining a presentation of the Yangian  $Y(\mathfrak{gl}_{2|1})$  using the Gauss decomposition above. We will then use this to give the more general result in the next section. We use the matrix relationship T(u) = F(u)D(u)E(u) to convert the defining relations (1.3) into relations between the generating series  $d_i(u)$ ,  $e_i(v)$  and  $f_i(v)$ . Note that in the Yangian  $Y(\mathfrak{gl}_{1|1})$ , and in the Yangian  $Y(\mathfrak{gl}_2)$ , we have the following

$$T(u) = \begin{pmatrix} d_1(u) & d_1(u) e_1(u) \\ f_1(u)d_1(u) & f_1(u)d_1(u)e_1(u) + d_2(u) \end{pmatrix}$$

$$T(v)^{-1} = \begin{pmatrix} d_1(v)^{-1} + e_1(v)d_2(v)^{-1}f_1(v) & -e_1(v)d_2(v)^{-1} \\ -d_2(v)^{-1}f_1(v) & d_2(v)^{-1} \end{pmatrix}.$$

$$(6.1)$$

$$T(v)^{-1} = \begin{pmatrix} d_1(v)^{-1} + e_1(v)d_2(v)^{-1}f_1(v) & -e_1(v)d_2(v)^{-1} \\ -d_2(v)^{-1}f_1(v) & d_2(v)^{-1} \end{pmatrix}.$$

$$(6.2)$$

whereas in the Yangian  $Y(\mathfrak{gl}_{2|1})$ ,

$$T(u) = \begin{pmatrix} d_1(u) & d_1(u)e_1(u) & d_1(u)e_{13}(u) \\ f_1(u)d_1(u) & f_1(u)d_1(u)e_1(u) + d_2(u) & f_1(u)d_1(u)e_{13}(u) + d_2(u)e_2(u) \\ f_{31}(u)d_1(u) & f_{31}(u)d_3(u)e_1(u) + f_2(u)d_3(u) & * \end{pmatrix},$$

$$T(v)^{-1} = \begin{pmatrix} * & * & (e_1(v)e_2(v) - e_{13}(v))d_3(v)^{-1} \\ * & d_2(v)^{-1} + e_2(v)d_3(v)^{-1}f_2(v) & -e_2(v)d_3(v)^{-1} \\ d_3(v)^{-1}(f_2(v)f_1(v) - f_{31}(v)) & -d_3(v)^{-1}f_2(v) & d_3(v)^{-1} \end{pmatrix}.$$

These expressions for the entries of T(u) allow us to derive the following relations.

**Lemma 6.1.** We have the following identities in  $Y(\mathfrak{gl}_{2|1})$ :

$$(u-v)[d_{i}(u), e_{j}(v)] = \begin{cases} (\delta_{i,j} - \delta_{i,j+1}) d_{i}(u)(e_{j}(v) - e_{j}(u)), & \text{if } j = 1; \\ (\delta_{i,j} + \delta_{i,j+1}) d_{i}(u)(e_{j}(v) - e_{j}(u)), & \text{if } j = 2; \end{cases}$$

$$(u-v)[d_{i}(u), f_{j}(v)] = \begin{cases} -(\delta_{ij} - \delta_{i,j+1})(f_{j}(v) - f_{j}(u))d_{i}(u), & \text{if } j = 1; \\ -(\delta_{ij} + \delta_{i,j+1})(f_{j}(v) - f_{j}(u))d_{i}(u), & \text{if } j = 2; \end{cases}$$

$$(u-v)[e_{j}(u), f_{k}(v)] = \begin{cases} \delta_{jk} \left(d_{j}(u)^{-1}d_{j+1}(u) - d_{j}(v)^{-1}d_{j+1}(v)\right), & \text{if } j = 1; \\ -\delta_{jk} \left(d_{j}(u)^{-1}d_{j+1}(u) - d_{j}(v)^{-1}d_{j+1}(v)\right), & \text{if } j = 2; \end{cases}$$

$$(u-v)[e_{j}(u), e_{j}(v)] = \begin{cases} (e_{j}(v) - e_{j}(u))^{2}, & \text{if } j = 1; \\ 0, & \text{if } j = 2; \end{cases}$$

$$(u-v)[f_{j}(u), f_{j}(v)] = \begin{cases} -(f_{j}(v) - f_{j}(u))^{2}, & \text{if } j = 1; \\ 0, & \text{if } j = 2; \end{cases}$$

$$(u-v)[e_{1}(u), e_{2}(v)] = e_{1}(u)e_{2}(v) - e_{1}(v)e_{2}(v) - e_{13}(u) + e_{13}(v),$$

$$(u-v)[f_{1}(u), f_{2}(v)] = -f_{2}(v)f_{1}(u) + f_{2}(v)f_{1}(v) + f_{31}(u) - f_{31}(v),$$

$$[[e_{i}(u), e_{j}(v)], e_{j}(w)] + [[e_{i}(u), e_{j}(w)], e_{j}(v)] = 0, & \text{if } |i-j| = 1;$$

$$[[f_{i}(u), f_{j}(v)], f_{i}(w)] + [[f_{i}(u), f_{i}(w)], f_{i}(v)] = 0, & \text{if } |i-j| = 1; \end{cases}$$

where unless otherwise indicated the indices i, j, k range over i = 1, 2, 3 and j, k = 1, 2.

*Proof.* We give a proof of just the first equation (6.3), since the rest are proven similarly. First, note that by the remarks at the end of the previous section,  $d_3(u) = \psi_2(d_1(u))$  commutes with  $e_1(v) = t_{11}(v)^{-1}t_{12}(v)$ . Similarly,  $e_2(v) = \psi_1(e_1(v))$  commutes with  $d_1(u)$ . Now consider the quasideterminants  $d_1(u), d_2(u)$  and  $e_1(v)$  in the algebra  $Y(\mathfrak{gl}_2)[[u^{-1}, v^{-1}]]$ . Here, we have the matrices T(u),  $T(v)^{-1}$  as in (6.1) and (6.2). By (1.5),

$$(u-v)[t_{11}(u), t'_{12}(v)] = t_{11}(u)t'_{12}(v) + t_{12}(u)t'_{22}(v),$$

but this is the same as

$$(u-v)[d_1(u), -e_1(v)d_2(v)^{-1}] = -d_1(u)e_1(v)d_2(v)^{-1} + d_1(u)e_1(u)d_2(v)^{-1}.$$

Cancelling  $d_2(v)$  on the right gives the desired equation when i = j = 1, but in  $Y(\mathfrak{gl}_2)[[u^{-1}, v^{-1}]]$ . We deduce the relation in  $Y(\mathfrak{gl}_{2|1}[[u^{-1},v^{-1}]])$  by following the natural inclusion  $Y(\mathfrak{gl}_2) \hookrightarrow$  $Y(\mathfrak{gl}_{2|1})$  which sends generators in  $Y(\mathfrak{gl}_2)$  to those of the same name in  $Y(\mathfrak{gl}_{2|1})$ .

For the result when i = 2, j = 1, we consider the commutator  $[t'_{22}(u), t_{12}(v)]$  in the algebra  $Y(\mathfrak{gl}_2)[[u^{-1},v^{-1}]]$  and make the same deduction. For the case j=2, we find the relations between  $d_1(u)$ ,  $d_2(u)$  and  $e_1(v)$  in the algebra  $Y(\mathfrak{gl}_{1|1})[[u^{-1}, v^{-1}]]$ , and map these into the algebra  $Y(\mathfrak{gl}_{2|1})[[u^{-1},v^{-1}]]$ , by following  $\psi_1:Y(\mathfrak{gl}_{1|1})\to Y(\mathfrak{gl}_{2|1})$ .

**Theorem 2.** The algebra  $Y(\mathfrak{gl}_{2|1})$  is generated by the even elements  $d_1^{(r)}$ ,  $d_2^{(r)}$ ,  $d_3^{(r)}$ ,  $d_1^{\prime}$ ,  $d_2^{\prime}$ ,  $d_3^{\prime(r)},~e_1^{(r)},~f_1^{(r)},~and~odd~elements~e_2^{(r)},f_2^{(r)}~with~r\geq 1,~subject~only~to~the~following~relations:$ 

$$d_{i}^{(0)} = 1,$$

$$\sum_{t=0}^{r} d_{i}^{(t)} d_{i}^{\prime(r-t)} = \delta_{r0},$$

$$[d_{i}^{(r)}, d_{l}^{(s)}] = 0,$$
(6.5)

$$[d_i^{(r)}, e_j^{(s)}] = \begin{cases} (\delta_{ij} - \delta_{i,j+1}) \sum_{t=1}^{r-1} d_i^{(t)} e_j^{(r+s-1-t)}, & \text{if } j = 1; \\ (\delta_{ij} + \delta_{i,j+1}) \sum_{t=1}^{r-1} d_i^{(t)} e_j^{(r+s-1-t)}, & \text{if } j = 2; \end{cases}$$
(6.6)

$$[d_{i}^{(r)}, e_{j}^{(s)}] = \begin{cases} (\delta_{ij} - \delta_{i,j+1}) \sum_{t=1}^{r-1} d_{i}^{(t)} e_{j}^{(r+s-1-t)}, & \text{if } j = 1; \\ (\delta_{ij} + \delta_{i,j+1}) \sum_{t=1}^{r-1} d_{i}^{(t)} e_{j}^{(r+s-1-t)}, & \text{if } j = 2; \end{cases}$$

$$[d_{i}^{(r)}, f_{j}^{(s)}] = \begin{cases} -(\delta_{i,j} - \delta_{i,j+1}) \sum_{t=1}^{r-1} f_{j}^{(r+s-1-t)} d_{i}^{(t)}, & \text{if } j = 1; \\ -(\delta_{i,j} + \delta_{i,j+1}) \sum_{t=1}^{r-1} f_{j}^{(r+s-1-t)} d_{i}^{(t)}, & \text{if } j = 2; \end{cases}$$

$$[e_{j}^{(r)}, f_{k}^{(s)}] = \begin{cases} -\delta_{jk} \sum_{t=0}^{r+s-1} d_{j}^{\prime} d_{j+1}^{(r+s-1-t)}, & \text{if } j = 1; \\ \delta_{jk} \sum_{t=0}^{r+s-1} d_{j}^{\prime} d_{j+1}^{(r+s-1-t)}, & \text{if } j = 2; \end{cases}$$

$$(6.8)$$

$$[e_j^{(r)}, f_k^{(s)}] = \begin{cases} -\delta_{jk} \sum_{t=0}^{r+s-1} d_j^{\prime(t)} d_{j+1}^{(r+s-1-t)}, & \text{if } j = 1; \\ \delta_{jk} \sum_{t=0}^{r+s-1} d_j^{\prime(t)} d_{j+1}^{(r+s-1-t)}, & \text{if } j = 2; \end{cases}$$
(6.8)

$$[e_1^{(r)}, e_1^{(s+1)}] - [e_1^{(r+1)}, e_1^{(s)}] = e_1^{(r)} e_1^{(s)} + e_1^{(s)} e_1^{(r)},$$

$$(6.9)$$

$$[f_1^{(r+1)}, f_1^{(s)}] - [f_1^{(r)}, f_1^{(s+1)}] = f_1^{(r)} f_1^{(s)} + f_1^{(s)} f_1^{(r)},$$

$$(6.10)$$

$$[e_{2}^{(r)}, e_{2}^{(s)}] = 0, \quad [f_{2}^{(r)}, f_{2}^{(s)}] = 0,$$

$$[e_{1}^{(r+1)}, e_{2}^{(s)}] - [e_{1}^{(r)}, e_{2}^{(s+1)}] = e_{1}^{(r)} e_{2}^{(s)},$$

$$[f_{1}^{(r+1)}, f_{2}^{(s)}] - [f_{1}^{(r)}, f_{2}^{(s+1)}] = -f_{2}^{(s)} f_{1}^{(r)},$$

$$[[e_{1}^{(r)}, e_{2}^{(s)}], e_{2}^{(t)}] + [[e_{1}^{(r)}, e_{2}^{(t)}], e_{2}^{(s)}] = 0,$$

$$[[f_{1}^{(r)}, f_{2}^{(s)}], f_{2}^{(t)}] + [[f_{1}^{(r)}, f_{2}^{(t)}], f_{2}^{(s)}] = 0,$$

$$[[e_{2}^{(r)}, e_{1}^{(s)}], e_{1}^{(t)}] + [[e_{2}^{(r)}, e_{1}^{(t)}], e_{1}^{(s)}] = 0,$$

$$[[f_{2}^{(r)}, f_{1}^{(s)}], f_{1}^{(t)}] + [[f_{2}^{(r)}, f_{1}^{(t)}], f_{1}^{(s)}] = 0$$

$$(6.12)$$

for all i, l = 1, 2, 3, j, k = 1, 2 and all admissible r, s, t.

**Remark 6.1.** Relations (6.9) and (6.10) are equivalent to the following relations:

$$[e_i^{(r)}, e_i^{(s)}] = \sum_{t=1}^{s-1} e_i^{(t)} e_i^{(r+s-1-t)} - \sum_{t=1}^{r-1} e_i^{(t)} e_i^{(r+s-1-t)}$$

$$[f_i^{(r)}, f_i^{(s)}] = \sum_{t=1}^{r-1} f_i^{(r+s-1-t)} f_i^{(t)} - \sum_{t=1}^{s-1} f_i^{(r+s-1-t)} f_i^{(t)}$$

*Proof.* First, we show that the corresponding coefficients of quasideterminants in the Yangian satisfy the relations in Theorem 2. The first three relations are obvious from the fact that the  $d_i(u)$ 's commute and the definition of the series  $d'_i(u) := (d_i(u))^{-1}$ . The rest follow from the relations in Lemma (6.1). We show the proof of only (6.6) and (6.9) since the rest are derived similarly.

Observe that for any formal series  $g(u) = \sum_{r>0} g^{(r)} u^{-r}$  we have the identity

$$\frac{g(u) - g(v)}{u - v} = -\sum_{r,s \ge 1} g^{(r+s-1)} u^{-r} v^{-s}.$$

Then, by (6.3),

$$[d_i(u),e_j(v)] = (\delta_{ij}-(-1)^{\delta_{i,2}}\delta_{i,j+1})(\sum_{t\geq 1}d_i^{(t)}u^{-t})(\sum_{p,s\geq 1}e_j^{(p+s-1)}u^{-p}v^{-s}).$$

Taking coefficients of  $u^{-r}v^{-s}$  gives (6.6).

Now consider (6.4). In the case where j = 1, this expands out as follows:

$$(u-v)[e_{1}(u), e_{1}(v)] = \left(\sum_{r\geq 1} e_{1}^{(r)} u^{-r} - \sum_{s\geq 1} e_{1}^{(s)} v^{-s}\right)^{2}$$

$$= -\sum_{r,s\geq 1} e_{1}^{(r)} e_{1}^{(s)} u^{-r} v^{-s} - \sum_{r,s\geq 1} e_{1}^{(s)} e_{1}^{(r)} u^{-r} v^{-s}$$

$$+ \sum_{r,s\geq 1} e_{1}^{(r)} e_{1}^{(s)} u^{-r-s} + \sum_{r,s\geq 1} e_{1}^{(r)} e_{1}^{(s)} v^{-r-s}.$$

Taking coefficients of  $u^{-r}v^{-s}$  on both sides gives the relation (6.9).

Now let  $\widehat{Y}$  be the algebra defined by the relations in Theorem 2. We have shown that there is an associative algebra homomorphism  $\widehat{Y} \to Y(\mathfrak{gl}_{2|1})$  taking each generator in  $\widehat{Y}$  to the quasideterminant coefficient of the same name in the Yangian. By (4.1) these elements generate the Yangian, so this homomorphism is surjective. We will now show that the algebra  $\widehat{Y}$  is spanned as a vector space by certain monomials, and that the images of these monomials form a basis for the Yangian  $Y(\mathfrak{gl}_{2|1})$ . It follows that the homomorphism is an isomorphism.

Let  $e_{13}^{(r)}$  and  $f_{31}^{(r)}$  be the elements of  $\widehat{Y}$  defined by

$$e_{13}^{(r)} = [e_1^{(r)}, e_2^{(1)}], \quad f_{31}^{(r)} = [f_1^{(r)}, f_2^{(1)}] \quad \text{(c.f. (4.1))}.$$

We want to show that the algebra  $\widehat{Y}$  is spanned by the set of ordered monomials in

$$\{f_{31}^{(r)},f_{2}^{(r)},f_{1}^{(r)},d_{1}^{(r)},d_{2}^{(r)},d_{3}^{(r)},e_{1}^{(r)},e_{2}^{(r)},e_{13}^{(r)}\mid r\geq 1\},$$

taken in order some order so that that the f's come before all the d's, which come before all the e's. It is clear from the relations (6.5), (6.6), (6.7) and (6.8) that the monomials in the above elements, where f's come before d's and d's come before e's, with the d's taken in some fixed order, do indeed span  $\widehat{Y}$ .

So our problem is to show that the subalgebra  $\widehat{Y}^+$  of  $\widehat{Y}$  generated by elements  $\{e_i^{(r)}\}_{i=1,2}$  is spanned by the monomials in  $\{e_1^{(r)}, e_2^{(r)}, e_{13}^{(r)}; r \geq 1\}$  taken in some fixed order, and similarly that the subalgebra  $\widehat{Y}^-$  generated by elements  $\{f_i^{(r)}\}_{i=1,2}$  is spanned by the monomials in  $\{f_{31}^{(r)}, f_2^{(r)}, f_1^{(r)}; r \geq 1\}$  taken in some fixed order. Consider  $\widehat{Y}^+$ . Define a filtration

$$L_0\widehat{Y}^+ \subseteq L_1\widehat{Y}^+ \subseteq \cdots$$

on  $\widehat{Y}^+$  by setting the degree of  $e_i^{(r)}$  equal to (r-1). Let  $gr^L\widehat{Y}^+$  be the associated graded algebra, and let  $\overline{e_i}^{(r)} := gr_{r-1}^L e_i^{(r)} \in gr^L \widehat{Y}^+$  for each i=1,2,13. Then we have the following:

$$\begin{split} & [\overline{e}_{1}^{(r)}, \overline{e}_{1}^{(s)}] &= 0, \quad [\overline{e}_{2}^{(r)}, \overline{e}_{2}^{(s)}] = 0, \\ & [\overline{e}_{13}^{(r)}, \overline{e}_{1}^{(s)}] &= 0, \quad [\overline{e}_{13}^{(r)}, \overline{e}_{2}^{(s)}] = 0, \\ & [\overline{e}_{13}^{(r)}, \overline{e}_{13}^{(s)}] &= 0, \quad [\overline{e}_{1}^{(r)}, \overline{e}_{2}^{(s)}] = \overline{e}_{13}^{(r+s-1)}. \end{split}$$

Indeed, the first two identities are clear by the relations in the remark above. For the next two, first note that

$$[\overline{e}_1^{(r+1)}, \overline{e}_2^{(s)}] = [\overline{e}_1^{(r)}, e_2^{(s+1)}].$$
 (6.13)

Then

$$\begin{split} [\overline{e}_{13}^{(r)}, \overline{e}_{12}^{(s)}] &= [[\overline{e}_{12}^{(r)}, e_{23}^{(1)}], e_{12}^{(s)}] = [[\overline{e}_{12}^{(1)}, e_{23}^{(r)}], e_{12}^{(s)}] \\ &= -[[e_{23}^{(r)}, \overline{e}_{12}^{(1)}], e_{12}^{(s)}] = -[[e_{23}^{(r)}, \overline{e}_{12}^{(s)}], e_{12}^{(1)}] \text{ (by (6.12))} \\ &= -[[e_{23}^{(r+s-1)}, \overline{e}_{12}^{(1)}], e_{12}^{(1)}] \\ &= 0 \text{ (by (6.12))}. \end{split}$$

Similarly,

$$\begin{aligned} [\overline{e}_{13}^{(r)}, \overline{e}_{23}^{(s)}] &= [[\overline{e}_{12}^{(r)}, e_{23}^{(1)}], e_{23}^{(s)}] \\ &= -[[\overline{e}_{12}^{(r)}, e_{23}^{(s)}], e_{23}^{(1)}] \text{ (by (6.11))} \\ &= [[\overline{e}_{12}^{(r+s-1)}, e_{23}^{(1)}], e_{23}^{(1)}] = 0 \end{aligned}$$

The fifth relation is an easy consequence of these and the super-Jacobi identity:

$$[\overline{e}_{13}^{(r)}, \overline{e}_{13}^{(s)}] = [[\overline{e}_{12}^{(r)}, \overline{e}_{23}^{(1)}], \overline{e}_{13}^{(s)}] = [[\overline{e}_{12}^{(r)}, \overline{e}_{13}^{(s)}, \overline{e}_{23}^{(1)}]] + [\overline{e}_{12}^{(r)}, [\overline{e}_{23}^{(1)}, \overline{e}_{13}^{(s)}]] = 0$$

The final relation is just another extended application of (6.13).

Given these calculations, it is clear that the graded algebra  $gr^L\widehat{Y}^+$  is spanned by the set of all ordered monomials in  $\{\overline{e}_{ij}^{(r)}\}_{1\leq i< j\leq 3; r\geq 1}$  taken in some fixed order. Hence  $\widehat{Y}^+$  is itself spanned by the corresponding monomials in  $\{e_{ij}^{(r)}\}_{1\leq i< j\leq 3; r\geq 1}$ . The result for the subalgebra  $Y^-$  is shown similarly.

Now we want to show that the monomials in

$$\{d_i^{(r)}\}_{1 \le i \le 3; r \ge 1} \cup \{e_{ij}^{(r)}, f_{ji}^{(r)}\}_{1 \le i < j \le 3; r \ge 1}$$

taken in some fixed order so that f's come before d's and d's come before e's form a basis for the Yangian  $Y(\mathfrak{gl}_{2|1})$ . By Corollary 3.1, we may identify the associated graded algebra  $\operatorname{gr}_2 Y(\mathfrak{gl}_{m|n})$  with  $U(\mathfrak{gl}_{m|n}[t])$ . By the definition of the quasideterminants, under this identification,  $\operatorname{gr}_2^r d_i^{(r+1)}$ ,  $\operatorname{gr}_2^r e_{ij}^{(r+1)}$ , and  $\operatorname{gr}_2^r f_{ji}^{(r+1)}$  are identified, respectively, with  $E_{ii}(-1)^{\overline{i}}t^r$ ,  $E_{ij}(-1)^{\overline{i}}t^r$ , and  $E_{ji}(-1)^{\overline{j}}t^r$ . Then the result follows from the Poincaré-Birkhoff-Witt theorem for Lie superalgebras.  $\square$ 

## 7 Gauss Decomposition of $Y(\mathfrak{gl}_{m|n})$

**Lemma 7.1.** The following relations hold in the algebra  $Y(\mathfrak{gl}_{m|n})[[u^{-1}, v^{-1}]]$ .

$$[d_i(u), d_j(v)] = 0 \text{ for all } 1 \leq i, j \leq m+n$$

$$(u-v)[d_i(u), e_j(v)] = \begin{cases} (\delta_{ij} - \delta_{i,j+1}) d_i(u)(e_j(v) - e_j(u)), & \text{if } 1 \leq j \leq m-1, \\ (\delta_{ij} + \delta_{i,j+1}) d_i(u)(e_j(v) - e_j(u)), & \text{if } j = m, \\ -(\delta_{ij} - \delta_{i,j+1}) d_i(u)(e_j(v) - e_j(u)), & \text{if } m+1 \leq j \leq m+n-1, \end{cases}$$

$$(u-v)[d_i(u), f_j(v)] = \begin{cases} -(\delta_{ij} - \delta_{i,j+1}) (f_j(v) - f_j(u)) d_i(u), & \text{if } 1 \leq j \leq m-1, \\ -(\delta_{ij} + \delta_{i,j+1}) (f_j(v) - f_j(u)) d_i(u), & \text{if } m+1 \leq j \leq m+n-1, \\ (\delta_{ij} - \delta_{i,j+1}) (f_j(v) - f_j(u)) d_i(u), & \text{if } m+1 \leq j \leq m+n-1, \\ (\delta_{ij} - \delta_{i,j+1}) (f_j(v) - f_j(u)) d_i(u), & \text{if } m+1 \leq j \leq m+n-1, \end{cases}$$

$$(u-v)[e_i(u), f_j(v)] = \begin{cases} \delta_{ij} (d_i(u)^{-1} d_{i+1}(u) - d_i(v)^{-1} d_{i+1}(v)), & \text{if } m \leq j \leq m+n-1, \\ -\delta_{ij} (d_i(u)^{-1} d_{i+1}(u) - d_i(v)^{-1} d_{i+1}(v)), & \text{if } m \leq j \leq m+n-1, \end{cases}$$

$$(u-v)[e_j(u), e_j(v)] = \begin{cases} (e_j(v) - e_j(u))^2, & \text{if } 1 \leq j \leq m-1, \\ 0, & \text{if } j = m, \\ -(e_j(v) - e_j(u))^2, & \text{if } m+1 \leq j \leq m+n-1, \end{cases}$$

$$(u-v)[f_j(u), f_j(v)] = \begin{cases} -(f_j(v) - f_j(u))^2, & \text{if } m+1 \leq j \leq m+n-1, \\ 0, & \text{if } j = m, \\ (f_j(v) - f_j(u))^2, & \text{if } m+1 \leq j \leq m+n-1, \end{cases}$$

$$(u-v)[e_i(u), e_{i+1}(v)] = \begin{cases} -e_{i+1}(v) - e_i(v) e_{i+1}(v) \\ -e_{i+1}(v) + e_i(v) e_{i+1}(v) \\ +e_{i+1}(v) - e_{i+1}(v), & \text{if } m \leq j \leq m+n-2, \end{cases}$$

$$(u-v)[f_i(u), f_{i+1}(v)] = \begin{cases} -f_{i+1}(v) f_i(u) + f_{i+1}(v) f_i(v) \\ +f_{i+2,i}(u) - f_{i+2,i}(v), & \text{if } m \leq j \leq m+n-2, \end{cases}$$

$$(u-v)[f_i(u), f_j(v)] = 0 \text{ for } |i-j| > 1;$$

$$[e_i(u), e_j(v)] = 0 \text{ for } |i-j| > 1.$$

*Proof.* The relations for i, j between 1 and m are an easy consequence of those already found for the Yangians  $Y(\mathfrak{gl}_m)$  in [2] and  $Y(\mathfrak{gl}_{2|1})$  in  $\S(6.1)$ , and the fact that the natural inclusions

 $Y(\mathfrak{gl}_m) \hookrightarrow Y(\mathfrak{gl}_{m|n})$  and  $Y(\mathfrak{gl}_{2|1}) \hookrightarrow Y(\mathfrak{gl}_{m|n})$  are homomorphisms. The remaining relations follow by applying the map  $\zeta_{n|m}$  to the corresponding relations in  $Y(\mathfrak{gl}_{n|m})$ .

**Theorem 3.** The Yangian  $Y(\mathfrak{gl}_{m|n})$  is isomorphic as an associative superalgebra to the algebra with even generators  $d_i^{(r)}$ ,  $d_i^{\prime(r)}$ ,  $f_j^{(r)}$ , and  $e_j^{(r)}$ , (for  $1 \le i \le m+n$ ,  $1 \le j \le m+n-1$ ,  $j \ne m$ ,  $r \ge 1$ ) and odd generators  $e_m^{(r)}$ ,  $f_m^{(r)}$  (where again  $r \ge 1$ ) and the following defining relations:

$$d_{i}^{(0)} = 1;$$

$$\sum_{t=0}^{r} d_{i}^{(t)} d_{i}^{\prime (r-t)} = \delta_{r,0};$$

$$[d_{i}^{(r)}, d_{l}^{(s)}] = 0;$$

$$[d_i^{(r)},e_j^{(s)}] \ = \ \begin{cases} (\delta_{i,j}-\delta_{i,j+1})\sum_{t=0}^{r-1}d_i^{(t)}e_j^{(r+s-1-t)}, & \text{for } 1\leq j\leq m-1, \\ (\delta_{i,j}+\delta_{i,j+1})\sum_{t=0}^{r-1}d_i^{(t)}e_j^{(r+s-1-t)}, & \text{for } j=m, \\ -(\delta_{i,j}-\delta_{i,j+1})\sum_{t=0}^{r-1}d_i^{(t)}e_j^{(r+s-1-t)}, & \text{for } m+1\leq j\leq m+n-1, \\ [d_i^{(r)},f_j^{(r)}] \ = \ \begin{cases} -(\delta_{i,j}-\delta_{i,j+1})\sum_{t=0}^{r-1}f_i^{(r+s-1-t)}d_i^{(t)}, & \text{for } 1\leq j\leq m-1; \\ -(\delta_{i,j}+\delta_{i,j+1})\sum_{t=0}^{r-1}f_j^{(r+s-1-t)}d_i^{(t)}, & \text{for } j=m; \\ (\delta_{i,j}-\delta_{i,j+1})\sum_{t=0}^{r-1}f_j^{(r+s-1-t)}d_i^{(t)}, & \text{for } m+1\leq j\leq m+n-1; \\ [e_j^{(r)},f_k^{(s)}] \ = \ \begin{cases} -\delta_{j,k}\sum_{t=0}^{r+s-1}d_j^{(t)}(d_{j+1}^{(r+s-1-t)}), & \text{for } 1\leq j\leq m-1; \\ +\delta_{j,k}\sum_{t=0}^{r+s-1}d_j^{(t)}(d_{j+1}^{(r+s-1-t)}), & \text{for } n\leq j\leq m+n-1; \\ +\delta_{j,k}\sum_{t=0}^{r+s-1}d_j^{(t)}(d_{j+1}^{(r+s-1-t)}), & \text{for } m\leq j\leq m+n-1; \end{cases} \\ [e_j^{(r)},e_j^{(s)}] \ = \ (-1)^{\overline{j}} \left(\sum_{t=1}^{s-1}e_j^{(t)}e_j^{(r+s-1-t)}-\sum_{t=1}^{r-1}e_j^{(t)}e_j^{(r+s-1-t)}\right), & \text{for } j\neq m; \end{cases} \\ [f_j^{(r)},f_j^{(s)}] \ = \ (-1)^{\overline{j}} \left(\sum_{t=1}^{r-1}f_j^{(t)}f_j^{(r+s-1-t)}-\sum_{t=1}^{s-1}f_j^{(t)}f_j^{(r+s-1-t)}\right), & \text{for } j\neq m; \end{cases} \\ [e_j^{(r)},e_j^{(s)}] \ = \ (-1)^{\overline{j}} \left(\sum_{t=1}^{r-1}f_j^{(t)}f_j^{(r+s-1-t)}-\sum_{t=1}^{s-1}f_j^{(t)}f_j^{(r+s-1-t)}\right), & \text{for } j\neq m; \end{cases} \\ [e_j^{(r)},e_j^{(s)}] \ = \ (-1)^{\overline{j}} \left(\sum_{t=1}^{r-1}f_j^{(r)}f_j^{(r+s-1-t)}-\sum_{t=1}^{s-1}f_j^{(t)}f_j^{(r+s-1-t)}\right), & \text{for } j\neq m; \end{cases} \\ [e_j^{(r)},e_j^{(s)}] \ = \ (-1)^{\overline{j}} \left(\sum_{t=1}^{r-1}f_j^{(t)}f_j^{(r+s-1-t)}-\sum_{t=1}^{s-1}f_j^{(t)}f_j^{(r+s-1-t)}\right), & \text{for } j\neq m; \end{cases} \\ [e_j^{(r)},e_j^{(s)}] \ = \ (-1)^{\overline{j}} \left(\sum_{t=1}^{r-1}f_j^{(t)}f_j^{(r+s-1-t)}-\sum_{t=1}^{s-1}f_j^{(t)}f_j^{(r+s-1-t)}\right), & \text{for } j\neq m; \end{cases} \\ [e_j^{(r)},e_j^{(s)}] \ = \ (-1)^{\overline{j}} \left(\sum_{t=1}^{r-1}f_j^{(t)}f_j^{(r+s-1-t)}-\sum_{t=1}^{s-1}f_j^{(t)}f_j^{(r+s-1-t)}\right), & \text{for } j\neq m; \end{cases} \\ [e_j^{(r)},e_j^{(s)}] \ = \ (-1)^{\overline{j}} \left(\sum_{t=1}^{r-1}f_j^{(t)}f_j^{(r+s-1-t)}-\sum_{t=1}^{s-1}f_j^{(t)}f_j^{(r+s-1-t)}\right), & \text{for } j\neq m; \end{cases}$$

*Proof.* We have already stated in  $\S(4)$  that the Yangian  $Y(\mathfrak{gl}_{m|n})$  is generated by the elements:

 $[[f_i^{(r)}, f_k^{(s)}], f_k^{(t)}] + [[f_i^{(r)}, f_k^{(t)}], f_k^{(s)}] = 0, \text{ if } j \neq k;$ 

$$\left\{ \left. d_i^{(r)}, e_j^{(r)}, f_j^{(r)} \right| 1 \leq i \leq m+n, 1 \leq j \leq m+n-1, r \geq 1 \right\}.$$

Also, by the properties of the map  $\psi_{m-1}: Y(\mathfrak{gl}_{1|n}) \to Y(\mathfrak{gl}_{m|n})$ , the elements of the set

$$\{d_m^{(r)}, \dots, d_{m+n}^{(r)}, e_m^{(r)}, \dots, e_{m+n-1}^{(r)}, f_m^{(r)}, \dots, f_{m+n-1}^{(r)}\}_{r \ge 1}$$

all commute with every element of the set

$$\{d_1^{(s)}, \dots, d_{m-1}^{(s)}, e_1^{(s)}, \dots, e_{m-2}^{(s)}, f_1^{(s)}, \dots, f_{m-2}^{(s)}\}_{s \ge 1},$$

and by the corresponding result in  $Y(\mathfrak{gl}_{n|m})$ , all the elements of the set

$$\{d_1^{(r)}, \dots, d_{m+1}^{(r)}, e_1^{(r)}, \dots, e_m^{(r)}, f_1^{(r)}, \dots, f_m^{(r)}\}_{r \ge 1}$$

commute with every element of the set

$$\{d_{m+2}^{(s)}, \dots, d_{m+n}^{(s)}, e_{m+2}^{(s)}, \dots, e_{m+n-1}^{(s)}, f_{m+2}^{(s)}, \dots, f_{m+n-1}^{(s)}\}_{s \ge 1}.$$

We have a natural inclusion  $\iota: Y(\mathfrak{gl}_m) \hookrightarrow Y(\mathfrak{gl}_{m|n})$ , which sends each generator of the Yangian  $Y(\mathfrak{gl}_m)$  to the generator of the same name in the super-Yangian  $Y(\mathfrak{gl}_{m|n})$ . By the Poincarè-Birkhoff-Witt theorem for super-Yangians, this map is an isomorphism between  $Y(\mathfrak{gl}_m)$  and the subalgebra of  $Y(\mathfrak{gl}_{m|n})$  generated by

$$\left\{ \left. d_i^{(r)}, e_j^{(r)}, f_j^{(r)} \right| 1 \leq i \leq m, 1 \leq j \leq m-1, r \geq 1 \right\}.$$

So we have automatically that this subalgebra is given by the presentation in [2].

Similarly, the map  $\psi_{m-2}$  provides an injective homomorphism from  $Y(\mathfrak{gl}_{2|1})$  into  $Y(\mathfrak{gl}_{m|1})$ , which gives an injective homomorphism  $Y(\mathfrak{gl}_{2|1}) \to Y(\mathfrak{gl}_{m|n})$ , sending  $d_1(u), d_2(u), d_3(u)$  to  $d_{m-1}(u), d_m(u), d_{m+1}(u)$ , and  $e_1(v), e_2(v), f_1(v), f_2(v)$  to  $e_{m-1}(v), e_m(v), f_{m-1}(v), f_m(v)$ , respectively. This gives the exact relations between these elements.

Finally, we may compose the natural inclusion  $\iota': Y(\mathfrak{gl}_{n|1}) \hookrightarrow Y(\mathfrak{gl}_{n|m})$  with the isomorphism  $\zeta_{n|m}: Y(\mathfrak{gl}_{n|m}) \to Y(\mathfrak{gl}_{m|n})$ , so that the Yangian  $Y(\mathfrak{gl}_{n|1})$  is isomorphic to the subalgebra of  $Y(\mathfrak{gl}_{m|n})$  generated by

$$\left\{ \left. d_i^{(s)}, e_j^{(s)}, f_j^{(s)} \right| m \leq i \leq m+n, m \leq j \leq m+n-1, s \geq 1 \right\},$$

with exactly the required defining relations.

## 8 The Centre of $Y(\mathfrak{gl}_{m|n})$

The quantum Berezinian was defined by Nazarov [15] as the following power series with coefficients in the Yangian  $Y(\mathfrak{gl}_{m|n})$ :

$$b_{m|n}(u) := \sum_{\tau \in S_m} \operatorname{sgn}(\tau) t_{\tau(1)1}(u) t_{\tau(2)2}(u-1) \cdots t_{\tau(m)m}(u-m+1)$$

$$\times \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) t'_{m+1,m+\sigma(1)}(u-m+1) \cdots t'_{m+n,m+\sigma(n)}(u-m+n)$$
(8.1)

Recall from [9] that we may also write the quantum Berezinian in the following form.

$$b_{m|n}(u) = d_1(u) d_2(u-1) \cdots d_m(u-m+1)$$

$$\times d_{m+1}(u-m+1)^{-1} \cdots d_{m+n}(u-m+n)^{-1}.$$
(8.2)

We shall prove that the coefficients of this formal power series generate the centre of the Yangian. This was a conjecture by Nazarov [15].


**Lemma 8.1.** Let  $\mathfrak{gl}_{m|n}[x]$  be the polynomial current algebra and  $Z = E_{11} + \ldots + E_{m+n,m+n}$ . The centre of  $U(\mathfrak{gl}_{m|n}[x])$  is generated by  $Z, Zx, Zx^2, \ldots$ 

Proof. When  $m \neq n$ , the proof of this result is the same as for Lemma 7.1 in [2], except that one needs to use the supersymmetrization map instead of the symmetrization map. This gives an isomorphism between the  $\mathfrak{gl}_{m|n}[x]$ -modules  $U(\mathfrak{gl}_{m|n}[x])$  and  $S(\mathfrak{gl}_{m|n}[x])$ , where  $S(\mathfrak{gl}_{m|n}[x])$  denotes the supersymmetric algebra of  $\mathfrak{gl}_{m|n}[x]$ . The natural action of  $\mathfrak{gl}_{m|n}[x]$  on  $S(\mathfrak{gl}_{m|n}[x])$  is obtained by extending the adjoint action. The proof relies on the fact that  $\mathfrak{sl}_{m|n}$  forms a  $\mathfrak{gl}_{m|n}$ -invariant complement to the centre  $\mathbb{C}Z$  of  $\mathfrak{gl}_{m|n}$ . In the case where m=n this is no longer true, but we have the following proof.

We begin by constructing a convenient basis for  $S(\mathfrak{gl}_{n|n}[x])$ . We consider a copy G of the dihedral group  $D_{2n}$  in the symmetric group  $S_{2n}$ . Recall that  $D_{2n}$  is the symmetry group of a regular n-sided polygon. Here we label the vertices of the regular n-gon by  $1, 2, \ldots, n$  on one side and by  $n+1, n+2, \ldots, 2n$  on the other side, so that 1 and n+1 sit back to back, as do 2 and n+2, 3 and n+3, etc. Then the rotations act on the letters  $1, 2, \ldots, n$  and  $n+1, n+2, \ldots, 2n$  in the same way, and the reflections flip the polygon over and thereby swap the sets  $1, 2, \ldots, n$  and  $n+1, n+2, \ldots, 2n$  completely. This subgroup G is generated by the permutations

$$\rho = (1, 2, \dots, n)(n+1, n+2, \dots, 2n);$$
 and  $\sigma = (1, 2n)(2, 2n-1)\cdots(n, n+1).$ 

For example, in the case when n = 5, the action of  $\sigma = (1, 10)(2, 9)(3, 8)(4, 7)(5, 6)$  is demonstrated in the diagram below.



For any  $\tau \in G$ , if we know the value of  $\tau(i)$  for any  $i \in \{1, 2, \dots, 2n\}$  then we know exactly which permutation  $\tau \in G$  must be. So the set  $\{E_{i,\tau(i)} \mid i=1,\dots,2n\,;\tau \in G\}$  is a basis for  $\mathfrak{gl}_{n|n}$ . In fact,  $E_{i,\tau(i)}$  is even if  $\tau=\rho^k$ , and odd if  $\tau=\rho^k\circ\sigma$ , for all i,k. So this is an homogeneous basis and the elements  $E_{i,\tau(i)}x^r$ ,  $1\leq i\leq 2n; r\geq 0$ ,  $\tau\in G$  form a basis for  $\mathfrak{gl}_{n|n}[x]$ , on which we may choose some order. A basis for the supercommutative algebra  $S(\mathfrak{gl}_{n|n}[x])$  is formed by the ordered monomials in  $(E_{i,\tau(i)}x^r)^d$  where  $d=0,1,2,\dots$  if  $\tau=\rho^j$  for some j and d=0,1 if  $\tau=\rho^j\circ\sigma$  for some j. In the following, we write  $\pi_j:=\rho^j\circ\sigma$  for  $j=0,1,\dots,n-1$ .

Suppose  $Y \in S(\mathfrak{gl}_{n|n})$  is a  $\mathfrak{gl}_{n|n}$ -invariant element and m is the maximal integer such that the element  $E_{i,\pi_j(i)}x^m$  occurs in Y for some  $j \in \{0,1,\ldots,n-1\}$  and some  $i \in \{1,\ldots,2n\}$ . Then

Y must be of the form:

$$Y = \sum_{d} Y_d \prod_{i,j} \left( E_{i,\pi_j(i)} x^m \right)^{d_j(i)},$$

where we sum over 2n(n-1)-tuples  $d = (d_j(i))_{0 \le j \le n-1; 1 \le i \le 2n}$  with  $d_j(i) = 0, 1$ ; and each  $Y_d$  is a polynomial in the variables  $E_{i,\rho^j(i)}x^r$ , with  $r \ge 0$ , and  $E_{i,\pi_j(i)}x^s$  with s < m. By assumption, we must have

$$[wx, Y] = 0, (8.3)$$

where  $w = \sum_{i=1}^{2n} (-1)^{\bar{i}} E_{i,i}$ . We calculate the sum of terms containing  $E_{k,\pi_l(k)} x^{m+1}$  in the left-hand-side of (8.3) for some fixed k, l. Now,

$$[wx, E_{i,\pi_j(i)}x^m] = 2(-1)^{\bar{i}}E_{i,\pi_j(i)}x^{m+1}.$$

So we find the coefficient of  $E_{k,\pi_l(k)}x^{m+1}$ , where we have shifted everything else to the left, is

$$0 = 2(-1)^{\overline{k} + \alpha_d} d_k(l) \sum_d Y_d \prod_{(i,j) \neq (k,l)} (E_{i,\pi_j(i)} x^m)^{d_i(j)}$$

for some  $\alpha_d=0,1$ . But the monomials  $\prod (E_{i,\pi_j(i)}x^m)^{d_i(j)}$  with different  $d_i(j)$  are linearly independent, which implies  $Y_d=0$  for all d such that  $d_k^{(l)}\neq 0$ . In other words,  $E_{k,\pi_l(k)}x^m$  does not occur in Y. So Y is a polynomial in even elements  $E_{i,\rho^j(i)}x^r$ . Then, since the action of  $\mathfrak{gl}_{n|n}[x]$  on even elements  $E_{i,\sigma_j(i)}x^r$  is the same as the action of  $\mathfrak{gl}_{2n}[x]$ , we have by Proposition 2.12 of [14] (or Lemma 7.2 of [2]) that Y is contained in the subalgebra of  $S(\mathfrak{gl}_{n|n})$  generated by the set  $\{Zx^r, r=0,1,2,\ldots\}$ .

**Theorem 4.** The coefficients of the quantum Berezinian generate the centre of  $Y(\mathfrak{gl}_{m|n})$ .

*Proof.* Write

$$b_{m|n}(u) = 1 + \sum_{k>1} c_k u^{-k}.$$

We showed in [9] that the coefficients  $c_k$  are central. It remains to show that they generate the centre. Our proof is based on that of Theorem 2.13. in [14].

Recall from Corollary 3.1 that the graded algebra  $\operatorname{gr}_2 Y(\mathfrak{gl}_{m|n})$  defined in Section (2) is isomorphic to  $U(\mathfrak{gl}_{m|n}[x])$ . We show that for any  $k=1,2,\ldots$ , the coefficient  $c_k$  has degree k-1 with respect to  $\deg_2(.)$  and that its image in the (k-1)th component of  $\operatorname{gr}_2 Y(\mathfrak{gl}_{m|n})$  coincides with  $Zx^{k-1}$ . Indeed, if we expand out the expression (8.2) for the quantum Berezinian, using the fact from [8] that

$$d_j(u) = t_{jj}(u) - \sum_{k,l \neq j} t_{jk}(u) (|T(u)^{jj}|_{lk})^{-1} t_{lj}(u),$$

we find

$$c_k = \sum_{l_1 + l_2 + \dots + l_{m+n} = k} t_{11}^{(l_1)} t_{22}^{(l_2)} \cdots t_{mm}^{(l_m)} \cdot \left( -t_{m+1,m+1}^{(l_{m+1})} \right) \cdots \left( -t_{m+n,m+n}^{(l_{m+n})} \right) + \text{ terms of lower degree.}$$

Then it is clear that the terms with  $l_i = k$  for some i = 1, ..., m + n have degree k - 1, and all else have lower degree. Then

$$c_k = t_{11}^{(k)} + \ldots + t_{mm}^{(k)} - t_{m+1,m+1}^{(k)} - \ldots - t_{m+n,m+n}^{(k)} + \text{ terms of lower degree.}$$

The result follows when we evaluate the image of the graded part of this under the isomorphism in Corollary (3.1).

## 9 A Subalgebra of the Yangian

Take  $\mu_f$  as defined as in [14]. In other words, for a formal power series

$$f = 1 + f_1 u^{-1} + f_2 u^{-2} + \dots \in \mathbb{C}[[u^{-1}]],$$

the map  $\mu_f$  is the automorphism of  $Y(\mathfrak{gl}_{m|n})$  given by

$$\mu_f: T(u) \mapsto f(u)T(u).$$

Then define a subalgebra of  $Y(\mathfrak{gl}_{m|n})$  as follows:

$$SY_{m|n} := \{ y \in Y(\mathfrak{gl}_{m|n}) \mid \mu_f(y) = y \text{ for all } f \}.$$

**Proposition 9.1.** Let  $Z_{m|n}$  denote the centre of the Yangian  $Y(\mathfrak{gl}_{m|n})$ . Then for  $m \neq n$ , we have

$$Y(\mathfrak{gl}_{m|n}) \cong Z_{m|n} \otimes SY_{m|n}.$$

*Proof.* We assume that m > n. (The result for n < m follows from this by the application of the map  $\zeta$ ). The proof of this result is very similar to that of Proposition 2.16 in [14]. We use the fact, stated there, that for any commutative associative algebra  $\mathcal{A}$  and any formal series,

$$a(u) = 1 + a_1 u^{-1} + a_2 u^{-2} + \dots \in \mathcal{A}[[u^{-1}]],$$

and any positive integer K there exists a unique series

$$\tilde{a}(u) = 1 + \tilde{a}_1 u^{-1} + \tilde{a}_2 u^{-2} + \ldots \in \mathcal{A}[[u^{-1}]]$$

such that

$$a(u) = \tilde{a}(u)\tilde{a}(u-1)\cdots\tilde{a}(u-K+1). \tag{9.1}$$

We take  $a(u) = b_{m|n}(u)$  and K = m-n in the commutative subalgebra  $Y^0 \subset Y(\mathfrak{gl}_{m|n})$  generated by the elements  $d_i^{(r)}$  for  $i = 1, \ldots, m+n$  and  $r \geq 1$ . Write

$$b_{m|n}(u) = \tilde{b}(u)\tilde{b}(u-1)\cdots\tilde{b}(u-m+n+1).$$

By the definition of the map  $\mu_f$  we have that

$$\mu_f(b_{m|n}(u)) = f(u)f(u-1)\cdots f(u-m+n+1)b_{m|n}(u).$$

It follows from the uniqueness of the expansion (9.1) that  $\mu_f(\tilde{b}(u)) = f(u)\tilde{b}(u)$  for all f. Also, the coefficients  $\tilde{b}_k$ ,  $(k \ge 1)$  of the series  $\tilde{b}(u)$  generate the centre  $Z_{m|n}$  since we may recover the coefficients of the series  $b_{m|n}(u)$  from them. The remaining parts of the proof are exactly the same as in [14].

**Lemma 9.2.** For any  $m, n \geq 0$ , the coefficients of the series

$$d_1(u)^{-1}d_{i+1}(u), e_i(u), f_i(u), \quad \text{for } 1 \le i \le m+n-1,$$
 (9.2)

generate the subalgebra  $SY_{m|n}$ .

Proof. It is clear that the coefficients of the series  $d_1(u)$  together with those of the series listed above generate the Yangian  $Y(\mathfrak{gl}_{m|n})$ . Also, for any f, the map  $\mu_f$  leaves the coefficients of the series in (9.2) fixed and maps  $\mu_f(d_1(u)) = f(u)d_1(u)$ . By the Poincaré-Birkhoff-Witt theorem, any element P of  $Y(\mathfrak{gl}_{m|n})$  is a polynomial in  $d_1^{(1)}, d_1^{(2)}, d_1^{(3)}, \ldots$  and the other generators that are fixed by  $\mu_f$  for all f. We can assume further that in each monomial in P the generators are ordered so that the  $f_i^{(r)}$ 's come before the  $d_i^{(r)}$ 's, which come before the  $e_i^{(r)}$ 's. Suppose that  $P \in SY_{m|n}$  and that R is the maximum r such that  $d_1^{(r)}$  occurs in P, and K is the maximum power of  $d_1^{(r)}$  occurring in P for any r. Fix  $f = 1 + \lambda u^{-R}$ , where  $\lambda$  is an arbitrary nonzero complex number. Then we can write:

$$P = \sum_{a_1, a_2, \dots, a_R} F_a D_a \left( d_1^{(1)} \right)^{a_1} \left( d_1^{(2)} \right)^{a_2} \dots \left( d_1^{(R)} \right)^{a_R} E_a,$$

where  $F_a, D_a$  and  $E_a$  are monomials in the generators fixed by  $\mu_f$ , and we sum over all R-tuples  $a = (a_1, a_2, \ldots, a_R)$  of positive integers not exceeding K. Then

$$\mu_f(P) = \sum_{a_1, a_2, \dots, a_R} F_a D_a \left( d_1^{(1)} \right)^{a_1} \left( d_1^{(2)} \right)^{a_2} \dots \left( d_1^{(R)} + \lambda \right)^{a_R} E_a = P.$$

By the linear independence of the different monomials and the fact that  $\lambda$  is an arbitrary complex number, we see that in fact  $d_1^{(R)}$  cannot occur in  $P \in SY_{m|n}$ .

## 10 The Yangian $Y(\mathfrak{sl}_{m|n})$

In this final section we recall the definition of the Yangian  $Y(\mathfrak{sl}_{m|n})$  given by Stukopin [17, 18] and show that this algebra is isomorphic to the subalgebra  $SY_{m|n}$  of  $Y(\mathfrak{gl}_{m|n})$ . Stukopin names his Yangian after the series of classical Lie superalgebras A(m-1,n-1) and defines it only for the case  $m \neq n$ , since in the case where m = n the Lie superalgebra  $\mathfrak{sl}_{m|n}[x]$  does not have a canonical Lie bi-superalgebra structure. It seems natural to adopt the resulting presentation as a definition for  $Y(\mathfrak{sl}_{m|n})$  even in the case m = n (see Proposition 10.1 below).

Recall (from [13]) that

$$A(m-1, n-1) = \mathfrak{sl}_{m|n} \quad \text{for } m \neq n; \quad m, n \ge 0; \tag{10.1}$$

$$A(n-1, n-1) = \mathfrak{sl}_{n|n}/\langle Z \rangle, \text{ for } n > 0,$$
(10.2)

where  $\langle Z \rangle$  is the one-dimensional ideal consisting of scalar matrices  $\lambda Z$ ,  $(\lambda \in \mathbb{C})$ . The Lie algebra A(m,n) is the contragredient Lie superalgebra with Cartan matrix  $A=(a_{ij})_{i,j=1}^{m+n-1}$ , where  $a_{ii}=2$  for all i < m;  $a_{mm}=0$ ;  $a_{ii}=-2$  for all i > m;  $a_{i+1,i}=a_{i,i+1}=-1$  for all i < m; and all other entries are 0.

The special linear Lie superalgebra  $\mathfrak{sl}_{m|n}$  may be defined explicitly by the following presentation [16, 10]. We take generators  $\{h_i, x_j^+, x_j^- | 1 \le i \le m+n-1\}$ . The generators  $h_i, x_j^\pm$  are declared even for all i and all  $j \ne m$ ; the generators  $x_m^\pm$  are declared odd. The defining relations are:

$$\begin{array}{rcl} [h_i,h_j] & = & 0; \\ [x_i^+,x_j^-] & = & \delta_{i,j}h_i; \\ [h_i,x_j^\pm] & = & \pm a_{ij}x_j^\pm; \\ [x_i^\pm,[x_i^\pm,x_j^\pm]] & = & 0, \quad i \neq j; \\ [x_m^\pm,x_m^\pm] & = & 0, \\ [[x_{m-1}^\pm,x_m^\pm],[x_{m+1}^\pm,x_m^\pm]] & = & 0, \end{array}$$

for all possible i, j. Stukopin derives the following relations for the Yangian  $Y(\mathfrak{sl}_{m|n})$  according to the definition of Yangian given in [7], as the quantization of the Lie bi-superalgebra  $\mathfrak{sl}_{m|n}[t]$ .

**Definition 10.1.** The Yangian  $Y(\mathfrak{sl}_{m|n})$  is the associative superalgebra over  $\mathbb{C}$  defined by the generators  $x_{i,s}^{\pm}$  and  $h_{i,s}$  for  $1 \leq i \leq m+n-1$  and  $s \in \mathbb{Z}_+$ , and by the relations

$$\begin{array}{rcl} [h_{i,r},h_{j,s}] & = & 0, \\ [x_{i,r}^+,x_{j,s}^-] & = & \delta_{ij}h_{i,r+s} \;, \\ [h_{i,0},x_{j,s}^\pm] & = & \pm a_{ij}x_{j,s}^\pm \;, \\ [h_{i,r+1},x_{j,s}^\pm] - [h_{i,r},x_{j,s+1}^\pm] & = & \frac{\pm a_{ij}}{2} \; (h_{i,r}\,x_{j,s}^\pm + x_{j,s}^\pm \, h_{i,r}), \; for \; i,j \; not \; both \; m, \\ [h_{m,r+1},x_{m,s}^\pm] & = & 0, \\ [x_{i,r+1}^\pm,x_{j,s}^\pm] - [x_{i,r}^\pm,x_{j,s+1}^\pm] & = & \frac{\pm a_{ij}}{2} (x_{i,r}^\pm x_{j,s}^\pm + x_{j,s}^\pm x_{i,r}^\pm), \; for \; i,j \; not \; both \; m, \\ [x_{m,r+1}^\pm,x_{m,s}^\pm] & = & 0, \\ [x_{i,r}^\pm,[x_{i,s}^\pm,x_{j,t}^\pm]] + [x_{i,s}^\pm,[x_{i,r}^\pm,x_{j,t}^\pm]] & = & 0, \; (i \neq j), \\ [[x_{m-1,q}^\pm,x_{m,r}^\pm],[x_{m+1,s}^\pm,x_{m,t}^\pm]] & = & 0, \end{array}$$

where q, r, s and t are arbitrary positive integers and  $a_{ij}$  are the elements of the Cartan matrix above. The generators  $x_{m,s}^{\pm}$  are odd and all other generators are even.

The Yangian  $Y(\mathfrak{sl}_{m|n})$  when  $m \neq n$  is a Hopf algebra (see [18]).

Proposition 10.1. We may define an associative algebra isomorphism

$$\alpha: Y(\mathfrak{sl}_{m|n}) \to SY_{m|n} \subset Y(\mathfrak{gl}_{m|n})$$

by:

$$1 + \sum_{s \ge 0} h_{i,s} u^{-s-1} := h_i(u) \quad \mapsto \quad \left\{ \begin{array}{l} d_i(u + \frac{m-i}{2})^{-1} d_{i+1}(u + \frac{m-i}{2}), & \text{for } i \le m, \\ d_i(u + \frac{i-m}{2})^{-1} d_{i+1}(u + \frac{i-m}{2}), & \text{for } i > m, \end{array} \right.$$

$$\sum_{s \ge 0} x_{i,s}^+ u^{-s-1} =: x_i^+(u) \quad \mapsto \quad \left\{ \begin{array}{l} f_i(u + \frac{m-i}{2}), & \text{for } i \le m, \\ f_i(u + \frac{i-m}{2}), & \text{for } i > m, \end{array} \right.$$

$$\sum_{s \ge 0} x_{i,s}^- u^{-s-1} := x_i^-(u) \quad \mapsto \quad \left\{ \begin{array}{l} e_i(u + \frac{m-i}{2}), & \text{for } i \le m, \\ -e_i(u + \frac{i-m}{2}), & \text{for } i > m. \end{array} \right.$$

for  $1 \le i \le m + n - 1$ .

*Proof.* Straightforward calculations show this map is a homomorphism. It is clearly injective, with image contained in  $SY_{m|n}$ . In order to show that the image is precisely  $SY_{m|n}$  we explain how to recover each power series in (9.2) as the image of some series in  $Y(\mathfrak{sl}_{m|n})$ . For the series  $e_i(v)$ ,  $f_i(v)$  this is easy. On the other hand,  $d_1(u)^{-1}d_i(u)$  is just the image under  $\alpha$  of

$$h_1(u-\frac{m-1}{2})h_2(u-\frac{m-2}{2})\cdots h_i(u-\frac{m-i}{2}).$$

The restriction of the map  $\zeta_{m|n}: Y(\mathfrak{gl}_{m|n}) \to Y(\mathfrak{gl}_{n|m})$  to the subalgebra  $Y(\mathfrak{sl}_{m|n})$  gives a nice map between the smaller Yangians:

Corollary 10.2. We may define an isomorphism  $\zeta_{m|n}: Y(\mathfrak{sl}_{m|n}) \to Y(\mathfrak{sl}_{n|m})$  as follows:

$$\zeta_{m|n}: \begin{cases}
h_{i,s} & \mapsto & h_{m+n-i,s}; \\
(-1)^{\overline{i}} x_{i,s}^{+} & \mapsto & -x_{m+n-i,s}^{-}; \\
(-1)^{\overline{i}} x_{i,s}^{-} & \mapsto & x_{m+n-i,s}^{+};
\end{cases}$$

for all  $1 \le i \le m+n-j$  and all  $s \ge 0$ .

#### Acknowledgements

I would like to thank my PhD supervisor Alex Molev for his patient assistance. He made many suggestions, both mathematical and stylistic, for improving this paper.

#### References

- [1] Changrim Ahn and Wai Ming Koo. gl(n|m) color Calogero-Sutherland models and super Yangian algebra. *Phys. Lett. B*, 365(1-4):105–112, 1996. hep-th/9505060.
- [2] Jonathan Brundan and Alexander Kleshchev. Parabolic presentations of the Yangian  $Y(\mathfrak{gl}_n)$ . Comm. Math. Phys., 254(1):191–220, 2005. math.QA/0407011.
- [3] Jin-fang Cai, Guo-xing Ju, Ke Wu, and Shi-kun Wang. Super Yangian double DY(gl(1|1)) and its Gauss decomposition. J. Phys. A, 30(11):L347–L350, 1997. hep-th/9612016.
- [4] V. Caudrelier and E. Ragoucy. Lax pair and super-Yangian symmetry of the nonlinear super-Schrödinger equation. *J. Math. Phys.*, 44(12):5706–5732, 2003. math.QA/0306115.
- [5] Vyjayanthi Chari and Andrew Pressley. A guide to quantum groups. Cambridge University Press, Cambridge, 1994.
- [6] N. Crampé. Hopf structure of the Yangian  $Y(sl_n)$  in the Drinfel'd realization. J. Math. Phys., 45(1):434-447, 2004. math.QA/0304254.

- [7] V. G. Drinfel'd. Quantum groups. In *Proceedings of the International Congress of Mathematicians*, Vol. 1, 2 (Berkeley, Calif., 1986), pages 798–820, Providence, RI, 1987. Amer. Math. Soc.
- [8] Israel Gelfand, Sergei Gelfand, Vladimir Retakh, and Robert Lee Wilson. Quasideterminants. Adv. Math., 193(1):56–141, 2005. math.QA/0208146.
- [9] Lucy Gow. On the Yangian  $Y(\mathfrak{gl}_{m|n})$  and its quantum Berezinian. Czechoslovak Journal of Physics, 55(11):1415–1420, 2005. math.QA/0501041.
- [10] Pavel Grozman and Dimitry Leites. Defining relations for Lie superalgebras with Cartan matrix. Czechoslovak J. Phys., 51(1):1–21, 2001. math.RT/0202152.
- [11] Machiko Hatsuda and Kentaroh Yoshida. Classical integrability and super Yangian of superstring on  $AdS_5 \times S^5$ . Adv. Theor. Math. Phys., 9(5):703–728, 2005. hep-th/0407044.
- [12] Guo-xing Ju, Shi-kun Wang, and Ke Wu. The algebraic structure of the gl(n|m) color Calogero-Sutherland models. J. Math. Phys., 39(5):2813–2820, 1998. q-alg/9710005.
- [13] V. G. Kac. Lie superalgebras. Advances in Math., 26(1):8–96, 1977.
- [14] A. Molev, M. Nazarov, and G. Olshanskii. Yangians and classical Lie algebras. Russian Math. Surveys, 51:205–282, 1996. hep-th/9409025.
- [15] M. L. Nazarov. Quantum Berezinian and the classical Capelli identity. *Lett. Math. Phys.*, 21(2):123–131, 1991.
- [16] M. Scheunert. The presentation and q deformation of special linear Lie superalgebras. J. Math. Phys., 34(8):3780–3808, 1993.
- [17] V. A. Stukopin. Yangians of Lie superalgebras of type A(m,n). Functional Analysis and its Applications, 28(3):217–219, 1994.
- [18] Vladimir Stukopin. Yangians of classical lie superalgebras: Basic constructions, quantum double and universal R-matrix. *Proceedings of the Institute of Mathematics of NAS of Ukraine*, 50(3):1195–1201, 2004.
- [19] R. B. Zhang. The gl(M|N) super Yangian and its finite-dimensional representations. Lett. Math. Phys., 37(4):419-434, 1996. q-alg/9507029.