

Bahadur-Kiefer theory for sample quantiles of weakly dependent linear processes

Rafał Kulik
University of Sydney
School of Mathematics and Statistics
F07, University of Sydney
NSW 2006, Australia
and
Mathematical Institute
University of Wrocław
Pl. Grunwaldzki 2/4
50-384 Wrocław
Poland
rkuli@math.uni.wroc.pl

March 2, 2019

Abstract

In this paper we establish the Bahadur-Kiefer representation for sample quantiles for a class of weakly dependent linear processes. The rate of approximation is the same as for i.i.d. sequences and thus it is optimal. Our results extend recent results by Wu, [24]. Also, we apply the Bahadur-Kiefer type result to strong asymptotics for Vervaat error processes.

Keywords: sample quantiles, linear processes, Bahadur representation, strong approximation, law of the iterated logarithms, empirical processes, Vervaat error process

Running title: Sample quantiles and weak dependence

1 Introduction

Let $\{X_i, i \geq 1\}$ be a stationary sequence with a marginal distribution function $F(x) = P(X_1 \leq x)$ and a density $f = F'$. Given the sample X_1, \dots, X_n ,

let $F_n(x) = \frac{1}{n} \sum_{i=1}^n 1_{\{X_i \leq x\}}$ be its corresponding empirical distribution function. Let $X_{1:n} \leq \dots \leq X_{n:n}$ be the order statistics. The empirical quantile function $Q_n(y)$, $y \in (0, 1)$ is defined as $Q_n(y) = X_{k:n}$ if $\frac{k-1}{n} < p \leq \frac{k}{n}$. Further, let

$$\begin{aligned}\beta_n(x) &= n^{1/2}(F_n(x) - F(x)), \quad x \in \mathbb{R}, \\ q_n(y) &= n^{1/2}(Q(y) - Q_n(y)), \quad y \in (0, 1),\end{aligned}$$

be the *general empirical* and the *general quantile* processes, respectively. IF F is continuous, define $U_i = F(X_i)$, $i \geq 1$. Let $E_n(x) = \frac{1}{n} \sum_{i=1}^n 1_{\{U_i \leq x\}} = F_n(Q(x))$ and $U_n(y)$ be the corresponding uniform empirical distribution and uniform empirical quantile functions. Let

$$\begin{aligned}\alpha_n(x) &= n^{1/2}(E_n(x) - x), \quad x \in (0, 1), \\ u_n(y) &= n^{1/2}(y - U_n(y)), \quad y \in (0, 1),\end{aligned}$$

be the corresponding *uniform empirical* and *uniform quantile* processes.

Our main purpose of this paper is to obtain Bahadur-Kiefer type representations for sample quantiles of weakly dependent random variables with the *optimal* rate. Assume for a while that X_i , $i \geq 1$ are i.i.d. Fix $y \in (0, 1)$. Let I_p be a neighborhood of $Q(y)$. Assuming that $\inf_{x \in I_p} f(x) > 0$ and $\sup_{x \in I_p} |f'(x)| < \infty$ Bahadur in [3] obtained the following *Bahadur representation*

$$f(Q(y))q_n(y) - \alpha_n(y) =: R_n(y), \quad (1)$$

where

$$R_n(y) = O_{a.s.}(n^{-1/4}(\log n)^{1/2}(\log \log n)^{1/4}). \quad (2)$$

The process $\{R_n(y), y \in (0, 1)\}$ is called the Bahadur-Kiefer process. Later, Kiefer proved in [18] that (2) can be strengthened to

$$R_n(y) = O_{a.s.}(n^{-1/4}(\log \log n)^{3/4}), \quad (3)$$

which is the optimal rate. Continuing his study, Kiefer obtained in [19] the uniform version of (1), referred later as the *Bahadur-Kiefer representation*:

$$\sup_{y \in (0, 1)} |f(Q(y))q_n(y) - \alpha_n(y)| =: R_n \quad (4)$$

and

$$R_n = O_{a.s.}(n^{-1/4}(\log n)^{1/2}(\log \log n)^{1/4}). \quad (5)$$

Once again, the above rate is optimal. Kiefer obtained his result assuming

$$(K1) \sup_{y \in (0,1)} |F''(Q(y))| < \infty,$$

$$(K2) \inf_{y \in (0,1)} f(Q(y)) > 0.$$

We shall refer to (K1)-(K2) as to the *Kiefer conditions*. Additionally, he assumed that a support of F is finite.

Further, Csörgő and Révész [8] obtained Kiefer's result (4) under the following, weaker conditions, which shall be referred later as the *Csörgő-Révész conditions*:

$$(CsR1) \ f' \text{ exists on } (a, b), \text{ where } a = \sup\{x : F(x) = 0\}, \ b = \inf\{x : F(x) = 1\}, \ -\infty \leq a < b \leq \infty,$$

$$(CsR2) \ \inf_{x \in (a,b)} f(x) > 0,$$

$$(CsR3) \ \sup_{y \in (0,1)} y(1-y) \left| \frac{f'(Q(y))}{f^2(Q(y))} \right| \leq \gamma \text{ with some } \gamma > 0,$$

$$(CsR4) \ f \text{ is nondecreasing (nonincreasing) on the interval to the right of } a \text{ (to the left of } b).$$

Precisely, they showed (among others) that under (CsR1), (CsR2), (CsR3) and with $\delta_n = n^{-1} \log \log n$,

$$\sup_{\delta_n \leq y \leq 1 - \delta_n} |f(Q(y))q_n(y) - u_n(y)| = O_{a.s.}(n^{-1/2} \log \log n). \quad (6)$$

Additionally, if (CsR4) holds, then

$$\sup_{0 < p < 1} |f(Q(y))q_n(y) - u_n(y)| = O_{a.s.}(n^{-1/2} \ell(n)), \quad (7)$$

where $\ell(n)$ is slowly varying at infinity. This, via

$$\sup_{y \in (0,1)} |q_n(y) - \alpha_n(y)| = O_{a.s.}(n^{-1/4} (\log n)^{1/2} (\log \log n)^{1/4})$$

yields the Bahadur-Kiefer representation (5) under less restrictive conditions compared to Kiefer's assumptions. In particular, Csörgő-Révész conditions are fulfilled if F is exponential or normal. We note that (CsR3) can be replaced by the following two conditions:

$$(CsR3(i)) \ f(Q(y)) \sim y^{\gamma_1} L_1(y^{-1}) \text{ as } y \downarrow 0;$$

$$(CsR3(ii)) \ f(Q(1-y)) \sim (1-y)^{\gamma_2} L_2((1-y)^{-1}) \text{ as } y \uparrow 1;$$

hold for some numbers $\gamma_1, \gamma_2 > 0$ and some slowly varying functions L_1, L_2 . Constants in (CsR3(i)), (CsR3(ii)) and (CsR3) are related as follows: $\gamma = \max\{\gamma_1, \gamma_2\}$. We refer to [7], [8], [9] and [10] for more discussion.

For weakly dependent random variables some results are available. For mixing random variables the best possible (in terms of the rates for R_n) results are included in [2]. In particular, for a class of ϕ -mixing sequences, under Kiefer conditions, they obtained (4) with the optimal rate (5). Those results were improved in [13] and [26] in terms of less restrictive mixing rates and Csörgő-Révész-type conditions. However, the rate was $R_n = O_{a.s.}((\log n)^{-\lambda})$ with some $\lambda > 0$. This rate is much worse compared to the optimal one in (5), on the other hand, however, sufficient for weak convergence purposes.

However, mixing is rather hard (if possible at all) to verify and requires some additional regularity assumptions. In particular, for linear processes

$$X_i = \sum_{k=0}^{\infty} c_k \epsilon_{i-k}, \quad (8)$$

in order to obtain strong mixing both regularity assumptions on a density of ϵ_1 and some constrains on c_k 's are required (cf. [23] or [12]). If c_k decay exponentially fast, then we are able to establish the strong with geometric rates. However, even in this case we do not attain the optimal rate in the Bahadur-Kiefer representation, see [2].

Another way of looking at linear processes is to approximate the sequence $\{X_i, i \geq 1\}$ by a sequence with finite memory and then use the classical Bernstein blocking technique. This method avoids some assumptions on a density of ϵ_1 , but it leads to some constrains on c_k 's and does not lead to the optimal rates. To be more specific, Hesse [15] obtained the Bahadur representation (1) with rate $R_n(y) = O_{a.s.}(n^{-1/4+\lambda})$ for some $\lambda > 0$.

Generally speaking, the blocking technique or mixing require some strong assumptions on the model (8), not only in the case of the Bahadur-Kiefer representation. To overcome such problems, Ho, Hsing, Mielniczuk and Wu (see [16], [17], [24], [25]) developed martingale based methods, which lead to optimal or almost optimal results, especially in the context of weak convergence. Based on these methods, Wu [24] obtained for a class of linear processes the exact rate (3) in the Bahadur representation (1). Also, he studied the Bahadur-Kiefer representation. However, he was not able to attain the optimal rate (5) due to a lack of an appropriate version of the law of iterated logarithm for empirical processes.

Wu [24] studied also the Bahadur-Kiefer representation for a class of

weakly dependent sequences satisfying a *geometric moment contraction* assumption. However, also in this case his rate is not optimal.

In this paper we shall combine the Bernstein's blocking technique together with Wu's method to obtain the optimal rate in the Bahadur-Kiefer representation (4) under quite mild conditions on c_k 's, as less restrictive as possible. Thus, in this case, both methods complement each other and lead to optimal rates. An appropriate result is stated in Theorem 2.1 and extends directly Kiefer's result (5). The methodology involves the recent strong approximation result of Berkes and Horváth [4]. Since the part of our computations follows exactly their proof, we include that part in the Appendix. Further, we shall obtain the optimal rate under general conditions on F , on the whole interval $(0, 1)$. A typical approach is to start with the distance between uniform empirical and uniform quantile processes (as Kiefer did) and then to obtain the distance between the normalized quantile process $f(Q(y))q_n(y)$ and the uniform quantile process, as in Csörgő and Révész [8]. However, we shall point out that there are some technical problems when dealing with increments of the uniform empirical processes: to exploit fully the linear structure of the model, we must deal with the behavior of the quantile function $Q(y)$ (see the proof of Lemma 3.6). To overcome such problem, we shall consider an appropriately weighted process $f(Q(y))q_n(y) - \alpha_n(y)$ as in Theorem 2.3.

Assuming much stronger conditions on c_k 's and instead of approach of Lemma 3.6 we may use an exponential inequality (see Lemma 3.1) to obtain the optimal rates for the distance between uniform empirical and quantile processes. Further, one could hope to obtain a Csáki-type law of the iterated logarithm and proceed exactly as in the proof in [8]. However, as mentioned before, our focus is on optimal rates together with optimal conditions on coefficients. Thus, we shall not follow this path.

The result of Theorem 2.1 will be applied to establish strong asymptotics of a *Vervaat error* processes. To be more specific, for $t \in (0, 1)$, let $V_n(y) = 2n^{1/2} \int_0^y (u_n(t) - \alpha_n(t)) dt$. In i.i.d. case it is known that $\sup_{y \in (0, 1)} |V_n(y) - \alpha_n^2(y)| \rightarrow 0$ in probability. Thus, in the same way as studying $u_n(y) - \alpha_n(y)$ we may study the process $\{I_n(y) = V_n(y) - \alpha_n^2(y)\}$, which is called the *Vervaat error* process. We shall refer to [6] and references therein for more details. In particular, in the i.i.d. case, one has

$$\sup_{t \in (0, 1)} |I_n(t)| = O_{a.s.}(n^{-1/4}(\log n)^{1/2}(\log \log n)^{3/4}) \quad (9)$$

and the above rate is optimal. We shall show that this result is still valid

for the model (8).

Finally, in Section 6 we shall compare our constraints on parameters of the model with those required when one wants to use other methods.

Summarizing, in order to obtain optimal rates in the Bahadur-Kiefer representation we combined the classical blocking method with the martingale approximation. It seems that this kind of approach should be helpful when one wants to prove almost sure results. As for weak convergence, the martingale approximation works sufficiently well.

In what follows C will denote a generic constant which may be different at each appearance. Also, we write $a_n \sim b_n$ if $\lim_{n \rightarrow \infty} a_n/b_n = 1$. For any stationary sequence $\{Z_i, i \geq 1\}$ of random variables, Z will be a random variable with the same distribution as Z_1 .

Throughout the paper we shall use:

$$b_n = n^{-3/4}(\log n)^{1/2}(\log \log n)^{1/4}$$

and

$$\lambda_n = n^{-1/2}(2 \log \log n)^{1/2}.$$

For any function $h(x)$ defined on \mathbb{R} we write for $x < y$, $h(x, y) := h(y) - h(x)$.

2 Results

Consider the class of linear processes

$$X_i = \sum_{k=0}^{\infty} c_k \epsilon_{i-k},$$

where $\{\epsilon_i, i \in \mathbb{Z}\}$ is an i.i.d. sequence and $\sum_{k=0}^{\infty} |c_k| < \infty$. Assume that for some $\alpha > 2$,

$$\mathbb{E}|\epsilon|^\alpha < \infty. \tag{10}$$

Suppose that for some $\rho \in (0, \frac{1}{2})$,

$$\sum_{k=i}^{\infty} |c_k|^\alpha = O(i^{-\alpha/\rho}(\log i)^{-\frac{3\alpha}{2}}). \tag{11}$$

The first result deals with the Bahadur-Kiefer representation in terms of the distance between the normalized sample quantile and uniform empirical processes.

Theorem 2.1 *Assume that (10) and (11) hold. Furthermore, assume that for f_ϵ , a density of ϵ , one has*

$$\sup_{x \in \mathbf{R}} (f_\epsilon(x) + |f'_\epsilon(x)| + |f''_\epsilon(x)|) < \infty. \quad (12)$$

Then, under Kiefer conditions,

$$\sup_{y \in (0,1)} |f(Q(y))q_n(y) - \alpha_n(y)| = O_{a.s.}(n^{-1/4}(\log n)^{1/2}(\log \log n)^{1/4}). \quad (13)$$

Remark 2.2 *Assume that $c_k \sim k^{-\tau}(\log k)^{-3/2}$.*

$$\sum_{k=i}^{\infty} |c_k|^\alpha = \sum_{k=i}^{\infty} k^{-\alpha\tau(\log k)^{-\frac{3\alpha}{2}}} = O(i^{-\alpha\tau+1}(\log i)^{-\frac{3\alpha}{2}}).$$

and (11) is fulfilled if $\tau > \frac{1}{\rho} + \frac{1}{\alpha}$. Further, if $c_k = O(k^{-\tau})$, $\tau > 2 + \frac{1}{\alpha}$, then we can find $\rho \in (0, \frac{1}{2})$ such that (11) holds. In particular, if X_1 has all moments finite then $\tau > 2$ provides the optimal bound. In such the case in [24], summability assumption $\sum_{k=0}^{\infty} |c_k| < \infty$ provides the bound $O(n^{-3/4}(\log n)^{\frac{1}{2}+\eta})$, $\eta > 0$. Generally speaking, our constrains (11) are more restrictive, but provide the better (optimal) rates compared to [24].

To obtain the bound without Kiefer conditions, note first that if F is arbitrary such that (K1), (K2) hold on (a, b) , then the result of Theorem 2.1 remains valid if one replaces $\sup_{y \in (0,1)}$ with $\sup_{y \in (a,b)}$. In general, we shall consider a weighted distance between the uniform empirical and normalized quantile processes.

Theorem 2.3 *Assume (CsR1), (CsR2), (CsR3(i)) and (CsR(ii)) with $\min\{\gamma_1, \gamma_2\} \geq 1$. As in Theorem 2.1, assume (10), (11) and (12). Let $\mu > 0$ be arbitrary. Then for arbitrary $\nu > \max\{2\gamma + \mu, 3\gamma - 2\}$,*

$$\sup_{y \in (0,1)} (y(1-y))^\nu |f(Q(y))q_n(y) - \alpha_n(y)| = O_{a.s.}(n^{-1/4}(\log n)^{1/2}(\log \log n)^{1/4}). \quad (14)$$

Remark 2.4 Let us discuss the meaning and conditions of Theorem 2.3. To obtain the result on the whole interval we need to put the weight function $h(y) = (y(1-y))^\nu$. Note first that $h(y)$ may be replaced with $f^{2+\mu}(Q(y))$. Thus, in particular, one would obtain for example

$$\limsup_{n \rightarrow \infty} (\log \log n)^{-1/2} f^{3+\mu}(Q(y))q_n(y) = C$$

instead of

$$\limsup_{n \rightarrow \infty} (\log \log n)^{-1/2} f(Q(y)) q_n(y) = C.$$

Moreover, if one wants to apply (14) to construct *direct simultaneous* confidence intervals for the quantile function $Q(y)$ like in [7], one needs to estimate $f^{3+\mu}(Q(y))$ instead of $f(Q(y))$. Thus, for most practical purposes the additional weight function does not increase complexity of a (estimation) problem, but makes proof work. IT should be pointed out that the additional weights required to study the distance between the uniform quantile and uniform empirical processes, not the distance between the normalized quantile and uniform quantile processes.

The restriction $\min\{\gamma_1, \gamma_2\} \geq 1$ is in fact relevant, since essentially the only distributions with either $\gamma_1 < 1$ or $\gamma_2 < 1$ are those with finite support from the left or from the right, respectively, see [20]. Such cases are covered by Theorem 2.1.

Further, it should be noted that (CsR4) is not needed. Therefore, we may apply Theorem 2.3 to densities like $f(x) \sim x^{-b}L(x)$, $b > 1$, L slowly varying. Theorem 3 in [8] cannot be applied for an arbitrary L .

From Theorem 2.1 we may derive the bound on the Vervaat error process.

Corollary 2.5 *Under conditions of Theorem 2.1 we have*

$$\sup_{y \in (0,1)} I_n(y) = O_{a.s.}(n^{-1/4}(\log n)^{1/2}(\log \log n)^{3/4}).$$

3 Proof of Theorem 2.1

Before giving the proof, we need some preliminary notations and results.

With ρ from (11), define

$$\hat{X}_i := \hat{X}_i(\rho) = \sum_{k=0}^{i^\rho-1} c_k \epsilon_{i-k}. \quad (15)$$

Without loss of generality we may assume that $c_0 = 1$. Let $X_{i,i-1} = \sum_{k=1}^{\infty} c_k \epsilon_{i-k}$ and define its truncated version $\hat{X}_{i,i-1} = \sum_{k=1}^{i^\rho-1} c_k \epsilon_{i-k}$.

By the Fatou's lemma, for any $\alpha > 2$,

$$\mathbb{E}|X_i - \hat{X}_i|^\alpha \leq C \mathbb{E} \left| \sum_{k=i^\rho}^{\infty} c_k \epsilon_{i-k} \right|^\alpha \leq C \lim_{m \rightarrow \infty} \mathbb{E} \left| \sum_{k=i^\rho}^m c_k \epsilon_{i-k} \right|^\alpha \leq C \sum_{k=i^\rho}^{\infty} |c_k|^\alpha. \quad (16)$$

This estimate is very well true for $\mathbb{E}|X_{i,i-1} - \hat{X}_{i,i-1}|^\alpha$ as well. Thus, replacing i with i^δ in (11),

$$\left\| \sum_{i=1}^n X_i - \sum_{i=1}^n \hat{X}_i \right\|_\alpha \leq \sum_{i=1}^n \left(\sum_{k=i^\delta}^{\infty} |c_k|^\alpha \right)^{1/\alpha} \leq \sum_{i=1}^{\infty} O(i^{-1}(\log i)^{-3/2}) < \infty \quad (17)$$

and the same estimate is valid for $\left\| \sum_{i=1}^n X_{i,i-1} - \sum_{i=1}^n \hat{X}_{i,i-1} \right\|_\alpha$.

Denote $\mathcal{F}_i = \sigma(\epsilon_i, \epsilon_{i-1}, \dots)$. Note that $F(x) = \mathbb{E}F_\epsilon(x - X_{i,i-1}) = \mathbb{E}F_\epsilon(x - X_{1,0})$ and $\hat{F}_i(x) := P(\hat{X}_i \leq x) = \mathbb{E}F_\epsilon(x - \hat{X}_{i,i-1})$, $i \geq 1$. Define

$$F_n^*(x) = \frac{1}{n} \sum_{i=1}^n F_\epsilon(x - X_{i,i-1}),$$

a conditional empirical distribution function, and

$$\hat{F}_n^*(x) = \frac{1}{n} \sum_{i=1}^n F_\epsilon(x - \hat{X}_{i,i-1}),$$

its corresponding version based on the truncated random variables.

Since f_ϵ exists, then so $f(x) = dF(x)/dx$, $\hat{f}_i(x) = d\hat{F}_i(x)/dx$ and consequently, we may define $f_n^*(x) = dF_n^*(x)/dx$, $\hat{f}_n^*(x) = d\hat{F}_n^*(x)/dx$, $\hat{f}_n(x) = \frac{1}{n} \sum_{i=1}^n d\hat{F}_i(x)/dx$.

Denote also $Y_i(x) = F_\epsilon(x - X_{i,i-1}) - F(x)$, $\hat{Y}_i(x) = F_\epsilon(x - \hat{X}_{i,i-1}) - \hat{F}_i(x)$. Then we have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n (1_{\{X_i \leq x\}} - F(x)) = \\ &= \frac{1}{n} \sum_{i=1}^n \left(1_{\{X_i \leq x\}} - \mathbb{E}(1_{\{X_i \leq x\}} | \mathcal{F}_{i-1}) \right) + \frac{1}{n} \sum_{i=1}^n \left(\mathbb{E}(1_{\{X_i \leq x\}} | \mathcal{F}_{i-1}) - F(x) \right) \\ &= \frac{1}{n} \sum_{i=1}^n \left(1_{\{X_i \leq x\}} - \mathbb{E}(1_{\{X_i \leq x\}} | \mathcal{F}_{i-1}) \right) + \frac{1}{n} \sum_{i=1}^n Y_i(x) =: M_n(x) + N_n(x). \end{aligned}$$

Then $nM_n(x)$, $n \geq 1$, is a martingale. Further, denote

$$\hat{N}_n(x) = \frac{1}{n} \sum_{i=1}^n (F_\epsilon(x - \hat{X}_{i,i-1}) - \hat{F}_i(x)) = \frac{1}{n} \sum_{i=1}^n \hat{Y}_i(x).$$

Now, the plan of the proof is following. First we shall obtain an uniform law of the iterated logarithm (ULIL) for the differentiable part N_n , which

together with ULIL for the martingale part M_n will imply ULIL for the empirical process β_n . To achieve this, we will use a strong approximation method from [4], which is refinement of the method from [5]. Specifically, we will show first

$$\|nN_n(x) - n\hat{N}_n(x)\|_\alpha \leq C_0 \quad (18)$$

with a specific constant C_0 . Then we will apply a blocking technique to \hat{N}_n , which together with (18) will provide a strong approximation of N_n by an appropriate Gaussian process. ULIL for β_n as well as for the uniform quantile process will easily follow. Then using [24, Lemma 13] we will be able to control increments of the empirical processes β_n and α_n , which together with ULIL will imply the result.

3.1 Approximation of the differentiable part by a Gaussian process

To follow our plan, we show first that we are able to replace the original sequence with its truncated version, i.e. we will establish (18). Then, we will compute variances of $N_n(x, y)$ and $\hat{N}_n(x, y)$. With help of this and the method from [4] we will obtain the approximation result.

With the help of differentiability and the comment following (16),

$$\begin{aligned} & \|F_\epsilon(x - X_{i,i-1}) - F_\epsilon(x - \hat{X}_{i,i-1})\|_\alpha \leq \\ & \leq \sup_{x \in \mathbf{R}} |f_\epsilon(x)| \|X_{i,i-1} - \hat{X}_{i,i-1}\|_\alpha = O \left(\left(\sum_{k=i^\delta}^{\infty} |c_k|^\alpha \right)^{1/\alpha} \right). \end{aligned} \quad (19)$$

Further,

$$|F(x) - \hat{F}_i(x)|^\alpha \leq \mathbb{E} |F_\epsilon(x - X_{i,i-1}) - F_\epsilon(x - \hat{X}_{i,i-1})|^\alpha.$$

Consequently,

$$\|Y_i(x) - \hat{Y}_i(x)\|_\alpha = O \left(\left(\sum_{k=i^\delta}^{\infty} |c_k|^\alpha \right)^{1/\alpha} \right). \quad (20)$$

Thus,

$$\|nN_n(x) - n\hat{N}_n(x)\|_\alpha \leq \sum_{i=1}^n \|Y_i(x) - \hat{Y}_i(x)\|_\alpha \leq O \left(\sum_{i=1}^n \left(\sum_{k=i^\delta}^{\infty} |c_k|^\alpha \right)^{1/\alpha} \right). \quad (21)$$

Therefore, like in (17), the estimate (18) is achieved with

$$C_0 = \sum_{i=1}^{\infty} i^{-1} (\log i)^{-3/2}. \quad (22)$$

Next, we obtain a bound on the variance of $nN_n(x, y)$ and its truncated version $n\hat{N}_n(x, y)$. We have for any $-\infty < x < y < \infty$,

$$\begin{aligned} \|nN_n(x, y)\|_2^2 &= n^2 \mathbf{E} \left| \int_x^y (f_n^*(u) - f(u)) du \right|^2 \\ &\leq n^2 (y-x)^2 \mathbf{E} \sup_{x \leq u \leq y} |f_n^*(u) - f(u)|^2 \leq Cn(y-x)^2 \end{aligned} \quad (23)$$

by Lemma 9 in [24]. Next,

$$\begin{aligned} \mathbf{E} \sup_{x \leq u \leq y} |\hat{f}_n^*(u) - \hat{f}_n(u)|^2 &\leq \mathbf{E} \sup_{x \leq u \leq y} |\hat{f}_n^*(u) - f_n^*(u)|^2 \\ &\quad + \mathbf{E} \sup_{x \leq u \leq y} |f_n^*(u) - f(u)|^2 + \sup_{x \leq u \leq y} |f(u) - \hat{f}_n(u)|^2. \end{aligned} \quad (24)$$

The second part is $O(n^{-1})$ as in (23). As for the first one we have

$$\begin{aligned} \mathbf{E} \sup_{x \leq u \leq y} |\hat{f}_n^*(u) - f_n^*(u)|^2 &= \frac{1}{n} \mathbf{E} \sup_{x \leq u \leq y} \left| \sum_{i=1}^n (f_\epsilon(u - X_{i,i-1}) - f_\epsilon(u - \hat{X}_{i,i-1})) \right|^2 \\ &= \frac{1}{n} \mathbf{E} \sup_{x \leq u \leq y} \left| \sum_{i=1}^n \int_{u - X_{i,i-1}}^{u - \hat{X}_{i,i-1}} f'_\epsilon(v) dv \right|^2 \leq \frac{1}{n} \sup_{x \in \mathbf{R}} |f'_\epsilon(x)|^2 \mathbf{E} \left(\sum_{i=1}^n |X_{i,i-1} - \hat{X}_{i,i-1}| \right)^2 \\ &\leq \frac{1}{n} \sup_{x \in \mathbf{R}} |f'_\epsilon(x)|^2 \left\| \sum_{i=1}^n |X_{i,i-1} - \hat{X}_{i,i-1}| \right\|_\alpha^2 \\ &\leq \frac{1}{n} \sup_{x \in \mathbf{R}} |f'_\epsilon(x)|^2 \left(\sum_{i=1}^n \|X_{i,i-1} - \hat{X}_{i,i-1}\|_\alpha \right)^2 \leq C \frac{1}{n} \end{aligned} \quad (25)$$

in view of $\alpha > 2$ and the comment following (17). Also,

$$\sup_{x \leq u \leq y} |f(u) - \hat{f}_n(u)|^2 \leq \sup_{x \leq u \leq y} \mathbf{E} |\hat{f}_n^*(u) - f_n^*(u)|^2 \leq C \frac{1}{n}. \quad (26)$$

Putting together (24), (25), (26) we arrive at

$$\|n\hat{N}_n(x, y)\|_2^2 \leq n^2 (y-x)^2 \mathbf{E} \sup_{x \leq u \leq y} |\hat{f}_n^*(u) - \hat{f}_n(u)|^2 \leq D_0 n (y-x)^2 \quad (27)$$

with some $D_0 > 0$.

Next, we derive an exponential inequality for $nN_n(x, y)$.

Lemma 3.1 *Assume (10), (11) and (12). Then for any $z > 0$*

$$P\left(\left|\sum_{i=1}^n Y_i(x, y)\right| > z\right) \leq C_1 z^{-\alpha} + C_2 \exp(-C_3 z^2 / (n(y-x)^2)) + C_4 \exp(-C_5 z / n^\rho),$$

where C_1, C_2, C_3, C_4, C_5 are positive constants.

Proof. From (18) and Markov inequality one gets

$$P(|nN_n(x, y)| > z) \leq C_0 z^{-\alpha} + P(|n\hat{N}_n(x, y)| > z/2).$$

To obtain the bound for the second part, divide $[1, n]$ into blocks $I_1, J_1, I_2, J_2, \dots, I_M, J_M$ with the same length n^ρ , ρ from (11). Thus, $M \sim n^{1-\rho}$. Let $\hat{U}_k = \sum_{i \in I_k} \hat{Y}_i(x, y)$, $\hat{V}_k = \sum_{i \in J_k} \hat{Y}_i(x, y)$ and $n\hat{N}_n^{(1)} = \sum_{k=1}^M \hat{U}_k$, $n\hat{N}_n^{(2)} = \sum_{k=1}^M \hat{V}_k$. Both $(\hat{U}_1, \dots, \hat{U}_M)$ and $(\hat{V}_1, \dots, \hat{V}_M)$ are vectors of independent random variables. Also, $\max_{k=1, \dots, M} \hat{U}_k \leq [I_k] \leq Cn^\rho$. The equation (27) yields $\|\hat{U}_k\|_2^2 \leq D_0 n^\rho (y-x)^2$. Recall that $\rho \in (0, \frac{1}{2})$. Applying the result in [21, p. 293] to the centered sequence $\hat{U}_1, \dots, \hat{U}_M$ with $M_n = n^\rho$, $B_n = D_0 n (y-x)^2$ one obtains

$$P(|n\hat{N}_n^{(1)}| > z) \leq \exp(-z^2 / (4D_0 n (y-x)^2)) + \exp(-z / (4n^\rho)).$$

The same applies to $P(|n\hat{N}_n^{(2)}| > z)$ and hence the result follows. \odot

Remark 3.2 Note that the proof of Lemma 2.8 in [4] has a gap. The result from [21] is applied to a non-centered sequence.

To state an approximation result, note that $Y_0(x)$ and $\hat{Y}_i(x)$, $i \geq 1$ are independent. Thus, we obtain

$$\begin{aligned} |\mathbf{E}Y_0(x)Y_i(y)| &\leq |\mathbf{E}Y_0(x)Y_i(y) - Y_0(x)\hat{Y}_i(y)| + |\mathbf{E}Y_0(x)\hat{Y}_i(y)| \\ &\leq \|Y_i(y) - \hat{Y}_i(y)\|_2 + 0 = O(i^{-\rho(\alpha\tau-1)/\alpha}). \end{aligned}$$

Then, clearly,

$$\Gamma(x, y) = \mathbf{E}Y_0(x)Y_0(y) + \sum_{i=1}^{\infty} (\mathbf{E}Y_0(x)Y_i(y) + \mathbf{E}Y_0(y)Y_i(x)) \quad (28)$$

is absolutely convergent for all $x, y \in \mathbb{R}$.

Having Lemma 3.1, (28) and $\alpha > 2$ we may proceed in the very same way as in [4] to obtain the following approximation result (see also Appendix).

Proposition 3.3 *Assume (10), (11) and (12). Then there exists a centered Gaussian process $K(x, t)$ with $\mathbb{E}K(x, t)K(y, t') = t \wedge t' \Gamma(x, y)$ such that*

$$\sup_{0 \leq t \leq 1} \sup_{x \in \mathbb{R}} |[nt]^{1/2} N_{[nt]}(x) - K(x, [nt])| = o(n^{1/2}(\log n)^{-\lambda}) \quad \text{almost surely}$$

with some $\lambda > 0$.

Remark 3.4 It should be pointed out that the process $K(x, t)$ in Proposition 3.3 approximates N_n , not β_n . We use this as a technical tool to achieve ULIL only. Using the blocking technique applied to β_n directly, we may establish Proposition 3.3 with $N_{[nt]}$ replaced with $\beta_{[nt]}$ and a different Gaussian process. However, it requires stronger than (11) assumptions on coefficients c_k . This is clear. Without exploiting differentiability, we have to compare indicators (non-smooth functions) applied to the original sequences X_i with the indicators applied to truncated sequence, see for example [4].

3.2 Laws of the iterated logarithm

The law of iterated logarithm follows from Proposition 3.3 and the ULIL for K . We have

$$\limsup_{n \rightarrow \infty} \frac{n^{1/2}}{(2 \log \log n)^{1/2}} \sup_{-\infty < x < \infty} |N_n(x)| = C \quad \text{almost surely.}$$

This, together with ULIL for the martingale M_n (see e.g. [24, Lemma 7]) yields

$$\limsup_{n \rightarrow \infty} \frac{n^{1/2}}{(2 \log \log n)^{1/2}} \sup_{-\infty < x < \infty} |\beta_n(x)| = C \quad \text{almost surely.} \quad (29)$$

If (29) holds then taking $U_i = F(X_i)$, $i \geq 1$, we obtain

$$\limsup_{n \rightarrow \infty} \frac{1}{(2 \log \log n)^{1/2}} \sup_{x \in (0,1)} |\alpha_n(x)| = C \quad \text{almost surely.}$$

Consequently,

$$\limsup_{n \rightarrow \infty} \frac{1}{(2 \log \log n)^{1/2}} \sup_{y \in (0,1)} |u_n(y)| = \limsup_{n \rightarrow \infty} \frac{1}{(2 \log \log n)^{1/2}} \sup_{x \in (0,1)} |\alpha_n(x)| = C, \quad (30)$$

almost surely. We may extend this result if Kiefer conditions are met.

Corollary 3.5 *If (29) holds for $\beta_n(x)$ and Kiefer conditions are fulfilled, then*

$$\limsup_{n \rightarrow \infty} \frac{1}{(2 \log \log n)^{1/2}} \sup_{y \in (0,1)} |q_n(y)| = C \quad \text{almost surely.} \quad (31)$$

Proof. Given X_1, \dots, X_n define $U_i = F(X_i)$, $i = 1, \dots, n$. Let $U_{1:n} \leq \dots \leq U_{n:n}$. With $\Delta_{n,y} = Q_n(y) - Q(y)$, we have

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \sup_{y \in (0,1)} \frac{n^{1/2} \Delta_{n,y}}{(\log \log n)^{1/2}} \\ & \leq \limsup_{n \rightarrow \infty} \sup_{1 \leq k \leq n} \frac{n^{1/2} |Q(F(X_{k:n})) - Q(\frac{k}{n})|}{(\log \log n)^{1/2}} + \limsup_{n \rightarrow \infty} \sup_{1 \leq k \leq n} \sup_{y \in (\frac{k-1}{n}, \frac{k}{n}]} \frac{n^{1/2} |Q(\frac{k}{n}) - Q(y)|}{(\log \log n)^{1/2}}. \end{aligned}$$

Now, if Kiefer conditions hold, then $\sup_{y \in (0,1)} |Q''(y)| < \infty$. Using Taylor's expansion one obtains $Q(\frac{k}{n}) = Q(y) + Q'(y)(\frac{k}{n} - y) + O(\frac{1}{n})$ and

$$Q(F(X_{k:n})) = Q(U_{k:n}) = Q\left(\frac{k}{n}\right) + Q'\left(\frac{k}{n}\right) \left(U_{k:n} - \frac{k}{n}\right) + O_{a.s.} \left(\left(U_{k:n} - \frac{k}{n}\right)^2 \right).$$

Thus, by (30),

$$\limsup_{n \rightarrow \infty} \sup_{y \in (0,1)} \frac{n^{1/2} \Delta_{n,y}}{(\log \log n)^{1/2}} \leq \sup_{y \in (0,1)} |Q'(y)| \limsup_{n \rightarrow \infty} \sup_{1 \leq k \leq n} \frac{n^{1/2} |U_{k:n} - \frac{k}{n}|}{(\log \log n)^{1/2}} \leq C.$$

◊

3.3 Controlling increments

Let $l_q(n) = (\log n)^{1/q} (\log \log n)^{2/q}$ and recall that $\sum_{k=0}^{\infty} |c_k| < \infty$.

First, we generalize [24, Lemma 13] to the whole \mathbb{R} .

Lemma 3.6 *Assume (10) and (12). Let $-\infty \leq a < b \leq \infty$. Then for any positive bounded sequence d_n such that $\log n = o(nd_n)$ one has*

$$\sup_{|x-y| \leq d_n, x, y \in [a,b]} |F_n(x) - F(x) - (F_n(y) - F(y))| = O_{a.s.} \left(\frac{\sqrt{d_n \log n}}{\sqrt{n}} + \frac{d_n l_q(n)}{\sqrt{n}} \right). \quad (32)$$

Proof. Define $U_i = F(X_i)$. Clearly, in order to show (32) it suffices to prove

$$\sup_{|u-v| \leq d_n, u, v \in (0,1)} |E_n(v) - v - (E_n(u) - u)| = O_{a.s} \left(\frac{\sqrt{d_n \log n}}{\sqrt{n}} + \frac{d_n l_q(n)}{\sqrt{n}} \right). \quad (33)$$

Let $y \in (0, 1)$. Decompose

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n (1_{\{U_i \leq y\}} - y) = \\ &= \frac{1}{n} \sum_{i=1}^n \left(1_{\{U_i \leq y\}} - \mathbb{E}(1_{\{U_i \leq y\}} | \mathcal{F}_{i-1}) \right) + \frac{1}{n} \sum_{i=1}^n \left(\mathbb{E}(1_{\{U_i \leq y\}} | \mathcal{F}_{i-1}) - y \right) \\ &=: \tilde{M}_n(x) + \tilde{N}_n(x), \end{aligned}$$

where

$$\tilde{N}_n(y) = \frac{1}{n} \sum_{i=1}^n \left(\mathbb{E}(1_{\{U_i \leq y\}} | \mathcal{F}_{i-1}) - y \right) = \frac{1}{n} \sum_{i=1}^n (F_\epsilon(Q(y) - X_{i,i-1}) - y)$$

so that

$$\tilde{N}'_n(y) = Q'(y) \frac{1}{n} \sum_{i=1}^n (f_\epsilon(Q(y) - X_{i,i-1}) - y) \quad (34)$$

and

$$\tilde{N}''_n(y) = Q''(y) \frac{1}{n} \sum_{i=1}^n (f_\epsilon(Q(y) - X_{i,i-1}) - y) + (Q'(y))^2 \frac{1}{n} \sum_{i=1}^n \left(f'_\epsilon(Q(y) - X_{i,i-1}) - y \right). \quad (35)$$

Define the projection operator $\mathcal{P}_k \xi = \mathbb{E}(\xi | \mathcal{F}_k) - \mathbb{E}(\xi | \mathcal{F}_{k-1})$. Let $g_y(\mathcal{F}_i) = f_\epsilon(Q(y) - X_{i,i-1})$. As in the proof of Lemma 3 in [24], with $\alpha = q$,

$$\|\mathcal{P}_0 g_y(\mathcal{F}_i)\|_\alpha = O(|c_i|)$$

uniformly in $y \in (0, 1)$. The same holds for $g_y(\mathcal{F}_i) = f'_\epsilon(Q(y) - X_{i,i-1})$. Further, under Kiefer conditions, $\sup_{y \in (0,1)} (|Q'(y)| + |Q''(y)|) < \infty$. Thus, applying Proposition 1 in [24], $\max_{y \in (0,1)} (\|\tilde{N}'_n(y)\|_\alpha + \|\tilde{N}''_n(y)\|_\alpha) = O(n^{-1/2})$. Consequently,

$$\mathbb{E} \left[\max_{y \in (0,1)} |\tilde{N}_n(y)|^\alpha \right] \leq \mathbb{E} \left[\int_0^1 |\tilde{N}'_n(y)| dy \right]^\alpha = O(n^{-\alpha/2}).$$

Similarly, $E \left[\max_{y \in (0,1)} |\tilde{N}'_n(y)|^\alpha \right] = O(n^{-\alpha/2})$. Thus, by the same argument as in [24, Lemma 9] one has $\sup_{y \in (0,1)} |\tilde{N}'_n(y)| = o_{a.s.}(l_q(n)/\sqrt{n})$. Further,

$$\sup_{|u-v| \leq d_n, u, v \in (0,1)} |\tilde{N}_n(v) - \tilde{N}_n(u)| \leq d_n \sup_{y \in (0,1)} |\tilde{N}'_n(y)| = o_{a.s.}(l_q(n)/\sqrt{n}).$$

This, together, with appropriate estimates for the martingale part yields (33). ◊

3.4 Conclusion of the proof

We apply Lemma 3.6 with $d_n = \lambda_n$. Then the second part in (32) is negligible. Thus, we have

$$\sup_{x \in \mathbf{R}} \sup_{|x-y| \leq \lambda_n} |\beta_n(x) - \beta_n(y)| = O_{a.s.}(b_n).$$

On account of (31) it yields

$$\sup_{y \in (0,1)} |\beta_n(Q_n(y)) - \beta_n(Q(y))| = O_{a.s.}(b_n). \quad (36)$$

Equivalently,

$$n^{1/2} \sup_{y \in (0,1)} |F_n(Q_n(y)) - F(Q_n(y)) - (F_n(Q(y)) - F(Q(y)))| = O_{a.s.}(b_n).$$

Since $|F_n(Q_n(y)) - F(Q(y))| \leq 1/n$ one obtains

$$n^{1/2} \sup_{y \in (0,1)} |F(Q_n(y)) - F(Q(y)) - (F(Q(y)) - F_n(Q(y)))| = O_{a.s.}(b_n).$$

Set $\Delta_{n,y} = Q_n(y) - Q(y)$. Using the Taylor's expansion $F(Q_n(y)) = F(Q(y)) + f(Q(y))\Delta_{n,y} + O_{a.s.}(\Delta_{n,y}^2)$ we finish the proof of Theorem 2.1. ◊

4 Proof of Theorem 2.3

Note that $Q'(y) = \frac{1}{f(Q(y))}$ and $Q''(y) = -\frac{f'(Q(y))}{f^3(Q(y))}$. Moreover,

$$(h(y)\tilde{N}_n(y))' = h'(y)\tilde{N}_n(y) + h(y)\tilde{N}'_n(y).$$

Thus, in view of (34), we need $h'(y)$ and $h(y)Q'(y)$ to be uniformly bounded, which is achieved by $\nu > 1$ and $\nu > \gamma + \mu$, respectively. Moreover,

$$(h(y)\tilde{N}_n(y))'' = h''(y)\tilde{N}_n(y) + 2h'(y)\tilde{N}'_n(y) + h(y)\tilde{N}''_n(y).$$

Thus, in view of (34), (35) we need $h''(y)$, $h'(y)Q'(y)$ and $h(y)Q''(y)$ to be uniformly bounded. The first claim is achieved by $\nu > 2$, the second by $\nu - 1 > \gamma + \mu$, respectively. As for the third one we have

$$h(y)|Q''(y)| = \left(\frac{|f'(Q(y))|}{f^2(Q(y))}(y(1-y)) \right) \left(\frac{(y(1-y))^{\nu-1}}{f(Q(y))} \right).$$

Since (CsR3) is fulfilled, the first part is $O(1)$ uniformly in $y \in (0, 1)$. The second part is $O(1)$ since $\nu - 1 > 2\gamma - 1 > \gamma$ (recall $\gamma > 1$).

Let $g(y) = h(y)\tilde{N}_n(y)$. Thus, proceeding exactly as in Lemma 3.6 we obtain $E \left[\max_{y \in (0,1)} |g'(y)|^\alpha \right] = O(n^{-\alpha/2})$. Thus, $\sup_{y \in (0,1)} |g'(y)| = o_{a.s.}(l_q(n)/\sqrt{n})$ and

$$\sup_{|u-v| \leq d_n, u, v \in (0,1)} |g(v) - g(u)| \leq d_n \sup_{y \in (0,1)} |g'(y)| = o_{a.s.}(l_q(n)/\sqrt{n}).$$

Further, by differentiability of h ,

$$\sup_{|u-v| \leq d_n, u, v \in (0,1)} |h(u) - h(v)| |\tilde{N}_n(u)| \leq d_n \sup_{y \in (0,1)} |\tilde{N}'_n(y)| = o_{a.s.}(l_q(n)/\sqrt{n}).$$

Consequently,

$$\sup_{|u-v| \leq d_n, u, v \in (0,1)} h(v) |\tilde{N}_n(v) - \tilde{N}_n(u)| = o_{a.s.}(l_q(n)/\sqrt{n}).$$

Therefore, taking into account the martingale part and $d_n = \lambda_n$

$$\sup_{|u-v| \leq \lambda_n, u, v \in (0,1)} h(v) |(E_n(v) - v) - (E_n(u) - u)| = O_{a.s.} \left(\frac{\sqrt{d_n \log n}}{\sqrt{n}} \right). \quad (37)$$

Note that

$$\begin{aligned} u_n(y) &= \sqrt{n}(E_n(U_n(y)) - U_n(y)) - \sqrt{n}(E_n(U_n(y)) - y) \\ &= \sqrt{n}(E_n(U_n(y)) - U_n(y)) + O_{a.s.}(n^{-1/2}). \end{aligned}$$

Consequently, by (37) and (30)

$$\sup_{y \in (0,1)} h(y) |u_n(y) - \alpha_n(y)| = \sup_{y \in (0,1)} h(y) |\alpha_n(U_n(y)) - \alpha_n(y)| + O_{a.s.}(n^{-1/2}) = O_{a.s.}(b_n).$$

Further, let $\delta_n = 2C^*n^{-1/2}(\log \log n)^{1/2}$, $C^* = C$, C from (30). As in [8], with $\theta = \theta_n(y)$ such that $|\theta - y| \leq n^{-1}u_n(y) = O_{a.s}(\delta_n)$,

$$\begin{aligned} & \sup_{y \in (\delta_n, 1-\delta_n)} h(y) |f(Q(y))q_n(y) - u_n(y)| = \\ & = n^{-1/2}u_n^2(y) \left(\frac{f'(Q(\theta))}{f^2(Q(\theta))} \theta(1-\theta) \right) \frac{f(Q(\theta))}{f(Q(y))} \frac{(y(1-y))^\nu}{\theta(1-\theta)}. \end{aligned} \quad (38)$$

In view of [8, Lemma 1] one has

$$\frac{f(Q(\theta))}{f(Q(y))} \leq \left\{ \frac{y \vee \theta}{y \wedge \theta} \frac{1-y \wedge \theta}{1-y \vee \theta} \right\}^\gamma. \quad (39)$$

Further on, if $y \geq 2\delta_n$,

$$\frac{y}{\theta} = \frac{y-\theta}{\theta} + 1 \leq \frac{\delta_n}{y - n^{-1/2}u_n(y)} + 1 \leq 2. \quad (40)$$

Consequently, by (CsR3), (39), (40) one has

$$\sup_{y \in (\delta_n, 1-\delta_n)} h(y) |f(Q(y))q_n(y) - u_n(y)| = O_{a.s}(n^{-1/2} \log \log n). \quad (41)$$

Let $(k-1)/n < y \leq k/n$. If $1 \geq U_{k:n} \geq y$ then as in [8, (3.14)]

$$\begin{aligned} & \sup_{y \in (0, \delta_n)} h(y) |f(Q(y))q_n(y)| = \sup_{y \in (0, \delta_n)} n^{1/2}h(y) \int_y^{U_{k:n}} \frac{f(Q(y))}{f(Q(u))} du \\ & \leq C \sup_{y \in (0, \delta_n)} n^{1/2}h(y) \int_y^{U_{k:n}} \left(\frac{u}{y} \right)^{\gamma_1} du \\ & \leq C \sup_{y \in (0, \delta_n)} n^{1/2}(U_{k:n} - y)h(y)y^{-\gamma_1} = O_{a.s}(n^{-1/2} \log \log n) \end{aligned}$$

by (30) as long as $\nu > \gamma_1 + 1$. Further, if $U_{k:n} \leq y$ then for $\gamma_1 > 1$

$$\sup_{y \in (0, \delta_n)} h(y) |f(Q(y))q_n(y)| \leq Cn^{1/2}\delta_n^{\nu+\gamma_1}U_{k:n}^{-(\gamma_1-1)} \quad (42)$$

and for $\gamma_1 = 1$,

$$\sup_{y \in (0, \delta_n)} h(y) |f(Q(y))q_n(y)| \leq Cn^{1/2}\delta_n^{\nu+1} \log(\delta_n/U_{k:n}). \quad (43)$$

Now,

$$P(U_{1:n} \leq n^{-2}(\log n)^{-3/2}) \leq \sum_{i=1}^n P(U_i \leq n^{-2}(\log n)^{-3/2}) \leq n^{-1}(\log n)^{-3/2}.$$

Consequently, via the Borel-Cantelli lemma, $U_{k:n} = o_{a.s.}(n^2(\log n)^{3/2})$. Therefore, the bound in (43) is $O_{a.s.}(n^{-1/2} \log \log n)$ and the bound in (42) is of the same order provided $\nu > 3\gamma_1 - 2$. The upper tail is treated in the similar way. Consequently, the result follows. \odot

5 Proof of Corollary 2.5

The proof follows the similar lines as in [6]. Let $A_n(y) = 2n^{1/2} \int_{Q_n(y)}^y (\alpha_n(u) - \alpha_n(y)) du$. Then like in [6, Lemma 3.1]

$$\sup_{y \in (0,1)} |I_n(y) - A_n(y)| = O_{a.s.}(n^{-1/2}(\log n)(\log \log n)^{1/2})$$

and it suffices to show that $\sup_{y \in (0,1)} A_n(y) = O_{a.s.}(n^{-1/4}(\log n)^{1/2}(\log \log n)^{3/4})$. Since

$$A_n(y) = -2u_n(y) \int_0^1 \left(\alpha_n \left(y + s \frac{u_n(y)}{n^{1/2}} \right) - \alpha_n(y) \right) ds \quad (44)$$

and $\sup_{y \in (0,1)} |\alpha_n(y)| = O_{a.s.}((\log \log n)^{1/2})$, it suffices to prove that the integral in (44) is $O_{a.s.}(n^{-1/4}(\log n)^{1/2}(\log \log n)^{1/4})$. This can be justified in the same way as in (36). \odot

6 Remarks

Remark 6.1 As mentioned before, the strong mixing, even with geometric rates, does not yield the optimal rates in the bahadur-Kiefer representation. However, the part of this paper is devoted to the law of the iterated logarithm for the empirical process. One may ask, whether imposing (11), one could obtain an uniform law of the iterated logarithm for empirical processes via strong mixing. From [23] one knows that if $c_k = O(k^{-\tau})$, $\tau > 1 + \frac{1}{\alpha} + \max\{1, \alpha^{-1}\}$, then $\alpha(n) = O(n^{-\lambda})$, where $\lambda = (\tau\alpha - \max\{\alpha, 1\}) / (1 + \alpha) - 1 > 0$ and $\alpha(n)$ is the strong mixing coefficients. In particular, if $\alpha > 2$, then his condition yields $\tau > 2 + \frac{1}{\alpha}$ and $\alpha(n) = O(n^{-\frac{\tau\alpha - 1 - 2\alpha}{1 + \alpha}})$. In view of [22] to obtain LIL for partial sums of bounded strongly mixing random variables one needs $\alpha(n)$ to be summable, which would require $\tau > 3 + \frac{2}{\alpha}$. Thus, first we need stronger conditions than (11), second, Rio's result would provide LIL for $\beta_n(x)$ at *fixed* x .

Remark 6.2 Assuming $\alpha > 8$ and $\tau > 4 + \frac{1}{\alpha}$ we may use the exponential inequality (see Lemma 3.1) instead of Lemma 3.6 and its modification at the beginning of Section 4 to obtain the optimal rates for the distance between uniform empirical and quantile processes without additional weight function. Then, coupling this with [26, Theorem 2.1] we may obtain under (CsR1), (CsR2)

$$\sup_{y \in (\delta_n, 1 - \delta_n)} |f(Q(y))q_n(y) - \alpha_n(y)| = O_{a.s.}(n^{-1/4}(\log n)^{1/2}(\log \log n)^{1/4}), \quad (45)$$

where $\delta_n = n^{-1/4}(\log n)^{-1/2}(\log \log n)^{3/4}$. Further, under somehow stronger conditions than (CsR3), (CsR4) one may obtain in (45) a bound on the whole interval, but with worse rates, see [26] for details.

Remark 6.3 As mentioned before, we obtain the strong approximation for the differentiable part N_n only in order to have as less restrictive conditions on c_k as possible. Of course, one could think about blocking technique and strong approximation applied to the original sequence.

Assume for simplicity $c_k = O(k^{-\tau})$ and $E|\epsilon|^\alpha$, $\alpha > 0$. Then, as in (16), $E|X_i - \hat{X}_i|^\alpha = O(i^{-\rho(\alpha\tau-1)})$. Further, with $X_i(x) = 1_{\{X_i \leq x\}} - F(x)$, $\hat{X}_i(x) = 1_{\{\hat{X}_i \leq x\}} - F(x)$ and some $\mu > 0$,

$$\begin{aligned} & P(X_i(x) \neq \hat{X}_i(x)) \\ & \leq P\left(|1_{\{X_i \leq x\}} - 1_{\{\hat{X}_i \leq x\}}| > 0, |X_i - \hat{X}_i| < i^{-\mu}\right) + P(|X_i - \hat{X}_i| \geq i^{-\mu}) \\ & \leq P(x - i^{-\mu} \leq X_i \leq x + i^{-\mu}) + O(i^{-\rho(\alpha\tau-1)+\mu\alpha}) = O(i^{-\mu}) + O(i^{-\rho(\alpha\tau-1)+\mu\alpha}). \end{aligned}$$

Next, for any $\kappa > 0$

$$\left(E \left| \sum_{i=1}^n (X_i(x) - \hat{X}_i(x)) \right|^\kappa\right)^{1/\kappa} = O\left(\sum_{i=1}^n i^{-\mu/\kappa} + \sum_{i=1}^n i^{-(\rho(\alpha\tau-1)-\mu\alpha)/\kappa}\right).$$

Then, κ would replace α in the exponential inequality of Lemma 6.6. Thus, we would require $\kappa > 2$. Consequently, to have both sums convergent we would require $\mu > \kappa > 2$ and $\tau > \frac{\kappa}{\rho\alpha} + \frac{1}{\alpha} + \frac{\mu}{\rho}$. The parameters μ and κ can be chosen as close to 2 as we wish. Further, ρ must be chosen between $(0, \frac{1}{2})$. Therefore, one would conclude $\gamma > 5 + \frac{4}{\alpha}$. This condition is much worse than ours, however, there is no restriction $\alpha > 2$. Thus, in this case we may state the result of Proposition 3.3 in terms of β_n . Additionally, we do not require assumptions (10), (11) and (12).

Appendix

Here, we re-prove strong approximation of Proposition 3.3 following the lines of [4].

We are keeping notation from Section 3.1.

For $\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{R}^d$ define random vectors in \mathbb{R}^d :

$$\xi_i = (Y_i(x_1), \dots, Y_i(x_d)), \quad \hat{\xi}_i = (\hat{Y}_i(x_1), \dots, \hat{Y}_i(x_d)).$$

Denote by $\Gamma_d = \Gamma_d(\mathbf{x}) = (\Gamma(x_i, x_j), 1 \leq i, j \leq d)$ the covariance matrix of ξ_i . We shall use the following implication which is valid for any vectors ξ_i, η_i in \mathbb{R}^d :

$$\mathbb{E}\|\xi_i - \eta_i\| \leq A \text{ implies } |\mathbb{E} \exp(i \langle \mathbf{u}, \xi_i \rangle) - \mathbb{E} \exp(i \langle \mathbf{u}, \eta_i \rangle)| \leq \|\mathbf{u}\| d^{1/2}. \quad (46)$$

Also, we will use the following bound: for $\mathbf{x}_i \in \mathbb{R}^d, i = 1, \dots, n$, one has $\|\sum_{i=1}^n \mathbf{x}_i\| \leq \sum_{i=1}^n \|\mathbf{x}_i\|$.

Lemma 6.4 *Under conditions of Proposition 3.3, for all $\mathbf{u} \in \mathbb{R}^d$*

$$\left| \mathbb{E} \exp(i \langle \mathbf{u}, n^{-1/2} \sum_{i=1}^n \xi_i \rangle) - \mathbb{E} \exp(i \langle \mathbf{u}, n^{-1/2} \sum_{i=1}^n \hat{\xi}_i \rangle) \right| \leq \|\mathbf{u}\| d^{1/2} n^{-1/2}.$$

Proof. Let $\beta = \rho(\alpha\tau - 1)/\tau$. From (20) we have $\|Y_i(x) - \hat{Y}_i(x)\| \leq \|Y_i(x) - \hat{Y}_i(x)\|_\alpha \leq C i^{-\beta}$. Thus

$$\mathbb{E}\|\xi_i - \hat{\xi}_i\|^2 \leq C d i^{-2\beta} \quad (47)$$

and (47) implies

$$\mathbb{E}\|\xi_i - \hat{\xi}_i\| \leq C d^{1/2} i^{-\beta}.$$

This implies $P(\|\xi_i - \hat{\xi}_i\| > d^{1/2}) \leq C i^{-\beta}$ which together with the Borel-Cantelli lemma yields

$$\sup_{n \geq 1} \sum_{i=1}^n \|\xi_i - \hat{\xi}_i\| \leq d^{1/2} \text{ almost surely.}$$

Therefore,

$$\mathbb{E} \left\| \sum_{i=1}^n (\xi_i - \hat{\xi}_i) \right\| \leq d^{1/2}.$$

This, together with (46) implies the result. \odot

Proposition 6.5 *Under conditions of Theorem 3.3, for all $\mathbf{u} \in \mathbb{R}^d$*

$$\begin{aligned} & \left| \mathbb{E} \exp(i \langle \mathbf{u}, n^{-1/2} \sum_{i=1}^n \xi_i \rangle) - \exp(-\frac{1}{2} \langle \mathbf{u}, \Gamma_d \mathbf{u} \rangle) \right| \\ & \leq Cd \|\mathbf{u}\| n^{-\delta_1} + C \|\mathbf{u}\|^2 (n^{-1/4} (\log n)^{d/2} \exp(Cd) + d^{d/2} \exp(-Cn^{1/2})) \end{aligned}$$

with some $\delta_1 > 0$.

Proof. Divide the interval $[1, n]$ into consecutive long and short blocks $I_1, J_1, I_2, J_2, \dots$ with length $\text{card}(I_k) = \lceil n^{\rho^*} \rceil$, $\text{card}(J_k) = \lceil n^\rho \rceil$, $1 \leq k \leq M$, $\rho < \rho^* < \frac{1}{2}$. Then $M = M_n \sim n^{1-\rho^*}$ (the last block is possibly incomplete). Define $U_k = \sum_{i \in I_k} \xi_i$, $\hat{U}_k = \sum_{i \in I_k} \hat{\xi}_i$, $\hat{V}_k = \sum_{i \in J_k} \hat{\xi}_i$. By (47) we have

$$\mathbb{E} \|U_k - \hat{U}_k\| \leq \sum_{i \in I_k} \mathbb{E} \|\xi_i - \hat{\xi}_i\| \leq Cd^{1/2} \sum_{i \in I_k} i^{-\beta}. \quad (48)$$

Note that \hat{U}_k , $k = 1, \dots, M$ are independent. Thus, we can construct independent random vectors $\tilde{U}_1, \dots, \tilde{U}_M$ such that $(\tilde{U}_k, \hat{U}_k) \stackrel{d}{=} (U_k, \hat{U}_k)$, $k = 1, \dots, M$.

By (48) and $(\tilde{U}_k, \hat{U}_k) \stackrel{d}{=} (U_k, \hat{U}_k)$ one obtains

$$\mathbb{E} \left\| \sum_{k=1}^M (\tilde{U}_k - \hat{U}_k) \right\| \leq \sum_{k=1}^M \mathbb{E} \|U_k - \hat{U}_k\| \leq Cd^{1/2} \sum_{k=1}^M \sum_{i \in I_k} i^{-\beta} \leq Cd^{1/2} \sum_{i=1}^n i^{-\beta} \leq Cd^{1/2}. \quad (49)$$

Further,

$$\mathbb{E} \|\hat{V}_k\| = \mathbb{E} \left\| \sum_{i \in J_k} \hat{\xi}_i \right\| \leq dn^{\rho/2}$$

which implies by independence

$$\mathbb{E} \left\| \sum_{k=1}^M \hat{V}_k \right\|^2 \leq \sum_{k=1}^M \mathbb{E} \|\hat{V}_k\|^2 \leq CdMn^\rho = dn^{\rho+1-\rho^*}. \quad (50)$$

We have

$$\sum_{i=1}^n \hat{\xi}_i = \sum_{k=1}^M (\hat{U}_k - \tilde{U}_k) + \sum_{k=1}^M \tilde{U}_k + \sum_{k=1}^M \hat{V}_k.$$

Thus, by (49), (50)

$$\mathbb{E} \left\| \sum_{i=1}^n \xi_i - \sum_{k=1}^M \tilde{U}_k \right\| \leq C(d^{1/2} + dn^{(\rho+1-\rho^*)/2}).$$

Consequently,

$$\left\| \mathbb{E} \exp(i \langle \mathbf{u}, n^{-1/2} \sum_{i=1}^n \xi_i \rangle) - \mathbb{E} \exp(i \langle \mathbf{u}, n^{-1/2} \sum_{k=1}^M \tilde{U}_k \rangle) \right\| \leq Cd \|\mathbf{u}\| n^{-(\rho^* - \rho)/2}. \quad (51)$$

Also, by [4, Lemma 2.9] and the same argument as at the end of Lemma 2.10 in the latter paper one obtains

$$\begin{aligned} & \left| \mathbb{E} \exp(i \langle \mathbf{u}, n^{-1/2} \sum_{k=1}^M \tilde{U}_k \rangle) - \exp(-\frac{1}{2} \langle \mathbf{u}, \Gamma_d \mathbf{u} \rangle) \right| \\ & \leq C \|\mathbf{u}\|^2 (n^{-(1-\rho^*)/2} (\log n)^{d/2} \exp(Cd) + d^{d/2} \exp(-Cn^{1-\rho^*})). \end{aligned} \quad (52)$$

By (51) and (52) the result follows. \odot

6.1 Approximation

Let $\varepsilon \in (0, 1/4)$. Denote $t_k = \exp(k^{1-\varepsilon})$, $p_k = \lceil t_k^{\rho/2} \rceil$, $d_k = k^{1/2}$ and $x_i = (i-1)/d_k$, $i \in \mathbb{Z}$, $M_k = t_{k+1} - t_k - p_k$ so that $M_k \sim Ck^{-\varepsilon} \exp(k^{1-\varepsilon})$ and $\log M_k \sim Ck^{1-\varepsilon}$. Define random vectors in \mathbb{R}^{d_k}

$$\eta_k = (\eta_{k1}, \dots, \eta_{kd_k}),$$

where

$$\eta_{ki} = R(x_i, t_{k+1}) - R(x_i, t_k + p_k) = \sum_{j=t_k+p_k+1}^{t_{k+1}} Y_j(x_i).$$

Also, $\eta_{ki} = \sum_{j=t_k+p_k+1}^{t_{k+1}} \hat{Y}_j(x_i)$.

By Proposition 6.5

$$\begin{aligned} & \left| \mathbb{E} \exp(i \langle \mathbf{u}, M_k^{-1/2} \eta_k \rangle) - \exp(-\frac{1}{2} \langle \mathbf{u}, \Gamma_{d_k} \mathbf{u} \rangle) \right| \\ & \leq C \|\mathbf{u}\|^2 \left(d_k M_k^{-\rho_1} + M_k^{-(1-\rho^*)/2} (\log M_k)^{d_k/2} \exp(Cd_k) + d_k^{d_k/2} \exp(-CM_k^{1-\rho^*}) \right) \\ & \leq C \|\mathbf{u}\|^2 \left(k^{1/2} k^{\varepsilon \rho_1} \exp(-C\rho_1 k^{1-\varepsilon}) \right) \\ & \quad + C \|\mathbf{u}\|^2 \left(k^{\varepsilon(1-\rho^*)/2} \exp(-C(1-\rho^*)k^{1-\varepsilon}/2) k^{(1-\varepsilon)\sqrt{k}/2} \exp(\sqrt{k}) \right) \\ & \leq C \exp(-Ck^{1-\varepsilon}) \|\mathbf{u}\|^2. \end{aligned}$$

Let ψ_{PL} be the Prokhorov-Levy distance. Choose $T := \exp(k^\varepsilon)$; for sufficiently large k , $T > 10^8 d_k$. By [5, Lemma 2.2] we obtain

$$\begin{aligned}
\psi_{PL}(M_k^{-1/2}\eta_k, N(\mathbf{0}, \Gamma_{d_k})) &\leq \frac{16d_k}{T} \log T + P(N(\mathbf{0}, \Gamma_{d_k}) > T/2) \\
&\quad + T^{d_k} \int_{\|\mathbf{u}\| \leq T} \left| \mathbb{E} \exp(i \langle \mathbf{u}, M_k^{-1/2}\eta_k \rangle) - \exp(-\frac{1}{2} \langle \mathbf{u}, \Gamma_{d_k} \mathbf{u} \rangle) \right| d\mathbf{u} \\
&\leq \frac{16d_k}{T} \log T + P(N(\mathbf{0}, \tau_{d_k}) > T/2) + CT^{d_k} \exp(-k^{1-\varepsilon}) \int_{\|\mathbf{u}\| \leq T} \|\mathbf{u}\|^2 d\mathbf{u} \\
&\leq \frac{16d_k}{T} \log T + k^{1/2} \exp(-cT^2 k^{-1/2}) + CT^{d_k} 2^{d_k} T^2 \exp(-ck^{1-\varepsilon}) \\
&\leq \frac{16\sqrt{k}}{\exp(k^\varepsilon)} k^\varepsilon + C \exp(C(k^{\frac{1}{2}+\varepsilon} - k^{1-\varepsilon})) + \exp(\frac{1}{2} \log k - \exp(k^\varepsilon k^{-1/2})) \\
&\leq C \exp(-Ck^\varepsilon)
\end{aligned}$$

since $\varepsilon < 1/4$.

Since $\hat{\eta}_k$, $k = 1, \dots, M$ are independent, we can define independent random vectors ζ_1, \dots, ζ_M such that $M_k^{-1/2}\zeta_k \sim N(\mathbf{0}, \Gamma_{d_k})$ and

$$P(\|M_k^{-1/2}\hat{\eta}_k - M_k^{-1/2}\zeta_k\| > C \exp(-k^\varepsilon)) \leq C \exp(-k^\varepsilon).$$

This yields

$$P(\|M_k^{-1/2}\eta_k - M_k^{-1/2}\zeta_k\| > C \exp(-k^\varepsilon)) \leq C \exp(-k^\varepsilon)$$

and then by Borel-Cantelli lemma

$$M_k^{-1/2}(\eta_k - \zeta_k) \leq \exp(-k^\varepsilon) \text{ almost surely.} \quad (53)$$

For $k \geq 1$, define random vectors Z_k and Y_k in \mathbb{R}^{d_k} by

$$Z_k = (R(x_i, t_{k+1}) - R(x_i, t_k), i = 1, \dots, d_k).$$

Since $Z_k - \eta_k$ is the sum of p_k random vectors in \mathbb{R}^{d_k} with coordinates bounded by 1 and since $M_k \leq t_k$, $d_k^{1/2} p_k^{1/2} \leq kt_k^{\rho/2} \leq \exp(ck^{1-\varepsilon})$, $\varepsilon < 1/2$ we have by (53)

$$\begin{aligned}
\|Z_k - \zeta_k\| &\leq \|Z_k - \eta_k\| + \|\eta_k - \zeta_k\| \\
&\leq C(d_k^{1/2} p_k + t_k^{1/2} \exp(-k^\varepsilon)) \\
&\leq C(\sqrt{k} t_k^{\rho/2} + t_k^{1/2} \exp(-k^\varepsilon)) \leq C t_k^{1/2} \exp(-Ck^\varepsilon). \quad (54)
\end{aligned}$$

Thus, the skeleton process $\{Z_k, k \geq 1\}$ can be approximated by the sequence $\{\zeta_k, k \geq 1\}$. The latter can be extended to a centered Gaussian process $\{K(x, t), x \in \mathbb{R}, t \geq 0\}$ with covariance $t \wedge t' \Gamma_{d_k}(s, x')$ such that

$$\zeta_{ki} = K(x, t_{k+1}) - K(x, t_k + p_k), \quad i = 1, \dots, d_k.$$

Define now

$$Y_k = (K(x_i, t_{k+1}) - K(x_i, t_k), i = 1, \dots, d_k).$$

Then

$$\|\zeta_k - Y_k\| \leq C t_k^{1/2} \exp(-C k^\varepsilon) \quad \text{almost surely} \quad (55)$$

by the last inequality on p. 807 in [4]. Thus, by (54)

$$\|Z_k - Y_k\| \leq C t_k^{1/2} \exp(-C k^\varepsilon) \quad \text{almost surely.} \quad (56)$$

6.2 Oscillations

The next lemma will be used to control oscillations of $R(x, y)$ as well as in the proof of the Bahadur-Kiefer representation.

Lemma 6.6 *Under conditions of Theorem 3.3, for $n \geq 1$, $\lambda \geq n^{\frac{1}{2}}$ and any $-\infty < a < b < \infty$ we have*

$$P \left(\sup_{a \leq x \leq x' \leq b, 0 \leq k \leq n} \sum_{i=1}^k Y_i(x, x') \geq \lambda \right) \leq C \exp \left(-C \lambda^2 / (n(b-a)^2) \right) + \frac{C}{n^\eta},$$

where $\eta > 0$.

Proof. Define

$$M_{u,v} = \max_{0 \leq i \leq 2^u, 0 \leq j \leq 2^v} \left| \sum_{n=nj2^{-v}}^{n(j+1)2^{-v}} Y_i(bi2^{-u}, b(i+1)2^{-u}) \right|.$$

Then for any integer $L \geq 1$,

$$\left| \sum_{i=1}^k Y_i(0, x) \right| \leq \sum_{1 \leq u, v \leq L} M_{u,v} + \frac{2n}{2^L}.$$

Choose

$$0 < \varepsilon < (\alpha - 2)/C_{12} \quad (57)$$

for some constant C_{12} to be specified later. Since $n^{1/2} \leq \lambda$, $u, v \leq L$ we have by Lemma 3.1

$$\begin{aligned}
& P(M_{u,v} \geq \lambda 2^{-\varepsilon(u+v)}) \\
& \leq 2^{u+v} C \left\{ (\lambda 2^{-\varepsilon(u+v)})^{-\alpha} + \exp\left(-C\lambda^2 2^{-2\varepsilon(u+v)} / (n 2^{-v} (b 2^{-u})^2)\right) + \right. \\
& \quad \left. \exp\left(-C\lambda 2^{-\varepsilon(u+v)} / (n 2^{-v})^\rho\right) \right\} \\
& \leq 2^{u+v} C \left\{ \lambda^{-\alpha} 2^{\varepsilon(u+v)\alpha} + \exp\left(-C\lambda^2 2^{-2\varepsilon(u+v)} 2^v 2^{2u} / (nb^2)\right) + \right. \\
& \quad \left. \exp\left(-C\lambda 2^{-\varepsilon(u+v)} 2^{v\rho} / n^\rho\right) \right\} \\
& \leq 2^{u+v} C \left\{ n^{-\frac{1}{2}\alpha} 2^{2\varepsilon L\alpha} + \exp\left(-C\lambda^2 2^{u(2-2\varepsilon)} 2^{v(1-2\varepsilon)} / (nb^2)\right) + \right. \\
& \quad \left. \exp\left(-Cn^{(\frac{1}{2}-\rho)} 2^{-\varepsilon(u+v)}\right) \right\} \\
& \leq 2^{u+v} C \left\{ n^{-\frac{1}{2}\alpha} 2^{2\varepsilon L\alpha} + \exp\left(-C\lambda^2 2^u 2^v / (nb^2)\right) + \right. \\
& \quad \left. \exp\left(-Cn^{(\frac{1}{2}-\rho)} 2^{-\varepsilon(u+v)}\right) \right\} \\
& \leq 2^{u+v} C \left\{ n^{-\frac{1}{2}\alpha} 2^{2\varepsilon L\alpha} + \exp\left(-C\lambda^2 2^u 2^v / (nb^2)\right) + \right. \\
& \quad \left. \left(n^{(\frac{1}{2}-\rho)} 2^{-\varepsilon(u+v)}\right)^{-r} \right\}
\end{aligned}$$

for any $r > 0$. Choose $r = \alpha$, then since $\alpha > 2$ and $u + v \leq 2L$,

$$\begin{aligned}
& P(M_{u,v} \geq \lambda 2^{-\varepsilon(u+v)}) \\
& \leq 2^{u+v} C \left\{ n^{-\alpha/2} 2^{2\varepsilon L\alpha} + \exp\left(-C\lambda^2 2^u 2^v / (nb^2)\right) + n^{-\alpha/2} 2^{\varepsilon(u+v)\alpha} \right\} \\
& \leq 2^{u+v} C \left\{ n^{-\alpha/2} 2^{2\varepsilon L\alpha} + \exp\left(-C\lambda^2 2^u 2^v / (nb^2)\right) \right\}.
\end{aligned}$$

For

$$A = \{M_{u,v} \geq \lambda 2^{-\varepsilon(u+v)} \text{ for some } 1 \leq u, v \leq L\}$$

one has

$$\begin{aligned}
P(A) & \leq \sum_{1 \leq u, v \leq L} 2^{u+v} \exp\left(-C\lambda^2 2^u 2^v / (nb^2)\right) + 2^{2L+2} 2^{C L \varepsilon} n^{-\alpha/2} \\
& \leq \sum_{1 \leq u, v \leq L} \exp(u+v) \exp\left(-C\lambda^2 2^u / (nb^2)\right) \exp\left(-C\lambda^2 2^v / (nb^2)\right) \\
& \quad + 2^{2L+2} 2^{C L \varepsilon} n^{-\alpha/2} \\
& \leq \left(\sum_{1 \leq u \leq L} \exp\left(u - \lambda^2 2^u / (nb^2)\right) \right)^2 + 2^{L(2+C\varepsilon)+2} n^{-\alpha/2} \\
& \leq C \exp\left(-C\lambda^2 / (nb^2)\right) + C 2^{L(2+C_{12}\varepsilon)} n^{-\alpha/2}.
\end{aligned}$$

Choose L so that $n^{1/2} < 2^L 2n^{1/2}$, then $\frac{2n}{2^L} \leq 2^L \leq 2\lambda$ and by (57) the second part in the latter expression is equal to $n^{-\eta}$ with some $\eta > 0$. Therefore,

$$\begin{aligned} & P\left(\sup_{0 \leq x \leq b, 0 \leq k \leq n} \left| \sum_{i=1}^k Y_i(0, x) \right| \geq \lambda\right) \\ & \leq P(A) + P\left(\sum_{1 \leq u, v \leq L} M_{u, v} + \frac{2n}{2^L} \geq \lambda; A^c\right) \leq C \exp\left(-C\lambda^2/(nb^2)\right) + Cn^{-\eta}. \end{aligned}$$

Thus, the result follows. \odot

Let

$$\tilde{R}_{i, k} = \sup_{s \in [x_i, x_{i+1}]} \sup_{0 \leq t \in [t_k, t_{k+1}]} |R(x, y) - R(x_i, t_k)|$$

and $\phi(k) = \log k((t_{k+1} - t_k)^{1/2} + t_{k+1}^{1/2} d_k^{-1}) > \max\{t_k^{1/2}, (t_{k+1} - t_k)^{1/2}\}$ for sufficiently large k .

Lemma 6.7 *Under conditions of Theorem 3.3,*

$$P\left(\max_{1 \leq i \leq d_k} |\tilde{R}_{i, k}| > \phi(k)\right) \leq C t_k^{1/2} (\log k)^{-\varepsilon^*} \quad \text{almost surely}$$

for some $\varepsilon^* > 0$.

Proof. Choose

$$r > 3/2. \tag{58}$$

Then by Lemma 6.6

$$\begin{aligned} & P\left(\max_{1 \leq i \leq d_k} |\tilde{R}_{i, k}| > \phi(k)\right) \\ & \leq d_k P\left(\sup_{s \in [x_i, x_{i+1}]} \sup_{0 \leq t \in [t_k, t_{k+1}]} |R(x, y) - R(s, t_k)| > \phi(k)/2\right) + \\ & \quad d_k P\left(\sup_{s \in [x_i, x_{i+1}]} |R(s, t_k) - R(x_i, t_k)| > \phi(k)/2\right) \\ & \leq C d_k \exp\left(-C \frac{\phi^2(k)}{(t_{k+1} - t_k)(x_{i+1} - x_i)^2}\right) + d_k \frac{C}{(t_{k+1} - t_k)^\eta} + \\ & \quad C d_k \exp\left(-C \frac{\phi^2(k)}{t_k(x_{i+1} - x_i)^2}\right) + d_k \frac{C}{t_k^\eta} \end{aligned}$$

$$\begin{aligned} &\leq Ck^{1/2} \left(\frac{\phi^2(k)}{(t_{k+1} - t_k)(x_{i+1} - x_i)^2} \right)^{-r} + \frac{Ck^{1/2}}{(t_{k+1} - t_k)^\eta} + \\ &Ck^{1/2} \left(C \frac{\phi^2(k)}{t_k(x_{i+1} - x_i)^2} \right)^{-r} + \frac{Ck^{1/2}}{t_k^\eta}. \end{aligned}$$

Using $t_{k+1} - t_k \sim M_k \sim k^{-\varepsilon} \exp(k^{1-\varepsilon}) \leq \exp(k^{1-\varepsilon}) = t_k$ one obtains

$$\begin{aligned} P \left(\max_{1 \leq i \leq d_k} |\tilde{R}_{i,k}| > \phi(k) \right) &\leq C \frac{k^{1/2} t_k^r}{\phi^{2r}(k) k^r} + \frac{Ck^{1/2}}{(t_{k+1} - t_k)^\eta} \\ &\leq C \frac{Ck^{1/2} t_k^r}{t_k^r k^r} + \frac{Ck^{1/2}}{(t_{k+1} - t_k)^\eta} \leq Ck^{(1-2r)/2} + C \exp(-Ck^{1-\varepsilon}) \end{aligned}$$

and the bound is summable by (58). The result follows by Borel-Cantelli Lemma. ◊

6.3 Conclusion of the proof

For $s \in [x, x_{i+1})$, $t \in [t_k, t_{k+1}]$ write

$$|R(x, y) - K(x, t)| \leq |R(x, y) - R(x_i, t_k)| + |K(x, t) - K(x_i, t_k)| + |R(x_i, t_k) - K(x_i, t_k)|.$$

Both the first and the second part are bounded almost surely by $Ct_k^{1/2}(\log k)^{-\varepsilon^*}$, the first one by Lemma 6.7, the second one by (2.55) in [4]. The second part is bounded in the same way as (2.56) in [4].

Acknowledgement.

This work was done during my stay at Carleton University. I am thankful to Professors Barbara Szyszkowicz and Miklos Csörgő for the support and helpful remarks.

References

- [1] Avram, F. and Taqqu, M. (1986). Weak convergence of moving averages with infinite variance. *Dependence in probability and statistics* (Oberwolfach, 1985), 399–415, Progr. Probab. Statist., 11, Birkhäuser Boston, Boston, MA.

- [2] Babu, G.J. and Singh, K. (1978). On deviations between empirical and quantile processes for mixing random variables. *Journal of Multivariate Analysis* **8**, 532–549.
- [3] Bahadur, R.R. (1966). A note on quantiles in large samples. *Ann. Math. Statist.* **37**, 577–580.
- [4] Berkes, I. and Horváth, L. (2001). Strong approximation of the empirical process of GARCH sequences. *Ann. Appl. Probab.* **11**, 789–809.
- [5] Berkes, I. and Philipp, W. (1977). An almost sure invariance principle for the empirical distribution function of mixing random variables. *Z. Wahr. und Verw. Gebiete* **41**, 115–137.
- [6] Csáki, E., Csörgő, M., Földes, A., Shi, Z. and Zitikis, R. (2002). Pointwise and uniform asymptotics of the Vervaat error process. *J. Theoret. Probab.* **15**, 845–875.
- [7] Csörgő, M. (1983). *Quantile Processes with Statistical Applications*. CBMS-NSF Regional Conference Series in Applied Mathematics.
- [8] Csörgő, M. and Révész, P. (1978). Strong approximation of the quantile process. *Ann. Statist.* **6**, 882–894.
- [9] Csörgő, M. and Révész, P. (1981). *Strong Approximation in Probability and Statistics*.
- [10] Csörgő, M. and Szyszkowicz, B. (1998). Sequential quantile and Bahadur-Kiefer processes. *Order statistics: theory & methods*, 631–688, Handbook of Statist., 16, North-Holland, Amsterdam.
- [11] Dehling, H., Mikosch, T. and Sorensen, M. (eds.) (2002). *Empirical process techniques for dependent data*, 3–113, Birkhuser Boston, Boston, MA.
- [12] Doukhan, P. (1984). *Mixing: Properties and Examples*. Lecture notes in Statistics. Springer-Verlag.
- [13] Fotopoulos, S.B. and Ahn, S.K. (1994). Strong Approximation of the Quantile Processes and Its Applications under Strong Mixing Properties. *J. Mult. Anal.* **51**, 17–45.
- [14] Freedman, D.A. (1975). On tail probabilities for martingales. *Ann. Probab.* **3**, 100–118.

- [15] Hesse, C.H. (1990). A Bahadur-Type Representation for Empirical Quantiles of a Large Class of Stationary, Possibly Infinite-Variance, Linear Processes. *Ann. Statist.* **18**, 1188–1202.
- [16] Ho, H.-C. and Hsing, T. (1997). Limit theorems for functionals of moving averages. *Ann. Probab.* **25**, 1636–1669.
- [17] Hsing, T. and Wu, W.B. (2004) On weighted U -statistics for stationary sequences. *Ann. Probab.* **32**, 1600–1631.
- [18] Kiefer, J. (1967). On Bahadur’s representation of sample quantiles. *Ann. Math. Statist.* **38**, 1323–1342.
- [19] Kiefer, J. (1970). Deviations between the sample quantile process and the sample df. *nonparametric Techniques in Statistical Inference*, 349–357, M.L. Puri, ed., Cambridge University Press.
- [20] Parzen, E. (1979). Nonparametric statistical data modeling. With comments by John W. Tukey, Roy E. Welsch, William F. Eddy, D. V. Lindley, Michael E. Tarter and Edwin L. Crow, and a rejoinder by the author. *J. Amer. Statist. Assoc.* **74**, 105–131.
- [21] Petrov, V.V. (1975). *Sums of Independent Random Variables*. Springer-Verlag.
- [22] Rio, E. (1995). The functional law of the iterated logarithm for stationary strongly mixing sequences. *Ann. Probab.* **23**, 1188–1203.
- [23] Withers, C. S. (1981). Conditions for linear processes to be strong-mixing. *Z. Wahrsch. Verw. Gebiete* **57**, 477–480.
- [24] Wu, W.B. (2005). On the Bahadur representation of sample quantiles for dependent sequences. *Ann. Statist.* **33**, 1934–1963.
- [25] Wu, W. B. and Mielniczuk, J. (2002). Kernel density estimation for linear processes. *Ann. Statist.* **30**, 1441–1459.
- [26] Yu, H. (1996). A note on strong approximation for quantile processes of strong mixing sequences. *Stat. Prob. Letters.* **30**, 1–7.