Toward the best constant factor for the Rademacher-Gaussian tail comparison

Iosif Pinelis

Department of Mathematical Sciences Michigan Technological University Houghton, Michigan 49931, USA E-mail: ipinelis@mtu.edu

Abstract: Let $S_n := a_1 \varepsilon_1 + \cdots + a_n \varepsilon_n$, where $\varepsilon_1, \ldots, \varepsilon_n$ are independent Rademacher random variables and a_1, \ldots, a_n are any real numbers such that $a_1^2 + \cdots + a_n^2 = 1$. Let $Z \sim N(0, 1)$. It is proved that the best constant factor c in inequality $\mathsf{P}(S_n \ge x) \le c \mathsf{P}(Z \ge x) \ \forall x \in \mathbb{R}$ is between two explicitly defined absolute constants c_1 and c_2 such that $c_1 < c_2 \approx 1.01 c_1$.

AMS 2000 subject classifications: Primary 60E15,62G10,62G15; secondary 60G50,62G35, 26D10.

Keywords and phrases: optimal upper bounds, probability inequalities, Rademacher random variables, sums of independent random variables, Student's test, self-normalized sums.

1. Introduction and summary

Let $\varepsilon_1, \ldots, \varepsilon_n$ be independent Rademacher random variables (r.v.'s), so that $\mathsf{P}(\varepsilon_i = 1) = \mathsf{P}(\varepsilon_i = -1) = \frac{1}{2}$ for all *i*. Let

$$S_n := a_1 \varepsilon_1 + \dots + a_n \varepsilon_n,$$

where a_1, \ldots, a_n are any real numbers such that

 $a_1^2 + \dots + a_n^2 = 1.$

The best upper exponential bound on the tail probability $\mathsf{P}(Z \ge x)$ for a standard normal random variable Z and a nonnegative number x is $\inf_{t\ge 0} e^{-tx} \mathsf{E} e^{tZ} = e^{-x^2/2}$. Thus, a factor of the order of magnitude of $\frac{1}{x}$ is "missing" in this bound, compared with the asymptotics $\mathsf{P}(Z \ge x) \sim \frac{1}{x} \varphi(x)$ as $x \to \infty$, where $\varphi(x) := e^{-x^2/2}/\sqrt{2\pi}$ is the density function of Z. Now it should be clear that any exponential upper bound on the tail probabilities for sums of independent random variables must be missing the $\frac{1}{x}$ factor.

Eaton [6] obtained an upper bound on $\mathsf{P}(S_n \ge x)$, which is asymptotic to $c_3 \mathsf{P}(Z \ge x)$ as $x \to \infty$, where

$$c_3 := \frac{2e^3}{9} \approx 4.46,$$

and he conjectured that $\mathsf{P}(S_n \ge x) \le c_3 \frac{1}{x} \varphi(x)$ for $x > \sqrt{2}$. The stronger form of this conjecture,

1

$$\mathsf{P}(S_n \ge x) \leqslant c \,\mathsf{P}(Z \ge x) \tag{1}$$

for all $x \in \mathbb{R}$ with $c = c_3$ was proved by Pinelis [11], along with a multidimensional extension.

Edelman [7] proposed inequality $P(S_n \ge x) \le P(Z \ge x - 1.5/x)$ for all x > 0, but his proof appears to have a gap. A more precise upper bound, with $\ln c_3 =$ 1.495... in place of 1.5, was recently shown [20] to be a rather easy corollary of the mentioned result of [11]. Various generalizations and improvements of inequality (1) as well as related results were given by Pinelis [12, 13, 15, 16, 17, 19, 20] and Bentkus [1, 2, 3].

Bobkov, Götze and Houdré (BGH) [4] gave a simple proof of (1) with a constant factor $c \approx 12.01$. Their method was based on the Chapman-Kolmogorov identity for the Markov chain (S_n) . Such an identity was used, e.g., in [14] to disprove a conjecture by Graversen and Peškir [9] on $\max_{k \leq n} |S_k|$.

In this paper, we shall show that a modification of the BGH method can be used to obtain inequality (1) with a constant factor $c \approx 1.01 c_*$, where c_* is the best (that is, the smallest) possible constant factor c in (1).

Let $\overline{\Phi}$ and r denote the tail function of Z and the inverse Mills ratio:

$$\overline{\Phi}(x) := \mathsf{P}(Z \ge x) = \int_x^\infty \varphi(u) \, du \quad \text{and} \quad r := \frac{\varphi}{\overline{\Phi}}.$$

Theorem 1 (Main). For the least possible absolute constant factor c_* in inequality (1) one has

$$c_* \in [c_1, c_2] \approx [3.18, 3.22], \quad where$$

$$c_1 := \frac{1}{4\overline{\Phi}(\sqrt{2})} \quad and \quad c_2 := c_1 \cdot \left(1 + \frac{1}{250} \left(1 + r(\sqrt{3})\right)\right) \approx c_1 \cdot 1.01.$$

Here we shall present just one application of Theorem 1, to self-normalized sums

$$V_n := \frac{X_1 + \dots + X_n}{\sqrt{X_1^2 + \dots + X_n^2}}$$

where, following Efron [8], we assume that the X_i 's satisfy the so-called orthant symmetry condition: the joint distribution of $\delta_1 X_1, \ldots, \delta_n X_n$ is the same for any choice of signs $\delta_1, \ldots, \delta_n \in \{1, -1\}$, so that, in particular, each X_i is symmetrically distributed. It suffices that the X_i 's be independent and symmetrically (but not necessarily identically) distributed. In particular, $V_n = S_n$ if $X_i = a_i \varepsilon_i$ $\forall i$. It was noted by Efron that (i) Student's statistic T_n is a monotonic function of the so-called self-normalized sum: $T_n = \sqrt{\frac{n-1}{n}} V_n / \sqrt{1 - V_n^2/n}$ and (ii) the orthant symmetry implies in general that the distribution of V_n is a mixture of the distributions of normalized Rademacher sums S_n . Thus, one obtains

Corollary 1. Theorem 1 holds with V_n in place of S_n .

2. Proof of Theorem 1

Theorem 1 follows immediately from Lemma 1, Theorem 2, and Lemma 3, stated in Subsection 2.1 below. In particular, Lemma 3 implies that the upper bound

 $h_1(x)$ on $\mathsf{P}(S_n \ge x)$ provided by Theorem 2 is somewhat better than the upper bound $c_2 \mathsf{P}(Z \ge x)$, implied by Theorem 1.

While S_n represents a simplest case of the sum of independent non-identically distributed r.v.'s, it is still very difficult to control in a precise manner. Figure 1 shows the graph of the ratio $R(x) := P(S_n \ge x)/P(Z \ge x)$ for n = 100 and $a_1 = \cdots = a_n$.



FIG 1. Ratio of the Rademacher and Gaussian tails for n = 100 and $a_1 = \cdots = a_{100} = \frac{1}{10}$.

One can see that even for such a fairly large value of n and equal coefficients a_1, \ldots, a_n , ratio R(x) oscillates rather wildly. In view of this, the existence of a high-precision inductive argument in the general setting with possibly unequal a_i 's may seem very unlikely. However, such an argument will be presented in this paper. A key idea in the proof of Theorem 1 is the construction of the upper bound h_1 and, in particular, the function g in (2) and (3), which allows an inductive argument to prove Theorem 2.

The proof of Theorem 2 is based on a number of lemmas. The proofs of all lemmas are deferred to Subsection 2.2.

2.1. Statements of lemmas and the proof of Theorem 2

Lemma 1. One has $c_* \ge c_1$.

For $a \in [0, 1)$ and $x \in \mathbb{R}$, introduce

$$g(x) := c_1 \cdot \left(1 + \frac{1}{250} (1 + r(x))\right) \overline{\Phi}(x) = \frac{c_1}{250} \cdot \left(251 \,\overline{\Phi}(x) + \varphi(x)\right); \quad (2)$$

$$h(x) := c_1 \cdot \left(1 + \frac{1}{250} (1 + r(\sqrt{3}))\right) \overline{\Phi}(x) = c_2 \cdot \overline{\Phi}(x);$$

$$h_1(x) := \begin{cases} 1 & \text{if } x \leqslant 0, \\ \frac{1}{2} & \text{if } 0 < x \leqslant 1, \\ \frac{1}{2x^2} & \text{if } 1 \leqslant x < \sqrt{2}, \\ g(x) & \text{if } \sqrt{2} \leqslant x \leqslant \sqrt{3}, \\ h(x) & \text{if } x \geqslant \sqrt{3}; \end{cases}$$
(3)

$$K(a, x) := h_1(u) + h_1(v) - 2h_1(x), \quad \text{where} \\ u := u(a, x) := \frac{x - a}{\sqrt{1 - a^2}} \quad \text{and}$$
(4)

$$v := v(a, x) := \frac{x+a}{\sqrt{1-a^2}}.$$
(5)

Theorem 2 (Refined). One has

$$\mathsf{P}(S_n \geqslant x) \leqslant h_1(x) \tag{6}$$

for all $x \in \mathbb{R}$.

Lemma 2. One has $g \leq h$ on $(-\infty, \sqrt{3}]$ and $g \geq h$ on $[\sqrt{3}, \infty)$.

Lemma 3. One has $h_1 \leq h$ on \mathbb{R} .

Lemma 4. One has $K(a, x) \leq 0$ for all $(a, x) \in [0, 1) \times [\sqrt{2}, \sqrt{3}]$.

Lemma 5. One has $K(a, x) \leq 0$ for all $(a, x) \in [0, 1) \times [\sqrt{3}, \infty)$.

Now we can present

Proof of Theorem 2. Theorem 2 will be proved by induction in n. It is obvious for n = 1. Let now $n \in \{2, 3, ...\}$ and assume that Theorem 2 holds with n - 1 in place of n.

Note that for $x \leq 0$ inequality (6) is trivial. For $x \in (0, \sqrt{2})$, it follows by the symmetry of S_n and Chebyshev's inequality. Therefore, assume without loss of generality that $x \geq \sqrt{2}$ and $0 \leq a_n < 1$. By the Chapman-Kolmogorov identity and induction,

$$P(S_n \ge x) = \frac{1}{2} P(S_{n-1} \ge x - a_n) + \frac{1}{2} P(S_{n-1} \ge x + a_n)$$

$$\leq \frac{1}{2} h_1 (u(a_n, x)) + \frac{1}{2} h_1 (v(a_n, x))$$

$$= h_1(x) + \frac{1}{2} K(a_n, x)$$

for all $x \in \mathbb{R}$. It remains to refer to Lemmas 4 and 5.

Lemma 4 is based on a series of other lemmas. To state those lemmas, more notation is needed:

$$\begin{aligned} x_* &:= \sqrt{\frac{5+2\sqrt{6}}{9-2\sqrt{6}}} = \sqrt{\frac{23+28\sqrt{\frac{2}{3}}}{19}} \approx 1.55; \end{aligned} \tag{7} \\ \overline{R} &:= \{(a,x) \in \mathbb{R}^2 \colon 0 \le a \le 1, \sqrt{2} \le x \le \sqrt{3}\}; \\ R &:= \{(a,x) \in \mathbb{R}^2 \colon 0 < a < 1, \sqrt{2} < x < \sqrt{3}\}; \\ \text{LIe} &:= \{(a,x) \in \mathbb{R} \colon u < \sqrt{2}, v \le \sqrt{3}\}; \\ \text{LG} &:= \{(a,x) \in \mathbb{R} \colon u < \sqrt{2}, v > \sqrt{3}\}; \\ \text{GL}_1 &:= \{(a,x) \in \mathbb{R} \colon u > \sqrt{2}, v < \sqrt{3}, x \le x_*\}; \\ \text{GL}_2 &:= \{(a,x) \in \mathbb{R} \colon u > \sqrt{2}, v < \sqrt{3}, x < x_*\}; \\ \text{GG}_1 &:= \{(a,x) \in \mathbb{R} \colon u > \sqrt{2}, v < \sqrt{3}, a < \frac{1}{\sqrt{3}}\}; \\ \text{GG}_1 &:= \{(a,x) \in \mathbb{R} \colon u > \sqrt{2}, v > \sqrt{3}, a < \frac{1}{\sqrt{3}}\}; \\ \text{GG}_2 &:= \{(a,x) \in \mathbb{R} \colon u > \sqrt{2}, v > \sqrt{3}, a < \frac{1}{\sqrt{3}}\}; \\ \text{GE} &:= \{(a,x) \in \mathbb{R} \colon u > \sqrt{2}, v < \sqrt{3}, a < \frac{1}{\sqrt{3}}\}; \\ \text{EIe} &:= \{(a,x) \in \mathbb{R} \colon u = \sqrt{2}, v < \sqrt{3}, a < \frac{1}{\sqrt{3}}\}; \\ \text{EG}_1 &:= \{(a,x) \in \mathbb{R} \colon u = \sqrt{2}, v > \sqrt{3}, a < \frac{1}{\sqrt{3}}\}; \\ \text{EG}_1 &:= \{(a,x) \in \mathbb{R} \colon u = \sqrt{2}, v > \sqrt{3}, a < \frac{1}{\sqrt{3}}\}; \\ \text{EG}_2 &:= \{(a,x) \in \mathbb{R} \colon u = \sqrt{2}, v > \sqrt{3}, a < \frac{1}{\sqrt{3}}\}; \\ \text{EG}_2 &:= \{(a,x) \in \mathbb{R} \colon u = \sqrt{2}, v > \sqrt{3}, a > \frac{1}{\sqrt{3}}\}; \\ \text{A}_1 &:= \{(a,x) \in \overline{\mathbb{R}} \colon a = 0\}; \\ \text{A}_2 &:= \{(a,x) \in \overline{\mathbb{R}} \colon x = \sqrt{2}, 0 < a < \frac{1}{\sqrt{2}}\}; \\ \text{X}_{1,1} &:= \{(a,x) \in \overline{\mathbb{R}} \colon x = \sqrt{2}, \frac{2\sqrt{2}}{3} \le a < 1\}; \\ \text{X}_{1,3} &:= \{(a,x) \in \overline{\mathbb{R}} \colon x = \sqrt{2}, \frac{2\sqrt{2}}{3} \le a < 1\}; \\ \text{X}_2 &:= \{(a,x) \in \overline{\mathbb{R}} \colon x = \sqrt{3}\}, \end{aligned}$$

where u and v are defined by (4) and (5). Here, for example, the L in the first position in symbol LIe refers to "less than" in inequality $u < \sqrt{2}$, while the ligature Ie in the second position refers to "less than or equal to" in inequality $v \leq \sqrt{3}$. Similarly, G and E in this notation refer to "greater than" and "equal to", respectively. Symbol A refers here to a fixed value of a, and X to a fixed value of x.

It will be understood that the function K is extended to \mathbf{A}_2 by continuity, so that

$$K(a, x) := -2g(x) \quad \forall (a, x) \in \mathcal{A}_2.$$
(8)

Lemma 6 (LLe). The function K does not attain a maximum on LLe.

Lemma 7 (LG). The function K does not attain a maximum on LG.

Lemma 8 (GL₁). The function K does not attain a maximum on GL_1 .

Lemma 9 (GL₂). The function K does not attain a maximum on GL_2 .

Lemma 10 (GG₁). The function K does not attain a maximum on GG_1 .

Lemma 11 (GG₂). The function K does not attain a maximum on GG_2 .

Lemma 12 (GE). One has $K \leq 0$ on GE.

Lemma 13 (ELe). One has $K \leq 0$ on ELe.

Lemma 14 (EG₁). One has $K \leq 0$ on EG₁.

Lemma 15 (EG₂). One has $K \leq 0$ on EG₂.

Lemma 16 (A₁). One has K = 0 on A₁.

Lemma 17 (A₂). One has $K \leq 0$ on A₂.

Lemma 18 (X_{1,1}). One has $K \leq 0$ on X_{1,1}.

Lemma 19 (X_{1,2}). One has $K \leq 0$ on X_{1,2}.

Lemma 20 (X_{1,3}). One has $K \leq 0$ on X_{1,3}.

Lemma 21 (X₂). One has $K \leq 0$ on X₂.

The proofs of some of these lemmas require a great amount of symbolic and numerical computation. We have done that with the help of MathematicaTM 5.2, which is rather effective and allows complete and easy control over the accuracy. The idea behind these calculations is to reduce a problem involving transcendental inequalities and/or equations to an algebraic setting, for it is well known that the existence of a real-valued solution to any system of algebraic equations and/or inequalities can be determined in a completely algorithmic manner, according to a result by Tarski [21, 10, 5]. This algorithm is implemented in Mathematica 5.2 via **Reduce** and related commands. However, execution of such commands may take a long time if the algebraic system is more than a little complicated; in such cases, Mathematica can use some human help.

2.2. Proofs of the lemmas

Proof of Lemma 1. Let n = 2 and $a_1 = a_2 = \frac{1}{\sqrt{2}}$. Then $\mathsf{P}(S_n \ge \sqrt{2}) = \frac{1}{4} = c_1 \mathsf{P}(Z \ge \sqrt{2})$.

Proof of Lemma 2. This follows from the well-known and easy-to-prove fact that the inverse Mills ratio r is increasing.

Proof of Lemma 3. On interval $(-\infty, 0]$, one has $h_1 = 1 < 1.6 < h(0) \leq h$, since h is decreasing.

On interval (0, 1], one similarly has $h_1 = \frac{1}{2} < 0.51 < h(1) \leq h$. On interval $[\sqrt{2}, \sqrt{3}]$, one has $h_1 = g \leq h$, by Lemma 2. On interval $[\sqrt{3}, \infty)$, one has $h_1 = h$.

It remains to consider the interval $(1,\sqrt{2})$. For $x \in (1,\sqrt{2})$, one has $h_1(x) = p(x) := \frac{1}{2x^2}$. Now let us apply a l'Hospital-type rule for monotonicity; see e.g. [18, Proposition 4.3] and the notation used therein. One has $p(\infty -) = h(\infty -) = 0$ and, for some constant C > 0, $h'(x)/p'(x) = Cx^3\varphi(x)$, so that $\frac{h'}{p'} \nearrow \omega$ on $(0,\infty)$. Hence, $\frac{h}{p} \searrow$ or $\nearrow \omega$ on $(0,\infty)$, and so, $\inf_{(1,\sqrt{2})} \frac{h}{h_1} = \min_{[1,\sqrt{2}]} \frac{h}{p} = \min_{\{1,\sqrt{2}\}} \frac{h}{p} \approx 1.01 > 1$, whence $h_1 < h$ on $(1,\sqrt{2})$.

Proof of Lemma 4. Since the function h_1 is upper-semicontinuous on \mathbb{R} and continuous on $[\sqrt{2}, \infty)$, the function K is upper-semicontinuous on the compact rectangle \overline{R} and therefore attains its maximum on \overline{R} . Now Lemma 4 follows from Lemmas 6–21 (proved below), since \overline{R} is the union of sets LLe,..., X₂.

Proof of Lemma 5. Let $0 \leq a < 1$ and $x \geq \sqrt{3}$. Then $u \geq \sqrt{2}$ and $v \geq \sqrt{3}$, where u and v are defined by (4) and (5), as before. Therefore and in view of Lemma 3 and (3), one has $h_1(u) \leq h(u)$, $h_1(v) = h(v)$, and $h_1(x) = h(x)$, so that

$$K(a,x) \leq 2c_2 \cdot \left(\frac{1}{2}\overline{\Phi}(u) + \frac{1}{2}\overline{\Phi}(v) - \overline{\Phi}(x)\right) \leq 0,$$

as shown in the mentioned proof in [4].

Proof of Lemma 6 (LLe). Expressing x and u in view of (4) and (5) in terms of a and v as $x(a, v) := \sqrt{1 - a^2} v - a$ and $\tilde{u}(a, v) := v - \frac{2a}{\sqrt{1 - a^2}}$, respectively, one has

$$K(a, x) = k(a, v) := k_{\text{LLe}}(a, v) := \frac{1}{2\tilde{u}(a, v)^2} + g(v) - 2g(x(a, v)) \quad \forall (a, x) \in \text{LLe},$$
(9)

since u > 1 and $v > \sqrt{2}$ on R. For $(a, x) \in R$, let

$$(D_a k)(a, v) := 4\overline{\Phi}(\sqrt{2}) \left(x(a, v) - a\right)^3 \frac{\partial k}{\partial a}(a, v); \tag{10}$$

$$(D_{a,a}k)(a,x) := \frac{125(1-a^2)^2}{(x-a)^2\varphi(x)} \frac{\partial(D_ak)}{\partial a} (a,v(a,x)).$$
(11)

Then $(D_{a,a}k)(a,x)$ is an algebraic expression (and even a polynomial) in a and x. With Mathematica 5.2, one can therefore use the command

Reduce[daakx<=0 && 0<a<1 && Sqrt[2]<x<Sqrt[3]]

(where daakx stands for $(D_{a,a}k)(a,x)$), which outputs False, meaning that $D_{a,a}k > 0$ on R. By (11), this implies $\frac{\partial (D_a k)}{\partial a}(a, v(a, x)) > 0$ for $(a, x) \in R$, so that $(D_a k)(a, v)$ is increasing in a for every fixed value of v; more exactly, $(D_a k)(a, v)$ is increasing in $a \in (a_1(v), a_2(v))$ for every fixed value of $v \in (\sqrt{2}, \sqrt{3}]$, where a_1 and a_2 are certain functions, such that

$$(a, x(a, v)) \in \text{LLe} \iff (v \in (\sqrt{2}, \sqrt{3}] \& a \in (a_1(v), a_2(v))).$$

Thus, for every fixed value of $v \in (\sqrt{2}, \sqrt{3}]$ the sign pattern of $(D_a k)(a, v)$ in $a \in (a_1(v), a_2(v))$ is - or + or -+; that is, $(D_a k)(a, v)$ may change sign

only from - to + as a increases. By (10), $\frac{\partial k}{\partial a}(a, v)$ has the same sign pattern. Hence, for every fixed value of $v \in (\sqrt{2}, \sqrt{3}]$ one has $k(a, v) \searrow$ or \nearrow or $\searrow \nearrow$ in $a \in (a_1(v), a_2(v))$. Now Lemma 6 follows.

Proof of Lemma 7 (LG). This proof is almost identical to that of Lemma 6, except that the term g(v) in (9) is replaced here by h(v), and the interval $(\sqrt{2}, \sqrt{3}]$ is replaced by $(\sqrt{3}, 5\sqrt{2})$. However, both terms g(v) and h(v) are constant for any fixed value of v.

Proof of Lemma 8 (GL_1). One has

$$(a, x) \in \mathrm{GL}_1 \iff \left(\sqrt{2} < x \leqslant x_* \& 0 < a < a_1(x)\right), \tag{12}$$

where $a_1(x) := \frac{x}{3} - \frac{1}{3}\sqrt{6 - 2x^2}$. Since $\sqrt{2} < u < v < \sqrt{3}$ on GL₁ (where again u and v are defined by (4) and (5)), one has

$$K(a,x) = k(a,x) := k_{\rm GL}(a,x) := g(u) + g(v) - 2g(x) \quad \forall (a,x) \in {\rm GL}_1.$$
(13)

For $(a, x) \in R$, let

$$(D_a k)(a, x) := \frac{\partial k}{\partial a}(a, x) \cdot \frac{125 \overline{\Phi}(\sqrt{2}) \left(1 - a^2\right)^{3/2}}{(1 + ax)(251 + v)\varphi(v)};$$
(14)

$$(D_{a,a}k)(s,x) := \frac{\partial (D_ak)}{\partial a}(a,x) \cdot (1-a^2)^3 (1+ax)^2 (251+v)^2 e^{-\frac{2ax}{1-a^2}}, \quad (15)$$

where $\sqrt{1-s^2}$ is substituted for *a* in the right-hand side of (15). Then $(D_{a,a}k)(s,x)$ is a polynomial in *s* and *x*. Using again the Mathematica command Reduce, namely

Reduce[daak[s,x]>0 && Sqrt[2]<x<Sqrt[3] && 0<s<1], (16)

where daak[s,x] stands for $(D_{a,a}k)(s,x)$, one sees that for every $(s,x) \in R$

$$(D_{a,a}k)(s,x) > 0 \iff \left(\sqrt{2} < x < x_{**} \& s_{*,1}(x) < s < s_{*,2}(x)\right),$$
(17)

where x_{**} is a certain number between $\sqrt{2}$ and $\sqrt{3}$, and $s_{*,1}$ and $s_{*,2}$ are certain functions. (In fact, $x_{**} \approx 1.678696$ is a root of a certain polynomial of degree 32 and, for each $x \in (\sqrt{2}, x_{**})$, the values $s_{*,1}(x)$ and $s_{*,2}(x)$ are two of the roots *s* of the polynomial $(D_{a,a}k)(s,x)$.) Next, Reduce [daak[95/100,x]<=0 && Sqrt[2]<x<=xx] produces False; here xx stands for x_* – recall the definition of GL₁ and (7); that is, $(D_{a,a}k)(\frac{95}{100}, x) > 0 \quad \forall x \in (\sqrt{2}, x_*]$. Hence and in view of (17),

$$s_{*,1}(x) < \frac{95}{100} < s_{*,2}(x) \quad \forall x \in (\sqrt{2}, x_*].$$
 (18)

On the other hand, setting

$$s_1(x) := \sqrt{1 - a_1(x)^2} \tag{19}$$

with a_1 as in (12), one has $s_1 > \frac{95}{100}$ on $(\sqrt{2}, x_*]$. Hence, by (18), $s_1 > s_{*,1}$ on $(\sqrt{2}, x_*]$. So, in view of (17), the sign pattern of $(D_{a,a}k)(s, x)$ in $s \in (s_1(x), 1)$ is - or +-, depending on whether $s_1(x) \ge s_{*,2}(x)$ or not, for each $x \in (\sqrt{2}, x_*]$. Now (15) and (19) imply that the sign pattern of $\frac{\partial (D_a k)}{\partial a}(a, x)$ in $a \in (0, a_1(x))$ is - or -+, so that $(D_a k)(a, x) \searrow$ or $\searrow \nearrow$ in $a \in (0, a_1(x))$, for each $x \in (\sqrt{2}, x_*]$. Also, $(D_a k)(0, x) = 0$ for all $x \in \mathbb{R}$. So, the sign pattern of $(D_a k)(a, x)$ in $a \in (0, a_1(x))$ is - or -+, for each $x \in (\sqrt{2}, x_*]$; in view of (14), $\frac{\partial k}{\partial a}(a, x)$ has the same sign pattern. Thus, $k(a, x) \searrow$ or $\searrow \nearrow$ in $a \in (0, a_1(x))$, for each $x \in (\sqrt{2}, x_*]$. Recalling (12), we complete the proof of Lemma 8.

Proof of Lemma 9 (GL_2). This proof is similar to that of Lemma 8. Here one has

$$(a,x) \in \mathrm{GL}_2 \iff \left(x_* < x < \sqrt{3} \& 0 < a < a_2(x)\right), \tag{20}$$

where $a_2(x) := -\frac{x}{4} + \frac{1}{4}\sqrt{12 - 3x^2}$. Relation (13) holds here, and we retain definitions (14) and (15). Definition (19) is replaced here by

$$s_2(x) := \sqrt{1 - a_2(x)^2}.$$
 (21)

Letting

$$s_* := s_2(x_*) = \frac{1}{12} \sqrt{\frac{1728 + 384\sqrt{6}}{19}} \approx 0.98761,$$

one can see that

$$s_2(x) \ge s_* \quad \forall x \in (x_*, \sqrt{3}).$$

$$(22)$$

On the other hand, using instead of (16) the Mathematica command

Reduce[daak[s,x]>0 && xx<x<Sqrt[3] && ss<s<1, Quartics->True],

where xx stands for x_* and ss stands for s_* , one sees that

$$\left((D_{a,a}k)(s,x) > 0 \& x_* < x < \sqrt{3} \& s_* < s < 1 \right)$$

$$\iff \left(x_* < x < x_{***} \& s_* < s < s_{*,2}(x) \right), \quad (23)$$

where $x_{***} \approx 1.678694$ is a root of a certain polynomial of degree 20 and $s_{*,2}(x)$ is the same root in s of the polynomial $(D_{a,a}k)(s,x)$ as $s_{*,2}(x)$ in (17). Hence, in view of (23) and (22), the sign pattern of $(D_{a,a}k)(s,x)$ in $s \in (s_2(x), 1)$ is - or +-, for each $x \in (x_*, \sqrt{3})$.

Now (15) and (21) imply that the sign pattern of $\frac{\partial (D_a k)}{\partial a}(a, x)$ in $a \in (0, a_2(x))$ is - or -+, so that $(D_a k)(a, x) \searrow$ or $\searrow \nearrow$ in $a \in (0, a_2(x))$, for each $x \in (x_*, \sqrt{3})$. Also, $(D_a k)(0, x) = 0$ for all $x \in \mathbb{R}$. So, the sign pattern of $(D_a k)(a, x)$ in $a \in (0, a_2(x))$ is - or -+, for each $x \in (x_*, \sqrt{3})$; in view of (14), $\frac{\partial k}{\partial a}(a, x)$ has the same sign pattern. Thus, $k(a, x) \searrow$ or $\searrow \nearrow$ in $a \in (0, a_2(x))$, for each $x \in (x_*, \sqrt{3})$. Recalling (20), we complete the proof of Lemma 9.

Proof of Lemma 10 (GG₁). Expressing x and v in view of (4) and (5) in terms of a and u as $\tilde{x}(a, u) := \sqrt{1 - a^2} u + a$ and $\tilde{v}(a, u) := u + \frac{2a}{\sqrt{1 - a^2}}$, respectively, and taking into account that $u < \sqrt{3}$ on GG₁, one has

$$K(a, x) = k(a, u) := k_{\text{GG}_1}(a, u) := g(u) + h(\tilde{v}(a, u)) - 2g(\tilde{x}(a, u)) \quad \forall (a, x) \in \text{GG}_1$$
(24)

For $(a, u) \in R$, let

$$d(a, u) := \frac{\partial k}{\partial a}(a, u) \cdot 125 \,\overline{\Phi}(\sqrt{2}) \,(1 - a^2)^{3/2} / \varphi(\tilde{v}(a, u)); \tag{25}$$
$$d_a(a, u) := \frac{\partial d}{\partial a}(a, u) \cdot (1 - a^2) \,\frac{\varphi(\tilde{v}(a, u))}{\varphi(\tilde{x}(a, u))};$$
$$d_u(a, u) := \frac{\partial d}{\partial u}(a, u) \cdot \frac{\varphi(\tilde{v}(a, u))}{\varphi(\tilde{x}(a, u))\sqrt{1 - a^2}}.$$

Then $d_a(a, u)$ and $d_u(a, u)$ are polynomials in a, u, and $\sqrt{1-a^2}$. Using

where da[a,u] and du[a,u] stand for $d_a(a, u)$ and $d_u(a, u)$, one sees that the system of equations $d_a(a, u) = 0 = d_u(a, u)$ has a unique solution $(a_*, u_*) \approx (0.11918, 1.57770)$ in $(a, u) \in (0, \frac{1}{\sqrt{3}}) \times (\sqrt{2}, \sqrt{3})$, and $d(a_*, u_*) \approx -0.44 < 0$.

Let us consider next the values of d on the boundary of the rectangle $(0, \frac{1}{\sqrt{3}}) \times (\sqrt{2}, \sqrt{3})$.

First, d(0, u) is an increasing affine function of u, and $d(0, \sqrt{3}) \approx -0.4 < 0$. Hence, $d(0, u) < 0 \ \forall u \in (\sqrt{2}, \sqrt{3})$.

Second,

$$\frac{27}{2 e^{(5+4\sqrt{2}u+u^2)/6}} \frac{\partial d}{\partial u} \left(\frac{1}{\sqrt{3}}, u\right)$$
$$= -\sqrt{2}u^3 - \left(3 + 251\sqrt{3}\right)u^2 - \sqrt{2}\left(3 + 251\sqrt{3}\right)u + 251\sqrt{3} + 7$$

is decreasing in u and takes on value $-9 - 753\sqrt{3} < 0$ at $u = \sqrt{2}$, so that $\frac{\partial d}{\partial u}(\frac{1}{\sqrt{3}}, u) < 0$ for $u \in (\sqrt{2}, \sqrt{3})$ and $d(\frac{1}{\sqrt{3}}, u) \searrow$ in $u \in (\sqrt{2}, \sqrt{3})$. Moreover, $d(\frac{1}{\sqrt{3}}, \sqrt{2}) < 0$. Thus, $d(\frac{1}{\sqrt{3}}, u) < 0 \ \forall u \in (\sqrt{2}, \sqrt{3})$. Third.

$$(1-a^2) \frac{\varphi(\tilde{v}(a,\sqrt{2}))}{\varphi(\tilde{x}(a,\sqrt{2}))} \frac{\partial d}{\partial a}(a,\sqrt{2}) = p_1(a) + \sqrt{1-a^2} p_2(a),$$

where $p_1(a)$ and $p_2(a)$ are certain polynomials in a. Therefore, the roots a of $\frac{\partial d}{\partial a}(a,\sqrt{2})$ are among the roots of the polynomial $p_{1,2}(a) := p_1(a)^2 - (1 - a^2)p_2(a)^2$, which has exactly two roots $a \in (0, \frac{1}{\sqrt{3}})$. Of the latter roots, one

is not a root of $\frac{\partial d}{\partial a}(a,\sqrt{2})$. Also, $\frac{\partial d}{\partial a}(0,\sqrt{2}) = 1 > 0$ and $\frac{\partial d}{\partial a}(\frac{1}{\sqrt{3}},\sqrt{2}) < 0$. Hence, $\frac{\partial d}{\partial a}(a,\sqrt{2})$ has exactly one root, $a_* \approx 0.2224$, in $a \in (0, \frac{1}{\sqrt{3}})$ and, moreover, $d(a,\sqrt{2}) \nearrow$ in $a \in (0,a_*]$ and $d(a,\sqrt{2}) \searrow$ in $a \in [a_*,\frac{1}{\sqrt{3}})$. Besides, $d(a_*,\sqrt{2}) \approx -0.088 < 0$. Thus, $d(a,\sqrt{2}) < 0 \ \forall a \in (0,\frac{1}{\sqrt{3}})$.

Fourth (very similar to third),

$$(1-a^2) \frac{\varphi(\tilde{v}(a,\sqrt{3}))}{\varphi(\tilde{x}(a,\sqrt{3}))} \frac{\partial d}{\partial a}(a,\sqrt{3}) = p_1(a) + \sqrt{1-a^2} p_2(a),$$

where $p_1(a)$ and $p_2(a)$ are certain polynomials in a, different from the polynomials $p_1(a)$ and $p_2(a)$ in the previous paragraph. Therefore, the roots a of $\frac{\partial d}{\partial a}(a,\sqrt{3})$ are among the roots of the polynomial $p_{1,2}(a) := p_1(a)^2 - (1-a^2)p_2(a)^2$, which has exactly two roots $a \in (0, \frac{1}{\sqrt{3}})$. Of the latter roots, one is not a root of $\frac{\partial d}{\partial a}(a,\sqrt{3})$. Also, $\frac{\partial d}{\partial a}(0,\sqrt{3}) = 1 > 0$ and $\frac{\partial d}{\partial a}(\frac{1}{\sqrt{3}},\sqrt{3}) < 0$. Hence, $\frac{\partial d}{\partial a}(a,\sqrt{3})$ has exactly one root, $a_* \approx 0.06651$, in $a \in (0, \frac{1}{\sqrt{3}})$ and, moreover, $d(a,\sqrt{3}) \nearrow$ in $a \in (0, a_*]$ and $d(a,\sqrt{3}) \searrow$ in $a \in [a_*, \frac{1}{\sqrt{3}})$. Besides, $d(a_*,\sqrt{3}) \approx -0.358 < 0$. Thus, $d(a,\sqrt{3}) < 0 \ \forall a \in (0, \frac{1}{\sqrt{3}})$.

We conclude that $d(a, u) < 0 \ \forall (a, u) \in [0, \frac{1}{\sqrt{3}}] \times [\sqrt{2}, \sqrt{3}]$. By (25), the same holds for $\frac{\partial k}{\partial a}(a, u)$. It remains to recall (24) and note that

$$(a,x) \in \mathrm{GG}_1 \implies \left(a,u(a,x)\right) \in \left(0,\frac{1}{\sqrt{3}}\right) \times \left(\sqrt{2},\sqrt{3}\right).$$

Proof of Lemma 11 (GG_2). In view of Lemma 2,

$$K(a,x) = (g \wedge h)(u(a,x)) + h(v(a,x)) - 2g(x) \quad \forall (a,x) \in \mathrm{GG}_2.$$
(26)

One has $\frac{\partial u}{\partial a} > 0$ on GG_2 and $\frac{\partial v}{\partial a} > 0$ on R. Since $g = \frac{c_1}{250} \left(251\overline{\Phi} + \varphi \right)$ and $h = c_2 \overline{\Phi}$ are decreasing on $[0, \infty)$, we conclude that K(a, x) is decreasing in a on GG_2 .

Proof of Lemma 12 (GE). One has

$$(a, x) \in GE \iff \left(x_* < x < \sqrt{3} \& a = a_2(x) := \frac{1}{4}\sqrt{12 - 3x^2} - \frac{x}{4}\right),$$

where, as before, x_* is defined by (7). Therefore, for all $(a, x) \in GE$

$$K(a, x) = k(x) := k_{\text{GE}}(x) := K(a_2(x), x) = g(u(a_2(x), x)) + g(\sqrt{3}) - 2g(x),$$

and it suffices to show that $k \leq 0$ on $[x_*, \sqrt{3}]$. For $x \in [x_*, \sqrt{3}]$, let

$$k_1(x) := \frac{k'(x)\overline{\Phi}(\sqrt{2})}{\varphi(x)(251+x)};$$

$$k_2(x) := k'_1(x) \cdot \frac{125(x+251)^2(4-x^2)^{3/2}\rho(x)^{13/2}}{16\sqrt{2}\varphi(u(a_2(x),x))/\varphi(x)},$$

imsart ver. 2005/10/19 file: arxiv.tex date: October 29, 2018

where $\rho(x) := x^2 + x\sqrt{3}\sqrt{4 - x^2} + 2$. Then $k_2(x) = p_1(x) + \sqrt{\rho(x)} p_2(x)$, where $p_1(x)$ and $p_2(x)$ are some polynomials in x and $\sqrt{4 - x^2}$. Hence, the roots of $k_2(x)$ are among the roots of

$$p_{1,2}(x) := p_1(x)^2 - \rho(x)p_2(x)^2 = p_{1,2,1}(x) + \sqrt{4 - x^2} p_{1,2,2}(x)$$

where $p_{1,2,1}(x)$ and $p_{1,2,2}(x)$ are some polynomials in x. Hence, the roots of $k_2(x)$ are among the roots of

$$\frac{p_{1,2,1}(x)^2 - (4 - x^2)p_{1,2,2}(x)^2}{1024 (x^2 - 1)^{14}},$$

which is a polynomial of degree 32 and has exactly one root in $[x_*, \sqrt{3}]$. Also, $k_2(x_*) \approx 1.39 \times 10^7 > 0$ and $k_2(\sqrt{3}) \approx -2.07 \times 10^6 < 0$. Therefore, k_2 and hence k'_1 have the sign pattern +- on $[x_*, \sqrt{3}]$. Next, $k_1(x_*) \approx -4.8494 < 0$ and $k_1(\sqrt{3}) = 0$, so that k_1 and hence k' have the sign pattern -+ on $[x_*, \sqrt{3}]$. It follows that k does not have a local maximum on $(x_*, \sqrt{3})$. At that, $k(x_*) \approx -3.0133 \times 10^{-6} < 0$ and $k_1(\sqrt{3}) = 0$. Thus, $k \leq 0$ on $[x_*, \sqrt{3}]$.

*Proof of Lemma 13 (*ELe). This proof is very similar to that of Lemma 12 (GE). One has

$$(a, x) \in \text{ELe} \iff \left(\sqrt{2} < x \leqslant x_* \& a = a_1(x) := \frac{x}{3} - \frac{1}{3}\sqrt{6 - 2x^2}\right),$$
 (27)

where, as before, x_* is defined by (7). Therefore, for all $(a, x) \in LLe$

$$K(a,x) = k(x) := k_{\text{ELe}}(x) := K(a_1(x), x) = g(\sqrt{2}) + g(v(a_1(x), x)) - 2g(x),$$

and it suffices to show that $k \leq 0$ on $[\sqrt{2}, x_*]$. For $x \in [\sqrt{2}, x_*]$, let

$$k_1(x) := \frac{500 \,\overline{\Phi}(\sqrt{2}) \,k'(x)}{\varphi(x)(251+x)};$$

$$k_2(x) := k_1'(x) \cdot \frac{\sqrt{2}(x+251)^2 \left(3-x^2\right)^{3/2} \rho(x)^{13/2}}{9\varphi(v(a_1(x),x))/\varphi(x)},$$

where $\rho(x) := x^2 + 2x\sqrt{2}\sqrt{3 - x^2} + 3$. Then $k_2(x) = p_1(x) + \sqrt{\rho(x)} p_2(x)$, where $p_1(x)$ and $p_2(x)$ are some polynomials in x and $\sqrt{3 - x^2}$. Hence, the roots of $k_2(x)$ are among the roots of

$$p_{1,2}(x) := p_1(x)^2 - \rho(x) p_2(x)^2 = p_{1,2,1}(x) + \sqrt{3 - x^2} p_{1,2,2}(x),$$

where $p_{1,2,1}(x)$ and $p_{1,2,2}(x)$ are some polynomials in x. Hence, the roots of $k_2(x)$ are among the roots of

$$\frac{p_{1,2,1}(x)^2 - (3-x^2)p_{1,2,2}(x)^2}{125524238436(x^2-1)^{14}},$$

which is a polynomial of degree 32 and has exactly one root in $[\sqrt{2}, x_*]$. Also, $k_2(\sqrt{2}) \approx -6.32 \times 10^7 < 0$ and $k_2(x_*) \approx 1.06 \times 10^8 > 0$. Therefore, k_2 and hence k'_1 have the sign pattern -+ on $[\sqrt{2}, x_*]$. Next, $k_1(\sqrt{2}) = 0$ and $k_1(x_*) \approx$ 0.000426 > 0, so that k_1 and hence k' have the sign pattern -+ on $[\sqrt{2}, x_*]$. It follows that k does not have a local maximum on $(\sqrt{2}, x_*)$. At that, $k(\sqrt{2}) = 0$ and $k(x_*) \approx -3.0133 \times 10^{-6} < 0$. Thus, $k \leq 0$ on $[\sqrt{2}, x_*]$.

Proof of Lemma 14 (EG₁). This proof is similar to that of Lemma 13. One has

$$(a, x) \in \mathrm{EG}_1 \iff \left(x_* < x < \sqrt{3} \& a = a_1(x) := \frac{x}{3} - \frac{1}{3}\sqrt{6 - 2x^2}\right).$$
 (28)

Therefore, for all $(a, x) \in EG_1$

$$K(a,x) = k(x) := k_{\mathrm{EG}_1}(x) := K(a_1(x), x) = g(\sqrt{2}) + h(v(a_1(x), x)) - 2g(x),$$

and it suffices to show that $k \leq 0$ on $[x_*, \sqrt{3}]$. For $x \in [x_*, \sqrt{3}]$, let

$$k_1(x) := \frac{500 \,\overline{\Phi}(\sqrt{2}) \,k'(x)}{\varphi(x)(251+x)};$$

$$k_2(x) := k_1'(x) \cdot \frac{\sqrt{2} \,(x+251)^2 \,(3-x^2)^{3/2} \,\rho(x)^{9/2}}{9 \,r(\sqrt{3}) \varphi\big(v(a_1(x),x)\big)/\varphi(x)},$$

where $\rho(x) := x^2 + 2x\sqrt{2}\sqrt{3-x^2} + 3$. Then $k_2(x) = p_1(x) + \sqrt{3-x^2}p_2(x)$, where $p_1(x)$ and $p_2(x)$ are some polynomials in x. Hence, the roots of $k_2(x)$ are among the roots of

$$p_{1,2}(x) := \frac{p_1(x)^2 - (3 - x^2)p_2(x)^2}{4374\left(1 + \frac{251}{r(\sqrt{3})}\right)^2 (x^2 - 1)^4},$$

which is a polynomial of degree 14 and has exactly one root in $[x_*, \sqrt{3}], x_{\#} \approx$ 1.6012. Also, $k_2(x_*) \approx 1.1722 \times 10^6 > 0$ and $k_2(\sqrt{3}) \approx -3.8778 \times 10^7 < 0$. Therefore, k_2 and hence k'_1 have the sign pattern +- on $[x_*, \sqrt{3}]$, so that $\max_{[x_*,\sqrt{3}]} k_1 = k_1(x_{\#}) \approx -0.00034907 < 0$. It follows that k' < 0 and hence $k \searrow$ on $[x_*,\sqrt{3}]$. At that, $k(x_*) \approx -3.0133 \times 10^{-6} < 0$. Thus, $k \leq 0$ on $[x_*, \sqrt{3}].$

Proof of Lemma 15 (EG_2). One has

$$(a, x) \in \mathrm{EG}_2 \iff \left(\sqrt{2} < x < \sqrt{3} \& a = a_2(x) := \frac{x}{3} + \frac{1}{3}\sqrt{6 - 2x^2}\right).$$
 (29)

Therefore, for all $(a, x) \in EG_2$

$$K(a,x) = k(x) := k_{\mathrm{EG}_2}(x) := K(a_2(x), x) = g(\sqrt{2}) + h(v(a_2(x), x)) - 2g(x),$$

and it suffices to show that $k \leq 0$ on $(\sqrt{2}, \sqrt{3})$.

For the functions a_1 (defined in (27) and (28)) and a_2 (defined in (29)), and for $x \in (\sqrt{2}, \sqrt{3})$, one has $a_2(x) \ge a_1(x)$; also, h(z) is decreasing in z and v(a, x)is increasing in a. Hence, $h(v(a_2(x), x)) \le h(v(a_1(x), x))$, so that $k_{\text{EG}_2} \le k_{\text{EG}_1}$ on $[x_*, \sqrt{3})$.

Similarly, in view of Lemma 2 one has $h(v(a_2(x), x)) \leq g(v(a_2(x), x)) \leq g(v(a_1(x), x)) \quad \forall x \in (\sqrt{2}, x_*]$, so that $k_{\text{EG}_2} \leq k_{\text{ELe}}$ on $(\sqrt{2}, x_*]$.

Now Lemma 15 follows, because it was shown in the proofs of Lemmas 13 and 14, respectively, that $k_{\text{ELe}} \leq 0$ on $[\sqrt{2}, x_*]$ and $k_{\text{EG}_1} \leq 0$ on $[x_*, \sqrt{3}]$.

Proof of Lemma 16 (A_1). This is trivial.

Proof of Lemma 17 (A₂). This is also trivial, in view of (8). \Box

Proof of Lemma 18 (X_{1,1}). On X_{1,1}, one has $u < \sqrt{2} \leq v$. Also, by Lemma 2, $g \geq h$ on $[\sqrt{3}, \infty)$. Therefore, for all $(a, x) \in X_{1,1}$

$$K(a,x) \leq k(a) := k_{X_{1,1}}(a) := \frac{1}{2u(a,\sqrt{2})^2} + g(v(a,\sqrt{2})) - 2g(\sqrt{2}), \quad (30)$$

and it suffices to show that $k \leq 0$ on $[0, \frac{1}{\sqrt{2}})$. For $a \in [0, \frac{1}{\sqrt{2}})$, let

$$k_1(a) := \frac{2000 \,\overline{\Phi}(\sqrt{2}) \, k'(a)}{\lambda(a)}, \quad \lambda(a) := \frac{1 - a\sqrt{2}}{\left(\sqrt{2} - a\right)^3} > 0;$$

$$k_2(a) := k_1'(a) \cdot \left(\sqrt{2} - a\right)^4 \left(1 - a^2\right)^4 \, \lambda(a)^2 / \left(\sqrt{2} \,\varphi\left(v(a,\sqrt{2})\right)\right).$$

Then $k_2(a) = \sqrt{1-a^2} p_1(a) + p_2(a)$, where $p_1(a)$ and $p_2(a)$ are some polynomials in a. Hence, the roots of $k_2(a)$ are among the roots of

$$p_{1,2}(a) := (1 - a^2)p_1(a)^2 - p_2(a)^2,$$

which is a polynomial of degree 12 and has exactly two roots in $[0, \frac{1}{\sqrt{2}})$. Of those two roots, one is not a root of $k_2(a)$, so that $k_2(a)$ has at most one root in $[0, \frac{1}{\sqrt{2}})$. Also, $k_2(0) = 251\sqrt{2} > 0$ and $k_2(\frac{1}{\sqrt{2}}) = -127 < 0$. Therefore, k_2 and hence k'_1 have the sign pattern +- on $[0, \frac{1}{\sqrt{2}}]$, so that $k_1 \nearrow \infty$ on $[0, \frac{1}{\sqrt{2}}]$. At that the values of k_1 at points 0, $\frac{6}{10}$, and $\frac{7}{10}$ are approximately -52 < 0, 48 > 0, and -344 < 0, respectively. Therefore, k_1 and hence k' have the sign pattern -+- on $[0, \frac{1}{\sqrt{2}})$ and, moreover, the only local maximum of k on $(0, \frac{1}{\sqrt{2}}]$ occurs only between $\frac{6}{10}$ and $\frac{7}{10}$; in fact, it occurs at $a \approx 0.67433$ and equals $\approx -0.00013578 < 0$. It remains to note that $k(0) \approx -0.0028660 < 0$.

Proof of Lemma 19 (X_{1,2}). This proof is similar to that of Lemma 11. In place of (26), here one still has relation (30) for all $(a, x) \in X_{1,2}$, since $u < \sqrt{2} \leq v$ on X_{1,2} as well. Since $u(a, \sqrt{2}) \nearrow$ and $v(a, \sqrt{2}) \nearrow$ in $a \in [\frac{1}{\sqrt{2}}, \frac{2\sqrt{2}}{3})$, one has $k \searrow$ on $[\frac{1}{\sqrt{2}}, \frac{2\sqrt{2}}{3})$, so that the maximum of k on $[\frac{1}{\sqrt{2}}, \frac{2\sqrt{2}}{3})$ equals $k(\frac{1}{\sqrt{2}})$, which is negative, in view of Lemma 18 and the continuity of k.

imsart ver. 2005/10/19 file: arxiv.tex date: October 29, 2018

Proof of Lemma 20 ($X_{1,3}$). This proof is similar to that of Lemma 19. In place of (30), here one has

$$K(a,x) \leq k(a) := k_{X_{1,3}}(a) := g(u(a,\sqrt{2})) + g(v(a,\sqrt{2})) - 2g(\sqrt{2}),$$

for all $(a, x) \in X_{1,3}$, since $u \ge \sqrt{2}$ and $v \ge \sqrt{2}$ on $X_{1,3}$. Since $k \searrow$ in $a \in [\frac{2\sqrt{2}}{3}, 1)$, the maximum of k on $[\frac{2\sqrt{2}}{3}, 1)$ equals $k(\frac{2\sqrt{2}}{3}) \approx -0.25287 < 0$.

Proof of Lemma 21 (X_2). This follows immediately from Lemma 5.

References

- BENTKUS, V. (2002) A remark on the inequalities of Bernstein, Prokhorov, Bennett, Hoeffding, and Talagrand. *Lithuanian Math. J.* 42, 262–269. MR1947624
- [2] BENTKUS, V. (2003) An inequality for tail probabilities of martingales with differences bounded from one side. J. Theoret. Probab. 16, 161–173. MR1956826
- [3] BENTKUS, V. (2004) On Hoeffding's inequalities. Ann. Probab. 32, 1650– 1673. MR2060313
- [4] BOBKOV, S. G.; GÖTZE, F.; HOUDRÉ, C. (2002) On Gaussian and Bernoulli covariance representations. *Bernoulli* 7, 439–451. MR1836739
- [5] COLLINS, G. E. (1975) Quantifier elimination for the elementary theory of real closed fields by cylindrical algebraic decomposition. *Lecture Notes In Computer Science*, 33, 134–183. MR0403962
- [6] EATON, M. L. (1974). A probability inequality for linear combinations of bounded random variables. Ann. Statist. 2, 609–614.
- [7] EDELMAN, D. (1990). An inequality of optimal order for the tail probabilities of the T statistic under symmetry. J. Amer. Statist. Assoc. 85, 120–122. MR1137357
- [8] EFRON, B. (1969). Student's t test under symmetry conditions. J. Amer. Statist. Assoc. 64, 1278–1302. MR0251826
- [9] GRAVERSEN, S. E.; PEŠKIR, G. (1998). Extremal problems in the maximal inequalities of Khintchine. *Math. Proc. Cambridge Philos. Soc.* 123, 169–177. MR1474873
- [10] LOJASIEWICZ, S. (1970). Sur les ensembles semi-analytiques. Actes du Congrès International des Mathématiciens. (Nice, 1970), Tome 2, pp. 237– 241. Gauthier-Villars, Paris. MR0425152
- [11] PINELIS, I. (1994). Extremal probabilistic problems and Hotelling's T^2 test under a symmetry condition. Ann. Statist. 22, 357–368. MR1272088
- [12] PINELIS, I. (1998). Optimal tail comparison based on comparison of moments. High dimensional probability (Oberwolfach, 1996), 297–314, Progr. Probab., 43, Birkhäuser, Basel. MR1652335
- [13] PINELIS, I. (1999). Fractional sums and integrals of r-concave tails and applications to comparison probability inequalities Advances in stochastic inequalities (Atlanta, GA, 1997), 149–168, Contemp. Math., 234, Amer. Math. Soc., Providence, RI. MR1694770

- [14] PINELIS, I. (2000). On exact maximal Khinchine inequalities. High dimensional probability, II (Seattle, WA, 1999), 49–63, Progr. Probab., 47, Birkhäuser Boston, Boston, MA. MR1857314
- [15] PINELIS, I. (2002). L'Hospital type rules for monotonicity: applications to probability inequalities for sums of bounded random variables. J. Inequal. Pure Appl. Math. 3, no. 1, Article 7, 9 pp. (electronic). MR1888922
- [16] PINELIS, I. (2005). Binomial upper bounds on generalized moments and tail probabilities of (super)martingales with differences bounded from above. *Preprint, to appear in the proceedings of the conference on high dimensional probability, New Mexico, 2005.* http://arxiv.org/abs/math.PR/0512301.
- [17] PINELIS, I. (2005). On normal domination of (super)martingales. *Preprint*, http://arxiv.org/abs/math.PR/0512382.
- [18] PINELIS, I. (2006). On l'Hospital-type rules for monotonicity. J. Inequal. Pure Appl. Math. 7, no. 2, to appear.
- [19] PINELIS, I. (2006). Exact inequalities for sums of asymmetric random variables, with applications. *Preprint*, http://arxiv.org/abs/math.PR/ 0602556.
- [20] PINELIS, I. (2006). On inequalities for sums of bounded random variables. Preprint, http://arxiv.org/abs/math.PR/ 0603030.
- [21] TARSKI, A. A. (1948). A Decision Method for Elementary Algebra and Geometry. RAND Corporation, Santa Monica, Calif. MR0028796