

RECURSIVE FORMULA FOR $\psi^g - \lambda_1\psi^{g-1} + \dots + (-1)^g\lambda_g$ IN $\overline{\mathcal{M}}_{g,1}$

D. ARCARA AND F. SATO

ABSTRACT. Mumford proved that $\psi^g - \lambda_1\psi^{g-1} + \dots + (-1)^g\lambda_g = 0$ in the Chow ring of $\mathcal{M}_{g,1}$ [Mum83]. We find an explicit recursive formula for $\psi^g - \lambda_1\psi^{g-1} + \dots + (-1)^g\lambda_g$ in the tautological ring of $\overline{\mathcal{M}}_{g,1}$ as a combination of classes supported on boundary strata.

1. INTRODUCTION

Mumford proved in [Mum83] that $\psi^g - \lambda_1\psi^{g-1} + \dots + (-1)^g\lambda_g = 0$ in the Chow ring of $\mathcal{M}_{g,1}$. Moreover, he showed that this class is supported on the boundary strata with a marked genus 0 component. Graber and Vakil proved in [GraVak05] that every codimension g class in the tautological ring of $\overline{\mathcal{M}}_{g,1}$ is supported on the boundary strata with at least one genus 0 component.

We complement these results by finding an explicit recursive formula for $\psi^g - \lambda_1\psi^{g-1} + \dots + (-1)^g\lambda_g$ in the tautological ring of $\overline{\mathcal{M}}_{g,1}$ as a combination of classes supported on boundary strata. It is clear from the formula being recursive that all the boundary strata have a genus 0 component in them, but it is not obvious from the formula that the marked point must be on a genus 0 component. We simplified the formula for $g < 5$ in Section 4, and checked that this is the case.

Theorem 1. *In the tautological ring of $\overline{\mathcal{M}}_{g,1}$,*

$$\sum_{i=0}^g (-1)^i \lambda_i \psi^{g-i} = \sum_{h=1}^g \left(1 - \frac{h}{g}\right) \iota_{h*}(c_h),$$

where

$$c_h := \sum_{i=0}^{g-1} (-1)^{h+i} \left[\left(\sum_{j=0}^h (-1)^j \lambda_j^0 \psi_0^{i-j} \right) \left(\sum_{j=0}^{g-h} (-1)^j \lambda_j^\infty \psi_\infty^{g-1-i-j} \right) \right],$$

ι_h is the natural boundary map

$$\iota_h : \overline{\mathcal{M}}_{h,2} \times \overline{\mathcal{M}}_{g-h,1} \longrightarrow \overline{\mathcal{M}}_{g,1},$$

ψ_0, ψ_∞ are descendents at the marked points glued by ι_h , and $\lambda^0, \lambda^\infty$ are the λ -classes on $\overline{\mathcal{M}}_{h,2}$ and $\overline{\mathcal{M}}_{g-h,1}$, respectively.

This formula is actually the first step of an algorithm which calculates each of the classes $\psi^g, \lambda_1\psi^{g-1}, \dots, \lambda_g$ in terms of classes supported on boundary strata. We want to single out the class $\psi^g - \lambda_1\psi^{g-1} + \dots + (-1)^g\lambda_g$, though, because it is the only class we found so far in the tautological ring of $\overline{\mathcal{M}}_{g,1}$ which has a nice recursive formula, and can therefore be easily calculated.

Acknowledgements. We would like to thank A. Bertram for many valuable discussions. This paper would have never been possible without his explanations, suggestions and guidance throughout the project.

2. VIRTUAL LOCALIZATION

The main tool we use to prove our theorems is the virtual localization theorem by Graber and Pandharipande [GraPan99].

Theorem (Virtual localization theorem). *Suppose $f: X \rightarrow X'$ is a \mathbb{C}^* -equivariant map of proper Deligne–Mumford quotient stacks with a \mathbb{C}^* -equivariant perfect obstruction theory. If $i': F' \hookrightarrow X'$ is a fixed substack and $c \in A_{\mathbb{C}^*}^*(X)$, let $f|_{F_i}: F_i \rightarrow F'$ be the restriction of f to each of the fixed substacks $F_i \subseteq f^{-1}(F')$. Then*

$$\sum_{F_i} f|_{F_i*} \frac{i_{F_i}^* c}{\epsilon_{\mathbb{C}^*}(F_i^{\text{vir}})} = \frac{i'^* f_* c}{\epsilon_{\mathbb{C}^*}(F'^{\text{vir}})},$$

where $i_{F_i}: F_i \rightarrow X$ and $\epsilon_{\mathbb{C}^*}(F^{\text{vir}})$ is the virtual equivariant Euler class of the “virtual” normal bundle F^{vir} .

Remark. The conditions in the theorem are satisfied for the Kontsevich–Manin spaces $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^m, d)$ of stable maps, and $\epsilon_{\mathbb{C}^*}(F^{\text{vir}})$ can be explicitly computed in terms of ψ and λ -classes [GraPan99] (see also [FabPan05]).

We define a \mathbb{C}^* -action on \mathbb{P}^1 by $a \cdot [x : y] = [x : ay]$ for $a \in \mathbb{C}^*$ and $[x : y] \in \mathbb{P}^1$. There are two fixed points, 0 and ∞ , and the torus acts with weight 1 on the tangent space at 0 and -1 on the tangent space at ∞ . This \mathbb{C}^* -action induces \mathbb{C}^* -actions on $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^1, d)$, and we shall consider the trivial \mathbb{C}^* -action on $\overline{\mathcal{M}}_{g,n}$.

3. PROOF OF THEOREM 1


We use virtual localization on the natural function $f: \overline{\mathcal{M}}_{g,3}(\mathbb{P}^1, 1) \rightarrow \overline{\mathcal{M}}_{g,3} \times (\mathbb{P}^1)^3$ defined by $f([g: (C, p_1, p_2, p_3) \rightarrow \mathbb{P}^1]) = ((C_{\text{stab}}, p_1, p_2, p_3), g(p_1), g(p_2), g(p_3))$. Consider the fixed locus

$$F' := \overline{\mathcal{M}}_{g,4} \times \{0\} \times \{\infty\} \times \{\infty\} \hookrightarrow \overline{\mathcal{M}}_{g,3} \times (\mathbb{P}^1)^3,$$

and apply the virtual localization theorem with $c = [1]^{\text{vir}}$ to obtain

$$\sum_{F_i} (f|_{F_i})_* \frac{[1]^{\text{vir}}}{\epsilon_{\mathbb{C}^*}(F_i^{\text{vir}})} = \frac{i'^* f_* [1]^{\text{vir}}}{t(-t)(-t)}.$$

There are $g+1$ fixed loci mapping to F' . One fixed locus has a marked point mapping to 0 and a curve in $\overline{\mathcal{M}}_{g,3}$ mapping to ∞ . We shall denote it by F_0 . Then there are g fixed loci which have a curve in $\overline{\mathcal{M}}_{h,2}$ mapping to 0 and a curve in $\overline{\mathcal{M}}_{g-h,3}$ mapping to ∞ (with $1 \leq h \leq g$). We shall denote these fixed loci by F_h . Note that $F_0 \simeq \overline{\mathcal{M}}_{g,3}$ and $F_h \simeq \overline{\mathcal{M}}_{h,2} \times \overline{\mathcal{M}}_{g-h,3}$ ($1 \leq h \leq g$).



Since $i'^* f_* [1]^{\text{vir}}$ is a polynomial in t , the sum of the contributions from the coefficient of t^{-4} on each fixed locus is 0. Call this contribution a_{-4} . We have that $\pi_{1,*}(\pi_{2,*}(a_{-4} \cdot \psi_3)) = 0$. We now calculate the contribution to the left hand side one fixed locus at the time.

- For F_0 , we obtain

$$\frac{[1]^{\text{vir}}}{\epsilon_{\mathbb{C}^*}(F_0^{\text{vir}})} = \frac{1}{t} \cdot (-1)^g \frac{t^g + \lambda_1 t^{g-1} + \dots + \lambda_g}{-t(-t - \psi_\infty)},$$

and the coefficient of t^{-4} is $-(\psi_\infty^{g+1} - \lambda_1 \psi_\infty^g + \dots + (-1)^g \lambda_g \psi_\infty)$. Under the isomorphism $F_0 \simeq \overline{\mathcal{M}}_{g,3}$, ψ_∞ gets identified with ψ_1 , and the contribution is therefore

$$-(\psi_1^{g+1} - \lambda_1 \psi_1^g + \dots + (-1)^g \lambda_g \psi_1).$$

- For F_h ($1 \leq h \leq g$), we obtain

$$\frac{[1]^{\text{vir}}}{\epsilon_{\mathbb{C}^*}(F_h^{\text{vir}})} = \frac{t^h - \lambda_1^0 t^{h-1} + \dots + (-1)^h \lambda_h^0}{t(t - \psi_0)} \cdot (-1)^{g-h} \frac{t^{g-h} + \lambda_1^\infty t^{g-h-1} + \dots + \lambda_{g-h}^\infty}{-t(-t - \psi_\infty)},$$

and the coefficient of t^{-4} is¹

$$c'_h := \sum_{i=0}^g (-1)^{h+i} \left[\left(\sum_{j=0}^h (-1)^j \lambda_j^0 \psi_0^{i-j} \right) \left(\sum_{j=0}^{g-h} (-1)^j \lambda_j^\infty \psi_\infty^{g-i-j} \right) \right].$$

This is a class of codimension g in $\overline{\mathcal{M}}_{h,2} \times \overline{\mathcal{M}}_{g-h,3}$ which maps to the codimension $g+1$ class $\iota_{h*}(c'_h)$ in $\overline{\mathcal{M}}_{g,3}$ under $(f|_{F_h})_*$.

To summarize, we obtain that

$$-(\psi_1^{g+1} - \lambda_1 \psi_1^g + \dots + (-1)^g \lambda_g \psi_1) + \sum_{h=1}^g \iota_{h*}(c'_h) = 0$$

in $\overline{\mathcal{M}}_{g,3}$. The first step is now to multiply by ψ_3 and push-forward to $\overline{\mathcal{M}}_{g,2}$.

- If $h = 0$, we obtain $-2g(\psi_1^{g+1} - \lambda_1 \psi_1^g + \dots + (-1)^g \lambda_g \psi_1)$ in $\overline{\mathcal{M}}_{g,2}$.
- If $1 \leq h < g$, note that, since the third marked point is on the curve at ∞ , we are really multiplying by ψ_3 in $\overline{\mathcal{M}}_{g-h,3}$ and pushing-forward to $\overline{\mathcal{M}}_{g-h,2}$. We therefore obtain, by Dilaton, the class $2(g-h)\iota_{h*}(c'_h)$, which is a class of codimension $g+1$ in $\overline{\mathcal{M}}_{g,2}$.
- If $h = g$, then $\psi_3 = 0$ because it is a descendent at a marked point of a genus 0 curve with 3 markings (the curve mapping to ∞).

Let us now suppose that $h < g$. The second and last step is to push-forward this class via the map that forgets the second marked point.

- If $h = 0$, we obtain, by String, $-2g(\psi_1^g - \lambda_1 \psi_1^{g-1} + \dots + (-1)^g \lambda_g)$.
- If $1 \leq h < g$, we obtain, by String, the class $2(g-h)\iota_{h*}(c_h)$, where c_h is just c'_h with every power of ψ_∞ lowered by 1 (with the convention that $\psi_\infty^{-1} = 0$), i.e.,

$$c_h = \sum_{i=0}^{g-1} (-1)^{h+i} \left[\left(\sum_{j=0}^h (-1)^j \lambda_j^0 \psi_0^{i-j} \right) \left(\sum_{j=0}^{g-h} (-1)^j \lambda_j^\infty \psi_\infty^{g-1-i-j} \right) \right].$$

Putting it all together, we obtain that

$$-2g(\psi_1^g - \lambda_1 \psi_1^{g-1} + \dots + (-1)^g \lambda_g) + \sum_{h=1}^g 2(g-h)\iota_{h*}(c_h) = 0,$$

from which we can derive the formula of Theorem 1. □

¹Note that c'_h is the summation (with the appropriate sign) of all possible products of codimension g of a class on the curve mapping to 0 with a class on the curve mapping to ∞ .

Remarks. (I) By taking the coefficient of t^{-3-j} with $j > 1$, it is possible to find a similar formula for $\psi^{g+j-1} - \lambda_1\psi^{g+j-2} + \dots + (-1)^g\lambda_g\psi^{j-1}$ in terms of classes supported on boundary strata.

(II) In [GraVak05], Graber and Vakil proved that a codimension g class in the tautological ring of $\overline{\mathcal{M}}_{g,1}$ can be written as a sum of classes supported on boundary strata with at least one genus 0 component. By induction on g , it is easy to see that this is the case for our c_h classes.

(III) Using the same function f as above, but with the fixed locus $\overline{\mathcal{M}}_{g,2} \times \{0\}^2 \times \{\infty\}$ instead of $\overline{\mathcal{M}}_{g,2} \times \{0\} \times \{\infty\}^2$, it is possible to obtain the following tautological relation on $\overline{\mathcal{M}}_{g,1}$:

$$\sum_{h=1}^{g-1} (2h)\iota_{h*}(c_h) + (2g)\pi_* (\psi_2^{g+1} - \lambda_1\psi_2^g + \dots + (-1)^g\lambda_g\psi_2) = 0.$$

4. EXPLICIT FORMULAS FOR LOW GENUS

The formula of Theorem 1 can be simplified recursively, and we calculated the answer for low values of g . Note that these formulas were already known for $g = 1$ and $g = 2$, but they were unknown for higher g 's.

Genus 1: In $\overline{\mathcal{M}}_{1,1}$,

$$\begin{array}{c} \psi - \lambda_1 \\ \text{---} \\ \text{---} \\ \bullet \\ \text{---} \\ \text{---} \\ \text{---} \end{array} = 0.$$

Genus 2: In $\overline{\mathcal{M}}_{2,1}$,

$$\begin{array}{c} \psi^2 - \lambda_1\psi + \lambda_2 \\ \text{---} \\ \text{---} \\ \bullet \\ \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \bullet \\ \text{---} \\ \text{---} \\ \text{---} \end{array}$$

Genus 3: In $\overline{\mathcal{M}}_{3,1}$,

$$\begin{array}{c} \psi^3 - \lambda_1\psi^2 + \lambda_2\psi - \lambda_3 \\ \text{---} \\ \text{---} \\ \bullet \\ \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \psi - \lambda_1 \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \bullet \\ \text{---} \\ \text{---} \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \bullet \\ \text{---} \\ \text{---} \\ \text{---} \end{array}$$

Genus 4: In $\overline{\mathcal{M}}_{4,1}$,

$$\begin{array}{c} \psi^4 - \lambda_1\psi^3 + \lambda_2\psi^2 - \lambda_3\psi + \lambda_4 \\ \text{---} \\ \text{---} \\ \bullet \\ \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \psi^2 - \lambda_1\psi + \lambda_2 \\ \text{---} \\ \text{---} \\ \bullet \\ \text{---} \\ \text{---} \\ \text{---} \end{array} + \begin{array}{c} \psi - \lambda_1 \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} + \begin{array}{c} \psi - \lambda_1 \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} + \begin{array}{c} \psi - \lambda_1 \\ \text{---} \\ \text{---} \\ \bullet \\ \text{---} \\ \text{---} \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \bullet \\ \text{---} \\ \text{---} \\ \text{---} \end{array} - \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \bullet \\ \text{---} \\ \text{---} \\ \text{---} \end{array}$$

We also have calculated the formula for $\psi^5 - \lambda_1\psi^4 + \lambda_2\psi^3 - \lambda_3\psi^2 + \lambda_4\psi - \lambda_5$ in $\overline{\mathcal{M}}_{5,1}$. We do not write it here because it was calculated via a (possibly incorrect) computer program and because it is rather long. Note that non-integer coefficients do appear in genus 5.

REFERENCES

- [FabPan05] C. Faber, R. Pandharipande, Relative maps and tautological classes, *J. Eur. Math. Soc. (JEMS)* 7 (2005), no. 1, 13–49.
- [GraPan99] T. Graber, R. Pandharipande, Localization of virtual classes, *Invent. Math.* 135, 1999, no. 2, pp. 487–518.
- [GraVak05] T. Graber, R. Vakil, Relative virtual localization and vanishing of tautological classes on moduli spaces of curves, *Duke Math. J.* 130 (2005), no. 1, 1–37.
- [Mum83] D. Mumford, Towards an Enumerative Geometry of the Moduli Space of Curves, in Arithmetic and geometry, Vol. II, *Progr. Math.* 36, Birkhuser Boston, Boston, MA, 1983, pp. 271–328.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, 155 S. 1400 E., ROOM 233, SALT LAKE CITY, UT 84112-0090, USA

E-mail address: arcara@math.utah.edu

SCHOOL OF MATHEMATICS, KOREAN INSTITUTE FOR ADVANCED STUDY, CHEONGNYANGNI 2-DONG, DONGDAEMUN-GU, SEOUL 130-722, SOUTH KOREA

E-mail address: fumi@kias.re.kr