

A COHOMOLOGICAL CHARACTERIZATION OF LEIBNIZ CENTRAL EXTENSIONS OF LIE ALGEBRAS

NAIHONG HU*, YUFENG PEI, AND DONG LIU

ABSTRACT. Motivated by Pirashvili's spectral sequences on a Leibniz algebra, some notions such as symmetric invariant bilinear forms, dual space derivations, as well as the Cartan-Koszul homomorphism are connected together in this note to give a description of the second Leibniz cohomology groups with trivial coefficients of Lie algebras (as Leibniz objects), which leads to a concise approach to determining one-dimensional Leibniz central extensions of Lie algebras. As applications, we contain the discussions for some interesting classes of infinite-dimensional Lie algebras. In particular, we generalize Gao's Theorem in [9] and answer a question in [16].

1. INTRODUCTION

Leibniz algebras, introduced by Loday ([17]) in the context of cyclic homology, are non-antisymmetric generalizations of Lie algebras. There is a (co)homology theory for these algebraic objects, whose properties are similar to those of the classical Chevalley-Eilenberg cohomology theory for Lie algebras. Naturally, each Lie algebra is a Leibniz algebra by definition. Therefore it is interesting to study Leibniz (co)homology of Lie algebras, which may produce new invariants for Lie algebras. Lodder ([19]) obtained the Godbillon-Vey invariants for foliations by a Leibniz cohomology computation of a certain Lie algebra and mentioned how a Leibniz algebra arises naturally in vertex (operator) algebras. Besides this, some relationships with classical geometry have recently been discovered which could allow ones to investigate Leibniz (co)homology in noncommutative geometry and its physical interpretations (see [11, 19], etc.).

As we know, the central extension theory plays an important role in structure and representation theory of Lie algebras (see [1, 2, 3, 6, 7, 8, 13, 14, 15, 20, 22], etc.). Regard a Lie algebra as a Leibniz algebra, it is natural to ask how to determine its Leibniz central extensions and compare the differences between Leibniz and Lie central extensions in question. Loday-Pirashvili ([18]) shown that the Virasoro algebra is a universal central extension of the Witt algebra in both settings of Lie algebras and of Leibniz algebras. Partially motivated by this, Gao ([9]) determined the universal Leibniz central extensions for symmetrizable Kac-Moody algebras and gave an interesting distinction between affine and non-affine Kac-Moody algebras by virtue of the second Leibniz homology group. The similar question for some other classes of infinite dimensional Lie algebras was discussed in

1991 *Mathematics Subject Classification.* Primary 17A32, 17B56; Secondary 17B65.

Key words and phrases. Leibniz central extensions, Leibniz cohomology, symmetric invariant bilinear forms, dual space derivations.

*N.H., Corresponding author. This work is supported in part by the NNSF (Grant 10431040), the TRAPOYT and the FUDP from the MOE of China, the SRSTP from the STCSM.

[16, 25], etc. The arguments involved unfortunately relied on some technical and tedious computations.

The aim of this note is to give a direct way to consider the one-dimensional Leibniz central extensions of Lie algebras. Our motivation is the Pirashvili's long exact sequence ([19, 21]), which is an essential technique for the calculation of Leibniz (co)homology of a Lie algebra. We observe the fact that the symmetric invariant bilinear forms and Leibniz central extensions can be well-connected by the Cartan-Koszul homomorphism (see [14]) in the sequence. While in practice, it is much easier to treat the symmetric invariant bilinear forms on a Lie algebra than to directly determine its Leibniz central extensions. On the other hand, we also notice that determining Leibniz central extensions of a Lie algebra is equivalent to treating dual space derivations of this Lie algebra (cf. [7]), which seemed lacking in enough attention for the Leibniz cases in the literature. We thus combine these two observations together to derive a concise and natural approach to studying the Leibniz central extensions of Lie algebras, which avoids complicated computations when applied to some interesting infinite dimensional Lie algebras including the Witt algebra, the twisted Heisenberg-Virasoro algebra, the Lie algebra of differential operators over a Laurant polynomial ring, Lie algebras of degenerate Block type (Virasoro-like algebra and its q -analog) and Kac-Moody algebras, etc. Our method is more conceptual in the cohomological sense.

Some notions on Leibniz algebras from [17] is recalled in Section 2. In Section 3, we present a direct elementary proof for the exact sequence in question — around the description of Leibniz central extensions of Lie algebras, in terms of the symmetric invariant bilinear forms and the Cartan-Koszul homomorphism. We obtain a description of the second Leibniz cohomology group with trivial coefficients by dual space derivations (Theorem 3.5). Section 4 contains some applications of our method. We generalize Gao's main result ([9]) for the symmetrizable Kac-Moody algebras to the general cases (Theorem 4.18). In Section 5, we study the Leibniz central extensions of the quadratic Lie algebras, and construct a counterexample to address a question in [16].

2. PREREQUISITES ON LEIBNIZ ALGEBRAS

2.1. Leibniz algebra. Let \mathbb{K} be an algebraically closed field with $\text{char } \mathbb{K} = 0$.

Definition 2.1. A Leibniz algebra is a \mathbb{K} -module L with a bilinear map $[-, -] : L \times L \rightarrow L$ satisfying the Leibniz identity

$$[x, [y, z]] = [[x, y], z] - [[x, z], y], \quad \forall x, y, z \in L. \quad (1.1)$$

The center of L is defined to be $\{z \in L \mid [z, L] = [L, z] = 0\}$. L is called *perfect* if $[L, L] = L$. If, in addition, $[x, x] = 0, \forall x \in L$, the Leibniz identity is equivalent to the Jacobi identity. In particular, Lie algebras are examples of Leibniz algebras.

Definition 2.2. Let L be a Leibniz algebra over \mathbb{K} . M is called a representation of L if M is a \mathbb{K} -vector space equipped with two actions (left and right) of L , i.e., $[-, -] : L \times M \rightarrow M$ and $[-, -] : M \times L \rightarrow M$ satisfying

$$\begin{aligned} (MLL) \quad & [m, [x, y]] = [[m, x], y] - [[m, y], x], \\ (LML) \quad & [x, [m, y]] = [[x, m], y] - [[x, y], m], \\ (LLM) \quad & [x, [y, m]] = [[x, y], m] - [[x, m], y], \end{aligned} \quad (1.2)$$

for any $m \in M$ and $x, y \in L$.

2.2. Cohomology of Leibniz algebras. Let L be a Leibniz algebra over \mathbb{K} and M a representation of L . Denote $C^n(L, M) := \text{Hom}_{\mathbb{K}}(L^{\otimes n}, M)$, $n \geq 0$. The Loday coboundary map $d^n : C^n(L, M) \rightarrow C^{n+1}(L, M)$ is defined by

$$\begin{aligned} (d^n f)(x_1, \dots, x_{n+1}) &= [x_1, f(x_2 \otimes \dots \otimes x_{n+1})] \\ &+ \sum_{i=2}^{n+1} (-1)^i [f(x_1, \dots, \hat{x}_i, \dots, x_{n+1}), x_i] \\ &+ \sum_{1 \leq i < j \leq n+1} (-1)^{j+1} f(x_1, \dots, x_{i-1}, [x_i, x_j], x_{i+1}, \dots, \hat{x}_j, \dots, x_{n+1}). \end{aligned}$$

Clearly, $d^{n+1}d^n = 0$, $n \geq 0$. $(C^*(L, M), d)$ is a well-defined cochain complex, whose cohomology is called the cohomology of Leibniz algebra L with coefficients in the representation M :

$$\text{HL}^*(L, M) := \text{H}^*((C^*(L, M), d)).$$

Similarly, we have a chain complex $(C_*(L, M), d)$, whose homology is called the homology of Leibniz algebra L with coefficients in the representation M :

$$\text{HL}_*(L, M) := \text{H}_*((C_*(L, M), d)).$$

2.3. Cohomology of Lie algebras. Let \mathfrak{g} be a Lie algebra over \mathbb{K} and M a \mathfrak{g} -module. Denote the Chevalley-Eilenberg cochain complex by

$$(\Omega^*(\mathfrak{g}, M), \delta) := (\text{Hom}(\wedge^* \mathfrak{g}, M), \delta),$$

where δ is the Chevalley-Eilenberg coboundary map defined by

$$\begin{aligned} (\delta^n f)(x_1, \dots, x_{n+1}) &= \sum_{i=1}^{n+1} (-1)^{i+1} x_i \cdot f(x_1, \dots, \hat{x}_i, \dots, x_{n+1}) \\ &+ \sum_{1 \leq i < j \leq n+1} (-1)^{i+j} f([x_i, x_j], x_1, \dots, \hat{x}_i, \dots, \hat{x}_j, \dots, x_{n+1}). \end{aligned}$$

Then,

$$\text{H}^*(\mathfrak{g}, M) := \text{H}^*((\Omega^*(\mathfrak{g}, M), \delta)),$$

is called the cohomology of Lie algebra \mathfrak{g} with coefficients in the \mathfrak{g} -module M .

3. LEIBNIZ COHOMOLOGY OF LIE ALGEBRAS AND LEIBNIZ CENTRAL EXTENSIONS

3.1. Central extensions of Leibniz algebras. A central extension of L is a pair (\hat{L}, π) , where \hat{L} is a Leibniz algebra and $\pi : \hat{L} \rightarrow L$ is a surjective homomorphism whose kernel lies in the center of \hat{L} . The pair (\hat{L}, π) is a universal central extension of L if for every central extension (\tilde{L}, τ) of L , there is unique homomorphism $\psi : \hat{L} \rightarrow \tilde{L}$ for which $\tau \circ \psi = \pi$.

The following result is known.

Proposition 3.1. ([18]) *There exists a one-to-one correspondence between the set of equivalent classes of one-dimensional Leibniz central extensions of L by \mathbb{K} and the second Leibniz cohomology group $\text{HL}^2(L, \mathbb{K})$.*

3.2. Invariant symmetric bilinear forms. Let \mathfrak{g} be a Lie algebra over \mathbb{K} and M a \mathfrak{g} -module. Recall that a symmetric bilinear forms ϕ on \mathfrak{g} is called \mathfrak{g} -invariant if ϕ satisfies $\phi([x, y], z) = \phi(x, [y, z])$, $\forall x, y, z \in \mathfrak{g}$. Let $B(\mathfrak{g}, \mathbb{K})$ stand for the set of all \mathbb{K} -valued symmetric \mathfrak{g} -invariant bilinear forms on Lie algebra \mathfrak{g} .

The following exact sequence is due to Pirashvili ([21]), but we give an elementary proof here.

Proposition 3.2. *For any Lie algebra \mathfrak{g} , there holds the following exact sequence*

$$0 \longrightarrow H^2(\mathfrak{g}, \mathbb{K}) \xrightarrow{f} HL^2(\mathfrak{g}, \mathbb{K}) \xrightarrow{g} B(\mathfrak{g}, \mathbb{K}) \xrightarrow{h} H^3(\mathfrak{g}, \mathbb{K}),$$

where

- f is the natural embedding map.
- g is defined by $g(\alpha)(x, y) = \alpha(x, y) + \alpha(y, x)$, $\forall \alpha \in HL^2(\mathfrak{g}, \mathbb{K})$, $x, y \in \mathfrak{g}$.
- h is the Cartan-Koszul map ([14]) defined by $h(\alpha)(x, y, z) = \alpha([x, y], z)$, $\forall \alpha \in B(\mathfrak{g}, k)$, $x, y, z \in \mathfrak{g}$.

Proof. We shall prove

- (1) The map g is well-defined, that is, $g(\alpha) = g(\alpha + d^1\beta) \in B(\mathfrak{g}, \mathbb{K})$ for any $\alpha \in HL^2(\mathfrak{g}, \mathbb{K})$ and $\beta \in HL^1(\mathfrak{g}, \mathbb{K})$ since $g(d^1\beta) \equiv 0$ by definition. Clearly, $g(\alpha)$ is symmetric. Since, for any $x, y, z \in \mathfrak{g}$,

$$\begin{aligned} g(\alpha)([x, y], z) &= \alpha([x, y], z) + \alpha(z, [x, y]) \\ &= \alpha(x, [y, z]) + \alpha([x, z], y) + \alpha([z, x], y) - \alpha([z, y], x) \\ &= \alpha(x, [y, z]) + \alpha([y, z], x) \\ &= g(\alpha)(x, [y, z]), \end{aligned}$$

$g(\alpha)$ is \mathfrak{g} -invariant.

- (2) $\text{Im}(f) = \ker(g)$. For any $\alpha \in H^2(\mathfrak{g}, \mathbb{K})$ and any $x, y \in \mathfrak{g}$, we have

$$g(f(\alpha))(x, y) = g(\alpha)(x, y) = \alpha(x, y) + \alpha(y, x) = 0,$$

since α is skew symmetric. Then $\text{Im}(f) \subseteq \ker(g)$. On the other hand, for any $\beta \in \ker(g)$, we have $g(\beta)(x, y) = 0$, which implies that $\alpha(x, y) + \alpha(y, x) = 0$ for any $x, y \in \mathfrak{g}$. Then α is a skew symmetric Leibniz 2-cocycle, namely, a Lie 2-cocycle in $H^2(\mathfrak{g}, k)$. Hence, we have $\ker(g) \subseteq \text{Im}(f)$.

- (3) $\text{Im}(g) = \ker(h)$. Now, $\forall \alpha \in HL^2(\mathfrak{g}, \mathbb{K})$, define a bilinear form α' on \mathfrak{g} by

$$\alpha'(x, y) = \alpha(y, x),$$

for any $x, y \in \mathfrak{g}$. Denote $\beta := \frac{\alpha + \alpha'}{2}$ and $\gamma := \frac{\alpha - \alpha'}{2}$. Then β is symmetric and γ is skew symmetric. Also, we have $\alpha = \beta + \gamma$.

Now, for any $x, y, z \in \mathfrak{g}$,

$$\begin{aligned} (h(g(\alpha)))(x, y, z) &= g(\alpha)([x, y], z) = \alpha([x, y], z) + \alpha(z, [x, y]) \\ &= \alpha([x, y], z) + \alpha([z, x], y) + \alpha([y, z], x) \\ &= \alpha(z, [x, y]) + \alpha(y, [z, x]) + \alpha(x, [y, z]) \\ &\quad - \alpha([x, y], z) - \alpha([z, x], y) - \alpha([y, z], x) \\ &= 2\gamma(z, [x, y]) + 2\gamma(y, [z, x]) + 2\gamma(x, [y, z]) \\ &= \delta^2(2\gamma)(x, y, z), \end{aligned}$$

which implies that $h(g(\alpha))$ is a 3-coboundary in $\Omega^3(\mathfrak{g}, \mathbb{K})$. Then $\text{Im}(g) \subseteq \ker(h)$. Conversely, for any $\omega \in \ker(h)$, there exists an element $\eta \in \Omega^2(\mathfrak{g}, \mathbb{K})$ such that $\delta^2(\eta) = h(\omega)$. Namely, for any $x, y, z \in \mathfrak{g}$, there holds

$$\eta(z, [x, y]) + \eta(y, [z, x]) + \eta(x, [y, z]) = \omega([x, y], z).$$

Then

$$\begin{aligned} \omega([x, y], z) + \eta([x, y], z) &= \omega(x, [y, z]) + \eta(x, [y, z]) + \omega([x, z], y) + \eta([x, z], y), \\ \text{since } \omega(x, [y, z]) + \omega([x, z], y) &= 0. \text{ Let } \rho = \frac{1}{2}(\omega + \eta). \text{ Then } g(\rho) = \omega \text{ and} \end{aligned}$$

$$\rho([x, y], z) = \rho(x, [y, z]) + \rho([x, z], y),$$

which means that $\rho \in \text{HL}^2(\mathfrak{g}, \mathbb{K})$. Hence $\ker(h) \subseteq \text{Im}(g)$.

We complete the proof. \square

Corollary 3.3.

$$\frac{\text{HL}^2(\mathfrak{g}, \mathbb{K})}{\text{H}^2(\mathfrak{g}, \mathbb{K})} = \ker(h).$$

In particular, $\text{HL}^2(\mathfrak{g}, \mathbb{K}) = \text{H}^2(\mathfrak{g}, \mathbb{K})$, $\iff \ker(h) = 0$.

Remark 3.4. Compared to our direct proof above, there is another argument for Proposition 3.2 in terms of spectral sequences, which is due to Pirashvili. Note that the natural embedding $\Omega^*(\mathfrak{g}, M) \hookrightarrow C^*(\mathfrak{g}, M)$ induces a short exact sequence in the category of cochain complexes of Lie algebra \mathfrak{g} :

$$0 \longrightarrow \Omega^*(\mathfrak{g}, M) \longrightarrow C^*(\mathfrak{g}, M) \longrightarrow C_{rel}^*(\mathfrak{g}, M)[2] \longrightarrow 0,$$

where

$$C_{rel}^*(\mathfrak{g}, M)[2] := \frac{C^*(\mathfrak{g}, M)}{\Omega^*(\mathfrak{g}, M)}$$

is the quotient cochain complex. One has the Pirashvili's long exact sequence below (see [19], [21]), which is the main tool to compare the Lie and the Leibniz cohomology of a Lie algebra in higher dimensions

$$0 \rightarrow \text{H}^2(\mathfrak{g}, M) \rightarrow \text{HL}^2(\mathfrak{g}, M) \rightarrow \text{H}_{rel}^0(\mathfrak{g}, M) \rightarrow \text{H}^3(\mathfrak{g}, M) \rightarrow \text{HL}^3(\mathfrak{g}, M) \rightarrow \dots$$

and isomorphisms $\text{HL}^i(\mathfrak{g}, M) = \text{H}^i(\mathfrak{g}, M)$, $i = 0, 1$.

Let $M = \mathbb{K}$. We have

$$0 \rightarrow \text{H}^2(\mathfrak{g}, \mathbb{K}) \rightarrow \text{HL}^2(\mathfrak{g}, \mathbb{K}) \rightarrow \text{H}_{rel}^0(\mathfrak{g}, \mathbb{K}) \rightarrow \text{H}^3(\mathfrak{g}, \mathbb{K}) \rightarrow \text{HL}^3(\mathfrak{g}, \mathbb{K}) \rightarrow \dots$$

Then, as a corollary of the spectral sequences, one has the Pirashvili's claim (see [21]) : $\text{H}_{rel}^0(\mathfrak{g}, \mathbb{K}) = \text{B}(\mathfrak{g}, \mathbb{K})$.

3.3. Dual space derivations. Let \mathfrak{g} be a Lie algebra over \mathbb{K} and M a \mathfrak{g} -module. Recall that a linear map $d : \mathfrak{g} \rightarrow M$ is called a derivation if

$$d([x, y]) = x \cdot d(y) - y \cdot d(x), \quad \forall x, y \in \mathfrak{g}.$$

The derivations of the form $x \mapsto x \cdot m$ for some $m \in M$ are called inner derivations. $\text{Der}(\mathfrak{g}, M)$ and $\text{Inn}(\mathfrak{g}, M)$ denote the spaces of derivations and inner derivations, respectively. It is clear that $\text{H}^1(\mathfrak{g}, M) = \text{Der}(\mathfrak{g}, M)/\text{Inn}(\mathfrak{g}, M)$ is the first cohomology group of \mathfrak{g} with coefficients in M .

Theorem 3.5. *For any Lie algebra \mathfrak{g} , there holds*

$$\mathrm{HL}^2(\mathfrak{g}, \mathbb{K}) = \mathrm{H}^1(\mathfrak{g}, \mathfrak{g}^*) = \mathrm{Der}(\mathfrak{g}, \mathfrak{g}^*) / \mathrm{Inn}(\mathfrak{g}, \mathfrak{g}^*),$$

where \mathfrak{g}^* is the dual \mathfrak{g} -module.

Proof. Let the map $\theta : \mathrm{H}^1(\mathfrak{g}, \mathfrak{g}^*) \rightarrow \mathrm{HL}^2(\mathfrak{g}, \mathbb{K})$ defined by $\theta(\alpha)(x, y) = \alpha(y)(x)$ for any $\alpha \in \mathrm{H}^1(\mathfrak{g}, \mathfrak{g}^*)$ and $x, y \in \mathfrak{g}$. Since

$$\begin{aligned} \theta(\alpha)(x, [y, z]) &= \alpha([y, z])(x) = (y \cdot \alpha(z))(x) - (z \cdot \alpha(y))(x) \\ &= \alpha(z)([x, y]) - \alpha(y)([x, z]) \\ &= \theta(\alpha)([x, y], z) + \theta(\alpha)([z, x], y), \end{aligned}$$

$\theta(\alpha) \in \mathrm{HL}^2(\mathfrak{g}, \mathbb{K})$. θ is well-defined. If $\theta(\alpha) = \bar{0}$ in $\mathrm{HL}^2(\mathfrak{g}, \mathbb{K})$, there exists an element β in $C^1(\mathfrak{g}, \mathbb{K})$ such that, for any $x, y \in \mathfrak{g}$, $d^1(\beta)(x, y) = \theta(\alpha)(x, y)$. Then

$$-\beta([x, y]) = \alpha(y)(x),$$

which implies α is a 1-coboundary in $\Omega^1(\mathfrak{g}, \mathfrak{g}^*)$, i.e. $\alpha = \bar{0} \in \mathrm{H}^1(\mathfrak{g}, \mathfrak{g}^*)$. Hence, θ is injective. On the other hand, for any $\beta \in \mathrm{HL}^2(\mathfrak{g}, \mathbb{K})$, we define a map $\alpha \in \mathrm{Hom}_{\mathbb{K}}(\mathfrak{g}, \mathfrak{g}^*)$ by $\alpha(x)(y) := \beta(y, x)$, for any $x, y \in \mathfrak{g}$. Since

$$\begin{aligned} \alpha([x, y])(z) &= \beta(z, [x, y]) = \beta([z, x], y) - \beta([z, y], x) = \alpha(y)([z, x]) - \alpha(x)([z, y]), \\ &= (x \cdot \alpha(x))(z) - (y \cdot \alpha(x))(z), \end{aligned}$$

for any $z \in \mathfrak{g}$, we get $\alpha([x, y]) = x \cdot \alpha(x) - y \cdot \alpha(x)$ for any $x, y \in \mathfrak{g}$, which means $\alpha \in \mathrm{H}^1(\mathfrak{g}, \mathfrak{g}^*)$. Hence, θ is surjective. \square

Denote $\mathrm{SDer}(\mathfrak{g}, \mathfrak{g}^*) := \{\phi \in \mathrm{Der}(\mathfrak{g}, \mathfrak{g}^*) \mid \phi(x)(y) + \phi(y)(x) = 0, \forall x, y \in \mathfrak{g}\}$. By Theorem 3.5 above and Proposition 1.3 (2) in [6], we have

Corollary 3.6. $\mathrm{HL}^2(\mathfrak{g}, \mathbb{K}) / \mathrm{H}^2(\mathfrak{g}, \mathbb{K}) = \mathrm{Der}(\mathfrak{g}, \mathfrak{g}^*) / \mathrm{SDer}(\mathfrak{g}, \mathfrak{g}^*) (\subset H_{rel}^0(\mathfrak{g}, \mathbb{K}))$.

Corollary 3.7. *If \mathfrak{g} is a finite dimensional Lie algebra over \mathbb{K} with a non-degenerate symmetric invariant bilinear form ψ , then $\mathrm{HL}^2(\mathfrak{g}, \mathbb{K}) = \mathrm{Der}(\mathfrak{g}, \mathfrak{g}) / \mathrm{Inn}(\mathfrak{g}, \mathfrak{g})$. If, in addition, \mathfrak{g} is simple, then $\mathrm{HL}^2(\mathfrak{g}, \mathbb{K}) = 0$.*

Proof. It follows from $\mathfrak{g} \simeq \mathfrak{g}^*$ as \mathfrak{g} -modules. \square

4. APPLICATIONS: LEIBNIZ CENTRAL EXTENSIONS OF SOME LIE ALGEBRAS

4.1. Lie algebras of Virasoro type. Let $\mathbb{K}[t, t^{-1}]$ be the Laurent polynomial algebra over \mathbb{K} and $\frac{d}{dt}$ be the differential operator on $\mathbb{K}[t, t^{-1}]$.

Definition 4.1. *Set $L_n = -t^{n+1} \frac{d}{dt}, n \in \mathbb{Z}$. The Witt algebra $\mathcal{W} = \bigoplus_{n \in \mathbb{Z}} \mathbb{K}L_n$ is defined by $[L_m, L_n] = (m - n)L_{m+n}$, $m, n \in \mathbb{Z}$.*

It is well-known that $\mathrm{H}^2(\mathcal{W}, \mathbb{K}) = \mathrm{H}_2(\mathcal{W}, \mathbb{K}) = \mathbb{K}\alpha$, where

$$\alpha(L_m, L_n) = \delta_{m+n}, 0 \frac{m^3 - m}{12}.$$

Definition 4.2. ([1]) *Set $L_n = -t^{n+1} \frac{d}{dt}, I_n = t^n, n \in \mathbb{Z}$. The Lie algebra*

$$\mathcal{H} = \bigoplus_{n \in \mathbb{Z}} \mathbb{C}L_n \bigoplus \bigoplus_{m \in \mathbb{Z}} \mathbb{C}I_m$$

is defined by $[L_m, L_n] = (m - n)L_{m+n}$, $[I_m, I_n] = 0$, $[L_m, I_n] = -nI_{m+n}$, $m, n \in \mathbb{Z}$.

In [1], authors proved that $\dim H^2(\mathcal{H}, \mathbb{K}) = \dim H_2(\mathcal{H}, \mathbb{K}) = 3$. In fact, as the universal central extension of \mathcal{H} , the twisted Heisenberg-Virasoro algebra H_{Vir} has a basis $\{I_m, L_m, C_I, C_L, C_{LI} \mid m \in \mathbb{Z}\}$ satisfying the following relations:

$$\begin{aligned} [I_m, I_n] &= n\delta_{m+n,0}C_I, \\ [L_m, L_n] &= (m-n)L_{m+n} + \delta_{m+n,0}\frac{1}{12}(m^3-m)C_L, \\ [L_m, I_n] &= -nI_{m+n} + \delta_{m+n,0}(m^2-m)C_{LI}, \\ [H_{Vir}, C_L] &= [H_{Vir}, C_I] = [H_{Vir}, C_{LI}] = 0. \end{aligned}$$

Definition 4.3. Set $D = t\frac{d}{dt}$. The Lie algebra of differential operators

$$\mathcal{D} = \bigoplus_{m \in \mathbb{Z}, n \in \mathbb{Z}_+} \mathbb{K}t^m D^n$$

is defined by $[t^m D^r, t^n D^s] = t^{m+n}((D+n)^r - (D+m)^s)$, $m, n \in \mathbb{Z}, r, s \in \mathbb{Z}_+$.

It is known [15] that $\dim H^2(\mathcal{D}, \mathbb{K}) = \dim H_2(\mathcal{D}, \mathbb{K}) = 1$ and the universal central extension of \mathcal{D} is a Lie algebra

$$\mathcal{W}_{1+\infty} = \bigoplus_{m \in \mathbb{Z}, n \in \mathbb{Z}_+} \mathbb{K}t^m D^n \bigoplus \mathbb{K}C$$

with relations: $[t^m D^r, t^n D^s] = t^{m+n}((D+n)^r - (D+m)^s) + \psi(t^m D^r, t^n D^s)C$, where $\psi(t^{m+r} D^r, t^{n+s} D^s) = \delta_{m+n,0}(-1)^r r! s! \binom{m+r}{r+s+1}$, for $m, n \in \mathbb{Z}, r, s \in \mathbb{Z}_+$.

Remark 4.4. It is clear that \mathcal{W} and \mathcal{H} are Lie subalgebras of \mathcal{D} .

Proposition 4.5. There is no non-trivial symmetric invariant bilinear form on Lie algebra \mathfrak{g} , where $\mathfrak{g} = \mathcal{W}, \mathcal{H}$, or \mathcal{D} .

Proof. Assume that f is a symmetric invariant bilinear form on \mathfrak{g} .

(1) $\mathfrak{g} = \mathcal{W}$.

Note that \mathcal{W} is generated as Lie algebra by the elements L_3, L_{-2} with

$$\begin{aligned} f(L_3, L_3) &= \frac{1}{3}f(L_3, [L_3, L_0]) = 0, \\ f(L_3, L_{-2}) &= \frac{1}{8}f(L_3, [L_3, L_{-5}]) = 0, \\ f(L_{-2}, L_{-2}) &= -\frac{1}{2}f(L_{-2}, [L_{-2}, L_0]) = 0. \end{aligned}$$

(2) $\mathfrak{g} = \mathcal{H}$.

Note that \mathcal{H} is generated as Lie algebra by the elements L_3, L_{-2} and I_1 with

$$\begin{aligned} f(L_m, L_n) &= 0, \quad n, m \in \mathbb{Z}, \\ f(I_1, I_1) &= f(I_1, [I_1, L_0]) = f([I_1, I_1], L_0) = 0, \\ f(I_1, L_3) &= -\frac{1}{2}f([I_{-2}, L_3], L_3) = -\frac{1}{2}f(I_{-2}, [L_3, L_3]) = 0, \\ f(I_1, L_{-2}) &= \frac{1}{3}f([I_3, L_{-2}], L_{-2}) = \frac{1}{3}f(I_3, [L_{-2}, L_{-2}]) = 0. \end{aligned}$$

(3) $\mathfrak{g} = \mathcal{D}$.

Note that \mathcal{D} can be generated as Lie algebra by t, t^{-1} and D^2 (see [23]).

By a direct computation, we have

$$t^2\left(\frac{d}{dt}\right)^2 = D^2 - D,$$

$$t^m\left(\frac{d}{dt}\right)^n = \frac{1}{n+1}\left[t^m\left(\frac{d}{dt}\right)^{n+1}, t\right], \quad m \in \mathbb{Z}, n \in \mathbb{Z}_+.$$

By (1) and (2), $f(t^{\pm 1}, t^{\pm 1}) = f(t^{\pm 1}, D) = f(D, D) = 0$.

$$\begin{aligned} f\left(t\left(\frac{d}{dt}\right)^2, t^2\left(\frac{d}{dt}\right)^2\right) &= f\left(t\left(\frac{d}{dt}\right)^2, -\frac{1}{3}[t, t^2\left(\frac{d}{dt}\right)^3]\right) = -\frac{1}{3}f\left([t\left(\frac{d}{dt}\right)^2, t], t^2\left(\frac{d}{dt}\right)^3\right) \\ &= -\frac{2}{3}f\left(t\frac{d}{dt}, t^2\left(\frac{d}{dt}\right)^3\right) = -\frac{2}{3}f\left(t\frac{d}{dt}, -\frac{1}{4}[t, t^2\left(\frac{d}{dt}\right)^4]\right) \\ &= \frac{1}{6}f\left([t\frac{d}{dt}, t], t^2\left(\frac{d}{dt}\right)^4\right) = \frac{1}{6}f\left(t, t^2\left(\frac{d}{dt}\right)^4\right) \\ &= \frac{1}{6}f\left(t, -\frac{1}{5}[t, t^2\left(\frac{d}{dt}\right)^5]\right) = -\frac{1}{30}f\left([t, t], t^2\left(\frac{d}{dt}\right)^5\right) \\ &= 0. \end{aligned}$$

Similarly, we have $f(D, t^2(\frac{d}{dt})^2) = 0$. Then,

$$\begin{aligned} f(D^2, D^2) &= f\left(D + t^2\left(\frac{d}{dt}\right)^2, D + t^2\left(\frac{d}{dt}\right)^2\right) \\ &= f(D, D) + 2f\left(D, t^2\left(\frac{d}{dt}\right)^2\right) + f\left(t^2\left(\frac{d}{dt}\right)^2, t^2\left(\frac{d}{dt}\right)^2\right) \\ &= 2f\left(D, t^2\left(\frac{d}{dt}\right)^2\right) + f\left(t^2\left(\frac{d}{dt}\right)^2, t^2\left(\frac{d}{dt}\right)^2\right) \\ &= 0. \end{aligned}$$

We complete the proof. \square

Corollary 4.6. $\text{HL}^2(\mathfrak{g}, \mathbb{K}) = \text{H}^2(\mathfrak{g}, \mathbb{K})$, for $\mathfrak{g} = \mathcal{W}$, \mathcal{H} , or \mathcal{D} .

Remark 4.7. By a different approach, for $\mathfrak{g} = \mathcal{W}$, Corollary 4.6 was obtained in [18]. For $\mathfrak{g} = \mathcal{H}$, Corollary 4.6 was obtained in [16].

4.2. Lie algebras of Block type.

Definition 4.8. ([5]) Let A be a torsion-free abelian group. $\phi : A \times A \rightarrow \mathbb{K}$ is a non-degenerate, skew-symmetric, \mathbb{Z} -bilinear function. Then we have the degenerate Block algebra

$$\mathcal{L}(A, \phi) = \bigoplus_{x \in A/\{0\}} \mathbb{K}e_x$$

with Lie bracket $[e_x, e_y] = \phi(x, y)e_{x+y}$.

Definition 4.9. Let $A = \mathbb{Z} \times \mathbb{Z}$ and $\phi((m, n), (m_1, n_1)) = nm_1 - mn_1$. It is clear that ϕ is a skew-symmetric bi-additive function. Then we have the Virasoro-like algebra (see [13])

$$\mathcal{V} = \bigoplus_{(m, n) \in \mathbb{Z} \times \mathbb{Z}/(0, 0)} \mathbb{K}e_{m, n}$$

with Lie bracket $[e_{m, n}, e_{m_1, n_1}] = (nm_1 - mn_1)e_{m+n, m_1+n_1}$.

Definition 4.10. Let $A = \mathbb{Z} \times \mathbb{Z}$, $\phi((m, n), (m_1, n_1)) = q^{nm_1} - q^{mn_1}$ with a fixed $q \in \mathbb{K}^*$ being non-root of unity. It is clear that ϕ is a skew-symmetric bi-additive function. Then we have the q -analogue Virasoro-like algebra (see [13])

$$\mathcal{V}_q = \bigoplus_{(m,n) \in \mathbb{Z} \times \mathbb{Z} / (0,0)} \mathbb{K}e_{m,n}$$

with Lie bracket $[e_{m,n}, e_{m_1, n_1}] = (q^{nm_1} - q^{mn_1})e_{m+n, m_1+n_1}$.

Lemma 4.11. ([25]) $B(\mathcal{L}(A, \phi), \mathbb{K}) = \mathbb{K}\alpha$, where $\alpha(e_x, e_y) = \delta_{x+y, 0}$.

Proposition 4.12. $HL^2(\mathcal{L}(A, \phi), \mathbb{K}) = H^2(\mathcal{L}(A, \phi), \mathbb{K})$.

Proof. By Corollary 3.3 and Lemma 4.11, it suffices to prove the image of α under the Cartan-Koszul homomorphism h is non-zero. If $h(\alpha) = \bar{0} \in H^3(\mathfrak{g}, \mathbb{K})$, there exists a skew-symmetric bilinear form ψ on \mathfrak{g} such that

$$\begin{aligned} h(\alpha)(e_x, e_y, e_z) &= \alpha([e_x, e_y], e_z) = d(\psi)(e_x, e_y, e_z) \\ &= \psi(e_x, [e_y, e_z]) + \psi(e_y, [e_z, e_x]) + \psi(e_z, [e_x, e_y]) \end{aligned}$$

for all $e_x, e_y, e_z \in \mathcal{L}(A, \phi)$. Let $x + y + z = 0$ and $\phi(x, y) \neq 0$. Then

$$\begin{aligned} \phi(x, y)\psi(e_x, e_{-x}) + \phi(x, y)\psi(e_y, e_{-y}) + \phi(x, y)\psi(e_z, e_{-z}) &= \phi(x, y), \\ \psi(e_x, e_{-x}) + \psi(e_y, e_{-y}) + \psi(e_z, e_{-z}) &= 1. \end{aligned}$$

On the other hand, since $-x - y - z = 0$, the above identity holds for $-x, -y, -z$ instead of x, y, z . Using the skew-symmetry of ψ , we have

$$\psi(e_x, e_{-x}) + \psi(e_y, e_{-y}) + \psi(e_z, e_{-z}) = -1.$$

However, this is impossible. Therefore, $h(\alpha) \neq \bar{0}$. \square

From [5], it follows

Corollary 4.13.

$$HL^2(\mathcal{L}(A, \phi), \mathbb{K}) = \{[\alpha_\mu] \mid \alpha_\mu(x, y) = \delta_{x+y, 0}\mu(x), \forall x, y \in A, \mu \in \text{Hom}_{\mathbb{K}}(A, \mathbb{K})\}.$$

Remark 4.14. Using a different approach, Corollary 4.13 was obtained in [25]. For (q -analogue) Virasoro-like algebras, Corollary 4.13 was obtained in [16].

4.3. Kac-Moody algebras.

Lemma 4.15. Let \mathfrak{g} be a Lie algebra satisfying the following conditions

- $\dim_{\mathbb{K}} B(\mathfrak{g}, \mathbb{K}) = 1$,
- \mathfrak{g} contains a subalgebra \mathfrak{a} , which is isomorphic to $sl(2, \mathbb{K})$,
- $\phi|_{\mathfrak{a}}$ is non-degenerate, where $\phi \in B(\mathfrak{g}, \mathbb{K})$.

Then $HL^2(\mathfrak{g}, \mathbb{K}) = H^2(\mathfrak{g}, \mathbb{K})$.

Proof. For nonzero $\phi \in B(\mathfrak{g}, \mathbb{K})$, $\phi|_{\mathfrak{a}}$ is nondegenerate, where \mathfrak{a} is isomorphic to $sl(2, \mathbb{K})$. So $\phi|_{\mathfrak{a}}$ is a multiple of the Killing form. Let h be the Cartan-Koszul homomorphism: $h(\phi)(a, b, c) = \phi([a, b], c)$ for any $a, b, c \in \mathfrak{g}$. If $h(\phi)(a, b, c) = \phi([a, b], c) = 0$, there exists a skew-symmetric bilinear form f on \mathfrak{g} such that

$$f(a, [b, c]) + f(b, [c, a]) + f(c, [a, b]) = \phi([a, b], c), \quad \forall a, b, c \in \mathfrak{g}.$$

We take $x, y, h \in \mathfrak{a}$ satisfying $[x, y] = h$, $[h, y] = -2y$, $[h, x] = 2x$. Then

$$\begin{aligned} f(x, [y, h]) + f(y, [h, x]) + f(h, [x, y]) &= \phi([x, y], h), \\ 2f(x, y) + 2f(y, x) + f(h, h) &= \phi(h, h). \end{aligned}$$

Therefore, $\phi(h, h) = 0$. It is contradict to the property of the Killing form. \square

Lemma 4.16. $\text{HL}^2(\mathfrak{g}, \mathbb{K}) = \text{Hom}(\text{HL}_2(\mathfrak{g}, \mathbb{K}), \mathbb{K})$, for any perfect Leibniz algebra \mathfrak{g} .

Proof. Following the universal coefficient theorem for Leibniz algebras in [4], one has the following short exact sequence

$$0 \longrightarrow \text{Ext}(\text{HL}_1(\mathfrak{g}, \mathbb{K}), \mathbb{K}) \longrightarrow \text{HL}^2(\mathfrak{g}, \mathbb{K}) \longrightarrow \text{Hom}(\text{HL}_2(\mathfrak{g}, \mathbb{K}), \mathbb{K}) \longrightarrow 0,$$

where $\text{Ext}(\text{HL}_1(\mathfrak{g}, \mathbb{K}), \mathbb{K})$ denotes the abelian extension of $\text{HL}_1(\mathfrak{g}, \mathbb{K})$ by \mathbb{K} . As a result, it is clear that $\text{Ext}(\text{HL}_1(\mathfrak{g}, \mathbb{K}), \mathbb{K}) = 0$ since $\text{HL}_1(\mathfrak{g}, \mathbb{K}) = \mathfrak{g}/[\mathfrak{g}, \mathfrak{g}] = 0$. \square

Let \mathfrak{g} be a perfect Lie algebra ($[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}$) over \mathbb{K} . Then, by [10], there exist a universal central extension $\pi : \hat{\mathfrak{g}} \rightarrow \mathfrak{g}$ in the category of Leibniz algebras, and a universal central extension $\tilde{\pi} : \tilde{\mathfrak{g}} \rightarrow \mathfrak{g}$ in the category of Lie algebras. And we have the following

Lemma 4.17. $\text{HL}_2(\tilde{\mathfrak{g}}, \mathbb{K}) = \ker\{\text{HL}_2(\mathfrak{g}, \mathbb{K}) \rightarrow \text{H}_2(\mathfrak{g}, \mathbb{K})\}$.

Proof. See 4.6 in [18] or Corollary 2.7 in [10]. \square

Let R be a unital, commutative and associative algebra over \mathbb{K} . The R -module of Kähler differentials $\Omega_{R|\mathbb{K}}^1$ is generated by \mathbb{K} -linear symbols da for $a \in R$ with the relation $d(ab) = a db + b da$, for any $a, b \in R$. In particular, if $R = \mathbb{K}[t, t^{-1}]$, then $\Omega_{R|\mathbb{K}}^1 = \bigoplus_{m \in \mathbb{Z}} \mathbb{K} t^m dt$. For any positive integer r , let

$$\Omega_{R|\mathbb{K}}^1(r) = \bigoplus_{i \in \mathbb{Z}} \mathbb{K} t^{ir-1} dt, \quad \Omega'_{R|\mathbb{K}}(r) = \bigoplus_{i \in \mathbb{Z} - \{0\}} \mathbb{K} t^{ir-1} dt$$

be two \mathbb{K} -subspaces of $\Omega_{R|\mathbb{K}}^1$, and $\Omega'_{R|\mathbb{K}}(r)$ is a subspace of $\Omega_{R|\mathbb{K}}^1(r)$ of codimension of 1.

Theorem 4.18. Let $A = (a_{ij})_{1 \leq i, j \leq n}$ denote an $n \times n$ -matrix of rank ℓ with entries in \mathbb{K} . Using the notations in [12], denote by $\mathfrak{g}(A)$ the Kac-Moody Lie algebra associated to A . Let $\mathfrak{g}'(A)$ be the derived algebra of $\mathfrak{g}(A)$, \mathfrak{c} the center of $\mathfrak{g}'(A)$, and $\bar{\mathfrak{g}}(A) = \mathfrak{g}'(A)/\mathfrak{c}$. Thus,

- (1) If $a_{ii} \neq 0$, $1 \leq i \leq n$, then $\dim \text{HL}^2(\mathfrak{g}(A), \mathbb{K}) = (n - \ell)^2$;
- (2) If A is an indecomposable generalized Cartan matrix, then

$$\begin{aligned} \text{HL}^2(\mathfrak{g}'(A), \mathbb{K}) &= (\Omega'_{R|\mathbb{K}}(r))^*, & A \text{ is of affine type } X_n^{(r)}, \\ \text{HL}^2(\mathfrak{g}'(A), \mathbb{K}) &= 0, & A \text{ is of non-affine type;} \end{aligned}$$

- (3) If A is an indecomposable generalized Cartan matrix, then

$$\begin{aligned} \text{HL}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) &= (\Omega'_{R|\mathbb{K}}(r))^*, & A \text{ is of affine type } X_n^{(r)}, \\ \text{HL}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) &= \mathfrak{c}^*, & A \text{ is of non-affine type.} \end{aligned}$$

Proof. (1) follows from Theorem 3.2 in [7] and Theorem 3.5.

(2), (3): First, we assume that A is an indecomposable generalized Cartan matrix of non-affine type. By Theorem 3.1 [2], $\bar{\mathfrak{g}}(A)$ possesses a non-degenerate invariant symmetric bilinear form if and only if A is symmetrizable. Moreover, such forms on $\bar{\mathfrak{g}}(A)$ are unique up to scalars. So $\bar{\mathfrak{g}}(A)$ satisfies the conditions in Lemma 4.15 if A is symmetrizable. We have

$$\text{HL}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \text{H}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \mathfrak{c}^*,$$

where the second “=” was proved by Theorem 2.3 [2]. If A is non-symmetrizable, then by Theorem 3.1 [2], together with A being indecomposable, we get $B(\bar{\mathfrak{g}}(A), \mathbb{K}) = 0$ and then, Corollary 3.3 above and Theorem 2.3 [2] give rise to

$$\mathrm{HL}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \mathrm{H}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \mathfrak{c}^*.$$

Thereby, in both cases, we get that $\mathfrak{g}'(A)$ is the universal Leibniz central extension of $\bar{\mathfrak{g}}(A) = \mathfrak{g}'(A)/\mathfrak{c}$. Consequently, we have

$$\mathrm{HL}^2(\mathfrak{g}'(A), \mathbb{K}) = 0.$$

Now, we assume that A is an indecomposable generalized Cartan matrix of affine type. Gao (Theorem 3.17 [9]) proved that $\mathrm{HL}_2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \Omega_{R|\mathbb{K}}^1(r)$. Then, by Lemma 4.16, we have

$$\mathrm{HL}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) = (\Omega_{R|\mathbb{K}}^1(r))^*.$$

Wilson’s well-known result in [22] told us that $\mathfrak{g}'(A)$ is the universal covering of $\bar{\mathfrak{g}}(A)$ in the category of Lie algebras, that is, we have the following exact sequence:

$$0 \longrightarrow \mathrm{H}_2(\bar{\mathfrak{g}}(A), \mathbb{K}) \longrightarrow \mathfrak{g}'(A) \longrightarrow \bar{\mathfrak{g}}(A) \longrightarrow 0,$$

where $\mathrm{H}_2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \Omega_{R|\mathbb{K}}^1(r)/\Omega_{R|\mathbb{K}}^1(r) = t^{-1} dt$. Therefore,

$$\ker\{\mathrm{HL}_2(\bar{\mathfrak{g}}(A), \mathbb{K}) \rightarrow \mathrm{H}_2(\bar{\mathfrak{g}}(A), \mathbb{K})\} = \Omega_{R|\mathbb{K}}^1(r).$$

By Lemma 4.17, we have $\mathrm{HL}_2(\mathfrak{g}'(A), \mathbb{K}) = \Omega_{R|\mathbb{K}}^1(r)$. Hence, by Lemma 4.16 again,

$$\mathrm{HL}^2(\mathfrak{g}'(A), \mathbb{K}) = (\Omega_{R|\mathbb{K}}^1(r))^*.$$

We complete the proof. \square

Corollary 4.19. *In particular, A is of finite type, then $\bar{\mathfrak{g}}(A) = \mathfrak{g}(A)$ is a finite dimensional simple Lie algebra. Hence, $\mathrm{HL}^2(\mathfrak{g}, \mathbb{K}) = \mathrm{H}^2(\mathfrak{g}, \mathbb{K}) = 0$.*

Remark 4.20. *Theorem 4.17 (2), (3) give a criterion to distinguish affine or nonaffine Kac-Moody algebras by means of the second Leibniz cohomology groups, where Theorem 4.17 (2) was obtained by Gao ([9]) under the assumption that A is symmetrizable.*

5. QUADRATIC LEIBNIZ ALGEBRAS AND THEIR CENTRAL EXTENSIONS

5.1. Quadratic Leibniz algebra.

Definition 5.1. *(\mathfrak{g}, ϕ) is called a quadratic Leibniz algebra if ϕ is a symmetric invariant bilinear form on the Leibniz algebra \mathfrak{g} .*

Lemma 5.2. *Let \mathfrak{g} be a Lie algebra. If (\mathfrak{g}, ϕ) is a quadratic Leibniz algebra and d is a derivation of \mathfrak{g} , then $f(x, y) := \phi(x, dy)$ is a Leibniz 2-cocycle on \mathfrak{g} .*

Proof. For any $x, y, z \in \mathfrak{g}$, one has

$$\begin{aligned} f(x, [y, z]) &= \phi(x, d[y, z]) = \phi(x, [dy, z] + [y, dz]) \\ &= \phi([x, y], dz) - \phi([x, z], dy) = f([x, y], z) - f([x, z], y). \end{aligned}$$

So f is a Leibniz 2-cocycle on \mathfrak{g} . \square

Corollary 5.3. *If d is a skew-derivation, i.e., $\phi(dx, y) + \phi(x, dy) = 0$, then $f(x, y) := \phi(x, dy)$ is a Lie 2-cocycle on \mathfrak{g} .*

5.2. A negative answer to a question in [16].

Proposition 5.4. *Assume that \mathfrak{g} is a finite dimensional simple Lie algebra over \mathbb{K} . Construct the Lie algebra $\mathfrak{g} \otimes \mathbb{K}((t))$ with bracket*

$$[x \otimes r, y \otimes s]' = [x, y] \otimes rs, \quad x, y \in \mathfrak{g}, r, s \in \mathbb{K}((t)).$$

Then $\mathfrak{g} \otimes \mathbb{K}((t))$ is a simple Lie algebra over \mathbb{K} and

$$H^2(\mathfrak{g} \otimes \mathbb{K}((t)), \mathbb{K}) \subsetneq HL^2(\mathfrak{g} \otimes \mathbb{K}((t)), \mathbb{K}).$$

Proof. Define $\phi(x \otimes r, y \otimes s) = (x, y) \text{Res}(rs)$, where Res is a linear function on $\mathbb{K}((t))$ and takes the coefficient of t^{-1} for every series, $(,)$ is the Killing form on \mathfrak{g} . Then ϕ is a symmetric invariant bilinear form on $\mathfrak{g} \otimes \mathbb{K}((t))$ and $t^k \frac{d}{dt}, k \in \mathbb{Z}/\{0\}$ is a derivation of $\mathbb{K}((t))$. By Lemma 5.2, we get a non-trivial Leibniz 2-cocycle of the Lie algebra $\mathfrak{g} \otimes \mathbb{K}((t))$:

$$f(x \otimes \sum_{m \geq N} a_m t^m, y \otimes \sum_{n \geq N} b_n t^n) = (x, y) \sum_{m, n \geq N} n a_m b_n \delta_{m+n+k, 0}, \quad (5.1)$$

where f is well-defined since the summation in (5.1) is finite, and f is not skew symmetric, that is, $H^2(\mathfrak{g} \otimes \mathbb{K}((t)), \mathbb{K}) \subsetneq HL^2(\mathfrak{g} \otimes \mathbb{K}((t)), \mathbb{K})$. \square

Remark 5.5. $\mathfrak{g} \otimes \mathbb{K}((t))$ is a simple Lie algebra (see [24]), and $\dim H^2(\mathfrak{g} \otimes \mathbb{K}((t)), \mathbb{K}) = 1$, but $\dim HL^2(\mathfrak{g} \otimes \mathbb{K}((t)), \mathbb{K}) = \infty$, which then leads to a negative answer to a question in [16].

ACKNOWLEDGMENT

The authors are grateful to Hagiwara and Papi for sending us their papers ([11], [20]). NH would like to express his thanks to the ICTP for the hospitality when he is visiting at the ICTP Mathematics Section, Trieste, Italy, during March 1st to August 31st, 2006.

REFERENCES

- [1] E. Arbarello, C. De Concini, V. Kac, C. Procesi, *Moduli spaces of curves and representation theory*, Comm. Math. Phys. **117** (1), (1988), 1–36.
- [2] S. Berman, *On the low-dimensional cohomology of some infinite-dimensional simple Lie algebras*, Pacific J. Math. **83** (1), (1979), 27–36.
- [3] R. Borcherds, *Central extensions of generalized Kac-Moody Lie algebras*, J. Algebra, **140**, (1991), 330–335.
- [4] J.M. Casas, E. Faro, A.M. Vieites, *Abelian extensions of Leibniz algebras*, Comm. Algebra, **27** (6), (1999), 2833–2846.
- [5] D. Dokovic, K. Zhao, *Some infinite-dimensional simple Lie algebras in characteristic 0 related to those of Block*, J. Pure and Applied Algebra, **127** (2), (1998), 153–165.
- [6] R. Farnsteiner, *Central extensions and invariant bilinear forms of graded Lie algebras*, Algebras, Groups and Geom. **3**, (1986), 431–455.
- [7] R. Farnsteiner, *Derivation and central extensions of finitely generated graded Lie algebras*, J. Algebra, **118**, (1988), 33–45.
- [8] Y. Gao, *Central extensions of nonsymmetrizable Kac-Moody algebras over commutative algebras*, Proc. Amer. Math. Soc. **121**, (1994), 67–76.
- [9] Y. Gao, *The second Leibniz homology group for Kac-Moody Lie algebras*, Bull. of Lond. Math. Soc. **32**, (2000), 25–33.
- [10] A.V. Gnedbaye, *Third homology groups of universal central extensions of a Lie algebra*, Afrika Matematika Serie 3, 10 (1998). (ICTP Preprint No.IC98059).

- [11] Y. Hagiwara, T. Mizutani, *Leibniz algebras associated with foliations*, Kodai Math. J. **25** (2), (2002), 151–165.
- [12] V. Kac, *Infinite Dimensional Lie Algebras*, Cambridge University Press, third edition, 1990.
- [13] E. Kirkman, C. Procesi, L. Small, *A q -analog for the Virasoro algebra*, Comm. Algebra. **22**, (1994), 3755–3774.
- [14] J.S. Koszul, *Homologie et cohomologie des algèbres de Lie*, Bull. Soc. Math. France **78** (1950) 65–127.
- [15] W. Li, *2-Cocycles on the algebras of differential operators*, J. Algebra. **122**, (1989), 64–80.
- [16] D. Liu, N. Hu, *Leibniz central extensions on some infinite-dimensional algebras*, Comm. Algebra. **32** (6), (2004), 2385–2405.
- [17] J. Loday, *Cyclic Homology*, vol. **301**, Grund. Math. Wiss., Springer-Verlag, 1992.
- [18] J. Loday, T. Pirashvili, *Universal enveloping algebras of Leibniz algebras and (co)homology*, Math. Ann. **296** (1), (1993), 139–158.
- [19] J.M. Lodder, *Leibniz cohomology for differentiable manifolds*, Ann. Inst. Fourier, Grenoble, **48** (1), (1998), 73–95.
- [20] P. Papi, *Cohomology of tensor product of quantum planes*, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei **9**, Mat. Appl. 3 (1), (1992), 5–13.
- [21] T. Pirashvili, *On Leibniz homology*, Ann. Inst. Fourier (Grenoble) **44** (2), (1994), 401–411.
- [22] R. L. Wilson, *Euclidean Lie algebras are universal central extensions*, *Lie Algebras and Related Topics*, Lecture Notes in Math. 933 (ed. David Winter, Springer, New York, 1982) 210–213.
- [23] K. Zhao, *Automorphism groups of the algebras of differential operators*, J. of Capital Normal Uni. **15**, (1994), 1–7.
- [24] L. Zhu, D. Meng, *Commutative associative algebras and quadratic Lie algebras*, Comm. Algebra **29**, (2001), 2249–2268.
- [25] L. Zhu, D. Meng, *Structure of degenerate Block algebras*, Algebra Colloq. **10**, (2003), 53–62.

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200062, PR CHINA

E-mail address: `nhhu@euler.math.ecnu.edu.cn` & `nhu@ictp.it` (until August 31, 2006)

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200062, PR CHINA

E-mail address: `peiyufeng@gmail.com`

DEPARTMENT OF MATHEMATICS, JIAOTONG UNIVERSITY, SHANGHAI 200030, PR CHINA

E-mail address: `dliu@sjtu.edu.cn`