

A COHOMOLOGICAL CHARACTERIZATION OF LEIBNIZ CENTRAL EXTENSIONS OF LIE ALGEBRAS

NAIHONG HU*, YUFENG PEI, AND DONG LIU

ABSTRACT. Motivated by Pirashvili's spectral sequences on a Leibniz algebra, a cohomological characterization of Leibniz central extensions of Lie algebras is given. Consequently, we obtain the cohomological version of Gao's main Theorem in [9] for Kac-Moody algebras and answer a question in [16].

1. INTRODUCTION

Leibniz algebras introduced by Loday ([17]) are a non-antisymmetric generalization of Lie algebras. There is a (co)homology theory for these algebraic objects whose properties are similar to those of the classical Chevalley-Eilenberg cohomology theory for Lie algebras. Since a Lie algebra is a Leibniz algebra, it is interesting to study Leibniz (co)homology of Lie algebras which may provide new invariants in return for Lie algebras. Lodder ([19]) obtained the Godbillon-Vey invariants for foliations by computing Leibniz cohomology of certain Lie algebras and mentioned how a Leibniz algebra arises naturally from vertex (operator) algebras. Recently, some interrelations with manifolds were investigated which could lead to possible applications of Leibniz (co)homology in geometry (see [11, 19], etc.).

Central extensions play a central role in the theory of Lie algebras (see [1, 2, 3, 6, 7, 8, 13, 14, 15, 20, 22], etc.). Viewing a Lie algebra as a Leibniz algebra, it is natural to determine its Leibniz central extensions and compare the differences between Leibniz and Lie central extensions in question. Loday and Pirashvili ([18]) shown that the Virasoro algebra is a universal central extension of the Witt algebra in the category of Leibniz algebras as well. It is well-known that any Kac-Moody Lie algebra is centrally closed. However, its universal central extension in the category of Leibniz algebras is not centrally closed for affine type as witnessed by Gao in ([9]). Some other cases of infinite dimensional Lie algebras were discussed in [16, 25], among others. Their approaches involve technical and lengthy computations.

The purpose of this note is to give a concise determination of the universal central extension of a class of Lie algebras in the category of Leibniz algebras. Our motivation comes from the long exact sequence used by Pirashvili ([19, 21]). Note that the invariant symmetric bilinear forms and Leibniz central extensions can be well-connected by the Cartan-Koszul homomorphism (see [14]) in the sequence. In practice, it is easier to handle the invariant symmetric bilinear forms on a Lie algebra than to directly determine its Leibniz central extensions. On the other hand, we

1991 *Mathematics Subject Classification.* Primary 17A32, 17B56; Secondary 17B65.

Key words and phrases. Leibniz central extensions, Leibniz cohomology, invariant symmetric bilinear forms, dual space derivations.

*N.H., Corresponding author. This work is supported in part by the NNSF (Grant 10431040), the TRAPOYT and the FUDP from the MOE of China, the SRSTP from the STCSM.

notice that determining Leibniz central extensions of a Lie algebra is equivalent to treating dual space derivations of this Lie algebra (cf. [7]). We combine these observations to study the (universal) Leibniz central extensions of Lie algebras. The current method avoids complicated computations when applied to some classes of infinite dimensional Lie algebras we are interested in.

The organization of this note is as follows. Section 2 introduces some basic notions on Leibniz algebras. In Section 3, we present a short exact sequence as a variation of the Pirashvili's long exact sequence. We then go on to describe the Leibniz central extensions of Lie algebras, in terms of the invariant symmetric bilinear forms and the Cartan-Koszul homomorphism, and to obtain the 2nd Leibniz cohomology group with trivial coefficients by dual space derivations (Theorem 3.5). Section 4 provides some applications. As a consequence, a cohomological version of Gao's result ([9]) in Kac-Moody algebras case (Theorem 4.18) is given. In Section 5, we study the Leibniz central extensions of the quadratic Lie algebras and construct a counterexample to address a question posed in [16].

2. PREREQUISITES ON LEIBNIZ ALGEBRAS

2.1. Leibniz algebra. Let \mathbb{K} be an algebraically closed field with $\text{char } \mathbb{K} = 0$.

Definition 2.1. A Leibniz algebra is a \mathbb{K} -module L with a bilinear map $[-, -] : L \times L \rightarrow L$ satisfying the Leibniz identity $[x, [y, z]] = [[x, y], z] - [[x, z], y]$, for $x, y, z \in L$.

The center of L is defined as $\{z \in L \mid [z, L] = [L, z] = 0\}$. L is called *perfect* if $[L, L] = L$. If, in addition, $[x, x] = 0, \forall x \in L$, the Leibniz identity is equivalent to the Jacobi identity. In particular, Lie algebras are examples of Leibniz algebras.

Definition 2.2. Let L be a Leibniz algebra over \mathbb{K} . M is called a representation of L if M is a \mathbb{K} -vector space equipped with two actions (left and right) of L , i.e., $[-, -] : L \times M \rightarrow M$ and $[-, -] : M \times L \rightarrow M$ satisfying

$$\begin{aligned} (MLL) \quad & [m, [x, y]] = [[m, x], y] - [[m, y], x], \\ (LML) \quad & [x, [m, y]] = [[x, m], y] - [[x, y], m], \\ (LLM) \quad & [x, [y, m]] = [[x, y], m] - [[x, m], y], \end{aligned}$$

for any $m \in M$ and $x, y \in L$.

2.2. Cohomology of Leibniz algebras. Let L be a Leibniz algebra over \mathbb{K} and M a representation of L . Denote $C^n(L, M) := \text{Hom}_{\mathbb{K}}(L^{\otimes n}, M)$, $n \geq 0$. The Loday coboundary map $d^n : C^n(L, M) \rightarrow C^{n+1}(L, M)$ is defined by

$$\begin{aligned} (d^n f)(x_1, \dots, x_{n+1}) &= [x_1, f(x_2, \dots, x_{n+1})] \\ &+ \sum_{i=2}^{n+1} (-1)^i [f(x_1, \dots, \hat{x}_i, \dots, x_{n+1}), x_i] \\ &+ \sum_{1 \leq i < j \leq n+1} (-1)^{j+1} f(x_1, \dots, x_{i-1}, [x_i, x_j], x_{i+1}, \dots, \hat{x}_j, \dots, x_{n+1}). \end{aligned}$$

Clearly, $d^{n+1}d^n = 0$, $n \geq 0$. $(C^*(L, M), d)$ is a well-defined cochain complex, whose cohomology is called the cohomology of Leibniz algebra L with coefficients in the representation M : $\text{HL}^*(L, M) := \text{H}^*((C^*(L, M), d))$. Similarly, we have a chain

complex $(C_*(L, M), d)$, whose homology is called the homology of Leibniz algebra L with coefficients in the representation M : $\text{HL}_*(L, M) := \text{H}_*((C_*(L, M), d))$.

2.3. Cohomology of Lie algebras. Let \mathfrak{g} be a Lie algebra over \mathbb{K} and M a \mathfrak{g} -module. Denote the Chevalley-Eilenberg cochain complex by

$$(\Omega^*(\mathfrak{g}, M), \delta) := (\text{Hom}(\wedge^* \mathfrak{g}, M), \delta),$$

where δ is the Chevalley-Eilenberg coboundary map defined by

$$\begin{aligned} (\delta^n f)(x_1, \dots, x_{n+1}) &= \sum_{i=1}^{n+1} (-1)^{i+1} x_i \cdot f(x_1, \dots, \hat{x}_i, \dots, x_{n+1}) \\ &+ \sum_{1 \leq i < j \leq n+1} (-1)^{i+j} f([x_i, x_j], x_1, \dots, \hat{x}_i, \dots, \hat{x}_j, \dots, x_{n+1}). \end{aligned}$$

Then $\text{H}^*(\mathfrak{g}, M) := \text{H}^*((\Omega^*(\mathfrak{g}, M), \delta))$ is called the cohomology of Lie algebra \mathfrak{g} with coefficients in the \mathfrak{g} -module M .

3. LEIBNIZ COHOMOLOGY AND LEIBNIZ CENTRAL EXTENSIONS OF LIE ALGEBRAS

3.1. Central extensions of Leibniz algebras. A central extension of L is a pair (\hat{L}, π) , where \hat{L} is a Leibniz algebra and $\pi : \hat{L} \rightarrow L$ is a surjective homomorphism whose kernel lies in the center of \hat{L} . The pair (\hat{L}, π) is a universal central extension of L if for every central extension (\tilde{L}, τ) of L , there is a unique homomorphism $\psi : \hat{L} \rightarrow \tilde{L}$ for which $\tau \circ \psi = \pi$. The following result is known.

Proposition 3.1. ([18]) *There exists a one-to-one correspondence between the set of equivalent classes of one-dimensional Leibniz central extensions of L by \mathbb{K} and the 2nd Leibniz cohomology group $\text{HL}^2(L, \mathbb{K})$.*

3.2. Invariant symmetric bilinear forms. Let \mathfrak{g} be a Lie algebra over \mathbb{K} and M a \mathfrak{g} -module. A symmetric bilinear form ϕ on \mathfrak{g} is called \mathfrak{g} -invariant if ϕ satisfies $\phi([x, y], z) = \phi(x, [y, z])$, $\forall x, y, z \in \mathfrak{g}$. Let $\text{B}(\mathfrak{g}, \mathbb{K})$ stand for the set of all \mathbb{K} -valued symmetric \mathfrak{g} -invariant bilinear forms on Lie algebra \mathfrak{g} .

The short exact sequence below is a variation of the first 5-terms of Pirashvili's long exact sequence (see [21]).

Proposition 3.2. *For any Lie algebra \mathfrak{g} , there holds the following exact sequence*

$$0 \longrightarrow \text{H}^2(\mathfrak{g}, \mathbb{K}) \xrightarrow{f} \text{HL}^2(\mathfrak{g}, \mathbb{K}) \xrightarrow{g} \text{B}(\mathfrak{g}, \mathbb{K}) \xrightarrow{h} \text{H}^3(\mathfrak{g}, \mathbb{K}),$$

where f is the natural embedding map, g is defined by $g(\alpha)(x, y) = \alpha(x, y) + \alpha(y, x)$, $\forall \alpha \in \text{HL}^2(\mathfrak{g}, \mathbb{K})$, $x, y \in \mathfrak{g}$, and h is the Cartan-Koszul map ([14]) defined by $h(\alpha)(x, y, z) = \alpha([x, y], z)$, $\forall \alpha \in \text{B}(\mathfrak{g}, \mathbb{K})$, $x, y, z \in \mathfrak{g}$. \square

Corollary 3.3. $\frac{\text{HL}^2(\mathfrak{g}, \mathbb{K})}{\text{H}^2(\mathfrak{g}, \mathbb{K})} = \ker(h)$. *In particular, $\text{HL}^2(\mathfrak{g}, \mathbb{K}) = \text{H}^2(\mathfrak{g}, \mathbb{K})$ if and only if $\ker(h) = 0$.*

Remark 3.4. *Note that the natural embedding $\Omega^*(\mathfrak{g}, M) \hookrightarrow C^*(\mathfrak{g}, M)$ induces a short exact sequence in the category of cochain complexes of Lie algebra \mathfrak{g} :*

$$0 \longrightarrow \Omega^*(\mathfrak{g}, M) \longrightarrow C^*(\mathfrak{g}, M) \longrightarrow C_{rel}^*(\mathfrak{g}, M)[2] \longrightarrow 0,$$

where $C_{rel}^*(\mathfrak{g}, M)[2] := \frac{C^*(\mathfrak{g}, M)}{\Omega^*(\mathfrak{g}, M)}$ is the quotient cochain complex. One has the Pirashvili's long exact sequence below (see [19], [21]), which is the main tool to compare the Lie and the Leibniz cohomology of a Lie algebra in higher dimensions

$$0 \rightarrow H^2(\mathfrak{g}, M) \rightarrow HL^2(\mathfrak{g}, M) \rightarrow H_{rel}^0(\mathfrak{g}, M) \rightarrow H^3(\mathfrak{g}, M) \rightarrow HL^3(\mathfrak{g}, M) \rightarrow \dots$$

and isomorphisms $HL^i(\mathfrak{g}, M) = H^i(\mathfrak{g}, M)$, $i = 0, 1$.

Let $M = \mathbb{K}$. We have

$$0 \rightarrow H^2(\mathfrak{g}, \mathbb{K}) \rightarrow HL^2(\mathfrak{g}, \mathbb{K}) \rightarrow H_{rel}^0(\mathfrak{g}, \mathbb{K}) \rightarrow H^3(\mathfrak{g}, \mathbb{K}) \rightarrow HL^3(\mathfrak{g}, \mathbb{K}) \rightarrow \dots$$

As a consequence (which is not clear for us) of the spectral sequences, Pirashvili claimed (with no proof, see [21]) : $H_{rel}^0(\mathfrak{g}, \mathbb{K}) \cong B(\mathfrak{g}, \mathbb{K})$, while a direct elementary proof of Proposition 3.2 is given in Appendix.

3.3. Dual space derivations. Let \mathfrak{g} be a Lie algebra over \mathbb{K} and M a \mathfrak{g} -module. A linear map $d : \mathfrak{g} \rightarrow M$ is a derivation if $d([x, y]) = x \cdot d(y) - y \cdot d(x)$, for $x, y \in \mathfrak{g}$. The derivations of the form $x \mapsto x \cdot m$ for some $m \in M$ are called inner derivations. $\text{Der}(\mathfrak{g}, M)$ and $\text{Inn}(\mathfrak{g}, M)$ denote the spaces of derivations and inner derivations, respectively. Clearly, $H^1(\mathfrak{g}, M) = \text{Der}(\mathfrak{g}, M)/\text{Inn}(\mathfrak{g}, M)$ is the first cohomology group of \mathfrak{g} with coefficients in M .

Theorem 3.5. $HL^2(\mathfrak{g}, \mathbb{K}) = H^1(\mathfrak{g}, \mathfrak{g}^*) = \text{Der}(\mathfrak{g}, \mathfrak{g}^*) / \text{Inn}(\mathfrak{g}, \mathfrak{g}^*)$, for any Lie algebra \mathfrak{g} , where \mathfrak{g}^* is the dual \mathfrak{g} -module.

Proof. Define a map $\theta : H^1(\mathfrak{g}, \mathfrak{g}^*) \rightarrow HL^2(\mathfrak{g}, \mathbb{K})$ as $\theta(\alpha)(x, y) := \alpha(y)(x)$ for any $\alpha \in H^1(\mathfrak{g}, \mathfrak{g}^*)$ and $x, y \in \mathfrak{g}$. Indeed, since

$$\begin{aligned} \theta(\alpha)(x, [y, z]) &= \alpha([y, z])(x) = (y \cdot \alpha(z))(x) - (z \cdot \alpha(y))(x) \\ &= \alpha(z)([x, y]) - \alpha(y)([x, z]) = \theta(\alpha)([x, y], z) + \theta(\alpha)([z, x], y), \end{aligned}$$

$\theta(\alpha) \in HL^2(\mathfrak{g}, \mathbb{K})$. θ is well-defined. If $\theta(\alpha) = \bar{0}$ in $HL^2(\mathfrak{g}, \mathbb{K})$, there exists an element β in $C^1(\mathfrak{g}, \mathbb{K})$ such that, for any $x, y \in \mathfrak{g}$, $d^1(\beta)(x, y) = \theta(\alpha)(x, y)$. That is, $-\beta([x, y]) = \alpha(y)(x)$. So α is a 1-coboundary in $\Omega^1(\mathfrak{g}, \mathfrak{g}^*)$, i.e., $\alpha = \bar{0} \in H^1(\mathfrak{g}, \mathfrak{g}^*)$. Hence, θ is injective. On the other hand, for any $\beta \in HL^2(\mathfrak{g}, \mathbb{K})$, we define a map $\alpha \in \text{Hom}_{\mathbb{K}}(\mathfrak{g}, \mathfrak{g}^*)$ by $\alpha(x)(y) := \beta(y, x)$, for any $x, y \in \mathfrak{g}$. Since

$$\begin{aligned} \alpha([x, y])(z) &= \beta(z, [x, y]) = \beta([z, x], y) - \beta([z, y], x) = \alpha(y)([z, x]) - \alpha(x)([z, y]), \\ &= (x \cdot \alpha(x))(z) - (y \cdot \alpha(x))(z), \quad \text{for any } z \in \mathfrak{g}, \end{aligned}$$

we get $\alpha([x, y]) = x \cdot \alpha(x) - y \cdot \alpha(x)$ for any $x, y \in \mathfrak{g}$, which means $\alpha \in H^1(\mathfrak{g}, \mathfrak{g}^*)$. Hence, θ is surjective. \square

Denote $\text{SDer}(\mathfrak{g}, \mathfrak{g}^*) := \{\phi \in \text{Der}(\mathfrak{g}, \mathfrak{g}^*) \mid \phi(x)(y) + \phi(y)(x) = 0, \forall x, y \in \mathfrak{g}\}$. By Theorem 3.5 above and Proposition 1.3 (2) in [6], we have

Corollary 3.6. $HL^2(\mathfrak{g}, \mathbb{K})/H^2(\mathfrak{g}, \mathbb{K}) = \text{Der}(\mathfrak{g}, \mathfrak{g}^*) / \text{SDer}(\mathfrak{g}, \mathfrak{g}^*) (\subset H_{rel}^0(\mathfrak{g}, \mathbb{K}))$.

Corollary 3.7. *If \mathfrak{g} is a finite dimensional Lie algebra over \mathbb{K} with a nondegenerate invariant symmetric bilinear form ψ , then $HL^2(\mathfrak{g}, \mathbb{K}) = \text{Der}(\mathfrak{g}, \mathfrak{g}) / \text{Inn}(\mathfrak{g}, \mathfrak{g})$. If, in addition, \mathfrak{g} is simple, then $HL^2(\mathfrak{g}, \mathbb{K}) = 0$. \square*

4. APPLICATIONS: LEIBNIZ CENTRAL EXTENSIONS OF SOME LIE ALGEBRAS

4.1. Lie algebras of Virasoro type. Let $\mathbb{K}[t, t^{-1}]$ be the Laurent polynomial algebra over \mathbb{K} and $\frac{d}{dt}$ be the differential operator on $\mathbb{K}[t, t^{-1}]$. Set $L_n = -t^{n+1}\frac{d}{dt}$ and $I_n = t^n$ for $n \in \mathbb{Z}$.

Definition 4.1. *The Witt algebra $\mathcal{W} = \bigoplus_{n \in \mathbb{Z}} \mathbb{K}L_n$ is defined by $[L_m, L_n] = (m-n)L_{m+n}$, $m, n \in \mathbb{Z}$.*

It is well-known that $H^2(\mathcal{W}, \mathbb{K}) = H_2(\mathcal{W}, \mathbb{K}) = \mathbb{K}\alpha$, $\alpha(L_m, L_n) = \delta_{m+n, 0}\frac{m^3-m}{12}$.

Definition 4.2. ([1]) *The Lie algebra $\mathcal{H} = \bigoplus_{n \in \mathbb{Z}} \mathbb{C}L_n \oplus \bigoplus_{m \in \mathbb{Z}} \mathbb{C}I_m$ is defined by $[L_m, L_n] = (m-n)L_{m+n}$, $[I_m, I_n] = 0$, $[L_m, I_n] = -nI_{m+n}$, $m, n \in \mathbb{Z}$.*

In [1], authors proved that $\dim H^2(\mathcal{H}, \mathbb{K}) = \dim H_2(\mathcal{H}, \mathbb{K}) = 3$. In fact, as the universal central extension of \mathcal{H} , the twisted Heisenberg-Virasoro algebra H_{Vir} has a basis $\{I_m, L_m, C_I, C_L, C_{LI} \mid m \in \mathbb{Z}\}$ satisfying the following relations:

$$\begin{aligned} [I_m, I_n] &= n\delta_{m+n, 0}C_I, \\ [L_m, L_n] &= (m-n)L_{m+n} + \delta_{m+n, 0}\frac{1}{12}(m^3 - m)C_L, \\ [L_m, I_n] &= -nI_{m+n} + \delta_{m+n, 0}(m^2 - m)C_{LI}, \\ [H_{Vir}, C_L] &= [H_{Vir}, C_I] = [H_{Vir}, C_{LI}] = 0. \end{aligned}$$

Definition 4.3. *Lie algebra $\mathcal{D} = \bigoplus_{m \in \mathbb{Z}, n \in \mathbb{Z}_+} \mathbb{K}t^m D^n$ of differential operators is defined by $[t^m D^r, t^n D^s] = t^{m+n}((D+n)^r - (D+m)^s)$, $m, n \in \mathbb{Z}$, $r, s \in \mathbb{Z}_+$, where $D = t\frac{d}{dt}$.*

By [15], $\dim H^2(\mathcal{D}, \mathbb{K}) = \dim H_2(\mathcal{D}, \mathbb{K}) = 1$, and the universal central extension of \mathcal{D} is a Lie algebra $\mathcal{W}_{1+\infty} = \bigoplus_{m \in \mathbb{Z}, n \in \mathbb{Z}_+} \mathbb{K}t^m D^n \oplus \mathbb{K}C$ with definition $[t^m D^r, t^n D^s] = t^{m+n}((D+n)^r - (D+m)^s) + \psi(t^m D^r, t^n D^s)C$, $\psi(t^{m+r} D^r, t^{n+s} D^s) = \delta_{m+n, 0}(-1)^r r! s! \binom{m+r}{r+s+1}$, for $m, n \in \mathbb{Z}$, $r, s \in \mathbb{Z}_+$.

Remark 4.4. \mathcal{W} and \mathcal{H} are Lie subalgebras of \mathcal{D} .

Proposition 4.5. *There is no non-trivial invariant symmetric bilinear form on Lie algebra \mathfrak{g} , where $\mathfrak{g} = \mathcal{W}$, \mathcal{H} , or \mathcal{D} .*

Proof. Assume that f is an invariant symmetric bilinear form on \mathfrak{g} . Note that

(1) $\mathfrak{g} = \mathcal{W}$ is generated as Lie algebra by L_{-2}, L_3 with $f(L_i, L_j) = 0$ for $i, j \in \{-2, 3\}$. So $f \equiv 0$.

(2) $\mathfrak{g} = \mathcal{H}$ is generated as Lie algebra by L_{-2}, L_3 and I_1 with $f(x, y) = 0$ for $x, y \in \{L_{-2}, L_3, I_1\}$. So $f \equiv 0$.

(3) $\mathfrak{g} = \mathcal{D}$ is generated as Lie algebra by t, t^{-1} and D^2 (see [23]). By a direct computation, we have

$$\begin{aligned} t^2\left(\frac{d}{dt}\right)^2 &= D^2 - D, \\ t^m\left(\frac{d}{dt}\right)^n &= \frac{1}{n+1}\left[t^m\left(\frac{d}{dt}\right)^{n+1}, t\right], \quad m \in \mathbb{Z}, n \in \mathbb{Z}_+. \end{aligned}$$

By (1) and (2), $f(t^{\pm 1}, t^{\pm 1}) = f(t^{\pm 1}, D) = f(D, D) = 0$.

$$\begin{aligned}
f\left(t\left(\frac{d}{dt}\right)^2, t^2\left(\frac{d}{dt}\right)^2\right) &= f\left(t\left(\frac{d}{dt}\right)^2, -\frac{1}{3}[t, t^2\left(\frac{d}{dt}\right)^3]\right) = -\frac{1}{3}f\left([t\left(\frac{d}{dt}\right)^2, t], t^2\left(\frac{d}{dt}\right)^3\right) \\
&= -\frac{2}{3}f\left(t\frac{d}{dt}, t^2\left(\frac{d}{dt}\right)^3\right) = -\frac{2}{3}f\left(t\frac{d}{dt}, -\frac{1}{4}[t, t^2\left(\frac{d}{dt}\right)^4]\right) \\
&= \frac{1}{6}f\left([t\frac{d}{dt}, t], t^2\left(\frac{d}{dt}\right)^4\right) = \frac{1}{6}f\left(t, t^2\left(\frac{d}{dt}\right)^4\right) \\
&= \frac{1}{6}f\left(t, -\frac{1}{5}[t, t^2\left(\frac{d}{dt}\right)^5]\right) = -\frac{1}{30}f\left([t, t], t^2\left(\frac{d}{dt}\right)^5\right) \\
&= 0.
\end{aligned}$$

Similarly, we have $f(D, t^2(\frac{d}{dt})^2) = 0$. Then,

$$\begin{aligned}
f(D^2, D^2) &= f\left(D + t^2\left(\frac{d}{dt}\right)^2, D + t^2\left(\frac{d}{dt}\right)^2\right) \\
&= f(D, D) + 2f\left(D, t^2\left(\frac{d}{dt}\right)^2\right) + f\left(t^2\left(\frac{d}{dt}\right)^2, t^2\left(\frac{d}{dt}\right)^2\right) \\
&= 2f\left(D, t^2\left(\frac{d}{dt}\right)^2\right) + f\left(t^2\left(\frac{d}{dt}\right)^2, t^2\left(\frac{d}{dt}\right)^2\right) \\
&= 0.
\end{aligned}$$

In summary, $B(\mathfrak{g}, \mathbb{K}) = 0$ for $\mathfrak{g} = \mathcal{W}, \mathcal{H}$, or \mathcal{D} . \square

Corollary 4.6. $\text{HL}^2(\mathfrak{g}, \mathbb{K}) = \text{H}^2(\mathfrak{g}, \mathbb{K})$, for $\mathfrak{g} = \mathcal{W}, \mathcal{H}$, or \mathcal{D} .

Remark 4.7. Corollary 4.6 for $\mathfrak{g} = \mathcal{W}$ was obtained by [18], for $\mathfrak{g} = \mathcal{D}$ by [16].

4.2. Lie algebras of Block type.

Definition 4.8. ([5]) Let A be a torsion-free abelian group. $\phi : A \times A \rightarrow \mathbb{K}$ is a non-degenerate, skew-symmetric, \mathbb{Z} -bilinear function. Then the degenerate Block algebra $\mathcal{L}(A, \phi) = \bigoplus_{x \in A - \{0\}} \mathbb{K}e_x$ is defined by $[e_x, e_y] = \phi(x, y)e_{x+y}$.

Definition 4.9. ([13]) Let $A = \mathbb{Z} \times \mathbb{Z}$, $\phi((m, n), (m_1, n_1)) = nm_1 - mn_1$ be a skew-symmetric bi-additive function. The Virasoro-like algebra

$$\mathcal{V} = \bigoplus_{(m, n) \in \mathbb{Z} \times \mathbb{Z} - \{(0, 0)\}} \mathbb{K}e_{m, n}$$

is defined by $[e_{m, n}, e_{m_1, n_1}] = (nm_1 - mn_1)e_{m+n, m_1+n_1}$.

Definition 4.10. ([13]) Let $A = \mathbb{Z} \times \mathbb{Z}$, $\phi((m, n), (m_1, n_1)) = q^{nm_1} - q^{mn_1}$ with a fixed $q \in \mathbb{K}^*$ (non-root of unity), where ϕ is a skew-symmetric bi-additive function. Then the q -analogue Virasoro-like algebra $\mathcal{V}_q = \bigoplus_{(m, n) \in \mathbb{Z} \times \mathbb{Z} - \{(0, 0)\}} \mathbb{K}e_{m, n}$ is defined by $[e_{m, n}, e_{m_1, n_1}] = (q^{nm_1} - q^{mn_1})e_{m+n, m_1+n_1}$.

Lemma 4.11. ([25]) $B(\mathcal{L}(A, \phi), \mathbb{K}) = \mathbb{K}\alpha$, where $\alpha(e_x, e_y) = \delta_{x+y, 0}$.

Proposition 4.12. $\text{HL}^2(\mathcal{L}(A, \phi), \mathbb{K}) = \text{H}^2(\mathcal{L}(A, \phi), \mathbb{K})$.

Proof. By Corollary 3.3 and Lemma 4.11, it suffices to prove the image of α under the Cartan-Koszul homomorphism h is non-zero. If $h(\alpha) = \bar{0} \in \text{H}^3(\mathfrak{g}, \mathbb{K})$, there is a $\psi \in \Lambda^2 \mathfrak{g}$ such that $h(\alpha)(e_x, e_y, e_z) = \alpha([e_x, e_y], e_z) = d(\psi)(e_x, e_y, e_z) = \psi(e_x, [e_y, e_z]) + \psi(e_y, [e_z, e_x]) + \psi(e_z, [e_x, e_y])$, for $e_x, e_y, e_z \in \mathcal{L}(A, \phi)$. Let $x+y+z = 0$ and $\phi(x, y) \neq 0$. Then $\phi(x, y)\psi(e_x, e_{-x}) + \phi(x, y)\psi(e_y, e_{-y}) + \phi(x, y)\psi(e_z, e_{-z}) =$

$\phi(x, y)$, that is, $\psi(e_x, e_{-x}) + \psi(e_y, e_{-y}) + \psi(e_z, e_{-z}) = 1$. On the other hand, since $-x - y - z = 0$, the above identity holds for $-x, -y, -z$. As ψ is skew-symmetric, we have $\psi(e_x, e_{-x}) + \psi(e_y, e_{-y}) + \psi(e_z, e_{-z}) = -1$. It is impossible. So $h(\alpha) \neq \bar{0}$. \square

It follows from [5] that

Corollary 4.13.

$$\mathrm{HL}^2(\mathcal{L}(A, \phi), \mathbb{K}) = \{[\alpha_\mu] \mid \alpha_\mu(x, y) = \delta_{x+y, 0}\mu(x), \forall x, y \in A, \mu \in \mathrm{Hom}_{\mathbb{K}}(A, \mathbb{K})\}.$$

Remark 4.14. *With a different method, Corollary 4.13 was given in [25]. For the (q -analogue) Virasoro-like algebras, the same result was given in [16].*

4.3. Kac-Moody algebras.

Lemma 4.15. *For a Lie algebra \mathfrak{g} with $\dim \mathrm{B}(\mathfrak{g}, \mathbb{K}) \leq 1$, in both cases below:*

- if $\mathrm{B}(\mathfrak{g}, \mathbb{K}) = 0$; or
- if $\mathrm{B}(\mathfrak{g}, \mathbb{K}) = \mathbb{K}\phi$ ($\phi \neq 0$), and there exists a subalgebra $\mathfrak{a} \cong \mathfrak{sl}(2, \mathbb{K})$ such that $\phi|_{\mathfrak{a}} \neq 0$,

then $\mathrm{HL}^2(\mathfrak{g}, \mathbb{K}) = \mathrm{H}^2(\mathfrak{g}, \mathbb{K})$.

Proof. The first case is clear by Corollary 3.3. Now for nonzero $\phi \in \mathrm{B}(\mathfrak{g}, \mathbb{K})$, if there is a subalgebra \mathfrak{a} as a $\mathfrak{sl}(2, \mathbb{K})$ -copy such that $\phi|_{\mathfrak{a}} \neq 0$, then $\phi|_{\mathfrak{a}}$ is a nonzero scalar multiple of the Killing form on \mathfrak{a} . Let h be the Cartan-Koszul homomorphism: $h(\phi)(a, b, c) = \phi([a, b], c)$ for any $a, b, c \in \mathfrak{g}$. If $h(\phi) = \bar{0}$, then there is a $\theta \in \Lambda^2 \mathfrak{g}$ such that $\delta^2(\theta) = h(\phi)$, i.e., $\theta(a, [b, c]) + \theta(b, [c, a]) + \theta(c, [a, b]) = \phi([a, b], c)$, for $a, b, c \in \mathfrak{g}$. Take $x, y, h \in \mathfrak{a}$ satisfying $[x, y] = h$, $[h, y] = -2y$, $[h, x] = 2x$. Then

$$\begin{aligned} \theta(x, [y, h]) + \theta(y, [h, x]) + \theta(h, [x, y]) &= \phi([x, y], h), \\ 2\theta(x, y) + 2\theta(y, x) + \theta(h, h) &= \phi(h, h). \end{aligned}$$

Therefore, $\phi(h, h) = 0$. It contradicts the property of the Killing form. \square

Lemma 4.16. $\mathrm{HL}^2(\mathfrak{g}, \mathbb{K}) = \mathrm{Hom}(\mathrm{HL}_2(\mathfrak{g}, \mathbb{K}), \mathbb{K})$, for any perfect Leibniz algebra \mathfrak{g} .

Proof. Following the universal coefficient theorem for Leibniz algebras in [4], one has the following short exact sequence

$$0 \longrightarrow \mathrm{Ext}(\mathrm{HL}_1(\mathfrak{g}, \mathbb{K}), \mathbb{K}) \longrightarrow \mathrm{HL}^2(\mathfrak{g}, \mathbb{K}) \longrightarrow \mathrm{Hom}(\mathrm{HL}_2(\mathfrak{g}, \mathbb{K}), \mathbb{K}) \longrightarrow 0,$$

where $\mathrm{Ext}(\mathrm{HL}_1(\mathfrak{g}, \mathbb{K}), \mathbb{K})$ denotes the abelian extension of $\mathrm{HL}_1(\mathfrak{g}, \mathbb{K})$ by \mathbb{K} . As a result, it is clear that $\mathrm{Ext}(\mathrm{HL}_1(\mathfrak{g}, \mathbb{K}), \mathbb{K}) = 0$ since $\mathrm{HL}_1(\mathfrak{g}, \mathbb{K}) = \mathfrak{g}/[\mathfrak{g}, \mathfrak{g}] = 0$. \square

Let \mathfrak{g} be a perfect Lie algebra ($[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}$) over \mathbb{K} . Then, by [10], there are a universal central extension $\pi : \hat{\mathfrak{g}} \rightarrow \mathfrak{g}$ in the category of Leibniz algebras, and a universal central extension $\tilde{\pi} : \tilde{\mathfrak{g}} \rightarrow \mathfrak{g}$ in the category of Lie algebras.

Lemma 4.17. $\mathrm{HL}_2(\tilde{\mathfrak{g}}, \mathbb{K}) = \ker\{\mathrm{HL}_2(\mathfrak{g}, \mathbb{K}) \rightarrow \mathrm{H}_2(\mathfrak{g}, \mathbb{K})\}$.

Proof. See 4.6 in [18] or Corollary 2.7 in [10]. \square

Let R be a unital, commutative and associative algebra over \mathbb{K} . The R -module of Kähler differentials $\Omega_{R|\mathbb{K}}^1$ is generated by \mathbb{K} -linear symbols da for $a \in R$ with the relation $d(ab) = a db + b da$, for any $a, b \in R$. In particular, if $R = \mathbb{K}[t, t^{-1}]$, then $\Omega_{R|\mathbb{K}}^1 = \bigoplus_{m \in \mathbb{Z}} \mathbb{K}t^m dt$. For any positive integer r , let

$$\Omega_{R|\mathbb{K}}^1(r) = \bigoplus_{i \in \mathbb{Z}} \mathbb{K}t^{ir-1} dt, \quad \Omega'_{R|\mathbb{K}}(r) = \bigoplus_{i \in \mathbb{Z} - \{0\}} \mathbb{K}t^{ir-1} dt$$

be two \mathbb{K} -subspaces of $\Omega_{R|\mathbb{K}}^1$, $\Omega_{R|\mathbb{K}}^1(r)$ is a subspace of $\Omega_{R|\mathbb{K}}^1(r)$ of codimension 1.

Theorem 4.18. *Let $A = (a_{ij})_{1 \leq i, j \leq n}$ denote an $n \times n$ -matrix of rank ℓ with entries in \mathbb{K} . Using the notations in [12], denote by $\mathfrak{g}(A)$ the Kac-Moody Lie algebra associated to A . Let $\mathfrak{g}'(A)$ be the derived algebra of $\mathfrak{g}(A)$, \mathfrak{c} the center of $\mathfrak{g}'(A)$, and $\bar{\mathfrak{g}}(A) = \mathfrak{g}'(A)/\mathfrak{c}$. Thus,*

- (1) *If $a_{ii} \neq 0$, $1 \leq i \leq n$, then $\dim \mathrm{HL}^2(\mathfrak{g}(A), \mathbb{K}) = (n - \ell)^2$;*
- (2) *If A is an indecomposable generalized Cartan matrix of affine $X_n^{(r)}$ type, then*

$$\mathrm{HL}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) = (\Omega_{R|\mathbb{K}}^1(r))^*, \quad \mathrm{HL}^2(\mathfrak{g}'(A), \mathbb{K}) = (\Omega_{R|\mathbb{K}}^1(r))^*.$$

- (3) *If A is an indecomposable generalized Cartan matrix of non-affine type, then*

$$\mathrm{HL}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \mathfrak{c}^*, \quad \mathrm{HL}^2(\mathfrak{g}'(A), \mathbb{K}) = 0.$$

Proof. (1) Theorem 3.2 [7] tells us that $\dim \mathrm{H}^1(\mathfrak{g}(A), (\mathfrak{g}(A))^*) = (n - \ell)^2$ under the assumption $a_{ii} \neq 0$ for $1 \leq i \leq n$. Then Theorem 3.5 gives the result.

(2) Let A be an indecomposable generalized Cartan matrix of affine $X_n^{(r)}$ type. By Gabber-Kac's radical Theorem (Theorem 9.11 and remarks in [12], pp. 159), $\mathfrak{g}'(A)$ and $\bar{\mathfrak{g}}(A)$ can be presented in term of $3n$ generators f_i, h_i, e_i ($1 \leq i \leq n$) and the Chevalley-Serre relations. By Theorem 3.17 in [9]¹, which says $\mathrm{HL}_2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \Omega_{R|\mathbb{K}}^1(r)$, together with Lemma 4.16, we get $\mathrm{HL}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) = (\Omega_{R|\mathbb{K}}^1(r))^*$.

Because $\mathfrak{g}'(A)$ is the universal covering of $\bar{\mathfrak{g}}(A)$ in the category of Lie algebras (see [22]), we have the following exact sequence:

$$0 \longrightarrow \mathrm{H}_2(\bar{\mathfrak{g}}(A), \mathbb{K}) \longrightarrow \mathfrak{g}'(A) \longrightarrow \bar{\mathfrak{g}}(A) \longrightarrow 0,$$

where $\mathrm{H}_2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \Omega_{R|\mathbb{K}}^1(r)/\Omega_{R|\mathbb{K}}^1(r) = t^{-1} dt$. Lemma 4.17 then yields

$$\mathrm{HL}_2(\mathfrak{g}'(A), \mathbb{K}) = \ker\{\mathrm{HL}_2(\bar{\mathfrak{g}}(A), \mathbb{K}) \rightarrow \mathrm{H}_2(\bar{\mathfrak{g}}(A), \mathbb{K})\} = \Omega_{R|\mathbb{K}}^1(r).$$

So Lemma 4.16 gives the second result.

(3) Let A be an indecomposable generalized Cartan matrix of non-affine type. Recall Theorem 3.1 [2], which says that $\bar{\mathfrak{g}}(A)$ possesses a non-degenerate invariant symmetric bilinear form ϕ if and only if A is symmetrizable. Moreover, such forms on $\bar{\mathfrak{g}}(A)$ are unique up to scalars.

If A is symmetrizable, then Berman's result above means $\dim \mathrm{B}(\bar{\mathfrak{g}}(A), \mathbb{K}) \geq 1$. On the other hand, for any $0 \neq \psi \in \mathrm{B}(\bar{\mathfrak{g}}(A), \mathbb{K})$, ψ is necessarily non-degenerate because $\bar{\mathfrak{g}}(A)$ is simple (by Theorem 4.3 [12], Exercise 1.4 [12]). So Berman's result above insures $\mathrm{B}(\bar{\mathfrak{g}}(A), \mathbb{K}) = \mathbb{K}\phi$. The non-degeneracy of ϕ naturally satisfies the nontrivial property of its restriction to any $\mathfrak{sl}(2, \mathbb{K})$ -copy. Lemma 4.15 shows $\mathrm{HL}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \mathrm{H}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \mathfrak{c}^*$ (the 2nd “=” was proved by Theorem 2.3 [2]).

If A is non-symmetrizable, there exists no non-degenerate invariant symmetric bilinear form on $\bar{\mathfrak{g}}(A)$. Since $\bar{\mathfrak{g}}(A)$ is simple (by Theorem 4.3 [12], Exercise 1.4 [12]), there is no invariant symmetric bilinear form on $\bar{\mathfrak{g}}(A)$, i.e., $\mathrm{B}(\bar{\mathfrak{g}}(A), \mathbb{K}) = 0$. Then Corollary 3.3 or Lemma 4.15 gives $\mathrm{HL}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \mathrm{H}^2(\bar{\mathfrak{g}}(A), \mathbb{K}) = \mathfrak{c}^*$ (the 2nd “=” was proved by Theorem 2.3 [2]).

Furthermore, since $\mathfrak{g}'(A)$ is the universal Leibniz central extension of $\bar{\mathfrak{g}}(A) = \mathfrak{g}'(A)/\mathfrak{c}$. Consequently, $\mathrm{HL}^2(\mathfrak{g}'(A), \mathbb{K}) = 0$. \square

¹Note that author in [9] used the different notations.

Remark 4.19. *Theorem 4.18 (2), (3) give a criterion to distinguish between affine and non-affine Kac-Moody algebras by means of the vanishing property of the 2nd Leibniz cohomology groups of $\mathfrak{g}'(A)$ with trivial coefficients, where the homological versions of Theorem 4.18 (2) and the 2nd statement of (3) were due to Gao ([9]). However, strictly speaking, owing to Gabber-Kac's Theorem (see [12]), the definition of $\mathfrak{g}'(A)$ for non-symmetrizable cases adopted by Gao is different from ours used here since our $\mathfrak{g}'(A)$ in Kac's notation is the quotient of the former. So in this sense, we get the same result for the quotient object.*

5. QUADRATIC LEIBNIZ ALGEBRAS AND THEIR CENTRAL EXTENSIONS

5.1. Quadratic Leibniz algebra.

Definition 5.1. *(\mathfrak{g}, ϕ) is called a quadratic Leibniz algebra if ϕ is a symmetric invariant bilinear form on Leibniz algebra \mathfrak{g} .*

Lemma 5.2. *Let \mathfrak{g} be a Lie algebra. If (\mathfrak{g}, ϕ) is a quadratic Leibniz algebra and d is a derivation of \mathfrak{g} , then $f(x, y) := \phi(x, dy)$ is a Leibniz 2-cocycle on \mathfrak{g} .*

Proof. Since $f(x, [y, z]) = \phi(x, d[y, z]) = f([x, y], z) - f([x, z], y)$ for $x, y, z \in \mathfrak{g}$, f is a Leibniz 2-cocycle on \mathfrak{g} . \square

Corollary 5.3. *([12]) If d is a skew-derivation, i. e., $\phi(dx, y) + \phi(x, dy) = 0$, then $f(x, y) := \phi(x, dy)$ is a Lie 2-cocycle on \mathfrak{g} .*

5.2. A negative answer to a question in [16].

Proposition 5.4. *Assume that \mathfrak{g} is a finite dimensional simple Lie algebra over \mathbb{K} . Construct the Lie algebra $\mathfrak{g} \otimes \mathbb{K}((t))$ with bracket*

$$[x \otimes r, y \otimes s]' = [x, y] \otimes rs, \quad x, y \in \mathfrak{g}, r, s \in \mathbb{K}((t)).$$

Then $\mathfrak{g} \otimes \mathbb{K}((t))$ is a simple Lie algebra over \mathbb{K} and

$$H^2(\mathfrak{g} \otimes \mathbb{K}((t)), \mathbb{K}) \subsetneq HL^2(\mathfrak{g} \otimes \mathbb{K}((t)), \mathbb{K}).$$

Proof. Define $\phi(x \otimes r, y \otimes s) = (x, y) \text{Res}(rs)$, where Res is a linear function on $\mathbb{K}((t))$ and takes the coefficient of t^{-1} for every series, $(,)$ is the Killing form on \mathfrak{g} . Then ϕ is an invariant symmetric bilinear form on $\mathfrak{g} \otimes \mathbb{K}((t))$ and $t^k \frac{d}{dt}$, $k \in \mathbb{Z} - \{0\}$ is a derivation of $\mathbb{K}((t))$. By Lemma 5.2, we get a non-trivial Leibniz 2-cocycle of the Lie algebra $\mathfrak{g} \otimes \mathbb{K}((t))$:

$$f(x \otimes \sum_{m \geq N} a_m t^m, y \otimes \sum_{n \geq N} b_n t^n) = (x, y) \sum_{m, n \geq N} n a_m b_n \delta_{m+n+k, 0}, \quad (5.1)$$

where f is well-defined since the summation in (5.1) is finite, and f is not skew symmetric, that is, $H^2(\mathfrak{g} \otimes \mathbb{K}((t)), \mathbb{K}) \subsetneq HL^2(\mathfrak{g} \otimes \mathbb{K}((t)), \mathbb{K})$. \square

Remark 5.5. *$\mathfrak{g} \otimes \mathbb{K}((t))$ is a simple Lie algebra (see [24]), and $\dim H^2(\mathfrak{g} \otimes \mathbb{K}((t)), \mathbb{K}) = 1$, but $\dim HL^2(\mathfrak{g} \otimes \mathbb{K}((t)), \mathbb{K}) = \infty$, which then leads to a negative answer to a question posed in [16].*

6. APPENDIX: AN ELEMENTARY PROOF OF THE EXACT SEQUENCE

Proof of Proposition 3.2. We shall prove the following

- (1) The map g is well-defined, that is, $g(\alpha) = g(\alpha + d^1\beta) \in \mathbf{B}(\mathfrak{g}, \mathbb{K})$ for any $\alpha \in \mathbf{HL}^2(\mathfrak{g}, \mathbb{K})$ and $\beta \in \mathbf{HL}^1(\mathfrak{g}, \mathbb{K})$ since $g(d^1\beta) \equiv 0$ by definition. Clearly, $g(\alpha)$ is symmetric. Since, for any $x, y, z \in \mathfrak{g}$,

$$\begin{aligned} g(\alpha)([x, y], z) &= \alpha([x, y], z) + \alpha(z, [x, y]) \\ &= \alpha(x, [y, z]) + \alpha([x, z], y) + \alpha([z, x], y) - \alpha([z, y], x) \\ &= \alpha(x, [y, z]) + \alpha([y, z], x) \\ &= g(\alpha)(x, [y, z]), \end{aligned}$$

$g(\alpha)$ is \mathfrak{g} -invariant.

- (2) $\text{Im}(f) = \ker(g)$. For any $\alpha \in \mathbf{H}^2(\mathfrak{g}, \mathbb{K})$ and any $x, y \in \mathfrak{g}$, we have

$$g(f(\alpha))(x, y) = g(\alpha)(x, y) = \alpha(x, y) + \alpha(y, x) = 0,$$

since α is skew symmetric. Then $\text{Im}(f) \subseteq \ker(g)$. On the other hand, for any $\beta \in \ker(g)$, we have $g(\beta)(x, y) = 0$, which implies that $\alpha(x, y) + \alpha(y, x) = 0$ for any $x, y \in \mathfrak{g}$. Then α is a skew symmetric Leibniz 2-cocycle, namely, a Lie 2-cocycle in $\mathbf{H}^2(\mathfrak{g}, k)$. Hence, we have $\ker(g) \subseteq \text{Im}(f)$.

- (3) $\text{Im}(g) = \ker(h)$. Now, $\forall \alpha \in \mathbf{HL}^2(\mathfrak{g}, \mathbb{K})$, define a bilinear form α' on \mathfrak{g} by

$$\alpha'(x, y) = \alpha(y, x),$$

for any $x, y \in \mathfrak{g}$. Denote $\beta := \frac{\alpha + \alpha'}{2}$ and $\gamma := \frac{\alpha - \alpha'}{2}$. Then β is symmetric and γ is skew symmetric. Also, we have $\alpha = \beta + \gamma$.

Now, for any $x, y, z \in \mathfrak{g}$,

$$\begin{aligned} (h(g(\alpha)))(x, y, z) &= g(\alpha)([x, y], z) = \alpha([x, y], z) + \alpha(z, [x, y]) \\ &= \alpha([x, y], z) + \alpha([z, x], y) + \alpha([y, z], x) \\ &= \alpha(z, [x, y]) + \alpha(y, [z, x]) + \alpha(x, [y, z]) \\ &\quad - \alpha([x, y], z) - \alpha([z, x], y) - \alpha([y, z], x) \\ &= 2\gamma(z, [x, y]) + 2\gamma(y, [z, x]) + 2\gamma(x, [y, z]) \\ &= \delta^2(2\gamma)(x, y, z), \end{aligned}$$

which implies that $h(g(\alpha))$ is a 3-coboundary in $\Omega^3(\mathfrak{g}, \mathbb{K})$. Then $\text{Im}(g) \subseteq \ker(h)$. Conversely, for any $\omega \in \ker(h)$, there exists an element $\eta \in \Omega^2(\mathfrak{g}, \mathbb{K})$ such that $\delta^2(\eta) = h(\omega)$. Namely, for any $x, y, z \in \mathfrak{g}$, there holds

$$\eta(z, [x, y]) + \eta(y, [z, x]) + \eta(x, [y, z]) = \omega([x, y], z).$$

Then

$$\omega([x, y], z) + \eta([x, y], z) = \omega(x, [y, z]) + \eta(x, [y, z]) + \omega([x, z], y) + \eta([x, z], y),$$

since $\omega(x, [y, z]) + \omega([x, z], y) = 0$. Let $\rho = \frac{1}{2}(\omega + \eta)$. Then $g(\rho) = \omega$ and

$$\rho([x, y], z) = \rho(x, [y, z]) + \rho([x, z], y),$$

which means that $\rho \in \mathbf{HL}^2(\mathfrak{g}, \mathbb{K})$. Hence $\ker(h) \subseteq \text{Im}(g)$. \square

ACKNOWLEDGMENT

The authors are indebted to the referee for the helpful comments. NH would like to express his thanks to the ICTP for the support and hospitality when he visited the ICTP Mathematics Group, Trieste, Italy, from March 1st to August 31st, 2006. YP is grateful to Hagiwara and Papi for sending him their papers ([11], [20]).

REFERENCES

- [1] E. Arbarello, C. De Concini, V. Kac, C. Procesi, *Moduli spaces of curves and representation theory*, Comm. Math. Phys. **117** (1), (1988), 1–36.
- [2] S. Berman, *On the low-dimensional cohomology of some infinite-dimensional simple Lie algebras*, Pacific J. Math. **83** (1), (1979), 27–36.
- [3] R. Borcherds, *Central extensions of generalized Kac-Moody Lie algebras*, J. Algebra, **140**, (1991), 330–335.
- [4] J.M. Casas, E. Faro, A.M. Vieites, *Abelian extensions of Leibniz algebras*, Comm. Algebra, **27** (6), (1999), 2833–2846.
- [5] D. Dokovic, K. Zhao, *Some infinite-dimensional simple Lie algebras in characteristic 0 related to those of Block*, J. Pure and Applied Algebra, **127** (2), (1998), 153–165.
- [6] R. Farnsteiner, *Central extensions and invariant bilinear forms of graded Lie algebras*, Algebras, Groups and Geom. **3**, (1986), 431–455.
- [7] R. Farnsteiner, *Derivation and central extensions of finitely generated graded Lie algebras*, J. Algebra, **118**, (1988), 33–45.
- [8] Y. Gao, *Central extensions of nonsymmetrizable Kac-Moody algebras over commutative algebras*, Proc. Amer. Math. Soc. **121**, (1994), 67–76.
- [9] Y. Gao, *The second Leibniz homology group for Kac-Moody Lie algebras*, Bull. of Lond. Math. Soc. **32**, (2000), 25–33.
- [10] A.V. Gnedbaye, *Third homology groups of universal central extensions of a Lie algebra*, Afrika Matematika Serie 3, 10 (1998). (ICTP Preprint No.IC98059).
- [11] Y. Hagiwara, T. Mizutani, *Leibniz algebras associated with foliations*, Kodai Math. J. **25** (2), (2002), 151–165.
- [12] V. Kac, *Infinite Dimensional Lie Algebras*, Cambridge University Press, third edition, 1990.
- [13] E. Kirkman, C. Procesi, L. Small, *A q -analog for the Virasoro algebra*, Comm. Algebra. **22**, (1994), 3755–3774.
- [14] J.S. Koszul, *Homologie et cohomologie des algèbres de Lie*, Bull. Soc. Math. France **78** (1950) 65–127.
- [15] W. Li, *2-Cocycles on the algebras of differential operators*, J. Algebra. **122**, (1989), 64–80.
- [16] D. Liu, N. Hu, *Leibniz central extensions on some infinite-dimensional algebras*, Comm. Algebra. **32** (6), (2004), 2385–2405.
- [17] J. Loday, *Cyclic Homology*, vol. **301**, Grund. Math. Wiss., Springer-Verlag, 1992.
- [18] J. Loday, T. Pirashvili, *Universal enveloping algebras of Leibniz algebras and (co)homology*, Math. Ann. **296** (1), (1993), 139–158.
- [19] J.M. Lodder, *Leibniz cohomology for differentiable manifolds*, Ann. Inst. Fourier, Grenoble, **48** (1), (1998), 73–95.
- [20] P. Papi, *Cohomology of tensor product of quantum planes*, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei **9**, Mat. Appl. 3 (1), (1992), 5–13.
- [21] T. Pirashvili, *On Leibniz homology*, Ann. Inst. Fourier (Grenoble) **44** (2), (1994), 401–411.
- [22] R. L. Wilson, *Euclidean Lie algebras are universal central extensions*, *Lie Algebras and Related Topics*, Lecture Notes in Math. 933 (ed. David Winter, Springer, New York, 1982) 210–213.
- [23] K. Zhao, *Automorphism groups of the algebras of differential operators*, J. of Capital Normal Uni. **15**, (1994), 1–7.
- [24] L. Zhu, D. Meng, *Commutative associative algebras and quadratic Lie algebras*, Comm. Algebra **29**, (2001), 2249–2268.
- [25] L. Zhu, D. Meng, *Structure of degenerate Block algebras*, Algebra Colloq. **10**, (2003), 53–62.

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200062, PR
CHINA

E-mail address: `nhhu@euler.math.ecnu.edu.cn` & `nhu@ictp.it` (until August 31, 2006)

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200062, PR
CHINA

E-mail address: `peiyufeng@gmail.com`

DEPARTMENT OF MATHEMATICS, JIAOTONG UNIVERSITY, SHANGHAI 200030, PR CHINA

E-mail address: `dliu@sjtu.edu.cn`