

Some Remarks on the Local Fundamental Group Scheme

V.B. Mehta and S. Subramanian

School of Mathematics

Tata Institute of Fundamental Research

Homi Bhabha Road

Mumbai 400 005

India.

e-mail: vikram@math.tifr.res.in

subramnn@math.tifr.res.in

Introduction

The fundamental group - scheme was introduced by M. Nori in [3,4]. In the same paper he had made 2 conjectures [4]. The first conjecture was proved by the present authors in [2]. In an attempt to prove the second conjecture, we had introduced the “local fundamental group-scheme” which is defined only in characteristic p . In this note, we gather together some necessary and sufficient conditions for it to base-change “correctly” from one algebraically closed field to another algebraically closed field of characteristic p .

Section I

We define the local fundamental group scheme, denoted by π^{loc} , as follows:

Let X be a smooth projective variety over an algebraically closed field of characteristic p , and let $F : X \rightarrow X$ be the Frobenius map. Let $FT(X)$ denote the category of all vector bundles V on X such that $F^{t*}(V)$ is *trivial* for some integer t . Fix a base point $x \in X$ and let $T : FT(X) \rightarrow \text{Vect } k$ be given by $V \rightarrow V_x$. One checks easily that $FT(X)$, with the fibre functor T , is a Tannaka category [1]. The corresponding affine group - scheme is defined to be the *local fundamental group-scheme* of X , denoted by $\pi^{loc}(X)$. Note that $\pi^{loc}(X)$ is defined only in characteristic p .

Remark 1 We have a canonical surjection $\pi(X) \rightarrow \pi^{loc}(X)$, where $\pi(X)$ is the fundamental group scheme as defined by Nori in [4].

Remark 2 Recall that all varieties are smooth and projective, and defined over algebraically closed fields of characteristic p . Now we have

Lemma 1.1 a) If $V \in FT(X)$, then V is semistable for any polarization H on X .

b) If V is stable for 1 polarization, then V is stable with respect to any polarization.

Proof a If $F^{t*}(V)$ is a trivial vector bundle on X for some t , then it is semistable for any polarization. It follows that V itself is semistable with respect to any polarization.

b) If $V \in FT(X)$, then there exist

1) a principal G -bundle

$$E \rightarrow X$$

where G is a finite local group-scheme, and E is *reduced* [4]

2)

and a representation

$$\sigma : G \rightarrow GL(n), n = \text{rank } V,$$

such that $V \cong V_\sigma$. It follows easily that subbundles W of V with $\mu(W) = \mu(V)$ are given by G subspaces W_1 of k^n , hence if $V \in FT(X)$ and stable with respect to one polarization, then it is stable with respect to any polarization.

Definition 1.2: Fix positive integers r and t . Let $S(X, r, t)$ denote the *set of isomorphism classes* of stable vector bundles V on X such that $\text{rank } V = r$ and $F^{t*}(V)$ is trivial. We shall show that $S(X, r, t)$ corresponds bijectively to the *closed* points of a scheme $M(X, r, t)$ defined over k .

Remark 3 : We recall that all fields are algebraically closed of characteristic p and all curves are assumed smooth and projective.

Theorem : The following statements are equivalent.

1)

$$g : \pi^{\text{loc}}(X_{k'}, x) \rightarrow \pi^{\text{loc}}(X_k, x) \times_k \text{spec } k'.$$

is an isomorphism, for every $k' \supset k$

2) $S(X; r, t)$ is a finite set, for all r and t .

3) For every $k' \supset k$, every $V \in S(X_{k'}; r, t,)$ descends to X , i.e. is defined over k .

Proof: Before we prove that the statements are equivalent, we discuss in brief some generalities on Tannaka categories. Let \mathcal{C}' be the Tannaka category of F -trivial vector bundles on X_k . Then \mathcal{C}' is isomorphic to the Tannaka category of rational, finite dimensional representations of $\pi^{\text{loc}}(X, x)$. We consider the category \mathcal{C}'_1 defined as follows:

a vector bundle V on $X_{k'}$ is an object of \mathcal{C}'_1 if there is an F -trivial vector bundle V_1 on X_k such that V is a degree zero subquotient of $V_1 \otimes_k k'$ on $X_{k'}$. The morphisms of \mathcal{C}'_1 are vector bundle homomorphisms and it is easily seen that with the tensor product of vector bundles as the tensor operation in the category and the fibre functor which associates to a bundle V its fibre over the base point x of $X_{k'}$, \mathcal{C}'_1 is a Tannaka category.

Let R' denote the Tannaka category of rational, finite dimensional representations of $\pi^{\text{loc}}(X_k, x) \times_k \text{spec } k'$ over k' . We can define a functor from R' to \mathcal{C}'_1 as follows:

If V is an object of R' namely a representation of $\pi^{\text{loc}}(X_k, x) \times_k \text{Spec } k'$, then there is a finite, local group scheme G over k and a surjection $\pi^{\text{loc}}(X_k, x) \rightarrow G$ such that V is actually a representation of $G_{k'} = G \times_k \text{spec } k'$. The homomorphism $\pi^{\text{loc}}(X_k, x) \rightarrow G$ defines a principal G -bundle $\pi : E \rightarrow X_k$ on X_k . Let $\pi : E_{k'} \rightarrow X_{k'}$ be the base change of this bundle to k' so that $E_{k'} \rightarrow X_{k'}$ is a principal $G_{k'}$ bundle. The representation V of $G_{k'}$ associates to this principal bundle a vector bundle (which is F -trivial) denoted by \tilde{V} . The $G_{k'}$ module V is a subquotient of $k'[G]^{\oplus n}$ where $k'[G]$ denotes the coordinate ring of $G_{k'}$ (see). As $k'[G] = k[G] \otimes_k k'$ where $k[G]$ denotes the coordinate ring of G over k , if we let V_1 be the vector bundle on X_k associated to the G -bundle $\pi : E \rightarrow X_k$ by the representation of G on $k[G]^{\oplus n}$, we obtain that \tilde{V} is a de-

gree zero subquotient of $V_1 \otimes_k k'$. Hence \tilde{V} is an object of \mathcal{C}'_1 . It is easily seen that the functor from R' to \mathcal{C}'_1 defined as above is an equivalence of Tannaka categories, so the affine group scheme associated to \mathcal{C}'_1 is $\pi^{\text{loc}}(X_k, x) \times_k \text{spec } k'$.

Let \mathcal{D}' denote the Tannaka category of all F -trivial vector bundles on $X_{k'}$. Since every object of \mathcal{C}'_1 is F -trivial \mathcal{C}'_1 is a Tannaka subcategory of \mathcal{D}' . The affine group scheme associated to \mathcal{D}' is $\pi^{\text{loc}}(X_{k'}, x)$ and the functor $\mathcal{C}'_1 \subset \mathcal{D}'$ induces the canonical homomorphism

$$g : \pi^{\text{loc}}(X_{k'}, x) \rightarrow \pi^{\text{loc}}(X_k, x) \times_k \text{spec } k'.$$

An application of Proposition 5, p.121 in [4] shows that g is surjective. Now we prove 1) ,2) and 3) are equivalent.

$1 \Rightarrow 3$ If g is an isomorphism, then it is, in particular, a closed immersion, and arguing as in Proposition (3.1) of [2], we see that any stable F -trivial bundle over $X_{k'}$ descends to X_k .

$3 \Rightarrow 1$ We now assume that every stable F -trivial bundle on $X_{k'}$ is defined over X_k . We have to show that g is a closed immersion. We use Proposition (2.21) (b) in [] for this purpose. We have to show that any object

of \mathcal{D}' , namely an F -trivial bundle on $X_{k'}$, is a subquotient of an object of \mathcal{C}'_1 . Since any F -trivial vector bundle on $X_{k'}$ is semistable, it has a filtration by stable F -trivial bundles on $X_{k'}$. By hypothesis, the stable F -trivial bundles descend to X_k , and we are left with an extension of stable bundles. For simplicity, let V be an object of \mathcal{D}' , which is an extension.

$$0 \rightarrow V_1 \rightarrow V \rightarrow V_2 \rightarrow 0$$

where V_1 and V_2 are stable F -trivial bundles on $X_{k'}$. The extension bundle V defines an element of $Ext_{\mathcal{O}_{X_{k'}}}^1(V_2, V_1)$. However, we have

$$Ext_{\mathcal{O}_{X_{k'}}}^1(V_2, V_1) = Ext_{\mathcal{O}_{X_k}}^1(V_2, V_1) \otimes_k k'.$$

Thus, the element representing V in $Ext_{\mathcal{O}_{X_{k'}}}^1(V_2, V_1)$ can be written as $\sum_i \alpha_i e_i$, where $\alpha_i \in k'$ and $e_i \in Ext_{\mathcal{O}_{X_k}}^1(V_2, V_1)$. By the construction of Baer sum in Ext^1 (see Definition (3.4.4), p.78, in [5]), we see that $\sum \alpha_i e_i$ represent a subquotient of extensions of V_2 by V_1 on X_k . Hence we obtain that V is a subquotient of a F -trivial bundle defined over X_k . The case where there are more than two stable factors in the stable filtration of V is similar.

2 \Leftrightarrow 3 Let O_t be the open subset of R^s (for the definition of R^s , see §4, p.141 in [6]) defined by

$O_t = \{x \in R^s \mid F^{t*}(U_x) \text{ is semistable on } X \text{ where } F \text{ is the Frobenius of } X\}$
. Let Y_t be the closed subset of O_t defined by $\{y \in O_t \mid F^{t*}U_y \text{ is trivial}\}$ and we give it the reduced subscheme structure. As O_t is open and $GL(N)$ invariant in R^s , the quotient $O_t/GL(W)$ exist as a scheme. Since Y_t is closed and $GL(N)$ invariant in O_t , the quotient $Y_t/GL(N)$ exists as a scheme, and let this scheme be denoted by $M(X, r, t)$. The k -rational points of $M(X, r, t)$ are in bijection with $S(X, r, t)$. It is further clear that $k' \supset k$ the schemes O_t, Y_t and $M(X, r, t)$ base change to the corresponding schemes over k' . In particular,

$$M(X_{k'}, r, t) \cong M(X_k, r, t) \times_k \text{spec } k'.$$

In general, one always has

$$M(X_{k'}, r, t)(k) \subset M(X_{k'}, r, t)(k').$$

If strict equality holds, $M(X, r, t)$ is the spectrum of an Artin ring on k . In

that case, every element of $S(X_{k'}, r, t)$ is defined over k . Further, we see that if every element of $S(X_{k'}, r, t)$ is defined over k , then $S(X, r, t)$ is finite. \square

References

- [1] P. Deligne, J. Milne: Tannakian categories in Hodge cycles, Motives and Shimura varieties, *Lecture Notes in Mathematics* 900, Springer.
- [2] V.B. Mehta, S. Subramanian: On the fundamental group scheme, *Inv. Math.* 148, 143-150 (2002).
- [3] M.V. Nori, Representations of the fundamental group, *Compositio Math.* 33, 1976.
- [4] M.V. Nori, The fundamental group scheme, *Proc. Indian Acad. Sci. (Math. Sci)* 91, 1982.
- [5] Charles A. Weibel, An introduction to homological algebra, *Cambridge university Press*, 1994.
- [6] P.E. Newstead, Introduction to Moduli problems and Orbit spaces, *TIFR Lecture Notes*, 1978.