

# Detailed Proof of Two Dimensional Jacobian Conjecture<sup>1</sup>

Yucai Su

*Department of Mathematics, University of Science and Technology of China*

*Hefei, Anhui 230026, China*

*Email: ycsu@ustc.edu.cn*

*Dedicated to my teacher Professor Zhexian Wan for his 80th birthday*

**Abstract.** We give a full proof of the two dimensional Jacobian conjecture.

## 1. INTRODUCTION

For any field  $\mathbb{F}$  of characteristic zero, it is a well known fact that if  $n$  polynomials  $f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n) \in \mathbb{F}[x_1, \dots, x_n]$  are generators of the polynomial ring  $\mathbb{F}[x_1, \dots, x_n]$ , then the *Jacobian determinant*

$$J(f_1, \dots, f_n) = \det A \in \mathbb{F}^* = \mathbb{F} \setminus \{0\}, \quad (1.1)$$

is a nonzero constant, where  $A = (\frac{\partial f_i}{\partial x_j})_{i,j=1}^n$  is the  $n \times n$  *Jacobian matrix* of  $f_1, \dots, f_n$ .

One of the major unsolved problems of mathematics [S] (see also [B, CM, V2]), viz. the *Jacobian conjecture*, states that the reverse of the above statement also holds, namely, if the Jacobian determinant  $J(f_1, \dots, f_n) \in \mathbb{F}^*$ , then  $f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n) \in \mathbb{F}[x_1, \dots, x_n]$  are generators of  $\mathbb{F}[x_1, \dots, x_n]$ . For convenience, if (1.1) holds, we shall refer  $f_1, \dots, f_n$  to as polynomials with *nonzero Jacobian determinant property* (or simply, *NJDP*).

This conjecture relates to many aspects of mathematics [A, ES, H, R, SW, SY] and has attracted great attention in mathematics and physics literature during the past 60 years and there have been a various ways of approaches toward the proof or disproof of this conjecture (here we simply give a short random list of references [BCW, CCS, D, J, K, Ki, KM, M1, V1, V2, W]). Hundreds of papers have appeared in connection with this conjecture, even for the simplest case  $n = 2$  [AO, N, No]. However this conjecture remains unsolved even for the case  $n = 2$ . The difficulty in solving this conjecture probably lies in that although the NJDP may contain much information, one is unable to use it.

In this self-contained paper, we give a proof of the Jacobian conjecture for the case  $n = 2$ . The main result is the following.

**Theorem 1.1** *Let  $\mathbb{F}$  be any field of characteristic zero.*

- (1) *Two polynomials  $F(x, y), G(x, y) \in \mathbb{F}[x, y]$  are generators of  $\mathbb{F}[x, y]$  if and only if its Jacobian determinant*

$$J(F, G) = \begin{vmatrix} \partial_x F & \partial_y F \\ \partial_x G & \partial_y G \end{vmatrix} \in \mathbb{F}^*, \quad (1.2)$$

---

<sup>1</sup>Supported by a NSF grant 10471096 of China, “One Hundred Talents Program” from University of Science and Technology of China and “Trans-Century Training Programme Foundation for the Talents” from National Education Ministry of China.

is a nonzero constant, where  $\partial_x, \partial_y$  stand for the partial derivatives  $\frac{\partial}{\partial x}, \frac{\partial}{\partial y}$  respectively.

- (2) The automorphism group  $\text{Aut } \mathbb{F}[x, y]$  is generated by  $\{\sigma_a, \phi_k, \tau \mid a \in \mathbb{F}^*, k \geq 1\}$ , where  $\sigma_a, \phi_k, \tau$  are automorphisms of  $\mathbb{F}[x, y]$  defined by

$$\begin{aligned} \sigma_a : x &\mapsto ax, & \phi_k : x &\mapsto x + y^k, & \tau : x &\mapsto y, \\ y &\mapsto y, & y &\mapsto y, & y &\mapsto x. \end{aligned} \tag{1.3}$$

Theorem 1.1(2) has been known before [V1] (however the known proofs were found to be not easy), we simply reproduce it as a by-product. The automorphism  $\tau$  is usually called the *twist automorphism*, and an automorphism of the form  $x \mapsto ax + f(y), y \mapsto y$  for some  $a \in \mathbb{F}^*, f(y) \in \mathbb{F}[y]$  is usually called a *triangular automorphism*. Thus  $\text{Aut } \mathbb{F}[x, y]$  is generated by triangular automorphisms and the twist.

Let us briefly explain the main points in the proof of Theorem 1.1 below: First by applying some automorphism, we can suppose  $F(x, y), G(x, y)$  are monic polynomials of  $y$  with coefficients in  $\mathbb{F}[x]$  (cf. (3.4)). We write  $G(x, y)$  as a *rational power series* of  $F(x, y)$  in (3.9). By introducing the *prime degree* of  $F(x, y)$  (cf. Definition 2.3), we are able to define the *leading polynomial*  $F_{\text{lead}}(x, y)$  and the *primary polynomial*  $F_{\text{prim}}(x, y)$  of  $F(x, y)$  (cf. Definitions 2.5 and 3.3). Then we introduce the *r-th components* of  $F(x, y)^{\frac{1}{m}}$  (cf. Definition 3.6) and prove that they are all rational functions under some condition (cf. Lemma 3.7). By showing that some component of  $G(x, y)$  must satisfy a differential equation (cf. (3.50) and Lemma 3.8), we can prove that the primary polynomial  $F_{\text{prim}}(x, y)$  has a form in Lemma 3.10. Then we prove in Lemma 3.11 (the key lemma) that Lemma 3.10(3) in fact cannot occur. Thus by applying some automorphism, we can reduce the degrees of  $F(x, y)$  and  $G(x, y)$  (cf. Lemma 3.9). Therefore the theorem is proved by induction on the degrees.

As a consequence of the proof of Lemma 3.11, we obtain

**Theorem 1.2** *Suppose  $F(x, y), G(x, y)$  satisfy (1.2) and  $\deg_y F(x, y) \geq \deg_x F(x, y)$ . Then  $F(x, y)$  is a monic polynomial of  $y$  (up to a nonzero scalar).*

## 2. PRELIMINARIES

Denote by  $\mathbb{Z}, \mathbb{Z}_+, \mathbb{N}, \mathbb{Q}$  the sets of integers, non-negative integers, positive integers, rational numbers respectively. Let  $A = \{a_1, \dots, a_n\} \subset \mathbb{Z}$ . We denote the greatest common divisor (g.c.d.) of  $A$  by  $(a_1, \dots, a_n)$  or  $(a \mid a \in A)$ , and the least common multiple (l.c.m) by  $[a_1, \dots, a_n]$  or  $[a \mid a \in A]$ . Denote  $\mathbb{F}(x, y) = \{\frac{P(x, y)}{Q(x, y)} \mid P(x, y), Q(x, y) \in \mathbb{F}[x, y]\}$ , the field of rational functions in two variables.

**Lemma 2.1** (cf. Remark 3.13) *Let  $H(x, y) \in \mathbb{F}[x, y] \setminus \mathbb{F}[x]$  (i.e., it is a polynomial on  $x$  and  $y$  but not a polynomial on  $x$ ). Let  $m \in \mathbb{N}$ . Suppose there exists a finite nonzero combination  $\sum_{i \in \mathbb{Z}} p_i(x) H(x, y)^{\frac{i}{m}} \in \mathbb{F}(x, y)$  (i.e., it is a rational function) for some  $p_i(x) \in \mathbb{F}[x]$ , where  $H(x, y)^{\frac{i}{m}}$  is regarded as a (possibly multi-valued) function on  $x$  and  $y$ . Then*

$H(x, y) = H_1(x, y)^{\frac{m}{d}}$  is the  $\frac{m}{d}$ -th power of some polynomial  $H_1(x, y) \in \mathbb{F}[x, y]$ , where  $d = (m, i \mid p_i(x) \neq 0)$  is the g.c.d. of  $\{m, i \mid p_i(x) \neq 0\}$ .

*Proof.* Let  $A = \{m, i \mid p_i(x) \neq 0\}$ . If  $A = \{m\}$ , there is nothing to prove. Suppose  $|A| > 1$ . If necessary by replacing  $i$  by  $\frac{i}{d}$ , we can suppose  $d = 1$ . If  $A = \{m, n\}$ , this means that  $H(x, y)^{\frac{n}{m}} = \frac{R(x, y)}{Q(x, y)}$  for some coprime polynomials  $R(x, y), Q(x, y)$ , then  $Q(x, y)^m H(x, y)^n = R(x, y)^m$ . Since  $\mathbb{F}[x, y]$  is a uniquely factorial domain, by decomposing each polynomial into the product of its irreducible polynomials, using  $(m, n) = 1$ , we see that  $H(x, y)$  is the  $m$ -th power of some polynomial.

Now suppose  $|A| > 2$ . Denote

$$P(x, y) = \sum_{i \in \mathbb{Z}} p_i(x) H(x, y)^{\frac{i}{m}}. \quad (2.1)$$

**Definition 2.2** Let  $n = \min\{i \mid p_i(x) \neq 0\}$ . We shall call  $P(x, y)$  in (2.1) a *combination of rational power of  $H(x, y)$  with leading term  $H(x, y)^{\frac{n}{m}}$*  or simply a *c.r.p. of  $H$  with l.t.  $H^{\frac{n}{m}}$* . For  $j, k \in \mathbb{Z}$ , we say the term  $p_j(x)H(x, y)^{\frac{j}{m}}$  is *higher than the term  $p_k(x)H(x, y)^{\frac{k}{m}}$*  if  $j > k$ .

Let us continue the proof of the lemma. Set

$$d_j(P) = (m, i \mid p_i(x) \neq 0, n \leq i \leq n + j) = \begin{cases} d_{j-1}(P) & \text{if } p_{n+j}(x) = 0, \\ (d_{j-1}(P), n + j) & \text{if } p_{n+j}(x) \neq 0, \end{cases} \quad (2.2)$$

for  $j = 0, 1, \dots$ , where we take  $d_{-1}(P) = 0$ . Then

$$(m, n) = d_0(P) > d_{j_1}(P) > d_{j_2}(P) > \dots > d_{j_s}(P) = 1, \quad (2.3)$$

where  $0 = j_0 < j_1 < \dots < j_s$  such that each  $j_r$  is the smallest integer satisfying  $d_{j_r}(P) \neq d_{j_r-1}(P)$ . In particular,

$$p_{n+j_r}(x) \neq 0 \quad \text{for } r = 0, 1, \dots, s. \quad (2.4)$$

**Claim 1.** For  $r = 0, 1, \dots, s$ , there exists a rational function  $P_r(x, y)$  which is a c.r.p. of  $H$  with l.t.  $H^{\frac{k_r m + j'_r}{m}}$  for some  $k_r, j'_r \in \mathbb{Z}_+$  such that  $j'_0 = n$ ,  $j'_r \leq j_r$  if  $r > 0$  and  $(m, j'_0, j'_1, \dots, j'_r) = d_{j_r}$ .

Obviously, for  $r = 0$ , we can take  $k_0 = 0$  and  $P_0(x, y) = P(x, y)$ . Suppose  $r > 0$ . We shall prove the claim by induction on  $j_r$ . Let us compute the coefficient  $\tilde{p}_{mn+j_r}(x)$  of  $H^{\frac{mn+j_r}{m}}$  in the rational function  $P(x, y)^m$ , which can be written as

$$P(x, y)^m = p_n(x)^m H(x, y)^{\frac{mn}{m}} + \sum_{i>0} \tilde{p}_{mn+i}(x) H(x, y)^{\frac{mn+i}{m}} \quad \text{for some } \tilde{p}_{mn+i}(x) \in \mathbb{F}[x], \quad (2.5)$$

i.e.,  $P(x, y)^m$  is a c.r.p. of  $H$  with l.t.  $H^{\frac{mn}{m}}$ . If a nonzero term  $p_{n+a_1}(x)H(x, y)^{\frac{n+a_1}{m}}$  of  $P(x, y)$  contributes to the computation of  $\tilde{p}_{mn+j_r}(x)$  for some  $0 < a_1 < j_r$ , then  $mn + j_r = (n + a_1) + \cdots + (n + a_m)$  for some  $0 \leq a_i < j_r$  with  $p_{n+a_i}(x) \neq 0$ , i.e.,

$$j_r = a_1 + \cdots + a_m. \quad (2.6)$$

But then  $d_{j_r}(P) = (d_{j_r-1}(P), a_1 + \cdots + a_m) = d_{j_r-1}(P)$ , a contradiction with the choice of  $j_r$ . Thus only two terms  $p_n(x)H(x, y)^{\frac{n}{m}}$  and  $p_{n+j_r}(x)H(x, y)^{\frac{n+j_r}{m}}$  contribute to  $\tilde{p}_{mn+j_r}(x)$  and in fact we have  $\tilde{p}_{mn+j_r}(x) = mp_n(x)^{m-1}p_{n+j_r}(x) \neq 0$  (cf. (2.4)). Now we take the rational function

$$\tilde{P}(x, y) = P(x, y)^m - p_n(x)^m H(x, y)^n = \sum_{i>0} \tilde{p}_{mn+i}(x) H(x, y)^{\frac{mn+i}{m}}. \quad (2.7)$$

Suppose the first nonzero term of  $\tilde{P}(x, y)$  is  $\tilde{p}_{mn+\tilde{n}}H(x, y)^{\frac{mn+\tilde{n}}{m}}$ , i.e.,  $\tilde{P}(x, y)$  is a c.r.p. of  $H$  with l.t.  $H^{\frac{mn+\tilde{n}}{m}}$ , and  $0 < \tilde{n} \leq j_r$  (cf. (2.5)). If  $\tilde{n} = j_r$ , we can take  $j'_r = j_r$  and the claim is proved. So suppose  $0 < \tilde{j}_r := j_r - \tilde{n} < j_r$ . Note that for  $1 \leq j < j_r$ , any nonzero term  $\tilde{p}_{mn+j}(x)H^{\frac{mn+j}{m}}$  of  $\tilde{P}(x, y)$  can be only contributed by nonzero terms  $p_{n+a_i}(x)H^{\frac{n+a_i}{m}}$  of  $P(x, y)$  with  $0 \leq a_i < j_r$  such that  $\sum_{i=1}^m a_i = j$ , thus as in the discussion of (2.6), we have

$$d_i(\tilde{P}) > d_{\tilde{j}_r}(\tilde{P}) \quad \text{if } i < \tilde{j}_r. \quad (2.8)$$

Using definitions (2.2) and (2.3), equation (2.8) means that if we replace  $P(x, y)$  by  $\tilde{P}(x, y)$  (and so  $n$  becomes  $\tilde{n}' := mn + \tilde{n}$ ) then the integer  $\tilde{j}_r$  is exactly the integer  $j_{r_1}$  for some  $r_1 \leq r$ . Thus the claim can be proved by induction on  $j_r$  (we remark that there may be more nonzero terms between  $H^{\frac{\tilde{n}'}{m}}$  and  $H^{\frac{\tilde{n}'+j'_r}{m}}$  in  $\tilde{P}(x, y)$  than nonzero terms between  $H^{\frac{n}{m}}$  and  $H^{\frac{n+j_r}{m}}$  in  $P(x, y)$ , but we only need the fact that  $j'_r < j_r$ ; furthermore,  $d_{j_r} = (d_{j_r-1}, j'_r)$ ).

From our choice of  $j_r$ , we have  $(m, j'_0, j'_1, \dots, j'_s) = d_{j_s} = 1$ . Thus there exist some integers  $a, a'_0, a'_1, \dots, a'_s$  such that  $am + \sum_{r=0}^s a'_r j'_r = 1$ . Then there exist positive integers  $a_r = b_r m + a'_r$  for some sufficient large integers  $b_r$ , such that  $\sum_{r=0}^s a_r (k_r m + j'_r) = km + 1$  for some  $k \in \mathbb{N}$ . By Claim 1, the rational function

$$Q(x, y) := \prod_{r=0}^s P_r(x, y)^{a_r} = q_1(x)H(x, y)^{\frac{km+1}{m}} + \sum_{i \geq 2} q_i(x)H(x, y)^{\frac{km+i}{m}}, \quad (2.9)$$

is a c.r.p. of  $H$  with l.t.  $H^{\frac{km+1}{m}}$  for some  $q_i(x) \in \mathbb{F}[x]$  with  $q_1(x) \neq 0$ . Now we re-denote the rational function  $P(x, y) = H(x, y)^{-k}Q(x, y)$ , which is a c.r.p. of  $H$  with l.t.  $H^{\frac{1}{m}}$ . Then we can always write  $P(x, y)$  as

$$P(x, y) = p_1(x)H(x, y)^{\frac{1}{m}} + p_2(x)H(x, y)^{\frac{2}{m}} + \cdots + p_{\ell m}(x)H(x, y)^{\frac{\ell m}{m}}, \quad (2.10)$$

for some  $p_i(x) \in \mathbb{F}[x]$  with  $p_1(x) \neq 0$ , where  $\ell$  is some sufficient large integer. For each  $1 \leq i \leq \ell m$ , we can express  $P(x, y)^i$  as (by writing  $H(x, y)^{\frac{\ell m + i}{m}}$  as  $H(x, y)^\ell H(x, y)^{\frac{i}{m}}$ )

$$P(x, y)^i = \sum_{j=1}^{\ell m} p_{i,j}(x, y) H(x, y)^{\frac{j}{m}}, \quad (2.11)$$

where  $p_{i,j}(x, y) \in \mathbb{F}[x, y]$  is some polynomial which is a combination of integral powers of  $H(x, y)$  with coefficients in  $\mathbb{F}[x]$ . Regarding (2.11) as a system of linear equations on the unknown variables  $H(x, y)^{\frac{j}{m}}$ ,  $j = 1, \dots, \ell m$ . If the determinant  $\Delta(x, y) = \det(p_{i,j}(x, y))$  of the matrix of coefficients is not the zero polynomial, then we can solve  $H(x, y)^{\frac{1}{m}}$  to obtain that it is a rational function, and we complete the proof of the lemma by the arguments in the first paragraph of the proof.

Thus suppose  $\Delta(x, y) = 0$ . Since  $\Delta(x, y) = \sum_{i \geq 0} \alpha_i(x) H(x, y)^i$  is a polynomial of  $H(x, y)$  with coefficients  $\alpha_i(x) \in \mathbb{F}[x]$  and the assumption of  $H(x, y) \notin \mathbb{F}[x]$  shows that  $\{H(x, y)^i \mid i \in \mathbb{Z}_+\}$  is  $\mathbb{F}[x]$ -linear independent, we must have  $\alpha_0(x) = 0$ . But by (2.10), one can easily compute that by modulo the subspace  $H(x, y)\mathbb{F}[x, y]$  (cf. definition “ $\equiv$ ” in (3.56)), we have  $P(x, y)^i \equiv p_1(x)^i H(x, y)^{\frac{i}{m}} + (\text{higher terms})$ , i.e., by modulo  $H(x, y)\mathbb{F}[x, y]$ , the matrix  $(p_{ij}(x, y))$  is upper-triangular with diagonals  $p_1(x)^i$ ,  $i = 1, \dots, \ell m$ . This shows that  $\alpha_0(x) = \prod_{i=1}^{\ell m} p_1(x)^i$ , which is a contradiction with  $\alpha_0(x) = 0$  and  $p_1(x) \neq 0$ . This proves the lemma.  $\square$

We shall work with the ring of meromorphic functions of  $y^{-1}$  over  $\mathbb{F}[x]$ :

$$\mathbb{F}[x]((y^{-1})) = \left\{ \sum_{i \in \mathbb{Z}} f_i(x) y^i \mid f_i(x) \in \mathbb{F}[x], f_i(x) = 0 \text{ if } i \gg 0 \right\}. \quad (2.12)$$

Any element of the form, where  $\ell \in \mathbb{Z}$ ,

$$H(x, y) = y^\ell + \sum_{i=1}^{\infty} h_i(x) y^{\ell-i}, \quad (2.13)$$

is called a *power series of  $y^{-1}$* .

**Definition 2.3** Let  $H(x, y)$  be as in (2.13). If  $\max_{1 \leq i < \infty} \frac{\deg h_i(x)}{i}$  exists, then the rational number

$$p(H) = \max_{1 \leq i < \infty} \frac{\deg h_i(x)}{i} \in \mathbb{Q}, \quad (2.14)$$

is called the *prime degree of  $H(x, y)$*  (obviously, we have  $p(H) = -\infty \iff H(x, y) = y^\ell$ ). Otherwise we set  $p(H) = +\infty$ .

The following arguments will illustrate the importance of introducing the notion of prime degree.

Let  $H(x, y)$  be as in (2.13). For any  $k \in \mathbb{Z}$ , we can uniquely expand  $H(x, y)^{\frac{k}{\ell}}$  as an element in  $\mathbb{F}[x]((y^{-1}))$ :

$$\begin{aligned} H(x, y)^{\frac{k}{\ell}} &= y^k \left( 1 + \left( \sum_{i=1}^{\infty} h_i(x) y^{-i} \right)^{\frac{k}{\ell}} \right) \\ &= y^k \sum_{j=0}^{\infty} \binom{\frac{k}{\ell}}{j} \left( \sum_{i=1}^{\infty} h_i(x) y^{-i} \right)^j = y^k + \sum_{i=1}^{\infty} h_{k, \ell, i}(x) y^{k-i} \in \mathbb{F}[x]((y^{-1})), \end{aligned} \quad (2.15)$$

where the coefficient of  $y^{k-i}$ , often denoted by  $\text{Coeff}(H(x, y)^{\frac{k}{\ell}}, y^{k-i})$  in this paper, is

$$h_{k, \ell, i}(x) = \sum_{\substack{r_1 + 2r_2 + \dots + ir_i = i \\ r_1, \dots, r_i \geq 0}} \binom{\frac{k}{\ell}}{r_1, r_2, \dots, r_i} h_1(x)^{r_1} \dots h_i(x)^{r_i}. \quad (2.16)$$

Here in general for any  $a \in \mathbb{F}$ ,  $r_1, \dots, r_i \in \mathbb{Z}_+$ ,

$$\binom{a}{r_1, r_2, \dots, r_i} = \frac{a(a-1) \dots (a - (r_1 + \dots + r_i) + 1)}{r_1! \dots r_i!}, \quad (2.17)$$

is a *multi-nomial coefficient*.

**Lemma 2.4** *Let  $H(x, y)$  be as in (2.13). Then for any  $k \in \mathbb{Z}$ ,  $p(H) = p(H^{\frac{k}{\ell}})$ .*

*Proof.* Let  $s = \min\{i \geq 1 \mid \deg h_i(x) = ip(H)\}$ . Then by (2.16), we see that  $\deg h_{k, \ell, i}(x) \leq ip(H)$  for  $i \geq 1$ , and  $\deg h_{k, \ell, s}(x) = sp(H)$ .  $\square$

For any  $H(x, y) \in \mathbb{F}[x, y] \setminus \mathbb{F}[y]$  as in (2.13), we always denote

$$H_{\text{lead}}(x, y) = y^\ell + \sum_{i=1}^{\infty} a_i x^{ip} y^{\ell-i} \quad (\text{which is clearly not equal to } y^\ell), \quad (2.18)$$

$$H_{\text{igno}}(x, y) = H(x, y) - H_{\text{lead}}(x, y) = \sum_{i=1}^{\infty} h_{0, i}(x) y^{\ell-i}, \quad \deg h_{0, i}(x) < ip, \quad (2.19)$$

where  $p = p(H)$  and  $a_i = \text{Coeff}(h_i(x), x^{ip})$  is the coefficient of  $x^{ip}$  in  $h_i(x)$  (here and below we always set  $a_i = 0$  if  $ip \notin \mathbb{Z}_+$ ), and  $h_{0, i}(x) = h_i(x) - a_i x^{ip}$ .

**Definition 2.5** Any polynomial (or any element in  $\mathbb{F}[x]((y^{-1}))$ ) of the form

$$\sum_{i=0}^{\infty} c_i x^{i_0 + ip} y^{\ell_0 - i} \quad \text{with} \quad c_i \in \mathbb{F}, \quad i_0, \ell_0 \in \mathbb{Z}_+,$$

is called a *quasi-homogenous polynomial of prime degree  $p$* . We shall call  $H_{\text{lead}}(x, y)$  the *leading polynomial of  $H(x, y)$*  and  $H_{\text{igno}}(x, y)$  the *ignored polynomial of  $H(x, y)$* .

We shall need the following easily verified results in the next section.

- Lemma 2.6** (1) *Any automorphism in (1.3) maps generators of  $\mathbb{F}[x, y]$  to generators of  $\mathbb{F}[x, y]$ , and maps polynomials with NJDP to polynomials with NJDP.*
- (2) *If  $F(x, y), G(x, y)$  satisfy (1.2), then  $F(x, y)$  is a square free polynomial, namely, there do not exist polynomials  $P(x, y), Q(x, y) \in \mathbb{F}[x, y]$  with  $P(x, y) \notin \mathbb{F}$  such that  $F(x, y) = P(x, y)^2 Q(x, y)$ .*

### 3. PROOF OF THE MAIN THEOREM

Let  $F(x, y), G(x, y)$  be two polynomials with NJDP. We shall prove Theorem 1.1(1) by induction on the pair  $(m, n)$  of positive integers, where

$$m = \deg F(x, y), \quad n = \deg G(x, y). \quad (3.1)$$

First we write  $F(x, y), G(x, y)$  as

$$F(x, y) = \sum_{i, j \in \mathbb{Z}_+ : i+j \leq m} f_{ij} x^i y^j, \quad G(x, y) = \sum_{i, j \in \mathbb{Z}_+ : i+j \leq n} g_{ij} x^i y^j, \quad (3.2)$$

for some  $f_{ij}, g_{ij} \in \mathbb{F}$ . Then

$$f(x) := \sum_{i=0}^m f_{i, m-i} x^i, \quad g(x) := \sum_{i=0}^n g_{i, n-i} x^i,$$

are nonzero polynomials on  $x$ . So we can choose some  $a \in \mathbb{F}$  such that  $f(a)g(a) \neq 0$ . Thus by applying the linear isomorphism of  $\mathbb{F}[x, y]$  (cf. Lemma 2.6(1)):

$$(x, y) \mapsto (x + ay, y), \quad (3.3)$$

$F(x, y), G(x, y)$  become polynomials of the forms in (3.2) with  $f_{0, m} g_{0, n} = f(a)g(a) \neq 0$ . Rescaling  $F(x, y), G(x, y)$ , we can assume  $f_{0, m} = g_{0, n} = 1$ . Thus we can rewrite  $F(x, y), G(x, y)$  as

$$F(x, y) = y^m + \sum_{i=1}^m f_i(x) y^{m-i}, \quad G(x, y) = y^n + \sum_{j=1}^n g_j(x) y^{n-j}, \quad (3.4)$$

for some  $f_i(x), g_j(x) \in \mathbb{F}[x]$ . For convenience, we denote  $f_0(x) = g_0(x) = 1$ .

We can assume  $m \leq n$ . By replacing  $y$  by  $y - \frac{1}{m} f_1(x)$ , i.e., by applying the automorphism

$$(x, y) \mapsto (x, y - \frac{1}{m} f_1(x)), \quad (3.5)$$

we can further suppose

$$f_1(x) = 0. \quad (3.6)$$

If either  $F(x, y)$  or  $G(x, y)$  does not depend on the variable  $x$ , then (1.2) forces  $F(x, y), G(x, y)$  to be generators of  $\mathbb{F}[x, y]$ . Thus we suppose

$$\text{both } F(x, y) \text{ and } G(x, y) \text{ depend on the variable } x. \quad (3.7)$$

If  $m = 1$ , then (3.6) shows that  $F(x, y) = y$  and (1.2) forces  $G(x, y) = ax + b$  for some  $a \in \mathbb{F}^*$ ,  $b \in \mathbb{F}$ . Thus  $F(x, y), G(x, y)$  are generators. Hence we can suppose

$$2 \leq m \leq n. \quad (3.8)$$

Note that for  $F(x, y), G(x, y)$  as in (3.4), we can expand  $G(x, y)$  as

$$G(x, y) = F(x, y)^{\frac{n}{m}} + \sum_{i=1}^{\infty} b_i(x) F(x, y)^{\frac{n-i}{m}} \text{ for some } b_i(x) \in \mathbb{F}[x], \quad (3.9)$$

where by comparing the coefficients of  $y^{n-i}$ , the polynomial  $b_i(x) \in \mathbb{F}[x]$  can be inductively determined by the following (cf. (2.16)):

$$b_i(x) = g_i(x) - \sum_{j=0}^{i-1} b_j(x) \sum_{\substack{r_1+2r_2+\dots+mr_m=i-j \\ r_1, \dots, r_m \geq 0}} \binom{\frac{n-j}{m}}{r_1, r_2, \dots, r_m} f_1(x)^{r_1} \cdots f_m(x)^{r_m} \text{ for } i \geq 1. \quad (3.10)$$

Here and below, we set  $b_0(x) = 1$  and  $g_i(x) = 0$  if  $i > n$ . Thus there exists a function on  $u$  and  $v$ :

$$\tilde{G}(u, v) = v^{\frac{n}{m}} + \sum_{i=1}^{\infty} b_i(u) v^{\frac{n-i}{m}}, \text{ such that } G(x, y) = \tilde{G}(x, F(x, y)). \quad (3.11)$$

Similarly, we can expand the polynomial  $y$  as

$$y = F(x, y)^{\frac{1}{m}} + \sum_{i=1}^{\infty} \bar{b}_i(x) F(x, y)^{\frac{1-i}{m}}, \quad (3.12)$$

where as in (3.10), we set  $\bar{b}_0(x) = 1$  and

$$\bar{b}_i(x) = - \sum_{j=0}^{i-1} \bar{b}_j(x) \sum_{\substack{r_1+2r_2+\dots+mr_m=i-j \\ r_1, \dots, r_m \geq 0}} \binom{\frac{1-j}{m}}{r_1, r_2, \dots, r_m} f_1(x)^{r_1} \cdots f_m(x)^{r_m} \text{ for } i > 0. \quad (3.13)$$

From (3.12), we obtain

$$\frac{1}{\partial_y F(x, y)} = \frac{1}{m} (F(x, y)^{\frac{1-m}{m}} + \sum_{i=1}^{\infty} \bar{f}_i(x) F(x, y)^{\frac{1-m-i}{m}}), \text{ where } \bar{f}_i(x) = (1-i)\bar{b}_i(x). \quad (3.14)$$

Throughout the paper, we denote  $p = p(F)$ , the prime degree of  $F(x, y)$ . Then (3.13) shows that  $\deg \bar{b}_i(x) \leq ip$  for  $i \geq 1$ . Thus,

$$\deg \bar{f}_i(x) \leq ip \text{ for } i \geq 1. \quad (3.15)$$

Using (3.11) in (1.2), we obtain

$$\begin{aligned} J(F, G) &= \begin{vmatrix} \partial_x F(x, y) & \partial_y F(x, y) \\ \partial_u \tilde{G}(u, v) + \partial_v \tilde{G}(u, v) \partial_x F(x, y) & \partial_v \tilde{G}(u, v) \partial_y F(x, y) \end{vmatrix} \\ &= -\partial_y F(x, y) \partial_u \tilde{G}(u, v) \in \mathbb{F}^*, \end{aligned}$$

where  $u = x$ ,  $v = F(x, y)$ . Thus  $\partial_u \tilde{G}(u, v) \Big|_{\substack{u=x \\ v=F(x,y)}} = \frac{c'}{\partial_y F(x, y)}$  for  $c' = -J(F, G) \in \mathbb{F}^*$ . This and (3.11) prove the following.

**Lemma 3.1**  $b_i := b_i(x) \in \mathbb{F}$  if  $i < n+m-1$ , and  $\frac{d}{dx} b_i(x) = c \bar{f}_{i-m-n+1}(x)$  if  $i \geq n+m-1$ , for  $c = -\frac{J(F, G)}{m} \in \mathbb{F}^*$ . In particular, by (3.15),

$$1 \leq \deg b_i(x) \leq 1 + (i - (m + n - 1))p \quad \text{if } i \geq m + n - 1. \quad (3.16)$$

Note that at least there exists  $i$  with  $0 \leq i < n$  and  $m \nmid (n-i)$  such that  $b_i \neq 0$ . Otherwise, by (3.9),

$$G(x, y) - \sum_{i=0}^n b_i F(x, y)^{\frac{n-i}{m}} = \sum_{i=n+1}^{\infty} b_i(x) F(x, y)^{\frac{n-i}{m}} \in \mathbb{F}[x, y] \cap y^{-1} \mathbb{F}[x, y^{-1}] = \{0\}, \quad (3.17)$$

which gives  $b_i(x) = 0$  for  $i > n$ , a contradiction with (3.16). Now we consider two cases.

*Case 1:  $m|n$ .*

Replace  $G(x, y)$  by  $G_1(x, y) = G(x, y) - F(x, y)^{\frac{n}{m}}$ , then  $n_1 := \deg_y \tilde{G}(x, y) < n$  and  $\text{Coeff}(G(x, y), y^{n_1}) = b_{n_1} \in \mathbb{F}^*$ . Rescaling  $G_1(x, y)$  to make  $b_{n_1} = 1$ , we obtain Theorem 1.1(1) by the inductive assumption on the pair  $(m, n)$ .

*Case 2:  $m \nmid n$ .*

For  $0 \leq i \leq n$ , if  $m \nmid (n-i)$ , by replacing  $G(x, y)$  by the polynomial  $G(x, y) - b_i F(x, y)^{\frac{n-i}{m}}$ , we can suppose  $b_i = 0$ .

Let  $d = (m, n)$ . We can write

$$m = dm_1, \quad n = dn_1, \quad \text{where } 2 \leq m_1 \leq n_1, \text{ and } m_1, n_1 \text{ are coprime integers.} \quad (3.18)$$

Write  $F(x, y) = F_{\text{lead}}(x, y) + F_{\text{igno}}(x, y)$  as in (2.18) and (2.19) such that  $F_{\text{lead}}(x, y) \neq y^m$  is the leading polynomial of  $F(x, y)$ . We have

**Lemma 3.2** (cf. Remark 3.13)  $F_{\text{lead}}(x, y) = \tilde{F}_1(x, y)^{m_1}$  for some polynomial

$$\tilde{F}_1(x, y) = y^d + \bar{c}_1 x^p y^{d-1} + \bar{c}_2 x^{2p} y^{d-2} + \cdots + \bar{c}_d x^{dp} \quad \text{for some } \bar{c}_i \in \mathbb{F}. \quad (3.19)$$

*Proof.* For  $i \geq 0$ , we expand  $F(x, y)^{\frac{n-i}{m}}$  as a power series of  $y^{-1}$  (cf. (2.15)),

$$F(x, y)^{\frac{n-i}{m}} = y^{n-i} + \sum_{k=1}^{\infty} f_{n-i, m, k}(x) y^{n-i-k} \quad \text{for some } f_{n-i, m, k}(x) \in \mathbb{F}[x]. \quad (3.20)$$

Thus we can use the right-hand side of (3.9) to express  $G(x, y)$  as a power series of  $y^{-1}$ ,

$$G(x, y) = y^n + \sum_{j=1}^{\infty} \tilde{g}_j(x) y^{n-j}, \quad \text{where } \tilde{g}_j(x) = \sum_{i=0}^{\infty} b_i(x) f_{n-i, m, j-i}(x). \quad (3.21)$$

Let us compute the coefficient  $\tilde{c}_j = \text{Coeff}(\tilde{g}_j(x), x^{jp})$  of  $x^{jp}$  in  $\tilde{g}_j(x)$  for  $j > 0$  (note that  $p(F^{\frac{n}{m}}) = p$ , the following arguments will also show that  $\deg \tilde{g}_j(x) \leq jp$ ). By (2.15) and (2.16),  $\deg f_{n-i,m,j-i}(x) \leq (j-i)p$ . Thus if  $1 \leq i < m+n-1$ , then  $b_i F(x, y)^{\frac{n-i}{m}}$  does not contribute to the computation of  $\tilde{c}_j$ . If  $i \geq m+n-1$ , by (3.16), noting that  $n > 2$  (by (3.8) and  $m \nmid n$ ) and the fact that  $p \geq \frac{1}{m}$  (cf. (3.7)), we have

$$\deg b_i(x) + \deg f_{n-i,m,j-i}(x) \leq 1 + (i - (m+n-1))p + (j-i)p < jp. \quad (3.22)$$

Thus  $b_i(x)F(x, y)^{\frac{n-i}{m}}$  does not contribute to  $\tilde{c}_j$  either. Similarly the ignored polynomial  $F_{\text{igno}}(x, y)$  does not contribute to  $\tilde{c}_j$  (cf. (2.19)). Therefore only  $F_{\text{lead}}(x, y)^{\frac{n}{m}}$  contributes to  $\tilde{c}_j$ , and in fact the above arguments prove

$$F_{\text{lead}}(x, y)^{\frac{n}{m}} = y^n + \sum_{i=1}^{\infty} \tilde{c}_j x^{pj} y^{n-j}. \quad (3.23)$$

Since  $G(x, y)$  is a polynomial, we must have  $\tilde{c}_j = 0$  if  $j > n$ . Then (3.23) shows that  $F_{\text{lead}}(x, y)^{\frac{n}{m}}$  is a polynomial. By Lemma 2.1, we have Lemma 3.2.  $\square$

Let  $m_2 \geq m_1$  be the largest divisor of  $m$  such that  $F_{\text{lead}}(x, y) = F_{\text{prim}}(x, y)^{m_2}$  for some

$$F_{\text{prim}}(x, y) = y^{d_2} + \sum_{i=1}^{d_2} c_i x^{ip} y^{d_2-i} \in \mathbb{F}[x, y] \text{ for some } c_i \in \mathbb{F} \text{ with some } c_i \neq 0, \quad (3.24)$$

where  $d_2 = \frac{m}{m_2}$ . If  $m_2 = m$ , then (3.24) shows that  $F_{\text{lead}}(x, y) = (y + c_1 x^p)^m$  for some  $c_1 \neq 0$ , and so  $f_1(x) = mc_1 x^p + (\text{lower terms}) \neq 0$ , a contradiction with (3.6). Thus

$$d_2 \geq 2. \quad (3.25)$$

If  $F_{\text{lead}}(x, y) = F_1(x, y)^{k_1} = F_2(x, y)^{k_2}$  for some  $F_i(x, y) \in \mathbb{F}[x, y]$  and  $k_i \in \mathbb{N}$ , then by Lemma 2.1,  $F_{\text{lead}}(x, y) = F_3(x, y)^{[k_1, k_2]}$  for some  $F_3(x, y) \in \mathbb{F}[x, y]$ . This shows that  $m_1 | m_2$  and thus  $d_2 | d$ .

**Definition 3.3** We shall call  $F_{\text{prim}}(x, y)$  in (3.24) the *primary polynomial* of  $F(x, y)$ .

The proof of Lemma 3.2 in fact shows that  $F_{\text{prim}}(x, y)$  is also the primary polynomial of  $G(x, y)$ , and  $F_{\text{prim}}(x, y)^{\frac{n}{d_2}} = F_{\text{lead}}(x, y)^{\frac{n}{m}}$  is the leading polynomial of  $G(x, y)$ .

**Lemma 3.4** (cf. Remark 3.13) *Let*

$$d_3 = (m, n - i \mid 0 \leq i \leq n, b_i \neq 0), \quad (3.26)$$

*be the g.c.d. of  $\{m, n - i \mid 0 \leq i \leq n, b_i \neq 0\}$ . There exists a polynomial of the form*

$$Q(x, y) = q_0(x)F(x, y)^{\frac{km+d_3}{m}} + \sum_{i=1}^{\infty} q_i(x)F(x, y)^{\frac{km+d_3-i}{m}}, \quad (3.27)$$

for some  $k \in \mathbb{N}$  (cf. (2.9)) and some  $q_i(x) \in \mathbb{F}[x]$  such that  $q_0(x) = 1$  and

$$\begin{aligned} q_i(x) &\in \mathbb{F} \text{ if } i < m + d_3 - 1, \text{ and} \\ \deg q_i(x) &\leq 1 + (i - (m + d_3 - 1))p \text{ if } i \geq m + d_3 - 1. \end{aligned} \quad (3.28)$$

*Proof.* We shall follow the arguments in the proof of Lemma 2.1 by regarding  $F(x, y), G(x, y)$  as  $H(x, y), P(x, y)$ . First note that  $G(x, y)$  has the form (3.9) with  $b_i(x)$  satisfying Lemma 3.1. Thus when we express  $G(x, y)^m$  (cf. (2.5)) as  $G(x, y)^m = \sum_{i=0}^{\infty} g_i^{(m)}(x)F(x, y)^{\frac{mn-i}{m}}$  for some polynomials  $g_i^{(m)}(x) \in \mathbb{F}[x]$ , we have

$$\begin{aligned} g_i^{(m)}(x) &\in \mathbb{F} \text{ if } i < m + n - 1, \text{ and} \\ \deg g_i^{(m)}(x) &\leq 1 + (i - (m + n - 1))p \text{ if } i \geq m + n - 1. \end{aligned} \quad (3.29)$$

Then we take  $\tilde{G}(x, y) = G(x, y)^m - F(x, y)^n$  and rewrite  $\tilde{G}(x, y) = \sum_{i=0}^{\infty} \tilde{g}_i(x)F(x, y)^{\frac{mn-\tilde{n}-i}{m}}$  as in (2.7) for some  $\tilde{n} \in \mathbb{N}$  and some  $\tilde{g}_i(x) \in \mathbb{F}[x]$  with  $\tilde{g}_0(x) \neq 0$ . Then (3.29) becomes

$$\begin{aligned} \tilde{g}_i(x) &\in \mathbb{F} \text{ if } i < m + n - \tilde{n} - 1 \text{ and} \\ \deg \tilde{g}_i(x) &\leq 1 + (i - (m + n - \tilde{n} - 1))p \text{ if } i \geq m + n - \tilde{n} - 1. \end{aligned}$$

Hence as in the proof of Claim 1 in Lemma 2.1, we can find some polynomial  $G_r(x, y) = \sum_{i=0}^{\infty} g_{r,i}(x)F(x, y)^{\frac{krm-j'_r-i}{m}}$ , here we define  $d_j(G) = (m, n-i \mid b_i(x) \neq 0, 0 \leq i \leq j)$  (cf. (2.2)) and define  $j_0, j_1, \dots, j_s$  as in (2.3), and  $j'_r \leq j_r$  for  $r > 0$ , such that  $g_{r,i}(x) \in \mathbb{F}[x]$  and

$$\begin{aligned} g_{r,i}(x) &\in \mathbb{F} \text{ if } i < m + n - j'_r - 1, \text{ and} \\ \deg g_{r,i}(x) &\leq 1 + (i - (m + n - j'_r - 1))p \text{ if } i \geq m + n - j'_r - 1. \end{aligned} \quad (3.30)$$

Finally, we take  $Q(x, y)$  as in (2.9). Then  $Q(x, y)$  has the form (3.27), and by (3.30),

$$\begin{aligned} q_i(x) &\in \mathbb{F} \text{ if } i < m + n - j'_s - 1, \text{ and} \\ \deg q_i(x) &\leq 1 + (i - (m + n - j'_s - 1))p \text{ if } i \geq m + n - j'_s - 1. \end{aligned} \quad (3.31)$$

Since  $d_3 \leq n - j'_s$ , (3.31) implies (3.28).  $\square$

Using (3.27) and (3.28), following the same arguments in the proof of Lemma 3.2, we have  $F_{\text{lead}}(x, y) = \tilde{F}_2(x, y)^{\frac{m}{d_3}}$  for some  $\tilde{F}_2(x, y) \in \mathbb{F}[x, y]$ . Thus  $\frac{m}{d_3} \mid m_2 = \frac{m}{d_2}$ . This proves

**Lemma 3.5** (cf. Remark 3.13)  $d_2 \mid d_3$ .

Denote

$$n_2 = \frac{n}{d_2}. \quad (3.32)$$

By Lemma 3.5 and (3.26), (3.9) becomes (cf. Remark 3.13)

$$G(x, y) = F(x, y)^{\frac{n_2}{m_2}} + \sum_{i=1}^{n_2} b_{d_2 i} F(x, y)^{\frac{n_2-i}{m_2}} + \sum_{i=n+1}^{\infty} b_i(x) F(x, y)^{\frac{n-i}{m}}. \quad (3.33)$$

Note that as in (3.9), we can also expand  $F(x, y)$  as

$$F(x, y) = F_{\text{prim}}(x, y)^{\frac{m}{d_2}} + \sum_{i=1}^{\infty} h_i(x) F_{\text{prim}}(x, y)^{\frac{m-i}{d_2}}, \quad (3.34)$$

for some  $h_i(x) \in \mathbb{F}[x]$ , which can be precisely determined by (cf. (3.10))

$$\begin{aligned} h_i(x) &= f_i(x) - \sum_{j=0}^{i-1} h_j(x) \sum_{\substack{r_1+2r_2+\dots+d_2 r_{d_2}=i-j \\ r_1, r_2, \dots, r_{d_2} \geq 0}} \binom{\frac{m-j}{d_2}}{r_1, r_2, \dots, r_{d_2}} (c_1 x^p)^{r_1} \dots (c_{d_2} x^{d_2 p})^{r_{d_2}} \\ &= f_i(x) - \sum_{j=0}^{i-1} h_j(x) \text{Coeff}(F_{\text{prim}}(x, y)^{\frac{m-j}{d_2}}, y^{m-i}), \end{aligned} \quad (3.35)$$

where  $h_0(x) = 1$  and  $f_i(x) = 0$  if  $i > m$ . Note that

$$\text{Coeff}(F_{\text{prim}}(x, y)^{\frac{m-j}{d_2}}, y^{m-i}) \text{ is a homogeneous polynomial of } x \text{ with degree } (i-j)p. \quad (3.36)$$

Thus one can prove

$$\deg h_i(x) < ip, \quad \text{and } h_j(x) \neq 0 \text{ for some } j > 0 \text{ (cf. Lemma 2.6(2)).} \quad (3.37)$$

*Subcase 2.1:*  $h_i(x) = 0$  for all  $1 \leq i \leq m$  with  $d_i \nmid (m-i)$  (cf. Remark 3.13).

Then as in (3.17), we deduce  $h_i(x) = 0$  for  $i > m$ . Thus

$$F(x, y) = F_{\text{prim}}(x, y)^{m_2} + \sum_{i=1}^{m_2} h_{d_2 i}(x) F_{\text{prim}}(x, y)^{m_2-i} = \hat{F}(x, F_{\text{prim}}(x, y)), \quad (3.38)$$

is in fact a polynomial on  $x$  and  $F_{\text{prim}}(x, y)$  for  $\hat{F}(u, v) = v^{m_2} + \sum_{i=1}^{m_2} h_{d_2 i}(u) v^{m_2-i} \in \mathbb{F}[u, v]$ . Using (3.38) in (3.33), by expanding  $G(x, y)$  as a c.r.p. of  $F_{\text{prim}}$ , we see that there exists a polynomial  $\hat{G}(u, v) \in \mathbb{F}[u, v]$  such that  $G(x, y) = \hat{G}(x, F_{\text{prim}}(x, y)) +$  (terms of negative powers of  $y$ ). Since  $G(x, y)$  is a polynomial, we have  $G(x, y) = \hat{G}(x, F_{\text{prim}}(x, y))$ . Then (1.2) gives

$$\begin{aligned} J(F, G) &= \begin{vmatrix} \partial_x F(x, y) & \partial_v \hat{F}(u, v) \partial_y F_{\text{prim}}(x, y) \\ \partial_x G(x, y) & \partial_v \hat{G}(u, v) \partial_y F_{\text{prim}}(x, y) \end{vmatrix} \\ &= \partial_y F_{\text{prim}}(x, y) \begin{vmatrix} \partial_x F(x, y) & \partial_v \hat{F}(u, v) \\ \partial_x G(x, y) & \partial_v \hat{G}(u, v) \end{vmatrix} \in \mathbb{F}^*, \end{aligned}$$

where  $u = x$ ,  $v = F_{\text{prim}}(x, y)$ . In particular,  $\partial_y F_{\text{prim}}(x, y) \in \mathbb{F}^*$ . This contradicts the fact that  $\deg_y F_{\text{prim}}(x, y) = d_2 \geq 2$  (cf. (3.25)). Thus this subcase does not occur.

*Subcase 2.2:*  $h_i(x) \neq 0$  for some  $1 \leq i \leq m$  with  $d_i \nmid (m - i)$  (the proof below in fact covers Subcase 2.1, cf. Remark 3.13).

First we need to introduce some new notations. Let  $K(x, y) \in \mathbb{F}[x]((y^{-1}))$  be any element with the form

$$K(x, y) = \sum_{i=0}^{\infty} k_i(x) F_{\text{prim}}(x, y)^{\frac{a_K - i}{d_2}} \text{ for some } k_i(x) \in \mathbb{F}[x] \text{ with } k_0(x) \neq 0 \text{ and } a_K \in \mathbb{Z}. \quad (3.39)$$

We always denote

$$A_K = \{r \in \mathbb{Q} \mid \exists i \in \mathbb{Z}_+ \text{ such that } ip - r \in \mathbb{Z}_+ \text{ and } \text{Coeff}(k_i(x), x^{ip-r}) \neq 0\} \subset \mathbb{Q}, \quad (3.40)$$

and denote

$$k_i^{[r]}(x) = \text{the homogeneous term of degree } ip - r \text{ of } k_i(x), \quad (3.41)$$

for  $r \in A_K$  (we take  $k_i^{[r]}(x) = 0$  if  $ip - r \notin \mathbb{Z}_+$ ). Denote

$$K(x, y)^{[r]} = \sum_{i=0}^{\infty} k_i^{[r]}(x) F_{\text{prim}}(x, y)^{\frac{a_K - i}{d_2}} \text{ for } r \in A_K. \quad (3.42)$$

**Definition 3.6** We call  $K(x, y)^{[r]}$  the  $r$ -th component of  $K(x, y)$ . Set  $K(x, y)^{[r]} = 0$  if  $r \notin A_K$ .

If we write  $K(x, y)$  in (3.39) as

$$K(x, y) = \sum_{i=0}^{\infty} k'_i(x) y^{a_K - i} \text{ for some } k'_i(x) \in \mathbb{F}[x], \quad (3.43)$$

then the relation between  $k_i(x)$  and  $k'_i(x)$  can be determined by (3.35) with  $m$ ,  $f_i(x)$ ,  $h_i(x)$  replaced by  $a_K$ ,  $k'_i(x)$ ,  $k_i(x)$  respectively. This and (3.36) show (cf. also (3.47))

$$K(x, y)^{[r]} = \sum_{i=0}^{\infty} k_i'^{[r]}(x) y^{a_K - i} \text{ for } r \in A_K, \quad (3.44)$$

where  $k_i'^{[r]}(x)$  is defined as in (3.41). In particular,

$$K(x, y)^{[r]} \text{ is always a polynomial if } K(x, y) \text{ is a polynomial.} \quad (3.45)$$

**Lemma 3.7** (1) Let  $K(x, y), L(x, y)$  be elements of  $\mathbb{F}[x]((y^{-1}))$  with the form of (3.39).

For any  $r \geq 0$ , the  $r$ -th component of  $K(x, y)L(x, y)$  is

$$(K(x, y)L(x, y))^{[r]} = \sum_{\substack{r_1 \in A_K, r_2 \in A_L \\ r_1 + r_2 = r}} K(x, y)^{[r_1]} L(x, y)^{[r_2]} \text{ for } r \in A_{KL}. \quad (3.46)$$

(2) For any  $\ell \in \mathbb{Z}$  with  $d_2|\ell$  and  $r \in A_{F^{\frac{\ell}{m}}}$ ,  $(F(x, y)^{\frac{\ell}{m}})^{[r]}$  is a rational function.

*Proof.* (1) Clearly, we have  $K(x, y)L(x, y) = \sum_{i=0}^{\infty} \chi_i(x) F_{\text{prim}}(x, y)^{\frac{(a_K+a_L)-i}{d_2}}$  with  $\chi_i(x) = \sum_{s_1+s_2=i} k_{s_1}(x)\ell_{s_2}(x)$ . Thus  $\chi_i^{[r]}(x) = \sum_{r_1+r_2=r} \sum_{s_1+s_2=i} k_{s_1}^{[r_1]}(x)\ell_{s_2}^{[r_2]}(x)$ . Hence we can obtain (3.46).

(2) Using (3.35) and (3.36), we have

$$h_i^{[r]}(x) = f_i^{[r]}(x) - \sum_{j=0}^{i-1} h_j^{[r]}(x) \text{Coeff}(F_{\text{prim}}(x, y)^{\frac{m-j}{d_2}}, y^{m-i}). \quad (3.47)$$

Thus the  $r$ -th component of  $F(x, y)$  (with  $F(x, y)$  being the form (3.34)) is in fact the polynomial

$$F(x, y)^{[r]} := \sum_{i=0}^{\infty} h_i^{[r]}(x) F_{\text{prim}}(x, y)^{\frac{m-i}{d_2}} = \sum_{i=0}^m f_i^{[r]}(x) y^{m-i} \text{ for all } r \in A_F.$$

Note from (3.34) and (3.37) that  $F(x, y)^{[0]} = F_{\text{prim}}(x, y)^{\frac{m}{d_2}}$ . We have

$$\begin{aligned} F(x, y)^{\frac{\ell}{m}} &= \left( \sum_{r \in A_F} F(x, y)^{[r]} \right)^{\frac{\ell}{m}} = F_{\text{prim}}(x, y)^{\frac{\ell}{d_2}} \left( 1 + \sum_{r \in A_F \setminus \{0\}} F(x, y)^{[r]} F_{\text{prim}}(x, y)^{-\frac{m}{d_2}} \right)^{\frac{\ell}{m}} \\ &= \sum_{r \in A_{F^{\frac{\ell}{m}}}} \sum_{\substack{s_1 t_1 + \dots + s_k t_k = r \\ 0 < s_1 < \dots < s_k \\ s_i \in A_F, t_i \in \mathbb{Z}_+, k \geq 0}} \binom{\frac{\ell}{m}}{t_1, \dots, t_k} \prod_{i=1}^k (F(x, y)^{[s_i]})^{t_i} F_{\text{prim}}(x, y)^{\frac{\ell - (t_1 + \dots + t_k)m}{d_2}}. \end{aligned}$$

By (1), the  $j$ -th component of  $(F(x, y)^{[s_i]})^{t_i}$  (which is the  $t_i$ -th power of the  $s_i$ -th component of  $F(x, y)$ ) is zero if  $j \neq s_i t_i$  for all  $i$ . Thus again by (1), the  $r$ -th component of  $F(x, y)^{\frac{\ell}{m}}$  is

$$(F(x, y)^{\frac{\ell}{m}})^{[r]} = \sum_{\substack{s_1 t_1 + \dots + s_k t_k = r \\ 0 < s_1 < \dots < s_k \\ s_i \in A_F, t_i \in \mathbb{Z}_+, k \geq 0}} \binom{\frac{\ell}{m}}{t_1, \dots, t_k} \prod_{i=1}^k (F(x, y)^{[s_i]})^{t_i} F_{\text{prim}}(x, y)^{\frac{\ell - (t_1 + \dots + t_k)m}{d_2}}, \quad (3.48)$$

which is a finite sum of rational functions (for any given  $r$ ) by noting that the powers of  $F_{\text{prim}}(x, y)$  in (3.48) are integers and  $A_F \subset \mathbb{Q}$  is a finite set since  $f_i(x) = 0$  if  $i > m$ .  $\square$

Note that  $G(x, y)$  has the form (3.9). We shall compute  $G(x, y)^{[r]}$  for some suitable chosen  $r$ . First we collect some basic facts:

**Fact (i)**  $G(x, y)^{[r]}$  is a polynomial since  $G(x, y)$  is a polynomial (cf. (3.45)).

**Fact (ii)** If  $0 \leq i < m + n - 1$ , then  $b_i(F(x, y)^{\frac{n-i}{m}})^{[r]}$  (which is a rational function if  $d_2|i$  by Lemma 3.7(2)) contributes to the  $(ip + r)$ -th component  $G(x, y)^{[ip+r]}$  of  $G(x, y)$ .

**Fact (iii)** Suppose  $i \geq m + n - 1$ . Denote  $q_i = (i - (m + n - 1))p$ , and  $b_{i,0} = \text{Coeff}(b_i(x), x^{1+q_i})$  (recall (3.16)). Then by (3.14) and Lemma 3.1,

$$(1 + q_i)b_{i,0} = c(1 - i')\bar{b}_{i',0}, \quad (3.49)$$

where  $i' = i - (m + n - 1)$ ,  $\bar{b}_{i',0} = \text{Coeff}(\bar{b}_{i'}(x), x^{i'p})$ , and  $c = -\frac{J(F,G)}{m} \in \mathbb{F}^*$ . Thus,

$$G_1(x, y) := \sum_{i=m+n-1}^{\infty} b_{i,0} x^{1+qi} F_{\text{prim}}(x, y)^{\frac{n-i}{d_2}} = c \sum_{i=0}^{\infty} \frac{1-i}{1+ip} \bar{b}_{i,0} x^{1+ip} F_{\text{prim}}(x, y)^{\frac{1-m-i}{d_2}}. \quad (3.50)$$

**Fact (iv)** Note that (3.50) contributes to  $G(x, y)^{[r]}$  for  $r = (m + n - 1)p - 1$ .

**Fact (v)** By computing the 0-th component of (3.12), one has

$$G_2(x, y) := \sum_{i=0}^{\infty} \bar{b}_{i,0} x^{ip} F_{\text{prim}}(x, y)^{\frac{1-i}{d_2}} = y. \quad (3.51)$$

**Lemma 3.8**  $G_1(x, y)$  defined in (3.50) satisfies the following differential equation, for  $c'_1 = -p^{-1}c \in \mathbb{F}^*$ ,

$$(m_2 - \frac{p+1}{d_2p}) F_{\text{prim}}(x, y)^{m_2-1} G_1(x, y) \partial_y F_{\text{prim}}(x, y) + F_{\text{prim}}(x, y)^{m_2} \partial_y G_1(x, y) = c'_1 x. \quad (3.52)$$

*Proof.* Denote  $k_1(z) = \sum_{i=0}^{\infty} \frac{1-i}{1+ip} \bar{b}_{i,0} z^{-ip-1}$ ,  $k_2(z) = \sum_{i=0}^{\infty} \bar{b}_{i,0} z^{p(1-i)}$ . We have

$$-p \frac{d}{dz} k_1(z) = z^{-p-1} \frac{d}{dz} k_2(z). \quad (3.53)$$

Fix  $x$  and let  $z = x^{-1} F_{\text{prim}}(x, y)^{\frac{1}{d_2p}}$ , using (3.50) and (3.51), we have

$$k_1(z) = c^{-1} F_{\text{prim}}(x, y)^{\frac{mp-p-1}{d_2p}} G_1(x, y) \quad \text{and} \quad k_2(z) = x^{-p} G_2(x, y) = x^{-p} y.$$

Using this in (3.53), we obtain

$$\begin{aligned} & \left( (m_2 - \frac{p+1}{d_2p}) F_{\text{prim}}(x, y)^{m_2-1-\frac{p+1}{d_2p}} G_1(x, y) \partial_y F_{\text{prim}}(x, y) + F_{\text{prim}}(x, y)^{m_2-\frac{p+1}{d_2p}} \partial_y G_1(x, y) \right) \frac{dy}{dz} \\ &= c'_1 x F_{\text{prim}}(x, y)^{-\frac{p+1}{d_2p}} \frac{dy}{dz}. \end{aligned}$$

Thus we have (3.52). □

Write  $p = \frac{p'}{q'}$  for some coprime positive integers  $p'$ ,  $q'$ . Note from (3.24) that at least one of  $p, 2p, \dots, d_2p$  is an integer, thus  $1 \leq q' \leq d_2$ .

**Lemma 3.9** *If  $p' = 1$  and  $q' = d_2$ , then  $F(x, y), G(x, y)$  are generators of  $\mathbb{F}[x, y]$ .*

*Proof.* In this case,  $p = \frac{1}{d_2}$ . By (3.24),  $F_{\text{prim}}(x, y) = y^{d_2} + c_{d_2} x$  with  $c_{d_2} \neq 0$ . Rescaling  $x$  we can suppose  $c_{d_2} = 1$ . Now by replacing  $x$  by  $x - y^{d_2}$ , the leading polynomial  $F_{\text{lead}}(x, y)$  of  $F(x, y)$  becomes  $x^{m_2}$ . By definition of the prime degree  $p$ , we have

$$d_2 i < j \quad \text{if a nonzero term } c_{i,j} x^i y^{m-j} \text{ with } c_{i,j} \in \mathbb{F}, \text{ appears in } F_{\text{igno}}(x, y). \quad (3.54)$$

Thus after the replacement,  $y^m$  does not appear in  $F_{\text{igno}}(x, y)$ . Thus  $F(x, y)$  becomes a new polynomial with  $\deg_y F(x, y) < m$ . Furthermore, if we let  $m'$  be the total degree of  $x$  and  $y$  in the new  $F(x, y)$ , then (3.54) also shows that  $m' < m$ . Say  $K(x, y) = \sum_{i=0}^s c'_i x^i y^{m'-i}$  is the polynomial of terms of  $F(x, y)$  with highest total degree of  $x$  and  $y$ , then the polynomial  $f(x) := \sum_{i=0}^s c'_i x^i \neq 0$ . Take  $a \in \mathbb{F}$  with  $f(a) \neq 0$ . Then by applying linear isomorphism (3.3), we obtain that  $F(x, y)$  becomes a polynomial with  $\deg_y F(x, y) = m'$  such that  $\alpha := \text{Coeff}(F(x, y), y^{m'}) = f(a) \in \mathbb{F}^*$ . By rescaling  $F(x, y)$ , we can suppose  $\alpha = 1$ .

The similar arguments show that  $G(x, y)$  becomes a polynomial with  $n' := \deg_y G(x, y) < n$  such that by a suitable choice of  $a$  in (3.3) and by rescaling  $G(x, y)$ , the coefficient  $\text{Coeff}(G(x, y), y^{n'}) = 1$  (cf. arguments before (3.4)). Thus by the inductive assumption on the pair  $(m, n)$ , the polynomials  $F(x, y)$  and  $G(x, y)$  are generators of  $\mathbb{F}[x, y]$ .  $\square$

Denote

$$B = \{i \mid 0 \leq i < m + n - 1, b_i \neq 0, d_2 \nmid i\}. \quad (3.55)$$

Now we consider the following cases:

*Subcase 2.2.1: Suppose  $B = \emptyset$  or  $(m + n - 1 - i)p < 1$  for all  $i \in B$ , or  $B \neq \emptyset$  and  $j = \min B$  satisfies  $(m + n - 1 - j)p = 1$ .*

First we remark that the case “ $B = \emptyset$  or  $(m + n - 1 - i)p < 1$  for all  $i \in B$ ” can be regarded as a special case of the case “ $B \neq \emptyset$  and  $j = \min B$  with  $b_j = 0$  in (3.57)”. Thus it suffices to consider the case “ $B \neq \emptyset$  and  $j = \min B$ ”. (This is the most nontrivial case. As stated in [M2, M3] (cf. [M1]), we only need to consider the case  $p = 1$ . However, we need some information for the general case, cf. proof of Lemma 3.11.)

We take  $r = jp = (m + n - 1)p - 1$ . Let us compute  $G(x, y)^{[r]}$  under modulo  $\mathbb{F}(x, y)$ . In the following, we use “ $\equiv$ ” to mean: If  $U, V$  are two vector spaces such that  $U \subset V$  and  $a, b \in V$  then

$$a \equiv b \pmod{U} \iff a - b \in U. \quad (3.56)$$

For  $0 \leq i < m + n - 1$ , if  $d_2 \mid i$ , then by Fact (ii), all components of  $b_i F(x, y)^{\frac{n-i}{m}}$  are rational functions of the form (3.48), thus does not contribute to  $G(x, y)^{[r]} \pmod{\mathbb{F}(x, y)}$ . If  $j \neq i \in B$ , then  $j < i$ , so  $r < ip$ , and again by Fact (ii),  $b_i F(x, y)^{\frac{n-i}{m}}$  does not contribute to  $G(x, y)^{[r]} \pmod{\mathbb{F}(x, y)}$  either since the top most component it can contribute is  $ip$ . Thus by Facts (i) and (iv),

$$0 \equiv G(x, y)^{[r]} \equiv b_j F_{\text{prim}}(x, y)^{\frac{n-j}{d_2}} + G_1(x, y) \pmod{\mathbb{F}(x, y)}, \quad (3.57)$$

i.e.,

$$G'_1(x, y) = b_j F_{\text{prim}}(x, y)^{\frac{n-j}{d_2}} + G_1(x, y) \quad (3.58)$$

is a rational function of the form in the right-hand side of (3.48).

Denote by  $\overline{\mathbb{F}(x)}$  the algebraic closure of the field  $\mathbb{F}(x)$  of rational functions of  $x$ . From now on, we regard polynomials in  $\mathbb{F}[x, y]$  as polynomials of  $y$  in  $\overline{\mathbb{F}(x)}[y]$  with coefficients in  $\overline{\mathbb{F}(x)}$ . By (3.58) and the right-hand side of (3.48), we can write  $G_1(x, y)$  as

$$G_1(x, y) = -b_j F_{\text{prim}}(x, y)^{\frac{n-j}{d_2}} + F_{\text{prim}}(x, y)^{-a} P(x, y), \quad (3.59)$$

for some  $a \in \mathbb{Z}$  and some  $P(x, y) \in \overline{\mathbb{F}(x)}[y]$  such that  $F_{\text{prim}}(x, y) \nmid P(x, y)$  (inside the ring  $\overline{\mathbb{F}(x)}[y]$ ). Using (3.59) in (3.52), we see that the term with  $F_{\text{prim}}(x, y)^{\frac{n-j}{d_2}}$  and the term with  $\partial_y(F_{\text{prim}}(x, y)^{\frac{n-j}{d_2}})$  are precisely canceled because of the fact that  $(m+n-1-j)p=1$ , and we obtain, for  $a' = m_2 - a$ ,  $c'_2 = d_2 p' c'_1 x \in \overline{\mathbb{F}(x)}$ ,

$$(a' d_2 p' - p' - q') F_{\text{prim}}(x, y)^{a'-1} P(x, y) \partial_y F_{\text{prim}}(x, y) + d_2 p' F_{\text{prim}}(x, y)^{a'} \partial_y P(x, y) = c'_2, \quad (3.60)$$

which is equivalent to, for some  $c'_3 \in \overline{\mathbb{F}(x)}$ ,

$$F_{\text{prim}}(x, y)^{a' d_2 p' - p' - q'} P(x, y)^{d_2 p'} = c'_2 \int F_{\text{prim}}(x, y)^{a' d_2 p' - p' - q' - a'} P(x, y)^{d_2 p' - 1} dy + c'_3. \quad (3.61)$$

For convenience, a pair  $(F(x, y), G(x, y))$  is called a *Jacobian pair* if  $F(x, y), G(x, y) \in \mathbb{F}[x, y]$  are monic polynomials of  $y$  such that  $J(F, G) \in \mathbb{F}^*$ . Here we do not assume  $F(x, y)$  satisfies condition (3.6). Note that in general  $m := \deg_y F$  is not necessarily equal to  $\deg F$  (we remark that  $\deg F$  always means the total degree of  $F(x, y)$ ); for instance, in case  $F(x, y) = y + x^3$ , we have  $m = 1$  and  $\deg F = 3$ .

**Lemma 3.10** *Let  $(F(x, y), G(x, y))$  be any Jacobian pair. We have one and only one of the following (up to a linear automorphism of  $\mathbb{F}[x, y]$ ):*

- (1)  $F_{\text{prim}}(x, y) = y + x^{p'}$ . In this case,  $d_2 = 1$ ,  $p = p'$ ,  $q' = 1$ . (We shall not consider this case since all our arguments above are based on the assumption that  $d_2 \geq 2$ . Anyway, if we require condition (3.6) then this case does not occur.)
- (2)  $F_{\text{prim}}(x, y) = y^{d_2} + x$ , for some  $d_2 \geq 2$ , is an irreducible polynomial in  $\mathbb{F}[x, y]$ . In this case,  $a' = 1$ ,  $p' = 1$ ,  $q' = d_2$ ,  $j = m + n - d_2 - 1$ ,  $P(x, y) = p_0 y$  and

$$b_j = p_0 = \frac{J(F, G)}{m_2}, \quad G'_1(x, y) = F_{\text{prim}}(x, y)^{-a} P(x, y) = p_0 (y^{q'} + x)^{-m_2+1} y. \quad (3.62)$$

- (3)  $F_{\text{prim}}(x, y) = (y^{q'} + x)^{i_1} y^{i_2}$  (if we require condition (3.6) and if  $q' = 1$ , then  $F_{\text{prim}}(x, y) = (y - i_2 x)^{i_1} (y + i_1 x)^{i_2}$ ) is a reducible polynomial in  $\mathbb{F}[x, y]$ , for some coprime positive integers  $i_1, i_2$  such that  $q' i_1 + i_2 = d_2$ . In this case,  $a' = 0$ ,  $a = -m_2$ ,  $p' = 1$ ,  $j = m + n - q' - 1$ ,  $P(x, y) = p_0 (y^{q'} + x) y$  and

$$b_j = p_0 = \frac{J(F, G)}{(i_1 - i_2) m_2}, \quad G'_1(x, y) = F_{\text{prim}}(x, y)^{-a} P(x, y) = p_0 (y^{q'} + x)^{-i_1 m_2 + 1} y^{-i_2 m_2 + 1}. \quad (3.63)$$

*Proof.* First note that (3.62) can be regarded as a special case of (3.63) with  $i_1 = 1$ ,  $i_2 = 0$ .

To avoid confusion on whether or not the inductive assumption on  $m$  is used, we want to remark that although the proof of Theorem 1.1 is based on induction on  $m$ , the proof of this lemma does not depend on induction on  $m$ , thus the lemma holds in general (since we shall see that Subcase 2.2.2 cannot occur, this lemma holds for all Jacobian pairs).

The proof is divided into three cases.

*Case (i):*  $a' > 1$ . Then  $F_{\text{prim}}(x, y)$  divides the left-hand side of (3.60), but does not divide the right-hand side. The contradiction shows that this case does not occur.

*Case (ii):*  $a' = 1$ . If  $d_2 = 1$ , we clearly have case (1) (up to a linear automorphism  $(x, y) \mapsto (\alpha x, y)$  for some  $\alpha \in \mathbb{F}^*$ ). Assume  $d_2 \geq 2$ . If  $d_2 p' - p' - q' - 1 \geq 0$ , then all functions in (3.61) are polynomials. By comparing the degrees of  $y$  in both sides, we see that the left-hand side has a higher degree than the right-hand side, thus we obtain a contradiction. Hence  $d_2 p' - p' - q' \leq 0$ . But  $d_2 p' - p' - q' \neq 0$  by (3.60). Thus  $d_2 p' - p' - q' \leq -1$ . So  $p' \leq \frac{q'-1}{d_2-1} \leq 1$ . This forces  $p' = 1$  and  $q' = d_2$ . Thus we can assume  $F_{\text{prim}}(x, y) = y^{d_2} + x$  (up to a linear automorphism  $(x, y) \mapsto (\alpha x, y)$  for some  $\alpha \in \mathbb{F}^*$ ). Assume  $P(x, y)$  has degree  $k$  on  $y$ . Then comparing the coefficients of  $y^{k+d_2-1}$  in (3.60) shows that  $k = 1$ . Thus we obtain  $P(x, y) = p_0 y$  for some  $p_0 \in \mathbb{F}^*$ . Noting that  $c'_2 = d_2 c'_1 x = -d_2 p^{-1} c x = d_2^2 \frac{J(F, G)}{m} x$ , we have  $p_0 = \frac{J(F, G)}{m_2}$ . We obtain

$$b_j = p_0, \quad (3.64)$$

as follows: Comparing the coefficients of  $y^{d_2+1-m}$  in both sides of (3.59) (expanding all terms as elements of  $\mathbb{F}[x]((y^{-1}))$  defined in (2.12)), the left-hand side has coefficient zero by (3.50), while the right-hand side has coefficient  $-b_j + p_0$  (noting that  $n - j = d_2 + 1 - m$ ). Thus we have case (2).

*Case (iii):*  $a'' = -a' \geq 0$ . Factorize  $F_{\text{prim}}(x, y), P(x, y) \in \overline{\mathbb{F}(x)}[y]$  as products of irreducible polynomials on  $y$ :

$$F_{\text{prim}}(x, y) = f_0 f_1^{i_1} \cdots f_\ell^{i_\ell}, \quad P(x, y) = p_0 f_1^{j_1} \cdots f_r^{j_r} p_{r+1}^{j_{r+1}} \cdots p_s^{j_s}, \quad (3.65)$$

for some  $\ell, s, i_1, \dots, i_\ell, j_1, \dots, j_s \in \mathbb{N}$  and  $0 \leq r \leq \min\{s, \ell\}$ ,  $f_0, p_0 \in \mathbb{F}^*$  (note that since  $F_{\text{prim}}(x, y), P(x, y) \in \mathbb{F}[x, y]$ , we have  $f_0, p_0 \in \mathbb{F}^*$  and in fact  $f_0 = 1$ ) and where

$$p_1 := f_1, \quad \dots, \quad p_r := f_r, \quad f_{r+1}, \quad \dots, \quad f_\ell, \quad p_{r+1}, \quad \dots, \quad p_s \in \overline{\mathbb{F}(x)}[y]$$

are different irreducible monic polynomials of  $y$  (thus, of degree 1). Multiplying (3.60) by  $F_{\text{prim}}(x, y)^{a''+1}$ , using (3.65), and canceling the common factor

$$f_1^{i_1+j_1-1} \cdots f_r^{i_r+j_r-1} f_{r+1}^{i_{r+1}-1} \cdots f_\ell^{i_\ell-1},$$

noting that  $\partial_y f_\mu = \partial_y p_\mu = 1$  for all  $\mu$ , we obtain

$$\begin{aligned} & \left( \alpha_1 p_{r+1} \cdots p_s \sum_{\mu=1}^{\ell} i_\mu \frac{f_1 \cdots f_\ell}{f_\mu} + \alpha_2 f_{r+1} \cdots f_\ell \sum_{\nu=1}^s j_\nu \frac{p_1 \cdots p_s}{p_\nu} \right) p_{r+1}^{j_{r+1}-1} \cdots p_s^{j_s-1} \\ &= \alpha_3 f_1^{i_1 a''+1-j_1} \cdots f_r^{i_r a''+1-j_r} f_{r+1}^{i_{r+1} a''+1} \cdots f_\ell^{i_\ell a''+1}, \end{aligned} \quad (3.66)$$

where

$$\alpha_1 = -(a'' d_2 p' + p' + q'), \quad \alpha_2 = d_2 p', \quad \alpha_3 = c'_2 f_0^{a''} p_0^{-1}. \quad (3.67)$$

If  $\ell > r$ , then  $f_\ell$  divides all terms except one term corresponding to  $\mu = \ell$  in (3.66), a contradiction. Thus  $\ell = r$ . Since  $p_{r+1}, \dots, p_s$  do not appear in the right-hand side of (3.66), we must have  $j_{r+1} = \dots = j_s = 1$ , and since the left-hand side is a polynomial, we have

$$i_k a'' + 1 - j_k \geq 0 \quad \text{for } k = 1, \dots, \ell. \quad (3.68)$$

If  $i_k a'' + 1 - j_k > 0$  for some  $k$ , then  $f_k$  divides all terms except two terms corresponding to  $\mu = k$  and  $\nu = k$  in (3.66), and the sum of these two terms is a term (not divided by  $f_k$ ) with coefficient  $\alpha_1 i_k + \alpha_2 j_k$ . This proves

$$i_k a'' + 1 - j_k = 0 \quad \text{or} \quad \alpha_1 i_k + \alpha_2 j_k = 0 \quad \text{for } k = 1, \dots, \ell. \quad (3.69)$$

If  $a'' > 0$ , then either case of (3.69) in particular shows that  $i_k \leq j_k$  (cf. (3.67)), thus,  $F_{\text{prim}}(x, y) | P(x, y)$ , a contradiction with our choice of  $P(x, y)$ . Thus  $a' = -a'' = 0$ . In particular  $j_k = 1$  for all  $k$  by (3.68). Then (3.66) is simplified to

$$p_{\ell+1} \cdots p_s \sum_{\mu=1}^{\ell} (\alpha_1 i_\mu + \alpha_2) \frac{f_1 \cdots f_\ell}{f_\mu} + \alpha_2 f_1 \cdots f_\ell \sum_{\nu=\ell+1}^s \frac{p_{\ell+1} \cdots p_s}{p_\nu} = \alpha_3. \quad (3.70)$$

If  $\ell = 1$ , then (3.65) shows that  $i_1 = d_2$  and  $F_{\text{prim}}(x, y) = f_1^{d_2}$ . Write  $f_1 = y + f_{11}$  for some  $f_{11} \in \overline{\mathbb{F}(x)}$ . Since  $d_2 f_{11} y^{d_2-1} = c_1 x^p y^{d_2-1}$  by (3.24), we have  $p = p' \in \mathbb{N}$  and  $f_{11} \in \mathbb{F}[x]$ . Thus  $f_1 \in \mathbb{F}[x, y]$  and  $d_2 = 1$  (since  $F_{\text{prim}}(x, y) \neq Q(x, y)^k$  for any  $Q(x, y) \in \mathbb{F}[x, y]$  and  $2 \leq k \in \mathbb{Z}$ ). However the fact  $d_2 = 1$  forces  $F_{\text{prim}}(x, y) | P(x, y)$ . Thus this case cannot occur.

Now suppose  $\ell > 1$ . In (3.70), computing the coefficient of the term with highest degree (i.e., degree  $s-1$ ) shows

$$-(p' + q') d_2 + s p' d_2 = \alpha_1 d_2 + s \alpha_2 = \sum_{\mu=1}^{\ell} (\alpha_1 i_\mu + \alpha_2) + \sum_{\nu=\ell+1}^s \alpha_2 = 0,$$

i.e.,  $s p' = p' + q'$ . Since  $(p', q') = 1$ , we obtain

$$p' = 1, \quad q' = s - 1, \quad j = m + n - q' - 1. \quad (3.71)$$

First suppose  $s > 2$ . Let  $H(x, y) \in \mathbb{F}[x, y]$ , a monic polynomial of  $y$ , be an irreducible factor of  $F_{\text{prim}}(x, y)$  (in the ring  $\mathbb{F}[x, y]$ ). Since  $y^{-d_2}F_{\text{prim}}(x, y)$  is in fact a polynomial of  $xy^{-q'}$ , if  $\deg_y H = k$  then

$$H(x, y) \text{ must have the form } \sum_{i \in \mathbb{Z}_+, i \leq \frac{k}{q'}} h_i x^i y^{k-q'i} \text{ for some } h_i \in \mathbb{F}, \quad (3.72)$$

i.e.,  $H(x, y)$  must be a quasi-homogenous polynomial of prime degree  $p$  by Definition 2.5. (This can also be proved as follows:  $F_{\text{prim}}(x, y)$  is a quasi-homogenous polynomial of prime degree  $p$ , and every quasi-homogenous polynomial of prime degree  $p$  must have only one nonzero component, then one can use (3.46) to prove that every factor of a quasi-homogenous polynomial of prime degree  $p$  must be a quasi-homogenous polynomial of prime degree  $p$ .) Thus (3.72) shows either

$$k = 1 \text{ (and } H(x, y) = y), \text{ or}$$

$$q'|k \text{ (in this case } H(x, y) \text{ has } k \text{ different irreducible factors in } \overline{\mathbb{F}(x)}[y]).$$

Since  $F_{\text{prim}}(x, y)$  has only  $\ell$  different irreducible factors in  $\overline{\mathbb{F}(x)}[y]$  and  $\ell \leq s = q' + 1$ , we see that  $F_{\text{prim}}(x, y)$  has to have the form (up to a linear automorphism  $(x, y) \mapsto (\alpha x, y)$  for some  $\alpha \in \mathbb{F}^*$ )

$$F_{\text{prim}}(x, y) = (y^{q'} + x)^{i_1} y^{i_2}, \quad P(x, y) = p_0(y^{q'} + x)y, \quad q'i_1 + i_2 = d_2, \quad i_1, i_2 \in \mathbb{N}, \quad (3.73)$$

such that  $i_1, i_2$  are coprime (cf. the fourth statement after (3.70)), where the second equation of (3.73) follows since each irreducible factor (in  $\overline{\mathbb{F}(x)}[y]$ ) of  $F_{\text{prim}}(x, y)$  must appear in  $P(x, y)$  and  $s = q' + 1$ . If  $i_1 = 0$  (then  $i_2 = 1$ ) or  $i_2 = 0$  (then  $i_1 = 1$ ), then  $F_{\text{prim}}(x, y) | P(x, y)$ , a contradiction. Thus  $i_1 \neq 0, i_2 \neq 0$ . Thus  $F_{\text{prim}}(x, y)$  has  $q' + 1$  different irreducible factors in  $\overline{\mathbb{F}(x)}[y]$ , namely  $\ell = q' + 1 = s$ .

If  $s = 2$ , then  $\ell = 2$  since  $2 \leq \ell \leq s$ . We can write  $F_{\text{prim}}(x, y) = (y + \beta_1)^{i_1} (y + \beta_2)^{i_2}$ , for some  $\beta_1, \beta_2 \in \overline{\mathbb{F}(x)}$  and  $i_1 + i_2 = d_2$ . We have  $i_1 \beta_1 + i_2 \beta_2 = 0$  from (3.6), and  $(\alpha_1 i_1 + \alpha_2) \beta_2 + (\alpha_1 i_2 + \alpha_2) \beta_1 = \alpha_3 \in \mathbb{F} \cdot x$  from (3.70). Thus  $\beta_1 = \beta'_1 x, \beta_2 = \beta'_2 x$  are different scalar multiples of  $x$ . Thus by applying the linear automorphism  $(x, y) \mapsto (\frac{1}{\beta_1 - \beta_2} x, y - \beta'_2 x)$ ,  $F_{\text{prim}}(x, y)$  still have the form in (3.73). Thus we can suppose (3.73) holds in general.

Using (3.73) in (3.60) or (3.52), we obtain the second equality of the first equation of (3.63) by noting that  $c'_2 = d_2 c'_1 x = -d_2 p^{-1} c x = d_2 q' \frac{J(F, G)}{m} x$ . The first equality follows as in (3.64). Thus we have case (3). This proves the lemma.  $\square$

The arguments in the proof of the following key lemma in fact will also prove Theorem 1.2. To convince readers that our approach works, we shall give two proofs in the final part of the proof of Lemma 3.11.

**Lemma 3.11 (Key Lemma)** *For any Jacobian pair  $(F(x, y), G(x, y))$ , the polynomials  $F(x, y), G(x, y)$  are generators of  $\mathbb{F}[x, y]$ .*

*Proof.* Suppose conversely that  $(F(x, y), G(x, y))$  is a Jacobian pair with minimal  $m = \deg_y F$  such that  $F(x, y), G(x, y)$  are not generators. (We claim that the proof in fact does not need to use the minimal assumption of  $m$  as long as the prime degree of  $F(x, y)$  is 1. See arguments after (3.122)).

Then we have  $m < n := \deg_y G$ . We can suppose Lemma 3.10(1) does not occur. This is because, if necessary, by applying some automorphism, we can add condition (3.6). Then the minimal choice of  $m$  and the proof of Lemma 3.9 show that Lemma 3.10(2) cannot occur. Thus we have Lemma 3.10(3). Since  $x = (y^{q'} + x) - y^{q'}$  and a term  $x^i y^j$  appears in  $F(x, y)$  with nonzero coefficient must satisfy  $q'i + j \leq m$ , we can write  $F(x, y)$  as

$$F(x, y) = \sum_{(i,j) \in S} f_{ij}(y^{q'} + x)^i y^j \text{ for some } f_{ij} \in \mathbb{F}, \text{ and} \quad (3.74)$$

$$S = \{(i, j) \mid i, j \in \mathbb{Z}_+, q'i + j \leq m, f_{ij} \neq 0\}. \quad (3.75)$$

Then

$$q'i + j = m \iff (i, j) = (i_1 m_2, i_2 m_2), \quad (3.76)$$

which corresponds to the leading polynomial  $F_{\text{lead}}(x, y) = (y^{q'} + x)^{i_1 m_2} y^{i_2 m_2}$  of  $F(x, y)$ . By (3.46), we have  $F(x, y)^{[r]} = 0$  if  $r \notin \frac{1}{q'}\mathbb{Z}$ , and

$$F(x, y)^{\lfloor \frac{r}{q'} \rfloor} = \sum_{q'i+j=m-r} f_{ij}(y^{q'} + x)^i y^j \text{ if } r \in \mathbb{Z}_+. \quad (3.77)$$

Denote  $I_n = \{(i, j) \in \mathbb{Z}^2 \mid 0 \leq i, j \leq n\}$ . Then  $S \subset I_n$  (the reason we use  $I_n$  instead of  $I_m$  is that we also need to consider  $G(x, y)$ ). We define the lexicographical order on  $\mathbb{Z}^2$ , and define

$$(i_0, j_0) = \max S, \quad (3.78)$$

i.e.,  $i_0 = \max\{i \mid f_{ij} \neq 0 \text{ for some } j\}$ ,  $j_0 = \max\{j \mid f_{i_0, j} \neq 0\}$ . We can suppose there exists some

$$(i, j) \in S \text{ with } i + j \geq 3. \quad (3.79)$$

In fact  $(i_1 m_2, i_2 m_2) \in S$  satisfies  $i_1 m_2 + i_2 m_2 \geq 4$  since  $m_2 \geq 2$ . If necessary, by applying the automorphism  $(y^{q'} + x, y) \mapsto (y, y^{q'} + x)$ , i.e.,  $(x, y) \mapsto (y - (y^{q'} + x)^{q'}, y^{q'} + x)$ , we can suppose

$$i_0 \geq j_0. \text{ Thus } i_0 \geq 2.$$

**Claim 1**  $j_0 \neq 0$ .

Otherwise by applying the automorphism  $\sigma : (x, y) \mapsto (y - x^{q'}, x)$ , the polynomial  $F(x, y)$  becomes the monic polynomial of  $y$  (up to the nonzero scalar  $f_{i_0, j_0}$ ):

$$\check{F}(x, y) = F(y - x^{q'}, x) = \sum_{(i, j) \in S} f_{ij} y^i x^j = f_{i_0, j_0} y^{i_0} + \sum_{(i_0, j_0) \neq (i, j) \in S} f_{ij} y^i x^j,$$

with  $\deg_y \check{F} = i_0 \leq q' i_0 + j_0 < m$  by (3.76) since  $(i_0, j_0) = (i_0, 0) \neq (i_1 m_2, i_2 m_2)$ . By the minimal choice of  $m$ , the polynomials  $\check{F}(x, y)$ ,  $\check{G}(x, y) = G(y - x^{q'}, x)$  are generators, and so  $F(x, y)$ ,  $G(x, y)$  are generators of  $\mathbb{F}[x, y]$ , a contradiction (we remark that in case  $q' = 1$  the total degree  $\deg \check{F}$  is not reduced, but what we want is that  $\deg_y \check{F}$  is reduced).

Now let

$$k > n \quad (\text{and } k \gg 0 \text{ whenever necessary}), \quad (3.80)$$

be any integer. Then

$$k i_0 + j_0 > k i + j \quad \text{for all } (i_0, j_0) \neq (i, j) \in S \subset I_n. \quad (3.81)$$

Take

$$\begin{aligned} F^{(k)}(x, y) &= f_{i_0, j_0}^{-1} F(y^k - y^{q'} + x, y) = \sum_{(i, j) \in S} f'_{ij} (y^k + x)^i y^j, \\ G^{(k)}(x, y) &= \alpha G(y^k - y^{q'} + x, y), \end{aligned} \quad (3.82)$$

for  $f'_{ij} = f_{ij} f_{i_0, j_0}^{-1}$  and some  $\alpha \in \mathbb{F}^*$  (with  $\alpha$  not depending on  $k$  such that  $G(x, y)$  becomes a monic polynomial of  $y$ ). Since  $F^{(k)}(x, y)$  contains the polynomial

$$(y^k + x)^{i_0} y^{j_0} = y^{k i_0 + j_0} + i_0 x y^{k(i_0 - 1) + j_0} + \dots, \quad (3.83)$$

and by (3.81) the first two terms in the right-hand side of (3.83) do not appear in all  $(y^k + x)^i y^j$  with  $(i_0, j_0) \neq (i, j) \in S$ , we see by (3.81),

$$m^{(k)} := \deg_y F^{(k)} = k i_0 + j_0.$$

**Convention 3.12** Here and below, we use the same symbols with superscript “ $\langle k \rangle$ ” to denote notations associated with the pair  $(F^{(k)}(x, y), G^{(k)}(x, y))$  which is also a Jacobian pair.

By Definition 2.3, the prime degree of  $F^{(k)}(x, y)$  is  $p^{(k)} \geq \frac{1}{k}$  (note from (3.83) that  $i_0 x$  appear as a nonzero term of  $h_k(x)$  in Definition 2.3). Without difficulty, one can easily show that a term

$x^{i'} y^{j'}$  cannot appear in  $F^{(k)}(x, y)$  if  $k i' + j' > m^{(k)}$ , and

$x^{i'} y^{j'}$  with  $k i' + j' = m^{(k)}$  can appear in  $(y^k + x)^i y^j$  with  $(i, j) \in S$  only when  $(i, j) = (i_0, j_0)$ .

Thus we have in fact proved

**Claim 2** The prime degree of  $F^{(k)}(x, y)$  is  $p^{(k)} = \frac{1}{k}$ , and the leading polynomial of  $F^{(k)}(x, y)$  is  $F_{\text{lead}}^{(k)}(x, y) = (y^k + x)^{i_0} y^{j_0}$ .

Set  $\bar{m}_2 = (i_0, j_0)$  (g.c.d of  $i_0, j_0$ ). Then

$$F_{\text{prim}}^{(k)}(x, y) = (y^k + x)^{\bar{i}_1} y^{\bar{i}_2}, \quad \text{where } \bar{i}_1 = \frac{i_0}{\bar{m}_2}, \quad \bar{i}_2 = \frac{j_0}{\bar{m}_2}. \quad \text{Thus } d_2^{(k)} = k\bar{i}_1 + \bar{i}_2. \quad (3.84)$$

So  $m_2^{(k)} = \bar{m}_2$  does not depend on  $k$  (recall definition of  $m_2$  in the statement before (3.24) and  $m_2^{(k)}$  is the “ $\langle k \rangle$ ” version of  $m_2$ ). Similarly,  $n_2^{(k)} := \frac{n^{(k)}}{d_2^{(k)}}$  should not depend on  $k$ , denoted it by  $\bar{n}_2$ .

In the following, we set

$$f'_{ij} = 0 \quad \text{if } (i, j) \notin S. \quad (3.85)$$

For each  $r \in \mathbb{Z}_+$ , there exists a unique pair  $(s_r, t_r) \in \mathbb{Z}_+^2$  such that  $ks_r + t_r = m^{(k)} - r$  and  $0 \leq t_r < k$ . Thus condition (3.80) and equation (3.77) show

$$F^{(k)}(x, y)^{\lfloor \frac{r}{k} \rfloor} = f'_{s_r, t_r} (y^k + x)^{s_r} y^{t_r}. \quad (3.86)$$

Set

$$(i'_0, j'_0) = \max(S \setminus \{(i_0, j_0)\}). \quad (3.87)$$

Such  $(i'_0, j'_0)$  must exist otherwise  $S = \{(i_0, j_0)\}$  and  $F^{(k)}(x, y) = (y^k + x)^{i_0} y^{j_0}$  cannot be an element of a Jacobian pair. Then (3.86) proves

**Claim 3** The first nonzero component  $F^{(k)}(x, y)^{\lfloor \frac{r}{k} \rfloor}$  with  $r \neq 0$  is the  $\frac{r^{(k)}}{k}$ -th component with

$$r^{(k)} := m^{(k)} - (ki'_0 + j'_0), \quad \text{and } F^{(k)}(x, y)^{\lfloor \frac{r^{(k)}}{k} \rfloor} = f'_{i'_0, j'_0} (y^k + x)^{i'_0} y^{j'_0}. \quad (3.88)$$

If necessary, by applying the automorphism  $(x, y) \mapsto (x + \alpha, y)$  for some  $\alpha \in \mathbb{F}$  (which then produces a term  $i_0\alpha(y^k + x)^{i_0-1}y^{j_0}$ ), we can always suppose  $f'_{i_0-1, j_0} \neq 0$  for some  $j_0$ . Thus

$$i'_0 \geq i_0 - 1 \geq 1. \quad (3.89)$$

We have the “ $\langle k \rangle$ ” version of (3.9), which is rewritten as

$$G^{(k)}(x, y) = H^{(k)}(x, y) + K^{(k)}(x, y), \quad \text{where} \quad (3.90)$$

$$H^{(k)}(x, y) = \sum_{0 \leq s < j^{(k)}} b_s^{(k)} F^{(k)}(x, y)^{\frac{n^{(k)}-s}{m^{(k)}}}, \quad K^{(k)}(x, y) = \sum_{t=j^{(k)}}^{\infty} b_t^{(k)}(x) F^{(k)}(x, y)^{\frac{n^{(k)}-t}{m^{(k)}}}, \quad (3.91)$$

where  $j^{(k)} = m^{(k)} + n^{(k)} - k - 1$ , and  $b_t^{(k)}(x) = b_t^{(k)} \in \mathbb{F}$  if  $t < m^{(k)} + n^{(k)} - 1$ . Since Subcase 2.2.2 cannot occur for our  $F^{(k)}(x, y)$ , we in fact have  $j^{(k)} = \min B^{(k)}$  (cf. (3.55)). Thus  $b_s^{(k)} = 0$  if  $s < j^{(k)}$  and if  $d_2^{(k)} \nmid s$ .

Set  $b'_s{}^{(k)} = b_{sd_2}{}^{(k)}$  for  $s < s^{(k)} := \frac{j^{(k)}}{d_2^{(k)}}$ . Then  $H^{(k)}(x, y)$  can be rewritten as

$$H^{(k)}(x, y) = \sum_{0 \leq s < s^{(k)}} b'_s{}^{(k)} F^{(k)}(x, y) \frac{n^{(k)} - sd_2^{(k)}}{m^{(k)}}. \quad (3.92)$$

(In fact one can prove  $b'_s{}^{(k)}$  does not depend on  $k$  for  $s < s^{(k)}$ , but we do not need this.) Write  $K^{(k)}(x, y)$  in terms of its components, we can rewrite (3.90) as

$$G^{(k)}(x, y) = H^{(k)}(x, y) + \sum_{r=0}^{\infty} L_{k,r}(x, y), \quad \text{where } L_{k,r}(x, y) = K^{(k)}(x, y)^{\lfloor \frac{r}{k} \rfloor}. \quad (3.93)$$

We want to compare  $L_{k,r}(x, y)$ 's for different  $k$ . Denote

$$z_k = y^k + x, \quad z = y, \quad (3.94)$$

which are generators of  $\mathbb{F}[x, y]$ . In (3.90), computing the  $\frac{j^{(k)}+r}{k}$ -th component (where the first equality follows from the fact that  $j^{(k)} + r > n^{(k)}$ )

$$0 = G^{(k)}(x, y)^{\lfloor \frac{j^{(k)}+r}{k} \rfloor} = H^{(k)}(x, y)^{\lfloor \frac{j^{(k)}+r}{k} \rfloor} + L_{k,r}(x, y) \quad (\text{cf. Fact (ii)}), \quad (3.95)$$

shows that  $L_{k,r}(x, y)$  is a rational function of the form (cf. arguments before (3.57), in fact  $L_{k,0}(x, y)$  is the “ $\langle k \rangle$ ” version of  $G'_1(x, y)$  in (3.58))

$$L_{k,r}(x, y) = -H^{(k)}(x, y)^{\lfloor \frac{j^{(k)}+r}{k} \rfloor} = z_k^{\alpha_r^{(k)}} z^{\beta_r^{(k)}} Q_r^{(k)}, \quad (3.96)$$

(cf. also the last term of (3.59) and (3.84), (3.94)) for some  $\alpha_r^{(k)}, \beta_r^{(k)} \in \mathbb{Z}$  and  $Q_r^{(k)} \in \mathbb{F}[x, y]$  such that  $z_k, z$  does not divide  $Q_r^{(k)}$ .

First, (3.50), (3.59), (3.84) and Lemma 3.10 show that (note that if a non-integer power of  $x$  appears, then the coefficient is zero)

$$\begin{aligned} L_{k,0}(x, y) &= b_{j^{(k)}}^{(k)} F_{\text{prim}}^{(k)}(x, y)^{\frac{k+1-m^{(k)}}{d_2^{(k)}}} + c^{(k)} \sum_{i=0}^{\infty} \frac{1-i}{1+\frac{i}{k}} \bar{b}_{i,0}^{(k)} x^{1+\frac{i}{k}} F_{\text{prim}}^{(k)}(x, y)^{\frac{1-i-m^{(k)}}{d_2^{(k)}}} \\ &= \bar{p}_0 z_k^{-\bar{i}_1 \bar{m}_2 + 1} z^{-\bar{i}_2 \bar{m}_2 + 1}, \end{aligned} \quad (3.97)$$

where by (3.82) and (3.63),

$$b_{j^{(k)}}^{(k)} = \bar{p}_0 = f_{i_0, j_0}^{-1} \alpha \frac{J(F, G)}{(\bar{i}_1 - \bar{i}_2) \bar{m}_2} \in \mathbb{F}^* \quad (\text{which does not depend on } k). \quad (3.98)$$

Computing the  $\frac{r}{k}$ -th component of the “ $\langle k \rangle$ ” version of (3.12) for  $0 < r \leq r^{(k)}$ , using induction on  $r$ , by Claim 3, we have (cf. (3.48))

$$\bar{b}_{i,r}^{(k)} := \text{Coeff}(\bar{b}_i^{(k)}(x), x^{\frac{i-r}{k}}) = 0 \quad \text{if } 0 < r < r^{(k)}, \quad \text{and} \quad (3.99)$$

$$\begin{aligned} 0 &= \sum_{i=0}^{\infty} \frac{1-i}{m^{(k)}} \bar{b}_{i,0}^{(k)} x^{\frac{i}{k}} F^{(k)}(x, y)^{[\frac{r^{(k)}}{k}]} F_{\text{prim}}^{(k)}(x, y)^{\frac{1-i-m^{(k)}}{d_2^{(k)}}} + \sum_{i=r^{(k)}}^{\infty} \bar{b}_{i,r^{(k)}}^{(k)} x^{\frac{i-r^{(k)}}{k}} F_{\text{prim}}^{(k)}(x, y)^{\frac{1-i}{d_2^{(k)}}} \\ &= (\text{denoted by}) F_{1,r^{(k)}} + F_{2,r^{(k)}}, \end{aligned} \quad (3.100)$$

(where (3.99) is obtained as follows: first by induction on  $r$ , we obtain (3.100) with  $r^{(k)}$  replaced by  $r$ , then use the fact that  $F^{(k)}(x, y)^{[\frac{r}{k}]} = 0$  if  $0 < r < r^{(k)}$ ). Using (3.91), (3.96) and (3.99) (recalling the relation (3.49)), we have

$$\begin{aligned} L_{k,r^{(k)}}(x, y) &= \bar{p}_0 \frac{k+1-m^{(k)}}{m^{(k)}} F^{(k)}(x, y)^{[\frac{r^{(k)}}{k}]} F_{\text{prim}}^{(k)}(x, y)^{\frac{k+1-2m^{(k)}}{d_2^{(k)}}} \\ &\quad + b_{j^{(k)}+r^{(k)}}^{(k)}(0) F_{\text{prim}}^{(k)}(x, y)^{\frac{k+1-r^{(k)}-m^{(k)}}{d_2^{(k)}}} \\ &\quad + c^{(k)} \sum_{i=0}^{\infty} \frac{1-i}{1+\frac{i}{k}} \frac{1-m^{(k)}-i}{m^{(k)}} \bar{b}_{i,0}^{(k)} x^{1+\frac{i}{k}} F^{(k)}(x, y)^{[\frac{r^{(k)}}{k}]} F_{\text{prim}}^{(k)}(x, y)^{\frac{1-i-2m^{(k)}}{d_2^{(k)}}} \\ &\quad + c^{(k)} \sum_{i=r^{(k)}}^{\infty} \frac{1-i}{1+\frac{i-r^{(k)}}{k}} \bar{b}_{i,r^{(k)}}^{(k)} x^{1+\frac{i-r^{(k)}}{k}} F_{\text{prim}}^{(k)}(x, y)^{\frac{1-i-m^{(k)}}{d_2^{(k)}}} \end{aligned}$$

(denote the last two terms by  $G_{1,r^{(k)}}$ ,  $G_{2,r^{(k)}}$ ). (3.101)

(Recall that if  $j^{(k)} + r < m^{(k)} + n^{(k)} - 1$  then  $b_{j^{(k)}+r}^{(k)} = b_{j^{(k)}+r}^{(k)}(x)$  does not depend on  $k$ .) From (3.100), we obtain

$$F_{2,r^{(k)}} = -F_{1,r^{(k)}} = -\frac{1}{\bar{m}_2} F_{\text{prim}}^{(k)}(x, y)^{1-\bar{m}_2} (\partial_y F_{\text{prim}}^{(k)}(x, y))^{-1} F^{(k)}(x, y)^{[\frac{r^{(k)}}{k}]}, \quad (3.102)$$

where the second equality follows by taking  $\partial_y$  in the “ $\langle k \rangle$ ” version of (3.51). Writing  $\frac{1-m^{(k)}-i}{m^{(k)}} = \frac{k+1-m^{(k)}}{m^{(k)}} - \frac{k}{\bar{m}_2} \frac{1+\frac{i}{k}}{d_2^{(k)}}$  in the expression of  $G_{1,r^{(k)}}$ , by using (3.97) and the equation which is obtained from taking  $\partial_y$  in the “ $\langle k \rangle$ ” version of (3.51), we can obtain

$$\begin{aligned} G_{1,r^{(k)}} &= \left( \frac{k+1-m^{(k)}}{m^{(k)}} \bar{p}_0 (z_k^{-\bar{i}_1 \bar{m}_2 + 1} z_k^{-\bar{i}_2 \bar{m}_2 + 1} - F_{\text{prim}}^{(k)}(x, y)^{\frac{k+1-m^{(k)}}{d_2^{(k)}}}) F_{\text{prim}}^{(k)}(x, y)^{-\bar{m}_2} \right. \\ &\quad \left. - \frac{k}{\bar{m}_2} c^{(k)} x F_{\text{prim}}^{(k)}(x, y)^{1-2\bar{m}_2} (\partial_y F_{\text{prim}}^{(k)}(x, y))^{-1} \right) F^{(k)}(x, y)^{[\frac{r^{(k)}}{k}]}. \end{aligned} \quad (3.103)$$

Similarly, writing  $\frac{1-i}{1+\frac{i-r^{(k)}}{k}} = -k + \frac{k+1-r^{(k)}}{1+\frac{i-r^{(k)}}{k}}$ , and making use of the expression of  $F_{2,r^{(k)}}$  in

(3.100), we can write  $G_{2,r^{(k)}}$  as

$$G_{2,r^{(k)}} = -kc^{(k)} x F_{2,r^{(k)}} F_{\text{prim}}^{(k)}(x, y)^{-\bar{m}_2} + (k+1-r^{(k)})c^{(k)} W F_{\text{prim}}^{(k)}(x, y)^{\frac{k+1-r^{(k)}}{d_2^{(k)}} - \bar{m}_2}, \quad (3.104)$$

where  $W = \sum \frac{1}{1+\frac{i-r^{(k)}}{k}} \bar{b}_{i,r^{(k)}}^{(k)} x^{1+\frac{i-r^{(k)}}{k}} F_{\text{prim}}^{(k)}(x, y)^{-\frac{k+i-r^{(k)}}{d_2^{(k)}}$  satisfies (using (3.102))

$$\partial_y W = \frac{k}{m^{(k)}} x F_{\text{prim}}^{(k)}(x, y)^{\frac{r^{(k)}-1-k}{d_2^{(k)}} - \bar{m}_2} F^{(k)}(x, y)^{\lfloor \frac{r^{(k)}}{k} \rfloor}. \quad (3.105)$$

Solving  $W$  from (3.104), using (3.84), (3.96), (3.100)–(3.103) and Claim 3, we obtain (where the terms marked with --(i)-- are canceled for  $i = 1, 2$ )

$$\begin{aligned} & (k+1-r^{(k)})c^{(k)}W \\ &= \left( z_k^{\alpha_{r^{(k)}}^{(k)}} z_{r^{(k)}}^{\beta_{r^{(k)}}^{(k)}} Q_{r^{(k)}}^{(k)} - \bar{p}_0 \frac{k+1-m^{(k)}}{m^{(k)}} F^{(k)}(x, y)^{\lfloor \frac{r^{(k)}}{k} \rfloor} F_{\text{prim}}^{(k)}(x, y)^{\frac{k+1-2m^{(k)}}{d_2^{(k)}}} \right. \\ & \quad \left. - b_{j^{(k)}+r^{(k)}}^{(k)}(0) F_{\text{prim}}^{(k)}(x, y)^{\frac{k+1-r^{(k)}-m^{(k)}}{d_2^{(k)}}} \right. \\ & \quad \left. - \frac{k+1-m^{(k)}}{m^{(k)}} \bar{p}_0 (z_k^{-\bar{i}_1 \bar{m}_2 + 1} z^{-\bar{i}_2 \bar{m}_2 + 1} - F_{\text{prim}}^{(k)}(x, y)^{\frac{k+1-m^{(k)}}{d_2^{(k)}}}) F_{\text{prim}}^{(k)}(x, y)^{-\bar{m}_2} F^{(k)}(x, y)^{\lfloor \frac{r^{(k)}}{k} \rfloor} \right. \\ & \quad \left. + \frac{k}{\bar{m}_2} c^{(k)} x F_{\text{prim}}^{(k)}(x, y)^{1-2\bar{m}_2} (\partial_y F_{\text{prim}}^{(k)}(x, y))^{-1} F^{(k)}(x, y)^{\lfloor \frac{r^{(k)}}{k} \rfloor} \right. \\ & \quad \left. - \frac{k}{\bar{m}_2} c^{(k)} x F_{\text{prim}}^{(k)}(x, y)^{1-2\bar{m}_2} (\partial_y F_{\text{prim}}^{(k)}(x, y))^{-1} F^{(k)}(x, y)^{\lfloor \frac{r^{(k)}}{k} \rfloor} F_{\text{prim}}^{(k)}(x, y)^{\frac{r^{(k)}-1-k}{d_2^{(k)}} + \bar{m}_2} \right. \\ & \quad \left. = z_k^{\alpha_{r^{(k)}}^{(k)} + \bar{i}_1 (\frac{r^{(k)}-1-k}{d_2^{(k)}} + \bar{m}_2)} z_{r^{(k)}}^{\beta_{r^{(k)}}^{(k)} + \bar{i}_2 (\frac{r^{(k)}-1-k}{d_2^{(k)}} + \bar{m}_2)} Q_{r^{(k)}}^{(k)} - b_{j^{(k)}+r^{(k)}}^{(k)}(0) \right. \\ & \quad \left. - \frac{k+1-m^{(k)}}{m^{(k)}} \bar{p}_0 f'_{i'_0, j'_0} z_k^{-2\bar{i}_1 \bar{m}_2 + 1 + i'_0 + \bar{i}_1 (\frac{r^{(k)}-1-k}{d_2^{(k)}} + \bar{m}_2)} z^{-2\bar{i}_2 \bar{m}_2 + 1 + j'_0 + \bar{i}_2 (\frac{r^{(k)}-1-k}{d_2^{(k)}} + \bar{m}_2)} \right). \quad (3.106) \end{aligned}$$

Using (3.106) in (3.105), we have

$$\begin{aligned} & \xi_1^{(k)} x z_k^{-\bar{i}_1 \bar{m}_2 + i'_0} z^{-\bar{i}_2 \bar{m}_2 + j'_0} + (\xi_2^{(k)} y^k + \xi_3^{(k)} x) z_k^{\bar{i}_1 \bar{m}_2 + \alpha_{r^{(k)}}^{(k)} - 1} z^{-\bar{i}_2 \bar{m}_2 + \beta_{r^{(k)}}^{(k)} - 1} Q_{r^{(k)}}^{(k)} \\ & + z_k^{\bar{i}_1 \bar{m}_2 + \alpha_{r^{(k)}}^{(k)}} z^{-\bar{i}_2 \bar{m}_2 + \beta_{r^{(k)}}^{(k)}} \partial_y Q_{r^{(k)}}^{(k)} = 0, \quad (3.107) \end{aligned}$$

where (note from Lemma 3.1 and (3.98) that  $c^{(k)} = -f_{i_0, j_0}^{-1} \alpha \frac{J(F, G)}{m^{(k)}}$ )

$$\begin{aligned} \xi_1^{(k)} &= \frac{m^{(k)} - k - 1}{m^{(k)}} \left( j'_0 + 1 - \frac{\bar{i}_2}{d_2^{(k)}} (k(i'_0 + 1) + j'_0 + 1) \right) \bar{p}_0 f'_{i'_0, j'_0} \\ &\quad + \frac{k}{m^{(k)}} c^{(k)} (m^{(k)} - k(i'_0 + 1) - j'_0 - 1) f'_{i'_0, j'_0}, \\ &= \frac{i'_0(1 - \bar{i}_2 \bar{m}_2) - j'_0(1 - \bar{i}_1 \bar{m}_2)}{(\bar{i}_1 - \bar{i}_2) \bar{m}_2 m^{(k)}} k f'_{i'_0, j'_0} f_{i_0, j_0}^{-1} \alpha J(F, G), \end{aligned} \quad (3.108)$$

$$\xi_2^{(k)} = 2m^{(k)} - k(i'_0 + 1) - j'_0 - 1 + \alpha_{r^{(k)}}^{(k)} k + \beta_{r^{(k)}}^{(k)}, \quad (3.109)$$

$$\xi_3^{(k)} = 2\bar{i}_2 \bar{m}_2 - (k(i'_0 + 1) + j'_0 + 1) \frac{\bar{i}_2}{d_2^{(k)}} + \beta_{r^{(k)}}^{(k)}. \quad (3.110)$$

Assume that  $(i'_0, j'_0)$  satisfies the condition

$$i'_0(1 - \bar{i}_2 \bar{m}_2) - j'_0(1 - \bar{i}_1 \bar{m}_2) \neq 0. \quad (3.111)$$

Then  $\xi_1^{(k)} \neq 0$ . Dividing (3.107) by  $z_k^{-\bar{i}_1 \bar{m}_2 + i'_0} z^{-\bar{i}_2 \bar{m}_2 + j'_0}$  shows

$$-\bar{i}_1 \bar{m}_2 + i'_0 \geq \bar{i}_1 \bar{m}_2 + \alpha_{r^{(k)}}^{(k)} - 1, \quad (3.112)$$

$$-\bar{i}_2 \bar{m}_2 + j'_0 \geq \bar{i}_2 \bar{m}_2 + \beta_{r^{(k)}}^{(k)} - 1. \quad (3.113)$$

If the equality in (3.113) does not hold, then canceling the common factor  $z^{\bar{i}_2 \bar{m}_2 + \beta_{r^{(k)}}^{(k)} - 1}$  in (3.107) shows that  $z$  must divide  $\xi_2^{(k)} y^k + \xi_3^{(k)} x$ , namely,  $\xi_3^{(k)} = 0$ . Then (3.110) shows that

$$u_k := \frac{k(i'_0 + 1) + j'_0 + 1}{k\bar{i}_1 + \bar{i}_2} \bar{i}_2 \quad \text{must be an integer.} \quad (3.114)$$

Hence  $u := \frac{i'_0 + 1}{\bar{i}_1} \bar{i}_2 = \lim_{k \rightarrow \infty} u_k \in \mathbb{Z}$ . Thus  $u_k = u$  does not depend on  $k$  when  $k \gg 0$ , this in particular shows  $(i'_0 + 1)\bar{i}_2 = (j'_0 + 1)\bar{i}_1$ . But then (3.113) become the equality. This contradiction shows that the equality in (3.113) holds when  $k \gg 0$ . Similarly, if the equality in (3.112) does not hold, then  $\xi_2^{(k)} = \xi_3^{(k)}$ , and we can again obtain a contradiction. This proves that the equalities must hold in (3.112), (3.113) when  $k \gg 0$ , i.e.,

$$\alpha_{r^{(k)}}^{(k)} = -2\bar{i}_1 \bar{m}_2 + i'_0 + 1, \quad \beta_{r^{(k)}}^{(k)} = -2\bar{i}_2 \bar{m}_2 + j'_0 + 1, \quad (3.115)$$

$$\xi_2^{(k)} = 0, \quad \xi_3 = \frac{\bar{i}_1(j'_0 + 1) - \bar{i}_2(i'_0 + 1)}{d_2^{(k)}} k, \quad (3.116)$$

where (3.116) follows from (3.115), (3.109) and (3.110). Note from (3.91) that

$$\deg_y L_{k,r}(x, y) \leq (n^{(k)} - j^{(k)}) - r = k + 1 - m^{(k)} - r. \quad (3.117)$$

For any  $A = \sum_{i,j} a_{ij}x^i y^j \in \mathbb{F}[x]((y^{-1}))$ ,  $a_{ij} \in \mathbb{F}$  with prime degree  $p$ , we define the *quasi-degree*  $\text{qdeg } A$  to be

$$\text{qdeg } A = \max\left\{\frac{i}{p} + j \mid a_{ij} \neq 0\right\}. \quad (3.118)$$

Then from (3.91), one can easily prove  $\text{qdeg } K^{(k)}(x, y) \leq n^{(k)} - j^{(k)} = k + 1 - m^{(k)}$ . Thus (cf. (3.117))

$$\text{qdeg } L_{k,r}(x, y) \leq k + 1 - m^{(k)} - r. \quad (3.119)$$

(In fact the equality holds.) Then (3.119) and (3.96) (cf. (3.88)) show  $\text{qdeg}_y Q_{r^{(k)}}^{(k)} \leq 0$ . This forces  $Q_{r^{(k)}}^{(k)} \in \mathbb{F}^*$  since  $Q_{r^{(k)}}^{(k)}$  is a polynomial. Then (3.107) shows and

$$\xi^{(k)} := Q_{r^{(k)}}^{(k)} = -\frac{\xi_1^{(k)}}{\xi_3^{(k)}} = -\frac{i'_0(1 - \bar{i}_2 \bar{m}_2) - j'_0(1 - \bar{i}_1 \bar{m}_2)}{(\bar{i}_1 - \bar{i}_2)(\bar{i}_1(1 + j'_0) - \bar{i}_2(1 + i'_0))\bar{m}_2^2} f'_{i'_0, j'_0} f_{i_0, j_0}^{-1} \alpha J(F, G), \quad (3.120)$$

$$L_{k,r^{(k)}}(x, y) = \xi^{(k)} z_k^{-2\bar{i}_1 \bar{m}_2 + i'_0 + 1} z^{-2\bar{i}_2 \bar{m}_2 + j'_0 + 1}. \quad (3.121)$$

Comparing the coefficients of  $y^{k+1-m^{(k)}-r}$  in (3.101) shows (in the right-hand side of (3.101), only the first two terms contain  $y^{k+1-m^{(k)}-r}$ , and the degrees of  $y$  for all other terms are  $\leq 1 - m^{(k)} - r$ )

$$\begin{aligned} b_{j^{(k)}+r^{(k)}}^{(k)}(0) &= \xi^{(k)} - \frac{k+1-m^{(k)}}{m^{(k)}} \bar{p}_0 \\ &= \frac{\bar{i}_2 \bar{m}_2 - j'_0 - 1 + k(\bar{i}_1 \bar{m}_2 - i'_0 - 1)}{(\bar{i}_1(j'_0 + 1) - \bar{i}_2(i'_0 + 1))\bar{m}_2 m^{(k)}} f'_{i'_0, j'_0} f_{i_0, j_0}^{-1} \alpha J(F, G). \end{aligned} \quad (3.122)$$

Now let us return to (3.74) to consider our original  $F(x, y)$ . As in the proof of Claim 1, we in fact have

$$q' = 1 \quad \text{and so} \quad m = (i_1 + i_2)m_2. \quad (3.123)$$

To convince readers that our approach works, we now give two proofs (the second proof, which in fact seems to be simpler, does not need to use the projection  $\tau_{\mathbf{t}}$  defined in (3.124)).

*The first proof.* Let  $\mathbf{t}$  be a variable in  $\mathbb{F}$ . Define the projection  $\tau_{\mathbf{t}} : \mathbb{F}[\mathbf{t}][x]((y^{-1})) \mapsto \mathbb{F}[x]((y^{-1})) \cdot \mathbf{t}$  by

$$\tau_{\mathbf{t}}(A) = A_1 \mathbf{t} \quad \text{if } A = A_0 + A_1 \mathbf{t} + A_2 \mathbf{t}^2 + \cdots, \quad \text{where } A_i \in \mathbb{F}[x]((y^{-1})). \quad (3.124)$$

In the following, for simplicity, we shall regard  $\mathbf{t}$  as a fixed element in  $\mathbb{F}$  so that elements in  $\mathbb{F}[\mathbf{t}]$  can be regarded as elements in  $\mathbb{F}$ .

Replacing  $F(x, y)$  by  $F(x - \mathbf{t}, y + \mathbf{t})$ , i.e., by applying the automorphism  $(x, y) \mapsto (x - \mathbf{t}, y + \mathbf{t})$ , we have

$$\begin{aligned} F(x, y) &= (y + \mathbf{t} + x - \mathbf{t})^{i_1 m_2} (y + \mathbf{t})^{i_2 m_2} + \dots \\ &= (y + x)^{i_1 m_2} y^{i_2 m_2} + i_2 m_2 \mathbf{t} (y + x)^{i_1 m_2} y^{i_2 m_2 - 1} + \dots, \end{aligned}$$

and we obtain (using notation  $f_{ij}$  as in (3.74))

**Claim 4**  $\tau_{\mathbf{t}}(f_{i_1 m_2, i_2 m_2 - 1}) = i_2 m_2 \mathbf{t}$ , and  $(i_1 m_2, i_2 m_2 - 1)$  is the unique pair  $(i, j)$  with  $i + j = m - 1$  satisfying  $\tau_{\mathbf{t}}(f_{ij}) \neq 0$ .

The proof of this claim is straightforward (or it is similar to the proof of the next claim, so we omit it).

Note that  $F(x, y)$  is in fact  $F^{(1)}(x, y)$ . From now on, we shall use the new order on  $S$  (cf. notation  $S$  in (3.75)) defined by

$$(i, j) < (i', j') \iff i + j < i' + j', \quad \text{or } i + j = i' + j' \text{ and } i < i'. \quad (3.125)$$

Now we define new  $(i_0, j_0)$  and  $(i'_0, j'_0)$  by

$$(i_0, j_0) := \max S = (i_1 m_2, i_2 m_2), \quad (3.126)$$

$$(i'_0, j'_0) := \max\{(i, j) \in S \mid \tau_{\mathbf{t}}(f_{ij}) \neq 0\} = (i_0, j_0 - 1). \quad (3.127)$$

Using notations as above, we shall be interested in computing  $\tau_{\mathbf{t}}(F^{(1)}(x, y)^{[1]})$ . Set

$$r^{(1)} = m - (i'_0 + j'_0) = 1.$$

All arguments from Claim 3 to (3.110) will be still valid if we apply  $\tau_{\mathbf{t}}$  to every expression, i.e., if we replace respectively

$$F^{(1)}(x, y)^{[1]}, \quad b_{j^{(1)} + r^{(1)}}^{(1)}(0), \quad \bar{b}_{i, r^{(1)}}^{(1)}, \quad Q_{r^{(1)}}^{(1)}, \quad \text{etc.},$$

by

$$\tau_{\mathbf{t}}(F^{(1)}(x, y)^{[1]}), \quad \tau_{\mathbf{t}}(b_{j^{(1)} + r^{(1)}}^{(1)}(0)), \quad \tau_{\mathbf{t}}(\bar{b}_{i, r^{(1)}}^{(1)}), \quad \tau_{\mathbf{t}}(Q_{r^{(1)}}^{(1)}), \quad \text{etc.}$$

(Remark: In the right-hand side of (3.95), the first term in general contains polynomials of  $\mathbf{t}$  with degree  $\geq 1$ , but the second term only contains polynomials of  $\mathbf{t}$  with degree 1. Thus when we apply  $\tau_{\mathbf{t}}$ , we see that any  $\mathbf{t}^i$  for  $i > 1$  which appears in the first term does not contribute to our computation.)

Note that the data  $(k, i'_0, j'_0, \bar{i}_1, \bar{i}_2)$  appearing in (3.111) is now  $(1, i_1 m_2, i_2 m_2 - 1, i_1, i_2)$ , and  $m_2 \geq 2$ . We have

$$i'_0(1 - i_2 m_2) - j'_0(1 - i_1 m_2) = i_0(1 - j_0) - (j_0 - 1)(1 - i_0) = 1 - i_2 m_2 < 0,$$

i.e., condition (3.111) holds. Also  $u_k$  defined in (3.114) is now

$$u_1 = \frac{m_2(i_1 + i_2) + 1}{i_1 + i_2} i_2, \quad \text{which is not an integer.} \quad (3.128)$$

So the arguments before and after (3.114) then show that equalities in (3.112) and (3.113) must hold. Hence (3.115) and (3.116) still hold. Thus (3.120) and (3.119) give (cf. Claim 4, note that the data  $(m^{(k)}, f'_{i'_0, j'_0}, f_{i_0, j_0}^{-1}, \alpha)$  is now  $(m, j_0 \mathbf{t}, 1, 1)$ )

$$\tau_{\mathbf{t}}(L_{1,1}(x, y)) = \xi^{(1)}(y + x)^{-i_0+1} y^{-j_0}, \quad \text{where } \xi^{(1)} = \frac{1 - j_0}{i_0 - j_0} \mathbf{t} J(F, G). \quad (3.129)$$

Now let  $\ell \gg 0$ , and take the Jacobian pair  $(\check{F}^{(\ell)}(x, y), \check{G}^{(\ell)}(x, y))$  with (cf. notations (3.94))

$$\begin{aligned} \check{F}^{(\ell)}(x, y) &= 2^{-i_0} F(y^\ell - y + x, y^\ell + x) = 2^{-i_0} \left( (2z_\ell - z)^{i_0} z_\ell^{j_0} + j_0 \mathbf{t} (2z_\ell - z)^{i_0} z_\ell^{j_0-1} + \dots \right) \\ &= z_\ell^m - \frac{i_0}{2} z_\ell^{m-1} z + j_0 \mathbf{t} z_\ell^{m-1} + \frac{1}{4} \frac{i_0(i_0-1)}{2} j_0 \mathbf{t} z_\ell^{m-2} z^2 - \frac{i_0}{2} j_0 \mathbf{t} z_\ell^{m-2} z + \dots, \end{aligned} \quad (3.130)$$

$$\check{G}^{(\ell)}(x, y) = 2^{-i_1 n_2} G(y^\ell - y + x, y^\ell + x). \quad (3.131)$$

Thus  $\check{m}^{(\ell)} = \ell m$  (we use the same symbols with a “ $\check{\cdot}$ ” to denote corresponding notations, cf. Convention 3.12) and

$$(\check{i}_0, \check{j}_0) := \max \check{S} = (m, 0), \quad (3.132)$$

$$(\check{i}'_0, \check{j}'_0) := \max \{(i, j) \in \check{S} \mid \tau_{\mathbf{t}}(\check{f}_{ij}) \neq 0\} = (m-1, 0), \quad (3.133)$$

$$\check{r}^{(\ell)} := \check{m}^{(\ell)} - (\check{i}'_0 \ell + \check{j}'_0) = \ell, \quad (3.134)$$

where (3.133) follows from the third term in the right-hand side of (3.130).

**Claim 5** (cf. Claim 3)  $\check{r}^{(\ell)}$  is the smallest positive integer  $r$  such that

$$\tau_{\mathbf{t}}(\check{F}^{(\ell)}(x, y)^{\lfloor \frac{r}{\ell} \rfloor}) = j_0 \mathbf{t} z_\ell^{m-1} \neq 0.$$

If  $f_{ij}(y+x)^i y^j$  appears as a term in  $F(x, y)$ , then the following polynomial appears as a part in  $\check{F}^{(\ell)}(x, y)$ :

$$\tau_{\mathbf{t}}(2^{-i_0} f_{ij} (2z_\ell - z)^i z_\ell^j) = 2^{i-i_0} \tau_{\mathbf{t}}(f_{ij}) \sum_{s=0}^i \binom{i}{s} \left(\frac{-1}{2}\right)^s z_\ell^{i+j-s} z^s. \quad (3.135)$$

Note from (3.77) that the term  $z_\ell^{i+j-s} z^s$  belongs to the  $\frac{r}{\ell}$ -th component with

$$r := \check{m}^{(\ell)} - \ell(i + j - s) - s.$$

Assume that  $r \leq \check{r}^{(\ell)}$  and  $\tau_t(f_{ij}) \neq 0$ . Then Claim 4 shows that  $i+j \leq m-1$ , and so  $r \geq \check{r}^{(\ell)}$ . If  $r = \check{r}^{(\ell)}$ , then Claim 4 also shows that  $i+j = m-1$  (and in fact  $(i, j) = (i_0, j_0 - 1)$ ) and  $s = 0$ . Thus Claim 5 follows from (3.135) and Claim 4.

Note that the data  $(k, i'_0, j'_0, \bar{i}_1, \bar{i}_2)$  appearing in (3.111) is now  $(\ell, m-1, 0, 1, 0)$ . As before, by applying the projection  $\tau_t$ , all arguments from Claim 3 to (3.110) are still valid. (Note that all arguments do not depend on whether or not  $\bar{i}_2 = 0$  since (3.62) can be regarded as a special case of (3.63) with  $i_1 = 1, i_2 = 0$ . But  $u_k$  defined in (3.114) is now zero, we need to use different approach below, cf. (3.128).)

In our case here, (3.107)–(3.110) become

$$\check{\xi}_1^{(\ell)} x z_\ell^{-1} + (\check{\xi}_2^{(\ell)} y^\ell + \check{\xi}_3^{(\ell)} x) z_\ell^{m+\check{\alpha}_{\check{r}^{(\ell)}}^{(\ell)} - 1} z^{\check{\beta}_{\check{r}^{(\ell)}}^{(\ell)} - 1} \tau_t(\check{Q}_{\check{r}^{(\ell)}}^{(\ell)}) + z_\ell^{m+\check{\alpha}_{\check{r}^{(\ell)}}^{(\ell)}} z^{\check{\beta}_{\check{r}^{(\ell)}}^{(\ell)}} \partial_y \left( \tau_t(\check{Q}_{\check{r}^{(\ell)}}^{(\ell)}) \right) = 0, \quad (3.136)$$

and (note that the data  $(m^{(k)}, f'_{i'_0, j'_0}, f_{i_0, j_0}^{-1}, \alpha)$  is now  $(\ell m, j_0 t, 1, 1)$ )

$$\check{\xi}_1^{(\ell)} = \frac{m-1}{\ell m^2} j_0 t J(F, G), \quad \check{\xi}_2^{(\ell)} = \ell m - 1 + \check{\alpha}_{\check{r}^{(\ell)}}^{(\ell)} \ell + \check{\beta}_{\check{r}^{(\ell)}}^{(\ell)}, \quad \check{\xi}_3^{(\ell)} = \check{\beta}_{\check{r}^{(\ell)}}^{(\ell)}, \quad (3.137)$$

such that (3.112) and (3.113) become

$$-1 = m + \check{\alpha}_{\check{r}^{(\ell)}}^{(\ell)} - 1, \quad 0 \geq \check{\beta}_{\check{r}^{(\ell)}}^{(\ell)} - 1, \quad (3.138)$$

where the first equation follows from the same arguments after (3.113): we see from (3.137) that  $\check{\xi}_2^{(\ell)} \neq \check{\xi}_3^{(\ell)}$  since  $\ell \nmid (\ell m - 1)$ .

If the equality does not hold in the second inequality of (3.138), then as before,  $\check{\xi}_3^{(\ell)}$  must be zero. But then (3.136) becomes (multiplying it by  $z_\ell$ , and simply denoting  $Q = \tau_t(\check{Q}_{\check{r}^{(\ell)}}^{(\ell)})$ )

$$\check{\xi}_1^{(\ell)} x - y^{\ell-1} Q + (y^\ell + x) \partial_y Q = 0. \quad (3.139)$$

Suppose  $\deg_y Q = s$ . Computing the coefficient of  $y^{\ell+s-1}$  of (3.139) shows  $s = 1$ . Then one can solve  $Q = -\check{\xi}_1^{(\ell)} y$  which is then divided by  $z = y$ . This is a contradiction (cf. the statement after (3.96)). Thus the equality holds in (3.138). Then again we have the “ $\tau_t$ ” version of (3.122), which is now has the form

$$\begin{aligned} \tau_t(\check{b}_{j^{(\ell)} + \check{r}^{(\ell)}}^{(\ell)}(0)) &= \frac{\check{j}_0 - \check{j}'_0 - 1 + \ell(\check{i}_0 - \check{i}'_0 - 1)}{(\check{i}_0(\check{j}'_0 + 1) - \check{j}_0(\check{i}'_0 + 1))\check{m}^{(k)}} j_0 t J(\check{F}, \check{G}) \\ &= -2^{-i_1(m_2+n_2)} \frac{1}{\ell m^2} j_0 t J(F, G). \end{aligned} \quad (3.140)$$

Noting from (3.91)–(3.93), we have

$$\begin{aligned}
\check{G}^{(\ell)}(x, y) &= 2^{-i_1 n_2} G^{(1)}(y^\ell + x - y, y^\ell + x) \\
&= \sum_{0 \leq s < j^{(1)}} 2^{-i_1 n_2} b_s^{(1)} F^{(1)}(y^\ell + x - y, y^\ell + x)^{\frac{n-s}{m}} + \sum_{r=0}^{\infty} 2^{-i_1 n_2} L_{1,r}(y^\ell + x - y, y^\ell + x) \\
&= \sum_{0 \leq s < j^{(1)}} 2^{-i_1 n_2 + i_1 m_2 \frac{n-s}{m}} b_s^{(1)} \check{F}^{(\ell)}(x, y)^{\frac{n-s}{m}} + \sum_{r=0}^{\infty} R_{\ell,r}(x, y), \tag{3.141}
\end{aligned}$$

where  $R_{\ell,r}(x, y) = 2^{-i_1 n_2} L_{1,r}(y^\ell + x - y, y^\ell + x)$ . We claim that the right-hand side of (3.141) is summable because from (3.119) we have

$$\deg_y R_{\ell,r}(x, y) \leq \ell(2 - m - r), \tag{3.142}$$

which tends to  $-\infty$  when  $r \rightarrow +\infty$ . To prove that (3.141) holds, it suffices to compare the coefficients of  $y^i$  in both sides for  $i \in \mathbb{Z}$ . To do this, we only need to consider (3.141) under modulo  $y^s \mathbb{F}[x][[y^{-1}]]$  for some  $s \ll 0$ . But in this case, (3.141) becomes a finite sum, and of course, variable changing  $(x, y) \mapsto (y^\ell + x - y, y^\ell + x)$  makes sense in a finite sum.

We shall use (3.141) to compute  $\tau_t(\check{b}_{j^{(\ell)} + \check{r}^{(\ell)}}^{(\ell)}(0))$  and compare it with (3.140). To do this, we need to rewrite  $R_{\ell,r}(x, y)$  into the form (recall (3.56))

$$R_{\ell,r}(x, y) \equiv \sum_{i \geq 1 - \ell m} \beta_{\ell,r,i}(x) \check{F}^{(\ell)}(x, y)^{\frac{i}{\ell m}} \pmod{y^{-\ell m} \mathbb{F}[x][[y^{-1}]]}, \tag{3.143}$$

for some  $\beta_{\ell,r,i}(x) \in \mathbb{F}[x]$ . Then

$$\tau_t(\check{b}_{j^{(\ell)} + \check{r}^{(\ell)}}^{(\ell)}(0)) = \tau_t(\beta_{\ell,0,1-\ell m}(0)) + \tau_t(\beta_{\ell,1,1-\ell m}(0)), \tag{3.144}$$

by noting that  $\check{b}_{j^{(\ell)} + \check{r}^{(\ell)}}^{(\ell)}(x)$  is the coefficient of  $\check{F}^{(\ell)}(x, y)^{\frac{1-\ell m}{\ell m}}$  when we expand  $\check{G}^{(\ell)}(x, y)$  as a combination of rational powers of  $\check{F}^{(\ell)}(x, y)$ , and noting from (3.142) that if  $r \geq 2$  then  $\deg_y R_{\ell,r}(x, y) \leq -\ell m$ , thus  $R_{\ell,r}(x, y)$  does not contribute to  $\tau_t(\check{b}_{j^{(\ell)} + \check{r}^{(\ell)}}^{(\ell)}(0))$ .

From (3.130), we have (we simply denote  $\check{F}^{(\ell)}(x, y)$  by  $\check{F}$ )

$$z_\ell = \left( \check{F} + \frac{i_0}{2} z_\ell^{m-1} z - j_0 \mathbf{t} z_\ell^{m-1} - \frac{1}{4} \frac{i_0(i_0-1)}{2} z_\ell^{m-2} z^2 + \frac{i_0}{2} j_0 \mathbf{t} z_\ell^{m-2} z + \dots \right)^{\frac{1}{m}}. \tag{3.145}$$

We shall use (3.145) to express  $R_{\ell,r}(x, y)$  into the form of (3.143). Note from (3.97) that (cf. (3.130))

$$R_{\ell,0}(x, y) = 2^{-i_1(m_2+n_2)+1} p_0 \left( z_\ell^{-m+2} + \frac{1}{2} (i_0 - 1) z_\ell^{-m+1} z + \frac{1}{4} \frac{i_0(i_0-1)}{2} z_\ell^{-m} z^2 + \dots \right), \tag{3.146}$$

where  $p_0 = \frac{J(F,G)}{i_0-j_0}$  by (3.63) and (3.126), and where the omitted terms are in  $y^{-\ell m}\mathbb{F}[x][[y^{-1}]]$ . Here and below the underlined element indicates that the corresponding term does not contribute to  $\tau_t(\check{b}_{j^{(\ell)}+\check{r}^{(\ell)}}^{(\ell)}(0))$ ; in the case in (3.146), one can see this from the following

$$z_\ell^{-m} z^2 \equiv \check{F}^{\frac{2-\ell m}{\ell m}} \pmod{y^{-\ell m}\mathbb{F}[x][[y^{-1}]]}. \quad (3.147)$$

Using (3.145), we obtain that the first term inside the bracket in (3.146) can be rewritten as

$$\begin{aligned} 2^{-i_1(m_2+n_2)+1} p_0 : z_\ell^{-m+2} \equiv & \check{F}^{\frac{2}{m}-1} + \left(\frac{2}{m} - 1\right) \left(\frac{i_0}{2} z_\ell^{m-1} z - j_0 \mathbf{t} z_\ell^{m-1} - \frac{1}{4} \frac{i_0(i_0-1)}{2} \underline{z_\ell^{m-2} z^2}\right. \\ & \left. + \frac{i_0}{2} j_0 \mathbf{t} \underbrace{z_\ell^{m-2} z}\right) \check{F}^{\frac{2}{m}-2} \pmod{y^{-\ell m}\mathbb{F}[x][[y^{-1}]]}, \end{aligned} \quad (3.148)$$

where we put a number before “:” in order to keep the record of the corresponding coefficient. Here and below the underbraced element indicates that its coefficient will contribute to  $\tau_t(\check{b}_{j^{(\ell)}+\check{r}^{(\ell)}}^{(\ell)}(0))$ ; in the case in (3.148), since the term  $z_\ell^{m-2} z \cdot \check{F}^{\frac{2}{m}-2}$  appears in (3.148) and

$$\deg_y(z_\ell^{m-2} z \check{F}^{\frac{2}{m}-2}) = 1 - \ell m,$$

we see that the coefficient

$$2^{-i_1(m_2+n_2)+1} p_0 \left(\frac{2}{m} - 1\right) \frac{i_0}{2} j_0 \mathbf{t} \quad (A.1)$$

contributes to  $\tau_t(\check{b}_{j^{(\ell)}+\check{r}^{(\ell)}}^{(\ell)}(0))$ . For the underlined element in (3.148), we have an expression of  $z_\ell^{m-2} z^2 \check{F}^{\frac{2}{m}-2}$  as in (3.147).

Using  $z = (z_\ell - x)^{\frac{1}{\ell}}$  and (3.145), the term  $z_\ell^{m-1} z \check{F}^{\frac{2}{m}-2}$  in (3.148) can be written as (we shall be only interested in computing coefficients which do not depend on  $x$ , thus in fact we can simply replace  $z = (z_\ell - x)^{\frac{1}{\ell}}$  by  $z_\ell^{\frac{1}{\ell}}$ ):

$$\begin{aligned} 2^{-i_1(m_2+n_2)+1} p_0 \left(\frac{2}{m} - 1\right) : & \check{F}^{\frac{2}{m}-2} \left( \check{F}^{\frac{1}{\ell m}} + \frac{1}{\ell m} \left( \frac{i_0}{2} \underline{z_\ell^{m-1} z} - j_0 \mathbf{t} \underbrace{z_\ell^{m-1}} \right) \check{F}^{\frac{1}{\ell m}-1} + \dots \right) \times \\ & \times \left( \check{F}^{1-\frac{1}{m}} + \left(1 - \frac{1}{m}\right) \left( \frac{i_0}{2} \underline{z_\ell^{m-1} z} - j_0 \mathbf{t} \underbrace{z_\ell^{m-1}} \right) \check{F}^{-\frac{1}{m}} + \dots \right). \end{aligned} \quad (3.149)$$

Thus

$$-2^{-i_1(m_2+n_2)+1} p_0 \left(\frac{2}{m} - 1\right) \frac{i_0}{2} j_0 \mathbf{t} \left(\frac{1}{\ell m} + 1 - \frac{1}{m}\right) \quad (A.2)$$

contributes to  $\tau_t(\check{b}_{j^{(\ell)}+\check{r}^{(\ell)}}^{(\ell)}(0))$  (for the underlined elements in (3.149), we have similar formulas as in (3.147)).

Now consider  $z_\ell^{m-1} \check{F}^{-2+\frac{2}{m}}$  in (3.148), which is written as

$$-2^{-i_1(m_2+n_2)+1} p_0 \left(\frac{2}{m} - 1\right) j_0 \mathbf{t} : \check{F}^{-2+\frac{2}{m}} \left( \check{F}^{1-\frac{1}{m}} + \frac{i_0}{2} \left(1 - \frac{1}{m}\right) \underbrace{z_\ell^{m-1} z \check{F}^{-\frac{1}{m}} + \dots} \right). \quad (3.150)$$

Thus

$$-2^{-i_1(m_2+n_2)+1} p_0 \left( \frac{2}{m} - 1 \right) \frac{i_0}{2} \left( 1 - \frac{1}{m} \right) j_0 \mathbf{t} \quad (\text{A.3})$$

contributes to  $\tau_{\mathbf{t}}(\check{b}_{j^{(\ell)} + \check{r}^{(\ell)}}^{(\ell)}(0))$ .

Next consider the term  $z_\ell^{-m+1} z$  in (3.146), which is written as in (3.149):

$$\begin{aligned} 2^{-i_1(m_2+n_2)+1} \frac{p_0}{2} (i_0 - 1) : & \left( \check{F}^{\frac{1}{m}-1} + \left( \frac{1}{m} - 1 \right) \left( \frac{i_0}{2} \underline{z_\ell^{m-1} z} - j_0 \mathbf{t} \underline{z_\ell^{m-1}} \right) \check{F}^{\frac{1}{m}-2} + \dots \right) \times \\ & \times \left( \check{F}^{\frac{1}{\ell m}} + \frac{1}{\ell m} \left( \frac{i_0}{2} \underline{z_\ell^{m-1} z} - j_0 \mathbf{t} \underline{z_\ell^{m-1}} \right) \check{F}^{\frac{1}{\ell m}-1} + \dots \right). \end{aligned} \quad (\text{3.151})$$

Thus

$$-2^{-i_1(m_2+n_2)+1} \frac{p_0}{2} (i_0 - 1) j_0 \mathbf{t} \left( \frac{1}{m} - 1 + \frac{1}{\ell m} \right) \quad (\text{A.4})$$

contributes to  $\tau_{\mathbf{t}}(\check{b}_{j^{(\ell)} + \check{r}^{(\ell)}}^{(\ell)}(0))$ .

Now consider  $R_{\ell,1}(x, y)$ , which is written as (cf. (3.146))

$$R_{\ell,1}(x, y) = 2^{-i_1(m_2+n_2)+1} \xi^{(1)} \left( z_\ell^{-m+1} - \frac{1}{2} (1 - i_0) \underline{z_\ell^{-m} z} + \dots \right), \quad (\text{3.152})$$

where  $\xi^{(1)}$  is defined in (3.129). Thus

$$-2^{-i_1(m_2+n_2)+1} \xi^{(1)} \frac{1}{2} (1 - i_0) \quad (\text{A.5})$$

contributes to  $\tau_{\mathbf{t}}(\check{b}_{j^{(\ell)} + \check{r}^{(\ell)}}^{(\ell)}(0))$ . Write the first term of (3.152) as

$$2^{-i_1(m_2+n_2)+1} \xi^{(1)} : \check{F}^{-1+\frac{1}{m}} + \frac{i_0}{2} \left( \frac{1}{m} - 1 \right) \underline{z_\ell^{m-1} z \check{F}^{-2+\frac{1}{m}}}. \quad (\text{3.153})$$

Thus

$$2^{-i_1(m_2+n_2)+1} \xi^{(1)} \frac{i_0}{2} \left( \frac{1}{m} - 1 \right) \quad (\text{A.6})$$

contributes to  $\tau_{\mathbf{t}}(\check{b}_{j^{(\ell)} + \check{r}^{(\ell)}}^{(\ell)}(0))$ .

Now taking the sum of (A.1)–(A.6) and subtracting the result by (3.140), and using  $J(F, G) = (i_0 - j_0) p_0$  (cf. (3.63)),  $\xi^{(1)} = (1 - j_0) p_0 \mathbf{t}$  (cf. (3.129)) and  $m = i_0 + j_0$  (cf. (3.123), (3.126)), from (3.144), we obtain

$$(m - 1)(m - 2) i_0 j_0 (i_0 - j_0) = 0. \quad (\text{3.154})$$

Since  $i_0 j_0 (i_0 - j_0) \neq 0$  (otherwise  $J(F, G) = p_0 (i_0 - j_0) = 0$ , a contradiction) and we assume  $m \geq 3$  (cf. (3.79)). This is a contradiction. This prove Lemma 3.11.

*The second proof.* (This proof does not need to use the projection  $\tau_t$ .) We start with our original  $F(x, y)$  in (3.74) satisfying (3.123):

$$F(x, y) = (y + x)^{i_0} y^{j_0} + \sum_{i+j=m-1} f_{ij} (y + x)^i y^j + \dots \quad (3.155)$$

Take  $\check{F}^{(\ell)}(x, y)$ ,  $\check{G}^{(\ell)}(x, y)$  as in (3.130) and (3.131). Then (again we simply denote  $\check{F} = \check{F}^{(\ell)}(x, y)$ )

$$\check{F} = z_\ell^m - \frac{i_0}{2} z_\ell^{m-1} z + \sum_{i+j=m-1} 2^{i-i_0} f_{ij} (z_\ell^{m-1} - \frac{i}{2} z_\ell^{m-2} z) + \frac{i_0(i_0-1)}{8} z_\ell^{m-2} z^2 + \dots \quad (3.156)$$

Similar to (3.132)–(3.134), we set

$$\check{m}^{(\ell)} = \ell m, \quad (\check{i}_0, \check{j}_0) = (m, 0), \quad (\check{i}'_0, \check{j}'_0) = (m-1, 1), \quad \check{r}^{(\ell)} = \check{m}^{(\ell)} - (\check{i}'_0 \ell + \check{j}'_0) = \ell - 1.$$

Then we have as in Claim 5:

**Claim 6**  $\check{r}^{(\ell)}$  is the first positive integer  $r$  such that  $\check{F}^{(\ell)}(x, y)^{[\frac{r}{\ell}]} \neq 0$ , and

$$\check{F}^{(\ell)}(x, y)^{[\frac{\check{r}^{(\ell)}}{\ell}]} = -\frac{i_0}{2} z_\ell^{m-1} z.$$

Now using the same arguments from (3.132) to (3.140), we have (cf. (3.136) and (3.137))

$$\check{\xi}_1^{(\ell)} x z_\ell^{-1} z + (\check{\xi}_2^{(\ell)} y^\ell + \check{\xi}_3^{(\ell)} x) z_\ell^{m+\check{\alpha}_{\check{r}^{(\ell)}}^{(\ell)}-1} z^{\check{\beta}_{\check{r}^{(\ell)}}^{(\ell)}-1} \check{Q}_{\check{r}^{(\ell)}}^{(\ell)} + z_\ell^{m+\check{\alpha}_{\check{r}^{(\ell)}}^{(\ell)}} z^{\check{\beta}_{\check{r}^{(\ell)}}^{(\ell)}} \partial_y \check{Q}_{\check{r}^{(\ell)}}^{(\ell)} = 0, \quad (3.157)$$

and (note that the data  $(m^{(k)}, f'_{i'_0, j'_0}, f_{i_0, j_0}^{-1}, \alpha)$  is now  $(\ell m, -\frac{i_0}{2}, 1, 1)$ )

$$\check{\xi}_1^{(\ell)} = \frac{1-m}{\ell m^2} i_0 J(F, G), \quad \check{\xi}_2^{(\ell)} = \ell m - 2 + \check{\alpha}_{\check{r}^{(\ell)}}^{(\ell)} \ell + \check{\beta}_{\check{r}^{(\ell)}}^{(\ell)}, \quad \check{\xi}_3^{(\ell)} = \check{\beta}_{\check{r}^{(\ell)}}^{(\ell)}, \quad (3.158)$$

such that (3.138) becomes

$$-1 = m + \check{\alpha}_{\check{r}^{(\ell)}}^{(\ell)} - 1, \quad 1 \geq \check{\beta}_{\check{r}^{(\ell)}}^{(\ell)} - 1. \quad (3.159)$$

Again as before, from (3.157) and (3.158), we see that the equality must hold in (3.159) (otherwise we would solve from (3.157) that  $\check{Q}_{\check{r}^{(\ell)}}^{(\ell)}$  is a polynomial of  $y$  with degree 2, which would be divided by  $z$ ). Then again we have (3.122), which is now has the form (cf. (3.140))

$$\begin{aligned} \check{b}_{\check{j}^{(\ell)} + \check{r}^{(\ell)}}^{(\ell)}(0) &= \frac{\check{j}_0 - \check{j}'_0 - 1 + \ell(\check{i}_0 - \check{i}'_0 - 1)}{(\check{i}_0(\check{j}'_0 + 1) - \check{j}_0(\check{i}'_0 + 1))\check{m}^{(k)}} \left(-\frac{i_0}{2}\right) J(\check{F}, \check{G}) \\ &= 2^{-i_1(m_2+n_2)-1} \frac{1}{\ell m^2} i_0 J(F, G) = 2^{-i_1(m_2+n_2)-1} \frac{1}{\ell m^2} i_0 (i_0 - j_0) p_0. \end{aligned} \quad (3.160)$$

We still have (3.141) and (3.142), while (3.143) and (3.144) should be rewritten as

$$R_{\ell,r}(x, y) \equiv \sum_{i \geq 2-\ell m} \beta_{\ell,r,i}(x) \check{F}^{(\ell)}(x, y)^{\frac{i}{\ell m}} \pmod{y^{1-\ell m} \mathbb{F}[x][[y^{-1}]],} \quad (3.161)$$

$$\check{b}_{j^{(\ell)}+\check{r}^{(\ell)}}^{(\ell)}(0) = \beta_{\ell,0,2-\ell m}(0) + \beta_{\ell,1,2-\ell m}(0). \quad (3.162)$$

**Claim 7**  $R_{\ell,1}(x, y)$  does not contribute to  $\check{b}_{j^{(\ell)}+\check{r}^{(\ell)}}^{(\ell)}(0)$ .

Recall notation  $R_{\ell,r}(x, y)$  in (3.141). Note that  $L_{1,r}(x, y)$  is a quasi-homogeneous polynomial of prime degree 1, which thus must be a linear combination of the form

$$(y+x)^i y^j \quad \text{for some } i, j \in \mathbb{Z} \text{ with } i+j=1-m \quad (\text{cf. (3.119)}).$$

Thus  $R_{\ell,1}(x, y)$  is a linear combination of the following (cf. the right-hand side of (3.156))

$$z_\ell^{1-m} - \frac{i}{2} \underline{z_\ell^{-m} z} + \dots, \quad (3.163)$$

where the underlined term has the same meaning as in (3.146); in our case here, it is in  $y^{1-\ell m} \mathbb{F}[x][[y^{-1}]]$ . Using (3.156) to express  $z_\ell$ , the first term in (3.163) can be written as

$$z_\ell^{1-m} = \check{F}^{\frac{1}{m}-1} + \left(\frac{1}{m} - 1\right) \left(\frac{i_0}{2} \underline{z_\ell^{m-1} z} + \dots\right) \check{F}^{\frac{1}{m}-2}, \quad (3.164)$$

since  $\deg_y(z_\ell^{m-1} z \check{F}^{\frac{1}{m}-2}) = 1 - \ell m < 2 - \ell m$ . Thus (3.163) does not contribute to  $\check{b}_{j^{(\ell)}+\check{r}^{(\ell)}}^{(\ell)}(0)$ , and the claim follows.

Using (3.97) and the definition of  $R_{\ell,r}(x, y)$  in (3.141), we have

$$R_{\ell,0}(x, y) = 2^{-i_1(m_2+n_2)+1} p_0 \left( z_\ell^{-m+2} + \frac{1}{2} (i_0 - 1) z_\ell^{-m+1} z + \frac{i_0(i_0-1)}{8} \underbrace{z_\ell^{-m} z^2 + \dots} \right). \quad (3.165)$$

Thus the coefficient

$$2^{-i_1(m_2+n_2)+1} p_0 \frac{i_0(i_0-1)}{8} \quad (\text{A'.1})$$

contributes to  $\check{b}_{j^{(\ell)}+\check{r}^{(\ell)}}^{(\ell)}(0)$ . The term  $z_\ell^{-m+2}$  in (3.165) is expressed as

$$\begin{aligned} 2^{-i_1(m_2+n_2)+1} p_0 : \quad & \check{F}^{\frac{2}{m}-1} - \left(\frac{2}{m} - 1\right) \left( -\frac{i_0}{2} z_\ell^{m-1} z + \sum_{i+j=m-1} f_{i,j} \left( \underline{z_\ell^{m-1}} - \frac{i}{2} \underline{z_\ell^{m-2} z} \right) \right. \\ & \left. + \frac{i_0(i_0-1)}{8} \underbrace{z_\ell^{m-2} z^2 + \dots} \right) \check{F}^{\frac{2}{m}-2}, \end{aligned} \quad (3.166)$$

where the fact that the first underlined term does not contribute to  $\check{b}_{\check{j}^{(\ell)}+\check{r}^{(\ell)}}^{(\ell)}(0)$  can be proved as in Claim 7, and the second can be proved as in (3.163) (also cf. (3.164)). We see that the coefficient

$$2^{-i_1(m_2+n_2)+1} p_0 \left(1 - \frac{2}{m}\right) \frac{i_0(i_0-1)}{8} \quad (\text{A'.2})$$

contributes to  $\check{b}_{\check{j}^{(\ell)}+\check{r}^{(\ell)}}^{(\ell)}(0)$ .

The term  $z_\ell^{m-1} z \check{F}_m^{\frac{2}{m}-2}$  in (3.166) can be written as (cf. (3.149))

$$2^{-i_1(m_2+n_2)+1} p_0 \left(\frac{2}{m} - 1\right) \frac{i_0}{2} : \check{F}_m^{\frac{2}{m}-2} \left( \check{F}_m^{1-\frac{1}{m}} + \left(1 - \frac{1}{m}\right) \left(\frac{i_0}{2} \underbrace{z_\ell^{m-1} z + \dots}_{\check{F}_m^{\frac{1}{m}}}\right) \check{F}_m^{-\frac{1}{m}} \right) \times \\ \times \left( \check{F}_m^{\frac{1}{m}} + \frac{1}{\ell m} \left(\frac{i_0}{2} \underbrace{z_\ell^{m-1} z + \dots}_{\check{F}_m^{\frac{1}{m}}}\right) \check{F}_m^{\frac{1}{m}-1} \right).$$

Thus the coefficient

$$2^{-i_1(m_2+n_2)+1} p_0 \left(\frac{2}{m} - 1\right) \frac{i_0}{2} \left( \left(1 - \frac{1}{m}\right) \frac{i_0}{2} + \frac{1}{\ell m} \frac{i_0}{2} \right) \quad (\text{A'.3})$$

contributes to  $\check{b}_{\check{j}^{(\ell)}+\check{r}^{(\ell)}}^{(\ell)}(0)$ .

Consider the term  $z_\ell^{-m+1} z$  in (3.165), which is written as

$$2^{-i_1(m_2+n_2)+1} p_0 \frac{1}{2} (i_0 - 1) : \left( \check{F}_m^{\frac{1}{m}-1} + \left(\frac{1}{m} - 1\right) \left(\frac{i_0}{2} \underbrace{z_\ell^{m-1} z + \dots}_{\check{F}_m^{\frac{1}{m}}}\right) \check{F}_m^{\frac{1}{m}-2} \right) \times \\ \times \left( \check{F}_m^{\frac{1}{m}} + \frac{1}{\ell m} \left(\frac{i_0}{2} \underbrace{z_\ell^{m-1} z + \dots}_{\check{F}_m^{\frac{1}{m}}}\right) \check{F}_m^{\frac{1}{m}-1} \right).$$

Thus the coefficient

$$2^{-i_1(m_2+n_2)+1} p_0 \frac{1}{2} (i_0 - 1) \left( \left(\frac{1}{m} - 1\right) \frac{i_0}{2} + \frac{1}{\ell m} \frac{i_0}{2} \right) \quad (\text{A'.4})$$

contributes to  $\check{b}_{\check{j}^{(\ell)}+\check{r}^{(\ell)}}^{(\ell)}(0)$ .

Now taking the sum of (A'.1)–(A'.4) and subtracting the result by (3.160), we again obtain (3.154), thus a contradiction. Lemma 3.11 is again proved.  $\square$

*Subcase 2.2.2: Suppose  $B \neq \emptyset$  and  $j = \min B$  satisfies  $(m+n-1-j)p > 1$ .*

Then  $b_i(x)F(x, y)^{\frac{n-i}{m}}$  does not contribute to  $G(x, y)^{[jp]}$  if  $i \geq m+n-1$ , and the computation of  $G(x, y)^{[jp]}$  shows that  $F_{\text{prim}}(x, y)^{\frac{n-j}{d_2}} = (F(x, y)^{\frac{n-j}{m}})^{[jp]} \equiv 0 \pmod{\mathbb{F}(x, y)}$  (cf. Fact (ii) and Subcase 2.2.1). By Lemma 2.1,  $F_{\text{prim}}(x, y) = H(x, y)^{\frac{d_2}{(d_2, j)}}$  for some polynomial  $H(x, y)$ . This contradicts the definition of primary polynomial in (3.24). Thus this subcase does not occur.

This completes the proof of Theorem 1.1(1).

The above proof in particular shows that any generators  $F(x, y), G(x, y)$  of  $\mathbb{F}[x, y]$  can be mapped to  $x, y$  by a sequence of automorphisms in (1.3). Thus any automorphism of  $\mathbb{F}[x, y]$  must be a product of automorphisms in (1.3). This proves Theorem 1.1(2).

*Proof of Theorem 1.2.* Suppose  $F(x, y)$ , written as in (3.2), is not a monic polynomial of  $y$  (up to a scalar). Take  $F_0(x, y) = \sum_{i+j=m} f_{ij}x^i y^j$ .

Since  $\deg_y F \geq \deg_x F$ , we see that  $F_0(x, y)$  contains  $y$ . Applying the automorphism  $(x, y) \mapsto (x + ay, y)$  for some  $a \neq 0$  so that  $F(x + ay, y)$  becomes a monic polynomial of  $y$  (up to a scalar) with prime degree  $p = 1$ , and  $F_0(x + ay, y)$  becomes the leading polynomial of  $F(x + ay, y)$ . We claim that the primary polynomial cannot have the forms in Lemma 3.10(1)(2). Otherwise  $F_0(x + ay, y)$  would take the form  $(\alpha y + \beta x)^m$  for some  $\alpha, \beta \in \mathbb{F}$ , and then  $F_0(x, y)$  would have the form  $(\alpha' y + \beta' x)^m$  for some  $\alpha', \beta' \in \mathbb{F}$ , i.e.,  $F(x, y)$  would, up to a scalar, be a monic polynomial of  $y$  (since the ignored polynomial of  $F(x, y)$  (cf. Definition 2.5) does not contain  $y^k$  for  $k \geq m$ ). Thus we have Lemma 3.10(3). Now the proof of Lemma 3.11 (cf. the first two statements in the proof of Lemma 3.11) shows that this is impossible.

**Remark 3.13** Note that the general result of Lemma 2.1 is not used in the proof of Theorem 1.1. We only use the simple fact that if  $H(x, y) \in \mathbb{F}[x, y]$  such that  $H(x, y)^{\frac{m}{n}} \in \mathbb{F}(x, y)$ , then  $H(x, y) = H_1(x, y)^{\frac{m}{(m,n)}}$  for some  $H_1(x, y) \in \mathbb{F}[x, y]$ . Also we can avoid using Lemmas 3.2–3.5 and avoid considering Subcase 2.1 by starting directly from the primary polynomial of  $F(x, y)$  in (3.24) and using (3.9) instead of (3.33). Thus the proof can be simplified. However, we prefer to give the above more natural and more easily followed way of the proof which also expresses our starting point on how to give a proof of the theorem.

### ACKNOWLEDGEMENTS

The author would like to thank Professor Zhexian Wan for encouragement. He would also like to thank Professor T.T. Moh for comments and suggestions in previous versions.

### REFERENCES

- [A] A. Abdesselam, The Jacobian conjecture as a problem of perturbative quantum field theory, *Ann. Henri Poincaré* **4** (2003), 199–215.
- [AO] H. Appelgate, H. Onishi, The Jacobian conjecture in two variables, *J. Pure Appl. Algebra* **37** (1985), 215–227.
- [B] H. Bass, The Jacobian conjecture, *Algebra and its Applications* (New Delhi, 1981), 1–8, Lecture Notes in Pure and Appl. Math. **91**, Dekker, New York, 1984.
- [BCW] H. Bass, E.H. Connell, D. Wright, The Jacobian conjecture: reduction of degree and formal expansion of the inverse, *Bull. Amer. Math. Soc.* **7** (1982), 287–330.
- [CCS] Z. Charzyński, J. Chadzyński, P. Skibiński, A contribution to Keller’s Jacobian conjecture. IV. *Bull. Soc. Sci. Lett. Łódź* **39** (1989), no. 11, 6 pp.
- [CM] M. Chamberland, G. Meisters, A mountain pass to the Jacobian conjecture, *Canad. Math. Bull.* **41** (1998), 442–451.
- [D] L.M. Drużkowski, The Jacobian conjecture: survey of some results, *Topics in Complex Analysis* (Warsaw, 1992), 163–171, Banach Center Publ., 31, Polish Acad. Sci., Warsaw, 1995.

- [DV] M. de Bondt, A. van den Essen, A reduction of the Jacobian conjecture to the symmetric case, *Proc. Amer. Math. Soc.* **133** (2005), 2201–2205.
- [ES] G.P. Egorychev, V.A. Stepanenko, The combinatorial identity on the Jacobian conjecture, *Acta Appl. Math.* **85** (2005), 111–120.
- [H] E. Hamann, Algebraic observations on the Jacobian conjecture, *J. Algebra* **265** (2003), 539–561.
- [J] Z. Jelonek, The Jacobian conjecture and the extensions of polynomial embeddings, *Math. Ann.* **294** (1992), 289–293.
- [K] S. Kaliman, On the Jacobian conjecture, *Proc. Amer. Math. Soc.* **117** (1993), 45–51.
- [KM] T. Kambayashi, M. Miyanishi, On two recent views of the Jacobian conjecture, *Affine Algebraic Geometry*, 113–138, Contemp. Math. **369**, Amer. Math. Soc., Providence, RI, 2005.
- [Ki] M. Kirezci, The Jacobian conjecture. I, II. *İstanbul Tek. Üniv. Bül.* **43** (1990), 421–436, 451–457.
- [M1] T.T. Moh, On the Jacobian conjecture and the configurations of roots, *J. Reine Angew. Math.* **340** (1983), 140–212.
- [M2] T.T. Moh, Comment on a paper by Yucai Su on Jacobian conjecture, math.RA/0512495.
- [M3] T.T. Moh, Comment on a paper by Yucai Su on Jacobian conjecture (Dec 30, 2005), math.RA/0604049.
- [N] M. Nagata, Some remarks on the two-dimensional Jacobian conjecture, *Chinese J. Math.* **17** (1989), 1–7.
- [No] A. Nowicki, On the Jacobian conjecture in two variables. *J. Pure Appl. Algebra* **50** (1988), 195–207.
- [R] K. Rusek, A geometric approach to Keller’s Jacobian conjecture, *Math. Ann.* **264** (1983), 315–320.
- [S] S. Smale, Mathematical problems for the next century, *Math. Intelligencer* **20** (1998), 7–15.
- [SW] M.H. Shih, J.W. Wu, On a discrete version of the Jacobian conjecture of dynamical systems, *Nonlinear Anal.* **34** (1998), 779–789.
- [SY] V. Shpilrain, J.T. Yu, Polynomial retracts and the Jacobian conjecture, *Trans. Amer. Math. Soc.* **352** (2000), 477–484.
- [V1] A. van den Essen, *Polynomial automorphisms and the Jacobian conjecture*, Progress in Mathematics **190**, Birkhäuser Verlag, Basel, 2000.
- [V2] A. van den Essen, The sixtieth anniversary of the Jacobian conjecture: a new approach, *Polynomial automorphisms and related topics*, Ann. Polon. Math. **76** (2001), 77–87.
- [W] D. Wright, The Jacobian conjecture: ideal membership questions and recent advances, *Affine Algebraic Geometry*, 261–276, Contemp. Math. **369**, Amer. Math. Soc., Providence, RI, 2005.