

# Rational Combinatorics

Rafael Díaz and Héctor Blandín

February 8, 2020

## Abstract

We propose a categorical setting for the study of the combinatorics of rational numbers. We find combinatorial interpretation for the Bernoulli and Euler numbers and polynomials.

## 1 Introduction

Let  $\mathbf{Cat}$  be the category whose objects are small categories (categories whose collection of objects are sets) and whose morphisms  $\mathbf{Cat}(\mathcal{C}, \mathcal{D})$  from category  $\mathcal{C}$  to category  $\mathcal{D}$  are functors  $F : \mathcal{C} \rightarrow \mathcal{D}$ . Let  $\mathbf{Set}$  be the category whose objects are sets and given  $x, y \in \mathbf{Ob}(\mathbf{Set})$  morphisms in  $\mathbf{Set}$  from  $x$  to  $y$  are maps  $f : x \rightarrow y$ . We define an equivalence relation  $\text{Iso}_{\mathcal{C}}$  on  $\mathbf{Ob}(\mathcal{C})$  as follows: two objects  $x$  and  $y$  are equivalent if and only if there exists an isomorphism  $f \in \mathcal{C}(x, y)$ . There is a natural functor  $D : \mathbf{Cat} \rightarrow \mathbf{Set}$  called *deategorification* given by  $D(\mathcal{C}) = \mathbf{Ob}(\mathcal{C})/\text{Iso}_{\mathcal{C}}$  for all  $\mathcal{C} \in \mathbf{Ob}(\mathbf{Cat})$ , see [3] for details. Given functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  then  $D(F) : \mathbf{Ob}(\mathcal{C})/\text{Iso}_{\mathcal{C}} \rightarrow \mathbf{Ob}(\mathcal{D})/\text{Iso}_{\mathcal{D}}$  is the induced map. If  $x$  is a set and  $\mathcal{C}$  is a category such that  $D(\mathcal{C}) = x$ , we say that  $\mathcal{C}$  is a *categorification* of  $x$ . For example a categorification of the natural numbers is the category  $\mathbb{B}$  whose objects are finite sets and whose morphisms are bijections. We are interested in the categorification of sets with additional properties. For example see [3], a partial categorification of a ring  $R$  is a symmetric monoidal category with negative objects  $\mathcal{C}$  together with a map  $|\cdot| : \mathbf{Ob}(\mathcal{C}) \rightarrow R$ , such that  $|x \oplus y| = |x| + |y|$ ,  $|x \otimes y| = |x||y|$ ,  $|1| = 1$  and  $|0| = 0$ . The main goal of this paper is to introduce a categorification of the ring  $\mathbb{Q}$  of rational numbers.

As explained in [8] the main topic of enumerative combinatorics is the following: given a infinite sequence  $x_0, x_1, \dots, x_n, \dots$  of finite sets, objects of  $\mathbb{B}$ , compute the corresponding sequence  $|x_0|, |x_1|, \dots, |x_n|, \dots$  of natural numbers. So one looks for a numerical representation of combinatorial objects. There is also an inverse problem in enumerative combinatorics: given a sequence of natural numbers  $a_0, a_1, \dots, a_n, \dots$  find an appropriated sequence of finite sets  $x_0, x_1, \dots, x_n, \dots$  such that  $|x_i| = a_i$  for all  $i \in \mathbb{N}$ . Here we look for a combinatorial interpretation of a sequence of natural numbers. Enumerative combinatorics can be extended to deal with integer sequences of numbers by considering sequences of pairs of sets  $(x_0, y_0), \dots, (x_n, y_n), \dots$  and demanding that  $|(x_i, y_i)| := |x_i| - |y_i| = a_i$  for  $a_i \in \mathbb{Z}$  and  $i \in \mathbb{N}$ . Our problem is the following: find a symmetric monoidal category  $\mathcal{C}$ , not too far from  $\mathbb{B}$ , provided with a natural valuation  $|\cdot| : \mathbf{Ob}(\mathcal{C}) \rightarrow \mathbb{Q}$ . Given such pair  $\mathcal{C}$  and  $|\cdot| : \mathbf{Ob}(\mathcal{C}) \rightarrow \mathbb{Q}$  one defines "rational combinatorics" as the problem of finding the sequence  $|x_0|, \dots, |x_n|, \dots$  for any sequence of

objects  $x_0, \dots, x_n, \dots$  in  $\text{Ob}(\mathcal{C})$ . Similarly the inverse problem in rational combinatorics would be the following: given a sequence  $a_0, \dots, a_n, \dots$  in  $\mathbb{Q}$ , find objects  $x_0, \dots, x_n, \dots$  in  $\text{Ob}(\mathcal{C})$  such that  $|x_i| = a_i$ ,  $i \in \mathbb{N}$ . Of course there are several categories  $\mathcal{C}$  provided with valuations  $|\cdot| : \text{Ob}(\mathcal{C}) \rightarrow \mathbb{Q}$ . So if we like our enumerative problem above to be called "combinatorial" the category  $\mathcal{C}$  must be close to  $\mathbb{B}$  and the valuation map  $|\cdot| : \text{Ob}(\mathcal{C}) \rightarrow \mathbb{Q}$  close to the notion of cardinality of finite sets. Our main proposal is that the category FGD of finite groupoids is a promising candidate for replacing  $\mathbb{B}$  when dealing with the combinatorial properties of rational numbers.

## 2 $\mathbb{B}^n$ -FGD species

We set  $[n] = \{1, 2, \dots, n\}$  and denote by  $\mathbb{B}^n$  the category whose objects are pairs  $(x, f)$  where  $x$  is a finite set and  $f : x \rightarrow [n]$  is a map. Morphisms in  $\mathbb{B}^n$  from  $(x, f)$  to  $(y, g)$  are given by

$$\mathbb{B}^n((x, f), (y, g)) = \{\alpha : x \rightarrow y \mid \alpha \text{ is a bijection such that } g \circ \alpha = f\}.$$

$\mathbb{B}^n$  is a symmetric monoidal category with the monoidal structure given by disjoint union

$$(x, f) \sqcup (y, g) = (x \sqcup y, f \sqcup g).$$

**Definition 1.** A groupoid  $G$  is a category such that all its morphisms are invertible. A groupoid  $G$  is called finite if  $\text{Ob}(G)$  is a finite set and  $G(x, y)$  is a finite set for all  $x, y \in \text{Ob}(G)$ .

**Example 2.** The category  $\mathbb{B}^n$  is a groupoid.

**Example 3.** Any group  $G$  may be regarded as a groupoid as follows:  $G$  has only one object 1 and  $G(1, 1) = G$ .

**Definition 4.** The category FGD of finite groupoids is the category whose objects are finite groupoids and whose morphisms from  $G$  to  $H$  are functors  $F : G \rightarrow H$ .

FGD is a symmetric monoidal category whose monoidal structure is given by disjoint union and Cartesian product, the groupoid 1 has only one object  $\text{Ob}(1) = \{1\}$  and are morphism  $1(1, 1) = 1$ . The empty groupoid  $\phi$  is such that  $\text{Ob}(\phi) = \phi$  the empty set. For the reader convenience we recall the notions of disjoint union and Cartesian product of categories. Given two categories  $\mathcal{C}$  and  $\mathcal{D}$  we define the disjoint union  $\mathcal{C} \sqcup \mathcal{D}$  as the category whose objects are  $\text{Ob}(\mathcal{C}) \sqcup \text{Ob}(\mathcal{D})$  and for  $x, y \in \text{Ob}(\mathcal{C} \sqcup \mathcal{D})$  the morphisms from  $x$  to  $y$  are given by

$$\mathcal{C} \sqcup \mathcal{D}(x, y) = \begin{cases} \mathcal{C}(x, y) & \text{if } x, y \in \text{Ob}(\mathcal{C}), \\ \mathcal{D}(x, y) & \text{if } x, y \in \text{Ob}(\mathcal{D}), \\ \phi & \text{otherwise.} \end{cases}$$

The Cartesian product of categories  $\mathcal{C}$  and  $\mathcal{D}$  is the category  $\mathcal{C} \times \mathcal{D}$  given by

$$\text{Ob}(\mathcal{C} \times \mathcal{D}) = \text{Ob}(\mathcal{C}) \times \text{Ob}(\mathcal{D})$$

and for all  $(x_1, y_1), (x_2, y_2) \in \text{Ob}(\mathcal{C} \times \mathcal{D})$  we have  $\mathcal{C} \times \mathcal{D}((x_1, y_1), (x_2, y_2)) = \mathcal{C}(x_1, x_2) \times \mathcal{D}(y_1, y_2)$ .

**Definition 5.** Let  $R$  be a commutative ring. A  $R$ -valuation on a symmetric monoidal category  $\mathcal{C}$  is a map  $|\cdot| : \text{Ob}(\mathcal{C}) \rightarrow R$  such that for  $x, y \in \text{Ob}(\mathcal{C})$  the following identities holds

$$\begin{aligned} (1) \quad |x| &= |y| \text{ if } x \cong y. & (2) \quad |x \oplus y| &= |x| + |y|. & (3) \quad |x \otimes y| &= |x| \cdot |y|. \\ (4) \quad |1_{\mathcal{C}}| &= 1_R. & (5) \quad |0_{\mathcal{C}}| &= 0_R. & (6) \quad |N(x)| &= -|x|. \end{aligned}$$

**Theorem 6.** The map  $|\cdot| : \text{Ob}(\text{FGD}) \rightarrow \mathbb{Q}$  given by  $|G| = \sum_{x \in D(G)} \frac{1}{|G(x, x)|}$  is a surjective  $\mathbb{Q}$ -valuation on FGD.

*Proof.* Let  $G, H \in \text{Ob}(\text{FGD})$ , Then

$$\begin{aligned} |G \sqcup H| &= \sum_{x \in D(G \sqcup H)} \frac{1}{|(G \sqcup H)(x, x)|} = \sum_{x \in D(G)} \frac{1}{|G(x, x)|} + \sum_{x \in D(H)} \frac{1}{|H(x, x)|} \\ &= |G| + |H|. \end{aligned}$$

$$\begin{aligned} |G \times H| &= \sum_{(x, y) \in D(G \times H)} \frac{1}{|(G \times H)(x, y)|} = \sum_{(x, y) \in D(G) \times D(H)} \frac{1}{|G(x, x)| |H(y, y)|} \\ &= \left( \sum_{x \in D(G)} \frac{1}{|G(x, x)|} \right) \left( \sum_{y \in D(H)} \frac{1}{|H(y, y)|} \right) = |G| |H|. \end{aligned}$$

$|1| = 1$  and  $|\phi| = 0$ . For each  $\frac{a}{b} \in \mathbb{Q}^+$  the groupoid  $\bigsqcup_{i=1}^a \mathbb{Z}_b$  is such that  $|\bigsqcup_{i=1}^a \mathbb{Z}_b| = \frac{a}{b}$ .  $\square$

The valuation map  $|\cdot| : \text{FGD} \rightarrow \mathbb{Q}$  above has been previously introduced by Baez and Dolan in [2]. Given two categories  $\mathcal{C}$  and  $\mathcal{D}$  we let  $\mathcal{C}^{\mathcal{D}}$  be the category whose objects are functors  $F : \mathcal{D} \rightarrow \mathcal{C}$  and morphisms from  $F$  to  $G$  in  $\mathcal{C}^{\mathcal{D}}$  are natural transformations  $T : F \rightarrow G$ .

**Definition 7.** The category of  $\mathbb{B}^n$ -FGD species is the category  $\text{FGD}^{\mathbb{B}^n}$ . An object  $F : \mathbb{B}^n \rightarrow \text{FGD}$  is called a  $\mathbb{B}^n$ -FGD species or a rational species.

We denote by  $\text{FGD}_0^{\mathbb{B}^n}$  the full subcategory of  $\text{FGD}^{\mathbb{B}^n}$  whose objects are functors  $F : \mathbb{B}^n \rightarrow \text{FGD}$  such that  $F(\phi) = 0$ . The following definition was extracted from [3]. We denote by  $\Pi[x]$  the set of all partitions of  $x \in \text{Ob}(\mathbb{B})$ .

**Definition 8.** Let  $F, G \in \text{Ob}(\text{FGD}^{\mathbb{B}^n})$ ,  $G_1, \dots, G_n \in \text{Ob}(\text{FGD}_0^{\mathbb{B}^n})$  and  $(x, f) \in \text{Ob}(\mathbb{B}^n)$ , the following defines, product, composition and derivative of  $\mathbb{B}^n$ -FGD species

1.  $(F + G)(x, f) = F(x, f) \sqcup G(x, f)$ .
2.  $(FG)(x, f) = \bigsqcup_{a \sqcup b = x} F(a, f|_a) \times G(b, f|_b)$ .
3.  $(F \times G)(x, f) = F(x, f) \times G(x, f)$ .
4.  $F(G_1, \dots, G_n)(x, f) = \bigsqcup_{\substack{\pi \in \Pi[x] \\ g : \pi \rightarrow [n]}} F(\pi, g) \times \prod_{b \in \pi} G_{g(b)}(b, f|_b)$ .

5.  $\partial_i : \text{FGD}^{\mathbb{B}^n} \rightarrow \text{FGD}^{\mathbb{B}^n}$  where  $\partial_i(F)$  is given by the formula

$$\partial_i F(x, f) = F(x \sqcup \{*\}, f \sqcup \{(*, i)\}).$$

For  $F \in \text{Ob}(\text{FGD}_0^{\mathbb{B}^n})$  we shall write  $F^n = FF \dots F$ , and if  $F \in \text{Ob}(\text{FGD}_0^{\mathbb{B}^n})$  then  $F^{(a)} = F \circ F \circ \dots \circ F$ . For  $a = (a_1, \dots, a_n) \in \mathbb{N}^n$  we shall write  $a! = a_1! \dots a_n!$  and  $x^a = x_1^{a_1} \dots x_n^{a_n}$ .

The class of a object  $(x, f) \in \text{Ob}(\mathbb{B}^n)$  is the tuple  $\text{cl}(x, f) = (|f^{-1}(i)|)_{i=1}^n \in \mathbb{N}^n$ . Notice that by functoriality  $|F(x, f)|$  only depends on the class  $\text{cl}(x, f)$  of  $(x, f) \in \text{Ob}(\mathbb{B}^n)$ .

**Theorem 9.** The map  $| \cdot | : \text{Ob}(\text{FGD}^{\mathbb{B}^n}) \rightarrow \mathbb{Q}[[x_1, \dots, x_n]]$  given by

$$|F|(x_1, \dots, x_n) = \sum_{a \in \mathbb{N}^n} |F(a)| \frac{x^a}{a!}.$$

is a  $\mathbb{Q}[[x_1, \dots, x_n]]$ -valuation on  $\text{FGD}^{\mathbb{B}^n}$ . Moreover  $|F \times G| = |F| \times |G|$ ,  $|\partial_i F| = \partial_i |F|$  and

$$|F(G_1, \dots, G_n)| = |F|(|G_1|, \dots, |G_n|).$$

Above  $\times$  is the Hadamard product of series.

The symmetric group is denoted by  $S_n = \{f : [n] \rightarrow [n] \mid f \text{ is bijective}\}$ . The cyclic group of order  $n$  is  $\mathbb{Z}_n = \{0, \dots, n-1\}$ .

**Example 10.** The singular species  $X_i : \mathbb{B}^n \rightarrow \text{FGD}$  for each  $1 \leq i \leq n$  such that

$$X_i(x, f) = \begin{cases} 1_{\text{FGD}} & \text{if } |x| = 1 \text{ and } f(x) = i, \\ 0 & \text{otherwise.} \end{cases}$$

for  $(x, f) \in \text{Ob}(\mathbb{B}^n)$ .

**Example 11.** Let  $S^{\times N} : \mathbb{B} \rightarrow \text{FGD}$  the species given as follows: for each  $x \in \text{Ob}(\mathbb{B})$ ,  $S^{\times N}(x)$  is the groupoid given by  $\text{Ob}(S^{\times N}(x)) = \{x\}$  for all  $x \in \text{Ob}(\mathbb{B})$  and the morphisms  $S^{\times N}(x)(x, x) = S_{|x|}^N$

$$|S^{\times N}| = \sum_{n=0}^{\infty} \frac{1}{(n!)^N} \frac{x^n}{n!}.$$

**Example 12.** Let  $\mathbb{Z}^{\times N} : \mathbb{B} \rightarrow \text{FGD}$  such that for each  $x \in \text{Ob}(\mathbb{B})$ ,  $\text{Ob}(\mathbb{Z}^{\times N}(x)) = \begin{cases} \{x\} & \text{if } x \neq \phi, \\ \phi & \text{if } x = \phi. \end{cases}$  and  $\mathbb{Z}^{\times N}(x)(x, x) = \mathbb{Z}_{|x|}^N$ , then

$$|\mathbb{Z}^{\times N}| = \sum_{n=1}^{\infty} \frac{1}{n^N} \frac{x^n}{n!}.$$

**Example 13.** Let  $E_N : \mathbb{B} \rightarrow \text{FGD}$  such that for each  $x \in \text{Ob}(\mathbb{B})$ ,  $\text{Ob}(E_N(x)) = \{x\}$ . and

$$E_N(x)(x, x) = \begin{cases} \mathbb{Z}_N^{|x|} & \text{if } x \neq \phi, \\ \{\phi\} & \text{if } x = \phi. \end{cases} \quad \text{then}$$

$$|E_N| = \sum_{n=0}^{\infty} \frac{1}{N^n} \frac{x^n}{n!} = e^{\frac{x}{N}}.$$

Let  $G$  be a group. Let  $\underline{G} : \mathbb{B} \rightarrow \text{FGD}$  the rational species such that for  $x \in \text{Ob}(\mathbb{B})$

$$\text{Ob}(\underline{G}(x)) = \begin{cases} \phi & \text{if } x \neq \phi, \\ \underline{G}(\phi) = G & \text{if } x = \phi. \end{cases}$$

Then we have

$$|\underline{G}| = \frac{1}{|G|}.$$

**Corolary 14.**

$$|\underline{\mathbb{Z}}_N| = \frac{1}{N}.$$

**Example 15.** Let  $(\ )^{(N)} : \mathbb{B} \rightarrow \text{FGD}$  the rational species such that for  $x \in \text{Ob}(\mathbb{B})$ ,

$$\text{Ob}\left((x)^{(N)}\right) = \begin{cases} \{x\} & \text{if } x \neq \phi, \\ \phi & \text{if } x = \phi. \end{cases}$$

and

$$(x)^N(x, x) = \mathbb{Z}_N \times \mathbb{Z}_{N+1} \times \dots \times \mathbb{Z}_{N+|x|-1}.$$

then

$$\left|(\ )^{(N)}\right| = \sum_{n=1}^{\infty} \frac{1}{(N)^{(n)}} \frac{x^n}{n!}.$$

where  $(x)^n = x(x+1) \dots (x+n-1)$  is the increasing factorial polynomial.

**Example 16.** Let  $P : \mathbb{B} \rightarrow \text{FGD}$  given by  $\text{Ob}(P(x)) = \{A \subseteq x \mid A \neq \phi\}$ , and morphisms in  $P(x)$  from  $A$  to  $B$  are given by  $P(x)(A, B) = \{\alpha : A \rightarrow B \mid \alpha \text{ is bijective}\}$ , then

$$|P| = \sum_{n=0}^{\infty} \left( \sum_{k=0}^n \frac{1}{k!} \right) \frac{x^n}{n!} = \sum_{n=0}^{\infty} \left( \sum_{k=0}^n \frac{\binom{n}{k}}{(n)_k} \right) \frac{x^n}{n!}.$$

If  $g$  is such that  $\frac{d^N g(x)}{dx^N} = f(x)$  then we shall write  $g(x) = \int^{(N)} f(x) dx$ .

Let  $\mathbb{Z} : \mathbb{B} \rightarrow \text{FGD}$  the species given in example 12 with  $N = 1$ , and let  $\mathbb{Z}^{(N)} : \mathbb{B} \rightarrow \text{FGD}$  be the increasing factorial rational species. It is such that  $x \in \text{Ob}(\mathbb{B})$  the groupoid  $\mathbb{Z}^{(N)}(x)$  is given by

$$\text{Ob}(\mathbb{Z}^{(N)}(x)) = \begin{cases} \{x\} & x \neq \phi, \\ \phi & \text{if } x = \phi. \end{cases}$$

and morphism  $\mathbb{Z}^{(N)}(x)(x, x) = \mathbb{Z}_{|x|} \times \mathbb{Z}_{|x|+1} \times \dots \times \mathbb{Z}_{|x|+N-1}$ .

**Theorem 17.**

$$\left|\mathbb{Z}^{(N)}\right| = x^{1-N} \int^{(N)} \frac{e^x - 1}{x} dx.$$

*Proof.*

$$\begin{aligned} \left| \mathbb{Z}^{(N)} \right| &= \sum_{n=1}^{\infty} \frac{1}{(n)^{(N)}} \frac{x^n}{n!} = \frac{1}{x^{N-1}} \sum_{n=1}^{\infty} \frac{1}{n} \int^{(N-1)} \frac{x^n}{n!} dx = \frac{1}{x^{N-1}} \int^{(N-1)} \left( \sum_{n=1}^{\infty} \frac{1}{n} \frac{x^n}{n!} \right) dx \\ &= \frac{1}{x^{N-1}} \int^{(N-1)} \left( \int \frac{e^x - 1}{x} dx \right) dx = x^{1-N} \int^{(N)} \frac{e^x - 1}{x} dx. \end{aligned}$$

□

Notice that  $\mathbb{Z}^{(1)} = \mathbb{Z}$  and

**Corolary 18.**

$$|\mathbb{Z}| = \int \frac{e^x - 1}{x} dx.$$

*Proof.*

$$|\mathbb{Z}| = \sum_{n=0}^{\infty} |\mathbb{Z}([n])| \frac{x^n}{n!} = \sum_{n=1}^{\infty} \frac{1}{n} \frac{x^n}{n!} = \int \frac{e^x - 1}{x} dx.$$

□

**Definition 19.** Given a groupoid  $G$  and an integer  $N$ . The increasing factorial groupoid  $G^{(N)}$  is given by

$$G^{(N)} = G \times (G \sqcup [1]) \times \cdots \times (G \sqcup [N-1]).$$

Explicitly

$$\text{Ob}(G^{(N)}) = \bigsqcup_{I \subseteq [n-1]} \left( \prod_{i \in I} G \right) \times \left( \prod_{i \in [n-1]-I} [i] \right).$$

Thus  $\text{Ob}(G^{(N)})$  is the set of all pairs  $(I, f)$  such that  $I \subseteq [n-1]$  and  $f$  is any map such that domain of  $f$  is  $[n-1]$  and  $f(i) \in G$  if  $i \in I$ , and  $f(i) \in [i]$  if  $i \notin I$ . Morphisms in  $G^{(N)}$  are given as follows

$$G^{(N)}((I, f); (J, g)) = \begin{cases} \phi & I \neq J \text{ or } f|_{I^c} \neq g|_{J^c}, \\ \prod_{i \in I} G(f(i), g(i)) & \text{otherwise.} \end{cases}$$

**Proposition 20.**  $|G^{(N)}| = |G|^{(N)}$ , for all finite groupoid  $G$ .

*Proof.*

$$\left| G^{(N)} \right| = \left| \prod_{i=0}^{n-1} (G \sqcup [i]) \right| = \prod_{i=0}^{n-1} |G \sqcup [i]| = \prod_{i=0}^{n-1} (|G| + i) = |G|^{(N)}$$

□

**Definition 21.** Let  $G$  be a finite groupoid such that  $|G| = \frac{a}{b}$ . The species  $\frac{1}{(1+X)^{\frac{a}{b}}} : \mathbb{B} \rightarrow \text{FGD}$  is given on  $x \in \text{Ob}(\mathbb{B})$  by

$$\frac{1}{(1+X)^{\frac{a}{b}}}(x) = (-1)^{|x|} G^{(|x|)}.$$

**Theorem 22.**

$$\left| \frac{1}{(1+X)^{\frac{a}{b}}} \right| = \frac{1}{(1+x)^{\frac{a}{b}}}.$$

*Proof.*

$$\left| \frac{1}{(1+X)^{\frac{a}{b}}} \right| = \sum_{n=0}^{\infty} (-1)^n \left( \frac{a}{b} \right)^{(n)} \frac{x^n}{n!} = \sum_{n=0}^{\infty} \left( -\frac{a}{b} \right)_n \frac{x^n}{n!} = \frac{1}{(1+x)^{\frac{a}{b}}}.$$

□

**Corolary 23.** *Let  $F : \mathbb{B} \rightarrow \text{FGD}$  be a rational species. Then the species given on  $x \in \text{Ob}(\mathbb{B})$  by*

$$\frac{1}{(1+F)^{\frac{a}{b}}}(x) = \bigsqcup_{\pi \in \Pi[x]} (-1)^{|\pi|} |G^{\pi}| \times \prod_{B \in \pi} F(B),$$

*is such that*

$$\left| \frac{1}{(1+F)^{\frac{a}{b}}} \right| = \frac{1}{(1+|F|)^{\frac{a}{b}}}.$$

*Proof.* Use Theorem 9. □

Let  $G$  be a group acting on a finite set  $X$ . The quotient groupoid is the groupoid  $X/G$  such that  $\text{Ob}(X/G) = X$  and  $X/G(x, y) = \{g \in G \mid gx = y\}$ , for all  $x, y \in X$ .

**Proposition 24.** *If  $G$  acts on  $X$ , then  $|X/G| = \frac{|X|}{|G|}$ .*

*Proof.*

$$|X/G| = \sum_{\bar{x} \in O(G)} \frac{1}{|\text{Iso}(x)|} = \sum_{\bar{x} \in O(G)} \frac{|O(x)|}{|G|} = \frac{1}{|G|} \sum_{\bar{x} \in O(G)} |O(x)| = \frac{|X|}{|G|}.$$

□

Let  $G = \{G_N\}$  be a sequence of groups such that  $G_N$  is a subgroup of  $S_N$ .

**Example 25.** *The species  $P_G : \mathbb{B} \rightarrow \text{FGD}$  is given by sending a finite set  $x$  to the quotient groupoid  $x^{[N]}/G$ . It should be clear that  $|P_{S_N}| = \sum_{n=1}^{\infty} \frac{n^N}{N!} \frac{x^n}{n!}$  and  $|P_{\mathbb{Z}_N}| = \sum_{n=1}^{\infty} \frac{n^N}{N} \frac{x^n}{n!}$ .*

We define the rational species  $\text{Isinh} : \mathbb{B} \rightarrow \text{FGD}$  as follows: for each  $x \in \text{Ob}(\mathbb{B})$  the groupoid  $\text{Isinh}(x)$  is given by

$$\begin{aligned} \text{Ob}(\text{Isinh}(x)) &= \begin{cases} \{x\} & \text{if } |x| \text{ is odd,} \\ \phi & \text{otherwise.} \end{cases} \\ \text{Isinh}(x)(x, x) &= \mathbb{Z}_{|x|} \text{ for } x \neq \phi. \end{aligned}$$

Thus we have

**Theorem 26.**

$$|\text{Isinh}| = \int \frac{\sinh(x)}{x} dx.$$

*Proof.*

$$|\text{Isinh}| = \sum_{n=0}^{\infty} \frac{1}{2n+1} \frac{x^{2n+1}}{(2n+1)!} = \int \frac{\sinh(x)}{x} dx.$$

□

The rational species  $\text{Icosh} : \mathbb{B} \rightarrow \text{FGD}$  is given on  $x \in \text{Ob}(\mathbb{B})$  by

$$\text{Ob}(\text{Icosh}(x)) = \begin{cases} \{x\} & \text{if } |x| \text{ is even,} \\ \phi & \text{otherwise.} \end{cases}$$

$$\text{Icosh}(x)(x, x) = \mathbb{Z}_{|x|} \text{ for } x \neq \phi.$$

Then we have the following

**Theorem 27.**

$$|\text{Icosh}| = \int \frac{\cosh(x)}{x} dx.$$

*Proof.*

$$|\text{Icosh}| = \sum_{n=0}^{\infty} \frac{1}{2n} \frac{x^{2n}}{(2n)!} = \int \frac{\cosh(x)}{x} dx.$$

□

The integral sine rational species  $\text{Si} : \mathbb{B} \rightarrow \text{FGD}$  is such that for each  $x \in \text{Ob}(\mathbb{B})$ ,

$$\text{Ob}(\text{Si}(x)) = \begin{cases} \{x\} & \text{if } |x| \text{ is odd,} \\ \phi & \text{otherwise.} \end{cases}$$

$$\text{Si}(x)(x, x) = (-1)^{|x|} \mathbb{Z}_{|x|} \text{ for } x \neq \phi.$$

**Theorem 28.**

$$|\text{Si}| = \int \frac{\sin(x)}{x} dx.$$

*Proof.*

$$|\text{Si}| = \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} \frac{x^{2n+1}}{(2n+1)!} = \int \frac{\sin(x)}{x} dx.$$

□

Consider the inertia functor  $I : \text{FGD}^{\mathbb{B}} \rightarrow \text{FGD}^{\mathbb{B}}$  such that for each specie  $F \in \text{FGD}^{\mathbb{B}}$  its image  $I(F)$  is given by

$$\text{Ob}(I(F)(x)) = \text{Ob}(F(x))$$

for all  $x \in \text{Ob}(\mathbb{B})$  and for objects  $a, b \in \text{Ob}(F(x))$  morphisms from  $a$  to  $b$  in  $I(F)(x)$  are given by

$$I(F)(x)(a, b) = \begin{cases} F(x)(a, a) & \text{if } a = b, \\ \phi & \text{otherwise.} \end{cases}$$



The groupoid  $I(X^{[N]}/S_N)$  is given by  $\text{Ob}(I(X^{[N]}/S_N)) = X^{[N]}$ , and for  $a, b \in X^{[N]}$ ,

$$I(X^{[N]}/S_N)(a, b) = \begin{cases} \{\sigma \in S_N \mid a\sigma = a\} & \text{if } a = b, \\ \phi & a \neq b. \end{cases}$$

One checks that

$$|\{\sigma \in S_N \mid a\sigma = a\}| = \prod_{i \in X} |a^{-1}(i)|!$$

$$|I(P_{S_N})| = \sum_{n=0}^{\infty} \left( \sum_{k_1 + \dots + k_n = N} \binom{N}{k_1, \dots, k_n} \frac{1}{k_1! \dots k_n!} \right) \frac{x^n}{n!}.$$

$$\left| I(X^{[N]}/S_N) \right| = \sum_{a: [N] \rightarrow X} \frac{1}{\prod_{i \in X} |a^{-1}(i)|!} = \sum_{k_1 + \dots + k_n = N} \binom{N}{k_1, \dots, k_n} \frac{1}{k_1! \dots k_n!}.$$

Recall that a set  $X$  can be consider as a groupoid as follows  $\text{Ob}(X) = X$  and morphisms in  $X$  from  $a$  to  $b$  are given by

$$X(a, b) = \begin{cases} 1 & \text{if } a = b, \\ \phi & \text{otherwise.} \end{cases}$$

**Example 29.** The specie  $1 \in \text{FGD}^{\mathbb{B}^n}$  is such that for each  $(x, f) \in \text{Ob}(\mathbb{B}^n)$

$$1(x, f) = \begin{cases} 1_{\text{FGD}} & \text{if } x = \phi, \\ \phi & \text{otherwise.} \end{cases}$$

**Example 30.** The exponential species  $\text{Exp}_n \in \text{FGD}^{\mathbb{B}^n}$  is given by  $\text{Exp}_n(x, f) = 1_{\text{FGD}}$ . It should be clear that  $|\text{Exp}_n| = e^{x_1 + \dots + x_n}$ . For  $n = 1$  we write  $\text{Exp}$  instead of  $\text{Exp}_1$ .

**Theorem 31.** Let  $F$  a  $\mathbb{B}^n$ -FGD species such that  $F(\phi) = \phi$  then the species  $(1 + F)^{-1}$  is given by

$$(1 + F)^{-1}(x, f) = \bigsqcup_{x_1 \sqcup \dots \sqcup x_n = x} (-1)^n F(x_1, f|_{x_1}) \times \dots \times F(x_n, f|_{x_n}).$$

*Proof.*

$$(1 + F)^{-1}(x, f) = \left( \sum_{n=0}^{\infty} (-1)^n F^n \right)(x, f) = \bigsqcup_{x_1 \sqcup \dots \sqcup x_n = x} (-1)^n F(x_1, f|_{x_1}) \times \dots \times F(x_n, f|_{x_n}).$$

□

**Definition 32.** Let  $F \in \text{Ob}(\text{FGD}^{\mathbb{B}^n})$  and  $a, b \in \mathbb{N}$ ,  $a \neq 0$ . The species  $\frac{b}{a}F$  is given as follows  $\frac{b}{a}F = \left( \bigsqcup_{i=1}^b Z_a \right) F$ .

**Theorem 33.** Let  $F \in \text{Ob}(\text{FGD}^{\mathbb{B}^n})$  be such that  $F(\phi) = \phi$ . Let  $G$  be the rational species  $\frac{a}{b} \frac{1}{(1 + \frac{b}{a}F)}$ , then

$$|G| = \frac{1}{\frac{a}{b} + |F|}.$$

*Proof.*

$$|G| = \left| \frac{a}{b} \frac{1}{(1 + \frac{b}{a}F)} \right| = \frac{a}{b} \left( \frac{1}{1 + \frac{b}{a}|F|} \right) = \frac{1}{\frac{a}{b} + |F|}.$$

□

### 3 Bernoulli numbers and polynomials

In this section Theorem 31 will be used to provide a combinatorial interpretation for Bernoulli numbers and polynomials.

**Definition 34.** Let  $N \in \mathbb{N}^+$ , the  $N$ -projection map  $\pi_N : \mathbb{Q}[[x]] \rightarrow \mathbb{Q}[[x]]/(x^N)$  is given by  $\pi_N \left( \sum_{n=0}^{\infty} f_n \frac{x^n}{n!} \right) = \sum_{n=0}^{N-1} f_n \frac{x^n}{n!}$ , for all  $f \in \mathbb{Q}[[x]]$ .

**Definition 35.** Let  $f \in \mathbb{Q}[[x]]$  and  $N \in \mathbb{N}^+$ . The sequence  $\{B_{N,n}^f\}_{n=0}^{\infty}$  is called the  $(f, N)$ -Bernoulli numbers sequence if it is such that

$$\frac{x^N/N!}{f(x) - \pi_N(f)(x)} = \sum_{n=0}^{\infty} B_{N,n}^f \frac{x^n}{n!}.$$

If  $N = 1$  and  $f(x) = e^x$ , then the sequence  $B_{1,n}^{e^x}$  is the Bernoulli numbers sequence  $B_n$  such that

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!}.$$

If  $N = 2$  and  $f(x) = e^x$  we have the  $B_{2,n}$  Bernoulli numbers, they satisfy

$$\frac{x^2/2!}{e^x - 1 - x} = \sum_{n=0}^{\infty} B_{2,n} \frac{x^n}{n!}.$$

**Corollary 36.**

$$\left| \frac{1}{\partial \mathbb{Z}} \right| = \sum_{n=1}^{\infty} B_n \frac{x^n}{n!}.$$

Notice that by Theorem 9 the valuation of the species  $\partial \mathbb{Z} = 1 + \overline{\partial \mathbb{Z}}$ , where  $\overline{\partial \mathbb{Z}}(x) = \partial \mathbb{Z}(x)$  if  $x \neq \phi$  and  $\overline{\partial \mathbb{Z}}(\phi) = \phi$  is  $\frac{e^x - 1}{x}$  and also

$$\begin{aligned} \frac{1}{\partial \mathbb{Z}}(x) &= \bigsqcup_{x_1 \sqcup \dots \sqcup x_k = x} (-1)^k \overline{\partial \mathbb{Z}}(x_1) \times \dots \times \overline{\partial \mathbb{Z}}(x_k) \\ &= \bigsqcup_{x_1 \sqcup \dots \sqcup x_k = x} (-1)^k \mathbb{Z}(x_1 \sqcup *_1) \times \dots \times \mathbb{Z}(x_k \sqcup *_k). \end{aligned}$$

This implies that

$$\frac{1}{\partial \mathbb{Z}}(x) = \bigsqcup_{x_1 \sqcup \dots \sqcup x_k = x} (-1)^k \mathbb{Z}_{|x_1|+1} \times \dots \times \mathbb{Z}_{|x_k|+1}.$$

That is

$$\text{Ob} \left( \frac{1}{\partial \mathbb{Z}}(x) \right) = (-1)^k \{(x_1, \dots, x_k) \mid x_1 \sqcup \dots \sqcup x_k = x\},$$

$$\frac{1}{\partial \mathbb{Z}}(x)((x_1, \dots, x_k), (y_1, \dots, y_k)) = \begin{cases} \mathbb{Z}_{|x_1|+1} \times \dots \times \mathbb{Z}_{|x_k|+1} & \text{if } x_i = y_i \text{ for all } i \in [k], \\ \phi & \text{otherwise.} \end{cases}$$

**Corolary 37.**

$$B_n = \sum_{a_1 + \dots + a_k = n} (-1)^k \frac{n!}{(a_1 + 1)! \dots (a_k + 1)!}.$$

*Proof.* By corollary 36 we have

$$B_n = \sum_{a_1 + \dots + a_k = n} (-1)^k \frac{1}{(a_1 + 1) \dots (a_k + 1)} \binom{n}{a_1, \dots, a_k}.$$

□

Let  $\mathbb{Z}_{(N)} : \mathbb{B} \rightarrow \text{FGD}$  be the decreasing factorial rational species which is such that for each  $x \in \text{Ob}(\mathbb{B})$ ,  $\text{Ob}(\mathbb{Z}_{(N)}(x)) = \text{Ob}(\mathbb{Z}(x))$  and

$$\mathbb{Z}_{(N)}(x, x) = \begin{cases} \mathbb{Z}_{|x|} \times \mathbb{Z}_{|x|-1} \times \dots \times \mathbb{Z}_{|x|-N+1} & \text{if } |x| \geq N, \\ \phi & \text{otherwise.} \end{cases}$$

**Theorem 38.**

$$|\mathbb{Z}_{(N)}| = \int^{(N)} \frac{e^x - \pi_N(e^x)}{x^N} dx.$$

*Proof.* It is easy to verify that

$$\partial^N |\mathbb{Z}_{(N)}| = \sum_{n=N}^{\infty} \frac{x^{n-N}}{n!} = \frac{e^x - \pi_N(e^x)}{x^N}.$$

This implies that

$$|\mathbb{Z}_{(N)}| = \sum_{n=N}^{\infty} \frac{1}{(n)_N} \frac{x^n}{n!} = \int^{(N)} \frac{e^x - \pi_N(e^x)}{x^N} dx.$$

□

More generally we have the following

**Theorem 39.** *Let  $F : \mathbb{B} \rightarrow \text{FGD}$  be a rational species, then*

$$|\partial^N (F \times \mathbb{Z}_{(N)})| = \frac{|F|(x) - \pi_N(|F|)(x)}{x^N}.$$

**Corolary 40.** *Let  $F : \mathbb{B} \rightarrow \text{FGD}$  be a rational species. The valuation of the species  $\frac{1}{F \times (\mathbb{Z})_N}$  is the generating function of the  $(|F|, N)$  Bernoulli numbers. That is*

$$\left| \frac{1}{\partial^N (F \times \mathbb{Z}_{(N)})} \right| = N! \sum_{n=0}^{\infty} B_{N,n}^{|F|} \frac{x^n}{n!}$$

**Definition 41.** *Let  $f \in \mathbb{Q}[[x]]$  and  $N \in \mathbb{N}^+$ . The sequence of functions  $B_{N,n}^f(x)$  are called the  $(f, N)$  Bernoulli polynomials sequence and are given by*

$$\sum_{n=0}^{\infty} B_{N,n}^f(x) \frac{z^n}{n!} = \frac{(z^N/N!) f(zx)}{f(z) - \pi_N(f)(z)}.$$

Notice that if  $N = 1$  and  $f(z) = e^z$  the Bernoulli polynomials  $B_n(x)$  are given by

$$\sum_{n=0}^{\infty} B_n(x) \frac{z^n}{n!} = \frac{ze^{zx}}{e^z - 1}.$$

If  $N = 2$  and  $f(z) = e^z$  the  $B_{2,n}$  polynomials are given by

$$\sum_{n=0}^{\infty} B_{2,n}(x) \frac{z^n}{n!} = \frac{(z^2/2!) e^{zx}}{e^z - 1 - z}.$$

Let  $A$  and  $B$  be finite sets, we use the following notation

$$\text{Bij}(A, B) = \{f : A \rightarrow B \mid f \text{ is bijective}\}.$$

Let  $\text{XZ} : \mathbb{B}^2 \rightarrow \mathbb{B}$  be the species such that for each pair  $(x, y) \in \text{Ob}(\mathbb{B}^2)$

$$\text{XZ}(x, y) = \begin{cases} 1 & \text{if } |x| = |y| = 1, \\ \phi & \text{otherwise.} \end{cases}$$

The valuation  $|\text{XZ}| \in \mathbb{Q}[x, z]$  is given by  $|\text{XZ}| = xz$ .

**Proposition 42.** *Let  $F \in \text{FGD}^{\mathbb{B}}$ . Then  $F(\text{XZ})$  is such that  $F(\text{XZ})(a, b) = F(a)\text{Bij}(a, b)$  and*

$$|F(\text{XZ})| = |F|(xz).$$

*Proof.*

$$F(\text{XZ})(a, b) = \bigsqcup_{\pi \in \prod [x]} F(\pi) \times \prod_{B \in \pi} \text{XZ}(B) = \bigsqcup_{\substack{f: a \rightarrow b \\ \text{bijective}}} F(a) = F(a) \times \text{Bij}(a, b).$$

The identity  $|F(\text{XZ})| = |F|(xz)$  follows from Theorem 9. □

**Corolary 43.**

$$\left| \frac{F(\text{XZ})}{\partial^N (F \times \mathbb{Z}_{(N)})} \right| = \frac{z^N |F|(zx)}{|F|(z) - \pi_N(|F|)(z)}.$$

## 4 Euler numbers and polynomials

In this section we provide a combinatorial interpretation for the Euler numbers and polynomials. It would be interesting to extend our results to the  $q$ -Euler numbers and polynomials discussed for example in [6].

The Euler numbers are denoted by  $E_n$ , so true satisfy

$$\frac{2}{e^x + 1} = \sum_{n=0}^{\infty} E_n \frac{x^n}{n!}.$$

Let  $\mathbb{F}_K : \mathbb{B} \rightarrow \text{FGD}$  the rational species given by  $\mathbb{F}_K = \overline{\mathbb{Z}_K} (1 + \text{Exp})$

**Theorem 44.**

$$|\mathbb{F}_K| = \frac{e^x + 1}{K}.$$

**Theorem 45.** *The species  $\frac{1}{\mathbb{F}_K}$  is such that  $\left| \frac{1}{\mathbb{F}_K} \right| = \frac{K}{e^x + 1}$ .*

Next we provide a combinatorial interpretation for the Euler polynomials. First let us recall the definition of the Euler polynomials also.

**Definition 46.** *The  $K$ -Euler polynomials sequence is given by*

$$\sum_{n=0}^{\infty} E_{n,K}(x) \frac{z^n}{n!} = \frac{K e^{xz}}{e^z + 1}.$$

**Theorem 47.**

$$\left| \frac{\text{Exp}(XZ)}{\mathbb{Z}_K(1 + \text{Exp})} \right| = \frac{K e^{xz}}{e^z + 1}.$$

## Acknowledgment

Our interest in the combinatorics of Bernoulli numbers arouse from a series of lectures on *Heap of Pieces* taught by Xavier Viennot in Caracas 2002.

## References

- [1] Bergeron, Labelle, and Leroux. *Combinatorial Species and Tree-Like Structures*. Encyclopedia of Mathematics and its applications. Editorial Board, Cambridge, 1994.
- [2] John C. Baez, James Dolan. *From finite Sets to Feynman Diagrams*. arXiv:math.QA/0004133.
- [3] Rafael Díaz and Eddy Pariguan. *Super, quantum and noncommutative species*. arXiv:math.CT/0509674.
- [4] Gian-Carlo Rota. *Gian-Carlo Rota on Combinatorics: Introductory papers and commentaries*. Joseph P. S. Kung, editor. Birkhaussär, 1995.
- [5] Hassen, Abdul and Hieu D. Nguyen, *Hypergeometric Zeta Functions*. arXiv:math.NT/0509637.
- [6] Kim, Taeyun. *q-Euler and Genocchi numbers*. Journal of Mathematical Analysis and Applications. In press.
- [7] S. Mac Lane, *Categories for the Working Mathematician*, vol 5, Graduate Texts in Mathematics, Springer Velarg, New York, NY, 1971.

- [8] Zeilberger, D., *Enumerative and Algebraic Combinatorics*. Princeton Companion to Mathematics.

Rafael Díaz. Universidad Central de Venezuela (UCV). `rdiaz@euler.ciens.ucv.ve`  
Héctor Blandín. Universidad Central de Venezuela (UCV). `hblandin@euler.ciens.ucv.ve`