## ON A NONHOMOGENEOUS QUASILINEAR EIGENVALUE PROBLEM IN SOBOLEV SPACES WITH VARIABLE EXPONENT<sup>\*</sup>

Mihai Mihăilescu and Vicențiu Rădulescu

Department of Mathematics, University of Craiova, 200585 Craiova, Romania

E-mail addresses: mmihailes@yahoo.com vicentiu.radulescu@math.cnrs.fr

ABSTRACT. We consider the nonlinear eigenvalue problem  $-\operatorname{div}\left(|\nabla u|^{p(x)-2}\nabla u\right) = \lambda |u|^{q(x)-2}u$  in  $\Omega$ , u = 0 on  $\partial\Omega$ , where  $\Omega$  is a bounded open set in  $\mathbb{R}^N$  with smooth boundary and p, q are continuous functions on  $\overline{\Omega}$  such that  $1 < \inf_{\Omega} q < \inf_{\Omega} p < \sup_{\Omega} q$ ,  $\sup_{\Omega} p < N$ , and q(x) < Np(x)/(N-p(x)) for all  $x \in \overline{\Omega}$ . The main result of this paper establishes that any  $\lambda > 0$  sufficiently small is an eigenvalue of the above nonhomogeneous quasilinear problem. The proof relies on simple variational arguments based on Ekeland's variational principle.

2000 Mathematics Subject Classification: 35D05, 35J60, 35J70, 58E05, 68T40, 76A02.

Key words: p(x)-Laplace operator, nonlinear eigenvalue problem, Sobolev space with variable exponent, Ekeland's variational principle.

## **1** Introduction and preliminary results

A basic result in the elementary theory of linear partial differential equations asserts that the spectrum of the Laplace operator in  $H_0^1(\Omega)$  is discrete, where  $\Omega$  is a bounded open set in  $\mathbb{R}^N$  with smooth boundary. More precisely, the problem

$$\begin{cases} -\Delta u = \lambda u & \text{in } \Omega \\ u = 0 & \text{on } \partial \Omega \end{cases}$$

has an unbounded sequence of eigenvalues  $0 < \lambda_1 < \lambda_2 \leq \ldots \leq \lambda_n \leq \ldots$  This celebrated result goes back to the Riesz-Fredholm theory of self-adjoint and compact operators on Hilbert spaces. The anisotropic case

$$\begin{cases} -\Delta u = \lambda a(x)u & \text{in } \Omega\\ u = 0 & \text{on } \partial\Omega \end{cases}$$

was considered by Bocher [3], Hess and Kato [12], Minakshisundaram and Pleijel [15, 17]. For instance, Minakshisundaram and Pleijel proved that the above eigenvalue problem has an unbounded sequence

<sup>\*</sup>Correspondence address: Vicențiu Rădulescu, Department of Mathematics, University of Craiova, 200585 Craiova, Romania. E-mail: vicentiu.radulescu@math.cnrs.fr

of positive eigenvalues if  $a \in L^{\infty}(\Omega)$ ,  $a \ge 0$  in  $\Omega$ , and a > 0 in  $\Omega_0 \subset \Omega$ , where  $|\Omega_0| > 0$ . Eigenvalue problems for homogeneous quasilinear problems have been intensively studied in the last decades (see, e.g., Anane [2]).

This paper is motivated by recent advances in elastic mechanics and electrorheological fluids (sometimes referred to as "smart fluids"), where some processes are modeled by nonhomogeneous quasilinear operators (see Diening [4], Halsey [11], Ruzicka [18], Zhikov [21], and the references therein). We refer mainly to the p(x)-Laplace operator  $\Delta_{p(x)}u := \operatorname{div}(|\nabla u|^{p(x)-2}\nabla u)$ , where p is a continuous non-constant function. This differential operator is a natural generalization of the p-Laplace operator  $\Delta_p u := \operatorname{div}(|\nabla u|^{p-2}\nabla u)$ , where p > 1 is a real constant. However, the p(x)-Laplace operator possesses more complicated nonlinearities than the p-Laplace operator, due to the fact that  $\Delta_{p(x)}$  is not homogeneous.

In this paper we are concerned with the nonhomogeneous eigenvalue problem

$$\begin{cases} -\operatorname{div}(|\nabla u|^{p(x)-2}\nabla u) = \lambda |u|^{q(x)-2}u, & \text{for } x \in \Omega\\ u = 0, & \text{for } x \in \partial\Omega, \end{cases}$$
(1)

where  $\Omega \subset \mathbb{R}^N$   $(N \ge 3)$  is a bounded domain with smooth boundary,  $\lambda > 0$  is a real number, and p, q are continuous on  $\overline{\Omega}$ .

The case p(x) = q(x) was considered by Fan, Zhang and Zhao in [10] who, using the Ljusternik-Schnirelmann critical point theory, established the existence of a sequence of eigenvalues. Denoting by  $\Lambda$ the set of all nonnegative eigenvalues, Fan, Zhang and Zhao showed that  $\sup \Lambda = +\infty$  and they pointed out that only under additional assumptions we have  $\inf \Lambda > 0$ . We remark that for the *p*-Laplace operator (corresponding to  $p(x) \equiv p$ ) we always have  $\inf \Lambda > 0$ .

In this paper we study problem (1) under the basic assumption

$$1 < \min_{x \in \overline{\Omega}} q(x) < \min_{x \in \overline{\Omega}} p(x) < \max_{x \in \overline{\Omega}} q(x).$$
(2)

Our main result establishes the existence of a continuous family of eigenvalues for problem (1) in a neighborhood of the origin. More precisely, we show that there exists  $\lambda^* > 0$  such that any  $\lambda \in (0, \lambda^*)$  is an eigenvalue for problem (1).

We start with some preliminary basic results on the theory of Lebesgue–Sobolev spaces with variable exponent. For more details we refer to the book by Musielak [16] and the papers by Edmunds et al. [5, 6, 7], Kovacik and Rákosník [13], Mihăilescu and Rădulescu [14], and Samko and Vakulov [19].

Assume that  $p \in C(\overline{\Omega})$  and p(x) > 1, for all  $x \in \overline{\Omega}$ . Set

$$C_{+}(\overline{\Omega}) = \{h; h \in C(\overline{\Omega}), h(x) > 1 \text{ for all } x \in \overline{\Omega} \}.$$

For any  $h \in C_+(\overline{\Omega})$  we define

$$h^+ = \sup_{x \in \Omega} h(x)$$
 and  $h^- = \inf_{x \in \Omega} h(x)$ 

For any  $p(x) \in C_+(\overline{\Omega})$ , we define the variable exponent Lebesgue space

$$L^{p(x)}(\Omega) = \{u; u \text{ is a measurable real-valued function such that } \int_{\Omega} |u(x)|^{p(x)} dx < \infty\}.$$

We define a norm, the so-called Luxemburg norm, on this space by the formula

$$|u|_{p(x)} = \inf\left\{\mu > 0; \ \int_{\Omega} \left|\frac{u(x)}{\mu}\right|^{p(x)} \ dx \le 1\right\}.$$

We remember that the variable exponent Lebesgue spaces are separable and reflexive Banach spaces. If  $0 < |\Omega| < \infty$  and  $p_1$ ,  $p_2$  are variable exponent so that  $p_1(x) \le p_2(x)$  almost everywhere in  $\Omega$  then there exists the continuous embedding  $L^{p_2(x)}(\Omega) \hookrightarrow L^{p_1(x)}(\Omega)$ .

We denote by  $L^{p'(x)}(\Omega)$  the conjugate space of  $L^{p(x)}(\Omega)$ , where 1/p(x) + 1/p'(x) = 1. For any  $u \in L^{p(x)}(\Omega)$  and  $v \in L^{p'(x)}(\Omega)$  the Hölder type inequality

$$\left| \int_{\Omega} uv \, dx \right| \le \left( \frac{1}{p^{-}} + \frac{1}{p'^{-}} \right) |u|_{p(x)} |v|_{p'(x)} \tag{3}$$

holds true.

An important role in manipulating the generalized Lebesgue-Sobolev spaces is played by the *modular* of the  $L^{p(x)}(\Omega)$  space, which is the mapping  $\rho_{p(x)} : L^{p(x)}(\Omega) \to \mathbb{R}$  defined by

$$\rho_{p(x)}(u) = \int_{\Omega} |u|^{p(x)} dx$$

If  $(u_n), u \in L^{p(x)}(\Omega)$  then the following relations hold true

$$|u|_{p(x)} > 1 \quad \Rightarrow \quad |u|_{p(x)}^{p^{-}} \le \rho_{p(x)}(u) \le |u|_{p(x)}^{p^{+}}$$

$$\tag{4}$$

$$|u|_{p(x)} < 1 \quad \Rightarrow \quad |u|_{p(x)}^{p^+} \le \rho_{p(x)}(u) \le |u|_{p(x)}^{p^-}$$
(5)

$$|u_n - u|_{p(x)} \to 0 \quad \Leftrightarrow \quad \rho_{p(x)}(u_n - u) \to 0.$$
 (6)

Next, we define  $W_0^{1,p(x)}(\Omega)$  as the closure of  $C_0^{\infty}(\Omega)$  under the norm

$$\|u\| = |\nabla u|_{p(x)}.$$

The space  $(W_0^{1,p(x)}(\Omega), \|\cdot\|)$  is a separable and reflexive Banach space. We note that if  $s(x) \in C_+(\overline{\Omega})$ and  $s(x) < p^*(x)$  for all  $x \in \overline{\Omega}$  then the embedding  $W_0^{1,p(x)}(\Omega) \hookrightarrow L^{s(x)}(\Omega)$  is compact and continuous, where  $p^*(x) = \frac{Np(x)}{N-p(x)}$  if p(x) < N or  $p^*(x) = +\infty$  if  $p(x) \ge N$ .

## 2 The main result

We say that  $\lambda \in \mathbb{R}$  is an eigenvalue of problem (1) if there exists  $u \in W_0^{1,p(x)}(\Omega) \setminus \{0\}$  such that

$$\int_{\Omega} |\nabla u|^{p(x)-2} \nabla u \nabla v \, dx - \lambda \int_{\Omega} |u|^{q(x)-2} uv \, dx = 0,$$

for all  $v \in W_0^{1,p(x)}(\Omega)$ . We point out that if  $\lambda$  is an eigenvalue of the problem (1) then the corresponding  $u \in W_0^{1,p(x)}(\Omega) \setminus \{0\}$  is a weak solution of (1).

Our main result is given by the following theorem.

**Theorem 1.** Assume that condition (2) is fulfilled,  $\max_{x\in\overline{\Omega}} p(x) < N$  and  $q(x) < p^*(x)$  for all  $x\in\overline{\Omega}$ . Then there exists  $\lambda^* > 0$  such that any  $\lambda \in (0, \lambda^*)$  is an eigenvalue for problem (1).

The above result implies

$$\inf_{u \in W_0^{1,p(x)}(\Omega) \setminus \{0\}} \frac{\int_{\Omega} |\nabla u|^{p(x)} dx}{\int_{\Omega} |u|^{q(x)} dx} = 0$$

Thus, for any positive constant C, there exists  $u_0 \in W_0^{1,p(x)}(\Omega)$  such that

$$C \int_{\Omega} |u_0|^{q(x)} dx \ge \int_{\Omega} |\nabla u_0|^{p(x)} dx$$

Let *E* denote the generalized Sobolev space  $W_0^{1,p(x)}(\Omega)$ .

For any  $\lambda > 0$  the energy functional corresponding to problem (1) is defined as  $J_{\lambda} : E \to \mathbb{R}$ ,

$$J_{\lambda}(u) = \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx - \lambda \int_{\Omega} \frac{1}{q(x)} |u|^{q(x)} dx.$$

Standard arguments imply that  $J_{\lambda} \in C^{1}(E, \mathbb{R})$  and

$$\langle J_{\lambda}'(u), v \rangle = \int_{\Omega} |\nabla u|^{p(x)-2} \nabla u \nabla v \, dx - \lambda \int_{\Omega} |u|^{q(x)-2} uv \, dx$$

for all  $u, v \in E$ . Thus the weak solutions of (1) coincide with the critical points of  $J_{\lambda}$ . If such an weak solution exists and is nontrivial then the corresponding  $\lambda$  is an eigenvalue of problem (1).

**Lemma 1.** There exists  $\lambda^* > 0$  such that for any  $\lambda \in (0, \lambda^*)$  there exist  $\rho$ , a > 0 such that  $J_{\lambda}(u) \ge a > 0$  for any  $u \in E$  with  $||u|| = \rho$ .

*Proof.* Since  $q(x) < p^{\star}(x)$  for all  $x \in \overline{\Omega}$  it follows that E is continuously embedded in  $L^{q(x)}(\Omega)$ . So, there exists a positive constant  $c_1$  such that

$$|u|_{q(x)} \le c_1 ||u||, \quad \forall \ u \in E.$$

$$\tag{7}$$

We fix  $\rho \in (0, 1)$  such that  $\rho < 1/c_1$ . Then relation (7) implies

$$|u|_{q(x)} < 1, \quad \forall \ u \in E, \text{ with } ||u|| = \rho.$$

Furthermore, relation (5) yields

$$\int_{\Omega} |u|^{q(x)} dx \le |u|^{q^{-}}_{q(x)}, \quad \forall \ u \in E, \text{ with } \|u\| = \rho.$$
(8)

Relations (7) and (8) imply

$$\int_{\Omega} |u|^{q(x)} dx \le c_1^{q^-} ||u||^{q^-}, \quad \forall \ u \in E, \text{ with } ||u|| = \rho.$$
(9)

Taking into account relations (5) and (9) we deduce that for any  $u \in E$  with  $||u|| = \rho$  the following inequalities hold true

$$J_{\lambda}(u) \geq \frac{1}{p^{+}} \int_{\Omega} |\nabla u|^{p(x)} dx - \frac{\lambda}{q^{-}} \int_{\Omega} |u|^{q(x)} dx$$
  
$$\geq \frac{1}{p^{+}} ||u||^{p^{+}} - \frac{\lambda}{q^{-}} c_{1}^{q^{-}} ||u||^{q^{-}}$$
  
$$= \frac{1}{p^{+}} \rho^{p^{+}} - \frac{\lambda}{q^{-}} c_{1}^{q^{-}} \rho^{q^{-}}$$
  
$$= \rho^{q^{-}} \left( \frac{1}{p^{+}} \rho^{p^{+} - q^{-}} - \frac{\lambda}{q^{-}} c_{1}^{q^{-}} \right).$$

By the above inequality we remark that if we define

$$\lambda^{\star} = \frac{\rho^{p^{+}-q^{-}}}{2p^{+}} \cdot \frac{q^{-}}{c_{1}^{q^{-}}} \tag{10}$$

then for any  $\lambda \in (0, \lambda^*)$  and any  $u \in E$  with  $||u|| = \rho$  there exists  $a = \frac{\rho^{p^+}}{2p^+} > 0$  such that

$$J_{\lambda}(u) \ge a > 0.$$

The proof of Lemma 1 is complete.

**Lemma 2.** There exists  $\varphi \in E$  such that  $\varphi \geq 0$ ,  $\varphi \neq 0$  and  $J_{\lambda}(t\varphi) < 0$ , for t > 0 small enough.

Proof. Assumption (2) implies that  $q^- < p^-$ . Let  $\epsilon_0 > 0$  be such that  $q^- + \epsilon_0 < p^-$ . On the other hand, since  $q \in C(\overline{\Omega})$  it follows that there exists an open set  $\Omega_0 \subset \Omega$  such that  $|q(x) - q^-| < \epsilon_0$  for all  $x \in \Omega_0$ . Thus, we conclude that  $q(x) \leq q^- + \epsilon_0 < p^-$  for all  $x \in \Omega_0$ .

Let  $\varphi \in C_0^{\infty}(\Omega)$  be such that  $\operatorname{supp}(\varphi) \supset \overline{\Omega}_0$ ,  $\varphi(x) = 1$  for all  $x \in \overline{\Omega}_0$  and  $0 \le \varphi \le 1$  in  $\Omega$ . Then

using the above information for any  $t \in (0, 1)$  we have

$$J_{\lambda}(t\varphi) = \int_{\Omega} \frac{t^{p(x)}}{p(x)} |\nabla\varphi|^{p(x)} dx - \lambda \int_{\Omega} \frac{t^{q(x)}}{q(x)} |\varphi|^{q(x)} dx$$
  
$$\leq \frac{t^{p^{-}}}{p^{-}} \int_{\Omega} |\nabla\varphi|^{p(x)} dx - \frac{\lambda}{q^{+}} \int_{\Omega} t^{q(x)} |\varphi|^{q(x)} dx$$
  
$$\leq \frac{t^{p^{-}}}{p^{-}} \int_{\Omega} |\nabla\varphi|^{p(x)} dx - \frac{\lambda}{q^{+}} \int_{\Omega_{0}} t^{q(x)} |\varphi|^{q(x)} dx$$
  
$$\leq \frac{t^{p^{-}}}{p^{-}} \int_{\Omega} |\nabla\varphi|^{p(x)} dx - \frac{\lambda \cdot t^{q^{-} + \epsilon_{0}}}{q^{+}} \int_{\Omega_{0}} |\varphi|^{q(x)} dx$$

Therefore

 $J_{\lambda}(t\varphi) < 0$ 

for  $t < \delta^{1/(p^- - q^- - \epsilon_0)}$  with

$$0 < \delta < \min\left\{1, \frac{\frac{\lambda \cdot p^{-}}{q^{+}} \int_{\Omega_{0}} |\varphi|^{q(x)} dx}{\int_{\Omega} |\nabla \varphi|^{p(x)} dx}\right\}.$$

Finally, we point out that  $\int_{\Omega} |\nabla \varphi|^{p(x)} dx > 0$ . Indeed, it is clear that

$$\int_{\Omega_0} |\varphi|^{q(x)} dx \le \int_{\Omega} |\varphi|^{q(x)} dx \le \int_{\Omega_0} |\varphi|^{q^-} dx.$$

On the other hand,  $W_0^{1,p(x)}(\Omega)$  is continuously embedded in  $L^{q^-}(\Omega)$  and thus, there exists a positive constant  $c_2$  such that

$$|\varphi|_{q^-} \le c_2 \|\varphi\|.$$

The last two inequalities imply that

 $\|\varphi\| > 0$ 

and combining that fact with relations (4) or (5) we deduce that

$$\int_{\Omega} |\nabla \varphi|^{p(x)} \, dx > 0.$$

The proof of Lemma 2 is complete.

PROOF OF THEOREM 1. Let  $\lambda^* > 0$  be defined as in (10) and  $\lambda \in (0, \lambda^*)$ . By Lemma 1 it follows that on the boundary of the ball centered at the origin and of radius  $\rho$  in E, denoted by  $B_{\rho}(0)$ , we have

$$\inf_{\partial B_{\rho}(0)} J_{\lambda} > 0. \tag{11}$$

On the other hand, by Lemma 2, there exists  $\varphi \in E$  such that  $J_{\lambda}(t\varphi) < 0$  for all t > 0 small enough. Moreover, relations (9) and (5) imply that for any  $u \in B_{\rho}(0)$  we have

$$J_{\lambda}(u) \ge \frac{1}{p^{+}} \|u\|^{p^{+}} - \frac{\lambda}{q^{-}} c_{1}^{q^{-}} \|u\|^{q^{-}}$$

It follows that

$$-\infty < \underline{c} := \inf_{\overline{B_{\rho}(0)}} J_{\lambda} < 0.$$

We let now  $0 < \epsilon < \inf_{\partial B_{\rho}(0)} J_{\lambda} - \inf_{B_{\rho}(0)} J_{\lambda}$ . Applying Ekeland's variational principle to the functional  $J_{\lambda} : \overline{B_{\rho}(0)} \to \mathbb{R}$ , we find  $u_{\epsilon} \in \overline{B_{\rho}(0)}$  such that

$$\begin{aligned} J_{\lambda}(u_{\epsilon}) &< \inf_{\overline{B_{\rho}(0)}} J_{\lambda} + \epsilon \\ J_{\lambda}(u_{\epsilon}) &< J_{\lambda}(u) + \epsilon \cdot \|u - u_{\epsilon}\|, \quad u \neq u_{\epsilon}. \end{aligned}$$

Since

$$J_{\lambda}(u_{\epsilon}) \leq \inf_{\overline{B_{\rho}(0)}} J_{\lambda} + \epsilon \leq \inf_{B_{\rho}(0)} J_{\lambda} + \epsilon < \inf_{\partial B_{\rho}(0)} J_{\lambda} ,$$

we deduce that  $u_{\epsilon} \in B_{\rho}(0)$ . Now, we define  $I_{\lambda} : \overline{B_{\rho}(0)} \to \mathbb{R}$  by  $I_{\lambda}(u) = J_{\lambda}(u) + \epsilon \cdot ||u - u_{\epsilon}||$ . It is clear that  $u_{\epsilon}$  is a minimum point of  $I_{\lambda}$  and thus

$$\frac{I_{\lambda}(u_{\epsilon} + t \cdot v) - I_{\lambda}(u_{\epsilon})}{t} \ge 0$$

for small t > 0 and any  $v \in B_1(0)$ . The above relation yields

$$\frac{J_{\lambda}(u_{\epsilon}+t\cdot v)-J_{\lambda}(u_{\epsilon})}{t}+\epsilon\cdot \|v\|\geq 0.$$

Letting  $t \to 0$  it follows that  $\langle J'_{\lambda}(u_{\epsilon}), v \rangle + \epsilon \cdot ||v|| > 0$  and we infer that  $||J'_{\lambda}(u_{\epsilon})|| \le \epsilon$ .

We deduce that there exists a sequence  $\{w_n\} \subset B_{\rho}(0)$  such that

$$J_{\lambda}(w_n) \to \underline{c} \quad \text{and} \quad J'_{\lambda}(w_n) \to 0.$$
 (12)

It is clear that  $\{w_n\}$  is bounded in E. Thus, there exists  $w \in E$  such that, up to a subsequence,  $\{w_n\}$  converges weakly to w in E. Since  $q(x) < p^*(x)$  for all  $x \in \overline{\Omega}$  we deduce that E is compactly embedded in  $L^{q(x)}(\Omega)$ , hence  $\{w_n\}$  converges strongly to w in  $L^{q(x)}(\Omega)$ . So, by relations (6) and (3),

$$\lim_{n \to \infty} \int_{\Omega} |w_n|^{q(x)-2} w_n(w_n - w) \, dx = 0.$$

On the other hand, relation (12) yields

$$\lim_{n \to \infty} \langle J'_{\lambda}(w_n), w_n - w \rangle = 0.$$

Using the above information we find

$$\lim_{n \to \infty} \int_{\Omega} |\nabla w_n|^{p(x)-2} \nabla w_n \nabla (w_n - w) \, dx = 0.$$
<sup>(13)</sup>

Relation (13) and the fact that  $\{w_n\}$  converges weakly to w in E enable us to apply Theorem 3.1 in Fan and Zhang [9] in order to obtain that  $\{w_n\}$  converges strongly to w in E. So, by (12),

$$J_{\lambda}(w) = \underline{c} < 0 \quad \text{and} \quad J_{\lambda}'(w) = 0.$$
(14)

We conclude that w is a nontrivial weak solution for problem (1) and thus any  $\lambda \in (0, \lambda^*)$  is an eigenvalue of problem (1).

The proof of Theorem 1 is complete.

Let us now assume that the hypotheses of Theorem 1 are fulfilled and, furthermore,

$$\max_{\overline{\Omega}} p(x) < \max_{\overline{\Omega}} q(x).$$

Then, using similar arguments as in the proof of Lemma 2, we find some  $\psi \in E$  such that

$$\lim_{t \to \infty} J_{\lambda}(t\psi) = -\infty.$$

That fact combined with Lemma 1 and the mountain pass theorem (see [1]) implies that there exists a sequence  $\{u_n\}$  in E such that

$$J_{\lambda}(u_n) \to \overline{c} > 0 \quad \text{and} \quad J'_{\lambda}(u_n) \to 0 \text{ in } E^{\star}.$$
 (15)

However, relation (15) is not useful because we can not show that the sequence  $\{u_n\}$  is bounded in E since the functional  $J_{\lambda}$  does not satisfy a relation of the Ambrosetti-Rabinowitz type. This enable us to affirm that we can not obtain a critical point for  $J_{\lambda}$  by using this method.

On the other hand, we point out that we will fail in trying to show that the functional  $J_{\lambda}$  is coercive since by relation (2) we have  $q^+ > p^-$ . Thus, we can not apply (as in the homogeneous case) a result as Theorem 1.2 in Struwe [20] in order to obtain a critical point of the functional  $J_{\lambda}$ .

## References

- A. Ambrosetti and P. H. Rabinowitz, Dual variational methods in critical point theory, J. Funct. Anal. 14 (1973), 349-381.
- [2] A. Anane, Simplicité et isolation de la première valeur propre du p-laplacien avec poids, C. R. Acad. Sci. Paris Sér. I 305 (1987), 725-728.
- [3] M. Bocher, The smallest characteristic numbers in a certain exceptional case, Bull. Amer. Math. Soc. 21 (1914), 6-9.
- [4] L. Diening, Theoretical and Numerical Results for Electrorheological Fluids, Ph.D. thesis, University of Frieburg, Germany, 2002.
- [5] D. E. Edmunds, J. Lang, and A. Nekvinda, On  $L^{p(x)}$  norms, Proc. Roy. Soc. London Ser. A 455 (1999), 219-225.
- [6] D. E. Edmunds and J. Rákosník, Density of smooth functions in  $W^{k,p(x)}(\Omega)$ , Proc. Roy. Soc. London Ser. A 437 (1992), 229-236.
- [7] D. E. Edmunds and J. Rákosník, Sobolev embedding with variable exponent, Studia Math. 143 (2000), 267-293.
- [8] I. Ekeland, On the variational principle, J. Math. Anal. Appl. 47 (1974), 324-353.
- [9] X. L. Fan and Q. H. Zhang, Existence of solutions for p(x)-Laplacian Dirichlet problem, Nonlinear Anal. 52 (2003), 1843-1852.

- [10] X. Fan, Q. Zhang and D. Zhao, Eigenvalues of p(x)-Laplacian Dirichlet problem, J. Math. Anal. Appl. **302** (2005), 306-317.
- [11] T. C. Halsey, Electrorheological fluids, Science 258 (1992), 761-766.
- [12] P. Hess and T. Kato, On some linear and nonlinear eigenvalue problems with an indefinite weight function, Comm. Partial Differential Equations 5 (1980), 999-1030.
- [13] O. Kováčik and J. Rákosník, On spaces  $L^{p(x)}$  and  $W^{1,p(x)}$ , Czechoslovak Math. J. 41 (1991), 592-618.
- [14] M. Mihăilescu and V. Rădulescu, A multiplicity result for a nonlinear degenerate problem arising in the theory of electrorheological fluids, Proc. Roy. Soc. London Ser. A, in press.
- [15] S. Minakshisundaram and A. Pleijel, Some properties of the eigenfunctions of the Laplace-operator on Riemannian manifolds, *Canadian J. Math.* 1 (1949), 242-256.
- [16] J. Musielak, Orlicz Spaces and Modular Spaces, Lecture Notes in Mathematics, Vol. 1034, Springer, Berlin, 1983.
- [17] A. Pleijel, On the eigenvalues and eigenfunctions of elastic plates, Comm. Pure Appl. Math. 3 (1950), 1-10.
- [18] M. Ruzicka, Electrorheological Fluids: Modeling and Mathematical Theory, Springer-Verlag, Berlin, 2002.
- [19] S. Samko and B. Vakulov, Weighted Sobolev theorem with variable exponent for spatial and spherical potential operators, J. Math. Anal. Appl. 310 (2005), 229-246.
- [20] M. Struwe, Variational Methods: Applications to Nonlinear Partial Differential Equations and Hamiltonian Systems, Springer, Heidelberg, 1996.
- [21] V. Zhikov, Averaging of functionals in the calculus of variations and elasticity, Math. USSR Izv. 29 (1987), 33-66.