On the Frobenius Number of Fibonacci Numerical Semigroups

J.M. Marín*

Dep. Matemática Aplicada I Universidad de Sevilla. Avda. Reina Mercedes s/n. 41012 Sevilla, Spain J. L. Ramírez Alfonsín[†] Equipe Combinatoire et Optimisation, Université Pierre et Marie Curie (Paris 6), Case 189 - 4 Place Jussieu, 75252 Paris, Cedex 05, France and M.P. Revuelta[‡]

Dep. Matemática Aplicada I Universidad de Sevilla. Avda. Reina Mercedes s/n. 41012 Sevilla, Spain

September 19, 2018

Abstract

In this paper we compute the Frobenius number of certain *Fibonacci numerical semigroups*, that is, numerical semigroups generated by a set of Fibonacci numbers, in terms of Fibonacci numbers.

^{*}e-mail: jmarin@us.es

 $^{^{\}dagger}e$ -mail: ramirez@math.jussieu

[‡]*e-mail*: pastora@us.es

1 Introduction

Let s_1, \ldots, s_n be positive integers such that their greatest commun divisor is one. Let $S = \langle s_1, \ldots, s_n \rangle$ be the numerical semigroup¹ generated by s_1, \ldots, s_n . A Fibonacci numerical semigroup is a numerical semigroup generated by a set of Fibonacci numbers F_{i_1}, \ldots, F_{i_r} , for some integers $3 \leq i_1 < \cdots < i_r$ where $g.c.d.(F_{i_1}, \ldots, F_{i_r}) = 1$.

The so-called *Frobenius number*, denoted by $g(s_1, \ldots, s_n)$, is defined as the largest integer not belonging to S, that is, the largest integer that is not representable as a nonnegative integer combination of s_1, \ldots, s_n . It is well known that $g(s_1, s_2) = s_1s_2 - s_1 - s_2$. In general, finding g(S) is a difficult problem and so formulas and upper bounds for particular sequences are of interest. For instance, it is known [3] the value of g(S) when S is an arithmetical sequence

$$g(a, a+d, \dots, a+kd) = a\left(\left\lfloor \frac{a-2}{k} \right\rfloor\right) + d(a-1)$$
(1)

We refer the reader to [2] where a complete account on the Frobenius problem can be found.

In this paper, we investigate the value of $g(F_i, F_j, F_l)$ for some triples $3 \leq i < j < l$ (we always assume that $g.c.d.(F_i, F_j, F_l) = 1$, recall that $g.c.d.(F_i, F_{i+l}) = 1$ if $i \not| l$).

We first notice that $g(F_i, F_{i+1}, F_l) = g(F_i, F_{i+1})$ for any integer $l \ge i + 2$. Indeed, since $F_l = F_{i+m} = F_m F_{i+1} + F_{m-1} F_i$ is a nonnegative integer combination of F_i and F_{i+1} then the semigroups $\langle F_i, F_{i+1}, F_l \rangle$ and $\langle F_i, F_{i+1} \rangle$ generate the same set of elements and thus they have the same Frobenius number.

¹Recall that a semigroup (S, *) consists of a nonempty set S and an associative binary operation * on S. If, in addition, there exists an element, which is usually denoted by 0, in S such that a + 0 = 0 + a = a for all $a \in S$, we say that (S, *) is a monoid. A numerical semigroup is a submonoid of \mathbb{N} such that the greatest common divisor of its elements is equal to one.

Let us consider then $g(F_i, F_{i+2}, F_l)$ with $l \ge i+3$. We notice that the case when l = i+3 is a consequence of equation (1) since the triple $\{F_i, F_{i+2}, F_{i+3}\} = \{F_i, F_i + F_{i+1}, F_i + 2F_{i+1}\}$ form an arithmetical sequence. However, it can be checked that $\{F_i, F_{i+2}, F_{i+k}\}$ do not form an arithmetical sequence when $k \ge 3$ and the calculation of $g(F_i, F_{i+2}, F_{i+k})$ is more complicated.

We state our main result.

Theorem 1.1 Let $i, k \geq 3$ be integers and let $r = \lfloor \frac{F_i - 1}{F_k} \rfloor$. Then,

$$g(F_i, F_{i+2}, F_{i+k}) = \begin{cases} (F_i - 1)F_{i+2} - F_i(rF_{k-2} + 1) & \text{if } r = 0 \text{ or } r \ge 1 \text{ and} \\ F_{k-2}F_i < (F_i - rF_k)F_{i+2}, \\ (rF_k - 1)F_{i+2} - F_i((r-1)F_{k-2} + 1) & \text{otherwise.} \end{cases}$$

Let $N(a_1, \ldots, a_n)$ be the number of positive integers with no representation by a nonnegative integer combination of a_1, \ldots, a_n . We refer the reader to [2, Chapter 5] for results related to $N(a_1, \ldots, a_n)$. Theorem 1.1 yields to the following result.

Corollary 1.2 Let $i, k \geq 3$ be integers and let $r = \lfloor \frac{F_i - 1}{F_k} \rfloor$. Then,

$$N(F_i, F_{i+2}, F_{i+k}) = \frac{(F_i - 1)(F_{i+2} - 1) - rF_{k-2}(2F_i - F_k(1 + r))}{2}$$

2 Fibonacci semigroups

In order to prove Theorem 1.1 we need the following result due to Brauer and Shockley [1].

Lemma 2.1 Let $1 < a_1 < \cdots < a_n$ be integers with $g.c.d.(a_1, \ldots, a_n) = 1$. Then,

$$g(a_1, \dots, a_n) = \max_{l \in \{1, 2, \dots, a_n - 1\}} \{t_l\} - a_1$$

where t_l is the smallest positive integer congruent to l modulo a_1 , that is representable as a nonnegative integer combination of a_2, \ldots, a_n .

Proof. Let L be a positive integer. If $L \equiv 0 \mod a_1$ then L is a non-negative integer combination of a_1 . If $L \equiv l \mod a_1$ then L is a nonnegative integer combination of a_1, \ldots, a_n if and only if $L \geq t_l$.

Proof of Theorem 1.1. Let $T^* = \{t_0^*, \ldots, t_{F_i-1}^*\}$ where t_l^* be the smallest positive integer congruent to l modulo F_i , that is representable as a nonnegative integer combination of F_{i+2} and F_{i+k} . We shall find t_l^* for each $l = 0, 1, \ldots, F_i - 1$. To this end, we consider all nonnegative integer combinations of F_{i+2} and F_{i+k} . We construct the following table, denoted by T_1 , having as entry $t_{x,y}$ the combination of the form $xF_{i+2} + yF_{i+k}$ with integers $x, y \ge 0$, see below.

$x \backslash y$	0	1	2	•••
0	0	F_{i+k}	$2F_{i+k}$	• • •
1	F_{i+2}	$F_{i+k} + F_{i+2}$	$2F_{i+k} + F_{i+2}$	•••
2	$2F_{i+2}$	$F_{i+k} + 2F_{i+2}$	$2F_{i+k} + 2F_{i+2}$	•••
3	$3F_{i+2}$	$F_{i+k} + 3F_{i+2}$	$2F_{i+k} + 3F_{i+2}$	•••
:	:	:	:	
$F_k - 1$	$(F_k - 1)F_{i+2}$	$F_{i+k} + (F_k - 1)F_{i+2}$	$2F_{i+k} + (F_k - 1)F_{i+2}$	• • •
:	•	÷	÷	

We notice that

 $F_{i+k} = F_{k-2}F_{i+1} + F_{k-1}F_{i+2} = F_{k-2}(F_{i+2} - F_i) + F_{k-1}F_{i+2} = F_{i+2}F_k - F_{k-2}F_i$ so, we obtain that

$$xF_{i+2} + yF_{i+k} = xF_{i+2} + y(F_{i+2}F_k - F_{k-2}F_i) = (x + yF_k)F_{i+2} - yF_{k-2}F_i.$$

Thus, T_1 can also be given by the following table, denoted by T_2 ,

x ackslash y	0	1	2		r	
0	0	$F_k F_{i+2} - F_{k-2} F_i$	$2F_kF_{i+2} - 2F_{k-2}F_i$		$rF_kF_{i+2} - rF_{k-2}F_i$	• • •
1	F_{i+2}	$(1+F_k)F_{i+2} - F_{k-2}F_i$	$(1+2F_k)F_{i+2} - 2F_{k-2}F_i$		$(1+rF_k)F_{i+2} - rF_{k-2}F_i$	• • •
2	$2F_{i+2}$	$(2+F_k)F_{i+2} - F_{k-2}F_i$	$(2+2F_k)F_{i+2} - 2F_{k-2}F_i$	• • •	$(2 + rF_k)F_{i+2} - rF_{k-2}F_i$	•••
:					:	
l	lF_{i+2}	$(l+F_k)F_{i+2} - F_{k-2}F_i$	$(l+2F_k)F_{i+2} - 2F_{k-2}F_i$		$(l+rF_k)F_{i+2} - rF_{k-2}F_i$	•••
:						
$F_k - 1$	$(F_k - 1)F_{i+2}$	$(2F_k - 1)F_{i+2} - F_{k-2}F_i$	$(3F_k - 1)F_{i+2} - 2F_{k-2}F_i$		••••	
÷	÷	÷	÷		÷	

Let S be the set formed by the first $F_k - 1$ entries of columns zero, one, two, and so on, that is,

$$S = \{t_{0,0}, t_{1,0}, \dots, t_{F_k-1,0}, t_{0,1}, t_{1,1}, \dots, t_{F_k-1,1}, \dots, t_{0,r}, t_{1,r}, \dots, t_{F_k-1,r}, \dots\}.$$

Remark 2.2 (a) Let $r = \lfloor \frac{F_i - 1}{F_k} \rfloor$ and set $F_i - 1 = rF_k + l$ for some integer $0 \le l \le F_k - 1$. Let

$$S' = \{t_{0,0}, t_{1,0}, \dots, t_{F_k-1,0}, t_{0,1}, t_{1,1}, \dots, t_{F_k-1,1}, \dots, t_{2,r}, t_{1,r}, \dots, t_{l,r}\},\$$

Then, for each $t_{x,y} = (x + yF_k)F_{i+2} - yF_{k-2}F_i \in S'$ we have that $0 \leq x + yF_k \leq F_i - 1$. Moreover, since g.c.d. $(F_{i+2}, F_i) = 1$ then S' forms a complete system of rests modulo F_i .

(b) The elements of S can be represented as $s_x = xF_{i+2} - \lfloor \frac{x}{F_k} \rfloor F_{k-2}F_i$ for $x = 0, 1, \ldots$ Indeed, it can be checked that $S = \bigcup_{q \ge 1} S_q$ where

$$S_q = \{s_{qF_k}, s_{qF_k+1}, \dots, s_{(q+1)F_k-1}\} = \{t_{0,q}, \dots, t_{F_k-1,q}\}$$

for each integer q = 0, 1, 2, ...

(c) By using table T_2 we have that $t_{i,j} < t_{k,l}$ for all $i \leq k$ and all $j \leq l$.

Lemma 2.3 Let $t_{u,v}$ be an entry of T_1 such that $t_{u,v} \notin S'$. There exists $t_{x,y} \in S'$ such that $t_{u,v} \equiv t_{x,y} \mod F_i$ and $t_{u,v} > t_{x,y}$.

Proof. We first notice that the set S can be written as follows

$$\begin{cases} s_0, \dots, s_{F_k-1}, s_{F_k}, \dots, s_{2F_k-1}, \dots, s_{rF_k}, \dots, s_{rF_k+l} = s_{F_l-1}, \\ s_{F_l}, \dots, s_{F_l+F_k-1}, s_{F_l+F_k}, \dots, s_{F_l+2F_k-1}, \dots, s_{F_l+rF_k}, \dots, s_{2F_l-1}, \\ s_{2F_l}, \dots, s_{2F_l+F_k-1}, s_{2F_l+F_k}, \dots, s_{2F_l+2F_k-1}, \dots, s_{2F_l+rF_k}, \dots, s_{3F_l-1}, \dots \end{cases}$$

where $S' = \{s_0, \ldots, s_{F_k-1}, s_{F_k}, \ldots, s_{2F_k-1}, \ldots, s_{rF_k}, \ldots, s_{F_{i-1}}\}$. We have two cases.

Case A) Suppose that $t_{u,v} \in S \setminus S'$. Then $t_{u,v}$ is of the form s_{pF_i+g} for some integers $p \ge 1$ and $0 \le g \le F_i - 1$. It is clear that,

$$s_g = gF_{i+2} - \left\lfloor \frac{g}{F_k} \right\rfloor F_i F_{k-2} \equiv (pF_i + g)F_{i+2} - \left\lfloor \frac{pF_i + g}{F_k} \right\rfloor F_i F_{k-2} = g_{pF_i + g} \mod F_i$$

We will show that $s_{pF_i+g} > s_g$. To this end, it suffice to prove that $s_{F_i+g} > s_g$ (since $s_{pF_i+g} \ge s_{F_i+g}$). Recall that $r = \lfloor \frac{F_i-1}{F_k} \rfloor$ and that $F_i - 1 = rF_k + l$ for some integer $0 \le l \le F_k - 1$. We have two subcases.

Subcase a) If r = 0 then $F_k \ge F_i$. If $F_k = F_i$ then $s_{F_i+g} = t_{g,1}$ and, by Remark 2.2 (c), $t_{g,0} < t_{g,1}$. If $F_k > F_i$ then $s_{F_i+g} = t_{q,0}$ for some integer $q \ge F_i$ and, by Remark 2.2 (c), $t_{g,0} < t_{q,0}$.

Subcase b) If $r \ge 1$ then $s_{F_i+g} > s_g$ holds if and only if

$$(F_i+g)F_{i+2} - \left\lfloor \frac{F_i+g}{F_k} \right\rfloor F_iF_{k-2} > gF_{i+2} - \left\lfloor \frac{g}{F_k} \right\rfloor F_iF_{k-2}$$

or equivalently if and only if

$$F_{i+2} > F_{k-2} \left(\left\lfloor \frac{F_i + g}{F_k} \right\rfloor - \left\lfloor \frac{g}{F_k} \right\rfloor \right)$$

Let $g = mF_k + n$ with $0 \le n \le F_k - 1$. Since $F_i - 1 = rF_k + l$ with $0 \le l \le F_k - 1$ then

$$\left\lfloor \frac{F_i - 1 + g + 1}{F_k} \right\rfloor = \left\lfloor \frac{rF_k + l + mF_k + n + 1}{F_k} \right\rfloor \le r + m + 1$$

and thus

$$\left\lfloor \frac{F_i + g}{F_k} \right\rfloor - \left\lfloor \frac{g}{F_k} \right\rfloor \le r + m + 1 - m = r + 1.$$

So, it is enough to show that $F_{i+2} > (r+1)F_{k-2}$ or equivalently to show that $F_i + F_{i+1} > (r+1)F_{k-2}$. Since $F_i = rF_k + l + 1$ then the latter inequality holds if and only if $rF_k + l + 1 + F_{i+1} > rF_{k-2} + F_{k-2}$, that is, if and only if

$$r(F_k - F_{k-2}) + l + 1 + F_{i+1} = r(F_{k-1}) + l + 1 + F_{i+1} > F_{k-2}$$

which is true since $r \ge 1$.

Case B) Suppose that $t_{u,v} \notin S$. Then we have that $0 \leq x \leq F_k - 1 < u$. If $v \geq y$ then, by Remark 2.2 (c), $t_{x,y} < t_{x,v} < t_{u,v}$. So, we suppose that v < y. Since, $t_{u,v} \equiv t_{x,y} \mod F_i$ then $u + vF_k \equiv x + yF_k \mod F_i$ but, by Remark 2.2 (a), $0 \leq x + yF_k \leq F_i - 1$ so $u + vF_k = d(x + yF_k)$ for some integer $d \geq 1$ and thus $u + vF_k \geq x + yF_k$. Also, since v < y, then $-vF_{k-2}F_i > -yF_{k-2}F_i$. So, combining the last two inequalities we have that

$$t_{u,v} = (u + vF_k)F_{i+2} - vF_{k-2}F_i > (x + yF_k)F_{i+2} - yF_{k-2}F_i = t_{x,y}.$$

Therefore, by the above lemma, we have that for each $x = 0, \ldots, F_i - 1$, s_x is the smallest positive integer congruent to l modulo F_i , for some integer $0 \le l \le F_i - 1$, that is representable as a nonnegative integer combination of F_{i+2} and F_{i+k} , that is, $S' = T^*$.

Now, by Remark 2.2 (c), if $r \ge 1$ then

$$t_{F_k-1,i} = \max_{0 \le x \le F_k-1} \{ t_{x,i} | t_{x,i} \in S' \} \text{ for each } i = 0, \dots, r-1,$$
$$t_{F_k-1,r-1} = \max_{0 \le i \le r-1} \{ t_{F_k-1,i} | t_{F_k-1,i} \in S' \},$$

and

$$t_{l,r} = \max_{0 \le x \le l} \{ t_{x,r} | t_{x,r} \in S' \}$$

Thus,

$$\max\{s|s \in S'\} = \begin{cases} t_{l,r} & \text{if } r = 0, \\ \max\{t_{F_k-1,r-1}, t_{l,r}\} & \text{otherwise.} \end{cases}$$

The result follows since $t_{l,r} > t_{F_k-1,r-1}$ if and only if

$$(rF_{k}+l)F_{i+2}-rF_{k-2}F_{i} = (F_{i}-1)F_{i+2}-rF_{k-2}F_{i} > (rF_{k}-1)F_{i+2}-(r-1)F_{k-2}F_{i}$$

or equivalently, if and only if $F_{i+2}(F_{i}-rF_{k}) > F_{k-2}F_{i}$.

We will use the following result due to Selmer [4] to show Corollary 1.2.

Lemma 2.4 Let $1 < a_1 < \cdots < a_n$ be integers with $g.c.d.(a_1, \ldots, a_n) = 1$. If $L = \{1, \ldots, a_1 - 1\}$ then

$$N(a_1, \dots, a_n) = \frac{1}{a_1} \sum_{l \in L} t_l - \frac{a_1 - 1}{2}$$

where t_l is the smallest positive integer congruent to l modulo a_1 , that is representable as a nonnegative integer combination of a_2, \ldots, a_n .

Proof. The number of $M \equiv l \not\equiv 0 \mod a_1$ with $0 < M < t_l$ is given by $\lfloor \frac{t_1}{a_1} \rfloor$. By assuming that $0 < l < a_1$, we have $\lfloor \frac{t_l}{a_1} \rfloor = \frac{t_l - l}{a_1}$. The result follows by summing over $l \in L$.

Proof of Corollary 1.2. Let
$$r = \lfloor \frac{F_i - 1}{F_k} \rfloor$$
 and set $F_i - 1 = rF_k + l$ for some

integer $0 \le l \le F_k - 1$. By Lemma 2.4 and Remark 2.2 (b), we have

$$\begin{split} N(F_i, F_{i+2}, F_{i+k}) &= \frac{1}{F_i} \sum_{s \in S'} s - \frac{F_i - 1}{2} \\ &= \frac{1}{F_i} \sum_{j=0}^{F_i - 1} (jF_{i+2} - F_{k-2} \lfloor \frac{j}{F_k} \rfloor F_i) - \frac{F_i - 1}{2} \\ &= \frac{1}{F_i} \left(F_{i+2} \frac{(F_i - 1)F_i}{2} \right) - \frac{1}{F_i} (F_{k-2}F_i) \sum_{j=0}^{F_i - 1} \lfloor \frac{j}{F_k} \rfloor - \frac{F_i - 1}{2}. \end{split}$$

By using the table T_1 , it is easy to verify that

$$\sum_{j=0}^{F_{k}-1} \left\lfloor \frac{j}{F_{k}} \right\rfloor = 0 + F_{k} + 2F_{k} + \dots + (r-1)F_{k} + r(l+1) = \frac{F_{k}(r-1)r}{2} + r(l+1)$$

and, since $l + 1 = F_i - rF_k$, then

$$N(F_i, F_{i+2}, F_{i+k}) = \frac{F_{i+2}(F_i-1)}{2} - F_{k-2} \left(\frac{F_k(r-1)r}{2} + r(F_i - rF_k) \right) - \frac{F_i-1}{2} \\ = \frac{(F_i-1)(F_{i+2}-1)}{2} - F_{k-2} \left(\frac{F_kr^2 - F_kr + 2F_ir - 2r^2F_k}{2} \right) \\ = \frac{(F_i-1)(F_{i+2}-1) - rF_{k-2}(2F_i - F_k(1+r))}{2}.$$

We end with the following problem.

Problem 2.5 Find upper (and lower) bounds (or formulas) for $g(F_i, F_j, F_k)$ for further triples $3 \le i < j < k$.

References

- A. Brauer and J.E. Shockley, On a problem of Frobenius, *Journal für Reine und Angewandte Mathematik* 211 (1962), 215-220.
- [2] J.L. Ramírez Alfonsín, The Diophantine Frobenius Problem, Oxford Lectures Series in Mathematics and its Applications 30, Oxford University Press, (2005).
- [3] J.B. Roberts, Note on linear forms, Proc. Amer. Math. Soc. 7 (1956), 465-469.
- [4] E.S. Selmer, On the linear diophantine Problem of Frobenius, Journal für Reine und Angewandte Mathematik 293/294(1) (1977), 1-17.

AMS Classification number: 11B39, 11D04, 11A07