RING-THEORETIC PROPERTIES OF PvMDS

SAID EL BAGHDADI AND STEFANIA GABELLI

ABSTRACT. We extend to Prüfer v-multiplication domains some distinguished ring-theoretic properties of Prüfer domains. In particular we consider the t##-property, the t-radical trace property, w-divisoriality and w-stability.

INTRODUCTION

A Prüfer v-multiplication domain, for short a PvMD, is a domain whose localizations at tmaximal ideals are valuation domains [22]. For this reason, the ideal-theoretic properties of valuation domains globalize to t-ideals of PvMDs and several properties of ideals of Prüfer domains hold for t-ideals of PvMDs: for example a domain is a PvMD if and only if each t-finite t-ideal is t-invertible. The aim of this paper is to show that, introducing suitable t-analogues of some distinguished properties of integral domains, Prüfer domains and PvMDs have a similar behaviour also from the ring-theoretic point of view. We recall that the class of PvMDs, besides Prüfer domains, includes Krull domains and GCD-domains.

As a matter of fact, the *t*-operation is not always as good as the *w*-operation for extending certain properties that hold in the classical case, that is in the *d*-operation setting. Thus in general it is often more convenient to consider the *w*-analogue of a given property (see for instance [43, 44, 9]). However in a P*v*MD the *w*-operation and the *t*-operation coincide [30] and one can use indifferently these two star operations.

In Section 1, we deal with the t##-property and the tRTP-property. PvMDs satisfying the t##-property have been studied in [15, Section 2]. Here we characterize PvMDs with the tRTP-property; getting for example that a PvMD is a tRTP-domain if and only if each v-finite divisorial ideal has at most finitely many minimal primes. Then, generalizing the Prüfer case, we show that the t##-property and the tRTP-property are strictly connected for PvMDs. Among other results, we prove that a PvMD satisfying the t##-property is a tRTP-domain and that the converse holds if each t-prime is branched. We also show that an almost Krull domain satisfying the t##-property is a Krull domain.

In Section 2 we introduce the notion of w-stability and relate it to w-divisoriality, a property defined and studied in [9]. First we show that the study of w-stability can be reduced to the t-local case. Then we use this result to extend to PvMDs some properties of stable and divisorial Prüfer domains. For example, we prove that w-stability of t-primes enforces a PvMD to be a generalized Krull domains and that an integrally closed w-stable domain is precisely a generalized Krull domain with t-finite character. We also characterize w-stable w-divisorial PvMDs and show that these domains behave like totally divisorial Prüfer domains.

We assume that the reader is familiar with the language of star operations [18, Sections 32 and 34]. We recall some definitions and basic properties.

Date: January 5, 2018.

¹⁹⁹¹ Mathematics Subject Classification. Primary: 13F05; Secondary: 13A15, 13A18.

Key words and phrases. Prüfer v-multiplication domain, #-property, radical trace property, divisoriality, stability.

Throughout this paper R will denote an integral domain with quotient field K and we will assume that $R \neq K$.

A star operation is a map $I \to I^*$ from the set F(R) of nonzero fractional ideals of R to itself such that:

- (1) $R^* = R$ and $(aI)^* = aI^*$, for all $a \in K \setminus \{0\}$;
- (2) $I \subseteq I^*$ and $I \subseteq J \Rightarrow I^* \subseteq J^*$;
- (3) $I^{**} = I^*$.

A star operation * is of *finite type* if $I^* = \bigcup \{J^*; J \subseteq I \text{ and } J \text{ is finitely generated}\}$, for each $I \in F(R)$. To any star operation *, we can associate a star operation $*_f$ of finite type by defining $I^{*_f} = \bigcup J^*$, with the union taken over all finitely generated ideals J contained in I. Clearly $I^{*_f} \subseteq I^*$.

An ideal $I \in F(R)$ is a *-*ideal* if $I = I^*$ and is *-*finite* if $I^* = J^*$ for some finitely generated ideal J. A *-finite *-ideal is also called a *-*ideal of finite type*.

A *-prime is a prime *-ideal and a *-maximal ideal is an ideal that is maximal in the set of the proper *-ideals. A *-maximal ideal (if it exists) is a prime ideal. If * is a star operation of finite type, an easy application of Zorn's Lemma shows that the set *-Max(R) of the *-maximal ideals of R is not empty. In this case, for each $I \in F(R)$, $I^* = \bigcap_{M \in *-Max(R)} I^*R_M$; in particular $R = \bigcap_{M \in *-Max(R)} R_M$ [22].

For any star operation *, the set of *-ideals of R is a semigroup under the *-multiplication, defined by $(I, J) \mapsto (IJ)^*$, with unity R. An ideal $I \in F(R)$ is called *-invertible if I^* is invertible with respect to the *-multiplication. In this case the *-inverse of I is (R : I). Thus I is *-invertible if and only if $(I(R : I))^* = R$.

The identity is a star operation, called the *d*-operation. The *v*-operation (or divisorial closure), the *t*-operation and the *w*-operation are the best known nontrivial star operations and are defined in the following way. For each $I \in F(R)$, we set $I_v := (R: (R: I))$ and $I_t := \bigcup J_v$ with the union taken over all finitely generated ideals J contained in I. Hence the *t*-operation is the finite type star operation associated to the *v*-operation. The *w*-operation is the star operation of finite type defined by setting $I_w := \bigcap_{M \in t-Max(R)} IR_M$. We have w-Max(R) = t-Max(R) and $IR_M = I_wR_M \subseteq I_tR_M$, for each $M \in t-Max(R)$. Thus $I_w \subseteq I_t \subseteq I_v$. Also, an ideal $I \in F(R)$ is *w*-invertible if and only if it is *t*-invertible.

A Prüfer domain is an integrally closed domain such that d = t [18, Proposition 34.12] and a PvMD is an integrally closed domain such that w = t [30, Theorem 3.1].

The v-, t- and w-operations on R can be extended to the set of nonzero R-submodules of K by setting $E^v := (R : (R : E)), E_t = \bigcup \{F_v; F \subseteq E; F \text{ finitely generated}\}$ and $E_w = \bigcap \{E_M; M \in t\text{-Max}(R)\}$, for each non zero R-submodule E of K. In this way, one obtains semistar operations on R. For more details, see for example [12]. By viewing w as a semistar operation on R, we can say that an overring T of a domain R is t-linked over R if $T_w = T$ [4, Proposition 2.13]. Each overring of R is t-linked precisely when d = w [4, Theorem 2.6].

We denote by t-Spec(R) the set of t-prime ideals of R. Each height-one prime is a t-prime and each prime minimal over a t-ideal is a t-prime. We say that R has t-dimension one if each t-prime ideal has height one.

We now define the ring-theoretic properties considered in this paper.

The t##-property. The #-property and the ##-property were introduced by R. Gilmer [19] and R. Gilmer and W. Heinzer [20] respectively. A domain R has the #-property, or it is a #-domain, if $\bigcap_{M \in \mathcal{M}_1} R_M \neq \bigcap_{M \in \mathcal{M}_2} R_M$, for any pair of distinct nonempty sets \mathcal{M}_1 and \mathcal{M}_2 of maximal ideals. Any overring of a one-dimensional Prüfer #-domain is a #-domain [19, Corollary 2], but in general

 $\mathbf{2}$

the #-property is not inherited by overrings [20, Section 2]. One says that R has the ##-property, or it is a ##-domain, if each overring of R is a #-domain.

The t#-property was introduced and studied in [15]. A domain R has the t#-property (or is a t#-domain) if $\bigcap_{M \in \mathcal{M}_1} R_M \neq \bigcap_{M \in \mathcal{M}_2} R_M$ for any two distinct subsets \mathcal{M}_1 and \mathcal{M}_2 of t-Max(R). Although in [15] it was explored the transfer of the t#-property to some distinguished classes of overrings, it was not given any definition for the t##-property. Here, we say that R has the t##property (or is a t##-domain) if $\bigcap_{P \in \mathcal{P}_1} R_P \neq \bigcap_{P \in \mathcal{P}_2} R_P$ for any two distinct subsets \mathcal{P}_1 and \mathcal{P}_2 of pairwise incomparable t-prime ideals of R. Our definition is motivated by the fact that for a PvMD this is equivalent to say that each t-linked overring has the t#-property [15, Proposition 2.10].

The tRTP-property. If R is an integral domain and M is a unitary R-module, the trace of M is the ideal of R generated by the set $\{f(m); f \in \operatorname{Hom}_R(M, R), m \in M\}$. An ideal J of R is called a trace ideal or a strong ideal if it is the trace of some R-module M. This happens if and only if J = I(R : I), for some nonzero ideal I of R, equivalently (J : J) = (R : J) [10, Lemmas 4.2.2. and 4.2.3]. If V is a valuation domain, a trace ideal is either equal to V or it is prime [10, Proposition 4.2.1]; this fact led to the consideration of several conditions related to trace ideals [28]. The radical trace property was introduced by W. Heinzer and I. Papick [24] and is particularly significant for Prüfer domains [24, 33]. R is a domain satisfying the radical trace property, or it is an RTP-domain, if each proper strong ideal is a radical ideal, that is, for each nonzero ideal I of R, either I(R : I) = R or I(R : I) is a radical ideal.

A. Mimouni studied trace properties in the setting of star operations, in particular he considered the t-operation [35]. As in [35], we say that a domain R has the t-radical trace property, or it is a tRTP-domain, if each proper strong t-ideal of R is a radical ideal. This is equivalent to say that, for each nonzero ideal I of R, either $(I(R:I))_t = R$ or $(I(R:I))_t$ is a radical ideal.

w-divisoriality. The class of domains in which each nonzero ideal is divisorial has been studied, independently and with different methods, by H. Bass [2], E. Matlis [34] and W. Heinzer [23] in the sixties. Following S. Bazzoni and L. Salce [3], a domain in which each nonzero ideal is divisorial is now called a *divisorial domain* and a domain such that each overring is divisorial is called *totally divisorial*.

The most suitable star analogue of divisoriality is the notion of w-divisoriality that was introduced and extensively studied in [9]. A *w*-divisorial domain is a domain such that each *w*-ideal is divisorial.

w-stability. Motivated by earlier work of H. Bass [2] an J. Lipman [32] on the number of generators of an ideal, in 1974 J. Sally and W. Vasconcelos defined a Noetherian ring R to be *stable* if each nonzero ideal of R is projective over its endomorphism ring $End_R(I)$ [41, 42]. When I is a nonzero ideal of a domain R, then $End_R(I) = (I : I)$; thus a domain R is stable if each nonzero ideal I of R is invertible in the overring (I : I). B. Olberding showed that stability and divisoriality are strictly connected and that stability is particularly significant in the context of Prüfer domains [10, 36, 37, 38, 39].

We introduce the notion of w-stability in Section 2. We say that a w-ideal I of a domain R is w-stable if I is w-invertible in the overring E(I) := (I : I), that is $(I(E(I) : I))_w = E(I)$, and say that R is w-stable if each w-ideal of R is w-stable. For a more general notion of stability with respect to a semistar operation we refer the reader to the forthcoming paper [16].

1. t##-Property and t-radical trace property

The ##-property and the radical trace property are closely related for a Prüfer domain. In this section we compare the *t*-analogues of these two properties for PvMDs.

Several characterizations of PvMDs satisfying the t##-property have been given in [15, Section 2]. For the study of the tRTP-property, we need some results on branched t-primes. Recall that a prime ideal P of a domain R is *branched* if there exists a P-primary ideal distinct from P. Clearly P is branched if and only if PR_P is branched. Since the localization of a PvMD at a t-prime is a valuation domain, the branched t-primes of PvMDs can be characterized by properties similar to those well known for the branched primes of Prüfer domains [18, Theorem 23.3 (e)].

Lemma 1.1. Let R be a PvMD and $J := x_1R + \cdots + x_nR$ a nonzero finitely generated ideal such that $J_v \neq R$. If P is a t-prime containing J, then P is minimal over J_v if and only if P is minimal over J, if and only if P is minimal over x_iR , for some $i, 1 \leq i \leq n$.

Proof. It is enough to observe that, since R_P is a valuation domain, we have $J_v R_P = J_t R_P = JR_P = x_i R_P$, for some $i, 1 \le i \le n$.

Proposition 1.2. Let R be a PvMD and P a t-prime of R. The following conditions are equivalent:

- (i) *P* is branched;
- (ii) P is a minimal prime of a principal ideal;
- (iii) *P* is a minimal prime of a finitely generated ideal;
- (iv) *P* is a minimal prime of a *v*-finite divisorial ideal;
- (v) P is not the union of the set of (t)-primes of R properly contained in P.

Proof. The equivalence of conditions (i), (ii) and (v) is obtained by localizing at P and using [18, Theorem 17.3 (e)]. The equivalence of conditions (ii), (iii) and (iv) follows from Lemma 1.1.

In any commutative ring with unity, if each minimal prime of an ideal I is the radical of a finitely generated ideal, then I has only finitely many minimal primes [21, Theorem 1.6]. By passing through the *t*-Nagata ring, we now show that a similar result holds for *t*-ideals of PvMDs.

If R is an integral domain, we set $R(X) := R[X]_N$, where $N = \{f \in R[X] : c(f) = R\}$ and $R\langle X \rangle := R[X]_{N_t}$, where $N_t = \{f \in R[X]; c(f)_t = R\}$. R(X) is called the *Nagata ring* of R and $R\langle X \rangle$ the t-Nagata ring of R [12, 29, 30].

B. G. Kang proved that R is a PvMD if and only if $R\langle X \rangle$ is a Prüfer (indeed a Bezout) domain [30, Theorem 3.7]. In addition, there is a lattice isomorphism between the lattice of *t*-ideals of R and the lattice of ideals of $R\langle X \rangle$ [30, Theorem 3.4]. More precisely, we have:

Proposition 1.3. Let R be a PvMD. Then the map $I_t \mapsto IR\langle X \rangle$ is an order-preserving bijection between the set of t-ideals of R and the set of nonzero ideals of $R\langle X \rangle$, whose inverse is the map $J \mapsto J \cap R$. Moreover, P is a t-prime (respectively, t-maximal) ideal of R if and only if $PR\langle X \rangle$ is a prime (respectively, maximal) ideal of $R\langle X \rangle$ and we have $R\langle X \rangle_{PR\langle X \rangle} = R[X]_{PR[X]} = R_P(X)$.

Proposition 1.4. Let R be a PvMD and I a proper t-ideal of R. If each minimal prime of I is the radical of a v-finite divisorial ideal, then I has finitely many minimal t-primes.

Proof. Each minimal prime of a *t*-ideal *I* is a *t*-ideal of *R*. By Proposition 1.3, the map $P \mapsto PR\langle X \rangle$ is a bijection between the set of minimal primes of *I* and the set of minimal primes of $IR\langle X \rangle$. Moreover if *J* is a nonzero finitely generated ideal of *R* such that $P = \operatorname{rad}(J_v)$, then $PR\langle X \rangle = \operatorname{rad}(JR\langle X \rangle)$. Hence each minimal prime of $IR\langle X \rangle$ is the radical of a finitely generated ideal. By [21, Theorem 1.6], $IR\langle X \rangle$ has finitely many minimal primes and the same holds for *I*. \Box

If T is an overring of R, the w-operation and the t-operation on R, viewed as semistar operations, induce two semistar operations of finite type on T, which here are still denoted by w and t respectively. If in addition T is t-linked over R, the w-operation is a star operation on T [8, Proposition 3.16]. Note that this star operation, being spectral and of finite type [12], is generally smaller than the w-operation on T, that we denote by w' to avoid confusion. **Proposition 1.5.** Let R be a PvMD and T a t-linked overring of R. Then T is a PvMD and w = t = t' = w' on T, where w' and t' denote respectively the w-operation and the t-operation on T.

Proof. When R is a PvMD also T is a PvMD [30, Theorem 3.8 and Corollary 3.9]. In addition, if R is a PvMD the two semistar operations w and t coincide [11, Theorem 3.1 $((i) \Rightarrow (vi))$]. Hence w = t and w' = t' as star operations on T.

We next show that t = t' on T. Let I be a nonzero ideal of T. Clearly, $I_t \subseteq I_{t'}$. On the other hand, we have $I_{t'} = \bigcap \{IT_M; M \in t' - \operatorname{Max}(T)\}$. Since $I_t = \bigcap \{I_tT_N; N \in t - \operatorname{Max}(T)\}$ [22, Proposition 4], to show that $I_{t'} \subseteq I_t$ it suffices to show that $t - \operatorname{Max}(T) \subseteq t' - \operatorname{Max}(T)$.

If $N \in t$ -Max(T), we have that $(N \cap R)_t \subseteq N_t \cap R_t = N \cap R$. Hence $N \cap R$ is a *t*-prime of R and $T_N \supseteq R_{N \cap R}$ are valuation domains. It follows that N is a *t*'-prime of T. In addition N is *t*'-maximal because it is *t*-maximal and each *t*'-prime of T is also a *t*-prime.

If I is a w-ideal of R, then it is easily shown that $E(I)_w = E(I)$. Thus E(I) is a t-linked overring of R. It follows that, when R is a PvMD, by Proposition 1.5, E(I) is a PvMD and w = t = t' = w'on E(I).

Proposition 1.6. Let R be a PvMD. Then:

- (1) If I is a strong t-ideal of R, then $E(I) = S \cap T$, where $S = \bigcap_{P \in Min(I)} R_P$ and $T := \bigcap_{M \in t-Max(R), M \not\supseteq I} R_M$.
- (2) If P is a t-prime of R which is not t-invertible, then $E(P) = (R : P) = R_P \cap T$, where $T := \bigcap_{M \in t-\operatorname{Max}(R), M \not\supseteq P} R_M$.
- (3) If P is a t-prime of R which is not t-invertible and R satisfies the ascending chain condition on radical t-ideals, then P is t-maximal in E(P).

Proof. (1) follows from [27, Theorem 4.5].

(2) If P is not t-invertible, then P is strong by [26, Proposition 2.3 and Lemma 1.2]. Whence (2) follows from (1).

(3) By part (2), E(P) = (R : P). Since E(P) is t-linked over R, then E(P) is t-flat on R (that is $E(P)_Q = R_{Q \cap R}$ for each t-prime ideal Q of E(P)) [31, Proposition 2.10] and P is a t-ideal of E(P) (Proposition 1.5). Let Q be a t-prime of E(P) properly containing P. By t-flatness we can write $Q = (P'E(P))_t$, where $P' = Q \cap R$ is a t-prime of R properly containing P [6, Proposition 2.4]. By the ascending chain condition on radical t-ideals, $P' = \operatorname{rad}(J_t)$ for some finitely generated ideal J [6, Lemma 3.7]. Since $P \subsetneq P'$, by checking t-locally, we get that $P \subsetneq J_t$. We have $R = (J(R : J))_t \subseteq (J(R : P))_t = (JE(P))_t \subseteq (P'E(P))_t = Q$. A contradiction. Hence P = Q and so P is a t-maximal ideal of E(P).

Lemma 1.7. Let R be a PvMD satisfying the tRTP-property. If P is a branched t-prime of R which is not t-invertible, then $R_P \not\supseteq T$, where $T = \bigcap_{M \in t-\operatorname{Max}(R), M \not\supseteq P} R_M$.

Proof. If $R_P \supseteq T$ then, by Proposition 1.6 (2), T = E(P). Let Q be a P-primary ideal of R. Since P is a t-ideal, we may assume that Q is a t-ideal. We have $QT \subseteq PT = P \subseteq R$ and so $QT \subseteq QR_P \cap R = Q$. Hence QT = Q. If M is a t-maximal ideal of R such that $P \nsubseteq M$, then $Q \nsubseteq M$. Thus $(R : Q) \subseteq R_M$ and it follows that $(R : Q) \subseteq T$. Hence Q(R : Q) = Q. Since R is a tRTP-domain, then we must have Q = P. It follows that P is not branched. \Box

Theorem 1.8. Let R be a PvMD. The following conditions are equivalent:

(i) R is a tRTP-domain;

- (ii) Each branched t-prime P contains a finitely generated ideal J such that $J \subseteq P$ and $J \nsubseteq M$, for each $M \in t$ -Max(R) not containing P;
- (iii) Each branched t-prime is the radical of a v-finite divisorial ideal;
- (iv) Each nonzero principal ideal has at most finitely many minimal (t)-primes;
- (v) Each nonzero finitely generated ideal has at most finitely many minimal t-primes;
- (vi) Each v-finite divisorial ideal has at most finitely many minimal (t-)primes.

Proof. (i) \Rightarrow (ii). Let P be a branched t-prime of R. If P is t-invertible, then P is v-finite and there is nothing to prove. If P is not t-invertible, then $E(P) = (R : P) = R_P \cap T$, where $T = \bigcap_{M \in t-\operatorname{Max}(R), M \not\supseteq P} R_M$ (Proposition 1.6 (2)). Since $R_P \not\supseteq T$ (Lemma 1.7), there exists a finitely generated ideal J such that $J \subseteq P$ and $J \not\subseteq M$, for each $M \in t-\operatorname{Max}(R)$ not containing P [7, Lemma 3.6].

 $(ii) \Rightarrow (iii)$ Let P be a branched t-prime of R and J as in the hypothesis. By Proposition 1.2, P is minimal over a finitely generated ideal H. Hence P is the radical of the v-finite divisorial ideal $(J+H)_v$.

 $(iii) \Rightarrow (iv)$. Let $x \in R$ be a nonzero nonunit and let $\{P_{\alpha}\}$ be the set of minimal primes of xR. By Proposition 1.2 each P_{α} is branched. Hence by hypothesis each P_{α} is the radical of a *v*-finite divisorial ideal. It follows from Proposition 1.4 that $\{P_{\alpha}\}$ is a finite set.

 $(iv) \Leftrightarrow (v) \Leftrightarrow (vi)$ follow from Lemma 1.1.

 $(iv) \Rightarrow (iii)$. Let P be a branched t-prime. By Proposition 1.2, P is minimal over a v-finite divisorial ideal I. Since I has finitely many minimal primes, then P is the radical of a v-finite divisorial ideal by [15, Lemma 2.13].

 $(iii) \Rightarrow (i)$. By [35, Theorem 15] it is enough to show that for each strong *t*-ideal *I* and each minimal prime *P* of *I* we have $IR_P = PR_P$.

Assume that $IR_P \subsetneq PR_P$. Then P is branched, because R_P is a valuation domain. Thus $P = \operatorname{rad}(H_v)$ for some finitely generated ideal H. Let $a \in P$ be such that $IR_P \subsetneq aR_P \subseteq PR_P$ and set J = H + aR. Then P is the radical of J_v . By checking t-locally, we have that $I \subseteq J_v$. In fact, let $M \in t$ -Max(R). If $P \nsubseteq M$, then $IR_M \subseteq R_M = J_vR_M$. If $P \subseteq M$, then $IR_M \subseteq IR_P$. Hence $a \notin IR_M$ and $IR_M \subsetneq J_vR_M$. Since I is strong, by Proposition 1.6(1), $(R:I) = E(I) \subseteq R_P$. Whence $J(R:J) \subseteq P(R:I) \subseteq PR_P$ and so $J(R:J) \subseteq P$. A contradiction because J is t-invertible.

Since in a Prüfer domain the *t*-operation is trivial, we get the following corollary, due to T. Lucas. The equivalence $(i) \Leftrightarrow (ii)$ is [33, Theorem 23], while $(i) \Leftrightarrow (iv)$ is, to our knowledge, unpublished.

Corollary 1.9. Let R be a Prüfer domain. The following conditions are equivalent:

- (i) R is a RTP-domain;
- (ii) Each branched prime is the radical of a finitely generated ideal;
- (iii) Each principal ideal has at most finitely many minimal primes;
- (iv) Each finitely generated ideal has at most finitely many minimal primes.

The following theorem was stated for Prüfer domains in [33, Corollaries 25 and 26].

Theorem 1.10. Let R be a PvMD.

- (1) If R is a t##-domain, then R is a tRTP-domain.
- (2) If R is a tRTP-domain and each t-prime is branched, then R is a t##-domain.

Proof. A PvMD R has the t##-property if and only if, for each t-prime ideal P, there exists a finitely generated ideal $J \subseteq P$ such that each t-maximal ideal containing J must contain P [15, Proposition 2.8]. Hence we can apply Theorem 1.8, $(i) \Leftrightarrow (ii)$.

Theorem 1.11. Let R be a PvMD. The following conditions are equivalent:

- (i) R satisfies the ascending chain condition on radical t-ideals;
- (ii) R is a tRTP-domain satisfying the ascending chain condition on prime t-ideals;
- (iii) R is a t##-domain satisfying the ascending chain condition on prime t-ideals;
- (iv) R is a t##-domain and each t-prime is branched.

Proof. $(i) \Leftrightarrow (iii) \Leftrightarrow (iv)$ by [15, Proposition 2.14].

 $(ii) \Leftrightarrow (iii)$ By Proposition 1.2, the ascending chain condition on prime *t*-ideals implies that each *t*-prime of *R* is branched. Hence we can apply Theorem 1.10.

Recall that a domain R has finite character (respectively, t-finite character) if each nonzero element of R belongs to at most finitely many maximal (respectively, t-maximal) ideals. A domain with finite character such that each nonzero prime ideal is contained in a unique maximal ideal was called by E. Matlis an *h*-local domain. Following [1], we say that R is a weakly Matlis domain if R has t-finite character and each t-prime ideal is contained in a unique t-maximal ideal.

Theorem 1.12. Let R be a PvMD and consider the following conditions:

- (i) R is a weakly Matlis domain;
- (ii) R has t-finite character;
- (iii) R has the t##-property;
- (iv) R is a tRTP-domain.

Then $(i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (iv)$.

If in addition each t-prime ideal of R is contained in a unique t-maximal ideal, all these conditions are equivalent.

Proof. $(i) \Rightarrow (ii)$ by definition.

 $(ii) \Rightarrow (iii)$. If R has t-finite character, for each $\Lambda \subseteq t$ -Max(R) the multiplicative system of ideals $\mathcal{F}(\Lambda) := \{I; I \notin M, \text{ for each } M \in \Lambda\}$ is finitely generated [14, Proposition 2.7]. We conclude that R is a t # #-domain by applying [15, Proposition 2.8].

 $(iii) \Rightarrow (iv)$ By Theorem 1.10 (1).

Now assume that each t-prime of R is contained in a unique t-maximal ideal. Then clearly conditions (i) and (ii) are equivalent.

 $(iv) \Rightarrow (ii)$ By Theorem 1.8, for each nonzero nonunit $x \in R$, the ideal xR has finitely many minimal (t)-primes. Since each t-prime is contained in a unique t-maximal ideal, then x is contained in finitely many t-maximal ideals.

When R is a Prüfer domain, for d = t, from Theorem 1.11 we get [24, Theorem 2.7] and from Theorem 1.12 we get [36, Proposition 3.4].

Remark 1.13. The hypothesis that R is a PvMD in Theorems 1.11 and 1.12 cannot be relaxed. In fact each Noetherian domain is a t##-domain [15, Proposition 2.4], but it is not necessarily a tRTP-domain [24, Corollary 2.2].

A strongly discrete valuation domain is a valuation domain such that each nonzero prime ideal is not idempotent [10, p. 145] and a strongly discrete Prüfer domain is a domain whose localizations at nonzero prime ideals are strongly discrete valuation domains; equivalently a domain such that $P \neq P^2$ for each nonzero prime ideal P [10, Proposition 5.3.5]. We say that a PvMD R is strongly discrete if R_P is a strongly discrete valuation domain for each t-prime ideal P of R; equivalently, if $(P^2)_t \neq P$, for each $P \in t$ -Spec(R) [9, Lemma 3.4]. Generalized Krull domains were introduced by the first author in [6] and can be defined as strongly discrete PvMDs satisfying the ascending chain condition on radical t-ideals [6, Theorem 3.5 and Lemma 3.7]. In the Prüfer case, that is for d = t, this class of domains coincides with the class of generalized Dedekind domains introduced by N. Popescu in [40]. A Krull domain is a generalized Krull domain of t-dimension one [6, Theorem 3.11].

Theorem 1.14. Let R be an integral domain. The following conditions are equivalent:

- (i) R is a generalized Krull domain;
- (ii) R is a strongly discrete PvMD satisfying the t##-property;
- (iii) R is a strongly discrete PvMD satisfying the tRTP-property.

Proof. $(i) \Rightarrow (ii)$ follows from Theorem 1.11 and $(ii) \Rightarrow (iii)$ follows from Theorem 1.10(1).

 $(iii) \Rightarrow (i)$. By Theorem 1.8, R is a strongly discrete PvMD such that each proper v-finite divisorial ideal has finitely many minimal primes. Hence R is a generalized Krull domain by [6, Theorem 3.9].

In the Prüfer case, for d = t, we recover from Theorem 1.14 a well known characterization of generalized Dedekind domains, see for example [10, Theorem 5.5.4].

Remark 1.15. Any *w*-divisorial domain is a t#-domain. In fact, clearly all the *t*-maximal ideals of a *w*-divisorial domain are divisorial; hence we can apply [15, Theorem 1.2].

If R is a domain such that $R_{\mathcal{F}(\Lambda)} := \bigcap_{P \in \Lambda} R_P$ is w-divisorial, for each set Λ of pairwise incomparable t-primes, then $\mathcal{F}(\Lambda)$ is v-finite by [9, Proposition 2.2]. Thus $t \operatorname{-Max}(R_{\mathcal{F}(\Lambda)}) = \{P_{\mathcal{F}(\Lambda)}; P \in \Lambda\}$ [9, Lemma 2.1]. It follows that, given two different sets Λ_1 and Λ_2 of pairwise incomparable t-primes of R, we have $R_{\mathcal{F}(\Lambda_1)} \neq R_{\mathcal{F}(\Lambda_2)}$. Therefore R is a t # #-domain.

Conversely, it is not true that a t##-domain is w-divisorial. In fact each Noetherian domain has the t##-property [15, Proposition 2.4], but a w-divisorial Noetherian domain must have t-dimension one [9, Theorem 4.2].

An integral domain R is an almost Krull domain if R_M is a rank-one discrete valuation domain for each t-maximal ideal M of R. Almost Krull domains were studied by Kang under the name of t-almost Dedekind domains in [30, Section IV]. A Krull domain is an almost Krull domain with t-finite character. In dimension one, the class of almost Krull domains coincides with the class of almost Dedekind domains introduced by R. Gilmer [17]. Gilmer showed that an almost Dedekind domain satisfying the #-property must be Dedekind [19, Theorem 3]. Next, we extend this result to almost Krull domains. First, we give the following characterization of almost Krull domains, which follows directly from the definitions.

Lemma 1.16. Let R be an integral domain. Then R is an almost Krull domain if and only if R is a strongly discrete PvMD of t-dimension one.

Theorem 1.17. Let R be an integral domain. Then the following conditions are equivalent:

- (i) R is an almost Krull domain satisfying the t#-property;
- (ii) R is an almost Krull domain satisfying the t##-property;
- (iii) R is a Krull domain.

Proof. $(i) \Rightarrow (ii)$ Since an almost Krull domain has t-dimension one (Lemma 1.16).

 $(ii) \Rightarrow (iii)$ By Lemma 1.16 and Theorem 1.14, if (ii) holds, R is a generalized Krull domain of t-dimension 1. Hence R is a Krull domain by [6, Theorem 3.11].

 $(iii) \Rightarrow (i)$ Follows from Theorem 1.14.

We end this Section by putting into evidence that the t##-property and the tRTP-property of a PvMD are related respectively to the ##-property and the RTP-property of its t-Nagata ring.

Theorem 1.18. Let R be a PvMD. Then:

- (1) R is a t#-domain if and only if $R\langle X \rangle$ is a #-domain.
- (2) R is a t##-domain if and only if $R\langle X \rangle$ is a ##-domain.
- (3) R is a tRTP-domain if and only if $R\langle X \rangle$ is a RTP-domain.

Proof. (1) follows from [15, Theorem 3.6]. The proof of (2) is similar and it is obtained by using Proposition 1.3 and the characterization of Prüfer ##-domains and PvMDs satisfying the t##-property proved respectively in [20, Theorem 3] and [15, Proposition 2.8 (7)].

(3) Follows from Proposition 1.3, Theorem 1.8 (v) and Corollary 1.9 (iv).

2. w-Divisoriality and w-stability

The notion of w-divisoriality has been studied in [9]. A domain R is an integrally closed wdivisorial domain if and only if it is a weakly Matlis PvMD such that each t-maximal ideal is t-invertible [9, Theorem 3.3] and R is an integrally closed domain such that each t-linked overring is w-divisorial if and only if it is a weakly Matlis strongly discrete PvMD, equivalently R is a wdivisorial generalized Krull domain [9, Theorem 3.5]. We now introduce the notion of w-stability and show that in PvMDs w-divisoriality and w-stability are strictly related; thus extending some results proved by B. Olberding for Prüfer domains.

As before, if T is a t-linked overring of R, we denote by w the star operation induced on T by the w-operation on R and by w' the w-operation on T. We say that a w-ideal I of R is w-stable if I is w-invertible in the (t-linked) overring E(I) := (I : I), that is if $(I(E(I) : I))_w = E(I)$, and we say that R is w-stable if each w-ideal of R is w-stable.

Our first result is a generalization of [39, Theorems 3.3 and 3.5] and shows in particular that the study of w-divisorial domains can be reduced to the t-local case. We recall that a valuation domain is stable if and only if it is strongly discrete [10, Proposition 5.3.8].

Lemma 2.1. Let R be a quasi-local domain. Then a nonzero ideal I of R is stable if and only if $I^2 = xI$ for some $x \in I$.

Proof. This follows from [39, Lemma 3.1] and [10, Lemma 7.3.4].

Theorem 2.2. Let R be an integral domain. The following conditions are equivalent:

- (i) R is w-stable;
- (ii) Each w-ideal I of R is divisorial in E(I);
- (iii) R has t-finite character and R_M is stable for each t-maximal ideal M of R.

Proof. $(i) \Rightarrow (ii)$. Let I be a w-ideal of R. Denote by v' the v-operation on E(I). Since I is w-stable, we have $E(I) = (I(E(I):I))_w \subseteq (I_{v'}(E(I):I))_w \subseteq E(I)$. Hence $(I_{v'}(E(I):I))_w = E(I)$ and $I = IE(I) = (I(E(I):I)I_{v'})_w = ((I(E(I):I))_w I_{v'})_w = I_{v'}$.

 $(ii) \Rightarrow (i)$. Let *I* be a *w*-ideal of *R* and set J = (E(I) : I). Proceeding like in the proof of [39, Theorem 3.5 $((2) \Rightarrow (1))$], we have (E(I) : IJ) = E(I) and hence $E((IJ)_w) = E(I)$. Thus $(IJ)_w$ is a divisorial ideal of E(I). It follows that $(IJ)_w = (E(I) : (E(I) : IJ)) = E(I) : E(I) = E(I)$, that is *I* is a *w*-stable ideal.

 $(i) \Rightarrow (iii)$ Let M be a t-maximal ideal of R and let $I = JR_M$ be a nonzero ideal of R_M , where J is an ideal of R which can be assumed to be a w-ideal (since $J_wR_M = JR_M$). By w-stability in R, $(J(E(J) : J))_w = E(J)$; in particular, $J(E(J) : J)R_M = E(J)R_M$. Since $1 \in E(J)R_M = J(E(J) : J)R_M \subseteq I(E(I) : I) \subseteq E(I)$, then I(E(I) : I) = E(I). Hence I is a stable ideal of R_M and therefore R_M is stable.

We next show that R has t-finite character. Let M be a t-maximal ideal of R. Since R_M is a quasi-local stable domain, then $M^2R_M = mMR_M$ for some $m \in M$ (Lemma 2.1). Set I(M) :=

 $mR_M \cap R$. The ideal I(M) is a t-ideal of R and $M^2 \subseteq I(M)$. Hence M is the only t-maximal ideal of R containing I(M). From this, and by checking t-locally, we get (I(M) : I(M)) = R. Since R is w-stable, I(M) is a w-invertible ideal of R. Thus I(M) is divisorial.

Now, let $\{M_{\alpha}\}$ be a family of t-maximal ideals of R such that $\bigcap M_{\alpha} \neq (0)$. We want to show that $\{M_{\alpha}\}$ is a finite family.

Set $I_{\alpha} := I(M_{\alpha})$ and let $J_{\alpha} := (\Sigma_{\beta \neq \alpha}(R : I_{\beta}))_w$. Note that J_{α} is a fractional ideal of R since $\bigcap_{\beta \neq \alpha} I_{\beta} \supseteq \bigcap_{\beta \neq \alpha} M_{\beta}^2 \neq 0$. We claim that $(J_{\alpha} : J_{\alpha}) = R$. To show this, we prove that $(J_{\alpha} : J_{\alpha}) \subseteq R_M$ for each t-maximal ideal M of R. Let $x \in (J_{\alpha} : J_{\alpha})$. We first assume that $M \notin \{M_{\beta}\}_{\beta \neq \alpha}$. Since M_{β} is the only t-maximal ideal of R containing I_{β} , then $(R : I_{\beta}) \subseteq R_M$ for each $\beta \neq \alpha$. Hence $x \in xJ_{\alpha} \subseteq J_{\alpha} \subseteq R_M$. If $M = M_{\gamma}$ for some $\gamma \neq \alpha$. We have $x(R : I_{\gamma}) \subseteq (\Sigma_{\beta \neq \alpha}(R : I_{\beta}))_w$, and since I_{γ} is w-invertible, then $x \in (\Sigma_{\beta \neq \alpha}(I_{\gamma}(R : I_{\beta}))_w)_w$. Moreover, for $\beta = \gamma$, we have $(I_{\gamma}(R : I_{\gamma}))_w = R$, and for $\beta \neq \gamma$, $(R : I_{\beta}) \subseteq R_{M_{\gamma}}$. Hence $(\Sigma_{\beta \neq \alpha}(I_{\gamma}(R : I_{\beta}))_w)_w \subseteq R_{M_{\gamma}}$. Thus $x \in R_M$, which prove the claim.

Now, for each α , set $T_{\alpha} := \bigcap_{\beta \neq \alpha} M_{\beta}$. We claim that $T_{\alpha} \not\subseteq N$ for each *t*-maximal ideal $N \notin \{M_{\beta}\}_{\beta \neq \alpha}$. By the *w*-stability, J_{α} is a *w*-invertible ideal of $(J_{\alpha} : J_{\alpha}) = R$. In particular, J_{α} is *w*-finite. Thus $(R : J_{\alpha})_N = (R_N : J_{\alpha N}) = R_N$ (since $I_{\beta} \not\subseteq N$ for each $\beta \neq \alpha$). On the other hand, we have $(R : J_{\alpha})_N = (R : \Sigma_{\beta \neq \alpha}(R : I_{\beta}))R_N = (\bigcap_{\beta \neq \alpha} I_{\beta})R_N$. Thus $\bigcap_{\beta \neq \alpha} I_{\beta} \not\subseteq N$. Since $\bigcap_{\beta \neq \alpha} I_{\beta} \subseteq T_{\alpha}$, then $T_{\alpha} \not\subseteq N$, in particular, $T_{\alpha} \not\subseteq M_{\alpha}$ for each α .

Now we proceed as in the proof of [25, Theorem 3.1]. Set $T := \Sigma T_{\alpha}$. By the above result T is not contained in any *t*-maximal ideal of R, hence $T_t = R$. Thus $(\sum_{i=1}^n T_i)_t = R$ for some finite subset $\{T_1, \ldots, T_n\}$ of $\{T_{\alpha}\}$. Let $\{M_1, \ldots, M_n\}$ be the corresponding set of *t*-maximal ideals. If $M_{\alpha} \notin \{M_1, \ldots, M_n\}$ for some α , then $\sum_{i=1}^n T_i \subseteq M_{\alpha}$, which is impossible. Hence $\{M_{\alpha}\}$ is finite. Therefore R has *t*-finite character.

 $(iii) \Rightarrow (i)$ Let I be a w-ideal of R and let M_1, \ldots, M_n be the t-maximal ideals of R containing I. Since I is t-locally stable then $IR_{M_i} = J_i E(I_{M_i})$ for some finitely generated ideal $J_i \subseteq I$, $i = 1, \ldots, n$. Choose $y \in I$ such that $y \notin M$ for each t-maximal ideal $M \neq M_i$ containing the ideal $H := \Sigma J_i$ and consider the ideal J := H + Ry of R. Clearly J is finitely generated. One can easily check that $IR_N = JE(I_N)$ for each t-maximal ideal N of R. We next show that $E(I_N) = E(I)_N$ for each t-maximal ideal N of R. Let $x \in E(I_N)$. Since $I_N = JE(I_N)$, then $xJ \subseteq I_N$. Hence $sxJ \subseteq I$ for some $s \in R \setminus N$. Let M be a t-maximal ideal of R. Then $sxI_M = sxJE(I_M) \subseteq IE(I_M) \subseteq I_M$. Thus $sxI_M \subseteq I_M$ for each t-maximal ideal M of R, so that $sxI \subseteq I$. Hence $x \in E(I)_N$. It follows that $E(I_N) = E(I)_N$, for each t-maximal ideal N, and $I = \bigcap_N I_N = \bigcap_N JE(I_N) = \bigcap_N JE(I)_N = (JE(I))_w$.

Finally, $(I(E(I) : I))_N = I_N(E(I) : JE(I))_N = I_N(E(I)_N : JE(I)_N) = I_N(E(I_N) : I_N) = E(I_N) = E(I)_N$, for each t-maximal ideal N. Therefore $(I(E(I) : I))_w = E(I)$ and so I is a w-stable ideal of R.

Proposition 2.3. Let R be a w-stable domain. Then:

- (1) Each t-maximal ideal of R is divisorial.
- (2) t-Spec(R) is treed.
- (3) R satisfies the ascending chain condition on prime t-ideals.

Proof. (1) Let M be a t-maximal ideal of R. If M is not divisorial, then $M_v = R$. Thus E(M) = (R:M) = R and M is t-invertible. Hence M is divisorial, which is impossible.

(2) and (3) follow from Theorem 2.2 because a quasi-local stable domains has these properties [39, Theorem 4.11]. \Box

The previous proposition shows that w-stable domains have some properties in common with generalized Krull domains [6]. We now prove that w-stability of t-primes enforces a PvMD to be a generalized Krull domain. For Prüfer domains, this follows from [13, Theorem 5] or [36, Theorem 4.7].

Theorem 2.4. Let R be a PvMD. The following conditions are equivalent:

- (i) R is a generalized Krull domain;
- (ii) Each radical t-ideal of R is divisorial and each divisorial ideal is w-stable;
- (iii) Each radical t-ideal of R is w-stable;
- (iv) Each t-prime ideal of R is w-stable.

Proof. We shall freely use Proposition 1.5.

 $(i) \Rightarrow (ii)$. Since t-Spec(R) is treed and a t-ideal of R has finitely many minimal primes, a radical t-ideal of R is a t-product of finitely many t-primes [7, Lemma 2.5]. Hence each radical t-ideal of R is divisorial [7, Proposition 3.1].

Let I be a divisorial ideal of R. If I is t-invertible, hence w-invertible, then E(I) = R and so I is w-stable. If I is not t-invertible, then consider the ideal $H := (I(R : I))_w$. By [7, Proposition 2.6], we have $H = (P_1 \cdots P_n)_w$, where $n \ge 1$ and each P_i is a strong t-prime. Thus $E(P_i) = (R :$ $P_i) \subseteq (R : H) = (R : I(R : I)) = ((R : (R : I)) : I) = E(I)$. Since P_i is t-maximal in $E(P_i)$ (Proposition 1.6) and $E(P_i)$ is t-linked over R then P_i is t-invertible in $E(P_i)$ by [6, Corollaries 3.2 and 3.6]. Hence $(P_i(E(P_i) : P_i))_w = (P_i(E(P_i) : P_i))_t = (P_i(E(P_i) : P_i))_{t'} = E(P_i)$, where t' denotes the t-operation on $E(P_i)$. Thus $P_i E(I)$ is w-invertible in E(I), for each i. It follows that H is w-invertible in E(I) and so I has the same property.

 $(ii) \Rightarrow (iii) \Rightarrow (iv)$ are clear.

 $(iv) \Rightarrow (i)$. Let P be a *t*-prime of R. Since P is *w*-invertible in E(P), then $P \neq (P^2)_w$ and so $P \neq (P^2)_t$. Thus R is a strongly discrete PvMD.

To prove that R is a generalized Krull domain, it is enough to show that R has the t##-property (Theorem 1.14). Let T be a t-linked overring of R and denote by t' the t-operation on T. Let M be a t'-maximal ideal of T. Since T is a PvMD, then T = E(M). The ideal $P = M \bigcap R$ is a t-prime of R and $M = (PT)_{t'} = (PT)_w$ (cf. [31, Proposition 2.10] and [6, Proposition 2.4]). Thus $R \subseteq E(P) \subseteq E(M) = T$. Since P is w-stable, then PT is w-invertible in T, and hence M is w-invertible in T. So, M is a t'-invertible t'-ideal of T (since w = t' in T). In particular M is a divisorial ideal of T. We conclude that T is a t#-domain by applying [15, Theorem 1.2].

It is known that a generalized Dedekind domain need not be stable [13, Example 10]. In fact an integrally closed domain is stable if and only if it is a strongly discrete Prüfer domain with finite character [36, Theorem 4.6]. Hence a generalized Dedekind domain is stable if and only if it has finite character. We now extend these results to generalized Krull domains.

Lemma 2.5. A domain with t-finite character is a strongly discrete PvMD if and only if it is a generalized Krull domain.

Proof. A strongly discrete PvMD is a generalized Krull domain if and only if each nonzero nonunit has finitely many minimal t-primes [6, Theorem 3.9]. We conclude by recalling that in a PvMD two incomparable t-primes are t-comaximal.

Theorem 2.6. Let R be an integral domain. The following conditions are equivalent:

- (i) R is integrally closed and w-stable;
- (ii) R is a w-stable PvMD;
- (iii) R is a strongly discrete PvMD with t-finite character;

- (iv) R is a generalized Krull domain with t-finite character;
- (v) R is a w-stable generalized Krull domain;
- (vi) R is a PvMD with t-finite character and each t-prime ideal of R is w-stable;
- (vii) R is w-stable and each t-maximal ideal of R is t-invertible.

Proof. $(i) \Rightarrow (ii)$. If M is a t-maximal ideal of R, then R_M is an integrally closed stable domain by Theorem 2.2. Hence R_M is a valuation domain [36, Theorem 4.6].

 $(ii) \Rightarrow (iii)$. For each t-maximal ideal M of R, R_M is a valuation stable domain (Theorem 2.2). Hence R_M is a strongly discrete valuation domain [10, Proposition 5.3.8]. The t-finite character follows again from Theorem 2.2.

 $(iii) \Leftrightarrow (iv)$ by Lemma 2.5.

 $(iv) \Rightarrow (v) R_M$ is stable, for each $M \in t$ -Max(R), because it is a strongly discrete valuation domain [10, Proposition 5.3.8]. By the *t*-finite character, R is *w*-stable (Theorem 2.2).

 $(v) \Rightarrow (vii)$ because each t-maximal ideal of a generalized Krull domain is t-invertible [6, Corollary 3.6].

 $(vii) \Rightarrow (i)$ By Theorem 2.2, R_M is a local stable domain, for each $M \in t$ -Max(R). Since M is t-invertible, MR_M is a principal ideal. Hence R_M is a valuation domain [39, Lemma 4.5] and R is integrally closed.

 $(iv) \Leftrightarrow (vi)$ follows from Theorem 2.4.

By Theorem 2.4, each divisorial ideal of a generalized Krull domain is w-stable. Hence a wdivisorial generalized Krull domain is w-stable. Several characterizations of w-divisorial generalized Krull domains were given in [9, Theorem 3.5]. The following theorem says something more in terms of w-stability; similar results for Prüfer domains were obtained by Olberding [36, 38].

Theorem 2.7. Let R be an integral domain. The following conditions are equivalent:

- (i) R is an integrally closed w-divisorial w-stable domain;
- (ii) R is a w-stable w-divisorial PvMD;
- (iii) R is a w-divisorial generalized Krull domain;
- (iv) R is a weakly Matlis w-stable PvMD;
- (v) R is a weakly Matlis strongly discrete PvMD;
- (vi) R is a weakly Matlis generalized Krull domain.

Proof. $(i) \Leftrightarrow (ii)$ by Theorem 2.6.

 $(ii) \Rightarrow (iv)$ because a *w*-divisorial domain is weakly Matlis [9, Theorem 1.5].

- $(iii) \Leftrightarrow (v)$ by [9, Theorem 3.5].
- $(v) \Leftrightarrow (vi)$ by Lemma 2.5.

 $(iii) + (vi) \Rightarrow (i)$ and $(iv) \Leftrightarrow (v)$ by Theorem 2.6, because a weakly Matlis domain has t-finite character.

From Theorems 2.6 and 2.7, we immediately get:

Corollary 2.8. Let R be an integrally closed w-stable domain. Then R is w-divisorial if and only if each nonzero t-prime ideal is contained in a unique t-maximal ideal.

A domain is stable and divisorial if and only if it is totally divisorial [38, Theorem 3.12]. The following is the *t*-analogue of this result in the integrally closed case.

Corollary 2.9. Let R be an integral domain. The following conditions are equivalent:

- (i) R is a w-stable w-divisorial PvMD;
- (ii) R is integrally closed and each t-linked overring of R is w-divisorial.

12

Proof. By [9, Theorem 3.5], if R is integrally closed, each t-linked overring of R is w-divisorial if and only if R is a weakly Matlis strongly discrete PvMD. We conclude by applying Theorem 2.7. \Box

We recall that each overring of a domain R is t-linked if and only if d = w on R [4, Theorem 2.6] and that each overring of a stable domain is stable [39, Theorem 5.1]. We now prove that w-stability is preserved by t-linked extension.

Theorem 2.10. Let R be an integral domain and T a t-linked overring of R. If R is w-stable then T is w'-stable, where w' denotes the w-operation on T.

Proof. We shall use Theorem 2.2. Since $R \subseteq T$ is t-linked, for each t'-maximal ideal M of T, there is a t-maximal ideal N of R such that $R_N \subseteq T_M$ [4, Proposition 2.1]. Hence T_M is an overring of a stable domain and is therefore stable [39, Theorem 5.1].

We next show that T has t-finite character. Let N be a t-maximal ideal of R and let $\{M_{\alpha}\}$ be a family of t-maximal ideals of T such that $\bigcap_{\alpha} M_{\alpha} \neq (0)$ and $M_{\alpha} \cap R \subseteq N$ for each α . Set $S := \bigcap_{\alpha} T_{M_{\alpha}}$. Then S is a stable domain since it is an overring of the stable domain R_N [39, Theorem 5.1]. The prime ideals $P_{\alpha} = M_{\alpha}T_{M_{\alpha}} \cap S$ of S are pairwise incomparable, since $S_{P_{\alpha}} = T_{M_{\alpha}}$ for each α . We have $(0) \neq \bigcap_{\alpha} M_{\alpha} \subseteq \bigcap_{\alpha} P_{\alpha}$, and, since S is treed [39, Theorem 4.11 (ii)] and has finite character [39, Theorem 3.3], then $\{P_{\alpha}\}$ must be finite. Hence $\{M_{\alpha}\}$ is also a finite set. Since R has t-finite character, it follows that T has t-finite character.

We do not know whether the integral closure of a *w*-stable domain is *w'*-stable. In fact the integral closure of a domain R is not always *t*-linked over R [5, Section 4] and we cannot apply Theorem 2.10. However, the *w*-integral closure $R^{[w]} := \bigcup \{ (J_w : J_w); J \text{ a finitely generated ideal of } R \}$ is integrally closed and *t*-linked over R [4, Proposition 2.2 (a)]. Thus we immediately get:

Corollary 2.11. The w-integral closure of a w-stable domain is a w'-stable PvMD.

We end by remarking that, in the integrally closed case, w-divisoriality and w-stability correspond to divisoriality and stability of the t-Nagata ring. We shall make use of Proposition 1.3.

Theorem 2.12. Let R be an integral domain. Then:

- (1) R has t-finite character if and only if $R\langle X \rangle$ has finite character.
- (2) R is a Weakly Matlis PvMD if and only if $R\langle X \rangle$ is an h-local Prüfer domain.
- (3) R is a strongly discrete PvMD if and only if $R\langle X \rangle$ is a strongly discrete Prüfer domain.
- (4) R is a generalized Krull domain if and only if $R\langle X \rangle$ is a generalized Dedekind domain.
- (5) R is an integrally closed w-divisorial domain if and only if $R\langle X \rangle$ is an integrally closed divisorial domain.
- (6) R is an integrally closed w-stable domain if and only if $R\langle X \rangle$ is an integrally closed stable domain.

Proof. Denote by c(f) the content of a polynomial $f(X) \in R[X]$.

(1) We have $Max(R\langle X \rangle) = \{MR\langle X \rangle; M \in t \operatorname{-Max}(R)\}$. Since $f(X) \in MR[X]$ if and only if $c(f)_v \subseteq M$, if R has t-finite character, then $R\langle X \rangle$ has t-finite character. The converse is clear.

(2) Follows from (1) and Proposition 1.3.

(3) For $M \in t$ -Max(R), we have that $R\langle X \rangle_{MR\langle X \rangle} = R[X]_{MR[X]} = R_M(X)$ is a strongly discrete valuation domain if and only if R_M has the same property.

(4) Follows from (3) and Proposition 1.3 by recalling the definitions.

(5) When R is integrally closed, R is divisorial if and only if R is an h-local Prüfer domain such that each maximal ideal is invertible [23, Theorem 5.1] and R is w-divisorial if and only if R is a weakly Matlis PvMD such that each t-maximal ideal is t-invertible [9, Theorem 3.3]. Hence we can

conclude by applying part (2) and recalling that, for each $M \in t$ -Max(R), $MR\langle X \rangle$ is invertible if and only if M is t-invertible [30, Theorem 2.4].

(6) Follows from [36, Theorem 4.6], Theorem 2.6 and statements (1) and (3).

References

- D. D. Anderson and M. Zafrullah, Independent locally-finite intersections of localizations, Houston J. Math. 25 (1999), 433-452.
- [2] H. Bass, On the ubiquity of Gorenstein rings, Math. Z. 82 (1963), 8-28.
- [3] S. Bazzoni and L. Salce, *Warfield Domains*, J. Algebra **185** (1996), 836-868.
- [4] D.E. Dobbs, E.G. Houston, T.G. Lucas, and M. Zafrullah, t-linked overrings and Pr
 üfer v-multiplication domains, Comm. Algebra, 17 (1989), 2835-2852.
- [5] D.E. Dobbs, E.G. Houston, T.G. Lucas, M. Roitman and M. Zafrullah, On t-linked overrings, Comm. Algebra 20 (1992), 1463-1488.
- [6] S. El Baghdadi, On a class of Prüfer v-multiplication domains, Comm. Algebra 30 (2002), 3723-3742.
- [7] S. El Baghdadi, Factorization of divisorial ideals in a generalized Krull domain, Rings, Modules, Algebras and Abelian Groups (Venice, 2002), 149-160, Lecture notes in Pure and appl. Math. 236, Marcel Dekker, New York, 2004.
- [8] S. El Baghdadi and M. Fontana, Semistar linkedness and flatness, Pr
 üfer semistar multiplication domains, Comm. Algebra 32 (2004), 1101-1126.
- [9] S. El Baghdadi and S. Gabelli, *w*-divisorial domains, J. Algebra **285** (2005), 335-355.
- [10] M. Fontana, J. Huckaba, and I. Papick, Prüfer domains, Dekker, New York, 1997.
- [11] M. Fontana, P. Jara and E. Santos, Prüfer *-multiplication domains and semistar operations, J. Algebra Appl. 2 (2003) 21-50.
- [12] M. Fontana and K. A. Loper, Nagata rings, Kronecker function rings and related semistar operations, Comm. Algebra 31 (2003), 4775-4805.
- [13] S. Gabelli, A class of Prüfer domains with nice divisorial ideals, Commutative Ring Theory (Fés, 1995), 313-318, Lecture Notes in Pure and Appl. Math. 185, M. Dekker, New York, 1997.
- [14] S. Gabelli, Prüfer (##)-domains and localizing systems of ideals, Advances in Commutative Ring Theory (Fés, 1997), 391-409, Lecture Notes in Pure and Appl. Math. 205, M. Dekker, New York, 1999.
- [15] S. Gabelli, E. Houston and T. Lucas The t#-property for integral domains, J. Pure Appl. Algebra, 194 (2004), 281-298.
- [16] S. Gabelli and G. Picozza, *Star stable domains*, manuscript.
- [17] R. Gilmer, Integral domains which are almost Dedekind, Proc. Am. Math. Soc. 15 (1964), 813-818.
- [18] R. Gilmer, *Multiplicative ideal theory*, Dekker, New York, 1972.
- [19] R. Gilmer, Overrings of Prüfer domains, J. Algebra 4 (1966), 331-340.
- [20] R. Gilmer and W. Heinzer, Overrings of Prüfer domains. II, J. Algebra 7 (1967), 281-302.
- [21] R. Gilmer and W. Heinzer, Primary ideals with finitely generated radical in a commutative ring, Manuscripta Math. 78 (1993), 201-221.
- [22] M. Griffin, Some results on Prüfer v-multiplication rings, Canad. J. Math. 19 (1967), 71722.
- [23] W. J. Heinzer, Integral domains in which each non-zero ideal is divisorial, Matematika 15 (1968), 164-170.
- [24] W. J. Heinzer and I. J. Papick, The radical trace property, J. Algebra 112 (1988), 110-121.
- [25] E. Houston and M. Zafrullah, Integral domains in which each t-ideal is divisorial, Michigan Math. J. 35 (1988), 291-300.
- [26] E. Houston and M. Zafrullah, On t-invertibility II, Comm. Algebra 17 (1989), 1955-1969.
- [27] E. Houston, S. Kabbaj, T. Lucas and A. Mimouni, When is the dual of an ideal a ring?, J. Algebra 225 (2000), 429-450.
- [28] J. A. Huckaba and I. J. Papick, *Connecting trace properties*, Non-Noetherian Commutative Ring Theory, 313-324, Math. Appl. **520**, Kluwer Acad. Publ., Dordrecht, 2000.
- [29] B.G. Kang, *-Operation on integral domains, Ph.D. Thesis, The University of Iowa, 1987.
- [30] B.G. Kang, Prüfer v-multiplication domains and the ring $R[X]_{N_n}$, J. Algebra **123** (1989), 151-170.
- [31] D.J. Kwak and Y.S. Park, On t-flat overrings, Ch. J. Math. 23 (1995), 17-24.
- [32] J. Lipman, Stable ideals and Arf rings J. Pure Appl. Algebra 4 (1974), 319–336.
- [33] T. Lucas, The radical trace property and primary ideals, J. Algebra 184 (1996), 1093-1112.
- [34] E. Matlis, *Reflexive domains*, J. Algebra 8 (1968), 1-33.

14

RING-THEORETIC PROPERTIES OF PvMDS

- [35] A. Mimouni, Star-operations and trace properties, International Journal of Commutative Rings, 1 (2002), 77-92.
- [36] B. Olberding, Globalizing local properties of Prüfer domains, J. Algebra **205** (1998), 480-504.
- [37] B. Olberding, On the classification of stable domains, J. Algebra 243 (2001), 177-197.
- [38] B. Olberding, Stability, duality, 2-generated ideals and a canonical decomposition of modules, Rend. Sem. Mat. Univ. Padova 106 (2001), 261-290.
- [39] B. Olberding, On the structure of stable domains, Comm. Algebra 30 (2002), 877-895.
- [40] N. Popescu, On a class of Prüfer domains, Rev. Roumanie Math. Pures Appl. 29 (1984), 777-786.
- [41] J.D. Sally and W.V. Vasconcelos, Stable rings and a problem of Bass, Bull. Am. Math. Soc. 79 (1973), 575-576.
- [42] J.D. Sally and W.V. Vasconcelos, *Stable rings*, J. Pure Appl. Alg. 40 (1974), 319-336.
- [43] Wang Fanggui and R.L. McCasland, On w-modules over Strong Mori domains, Comm. Algebra 25 (1997), 1285-1306.
- [44] Wang Fanggui and R.L. McCasland, On Strong Mori domains, J. Pure Appl. Algebra 135 (1999), 155-165.

DEPARTMENT OF MATHEMATICS, FACULTÉ DES SCIENCES ET TECHNIQUES, P.O. BOX 523, BENI MELLAL, MOROCCO

E-mail address: baghdadi@fstbm.ac.ma

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI "ROMA TRE", LARGO S. L. MURIALDO, 1, 00146 Roma, Italy

E-mail address: gabelli@mat.uniroma3.it