# On the Centralizer of K in $U(\mathfrak{g})$

#### BERTRAM KOSTANT\*

Dedicated with respect to Ernest Vinberg on the occasion of his seventieth birthday

Abstract. Let  $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$  be a complexified Cartan decomposition of a complex semisimple Lie algebra  $\mathfrak{g}$  and let K be the subgroup of the adjoint group of  $\mathfrak{g}$  corresponding to  $\mathfrak{k}$ . If H is an irreducible Harish-Chandra module of  $U(\mathfrak{g})$ , then H is completely determined by the finite-dimensional action of the centralizer  $U(\mathfrak{g})^K$  on any one fixed primary  $\mathfrak{k}$  component in H. This original approach of Harish-Chandra to a determination of all H has largely been abandoned because one knows very little about generators of  $U(\mathfrak{g})^K$ . Generators of  $U(\mathfrak{g})^K$  may be given by generators of the symmetric algebra analogue  $S(\mathfrak{g})^K$ . Let  $S_m(\mathfrak{g})^K$ ,  $m \in \mathbb{Z}_+$ , be the subalgebra of  $S(\mathfrak{g})^K$  defined by K-invariant polynomials of degree at most m. For convenience write  $A = S(\mathfrak{g})^K$  and  $A_m$  for the subalgebra of A generated by  $S_m(\mathfrak{g})^K$ . Let Q and  $Q_m$  be the respective quotient fields of A and  $A_m$ . We prove that if  $n = \dim \mathfrak{g}$  one has  $Q = Q_{2n}$ .

We also determine the variety,  $Nil_K$ , of unstable points with respect to the action K on  $\mathfrak{g}$  and show that  $Nil_K$  is already defined by  $A_{2n}$ . As pointed out to us by Hanspeter Kraft this fact together with a result of Harm Derksen (See [D]) implies, indeed, that  $A = A_r$  where  $r = \binom{2n}{2} \dim \mathfrak{p}$ .

#### 1. Introduction

**1.1.** Let  $\mathfrak{g}$  be a complex semisimple Lie algebra. The value taken by the Killing form, B, on  $w, z \in \mathfrak{g}$  will be denoted by (w, z). Let

$$\mathfrak{g} = \mathfrak{k} + \mathfrak{p} \tag{1.1}$$

be a complexified Cartan decomposition and let  $\theta$  be the corresponding complexified Cartan involution. One has that  $[\mathfrak{p},\mathfrak{p}]$  is an ideal of  $\mathfrak{k}$  (and  $\mathfrak{p} + [\mathfrak{p},\mathfrak{p}]$  is an ideal of  $\mathfrak{g}$ ). We will assume that (1.1) is proper in the sense that

$$\mathfrak{k} = [\mathfrak{p}, \mathfrak{p}] \tag{1.2}$$

<sup>\*</sup> Research supported in part by the KG&G Foundation.

(i.e., (1.1) arises from the Cartan decomposition of a real form of  $\mathfrak{g}$  without "compact components"). Let G be the adjoint group of  $\mathfrak{g} = Lie\,\mathfrak{g}$  and let  $K \subset G$  be the subgroup corresponding to  $\mathfrak{k}$ . Of course G has trivial center.

We recall that the centralizer  $U(\mathfrak{g})^K$  of K in the universal enveloping algebra  $U(\mathfrak{g})$  of g played a key role in Harish-Chandra's original approach to the study of certain infinite dimensional representations of g. A critical end product of the theory is the existence of irreducible Harish-Chandra modules. Such a module M is an irreducible  $U(\mathfrak{g})$ -module which not only is completely reducible as a \( \mathbb{t}\)-module but also the primary components are finite dimensional. Any such primary component then defines a finite-dimensional  $U(\mathfrak{g})^K$ module and, remarkably, the entire  $U(\mathfrak{g})$ -module M is completely determined by the action of  $U(\mathfrak{g})^K$  on any one fixed primary component. An early consequence of all of this is Harish-Chandra's subquotient theorem. (For a considerable simplification and clarification of Harish-Chandra's proof see Lepowsky [L] and Lepowsky-McCollum [L-M]. See also Wallach [W] and Vogan [V-1]). With the determination of Harish-Chandra modules reduced to a determination of the finite-dimensional representation theory of  $U(\mathfrak{g})^K$  one might have expected a subsequent development of representation theory along these lines. However this has not been the case although a considerable effort in this direction is seen in [V-1]. The main result of [V-1] is a classification theorem. One major obstacle to making progress with this approach is that the algebra  $U(\mathfrak{g})^K$  is poorly understood. This is more or less attested to by Vogan in [V-2] where he remarks that  $U(\mathfrak{g})^K$  is "hideously complicated". See p. 17 in [V-2]. Also see [K-T] for a glimpse into this complication.

It is not difficult to construct a linear basis of  $U(\mathfrak{g})^K$ . The difficulty lies with its ring structure. Progress would be made if we could pin down a set of (algebra) generators of  $U(\mathfrak{g})^K$ . Indeed focusing on the primary component, given by Vogan's minimal  $\mathfrak{k}$ -type, the corresponding representation of  $U(\mathfrak{g})^K$  is given by a one-dimensional character. Consequently the whole  $U(\mathfrak{g})$ -module M is known as soon as one knows the scalar values assigned to these generators by the character.

The algebra  $U(\mathfrak{g})^K$  has a natural filtration and PBW implies an algebra isomorphism

$$Gr U(\mathfrak{g})^K \cong S(\mathfrak{g})^K$$
 (1.3)

where  $S(\mathfrak{g})^K$  is the finitely generated integral domain of AdK invariants in the symmetric algebra  $S(\mathfrak{g})$ . A set of homogeneous generators of  $S(\mathfrak{g})^K$  then yields a set of generators of  $U(\mathfrak{g})^K$ . The main results of this paper together with a result of Derksen in [D] yields generators of  $S(\mathfrak{g})^K$ .

**1.2.** The adjoint action of  $k \in K$  on  $z \in \mathfrak{g}$  will be denoted by  $k \cdot z$ . If  $z \in \mathfrak{g}$ , then  $K \cdot z$  is Zariski closed if and only if it is closed in the usual Hausdorff topology. Let

$$Cl \mathfrak{g} = \{ z \in \mathfrak{g} \mid K \cdot z \text{ is closed} \}$$

For notational convenience we will put  $A = S(\mathfrak{g})^K$ . Also for notational convenience we identify  $S(\mathfrak{g})$  with the algebra of polynomial functions on  $\mathfrak{g}$  where for any  $x, y \in \mathfrak{g}$  and  $m \in \mathbb{Z}_+$ , one has  $x^m(y) = (x, y)^m$ . Then A is the affine algebra of the affine variety V of all homomorphisms  $A \to \mathbb{C}$ , i.e., all closed points in Spec A. Then, from invariant theory, one knows that

$$V \cong Cl \mathfrak{g}/K$$

i.e.,

$$V$$
 identifies with the set of all closed  $K$ -orbits in  $\mathfrak{g}$ . (1.4)

For any  $z \in Cl \mathfrak{g}$  we will let

$$v_z \in V$$
 be the point corresponding to  $K \cdot z$ . (1.5)

The symmetric algebra  $S(\mathfrak{g})$  is filtered by the subspaces  $S_m(\mathfrak{g})$ ,  $m \in \mathbb{Z}_+$ , where  $S_m(\mathfrak{g}) = \sum_{j=0}^m S^j(\mathfrak{g})$ . Obviously

$$S_m(\mathfrak{g})^K = \sum_{j=0}^m S^j(\mathfrak{g})^K \tag{1.6}$$

But then A is filtered by the subalgebras  $A_m$ ,  $m \in \mathbb{Z}_+$ , where we let

$$A_m$$
 be the subalgebra of  $A$  generated by  $S_m(\mathfrak{g})^K$  (1.7)

Let  $V_m$  be the affine variety corresponding to  $A_m$ . The injection

$$0 \longrightarrow A_m \longrightarrow A \tag{1.8}$$

defines a dominant morphism

$$\gamma_m: V \to V_m \tag{1.9}$$

Let Q (resp. $Q_m$ ) be the quotient field of A (resp. $A_m$ ) and let

$$n = \dim \mathfrak{g} \tag{1.10}$$

The first main result is

**Theorem 1.1**. The dominant morphism  $\gamma_{2n}$  is birational so that

$$Q = Q_{2n} \tag{1.11}$$

In particular any  $h \in A$  is of the form

$$h = f/g (1.12)$$

where  $f, g \in A_{2n}$  and of course  $g \neq 0$ .

**1.3.** Let  $z \in \mathfrak{g}$  be arbitrary. Then z can be uniquely written

$$z = x + y$$
, where  $x \in \mathfrak{k}$  and  $y \in \mathfrak{p}$  (1.13)

Let  $\mathfrak{g}(z)$  be the Lie subalgebra of  $\mathfrak{g}$  generated by x and y. We will use this notation throughout the paper.

In constrast to the closed K-orbits in  $\mathfrak{g}$ , consider the cone of K-unstable points in  $\mathfrak{g}$ . Let

$$Nil_K = \{z \in \mathfrak{g} \mid f(z) = 0, \ \forall \text{ homogeneous } f \in S(\mathfrak{g})^K \text{ of positive degree} \}$$

Since  $S(\mathfrak{g})^G \subset S(\mathfrak{g})^K$  obviously  $Nil_K$  is a subvariety of the nilcone of  $\mathfrak{g}$ .

**Theorem 1.2.** Let  $z \in \mathfrak{g}$ . Then  $z \in Nil_K$  if and only if  $\mathfrak{g}(z)$  is a (nilpotent) Lie algebra of nilpotent elements.

For a number of results about the nilcones of the actions of K, or rather  $K_{\theta}$ , (defined in (2.32) below) on multiple copies of  $\mathfrak{p}$  see [K-W]. Also see [P-3]. For the case we are considering here, Wallach raised the question for a determination of some value of  $m \in \mathbb{Z}_+$  with the property that  $Nil_K$  is given already by the homogeneous elements in  $A_m$  of positive degree. The following result answers this question with the same value of m appearing in Theorem 1.1, namely m = 2n.

**Theorem 1.3.** Let  $z \in \mathfrak{g}$ . Then  $z \in Nil_K$  if and only if

$$f(z) = 0, \ \forall f \in A_{2n} \ of \ positive \ degree.$$

The idea of using a degree which defines  $Nil_K$  (in this case 2n) to determine r such that  $A=A_r$ , goes back to Popov. See [P-1] and [P-2]. Harm Derksen in [D] has sharply reduced Popov's estimate of r. Thus combining Theorem 1.3 with the result in [D] one has

Theorem 1.4 One has

$$A = A_r \tag{1.14}$$

where

$$r = \binom{2n}{2} \dim \mathfrak{p} \tag{1.15}$$

where, we recall  $n = \dim \mathfrak{g}$ .

I thank Hanspeter Kraft for informing me about Derksen's result. Kraft formulated Theorem 1.4, seeing it as an immediate consequence of my Theorem 1.3 and Derksen's result. I also thank Nolan Wallach for motivating me to think about finding an integer m such that  $A_m$  defines  $Nil_K$  (see Theorem 1.3). I also thank him for many conversations about invariant theory.

## 2. The proof of Theorems 1.1, 1.2, 1.3 and 1.4

**2.1.** Let  $\Phi = \Phi(X, Y)$  be the free Lie algebra, over  $\mathbb{C}$  on two generators X, Y. The Lie algebra  $\Phi$  is naturally graded over  $\mathbb{Z}_+$  with homogeneous spaces  $\Phi^j$ . It is then clearly filtered by the subspaces  $\Phi_m$ ,  $m \in \mathbb{Z}_+$ , where

$$\Phi_m = \sum_{j=0}^m \Phi^j \tag{2.1}$$

Clearly

$$\Phi_{m+1} = \Phi_m + [X, \Phi_m] + [Y, \Phi_m] \tag{2.2}$$

Using notation introduced in §1.3 one then has a Lie algebra epimorphism,

$$\xi_z : \Phi \to \mathfrak{g}(z), \text{ where } \xi_z(X) = x \text{ and } \xi_z(Y) = y$$
 (2.3)

The Lie subalgebra  $\mathfrak{g}(z)$  of  $\mathfrak{g}$  is filtered by the subspaces  $\mathfrak{g}_m(z)$  where we put  $\mathfrak{g}_m(z) = \xi_z(\Phi_m)$ . By (2.2) one has

$$\mathfrak{g}_{m+1}(z) = \mathfrak{g}_m(z) + [x, \mathfrak{g}_m(z)] + [y, \mathfrak{g}_m(z)]$$
(2.4)

**Proposition 2.1.** For any  $z \in \mathfrak{g}$  one has

$$\mathfrak{g}_{n-1}(z) = \mathfrak{g}(z) \tag{2.5}$$

**Proof.** It follows immediately from (2.4) that  $\mathfrak{g}(z) = \mathfrak{g}_m(z)$  in case

$$\mathfrak{g}_m(z) = \mathfrak{g}_{m+1}(z) \tag{2.6}$$

Indeed (2.6) implies that  $\mathfrak{g}_k(z) = \mathfrak{g}_m(z)$  for all  $k \in \mathbb{Z}_+$  where  $k \geq m$ .

The statement of the proposition is obviously true if  $\dim \mathfrak{g}_1(z) \leq 1$ . We can therefore assume  $\dim \mathfrak{g}_1(z) = 2$ . We refer to the equality (2.6) as "stability at m". If one does not have stability at m then clearly

$$\dim \mathfrak{g}_{m+1}(z) > m+1 \tag{2.7}$$

But then nonstability at n-1 yields the contradictory statement  $\dim \mathfrak{g}_n(z) > n = \dim \mathfrak{g}$ . Hence one necessarily has stability at n-1. QED

**2.2.** If z = x + y is the decomposition (1.13) for  $z \in \mathfrak{g}$ , then obviously  $k \cdot z = k \cdot x + k \cdot y$  is the decomposition (1.13) for  $k \cdot z$  for any  $k \in K$ . The following simple statement is important for us.

**Proposition 2.2.** Let  $T, T' \in \Phi_n$ . Then  $f_{T,T'} \in S_{2n}(\mathfrak{g})^K$  where, for  $z \in \mathfrak{g}$ ,

$$f_{T,T'}(z) = (\xi_z(T), \xi_z(T'))$$
 (2.8)

**Proof.** We only have to observe that  $f_{T,T'} \in S_{2n}(\mathfrak{g})$ . The remainder follows from invariance of the Killing form and the fact that for  $W \in \Phi$ ,  $z \in \mathfrak{g}$  and  $k \in K$ ,

$$k \cdot \xi_z(W) = \xi_{k \cdot z}(W) \tag{2.9}$$

QED

Let

$$\mathfrak{g}^{K \, reg} = \{ z \in \mathfrak{g} \mid \mathfrak{g}(z) = \mathfrak{g} \}$$

Thus, by Proposition 2.1,  $z \in \mathfrak{g}^{K reg}$  if and only if

$$\mathfrak{g}_{n-1}(z) = \mathfrak{g}_n(z) \\
= \mathfrak{g} \tag{2.10}$$

One readily constructs some  $z \in \mathfrak{g}$  to show that  $\mathfrak{g}^{K reg}$  is not empty. See Appendix for a proof that  $\mathfrak{g}^{K reg}$  is not empty.

Let  $d(n) = \dim \Phi_n$ . Let  $T_j$ , j = 1, ..., d(n), be a basis of  $\Phi_n$ . The following is a restatement of Proposition 2.1 and (2.10).

**Proposition 2.3.** Let  $z \in \mathfrak{g}$ . Then  $\xi_z(T_j)$ , j = 1, ..., d(n), spans  $\mathfrak{g}(z)$ . In particular  $z \in \mathfrak{g}^{K \ reg}$  if and only if  $\xi_z(T_j)$ , j = 1, ..., d(n), spans  $\mathfrak{g}$ .

As functions on  $\mathfrak{g}$  the entries of the  $d(n) \times d(n)$  matrix M(z) given by

$$M_{i,i}(z) = (\xi_z(T_i), \xi_z(T_i))$$

are in  $S_{2n}(\mathfrak{g})^K$ .

For any  $z\in \mathfrak{g}$  let  $K_z$  be the stabilizer of z with respect to the adjoint action K on  $\mathfrak{g}$ . Let  $\mathfrak{k}_z=Lie\,K_z$ . Clearly

$$\mathfrak{k}_z$$
 is the centralizer of  $\mathfrak{g}(z)$  in  $\mathfrak{k}$  (2.11)

From the semisimplicity of  $\mathfrak{g}$  one then has

$$\mathfrak{k}_z = 0 \text{ for any } z \in \mathfrak{g}^{K reg}$$
 (2.12)

**Theorem 2.4.**  $\mathfrak{g}^{K reg}$  is a nonempty Zariski open subset of  $\mathfrak{g}$ . Furthermore if  $z \in \mathfrak{g}^{K reg}$  then the K-orbit  $K \cdot z$  is closed. That is,

$$\mathfrak{g}^{K\,reg} \subset Cl(\mathfrak{g}) \tag{2.13}$$

Put

$$V^{K reg} = \{ v \in V \mid v = v_z \text{ for some } z \in \mathfrak{g}^{K reg} \}$$
 (2.14)

Then  $V^{K reg}$  is a nonempty Zariski open (and hence dense) subset of V.

**Proof.** Let  $z \in \mathfrak{g}$ . Then clearly

$$rank M(z) \le \dim g(z)$$
 (2.15)

But since the Killing form is nonsingular on  $\mathfrak{g}$  it follows that

$$rank M(z) = dim \mathfrak{g} \iff z \in \mathfrak{g}^{K reg}$$

Let  $z \in \mathfrak{g}^{K \, reg}$  and let  $z' \in \overline{K \cdot z}$ . But then clearly M(z) = M(z') so that  $z' \in \mathfrak{g}^{K \, reg}$ . But then  $k_{z'} = 0$  by (2.12). Thus  $\dim K \cdot z = \dim K \cdot z'$ . This implies that  $K \cdot z$  is closed since the K-orbits on the boundary of  $K \cdot z$  must have dimension smaller than  $\dim K \cdot z$ . But now the determinants of all the  $\dim \mathfrak{g} \times \dim \mathfrak{g}$  minors of M(z) are in A. It is an easy exercise to show that  $\mathfrak{g}^{K \, reg}$  is not empty. (As mentioned above a proof that  $\mathfrak{g}^{K \, reg}$  is not empty is given in the Appendix.) This proves that  $\mathfrak{g}^{K \, reg}$  is a nonempty Zariski open subset of  $\mathfrak{g}$  and  $V^{K \, reg}$  is a nonempty Zariski open subset of V. QED

**Remark 2.5.** Note that since the entries of M(z) are in  $S_{2n}(\mathfrak{g})^K$  the determinants of all the  $\dim \mathfrak{g} \times \dim \mathfrak{g}$  minors of M(z) are, in fact, in  $A_{2n}$ .

**2.3. Proof of Theorem 1.1.** To show that  $\gamma_{2n}$  is birational it suffices, by Theorem 2.4, to prove that there exists a nonempty Zariski open subset  $V_* \subset V$  such that the restriction

$$\gamma_{2n}: V_* \to V_{2n} \tag{2.16}$$

is injective. Theorem 2.4 asserts that  $V^{K reg}$  is a nonempty open subvariety of V. The variety  $V_*$ , to be constructed, will in fact be a nonempty open subvariety of  $V^{K reg}$ . Before constructing  $V_*$  we will first establish certain properties of the restriction

$$\gamma_{2n}: V^{K\,reg} \to V_{2n} \tag{2.17}$$

Let  $z, z' \in \mathfrak{g}^{K reg}$  be such that

$$f(z) = f(z'), \ \forall f \in A_{2n}$$

$$(2.18)$$

We will prove that there exists an automorphism  $\pi$  of  $\mathfrak{g}$ , which commutes with  $\theta$  such that  $z' = \pi(z)$ .

Assume (2.18) is satisfied. For  $T \in \Phi_m$  and j = 1, ..., d(n), let  $f_{T,j} \in A_{2n}$  be defined by putting, for any  $w \in \mathfrak{g}$ ,

$$f_{T,j}(w) = (\xi_w(T), \xi_w(T_j))$$
 (2.19)

But since  $f_{T,j} \in A_{2n}$ , one has

$$f_{T,j}(z) = f_{T,j}(z')$$
 (2.20)

We construct a linear isomorphism

$$\pi: \mathfrak{g} \to \mathfrak{g} \tag{2.21}$$

as follows: Let  $w \in \mathfrak{g}$ . Then, by (2.10), there exists  $T \in \Phi_m$  (obviously not necessarily unique) such that  $\xi_z(T) = w$ . Define (to be shown to be well-defined)

$$\pi(w) = w', \text{ where } w' = \xi_{z'}(T)$$
 (2.22)

To see that  $\pi$  is well-defined we have only to establish that if  $T \in \Phi_m$ , then

$$\xi_z(T) = 0, \iff \xi_{z'}(T) = 0 \tag{2.23}$$

But one has

$$\xi_z(T) = 0, \iff f_{T,j}(z) = 0, \ \forall \ j = 1, \dots, d(n)$$
 (2.24)

The same statement holds when z' replaces z. But then one has (2.23) so that the linear isomorphism  $\pi$  is well-defined, noting also that

$$\pi(z) = z' \tag{2.25}$$

**Lemma 2.6.**  $\pi$  is a Lie algebra automorphism which also commutes with  $\theta$ . That is,  $\pi$  stabilizes both  $\mathfrak{k}$  and  $\mathfrak{p}$ .

**Proof.** Let

$$\mathfrak{u} = \{t \in \mathfrak{g} \mid \pi([t,w]) = [\pi(t),\pi(w)], \ \forall w \in \mathfrak{g}\}$$

Then the Jacobi identity immediately implies that  $\mathfrak{u}$  is a Lie subalgebra of  $\mathfrak{g}$ . Let  $w \in \mathfrak{g}$  be arbitrary. By (2.10) there exists  $T \in \Phi_{n-1}$  such that  $\xi_z(T) = w$ . Let  $T_X = [X, T]$  so that  $T_X \in \Phi_n$ . Define  $T_Y \in \Phi_n$  similarly where Y replaces X. Then

$$\xi_z(T_X) = [x, w]$$

$$\xi_z(T_Y) = [y, w]$$

Let  $\xi_{z'}(T) = w'$  so that  $\pi(w) = w'$ . Also let z' = x' + y' be the decomposition (1.13) when z' replaces z. Then

$$\xi_{z'}(T_X) = [x', w']$$

$$\xi_{z'}(T_Y) = [y', w']$$

Thus the Lie subalgebra  $\mathfrak u$  of  $\mathfrak g$  contains x and y. But then  $\mathfrak u = \mathfrak g$  since x and y generate  $\mathfrak g$ . Hence  $\pi$  is an automorphism. Now let  $m \leq n$  where  $m \in \mathbb Z_+$ . Let  $t_i \in \mathfrak g$ ,  $i = 1, \ldots, m$ , where  $t_i \in \{x,y\}$ . Let

$$w = [t_1, [t_2, [\cdots [t_{m-1}, t_m] \cdots]]$$

Then note that  $w \in \mathfrak{k}$  or  $\mathfrak{p}$  according as the number indices j such that  $t_j = y$  is even or odd. It follows immediately that  $\pi$  stabilizes both  $\mathfrak{k}$  and  $\mathfrak{p}$ . QED

We will next restrict  $\gamma_{2n}$  to a nonempty Zariski open subset  $V_1$  of  $V^{K reg}$  to guarantee that  $\pi$  is an inner automorphism.

One knows the degrees of the generators of  $S(\mathfrak{g})^G$ . The maximum degree is the Coxeter number of some simple component of  $\mathfrak{g}$ . This number is certainly less than n and hence

$$S(\mathfrak{g})^G \subset A_{2n} \tag{2.26}$$

Let  $\Gamma$  be the quotient of the group  $Out \mathfrak{g}$  of outer automorphisms of  $\mathfrak{g}$  by the normal subgroup Inn g = G of inner automorphisms. The group  $\Gamma$  is finite. The image, in  $\Gamma$ , of any  $\alpha \in Out g$  will be denoted by  $\sigma_{\alpha}$ . Clearly  $S(\mathfrak{g})^G$  is stable under the action of Out G on  $S(\mathfrak{g})$ . But this clearly defines a representation of

$$\Gamma \to Aut \, S(\mathfrak{g})^G$$
 (2.27)

The following is well known but we will give a proof for completeness.

**Lemma 2.7.** The representation (2.27) is faithful.

**Proof.** Let  $\alpha \in Out \mathfrak{g}$  and assume that  $\alpha \notin Inn \mathfrak{g}$ . Let  $g \in G$  and put  $\alpha' = Ad g \circ \alpha$ . Then  $\sigma_{\alpha} = \sigma_{\alpha'} \neq 1$ . However g can be chosen so that  $\alpha'$  stabilizes the Weyl chamber C of a split Cartan subalgebra of a split real form of  $\mathfrak{g}$  and  $\alpha'|C$  does not reduce to the identity. However from Weyl group theory one knows that  $S(\mathfrak{g})^G$  separates the points of C. This proves that the image of  $\sigma_{\alpha}$  in (2.27) is not the identity. QED

For any  $1 \neq \sigma \in \Gamma$  choose  $f_{\sigma} \in S(\mathfrak{g})^G$  such that  $f \neq f_{\sigma}$  and let

$$F = \prod_{\sigma \in \Gamma/\{1\}} (f_{\sigma} - \sigma(f_{\sigma}))$$
 (2.28)

putting F = 1 if  $\Gamma$  reduces to the identity. Obviously  $F \in S(\mathfrak{g})^G \subset A_{2n}$ . Let

$$\mathfrak{g}_1^{K \, reg} = \{ z \in \mathfrak{g}^{K \, reg} \mid F(z) \neq 0 \} \tag{2.29}$$

so that  $\mathfrak{g}_1^{K \, reg}$ , by Theorem 2.4, is a nonempty Zariski open subset of  $\mathfrak{g}^{K \, reg}$  and

$$V_1^{K reg} = \{ v \mid v = v_z \text{ for some } z \in \mathfrak{g}_1^{K reg} \}$$
 (2.30)

is a nonempty Zariski open subset of  $V^{K reg}$ . Here we are implicitly using the fact that the intersection of two nonempty Zariski open subsets of an irreducible variety is again a nonempty Zariski open set.

**Lemma 2.8.** Let  $z, z' \in \mathfrak{g}_1^{K \, reg}$  and assume that (2.18) is satisfied. Let  $\pi$  be the  $\mathfrak{g}$ -automorphism of Lemma 2.6. Then  $\pi$  is inner. That is,  $\pi = Ad \, g$  for some  $g \in G$  such that  $Ad \, g$  stabilizes both  $\mathfrak{k}$  and  $\mathfrak{g}$ .

**Proof.** If  $\pi$  is inner there is nothing to prove. Assume  $\pi$  is not inner and let  $1 \neq \sigma \in \Gamma$  be defined by putting  $\sigma = \sigma_{\pi^{-1}}$ . But by (2.25) one has

$$f_{\sigma}(\pi(z)) = f_{\sigma}(z) \tag{2.31}$$

But

$$f_{\sigma}(\pi(z)) = (\pi^{-1}f_{\sigma})(z)$$
$$= (\sigma f_{\sigma})(z)$$

But  $(\sigma f_{\sigma})(z) \neq f_{\sigma}(z)$  since  $F(z) \neq 0$ . This contradicts (2.31). Thus  $\pi$  is inner. QED

Let the notation be as in Lemma 2.8. We will now restrict  $\gamma_{2n}$  even further to finally guarantee that  $g \in K$ .

Taking notation from [K-R] let

$$K_{\theta} = \{ g \in G \mid Ad \, g \text{ stabilizes both } \mathfrak{k} \text{ and } \mathfrak{p} \}$$
 (2.32)

so that, in the notation of Lemma 2.8,  $g \in K_{\theta}$ . Obviously  $K \subset K_{\theta}$ . Let  $Out_{G}^{\mathfrak{k}}$  be the group of all automorphisms of  $\mathfrak{k}$  of the form  $Adg|\mathfrak{k}$  for  $g \in K_{\theta}$  and let  $Inn\mathfrak{k}$  be the group of

all inner automorphisms of  $\mathfrak{k}$ . Obviously  $Inn\,\mathfrak{k}$  is a normal subgroup of  $Out_G\mathfrak{k}$ . One knows that the quotient group  $\Gamma_K = Out_G\mathfrak{k}/Inn\,\mathfrak{k}$  is finite. See Proposition 1, p. 761 in [K-R]. The argument yielding (2.26) readily also implies

$$S(\mathfrak{k})^K \subset A_{2n} \tag{2.33}$$

Also the natural action of  $Out_G \mathfrak{k}$  on  $S(\mathfrak{k})^K$  descends to a representation

$$\Gamma_K \to Aut \, S(\mathfrak{k})^K$$
 (2.34)

The argument establishing Lemma 2.7 is readily modified (to deal with the case where  $\mathfrak{t}$  is only reductive but not semisimple) so that one has

**Lemma 2.9.** The representation (2.34) is faithful.

For each  $1 \neq \tau \in \Gamma_K$  let  $f_{\tau} \in S(\mathfrak{k})^K$  be such that  $f_{\tau} \neq \tau f_{\tau}$ . If  $\Gamma_K$  reduces to the identity put  $F_K = 1$ , otherwise let

$$F_K = \prod_{\tau \in \Gamma_K / \{1\}} (f_\tau - \tau f_\tau)$$
 (2.35)

Let

$$\mathfrak{g}_* = \{ z \in \mathfrak{g}_1^{K \, reg} \mid F_K(z) \neq 0 \} \tag{2.36}$$

and let

$$V_* = \{ v \in V \mid v = v_z \text{ for some } z \in \mathfrak{g}_* \}$$
 (2.37)

Again, since the intersection of two nonempty Zariski open subsets of an irreducible variety is again a nonempty Zariski open set, it follows that  $\mathfrak{g}_*$  is a nonempty Zariski open subset of  $\mathfrak{g}$  and  $V_*$  is a nonempty Zariski open subset of V. The following lemma establishes Theorem 1.1.

**Lemma 2.10.** Let  $z, z' \in \mathfrak{g}^*$  be such that

$$f(z) = f(z') \tag{2.38}$$

for all  $f \in A_{2n}$ . Let  $g \in G$  be given by Lemma 2.8 so that

$$Ad g(z) = z' (2.39)$$

and  $g \in K_{\theta}$  using the notation of (2.32). Then  $g \in K$  so that

$$z' \in K \cdot z \tag{2.40}$$

proving the injectivity of (2.16) and as, noted in the beginning of  $\S 2.3$ , proving Theorem 1.1.

**Proof.** We first prove that  $Adg|\mathfrak{k} \in Inn\mathfrak{k}$ . Assume this is not the case and let  $1 \neq \tau$  be the image of  $Adg^{-1}|\mathfrak{k}$  in  $\Gamma_K$ . Then, by (2.38),

$$f_{\tau}(Ad\,g\,(z)) = f_{\tau}(z) \tag{2.41}$$

But, recalling (2.2),

$$f_{\tau}(Ad g(z)) = f_{\tau}(Ad g(x))$$

$$= (Ad g^{-1} f_{\tau})(x)$$

$$= (\tau f_{\tau})(x)$$

$$= (\tau f_{\tau})(z)$$

But this contradicts (2.41) since  $F_K(z) \neq 0$ . Hence there exists  $k \in K$  such that if  $b = k^{-1} g$ , then b centralizes  $\mathfrak{k}$ . But then both the semisimple element  $b_s$  and the unipotent element  $b_u$  centralize  $\mathfrak{k}$  where  $b = b_s b_u$  is the Jordan decomposition of b. But, as one knows, the centralizer of  $\mathfrak{k}$  in  $\mathfrak{g}$  is commutative, reductive and contained in  $\mathfrak{k}$ . This readily implies that  $b_u = 1$  since the nilpotent element  $log \ b_u$  must commute with  $\mathfrak{k}$ . Thus b is semisimple. Hence b centralizes a Cartan subalgebra  $\mathfrak{h}$  of  $\mathfrak{g}$ . Let  $\mathfrak{g}^b$  be the centralizer of b in  $\mathfrak{g}$  so that  $\mathfrak{h} + \mathfrak{k} \subset \mathfrak{g}^b$ . For any simple component  $\mathfrak{g}_i$  of  $\mathfrak{g}$  let  $\mathfrak{k}_i = \mathfrak{g}_i \cap \mathfrak{g}^b$  and let  $\mathfrak{p}_i$  be the Killing form orthocomplement of  $\mathfrak{k}_i$  in  $\mathfrak{g}_i$ . Since  $\mathfrak{g}^b$  contains  $\mathfrak{h}$  it is clear that  $\mathfrak{g}^b$  is the sum of its intersections with all the simple components of  $\mathfrak{g}$ . It follows then that  $\mathfrak{p}_i$  is Killing form orthogonal to  $\mathfrak{k}$  so that  $\mathfrak{p}_i \subset \mathfrak{p}$ . Hence  $\mathfrak{p}_i + [\mathfrak{p}_i, \mathfrak{p}_i]$  is an ideal in  $\mathfrak{g}_i$ . By simplicity either  $\mathfrak{p}_i = 0$  in which case  $\mathfrak{g}_i = \mathfrak{k}_i$  so that  $\mathfrak{g}_i$  makes no nontrivial contribution to b or  $[\mathfrak{p}_i, \mathfrak{p}_i] = \mathfrak{k}_i \subset \mathfrak{k}$ . Since b is in the subgroup of G corresponding to  $\mathfrak{h}$  it is then clear that  $b \in K$  and hence  $g \in K$ . QED

**2.4. Proof of Theorem 1.2.** Let  $z \in \mathfrak{g}$ . Then one knows from invariant theory that  $K \cdot z$  has a unique closed K-orbit in its closure (this is immediate from (1.4)). Consequently  $z \in Nil_K$  if and only if

$$0 \in \overline{K \cdot z} \tag{2.42}$$

Assume that  $z \in Nil_K$  and let  $k_m \in K$ ,  $m \in \mathbb{Z}_+$ , be a sequence such that  $k_m \cdot z$  converges to 0. But then recalling the decomposition (1.13) one must have that both  $k_m \cdot x$  and  $k_m \cdot y$  also converge to 0. But then obviously  $k_m \cdot w$  converges to 0 for any  $w \in \mathfrak{g}(z)$ . But then (recalling that  $S(\mathfrak{g})^G \subset S(\mathfrak{g})^K$ ) w is nilpotent for any  $w \in \mathfrak{g}(z)$ .

Conversely, assume that every element in  $\mathfrak{g}(z)$  is nilpotent. Then there exists a Borel subalgebra  $\mathfrak{b}$  of  $\mathfrak{g}$  such that  $\mathfrak{g}(z) \subset \mathfrak{n}$  where  $\mathfrak{n}$  is the nilradical of  $\mathfrak{b}$ . Put  $\mathfrak{b}' = \theta(\mathfrak{b})$  so that  $\theta(\mathfrak{n}) = \mathfrak{n}'$  where  $\mathfrak{n}'$  is the nilradical of  $\mathfrak{b}'$ . Let  $\mathfrak{s} = \mathfrak{b} \cap \mathfrak{b}'$  so that  $\mathfrak{s}$  is a solvable subalgebra of  $\mathfrak{g}$  which is stable under  $\theta$ , since  $\theta$  is involutory. Moreover there exists a Cartan subalgebra  $\mathfrak{h}$  of  $\mathfrak{g}$  which is contained in  $\mathfrak{s}$  since the intersection of any two Borel subalgebras contains a Cartan subalgebra. Furthermore from Weyl group theory

$$\mathfrak{s} = \mathfrak{h} + \mathfrak{n} \cap \mathfrak{n}' \tag{2.43}$$

is a Levi decomposition of  $\mathfrak{s}$ . But since  $\mathfrak{g}(z)$  is stable under  $\theta$  one also has

$$\mathfrak{g}(z) \subset \mathfrak{n} \cap \mathfrak{n}' \tag{2.44}$$

But now there exists a regular semisimple element  $u \in \mathfrak{h}$  such that the spectrum of  $ad\mathfrak{u}|\mathfrak{n}$  is a set of positive numbers. In particular the spectrum of  $ad\mathfrak{u}|\mathfrak{n}\cap\mathfrak{n}'$  is again strictly positive. Now let  $u' = \theta(u)$  so that  $u' \in \mathfrak{h}'$  where  $\mathfrak{h}' = \theta(\mathfrak{h})$ . But since  $\mathfrak{s}$  is stable under  $\theta$  one has  $\mathfrak{h}' \subset \mathfrak{s}$ . Interchanging the roles of  $\mathfrak{h}$  and  $\mathfrak{h}'$  it follows that the spectrum of  $ad\mathfrak{u}'|\mathfrak{n}\cap\mathfrak{n}'$  is again strictly positive. But, by Lie's theorem, the adjoint action of  $\mathfrak{s}$  on  $\mathfrak{n}\cap\mathfrak{n}$  may be triangularized. The diagonal entries of both  $ad\mathfrak{u}$  and  $ad\mathfrak{u}'$  on  $\mathfrak{n}\cap\mathfrak{n}$  are positive. Hence the same is true of  $ad\mathfrak{v}$  where  $\mathfrak{v}=\mathfrak{u}+\mathfrak{u}'$ . This however implies that for any  $\mathfrak{w}\in\mathfrak{n}\cap\mathfrak{n}'$ ,

$$exp(-t) v \cdot w$$
 converges to 0 as  $t$  goes to  $+\infty$  (2.45)

(noting that even though v may not be semisimple the nilpotent component of v relative to its Jordan decomposition contributes only polynomial terms in t). But this implies that

$$\mathfrak{n} \cap \mathfrak{n}' \subset Nil_K \tag{2.46}$$

since  $v \in \mathfrak{k}$ . Hence  $z \in Nil_K$  proving Theorem 1.2.

**2.5. Proof of Theorem 1.3.** That is, we prove that if  $z \in \mathfrak{g}$  then  $z \in Nil_K$  if and only if f(z) = 0 for all homogeneous  $f \in A_{2n}$  of positive degree. Of course the "only if"

is obvious since  $A_{2n} \subset A$ . Assume then that  $z \in \mathfrak{g}$  and f(z) = 0 for all homogeneous  $f \in A_{2n}$  of positive degree. But then recalling the  $d(n) \times d(n)$  matrix M(z) of §2.2 one has

$$(\xi_z(T_i), \xi_z(T_j)) = 0$$
 (2.47)

for all  $i, j \in \{1, \dots, d(n)\}$ . But then, by Proposition 2.3, one has

$$tr\,ad\,u\,ad\,v = 0\tag{2.48}$$

for all  $u, v \in \mathfrak{g}(z)$ . Thus, since ad is faithful,  $\mathfrak{g}(z)$  is solvable and hence its adjoint action on  $\mathfrak{g}$  can be triangularized. The nilcone of  $\mathfrak{g}$  intersected with  $\mathfrak{p}$  is just the set of zeros of the polynomials in  $S(\mathfrak{p})^K$  of positive degree (see Proposition 11 in [K-R]). But as one knows the homogeneous generators of  $S(\mathfrak{p})^K$  have the same degrees as the homogeneous generators of the polynomial invariants of the restricted Weyl group operating on a Cartan subspace of  $\mathfrak{p}$  (the symmetric space analogue of Chevalley's theorem). But then one easily has  $S(\mathfrak{p})^K \subset A_{2n}$ . (This follows, for example, from Proposition 23 in [K-R].) But since  $S(\mathfrak{t})^K \subset A_{2n}$  and  $S(\mathfrak{p})^K \subset A_{2n}$  one has that x and y are nilpotent where z = x + y is the decomposition (1.13). Thus the diagonal entries of ad x and ad y are zero. But since x and y generate  $\mathfrak{g}(z)$  the diagonal entries of any element in  $\mathfrak{g}(z)$  are zero. Thus any element in  $\mathfrak{g}(z)$  is nilpotent. Theorem 1.3 then follows from Theorem 1.2. QED

**2.6. Proof of Theorem 1.4.** Theorem 1.3 above and Theorem 1.1 in [D] assert that there exists r such that  $A = A_r$  where  $r \le \max\{2, \frac{3}{8} \dim \mathfrak{p} (2n)^2\}$ . But then Theorem 1.4 follows since  $\frac{1}{2}(x(x-1)) \ge \frac{3}{8} x^2$  for  $x \ge 4$ , and (assuming  $\mathfrak{g} \ne 0$ ), one surely has n > 2. QED

### Appendix

The purpose of this appendix is to show that  $\mathfrak{g}^{K reg}$  is not empty.

**1.1A**. Let  $\mathfrak{g} = \mathfrak{k} + \mathfrak{a} + \mathfrak{n}$  be a complexified Iwasawa decomposition of  $\mathfrak{g}$ , consistent with the complexified Cartan decomposition

$$\mathfrak{g} = \mathfrak{k} + \mathfrak{p} \tag{1.1A}$$

(e.g.  $\mathfrak{a}$  is a complexified Cartan subspace of  $\mathfrak{p}$ ). Let  $R \subset \mathfrak{a}^*$  be the set of restricted roots, and for any  $\nu \in R$  let  $\mathfrak{g}_{\nu} \subset \mathfrak{g}$  be the corresponding restricted root space. Let  $R_+ \subset R$ 

be the set of positive restricted roots defined so that

$$\mathfrak{n} = \oplus_{\nu \in R_+} \mathfrak{g}_{\nu}$$

Let  $\zeta$  be the nonvanishing polynomial function on  $\mathfrak{a}$  defined by putting

$$\zeta = \prod_{\nu,\nu' \in R, \ \nu \neq \nu'} (\nu - \nu') \tag{1.2A}$$

Let  $y \in \mathfrak{a}$  be defined so that

$$\zeta(y) \neq 0$$

Let  $\mathfrak{m}$  be the centralizer of  $\mathfrak{a}$  in  $\mathfrak{k}$ . We recall that  $\theta$  is the complexified Cartan involution corresponding to (1.1A). For  $\nu \in R_+$  let  $x_{\nu} \in \mathfrak{g}_{\nu}$ . Let  $x_{-\nu} \in \mathfrak{g}_{-\nu}$  be defined by putting  $x_{-\nu} = \theta \ x_{\nu}$ . Let  $\widetilde{R} = R \cup \{0\}$  where, here, we regard 0 as the zero linear functional on  $\mathfrak{a}$ . Then  $\widetilde{R}$  is the set of weights for the adjoint action of  $\mathfrak{a}$  on  $\mathfrak{g}$ . Let  $\mathfrak{r}$  be the  $\mathbb{C}$ -span of the set  $\{x_{\nu}\} \nu \in \widetilde{R}$ . Also let  $x = \sum_{\nu \in \widetilde{R}} x_{\nu}$  so that  $x \in \mathfrak{k}$  and also  $x \in \mathfrak{r}$ .

**Remark 1.1A.** Note that, for any  $\nu \in R$ ,  $2\nu$  is a factor of  $\zeta$ , so that  $\nu(y) \neq 0$ .

Let z = x + y and let  $\mathfrak{g}(z)$  be the Lie subalgebra of  $\mathfrak{g}$  generated by x and y. One notes that  $\mathfrak{r}$  is stable under ady and that  $ady|\mathfrak{r}$  is diagonalizable with distinct eigenvalues. In fact clearly  $\mathfrak{r}$  is a cyclic ady module with x as cyclic generator and hence

**Proposition 1.2A.** One has  $x_{\nu} \in \mathfrak{g}(z)$  for any  $\nu \in \widetilde{R}$ .

**1.2A.** The element  $y \in \mathfrak{p}$  will be fixed as in §1.1A. It will be our objective in this section to show that  $x_0$  and  $x_{\nu}$ ,  $\nu \in R_+$  can be chosen, consequently x can chosen, so that  $\mathfrak{g}(z) = \mathfrak{g}$ , i.e.  $z \in \mathfrak{g}^{K reg}$ . This will establish that  $\mathfrak{g}^{K reg}$  is not empty.

Let  $\mathfrak{h}_{\mathfrak{m}}$  be a Cartan subalgebra of  $\mathfrak{m}$  so that  $\mathfrak{h} = \mathfrak{m} + \mathfrak{a}$  is a Cartan subalgebra of  $\mathfrak{g}$ . Let  $\Delta \subset \mathfrak{h}^*$  be the set of roots for  $(\mathfrak{h}, \mathfrak{g})$ , and for each  $\varphi \in \Delta$ , let  $e_{\varphi} \in \mathfrak{g}$  be a corresponding root vector. Obviously  $\mathfrak{g}_{\nu}$  is stable under  $ad\mathfrak{h}$  for any  $\nu \in R$ . Hence there exists a subset  $\Delta_{\nu} \subset \Delta$  such that

$$\mathfrak{g}_{\nu} = \sum_{\varphi \in \Delta_{\nu}} \mathbb{C} e_{\varphi} \tag{1.3A}$$

It is immediate that

$$\Delta_{-\nu} = -\Delta_{\nu} \tag{1.4A}$$

For any  $\nu \in R$  let  $h_{\nu} \in \mathfrak{a}$  be such that, with respect to the Killing form,  $(h, h_{\nu}) = \nu(h)$  for any  $h \in \mathfrak{a}$ . It is clear of course that  $\mathfrak{a}$  is spanned by  $\{h_{\nu} \mid \nu \in R_{+}\}$ .

Let  $P = \frac{1}{2}(1 - \theta)$  so that  $P : \mathfrak{g} \to \mathfrak{p}$  is the projection of  $\mathfrak{g}$  on  $\mathfrak{p}$  with respect to (1.1A). Since  $\mathfrak{g}(z)$  is clearly stable under  $\theta$  for any  $x \in \mathfrak{k}$  it is also stable under P. One easily has

**Lemma 1.3A.** Let  $\nu \in R$  and let  $\varphi \in \Delta_{\nu}$  so that  $-\varphi \in \Delta_{-\nu}$ . Then

$$P[e_{\varphi}, e_{-\varphi}] = c h_{\nu} \tag{1.5A}$$

for some  $c \in \mathbb{C}^{\times}$ .

A useful criterion for K-regularity is given in

**Proposition 1.4A.** For z to be in  $\mathfrak{g}^{K reg}$  it is necessary and sufficient that  $\mathfrak{n} \subset \mathfrak{g}(z)$ .

**Proof.** The necessity is by definition. Assume  $\mathfrak{n} \subset \mathfrak{g}(z)$ . Then  $\mathfrak{g}_{\nu} \in g(z)$  for any  $\nu \in R_+$ . But clearly  $\theta(\mathfrak{g}_{\nu}) = \mathfrak{g}_{-\nu}$  so that  $\mathfrak{g}_{-\nu} \subset \mathfrak{g}(z)$ . But then  $h_{\nu} \in \mathfrak{g}(z)$  for any  $\nu \in R_+$  by Lemma 1.3. Hence  $\mathfrak{a} + \mathfrak{n} \subset \mathfrak{g}(z)$ . But from the Iwasawa decomposition  $P(\mathfrak{a} + \mathfrak{n}) = \mathfrak{p}$ . Thus  $\mathfrak{p} \subset \mathfrak{g}(z)$ . However  $\mathfrak{g} = \mathfrak{p} + [\mathfrak{p}, \mathfrak{p}]$ . Thus  $\mathfrak{g}(z) = \mathfrak{g}$ . QED

Let  $R_+^1$  be the set of all  $\nu \in R_+$  such that  $\dim g_{\nu} = 1$  and let  $R_+^2$  be the complement of  $R_+^1$  in  $R_+$ . Assume  $\nu \in R_+^2$ . Then the weights of  $ad \, \mathfrak{h}_{\mathfrak{m}}$  on  $\mathfrak{g}_{\nu}$  are of the form  $\varphi | \mathfrak{h}_{\mathfrak{m}}$ where  $\varphi \in \Delta_{\nu}$ . Since roots, as weights of  $ad \, \mathfrak{h}$  acting on  $\mathfrak{g}$ , have multiplicity 1 it follows immediately that the weights of  $ad \, \mathfrak{h}_{\mathfrak{m}}$  on  $\mathfrak{g}_{\nu}$  have multiplicity one. Thus if  $\eta_{\nu}$  is the polynomial function on  $\mathfrak{h}_{\mathfrak{m}}$  defined by putting

$$\eta_{\nu} = \prod_{\varphi, \varphi' \in \Delta_{\nu}, \, \varphi \neq \varphi'} (\varphi - \varphi') | \mathfrak{h}_{\mathfrak{m}}$$

$$(1.6A)$$

then  $\eta_{\nu}$  is nonvanishing. One immediately has

**Proposition 1.5A.** Assume  $\nu \in R^2_+$ . Let  $x' \in \mathfrak{h}_{\mathfrak{m}}$  be such that  $\eta_{\nu}(x') \neq 0$ . (Such an element x' exists since  $\eta_{\nu}$  is nonvanishing.) Then  $\mathfrak{g}_{\nu}$  is a cyclic module for ad x'.

We can now exhibit an element  $z \in \mathfrak{g}^{K reg}$ . Recall the notation of §1.1A.

**Theorem 1.6A.** For any  $\nu \in R^1_+$  let  $0 \neq x_{\nu} \in \mathfrak{g}_{\nu}$ . If  $R^2_+$  is empty let  $x_0 = 0$ . If  $R^2_+$ 

is not empty let  $\eta$  be the nonvanishing function on  $\mathfrak{h}_{\mathfrak{m}}$  defined by putting

$$\eta = \prod_{\nu \in R_+^2} \eta_{\nu} \tag{1.7A}$$

Let  $x_0 \in \mathfrak{h}_{\mathfrak{m}}$  be such that  $\eta(x_0) \neq 0$  so that (by Proposition 1.5A)  $\mathfrak{g}_{\nu}$  is a cyclic module for  $ad\ x_0$  for any  $\nu \in R^2_+$ . For  $\nu \in R^2_+$  let  $x_{\nu} \in \mathfrak{g}_{\nu}$  be a cyclic generator of  $\mathfrak{g}_{\nu}$  with respect to the action of  $ad\ x_0$ . Now let  $y \in \mathfrak{a}$  be as in §1.1, and as in §1.1, let  $x = \sum_{\nu \in \widetilde{R}} x_{\nu}$  where we recall  $x_{-\nu} = \theta(x_{\nu})$  for  $\nu \in R_+$  so that  $x \in \mathfrak{k}$ . Then  $\mathfrak{g}(z) = \mathfrak{g}$  where z = x + y.

**Proof.** One has  $x_{\nu} \in \mathfrak{g}(z)$  for any  $\nu \in \widetilde{R}$  by Proposition 1.2A. Thus  $\mathfrak{g}_{\nu} \subset \mathfrak{g}(z)$  for any  $\nu \in R^1_+$ . On the other hand if  $R^2_+$  is not empty then  $\mathfrak{g}_{\nu} \subset \mathfrak{g}(z)$  for  $\nu \in R^2_+$  since the Lie algebra generated by  $x_0$  and  $x_{\nu}$  contains  $\mathfrak{g}_{\nu}$ . Thus  $\mathfrak{n} \subset \mathfrak{g}(z)$  and hence  $z \in \mathfrak{g}^{K \, reg}$  by Proposition 1.4A. QED

### References

- [D] H. Derksen, Polynomial bounds for rings of invariants, *Proc. Amer. Math. Soc.*, **129**, no.4, 955–963
- [K-T] B. Kostant and Juan Tirao, On the structure of certain subalgebras of a universal enveloping algebra, *Trans. Amer. Math. Soc.*, **218**(1976), 133–154
- [K-R] B. Kostant and S. Rallis, Orbits and Representations associated with Symmetric Spaces, Amer. J. Math., 93(1971), No. 3, 753–809
- [K-W] H. Kraft and N. Wallach, On the nullcone of representations of Reductive Groups, Pacific J. Math., 224(2006), 119–140
  - [L] J. Lepowsky, Algebraic results on representations of semisimple Lie groups, Trans.
     Amer. Math. Soc., 176(1973), 1–43
- [L-M] J. Lepowsky and G. McCollum, On the determination of irreducible modules by restriction to a subalgebra, *Trans. Amer. Math. Soc.*, **176**(1973), 44–57
- [P-1] V. Popov, Constructive invariant theory, Astérisque, 87-88(1981), 303-334
- [P-2] V. Popov, The constructive theory of invariants, Math. USSR Izvest., 10(1982), 359-376
- [P-3] V. Popov, The cone of Hilbert nullforms, Steklov Inst. Math. 241(2003), 177–194
- [V-1] D. Vogan, The algebraic structure of representations of semi-simple Lie groups, I, Ann. of Math., **109**(1979), 1–60

[V-2] D. Vogan, Representations of Real Reductive Lie Groups, Birkhäuser, PM 15(1981)

[W] N. Wallach, Real Reductive Groups, I, Academic Press Inc, 132, 1988

Bertram Kostant Dept. of Math. MIT Cambridge, MA 02139

 $E\text{-}mail\ kostant@math.mit.edu$